Binge-like Acquisition of α-pyrrolidinopentiophenone (α-PVP) Self-Administration in Female Rats

Mehrak Javadi-Paydar[1], Yanabel Grant[1], Sophia A. Vandewater[1], Kevin M. Creehan[1], Jacques D. Nguyen[1], Tobin J. Dickerson[2] and Michael A. Taffe[1]

[1]Department of Neuroscience; [2]Department of Chemistry; The Scripps Research Institute; La Jolla, CA, USA

Running Head: α-PVP Binge IVSA

Address Correspondence to: Dr. Michael A. Taffe, Department of Neuroscience, SP30-2400; 10550 North Torrey Pines Road; The Scripps Research Institute, La Jolla, CA 92037; USA; Phone: +1.858.784.7228; Fax: +1.858.784.7405; Email: mtaffe@scripps.edu

Abstract

Rationale: The synthetic cathinone α-pyrrolidinopentiophenone (α-PVP) recently drew attention due to reports of violent and/or bizarre public behavior in users. Association of such behavior with extended binges of drug use motivates additional investigation, particularly since a prior study found that half of male rats experience a binge of exceptionally high intake, followed by sustained lower levels of self-administration during the acquisition of intravenous self-administration (IVSA) of a related drug, 3,4-methylenedioxypyrovalerone.

Objectives: The binge-like acquisition pattern appeared to be novel for rat IVSA, thus the present study was designed to determine if this effect generalizes to IVSA of α-PVP in female rats.

Methods: Female Wistar rats were trained in IVSA of α-PVP (0.05 mg/kg/inf) in experimental chambers that contained an activity wheel. Groups (N=12 per group) of animals were trained with the wheels fixed (No-Wheel group) or free to move (Wheel group). Following acquisition, the groups were subjected to a wheel-access switch and then all animals to dose-substitution (0.0125-0.3 mg/kg/inf) with the wheels alternately fixed and free to move.

Results: Half of the rats initiated their IVSA pattern with a binge day of exceptionally high levels of drug intake, independent of wheel access condition. Wheel activity declined as acquisition progressed in the Wheel group and was much lower in the No-Wheel group in the wheel switch post-acquisition. Dose-effect curves were similar for binge/no binge subgroups and were not altered with wheel access during the dose-substitution.

Conclusions: This confirms the high reinforcer efficacy of α-PVP in female rats and the accompanying devaluation of wheel activity as a naturalistic reward.

Introduction

The structurally related (see **Figure 1**) cathinone derivative psychostimulant drugs 3,4-methylenedioxypyrovalerone (MDPV) and α-pyrrolidinopentiophenone (α-PVP) are monoamine transporter inhibitors with high selectivity for the dopamine transporter over the serotonin or norepinephrine transporters (Baumann et al. 2013; Marusich et al. 2014). These compounds are highly effective and potent reinforcers in male rats when assessed with the intravenous self-administration (IVSA) model (Aarde et al. 2013; Schindler et al. 2016; Watterson et al. 2014), and MDPV and α-PVP have been shown to be approximately equipotent when directly compared under a Progressive Ratio schedule of reinforcement in IVSA (Aarde et al. 2015a; Gannon et al. 2017a). A binge-like acquisition pattern of MDPV IVSA has been previously reported for male rats (Aarde et al. 2015b) in a study which also contrasted the effect of concurrent access to an activity

Figure 1: Chemical structures of MDPV and α-PVP.

wheel. In this pattern, a session was observed early in acquisition for about half of the individual rats wherein drug intake was particularly high on one day, but this was followed by a subsequent pattern of lower, but sustained intake for subsequent sessions. This binge-like pattern was independent of concurrent wheel-access since it occurred in equal proportions of the wheel-access groups. The pattern has not been reported for the acquisition of IVSA of other psychostimulants; however it is unclear if such individual patterns occurred but were overlooked in typical group-mean analysis or if such patterns do not typically occur with other drugs. The present study was therefore designed to determine if the binge-like acquisition would occur during acquisition of α-PVP IVSA, i.e., if the phenomenon would *generalize* to another drug.

The prior study of MDPV IVSA included an examination of the effect of concurrent access to an activity wheel because of prior findings that availability of an activity wheel in the home cage reduced IVSA of cocaine, methamphetamine, MDMA or mephedrone on the following session (Aarde et al. 2015c; Smith and Witte 2012), that wheel access provided concurrently with the IVSA session reduced

intake of methamphetamine in male rats previously trained to press the lever for food reward (Miller et al. 2012) and reduced cocaine IVSA in female rats with previous wheel experience (Cosgrove et al. 2002). In our MDPV study concurrent wheel access did not affect the acquisition of MDPV IVSA in behaviorally naïve male rats (Aarde et al. 2015b), however the ongoing IVSA of MDPV was accompanied by a gradual group mean decrease in wheel activity; this was similar to effects observed in male Wistar and Sprague-Dawley rats allowed to IVSA methamphetamine (Miller et al. 2012). Furthermore, the group mean trend for the MDPV study obscured the essentially immediate transition within-animal whereby IVSA of MDPV supplanted wheel activity within a single session for many of the subjects. One secondary goal of this study was therefore to determine if a similar co-option of wheel reinforcement would generalize to rats trained to IVSA α-PVP.

Female rats were used for this study as a further test of the generality of the binge-acquisition phenomenon and to address a general dearth of information on the self-administration of α-PVP, MDPV or indeed most cathinones in female animals (Gannon et al. 2017b; Hadlock et al. 2011; Schindler et al. 2016; Vendruscolo et al. 2017; Watterson et al. 2014). Female rats will self-administer more cocaine (Roth and Carroll 2004b; Smith et al. 2011) and more methamphetamine (Reichel et al. 2012; Roth and Carroll 2004a) than male rats (see (Anker and Carroll 2011; Lynch 2017) for reviews), however the IVSA of entactogen stimulants such as mephedrone, methylone and 3,4-methylenedioxymethamphetamine (MDMA) does not differ substantially between male and female rats (Creehan et al. 2015; Vandewater et al. 2015). Other studies have failed to find any sex-difference in IVSA of cocaine in rats (Miller et al. 2017; Perry et al. 2013) or the oral self-administration of cocaine in mice (DePoy et al. 2016) and in some cases male rats may acquire cocaine IVSA more rapidly and in higher proportion compared with female rats (Swalve et al. 2016a). These findings caution against simple assumptions about sex differences in drug IVSA and recommend investigation of the abuse liability of novel drugs in female as well as male rats to establish similarities and differences that might exist.

Finally, this study was designed to examine any association of the binge-like acquisition phenotype with reinforcer potency or efficacy after acquisition. Some of the popular media reports of

disturbing behavior (Alvarez et al. 2012; Anderson 2015; D'Oench 2015; Milian 2015) and medical emergency/fatality (Antonowicz et al. 2011; Benzie et al. 2011; Borek and Holstege 2012; Penders and Gestring 2011; Spiller et al. 2011; Umebachi et al. 2016; Wurita et al. 2014) subsequent to MDPV or α-PVP use appear to be associated with binge-like use. Our prior study found that stable post-acquisition MDPV IVSA was higher in the binge-acquisition animals compared with the non-bingers, suggesting either a pre-existing liability or a change caused by the binge-like acquisition. Therefore a dose-substitution experiment was included to determine if α-PVP was more potent or efficacious in the binge-acquisition individuals.

Materials and Methods

Subjects

Female (N=32) Wistar rats (Charles River, New York) entered the laboratory at 10 weeks of age and were housed in humidity and temperature-controlled (23±1 °C) vivaria on 12:12 hour light:dark cycles. Animals had ad libitum access to food and water in their home cages and remained gonadally intact for this study. All experimental procedures took place in scotophase and were conducted under protocols approved by the Institutional Care and Use Committees of The Scripps Research Institute and in a manner consistent with the Guide for the Care and Use of Laboratory Animals (National Research Council (U.S.). Committee for the Update of the Guide for the Care and Use of Laboratory Animals. et al. 2011).

Drugs

The α-pyrrolidinopentiophenone HCl (α-PVP) was obtained from Cayman Chemical and dissolved in physiological saline for the i.v. route of administration. Dosing is expressed as the salt.

Intravenous catheter implantation

Rats were anesthetized with an isoflurane/oxygen vapor mixture (isoflurane 5 % induction, 1-3 % maintenance) and prepared with chronic intravenous catheters as described previously (Aarde et al. 2015a; Aarde et al. 2013; Miller et al. 2013; Nguyen et al. 2017). Briefly, the catheters consisted of a 14-cm length polyurethane based tubing (MicroRenathane®, Braintree Scientific, Inc, Braintree MA, USA) fitted to a guide cannula (Plastics one, Roanoke, VA) curved at an angle and encased in dental cement anchored to an ~3-cm circle of durable mesh. Catheter tubing was passed subcutaneously from the animal's back to the right jugular vein. Catheter tubing was inserted into the vein and secured gently with suture thread. A liquid tissue adhesive was used to close the incisions (3M™ Vetbond™ Tissue Adhesive; 1469S B). A minimum of 4 days was allowed for surgical recovery prior to starting an experiment. For the first 3 days of the recovery period, an antibiotic (cephazolin) and an analgesic (flunixin) were administered daily. During testing and training, intra-venous catheters were flushed with ~0 .2–0.3 ml heparinized (32.3 USP/ml) saline before sessions and ~0.2–0.3 ml heparinized saline containing cephazolin (100 mg/ml) after sessions. Catheter patency was assessed once a week, beginning in the third week of training, via administration through the catheter of ~0.2 ml (10 mg/ml) of the ultra-short-acting barbiturate anesthetic, Brevital sodium (1 % methohexital sodium; Eli Lilly, Indianapolis, IN). Animals with patent catheters exhibit prominent signs of anesthesia (pronounced loss of muscle tone) within 3 s after infusion. Animals that failed to display these signs were considered to have faulty catheters and any data that were collected after the previous passing of the test were excluded from analysis.

Experimental apparatus

Operant conditioning chambers (Med Associates Model ENV-045; Med-PC IV software) enclosed in sound-attenuating cubicles were used for concurrent wheel access and self-administration, as previously described (Aarde et al. 2015b). Chambers were equipped with slanted activity wheels (~100 cm inner circumference, 30 degree tilt of running surface relative to the disk/back), which allowed wheel

access while tethered for self-administration. Eight total chambers were available thus animals were run in three sequential daily (M-F) cohorts with N=4 Wheel and N=4 No-Wheel per run.

Experiments

Acquisition and Wheel Switch: Female Wistar rats were prepared with intravenous catheters and trained to self-administer α-PVP (0.05 mg/kg per infusion) using a fixed-ratio 1 (FR1) response contingency in one hour daily (M-F) sessions. A pump pulse calculated to clear non-drug saline through the catheter started the session to ensure the first reinforcer delivery was not diluted, and a single priming infusion was delivered non-contingently if no response was made in the first 30 minutes of the session. The Wheel group (N=12) had the wheels unlocked and the No-Wheel group (N=12) had the wheels locked for the first 20 sessions. The wheel-access conditions were then switched for Sessions 21-27.

Dose Substitution: Thereafter, animals were subjected to dose substitution (0.0125, 0.025, 0.05, 0.1, 0.3 mg/kg/infusion) sessions in a randomized order (one session per dose). The response contingency was FR1,the sessions length 1 h. Half of the rats first completed a dose-substitution series with the wheel locked and half of the rats with the wheel unlocked (~half of each original wheel Group in each wheel condition). The dose-substitution was then repeated with the wheel-access conditions switched such that all rats completed one series with the wheel locked and one series with it unlocked.

Estrous Phase: Rats (N=14) were returned to standard training conditions (1 h sessions; 0.05 mg/kg/infusion dose) with the wheels locked for the assessment of any effects of estrous phase. A group of 5 completed the estrous studies with their original catheters intact and a group of 9 had been re-catheterized prior to this study. These studies followed dose-substitution studies with pentedrone and 4-methylpentedrone under FR and PR response contingencies not reported here. Estrous status was assessed via vaginal lavage and cytology using published criteria (Hubscher et al. 2005; Marcondes et al. 2002) after each session (M-F). Self-administration data were assessed over 11-16 consecutive sessions resulting in 1-5 days at each stage for N=14 subjects based on cytology alone. Secondary

analysis was conducted on a subset of the data in which a consistent four day progression of the estrous cycle could be identified (N=7; 1-2 sessions per estrous phase).

Data Analysis

The number of infusions obtained in the acquisition and wheel switch intervals was analyzed by Analysis of Variance (ANOVA) with Sessions as a within-subjects (i.e., repeated measures) factor and wheel access Group as a between-subjects factor. The dose-substitution data were analyzed with Dose and Wheel Access as within-subjects factors and the estrous data were analyzed with estrous phase as a within-subjects factor. Significant main effects from the ANOVAs were further analyzed with post hoc multiple comparisons analysis using the Tukey procedure for multi-level factor, and the Sidak procedure for two-level factor, comparisons. Survival analysis (Mantel-Cox log-rank) was used to compare the rate at which the Groups met an acquisition criterion of 3 sequential sessions with at least 6 infusions obtained; see (Aarde et al. 2015b). Review of the relevant literature (Anker and Carroll 2011; Lynch 2017) supports the hypothesis that drug intake during estrus would be higher than during proestrus thus pre-planned comparisons of estrus versus the other phases was used as the strongest test this hypothesis. The criterion for significant results was at $P < 0.05$ and all analyses were conducted using Prism 7 for Windows (v. 7.03; GraphPad Software, Inc, San Diego CA).

Results

Acquisition

A total of N=18 (N=7 No-Wheel, N=11 Wheel) rats retained patent catheters throughout the acquisition and wheel switch parts of the study. Examination of individual patterns of drug self-administration during the acquisition and wheel switch intervals under fixed per-infusion dose conditions confirmed the binge acquisition pattern in 7 / 11 Wheel animals and 4 / 7 No-Wheel animals (**Figure 2**). The wheels in two of the experimental chambers were found to be partially malfunctioning to the extent they did not move normally with an animal's weight and were adjusted after 6 sessions. This affected half

Figure 2: Individual infusions and wheel activity across the acquisition and wheel switch intervals for A) Wheel (N=11) and B) No-Wheel (N=7), acquisition groups. Grey symbols highlight the sessions in which 8 or more infusions were obtained in a five-minute interval and the arrow points to the binge-like acquisition sessions.

of the "Wheel" group (#2, 6, 10, 14 and 18 of those that remained patent) and created a new Wheel

subgroup which effectively was a No-Wheel group for the initial 6 sessions. All of the wheels were

moveable by rats after this point, however it appeared that one of them (affecting #2, 10 and 18) was not

fully free and a lubrication on the final session led to further increased activity for all of the affected animals.

Overall, the two groups of rats self-administered similar numbers of infusions of α-PVP (0.05 mg/kg/inf) during both acquisition and the wheel switch (**Figure 3A**). The ANOVA confirmed a significant effect of Sessions of acquisition [F (26, 416) = 4.51; P<0.0001], but not of Wheel access condition or of the interaction of factors. Similarly, the analysis of lever discrimination (**Figure 3B**) confirmed only a significant effect of Session [F (26, 416) = 6.72; P<0.0001]. The Mantel-Cox log-rank test failed to confirm (χ^2 = 2.34, df = 1, P = 0.127) any difference between the Groups in terms of the rate or proportion of subjects that reached acquisition criteria during the acquisition interval (**Figure 3C**). The rats in the Wheel group with functional wheel access from the start of acquisition were initially delayed in starting IVSA but ended up with a similar sustained mean intake compared with those which only had functional wheels after session 5 (**Figure 4A,B**). There was no systematic difference in wheel activity across acquisition sessions confirmed for either the entire Wheel cohort or for the subset that had fully functioning wheels from the beginning.

Figure 3: Mean (±SEM) infusions (A) and percent drug-associated lever responses (B) for Wheel (N=11) and No-Wheel (N=7) acquisition groups. C) Percent of each group meeting acquisition criteria. A significant difference from the first session, across groups, is indicated by *.

The Wheel switch produced no significant change in infusions obtained by the Wheel group upon fixing the wheels in place or by the No-Wheel group upon unlocking their wheels. Three of four Wheel individuals with essentially no IVSA intake during acquisition did start to self-administer drug (see **Figure 1**) but this produced no significant mean increase. Since wheel activity in the No-Wheel group was low after they gained access (**Figure 4C**), an additional

Figure 4: Mean (±SEM) infusions and wheel activity for A) Good Wheel (N=6); B) Bad Wheel (N=5); and C) No-Wheel (N=7) Groups.

experiment was conducted wherein a different group (N=8) of female Wistar rats were placed in the set of 8 experimental chambers with unlocked wheels for four one-hour sessions. The rats were in different boxes each session, thus data for each wheel were derived from 4 different rats. This study confirmed that the wheels in the chambers of the No-Wheel group were not dysfunctional, i.e., the test group produced mean activity levels in the No-Wheel chambers that was similar to the range for the Wheel group chambers. In addition, the No-Wheel chamber producing the lowest mean activity in this validation test was the one used for the No-Wheel rat that exhibited the most activity of that group in the wheel switch sessions.

Dose-Substitution

Self-Administration: A total of N=14 (N=7 No-Wheel, N=7 Wheel) rats retained patent catheters throughout the α-PVP dose-substitution studies. Wheel access had no effect on the dose-effect function for α-PVP IVSA (**Figure 5A**). The ANOVA confirmed a significant effect of Dose [F (4, 52) = 161.1; P<0.0001] but not of Wheel Access condition or the interaction of factors. The post-hoc test confirmed that each dose produced significantly different numbers of infusions compared with each of the other doses, collapsed across wheel-access condition. There was similarly no effect of Binge (N=10) / No Binge (N=4) phenotype (as expressed during acquisition) on the dose-effect relationship (**Figure 5B**). The ANOVAs confirmed a significant effect of Dose in the Wheel [F (4, 48) = 66.77; P<0.0001] and No-Wheel [F (4, 48) = 72.89; P<0.0001] dose-effect series but no effect of Binge phenotype or the interaction. The post-hoc tests

confirmed that, collapsed across Binge groups, all dose conditions differed from each other within Wheel conditions save that in the No-Wheel condition intake at the 0.025 mg/kg/infusion dose did not differ from 0.05 mg/kg/infusion conditionand intake in the 0.1 mg/kg/infusion condition did not differ from 0.3 mg/kg/infusion condition.

Wheel Activity: Wheel activity during the dose-substitution with the unlocked wheel followed an approximately monotonic trend inversely related to self-administered dose in 13 of the animals (**Figure 6A**). One subject used the

Figure 5: Mean (±SEM) infusions obtained during a dose-substitution. A) All rats with patent catheters (N=14) completed the dose series with the wheel unlocked (Wheel) or locked (No-Wheel). All doses differ significantly from all other doses. B) The results are divided by Binge (N=10) and No Binge (N=4) acquisition phenotype subgroups for the locked and unlocked wheel dose series.

wheel substantially more than the other animals, despite cumulative drug intake that was similar to the group mean and was thus treated as a significant outlier (**Figure 6B**). Analysis for the activity of the remaining 13 animals confirmed a significant effect of dose [F (2.39, 28.72) = 8.43; P<0.001] with the post-hoc test confirming significantly more activity after the 0.0125 mg/kg/infusion dose compared with 0.05-0.3 mg/kg/infusion doses.

Estrous Phase

A total of N=5 rats retained their original catheter patency through the estrous test. An additional N=9 were recatheterized prior to this phase. There were no differences in the self-administration of α-PVP (0.05 mg/kg/infusion) across the estrous phases (**Figure 7**) and statistical analysis of the full dataset (N=14) confirmed no effects of estrous phase on infusions obtained [F (1.782, 23.16) = 1.12;

P=0.34].Analysis of the subset (N=7) with cycle verified by both post-session cytology and observation of a standard four-day cycle likewise found no effect of estrous stage [F (1.634, 9.802) = 0.41; P=0.63].

Figure 6: A) Mean (N=13; ±SEM) drug intake and wheel activity during the dose-substitution procedure. A significant difference from the 0.0125 mg/kg/inf condition is indicated by *. B) Cumulative drug intake and wheel activity for one individual rat that was a significant outlier in wheel activity.

Discussion

This study confirms that α-PVP is a potent reinforcer in female rats, just as has been previously shown in male rats (Aarde et al. 2015a; Gannon et al. 2017a). Mean cumulative drug intake after 13 sessions was also similar to that found in a single prior α-PVP IVSA study in which female rats were trained for 13 acquisition sessions (Javadi-Paydar et al. 2017). The amount of α-PVP that was self-administered by the female rats by the end of acquisition in this study was also very similar to the amount obtained by males in a prior study of α-PVP IVSA using similar methods (Aarde et al. 2015a), thus supporting a conclusion of minimal sex- differences in the IVSA of α-PVP. As with a prior study of MDPV IVSA in male rats (Aarde et al. 2015b), the acquisition of IVSA of α-PVP in this study was not altered by concurrent access to an activity wheel on a group basis. Correspondingly, the introduction of wheel access after the initial acquisition period did not decrease IVSA, similar to a prior result for methamphetamine IVSA (Miller et al. 2012). In addition, about half of the rats exhibited a binge-like session during acquisition after which intake subsided to a stable level that was higher than that observed prior to the binge day. As such, this study shows that the binge-like pattern of acquisition that was previously reported for

Figure 7: Mean (±SEM) infusions obtained across estrous phases for A) the full sample of 14 rats, or B) the subset of 7 rats with standard 4 day cycles verified . P, proestrus; E, estrus; M, metestrus; D, diestrus.

MDPV IVSA in male rats (Aarde et al. 2015b) generalizes to female rats self-administering α-PVP. The binge session was observed in half of the Wheel and No-Wheel groups in the prior MDPV study (Aarde et al. 2015b) and this was matched by the relative proportions of binge-like acquisition female rats in the Wheel and No-Wheel groups in this study, further supporting the conclusion that this binging pattern is independent of wheel access during training.

The dose-substitution data indicated first that there were no lasting consequences of the binge-like acquisition phenotype. This shows that the binge is not likely a marker for a priori individual-preference differences that would have significant lasting effects on the potency or efficacy of α-PVP in IVSA. Similarly, this lack of any difference in the dose-substitution study suggests that the day or days of peak brain exposure to α-PVP during the binge did not induce any lasting neuronal plasticity related to rewarding value. Although data for α-PVP are not available, repeated non-contingent dosing with MDPV does not induce significant changes in brain monoamine function (Angoa-Perez et al. 2017; Anneken et al. 2015; Miner et al. 2017) as is the case for amphetamine derivative transporter substrates / monoamine releasers such as methamphetamine and 3,4-methylenedioxymethamphetamine (Green et al. 2003; McFadden et al. 2011; Ricaurte and McCann 1992; Sarkar and Schmued 2010; Taffe et al. 2002). Interestingly a recent study found that five repeated 96 hour MDPV self-administration sessions separated by 72 hours produced only subtle cognitive changes in rats (Sewalia et al. 2017).

Originally undetected issues with the movement of wheels in two of the chambers used for Wheel access animals precluded a robust test of the group mean reduction in wheel activity that was observed in prior studies of MDPV and methamphetamine IVSA in male rats (Aarde et al. 2015b; Miller et al. 2012). Nevertheless, wheel activity declined in concert with the onset of sustained IVSA behavior for the 6 rats with wheels that functioned normally from the start of acquisition. The fact that the female rats exhibited about twice as much wheel activity than the prior male group of Wistar rats did (Aarde et al. 2015b) emphasizes the competition of alternate reinforcers, rather than physical activity as measured by distance-traveled, in this model. This is a further expansion of the finding that male rats strains with a six-fold difference in wheel activity (Wistar > Sprague-Dawley) exhibited similar suppression of methamphetamine IVSA (Miller et al. 2012). In addition, the wheel activity of the No-Wheel group after

the wheel switch, and in the Wheel group with poorly adjusted wheels after adjustment, was lower than the initial activity in the Wheel subgroup with functional wheels from the beginning of training. This finding parallels an outcome for groups given wheel access after 7 or 14 days of methamphetamine IVSA (Miller et al. 2012), and further underlines the phenomenon by which drug IVSA co-opts and devalues more naturalistic sources of reinforcement.

This study failed to find any effect of wheel access on established IVSA both in the switched condition for the original No-Wheel group and in the dose-effect functions generated with and without wheel access during the dose-substitution study. This outcome contrasts with the foundational report of Cosgrove and colleagues in which cocaine IVSA and wheel activity were mutually inhibited (Cosgrove et al. 2002). The present finding confirms that our prior finding in which wheel access had no effect on established MA IVSA (Miller et al. 2012) was not a consequence of study in male rats. This is a critical confirmation given that the effect of wheel on cocaine IVSA in Cosgrove et al. (2002) was present in female rats to a much greater extent than in male rats. This study also shows the result is not likely attributable to pharmacological class, since α-PVP and cocaine are both restricted transporter inhibitors whereas methamphetamine is a transporter substrate and releaser.

There is one caveat to the overall conclusion that wheel access had minimal mean effect on the IVSA of α-PVP. Despite the fact that discontinuing wheel access had no significant mean effect on IVSA in the Wheel group (as in our prior study of methamphetamine IVSA), it was the case that three of four individuals who sustained high levels of wheel activity and minimal IVSA behavior during the 20 session acquisition interval started to respond for drug when the wheel was subsequently fixed in place. Thus it may be the case that a subset of rats would fail to ever acquire IVSA if wheel access was always available as an alternative.

This is one of the first available studies of the IVSA of the restricted transporter inhibitor cathinone derivatives in female rats and post-acquisition behavior was both highly stable under a fixed per-infusion dose and parametrically sensitive to dose-substitution. Thus the model offered the opportunity to determine potential effects of estrous phase since some prior reports indicate that cocaine and methamphetamine IVSA change significantly in female rats across the cycle, as reviewed previously

(Anker and Carroll 2011; Lynch 2017). No differences in IVSA were found across estrus phases in this study whether phase was determined solely by cytology or when the analysis was restricted to only a subset of animals for which a 4-day cycle could be clearly established. Additional study will be required to determine if there is something unique about α-PVP relative to either cocaine or methamphetamine with respect to differences in self-administration caused by hormonal fluctuation in female rats. Alternately, it is possible that our choice to run animals in the dark cycle, contrasted with prior models which run animals in the light (inactive) part of the day (Lynch et al. 2000; Lynch and Carroll 1999; Ramoa et al. 2013; Smethells et al. 2016; Swalve et al. 2016b), may explain the observed differences.

In conclusion, this study further confirms the likely high abuse liability of the second-generation cathinone α-PVP. The establishment of IVSA by female rats was similar to that of α-PVP or MDPV IVSA in male rats reported in prior studies, including a similar cumulative intake, a reduction in the value of wheel activity as an alternate reinforcer and a binge-like initial acquisition session in about half of the rats. Nevertheless, this study went on to show that the binge-like acquisition phenotype did not have lasting consequences for self-administration behavior, thus it may be the case that the pattern of acquisition is not predictive of individual differences in sustained drug seeking patterns of behavior.

Funding and Disclosure:

This work was funded by support from the United States Public Health Service National Institutes of Health (R01DA042211) which had no direct input on the design, conduct, analysis or publication of the findings. The authors declare no competing financial interests. This is manuscript #29574 from The Scripps Research Institute.

References

Aarde SM, Creehan KM, Vandewater SA, Dickerson TJ, Taffe MA (2015a) In vivo potency and efficacy of the novel cathinone alpha-pyrrolidinopentiophenone and 3,4-methylenedioxypyrovalerone: self-administration and locomotor stimulation in male rats. Psychopharmacology 232: 3045-55.

Aarde SM, Huang PK, Creehan KM, Dickerson TJ, Taffe MA (2013) The novel recreational drug 3,4-methylenedioxypyrovalerone (MDPV) is a potent psychomotor stimulant: self-administration and locomotor activity in rats. Neuropharmacology 71: 130-40.

Aarde SM, Huang PK, Dickerson TJ, Taffe MA (2015b) Binge-like acquisition of 3,4-methylenedioxypyrovalerone (MDPV) self-administration and wheel activity in rats. Psychopharmacology 232: 1867-77.

Aarde SM, Miller ML, Creehan KM, Vandewater SA, Taffe MA (2015c) One day access to a running wheel reduces self-administration of D-methamphetamine, MDMA and methylone. Drug and alcohol dependence 151: 151-8.

Alvarez SC, Benz GR, Elder R, Lew M, Sneider J (2012) Intervention Season 12, Episode 8: Skyler & Jessa. In: Mettler S (ed) Intervention. A & E, aetv.com

Anderson C (2015) Flakka, Synthetic Drug Behind Increasingly Bizarre Crimes Associated Press. The Associated Press, New York, NY
http://hosted.ap.org/dynamic/stories/U/US_FLAKKAS_EMERGENCE

Angoa-Perez M, Anneken JH, Kuhn DM (2017) Neurotoxicology of Synthetic Cathinone Analogs. Current topics in behavioral neurosciences 32: 209-230.

Anker JJ, Carroll ME (2011) Females are more vulnerable to drug abuse than males: evidence from preclinical studies and the role of ovarian hormones. Current topics in behavioral neurosciences 8: 73-96.

Anneken JH, Angoa-Perez M, Kuhn DM (2015) 3,4-Methylenedioxypyrovalerone prevents while methylone enhances methamphetamine-induced damage to dopamine nerve endings: beta-

ketoamphetamine modulation of neurotoxicity by the dopamine transporter. Journal of neurochemistry 133: 211-22.

Antonowicz JL, Metzger AK, Ramanujam SL (2011) Paranoid psychosis induced by consumption of methylenedioxypyrovalerone: two cases. Gen Hosp Psychiatry 33: 640 e5-6.

Baumann MH, Partilla JS, Lehner KR, Thorndike EB, Hoffman AF, Holy M, Rothman RB, Goldberg SR, Lupica CR, Sitte HH, Brandt SD, Tella SR, Cozzi NV, Schindler CW (2013) Powerful cocaine-like actions of 3,4-Methylenedioxypyrovalerone (MDPV), a principal constituent of psychoactive 'bath salts' products. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 38: 552-62.

Benzie F, Hekman K, Cameron L, Wade DR, Smolinske S (2011) Emergency department visits after use of a drug sold as "bath salts" --- michigan, november 13, 2010--march 31, 2011. MMWR Morb Mortal Wkly Rep 60: 624-7.

Borek HA, Holstege CP (2012) Hyperthermia and Multiorgan Failure After Abuse of "Bath Salts" Containing 3,4-Methylenedioxypyrovalerone. Annals of emergency medicine Jul;60(1):103-5.

Cosgrove KP, Hunter RG, Carroll ME (2002) Wheel-running attenuates intravenous cocaine self-administration in rats: sex differences. Pharmacology, biochemistry, and behavior 73: 663-71.

Creehan KM, Vandewater SA, Taffe MA (2015) Intravenous self-administration of mephedrone, methylone and MDMA in female rats. Neuropharmacology 92: 90-7.

D'Oench P (2015) Exclusive: Surveillance Tape Shows Mom Accused Of Abandoning Baby After Taking Flakka CBSMiami. CBSMiami, Miami, FL http://miami.cbslocal.com/2015/05/04/exclusive-surveillance-tape-shows-mom-accused-of-abandoning-baby-after-taking-flakka/

DePoy LM, Allen AG, Gourley SL (2016) Adolescent cocaine self-administration induces habit behavior in adulthood: sex differences and structural consequences. Transl Psychiatry 6: e875.

Gannon BM, Galindo KI, Mesmin MP, Sulima A, Rice KC, Collins GT (2017a) Relative reinforcing effects of second-generation synthetic cathinones: Acquisition of self-administration and fixed ratio dose-response curves in rats. Neuropharmacology Aug 12. pii: S0028-3908(17)30385-4. doi: 10.1016/j.neuropharm.2017.08.018.

Gannon BM, Galindo KI, Rice KC, Collins GT (2017b) Individual Differences in the Relative Reinforcing

Effects of 3,4-Methylenedioxypyrovalerone under Fixed and Progressive Ratio Schedules of

Reinforcement in Rats. The Journal of pharmacology and experimental therapeutics 361: 181-

189.

Green AR, Mechan AO, Elliott JM, O'Shea E, Colado MI (2003) The pharmacology and clinical

pharmacology of 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy"). Pharmacol Rev 55:

463-508.

Hadlock GC, Webb KM, McFadden LM, Chu PW, Ellis JD, Allen SC, Andrenyak DM, Vieira-Brock PL,

German CL, Conrad KM, Hoonakker AJ, Gibb JW, Wilkins DG, Hanson GR, Fleckenstein AE

(2011) 4-Methylmethcathinone (mephedrone): neuropharmacological effects of a designer

stimulant of abuse. The Journal of pharmacology and experimental therapeutics 339: 530-6.

Hubscher CH, Brooks DL, Johnson JR (2005) A quantitative method for assessing stages of the rat

estrous cycle. Biotechnic & histochemistry : official publication of the Biological Stain Commission

80: 79-87.

Javadi-Paydar M, Nguyen JD, Vandewater SA, Dickerson TJ, Taffe MA (2017) Locomotor and

reinforcing effects of pentedrone, pentylone and methylone in rats. Neuropharmacology Sep 4.

pii: S0028-3908(17)30419-7. doi: 10.1016/j.neuropharm.2017.09.002.

Lynch WJ (2017) Modeling the development of drug addiction in male and female animals.

Pharmacology, biochemistry, and behavior Jun 15. pii: S0091-3057(17)30169-7. doi:

10.1016/j.pbb.**2017**.06.006.

Lynch WJ, Arizzi MN, Carroll ME (2000) Effects of sex and the estrous cycle on regulation of

intravenously self-administered cocaine in rats. Psychopharmacology 152: 132-9.

Lynch WJ, Carroll ME (1999) Sex differences in the acquisition of intravenously self-administered

cocaine and heroin in rats. Psychopharmacology 144: 77-82.

Marcondes FK, Bianchi FJ, Tanno AP (2002) Determination of the estrous cycle phases of rats: some

helpful considerations. Brazilian journal of biology = Revista brasileira de biologia 62: 609-14.

Marusich JA, Antonazzo KR, Wiley JL, Blough BE, Partilla JS, Baumann MH (2014) Pharmacology of novel synthetic stimulants structurally related to the "bath salts" constituent 3,4-methylenedioxypyrovalerone (MDPV). Neuropharmacology 87: 206-13.

McFadden LM, Hoonakker AJ, Vieira-Brock PL, Stout KA, Sawada NM, Ellis JD, Allen SC, Walters ET, Nielsen SM, Gibb JW, Alburges ME, Wilkins DG, Hanson GR, Fleckenstein AE (2011) Methamphetamine treatment during development attenuates the dopaminergic deficits caused by subsequent high-dose methamphetamine administration. Synapse 65: 771-7.

Milian J (2015) Riviera Beach Police: Man smoking flakka wielded machete at stepdad PalmBeachPostcom. Palm Beach Post, Palm Beach, FL http://www.palmbeachpost.com/news/news/crime-law/riviera-beach-police-man-smoking-flakka-wielded-ma/nmW5k/

Miller ML, Moreno AY, Aarde SM, Creehan KM, Vandewater SA, Vaillancourt BD, Wright MJ, Jr., Janda KD, Taffe MA (2013) A methamphetamine vaccine attenuates methamphetamine-induced disruptions in thermoregulation and activity in rats. Biological psychiatry 73: 721-8.

Miller ML, Vaillancourt BD, Wright MJ, Jr., Aarde SM, Vandewater SA, Creehan KM, Taffe MA (2012) Reciprocal inhibitory effects of intravenous d-methamphetamine self-administration and wheel activity in rats. Drug and alcohol dependence 121: 90-6.

Miller MM, Meyer AE, Sprowles JL, Sable HJ (2017) Cocaine self-administration in male and female rats perinatally exposed to PCBs: Evaluating drug use in an animal model of environmental contaminant exposure. Experimental and clinical psychopharmacology 25: 114-124.

Miner NB, O'Callaghan JP, Phillips TJ, Janowsky A (2017) The combined effects of 3,4-methylenedioxymethamphetamine (MDMA) and selected substituted methcathinones on measures of neurotoxicity. Neurotoxicology and teratology 61: 74-81.

National Research Council (U.S.). Committee for the Update of the Guide for the Care and Use of Laboratory Animals., Institute for Laboratory Animal Research (U.S.), National Academies Press (U.S.) (2011) Guide for the care and use of laboratory animals: Eigth Edition. National Academies Press,, Washington, D.C., pp xxv, 220 p

Nguyen JD, Grant Y, Creehan KM, Vandewater SA, Taffe MA (2017) Escalation of intravenous self-administration of methylone and mephedrone under extended access conditions. Addiction biology 22: 1160-1168.

Penders TM, Gestring R (2011) Hallucinatory delirium following use of MDPV: "Bath Salts". Gen Hosp Psychiatry 33: 525-6.

Perry AN, Westenbroek C, Becker JB (2013) The development of a preference for cocaine over food identifies individual rats with addiction-like behaviors. PloS one 8: e79465.

Ramoa CP, Doyle SE, Naim DW, Lynch WJ (2013) Estradiol as a mechanism for sex differences in the development of an addicted phenotype following extended access cocaine self-administration. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 38: 1698-705.

Reichel CM, Chan CH, Ghee SM, See RE (2012) Sex differences in escalation of methamphetamine self-administration: cognitive and motivational consequences in rats. Psychopharmacology 223: 371-80.

Ricaurte GA, McCann UD (1992) Neurotoxic amphetamine analogues: effects in monkeys and implications for humans. Annals of the New York Academy of Sciences 648: 371-82.

Roth ME, Carroll ME (2004a) Sex differences in the acquisition of IV methamphetamine self-administration and subsequent maintenance under a progressive ratio schedule in rats. Psychopharmacology 172: 443-449.

Roth ME, Carroll ME (2004b) Sex differences in the escalation of intravenous cocaine intake following long- or short-access to cocaine self-administration. Pharmacology, biochemistry, and behavior 78: 199-207.

Sarkar S, Schmued L (2010) Neurotoxicity of ecstasy (MDMA): an overview. Curr Pharm Biotechnol 11: 460-9.

Schindler CW, Thorndike EB, Goldberg SR, Lehner KR, Cozzi NV, Brandt SD, Baumann MH (2016) Reinforcing and neurochemical effects of the "bath salts" constituents 3,4-

methylenedioxypyrovalerone (MDPV) and 3,4-methylenedioxy-N-methylcathinone (methylone) in

    male rats. Psychopharmacology 233: 1981-90.

Sewalia K, Watterson LR, Hryciw A, Belloc A, Ortiz JB, Foster Olive M (2017) Neurocognitive dysfunction

    following repeated binge-like self-administration of the synthetic cathinone 3,4-

    methylenedioxypyrovalerone (MDPV). Neuropharmacology Nov 26. pii: S0028-3908(17)30561-0.

    doi: 10.1016/j.neuropharm.2017.11.034. [Epub ahead of print]

Smethells JR, Swalve NL, Eberly LE, Carroll ME (2016) Sex differences in the reduction of impulsive

    choice (delay discounting) for cocaine in rats with atomoxetine and progesterone.

    Psychopharmacology 233: 2999-3008.

Smith MA, Walker KL, Cole KT, Lang KC (2011) The effects of aerobic exercise on cocaine self-

    administration in male and female rats. Psychopharmacology 218: 357-69.

Smith MA, Witte MA (2012) The effects of exercise on cocaine self-administration, food-maintained

    responding, and locomotor activity in female rats: Importance of the temporal relationship

    between physical activity and initial drug exposure. Experimental and clinical

    psychopharmacology 20: 437-446.

Spiller HA, Ryan ML, Weston RG, Jansen J (2011) Clinical experience with and analytical confirmation of

    "bath salts" and "legal highs" (synthetic cathinones) in the United States. Clinical toxicology

    (Philadelphia, Pa 49: 499-505.

Swalve N, Smethells JR, Carroll ME (2016a) Sex differences in the acquisition and maintenance of

    cocaine and nicotine self-administration in rats. Psychopharmacology 233: 1005-13.

Swalve N, Smethells JR, Zlebnik NE, Carroll ME (2016b) Sex differences in reinstatement of cocaine-

    seeking with combination treatments of progesterone and atomoxetine. Pharmacology,

    biochemistry, and behavior 145: 17-23.

Taffe MA, Davis SA, Yuan J, Schroeder R, Hatzidimitriou G, Parsons LH, Ricaurte GA, Gold LH (2002)

    Cognitive performance of MDMA-treated rhesus monkeys: sensitivity to serotonergic challenge.

    Neuropsychopharmacology : official publication of the American College of

    Neuropsychopharmacology 27: 993-1005.

Umebachi R, Aoki H, Sugita M, Taira T, Wakai S, Saito T, Inokuchi S (2016) Clinical characteristics of alpha-pyrrolidinovalerophenone (alpha-PVP) poisoning. Clinical toxicology (Philadelphia, Pa 54: 563-7.

Vandewater SA, Creehan KM, Taffe MA (2015) Intravenous self-administration of entactogen-class stimulants in male rats. Neuropharmacology 99: 538-545.

Vendruscolo JC, Tunstall BJ, Carmack SA, Schmeichel BE, Lowery-Gionta EG, Cole M, George O, Vandewater S, Taffe M, Koob GF, Vendruscolo LF (2017) Compulsive-Like Sufentanil Vapor Self-Administration in the Rat. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology. Aug 16. doi: 10.1038/npp.2017.172. [Epub ahead of print]

Watterson LR, Kufahl PR, Nemirovsky NE, Sewalia K, Grabenauer M, Thomas BF, Marusich JA, Wegner S, Olive MF (2014) Potent rewarding and reinforcing effects of the synthetic cathinone 3,4-methylenedioxypyrovalerone (MDPV). Addiction biology 19: 165-74.

Wurita A, Hasegawa K, Minakata K, Gonmori K, Nozawa H, Yamagishi I, Suzuki O, Watanabe K (2014) Postmortem distribution of alpha-pyrrolidinobutiophenone in body fluids and solid tissues of a human cadaver. Leg Med (Tokyo). 2014 Sep;16(5):241-6.