Novel Compensatory Mechanisms Enable the Mutant KCNT1 Channels to Induce

Seizures

Salleh N. Ehaideb[1,2#,*], Gentry T. Decker[1*], Petrina Smith[1], Daniel Davis[3], <u>Bing</u> <u>Zhang</u>[1@]

[1] Division of Biological Sciences, University of Missouri, Columbia, MO

[2] King Abdullah International Medical Research Center, Department of Medical

Genomic Research, Riyadh, Saudi Arabia

[3] Animal Modeling Core, University of Missouri, Columbia, MO

*These authors contributed equally to this work.

[@]To whom correspondence should be addressed (zhangbing@missouri.edu)

Words in Abstract: **150**

Words in main text: **3419**

Words in Introduction: **519**

Words in Figure legends: **821**

Abstract:

Mutations in the sodium-activated potassium channel (KCNT1) gene are linked to epilepsy. Surprisingly, all KCNT1 mutations examined to date increase K^+ current amplitude. These findings present a major neurophysiological paradox: how do gain-of-function KCNT1 mutations expected to silence neurons cause epilepsy? Here, we use *Drosophila* to show that expressing mutant KCNT1 in GABAergic neurons leads to seizures, consistent with the notion that silencing inhibitory neurons tips the balance towards hyperexcitation. Unexpectedly, mutant KCNT1 expressed in motoneurons also causes seizures. One striking observation is that mutant KCNT1 causes abnormally large and spontaneous EJPs (sEJPs). Our data suggest that these sEJPs result from local depolarization of synaptic terminals due to a reduction in Shaker channel levels and more active Na^+ channels. Hence, we provide the first *in vivo* evidence that both disinhibition of inhibitory neurons and compensatory plasticity in motoneurons can account for the paradoxical effects of gain-of-function mutant KCNT1 in epilepsy.

Introduction

Ion channels, such as potassium (K^+), sodium ($Na^{+)}$, and calcium (Ca^{2+}) channels, play important roles in neuronal excitability via regulation of the resting potential, the amplitude and the duration of action potentials (APs), the firing rates or patterns of APs, and synaptic transmission. Mutations in voltage-gated cation channels (Na^+, Ca^{2+}, and K^+) that result in neuronal hyper-excitability are often linked to epilepsy[1-3]. One of the distinct exceptions is the sodium-activated potassium channel (KCNT1, also called *Slack or Slo2.2*), which is encoded by the KCNT1 gene and unique in that they are activated by intracellular Na^+ and Cl^-[4-8]. At least 24 point mutations have been found in the human *KCNT1* gene in patients with autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), malignant migrating partial seizures of infancy (MMIPSI), early onset epileptic encephalopathy (EOEE), and West syndrome[9-15]. The number of patients affected by KCNT1 mutations is not known but many of these patients do not respond well to current anti-epilepsy drugs[13,16-18].

At present, the cellular mechanisms of KCNT1-related epilepsy remain poorly understood. Electrophysiological studies using heterologous expression systems such as the *Xenopus* oocytes or Chinese hamster ovary (CHO) cells have shown that all of the mutant KCNT1 channels studied to date significantly increase the magnitude of K^+ currents[9,11,12,19]. These findings are important but they present an interesting neurophysiological conundrum in that increased K^+ currents are expected to hyperpolarize or silence neurons and reduce the possibility for neurons to fire APs, a condition paradoxically unfavorable of triggering seizure or epilepsy.

Two different hypotheses have been proposed to explain this puzzle. The

'repolarization hypothesis'[5,20-23] states that increased Na^+-dependent K^+ currents might

accelerate the rate of AP repolarization, thus enhancing the firing rate. Studies of

KCNT1 KO mice showed, however, that the lack of the KCNT1 K^+ current sped up

repolarization and enhanced AP firing[24]. This result is not in agreement with the

'repolarization hypothesis', suggesting that KCNT1 K^+ current hinders repetitive firing

normally. Another hypothesis is the 'disinhibition hypothesis'[19], which states that

increased K^+ currents in inhibitory interneurons reduce GABA release and consequently

cause hyperexcitability of postsynaptic neurons. This hypothesis is plausible; however,

it has not been tested directly.

To test the disinhibition hypothesis and to discover new *in vivo* function of mutant

KCNT1, we generated a fruit fly model of human mutant KCNT1. Our data show that

expression of the mutant KCNT1 channels in GABAergic interneurons indeed causes

bang-sensitive seizures in flies, the first *in vivo* observation in support of the disinhibition

hypothesis. To our surprise, however, expression of the mutant channels in

motoneurons results in uncoordinated larval locomotion, enhanced synaptic

transmission, and also bang-sensitive seizures in adult flies. These results suggest that

neuronal excitability is enhanced rather than reduced in motoneurons by the mutant

KCNT1. We further show that neuronal hyperexcitability is achieved in part by a

compensatory downregulation of endogenous *Drosophila* K^+ channels such as Shaker

and enhancement of voltage-gated Na^+ channels, in an attempt to counter balance the

effect of increased KCNT1 K^+ current. Hence, our study supports two complementary

hypotheses, the 'disinhibition hypothesis' and a new 'compensatory plasticity

hypothesis', to help better understand the neurobiology underlying KCNT1-associated

epilepsy.

Results

Mutant hKCNT1 channels increase K$^+$ conductance in muscles

To date, nearly all electrophysiological studies of mutant KCNT1 channels

associated with epilepsy have been conducted in heterologous expression systems,

such as the Xenopus oocytes or CHO cells[9,11,12,19]. These studies show that mutant

hKCNT1 channels display a significant increase in K$^+$ current magnitude compared to

control[9,11,12,19]. To determine whether hKCNT1 mutations also exhibit gain-of-function

(an increase in K$^+$ conductance) phenotypes *in vivo*, we used the *GAL4/UAS* system to

overexpress wild type (WT) human KCNT1 (hKCNT1) and two mutants, G288S (a point

mutation in S5; next to the channel pore) and R928C (a point mutation near the NAD$^+$

domain) (Suppl Fig. 1a). Our data show that the resting potential is significantly

hyperpolarized and the input resistance dramatically reduced in muscles expressing

either G288S or R928C *hKCNT1* mutations by *Mhc-GAL4* (Suppl Fig. 1b-e). The

hKCNT1 mutant larvae were not able to crawl very far from its starting point and for the

most part of the recording period the larvae were stationary and inactive (Suppl Movie 1

and Suppl Fig. 2). These results are similar to the effects of human inward-rectifier

potassium channel 2.1 (*Kir2.1*)[25] expressed in muscles (Suppl Fig. 1b-e). Thus, our *in*

vivo data illustrate that *hKCNT1* G288S and R928C mutations are gain-of-function

resulting in an increase in K$^+$ conductance, and subsequently leading to impaired larvae

locomotion activities, hyperpolarized resting potential, and lower muscle resistance.

Mutant KCNT1 causes seizures in adult flies

Epilepsy is neurological disease defined as involuntary muscle convulsions

resulting in seizures. Drosophilists have been using adult flies to determine the genetic

bases of epilepsy and whether a particular mutation will predispose the flies to seizures.

Several behavior and electrophysiological techniques have been developed to assess

seizures in adult flies[26-28]. The bang-sensitivity behavioral assay is used to determine

whether subjecting flies to mechanical stimulation can elicit seizure activities[29,30].

To directly test the disinhibition hypothesis, we expressed the mutant and

wildtype human KCNT1 in GABAergic neurons using a cell-specific *Gal4, Gad-Gal4*[31].

Adult flies were then subjected to the bang-sensitive test in which the flies were placed

in an empty fly vial, vortexed for 20 sec, and then observed for seizing behavior (often

flip on their backs and unable to stand up or buzz around for at least one second). We

noted that the flies expressing mutant R928C in GABAergic neurons had a higher

propensity to display seizure (Fig. 1a-e; Suppl Movie 2). The frequency of seizures is

significantly higher and duration significantly longer compared to the two control groups

(*Gad-Gal4 >WT*; *Gad-Gal4 > WT human KCNT1*). The mean seizing frequency is 26%

compared to the controls (10%, 5%, respectively). The mean seizing duration of these

flies is significantly longer (p<0.01), averaging 28 Sec compared to 2 sec and 1.5 sec in

Gad-Gal4>WT and *Gad-Gal4>WT human KCNT1*, respectively. These results support

the 'disinhibition hypothesis', indicating that silencing inhibitory neurons tips the balance

towards excitation and thus implicating a role for GABAergic neurons in epilepitogenesis

in human patients.

We were also curious about whether expression of the mutant KCNT1 channels

in motoneurons would have a similar effect as they did in GABAergic neurons. In

motoneurons mutant KCNT1 driven by *D42-Gal4*[32,33] causes embryonic lethality at 25

°C, consistent with the silencing effect of enhanced K^+ current and pan neuronal

expression of D42 in early embryos. To overcome this obstacle, we used *GAL80*[ts][34]

(this form of *GAL80* become inactive when flies are reared at higher temperature) in the

context of the *D42, UAS hKCNT1* mutation to prevent the expression of hKCNT1

channels during development. Flies were kept at 18 °C until eclosion, then adult flies

were collected and kept at 25 °C to age, allowing the expression of hKCNT1 mutant

channels in adult flies. Mechanical stimulation of flies expressing R928C mutation by

vortexing showed a significant increase in the percentage of seizing flies and seizing

duration (25.7 ± 3.5 S and 11.3 ± 2.1 S, respectively) compared to *D42/+* (1.25 ± 0.9 S

and 1 S, respectively) and *D42/hKCNT1* (0.7 ± 0.7 S and 1 S, respectively, Fig. 1f-j and

Suppl Movie 3). These data suggest a new mode by which mutant hKCNT1 channels

cause seizure and epilepsy.

Figure 1: Neuronal expression of mutant hKCNT1 channel leads to seizures in adult flies.
(**a**, **f**) Representative image of flies after twenty seconds of mechanical stimulation (vortexing). Close-up images of the bottom of the vials are shown in (**b**, **c**, **g**, and **h**). Control flies show no sign of seizure activity after vortexing whereas a large number of mutant KCNT1-expressing flies exhibit seizure. (**d** and **e**) GABAergic expression of mutant hKCNT1 channel in adult flies show a significant increase in both number of seizing flies and seizing duration. (**i** and **j**) Motoneuron expression of mutant hKCNT1 channel predispose flies to seizure and very few control flies seized after vortex. If control flies seized they did not seize more than one second while flies expressing mutant hKCNT1 channels showed significantly longer seizure durations. At least 120 and 40 flies were tested for Motoneuron and GABAergic expression per genotype. Error bars show S.E.M. **$P < 0.01$ one-way ANOVA, Turkey HSD *post hoc test*.

Mutant hKCNT1 channels do not silence motoneurons but disrupt coordinated

crawling and motoneuronal firing

The findings from expressing mutant KCNT1 in muscle, GABAergic neurons, and

motoneurons appear to contradict one another in that muscles and GABAergic neurons

are silenced whereas motoneurons in adult flies are not. In order to better understand

the mechanisms by which mutant KCNT1 in motoneurons causes seizures, we decided

to examine the impact of mutant KCNT1 on larval behavior and physiology. To

overcome the embryonic lethality, we raised the flies with motoneuronal expression of

mutant KCNT1 at 18°C, which reduces the expression levels of the mutant hKCNT1

channels. *D42>G288S* reached pupal stage and few of the *D42>R928C* flies were able

to reach the adult stage. We noted that larvae expressing the mutant KCNT1 channels

showed severe crawling defects compared to control larvae (wildtype or larvae

expressing the wildtype KCNT1) (Suppl Fig. 3a,b; Suppl Movie 4). Because of the

uncoordinated crawling, the pupae became bended (crescent shape, Suppl Fig. 3c).

However, after careful observation we noted that unlike muscle expression of mutant

hKCNT1 channels, which were inactive for the most part of the observation, larvae

expressing G288S or R928C in motor neurons showed a strong and active but one

sided larval contractions (Suppl Movie 4). This finding suggests that the reduction in

larvae crawling distance is not due to neuronal inactivity but rather uncoordinated

neuronal firing.

Our behavioral observations of both larvae and adults strongly suggest that

neuronal expression of mutant KCNT1 at low levels does not silence motoneurons. On

the contrary, they cause hyperactivity in larvae (see below) and seizures in adult. To

determine the effects of *hKCNT1* mutations on neuronal activity, we utilized the well-

established neuromuscular junction (NMJ) preparation to assess their effects on

synaptic transmission in third-instar larvae[35,36]. Surprisingly, electrical stimulation of the

segmental nerve did not alter the amplitude of excitatory junction potentials (EJPs) of

larvae expressing G288S or R928C mutations in motor neurons compared to control

larvae. The mean EJP amplitude for *D42-GAL4>G288S*, *D42>R928C* was 49.1 ± 1.7

mV and 49.5 ± 1.7 mV, respectively, which were not significantly different from *D42/+*

(50.6 ± 1.1 mV) and *D42/hKCNT1* (49.5 ± 1.3 mV) (Fig. 2a,b). In addition, there was no

significant difference in membrane resting potentials between of G288S and R928C

mutations and controls (*D42/+* and *D42/hKCNT1*) (Suppl Fig. 4). These data indicate

that reduction of larval crawling activity caused by motoneuronal expression of mutant

hKCNT1 channels is not due to neuronal silencing but most likely due to inability to

coordinate neuronal firing, which hinders larvae crawling ability.

To more directly investigate whether *hKCNT1* mutations compromise

coordinated neuronal firing, we examined ventral nerve cord (VNC) neuronal firing in

third-instar larvae using Ca^{2+} imaging. Larvae VNC (equivalent to spinal cord in

humans) have eight pairs of abdominal ganglia, each right and left ganglion controls the

corresponding larval body wall segment[37-40]. For example, the first pair of ganglia

control the right and left abdominal muscle segment 1. During larvae crawling the first

pair of ganglia will fire together then the next pair and signal will propagate down until

the last pair fired[37-40]. To visualize ganglia firing we used the calcium sensor *GCaMP6S*

expressed in motoneurons. Larvae expressing WT hKCNT1 channels showed

coordinated ganglion firing, where right and left ganglia fired simultaneously (Fig 2c,d

and Suppl Movie 5). On the other hand, expression of hKCNT1 mutated channels

resulted in a significant number of unsynchronized motoneuronal firing, and sometimes

only ganglion in one side fired without the other pair firing at all (Fig. 2c,d,e and Suppl

Movie 5). In addition, oftentimes calcium waves were not able to pass through all

ganglia and usually terminated after reaching the fourth ganglion in larvae expressing

mutant hKCNT1 channels. However, both larvae expressing WT and mutated hKCNT1

channels showed a similar peak amplitude of Ca^{2+} waves (Fig. 2f). Taken together, our

data on both EJP and calcium waves suggest that hKCNT1 mutations do not silence

neuronal firing but rather influence firing synchrony.

Figure 2: Expression of *hKCNT1* mutation did not silence neurons, but caused uncoordinated ventral ganglion firing.
(**a**) Representative traces of excitatory junction potential (EJPs) under physiological conditions. (**b**) Average EJPs amplitude of ten recordings (two per larva) per genotype. Recordings were taken from muscle 6 segment A3 or A4. (**c**) Representative image of calcium waves in larvae expressing WT hKCNT1 channels (top) in motor neurons and larvae expressing mutated hKCNT1 channel (bottom). The calcium waves were visualized using *GCaMP6S* calcium sensor expressed in motoneuron. **L** and **R** indicate left and right sides, respectively, of ventral nerve cord (VNC) in third-instar larvae. White and yellow arrow heads indicate coordinated and uncoordinated left and right VNC ganglion firing respectively. (**d**) Representative calcium waves signals from left and right VNC ganglion pairs, showing coordinated (top) and uncoordinated (bottom) waves in larvae expressing WT and mutated hKCNT1 channels during active crawling respectively. (**e**) Quantification of percent of uncoordinated left and right VNC ganglion calcium waves. (**f**) Quantification of average calcium wave amplitude. Five larvae were used for calcium waves analysis. All larvae were reared at 22 °C for 5 days and 2 days at 25 °C prior to EJP recordings and calcium imaging. Error bars show S.E.M. **P < 0.01, one-way ANOVA, *Student's T test*.

Mutant hKCNT1 channels cause large spontaneous postsynaptic potentials at the larval NMJ

Spontaneous miniature excitatory junction potentials (mEJPs or minis) play a role in synaptic plasticity and function, evoked transmitter release, neuronal excitability, and postsynaptic membrane resistance[41-45]. It has been shown that in somatostatin cells taken from epileptic mice exhibit an increase in miniature excitatory postsynaptic synaptic current (mEPSC) frequency[46]. Thus, we investigated whether *hKCNT1* mutations alter the mini properties at the NMJ of third-instar larvae. We severed the segmental nerves posterior to the ventral ganglion and monitored mEJP activity (usually spontaneous release from single vesicles) at the NMJ.

Control larvae (both *D42/+* and *D42/hKCNT1*) displayed typical spontaneous minis, with an average frequency of 1.4 Hz and amplitude of 1.7 ± 0.1 mV (for both genotypes). Remarkably, larvae expressing mutant hKCNT1 channels showed very large spontaneous synaptic potentials, up to 14 mV (Fig. 3a). As shown below, these large synaptic events are spontaneous EJPs. Hence, we will refer them as spontaneous EJPs (sEJPs) rather than minis or mEJPs. The average amplitude of spontaneous synaptic potentials (note that this refers to both minis and sEJPs) in larvae expressing G288S and R928C mutations was significantly higher (2.6 ± 0.2 mV and 3 ± 0.2 mV, respectively, ANOVA, *P* < 0.01). The counts of minis plus sEJPs and their cumulative probability plots are shifted to the right in larvae expressing mutant hKCNT1 channels, due to the presence of significantly high number of large sEJPs in these larvae compared to *D42/+* and *D42/hKCNT1* larvae (Fig. 3b,c).

Figure 3: Third-instar larvae expressing *hKCNT1* mutations in motor neurons showed spontaneous EJPs.
(**a**) Representative traces of miniature excitatory junction potentials (mEJP or minis) under physiological condition. Both *hKCNT1* mutants show significantly unusually larger spontaneous synaptic events in addition to minis compare to *D42/+* and *D42/hKCNT1* traces. We call these large synaptic potentials spontaneous EJPs (sEJPs). (**b**) Histograms of spontaneous synaptic potentials (minis plus sEJPs) of the four different genotypes. Synaptic potential counts are sorted into 0.125 mV bins. Note the right-shift distribution of spontaneous synaptic potentials in the two mutant KCNT1 larvae. (**c**) Cumulative probability plot of spontaneous synaptic potentials from larvae expressing mutant hKCNT1 channels illustrate the present of significantly higher number of large sEJPs compare to *D42/+* and *D42/hKCNT1*. Recordings were taken from muscle 6 segment A3 or A4 and all larvae were reared at 22 °C for 5 days and 2 days at 25 °C prior to synaptic recordings. A total of ten recordings (two per larva) were made per genotype.

These sEJPs are highly unusual as the axons of the motoneurons are cut free

from the soma and it is not possible for action potentials to be propagated from the

soma. We reasoned that they were caused by local depolarization at the synaptic

terminal. This could be possible only if there was a compensatory enhancement in

voltage-gated cation (Na^+ or Ca^{2+}) channel levels and activities or a reduction in K^+

channel levels and activities. To test this hypothesis, we used tetrodotoxin (TTX) to

block voltage-gated sodium channel activity and surprisingly observed that TTX only

reduced the number of 'extremely' large sEJPs (>10 mV) but did not abolish sEJPs in

larvae expressing mutant hKCNT1 channels (Fig. 4a,b,c; Suppl Fig. 5). However, we

noticed a slight decrease in average amplitude of spontaneous synaptic potentials in the

G288S and R928C expressing larvae (2.1 ± 0.1 mV and 2.5 ± 0.2 mV, respectively,

ANOVA, P < 0.01), but TTX had no significant effect on mini amplitude in *D42/+* and

D42/hKCNT1 larvae (1.6 ± 0.1 mV and 1.5 ± 0.1 mV, respectively). These data suggest

that there is an upregulation of voltage-gated Na^+ channels but they are not responsible

for eliciting the sEJPs.

We then added the Ca^{2+} channel blocker cadmium to the TTX-containing saline

and effectively eliminated the sEJP in the mutant KCNT1-expressing larvae (Fig. 4d,e,f).

On the other hand, no changes in mEJP properties were observed in the *D42/+* and

D42/hKCNT1 larvae. These results indicate that depolarization of voltage-gated Ca^{2+}

channels at the NMJ likely results in more transmitter release and gives rise to sEJPs.

Figure 4: Blocking both voltage-gated sodium and calcium channels in larvae expressing mutant hKCNT1 channels eliminated the spontaneous EJPs.
(**a**) Representative traces of spontaneous synaptic potentials (minis + sEJPs) treated with voltage-gated sodium channel blocker TTX (1 µM) in HL-3 saline. Note that the large sEJPs persist in TTX-saline. (**b** and **c**) Histograms and cumulative probability plots show that the extremely large sEJPS (>10 mV) are blocked by TTX but TTX does not eliminate all sEJPs. (**d**) Representative traces of spontaneous synaptic potentials (minis + sEJPs) treated with both voltage-gated Na^+ channel blocker TTX (1 µM) and voltage-gated K^+ channel blocker $CdCl_2$ (10 µM) in HL-3 saline (This treatment successfully reduces or abolishes the occurrence of spontaneous EJPs. (**e** and **f**) Histograms and cumulative probability plots show similar distributions of spontaneous synaptic potentials in TTX and Cd^{2+}-treated larval NMJs. Recordings were taken from muscle 6 segment A3 or A4 and all larvae were reared at 22 °C for 5 days and 2 days at 25 °C prior to mini recordings. A total of ten recordings (two per larva) were made per genotype.

Immunocytochemistry reveals compensatory decreases in endogenous K^+

channels in larvae expressing mutant KCNT1

Our behavioral, imaging, and electrophysiological studies collectively suggest

that mutant KCNT1 channels trigger compensatory mechanisms by which other ion

channels are down or up regulated to counter balance the silencing effect of KCNT1 K^+

channels. In other words, the large increase in K^+ currents through mutant KCNT1

channels has the potential to fully or partially silence neurons, like Kir2.1 does, to

reduce the possibility of firing and impair synaptic transmission. In contrast, the

motoneurons appear to fire normally (albeit not synchronized) and the EJP amplitude is

largely unchanged. What enables the motoneurons to fire action potentials and the

synaptic terminal to produce local and spontaneous auto-depolarization? We

hypothesize that there is a compensatory change of excitability at both cell bodies and

axons to ensure that action potentials can be produced and propagated. Furthermore,

we hypothesize that there is a local compensation at the NMJ to enhance synaptic

activity. We used antibodies specific to Shaker (Sh) to stain the CNS and the

neuromuscular preparation of larvae and showed that Shaker channels are reduced in

levels (Fig. 5a,b).

Figure 5: Neuronal expression of mutant hKCNT1 channel leads to significant reduction in Shaker channels staining at the NMJ.
(**a**) Representative NMJ confocal images of WT and larvae expressing mutant hKCNT1 channel in motoneurons. Larvae expressing mutant hKCNT1 channel show weak or no Shaker staining. Scale = 5 μM. (**b**) Quantification of Shaker staining intensity show that Larvae expressing mutant hKCNT1 channel have a significant reduction in Shaker signal compared to control. **$P < 0.01$, *Student's T test*.

Discussion

The discovery of the link between mutant KCNT1 and epilepsy is important but it

also presents challenges understanding the role of the K^+ channel in epileptogenesis. *In*

vitro studies to date all show that mutant KCNT1 channels significantly increases K^+

current magnitude[9,11,12,19]. It is easier to understand why mutations enhancing Na^+ and

Ca^{2+} channel activities or reducing K^+ channel activities can cause hyperexcitability and

increase the probability for seizures or epilepsy. Similarly, mutant Cl^- channels in

skeletal muscles both in the fainting goat and humans cause myotonia[47-49].

Neurophysiologically speaking, enhancing K^+ currents is expected to hyperpolarize the

resting potential or truncate action potentials and thereby reducing the possibility for

neuronal firing. What then accounts for the neuropathology in KCNT1-associated

epilepsy?

Two different hypotheses have been proposed to explain why enhanced K^+

channel activities in mutant KCNT1 could cause hyperexcitability and generate

conditions in favor of seizures. One hypothesis, which we name 'repolarization

hypothesis', states that following Na^+ influx the activation of KCNT1 K^+ channels

shortens the duration of APs by repolarizing it at a faster rate. This in turn enables the

neuron to fire more APs per unit time, resulting in hyperexcitability. This possibility is

plausible, as shown in electrocytes of some electrical fish[20] and BK channel and β

subunit -linked epilepsy[50-52]. However, it will depend on the mode by which Na^+

activates KCNT1 channels and the kinetics of the $K_{(Na)}$ current[53]. KCNT1 channels are

unique in that is activated by intracellular Na^+ and Cl^-. Salkof and colleagues showed

that activation of KCNT1 channels does not need high intracellular $[Na^+]_i$[54], implying that

KCNT1 may also be important for contributing to resting potential as well as repolarizing

AP. Furthermore, $K_{(Na)}$ following Na^+ influx could outlast the duration of an AP,

especially during afterhyperpolarization period, and therefore prevent or delay the onset

of the next AP. Finally, genetic studies of KCNT1 KO in mice showed that AP

repolarization is faster and neurons fire more APs[24]. This *in vivo* study indicates that the

presence of KCNT1 K current normally hinders excitability or silence neurons. Our data

obtained from muscle expression of the mutant KCNT1 provide strong evidence that the

mutant KCNT1 channels are similar to inward rectifier K^+ channel 2.1 in reducing the

muscle input resistance and hyperpolarizing the muscle resting potential. Further,

neuronal expression of the mutant KCNT1 at high levels causes embryonic lethality or

folded wings (Suppl Fig. 6) if the flies live to adult. These observations are consistent

with the notion that mutant KCNT1 are gain of function mutations that silence both

muscles and neurons. Hence, studies in both mice and fruit flies do not support the

'repolarization hypothesis'.

The second hypothesis, which we call the 'disinhibition hypothesis'[19], postulates

that the enhanced mutant KCNT1 K^+ current silences inhibitory neurons and thereby

removing inhibition of neural circuits and tipping the balance towards hyperexcitability

and seizures. This is an exciting hypothesis, but up to this point this hypothesis has not

been directly tested. Our data showing that expression of mutant KCNT1 in GABAergic

neurons induces seizures in adult flies lends a strong support to this hypothesis.

However, the disinhibition hypothesis alone may not be sufficient to account for

the mutant KCNT1 actions in the nervous system. This is because KCNT1 is broadly

expressed in a variety of neuronal types and regions in the human brain[55,56]. The broad

and complex expression pattern of KCNT1 begs for additional mechanisms other than

the disinhibition hypothesis to account for the effect of mutant KCNT1 in other neurons.

By expressing the mutant KCNT1 in motoneurons we have learned that these neurons

can be silenced if the expression levels are high or become hyperexcited if the

expression levels are low. When specifically expressed in adult motoneurons, mutant

KCNT1 also causes bang-induced seizures, suggesting that these neurons are not

silenced. This is consistent with the observations of normal EJPs and Ca^{2+} peaks in

motoneurons in larvae. More strikingly, we reveal novel changes at the synapse where

local depolarization leads to spontaneous synaptic potentials, which is partially sensitive

to TTX blockade and fully sensitive to Cd^+. At the NMJ, the major endogenous K^+

channel Shaker is significant reduced in levels, and TTX can block some of the sEJPs,

providing a molecular explanation for the enhanced excitability in motoneurons. Our

finding of sEJPs is consistent with previous findings that sEJPs also occurred in flies

with mutations in Sh eag K^+ channels[57]. Based on these observations, we propose a

'compensatory plasticity hypothesis' as a novel mechanism to counter balance the

silencing effect of mutant KCNT1 currents as an additional means to produce neuronal

hyperexcitability and seizures. This work adds to the growing list of examples of

homeostatic regulation of neuronal excitability[58], although in our KCNT1 fly model it fails

to fully compensate to restore normal function. The disinhibition and compensatory

mechanisms are complementary to each other, suggesting that mutant KCNT1

channels likely have different effects on different neurons. Further studies are expected

to pinpoint the regulatory mechanisms by which mutant KCNT1 regulates Sh, Na^+ and

perhaps ion channel gene expression or trafficking.

  Supplementary Figure legends

Supplementary Figure 1: Muscle expression of mutant hKCNT1 protein in third-

instar larvae reduce crawling distance, membrane resting potential, and muscle

resistance.

(**a**) Schematic diagram of human KCNT1 protein showing protein structure, location of

the mutations studied in this paper, and type of epilepsy syndrome caused by these

mutations. (**b**) Third-instar larvae expressing either mutant *hKCNT1* or *Kir2.1* in muscles

show a significant reduction in average crawling distance compare to controls. At least

twenty larvae were used per genotype (10 males and 10 females). (**c** and **d**) In addition,

overexpression of mutant KCNT1 channels significantly hyperpolarized the average

resting membrane potential (**c**) and average muscle input resistance (**d**). (**e**)

Representative traces for muscle input resistance in control and experimental larvae.

Note the dramatic and similar reduction in muscle input resistance in both *MHC>R982C*

and *MHC>Kir2.1* larvae. Muscle 6 of abdominal segment 3 or 4 was used for assisting

the resting membrane potential and muscle input resistance. Ten larvae were used per

genotype. All larvae were kept at 25 °C until the time of testing. Error bars show S.E.M.

**P* < 0.01, one-way ANOVA, Turkey HSD *post hoc test*.

Supplementary Figure 2: Representative larvae crawling tracks. Larvae expressing

mutant hKCNT1 channels in muscles show significant reduction in crawling distance

compared to controls.

Supplementary Figure 3: Motor neuronal expression of mutant hKCNT1 channels

alter pupal and adult wings morphology and reduces third-instar larval crawling

activity.

(**a**) Representative larvae crawling tracks. (**b**). (**c**) Motor neuron expression of mutant

hKCNT1 channels significantly reduce average crawling distance compare to controls.

(**c**) Representative images of pupal morphology. Larvae expressing mutant hKCNT1

channels show bent pupa shape. At least twenty larvae were used per genotype (10

387 males and 10 females). All larvae were reared at 22 °C for 5 days then at 25 C° for 2

388 days. Error bars show S.E.M. **$P < 0.01$ one-way ANOVA, Turkey HSD *post hoc test*.

391 **Supplementary Figure 4: Motoneuronal expression of mutant hKCNT1 channels**

392 **did not affect membrane resting potential.** Both controls and larvae expressing

393 mutant hKCNT1 channels in motoneurons have similar membrane resting potentials.

Supplementary Figure 5: Blocking voltage-gated Na⁺ channels with TTX reduced the number of extremely large sEJPS but did not abolish sEJPs in larvae expressing mutant hKCNT1 channels. Cumulative probability plot of minis plus sEJPs shows that TTX reduced the number of the extremely large sEJPs (>10 mV), but did not eliminate the sEJPs.

Supplementary Figure 6: neuronal expression of the mutant KCNT1 Channels causes folded wing phenotype. Representative images of adult fly wing morphology. All larvae were reared at 18 °C until eclosion.

Supplementary Movie legends

Movie 1: Muscle expression of mutant hKCNT1 channels hinders larvae crawling

activity. Representative crawling activity assay videos of controls and mutant KCNT1

larvae. Larvae expressing mutant hKCNT1 channels in muscle show significant

reduction in crawling activity compared to controls. Twenty larvae (ten males and ten

females) were used per genotype, and all larvae were aged at 25 °C prior to testing.

Movie 2: GABAergic expression of mutant hKCNT1 channels increase the number

of seizing flies and seizing duration in adult flies. Representative bang-sensitivity

behavioral assay videos. Flies expressing mutant hKCNT1 channels in inhibitory

neurons exhibit seizure phenotypes, such as spinning in the bottom of the vial or flip on

their back and remain at that state for one second or more. At least 40 flies were used

per genotype, and all larvae were aged at 18 °C until they enclosed then adult flies are

aged at 25 °C prior to testing.

Movie 3: Motoneuronal expression of mutant hKCNT1 channels increase the

number of seizing flies and seizing duration in adult flies. Representative bang-

sensitivity behavioral assay videos. Flies expressing mutant hKCNT1 channels in

motoneurons exhibit seizure phenotypes, such as spinning in the bottom of the vial or

flip on their back and remain at that state for one second or more. 120 flies were used

per genotype, and all larvae were aged at 18 °C until they enclosed then adult flies are

aged at 25 °C prior to testing.

Movie 4: Larvae expressing mutant hKCNT1 channels in motoneurons shows

uncoordinated contraction of left and right abdominal muscle hemisegments.

Representative crawling activity assay videos of controls and mutant KCNT1 larvae.

Larvae expressing mutant hKCNT1 channels in motoneurons show significant reduction

in crawling activity compared to controls, due to unsynchronized muscle contractions.

Twenty larvae (ten males and ten females) were used per genotype, and all larvae were

aged at 22 °C for 5 days and at 25 °C for two days prior to testing.

Movie 5: Ca^{2+} waves in larvae expressing mutant hKCNT1 channels in

motoneurons shows uncoordinated VNC abdominal ganglia firing. Representative

Ca2+ waves assay videos of controls and mutant hKCNT1 larvae. Larvae expressing

mutant KCNT1 channels in motoneurons show significant increase in unsynchronized

abdominal ganglia firing. Ca^{2+} waves recording was at 2 frames per second and movie

is showing 4X the original speed. Four larvae were used per genotype, and all larvae

were aged at 22 °C for 5 days and at 25 °C for two days prior to testing.

Methods

Fly Strains.

Fly cultures kept at 12-h light/dark cycle on standard cornmeal food. The following

Drosophila strains were used: *D42-GAL4*, *GAD1-GAL4*, *GAL80^{ts}*, *Mhc-GAL4*, *8622*, and

UAS-GCaMP6S were obtained from Bloomington Stock Center. *UAS-Kir2.1* was gift

from G. Davis, *UAS-EKOX2*, gift from B. White, and UAS-KCNT1 transgenic lines, gift

from L. Dibbens.

Larvae crawling behavior assay and analysis.

Wandering third-instar larvae were washed with double distilled water and placed on

150mm petri dish containing 1% agarose. The larvae locomotion behavior was recorded

for 1 min at 30 frames/s by Fujinon DF6HA-1B camera and FlyCapture software. Videos

were analyzed using ImagJ and wrMTrck plugin as previously descried[59].

Calcium imaging.

Third-instar larvae were dissected and ventral nerve cord, which contains motoneurons,

was viewed using Olympus BX61 compound microscope with 10X air lens. Calcium

waves were recorded using ORCA-R^2 CCD camera (Hamamatsu) and CellSens

Dimension 1.9 software. ImageJ software was used for calcium waves analysis.

Electrophysiology

Recordings were made from muscle 6 in abdominal segments 3 and 4 from third-instar

larvae as previously discribed[35,36]. Recordings were made with a modified HL3 saline[60]

containing 70mM NaCl, 5mM KCl, 10mM $MgCl_2$, 10mM $NaHCO_3$, 5mM trehalose,

115mM sucrose, 5mM HEPES, and 1.2mM $CaCl_2$. For true mini was recorded with HL3

solution plus 1 µM TTX and 10 µM $CdCl_2$. Recording microelectrodes were fabricated

using P2000g puller (Sutter instruments) and filled with 3M KCl. Signals were recorded

by Axoclamp B2 amplifier and pCLAMP 6 software (Molecular devices). Severed

segmental nerve was sucked by stimulation microelectrode and stimulated by Master-8

pulse generator and iso-flex stimulus isolator (A.M.P.I.).

Fly bang-sensitive assay

Experiment was done as previously descried with some modification[29,30]. Briefly, flies

were collected immediately after eclosion, kept at 25 °C, and mechanically stimulated

with mini vortexer (VWR Scientific Products) at power setting 10. Flies behavior was

recorded by SONY (HDR-SR11) camcorder. Flies that flipped on their back and

remained at that position for at least one second were considered to be exhibiting

seizure activity.

Immunocytochemistry.

Wandering third-instar larvae were dissected in HL3 solution and fixed for 2h in

4%paraformaldehyde. Fixed larvae were washed 3X 15 min each with PBST (0.2

TritonX-100). Larvae were incubated overnight at 4 °C with primary antibody with the

appropriate dilution. The following antibodies were used: guinea pig anti-Shaker 1:500

(gift from Kyunghee Koh), rabbit HRP-488 1:500. Alexa-conjugated secondary

antibodies (1:500) were used. Images were acquired with Leica SP2 confocal

microscope using a 63X oil immersion objective.

Statistical analysis.

Data were analyzed as described in the figure legends using Origin software version

6.0.

References

1   Cain, S. M. & Snutch, T. P. in *Jasper's Basic Mechanisms of the Epilepsies*   (eds J. L.

502   Noebels *et al.*) (2012).

503   2   Catterall, W. A. in *Jasper's Basic Mechanisms of the Epilepsies*   (eds J. L. Noebels *et al.*)

504   (2012).

505   3   Cooper, E. C. in *Jasper's Basic Mechanisms of the Epilepsies*   (eds J. L. Noebels *et al.*)

506   (2012).

507   4   Yuan, A. *et al.* The sodium-activated potassium channel is encoded by a member of the

508   Slo gene family. *Neuron* **37**, 765-773 (2003).

509   5   Bhattacharjee, A. & Kaczmarek, L. K. For K+ channels, Na+ is the new Ca2+. *Trends*

510   *Neurosci* **28**, 422-428, doi:10.1016/j.tins.2005.06.003 (2005).

511   6   Kim, G. E. & Kaczmarek, L. K. Emerging role of the KCNT1 Slack channel in intellectual

512   disability. *Front Cell Neurosci* **8**, 209, doi:10.3389/fncel.2014.00209 (2014).

513   7   Villa, C. & Combi, R. Potassium Channels and Human Epileptic Phenotypes: An Updated

514   Overview. *Front Cell Neurosci* **10**, 81, doi:10.3389/fncel.2016.00081 (2016).

515   8   Joiner, W. J. *et al.* Formation of intermediate-conductance calcium-activated potassium

516   channels by interaction of Slack and Slo subunits. *Nat Neurosci* **1**, 462-469,

517   doi:10.1038/2176 (1998).

518   9   Barcia, G. *et al.* De novo gain-of-function KCNT1 channel mutations cause malignant

519   migrating partial seizures of infancy. *Nat Genet* **44**, 1255-1259, doi:10.1038/ng.2441

520   (2012).

521   10   Heron, S. E. *et al.* Missense mutations in the sodium-gated potassium channel gene

522   KCNT1 cause severe autosomal dominant nocturnal frontal lobe epilepsy. *Nat Genet* **44**,

523   1188-1190, doi:10.1038/ng.2440 (2012).

524   11   Kim, G. E. *et al.* Human slack potassium channel mutations increase positive

525   cooperativity between individual channels. *Cell Rep* **9**, 1661-1672,

526   doi:10.1016/j.celrep.2014.11.015 (2014).

527   12   Martin, H. C. *et al.* Clinical whole-genome sequencing in severe early-onset epilepsy

528   reveals new genes and improves molecular diagnosis. *Hum Mol Genet* **23**, 3200-3211,

529   doi:10.1093/hmg/ddu030 (2014).

530   13   Moller, R. S. *et al.* Mutations in KCNT1 cause a spectrum of focal epilepsies. *Epilepsia* **56**,

531   e114-120, doi:10.1111/epi.13071 (2015).

532   14   Vanderver, A. *et al.* Identification of a novel de novo p.Phe932Ile KCNT1 mutation in a

533   patient with leukoencephalopathy and severe epilepsy. *Pediatr Neurol* **50**, 112-114,

534   doi:10.1016/j.pediatrneurol.2013.06.024 (2014).

535   15   Lim, C. X., Ricos, M. G., Dibbens, L. M. & Heron, S. E. KCNT1 mutations in seizure

536   disorders: the phenotypic spectrum and functional effects. *J Med Genet* **53**, 217-225,

537   doi:10.1136/jmedgenet-2015-103508 (2016).

538   16   Bearden, D. *et al.* Targeted treatment of migrating partial seizures of infancy with

539   quinidine. *Ann Neurol* **76**, 457-461, doi:10.1002/ana.24229 (2014).

540   17   Fukuoka, M. *et al.* Quinidine therapy for West syndrome with KCNTI mutation: A case

541   report. *Brain Dev* **39**, 80-83, doi:10.1016/j.braindev.2016.08.002 (2017).

18   Mikati, M. A. *et al.* Quinidine in the treatment of KCNT1-positive epilepsies. *Ann Neurol*

78, 995-999, doi:10.1002/ana.24520 (2015).

19   Tang, Q. Y. *et al.* Epilepsy-Related Slack Channel Mutants Lead to Channel Over-Activity

by Two Different Mechanisms. *Cell Rep* **14**, 129-139, doi:10.1016/j.celrep.2015.12.019

(2016).

20   Markham, M. R., Kaczmarek, L. K. & Zakon, H. H. A sodium-activated potassium channel

supports high-frequency firing and reduces energetic costs during rapid modulations of

action potential amplitude. *J Neurophysiol* **109**, 1713-1723, doi:10.1152/jn.00875.2012

(2013).

21   Franceschetti, S. *et al.* Na+-activated K+ current contributes to postexcitatory

hyperpolarization in neocortical intrinsically bursting neurons. *J Neurophysiol* **89**, 2101-

2111, doi:10.1152/jn.00695.2002 (2003).

22   Yang, B., Desai, R. & Kaczmarek, L. K. Slack and Slick K(Na) channels regulate the

accuracy of timing of auditory neurons. *J Neurosci* **27**, 2617-2627,

doi:10.1523/JNEUROSCI.5308-06.2007 (2007).

23   Lu, R. *et al.* Slack channels expressed in sensory neurons control neuropathic pain in

mice. *J Neurosci* **35**, 1125-1135, doi:10.1523/JNEUROSCI.2423-14.2015 (2015).

24   Martinez-Espinosa, P. L. *et al.* Knockout of Slo2.2 enhances itch, abolishes KNa current,

and increases action potential firing frequency in DRG neurons. *Elife* **4**,

doi:10.7554/eLife.10013 (2015).

25   Paradis, S., Sweeney, S. T. & Davis, G. W. Homeostatic control of presynaptic release is

triggered by postsynaptic membrane depolarization. *Neuron* **30**, 737-749 (2001).

26   Pandey, U. B. & Nichols, C. D. Human disease models in Drosophila melanogaster and

the role of the fly in therapeutic drug discovery. *Pharmacol Rev* **63**, 411-436,

doi:10.1124/pr.110.003293 (2011).

27   Song, J. & Tanouye, M. A. From bench to drug: human seizure modeling using

Drosophila. *Prog Neurobiol* **84**, 182-191, doi:10.1016/j.pneurobio.2007.10.006 (2008).

28   Stilwell, G. E., Saraswati, S., Littleton, J. T. & Chouinard, S. W. Development of a

Drosophila seizure model for in vivo high-throughput drug screening. *Eur J Neurosci* **24**,

2211-2222, doi:10.1111/j.1460-9568.2006.05075.x (2006).

29   Benzer, S. From the gene to behavior. *JAMA* **218**, 1015-1022 (1971).

30   Ganetzky, B. & Wu, C. F. Indirect Suppression Involving Behavioral Mutants with Altered

Nerve Excitability in DROSOPHILA MELANOGASTER. *Genetics* **100**, 597-614 (1982).

31   Jackson, F. R., Newby, L. M. & Kulkarni, S. J. Drosophila GABAergic systems: sequence

and expression of glutamic acid decarboxylase. *J Neurochem* **54**, 1068-1078 (1990).

32   Sanyal, S. Genomic mapping and expression patterns of C380, OK6 and D42 enhancer

trap lines in the larval nervous system of Drosophila. *Gene Expr Patterns* **9**, 371-380,

doi:10.1016/j.gep.2009.01.002 (2009).

33   Yeh, E., Gustafson, K. & Boulianne, G. L. Green fluorescent protein as a vital marker and

reporter of gene expression in Drosophila. *Proc Natl Acad Sci U S A* **92**, 7036-7040

(1995).

34   McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K. & Davis, R. L. Spatiotemporal

rescue of memory dysfunction in Drosophila. *Science* **302**, 1765-1768,

doi:10.1126/science.1089035 (2003).

35   Zhang, B. & Stewart, B. Electrophysiological recording from Drosophila larval body-wall

muscles. *Cold Spring Harb Protoc* **2010**, pdb prot5487, doi:10.1101/pdb.prot5487

(2010).

36   Jan, L. Y. & Jan, Y. N. Properties of the larval neuromuscular junction in Drosophila

melanogaster. *J Physiol* **262**, 189-214 (1976).

37   Lemon, W. C. *et al.* Whole-central nervous system functional imaging in larval

Drosophila. *Nat Commun* **6**, 7924, doi:10.1038/ncomms8924 (2015).

38   Streit, A. K., Fan, Y. N., Masullo, L. & Baines, R. A. Calcium Imaging of Neuronal Activity in

Drosophila Can Identify Anticonvulsive Compounds. *PLoS One* **11**, e0148461,

doi:10.1371/journal.pone.0148461 (2016).

39   Pulver, S. R. *et al.* Imaging fictive locomotor patterns in larval Drosophila. *J Neurophysiol*

114, 2564-2577, doi:10.1152/jn.00731.2015 (2015).

40   Kohsaka, H., Okusawa, S., Itakura, Y., Fushiki, A. & Nose, A. Development of larval motor

circuits in Drosophila. *Dev Growth Differ* **54**, 408-419, doi:10.1111/j.1440-

169X.2012.01347.x (2012).

41   Frank, C. A., Kennedy, M. J., Goold, C. P., Marek, K. W. & Davis, G. W. Mechanisms

underlying the rapid induction and sustained expression of synaptic homeostasis.

Neuron **52**, 663-677, doi:10.1016/j.neuron.2006.09.029 (2006).

42   Gertner, D. M., Desai, S. & Lnenicka, G. A. Synaptic excitation is regulated by the

postsynaptic dSK channel at the Drosophila larval NMJ. *J Neurophysiol* **111**, 2533-2543,

doi:10.1152/jn.00903.2013 (2014).

43 Kombian, S. B., Hirasawa, M., Mouginot, D., Chen, X. & Pittman, Q. J. Short-term

potentiation of miniature excitatory synaptic currents causes excitation of supraoptic

neurons. *J Neurophysiol* **83**, 2542-2553 (2000).

44 Powers, A. S., Grizzaffi, J. & Lnenicka, G. A. Increased postsynaptic Ca2+ reduces mini

frequency at the Drosophila larval NMJ. *Synapse* **71**, doi:10.1002/syn.21971 (2017).

45 Sutton, M. A. *et al.* Miniature neurotransmission stabilizes synaptic function via tonic

suppression of local dendritic protein synthesis. *Cell* **125**, 785-799,

doi:10.1016/j.cell.2006.03.040 (2006).

46 Halabisky, B., Parada, I., Buckmaster, P. S. & Prince, D. A. Excitatory input onto hilar

somatostatin interneurons is increased in a chronic model of epilepsy. *J Neurophysiol*

104, 2214-2223, doi:10.1152/jn.00147.2010 (2010).

47 Beck, C. L., Fahlke, C. & George, A. L., Jr. Molecular basis for decreased muscle chloride

conductance in the myotonic goat. *Proc Natl Acad Sci U S A* **93**, 11248-11252 (1996).

48 Meyer-Kleine, C., Steinmeyer, K., Ricker, K., Jentsch, T. J. & Koch, M. C. Spectrum of

mutations in the major human skeletal muscle chloride channel gene (CLCN1) leading to

myotonia. *Am J Hum Genet* **57**, 1325-1334 (1995).

49 Pusch, M. Myotonia caused by mutations in the muscle chloride channel gene CLCN1.

Hum Mutat **19**, 423-434, doi:10.1002/humu.10063 (2002).

50 Du, W. *et al.* Calcium-sensitive potassium channelopathy in human epilepsy and

paroxysmal movement disorder. *Nat Genet* **37**, 733-738, doi:10.1038/ng1585 (2005).

51   Yang, J. *et al.* An epilepsy/dyskinesia-associated mutation enhances BK channel

activation by potentiating Ca2+ sensing. *Neuron* **66**, 871-883,

doi:10.1016/j.neuron.2010.05.009 (2010).

52   Brenner, R. *et al.* BK channel beta4 subunit reduces dentate gyrus excitability and

protects against temporal lobe seizures. *Nat Neurosci* **8**, 1752-1759,

doi:10.1038/nn1573 (2005).

53   Dryer, S. E. Na(+)-activated K+ channels: a new family of large-conductance ion

channels. *Trends Neurosci* **17**, 155-160 (1994).

54   Budelli, G. *et al.* Na+-activated K+ channels express a large delayed outward current in

neurons during normal physiology. *Nat Neurosci* **12**, 745-750, doi:10.1038/nn.2313

(2009).

55   Bhattacharjee, A., Gan, L. & Kaczmarek, L. K. Localization of the Slack potassium channel

in the rat central nervous system. *J Comp Neurol* **454**, 241-254, doi:10.1002/cne.10439

(2002).

56   Rizzi, S., Knaus, H. G. & Schwarzer, C. Differential distribution of the sodium-activated

potassium channels slick and slack in mouse brain. *J Comp Neurol* **524**, 2093-2116,

doi:10.1002/cne.23934 (2016).

57   Ganetzky, B. & Wu, C. F. Neurogenetic analysis of potassium currents in Drosophila:

synergistic effects on neuromuscular transmission in double mutants. *J Neurogenet* **1**,

17-28 (1983).

58   Turrigiano, G. Homeostatic synaptic plasticity: local and global mechanisms for

stabilizing neuronal function. *Cold Spring Harb Perspect Biol* **4**, a005736,

doi:10.1101/cshperspect.a005736 (2012).

59   Brooks, D. S., Vishal, K., Kawakami, J., Bouyain, S. & Geisbrecht, E. R. Optimization of

wrMTrck to monitor Drosophila larval locomotor activity. *J Insect Physiol* **93-94**, 11-17,

doi:10.1016/j.jinsphys.2016.07.007 (2016).

60   Frank, C. A., Pielage, J. & Davis, G. W. A presynaptic homeostatic signaling system

composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels. *Neuron*

61, 556-569, doi:10.1016/j.neuron.2008.12.028 (2009).

Acknowledgements: We thank an internal fund for supporting this line of research

from the University of Missouri and Division of Biological Sciences and a postdoctoral

fellowship from King Abdullah International Medical Research Center, Riyadh, Saudi

Arabia for the support of SNE. We thank Kyunghee Koh for the gift of the Shaker

antibody, Leanne Dibbens for sharing fly lines, Weijie Liu for her initial contributions to

this project, and the members of the Zhang lab for helpful discussions.

Author contributions: SNE, GTD, BZ designed the experiments; DD helped design

the molecular cloning of KCNT1 channels; SNE, GTD, PS, and BZ collected and

analyzed data; SNE and BZ wrote the manuscript, GTD, SP, and DD revised or made

comments on the manuscript.

Competing financial interests: This work described her has been filed for a

provisional patent application.

Materials & Correspondence: zhangbing@missouri.edu