CLASSIFICATION

BIOLOGICAL SCIENCES: Population Biology

TITLE (112; 135 CHARACTERS MAX.)

Ancient genomes from North Africa evidence prehistoric migrations to the Maghreb from

both the Levant and Europe

SHORT TITLE (48; 50 CHARACTERS MAX.)

Paleogenomics of Moroccan Neolithic populations

AUTHORS

Rosa Fregel[a*], Fernando L. Méndez[a], Youssef Bokbot[b], Dimas Martín-Socas[c], María D.

Camalich-Massieu[c], Jonathan Santana[d], Jacob Morales[e], María C. Ávila-Arcos[a,f], Peter A.

Underhill[a], Beth Shapiro[g], Genevieve Wojcik[a], Morten Rasmussen[a], Andre E. R. Soares[g§],

Joshua Kapp[g], Alexandra Sockell[a], Francisco J. Rodríguez-Santos[h], Abdeslam Mikdad[b],

Aioze Trujillo-Mederos[c] & Carlos D. Bustamante[a¶]

AFFILIATIONS

[a]Department of Genetics, School of Medicine, Stanford University, 365 Lasuen Street, CA

94305, Stanford, US.

[b]Institut National des Sciences de l'Archéologie et du Patrimoine, Avenue Allal el Fassi,

6828, Rabat, Morocco.

[c]Departmento de Prehistoria, Universidad de La Laguna, Calle Prof. Jose Luis Moreno

Becerra, 38320, San Cristóbal de La Laguna, Spain.

[d]Department of Archaeology, Durham University, South Rd, Durham DH1 3LE, United

Kingdom.

[e]Departamento de Ciencias Históricas, Universidad de Las Palmas de Gran Canaria, Pérez

del Toro 1, 35003, Las Palmas de Gran Canaria, Spain.

[f]International Laboratory for Human Genome Research, National Autonomous University of

Mexico, Blvd. Juriquilla 3001, 76230, Querétaro, Mexico.

[g]Department of Ecology and Evolutionary Biology, University of California, 1156 High

Street, CA 95064, Santa Cruz, US.

[h]Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Avda. de los Castros,

39005, Cantabria, Spain.

CORRESPONDING AUTHOR

Rosa Fregel

Departamento de Genética

Universidad de La Laguna

Avda. Astrofísico Fco. Sánchez, SN

San Cristóbal de La Laguna 38206

Spain

rfregel@ull.edu.es

KEYWORDS

ancient DNA; paleogenomics; Neolithic; North Africa

ABSTRACT (249; 250 WORDS MAX.)

The extent to which prehistoric migrations of farmers influenced the genetic pool of western North Africans remains unclear. Archaeological evidence suggests the Neolithization process may have happened through the adoption of innovations by local Epipaleolithic communities, or by demic diffusion from the Eastern Mediterranean shores or Iberia. Here, we present the first analysis of individuals' genome sequences from early and late Neolithic sites in Morocco, as well as Early Neolithic individuals from southern Iberia. We show that Early Neolithic Moroccans are distinct from any other reported ancient individuals and possess an endemic element retained in present-day Maghrebi populations, confirming a long-term genetic continuity in the region. Among ancient populations, Early Neolithic Moroccans are distantly related to Levantine Natufian hunter-gatherers (~9,000 BCE) and Pre-Pottery Neolithic farmers (~6,500 BCE). Although an expansion in Early Neolithic times is also plausible, the high divergence observed in Early Neolithic Moroccans suggests a long-term isolation and an early arrival in North Africa for this population. This scenario is consistent with early Neolithic traditions in North Africa deriving from Epipaleolithic communities who adopted certain innovations from neighbouring populations. Late Neolithic (~3,000 BCE) Moroccans, in contrast, share an Iberian component, supporting theories of trans-Gibraltar gene flow. Finally, the southern Iberian Early Neolithic samples share the same genetic composition as the Cardial Mediterranean Neolithic culture that reached Iberia ~5,500 BCE. The cultural and genetic similarities of the Iberian Neolithic cultures with that of North African Neolithic sites further reinforce the model of an Iberian migration into the Maghreb.

SIGNIFICANCE STATEMENT (119; 120 WORDS MAX.)

The acquisition of agricultural techniques during the so-called Neolithic revolution has been one of the major steps forward in human history. Using next-generation sequencing and ancient DNA techniques, we directly test if Neolithization in North Africa occurred through the transmission of ideas or by demic diffusion. We show that Early Neolithic Moroccans are

composed of an endemic Maghrebi element still retained in present-day North African

populations and distantly related to Epipaleolithic communities from the Levant. However,

late Neolithic individuals from North Africa are admixed, with a North African and a

European component. Our results support the idea that the Neolithization of North Africa

might have involved both the development of Epipaleolithic communities and the migration

of people from Europe.

INTRODUCTION

One of the greatest transitions in human history was the transition from hunter-gatherer to farming lifestyle. How farming traditions expanded from their birthplace in the Fertile Crescent has been a matter of contention. Two models were proposed: one involving the movement of people and the other based on the transmission of ideas. Over the last decade, paleogenomics has been instrumental in settling long-disputed archaeological questions[1], including those surrounding the Neolithic revolution[2]. Compared to the extensive genetic work done on Europe and the Near East, the Neolithic transition in North Africa, including the Maghreb, remains largely uncharacterized. Archaeological evidence suggests that some of the major innovations associated with the Neolithic, such as farming and pottery production, could have been introduced into northern Morocco through sea voyaging by people from Iberia or the central Mediterranean as early as *ca*. 5400 BCE[3,4]. In fact, some of the Neolithic pottery recorded in North Africa strongly resembles that of European cultures like the Cardial Early Neolithic, the Mediterranean early farmer culture located in Iberia[5]. However, other innovations such as some pottery traditions and bone and lithic technical customs could be the result of *in situ* development from Epipaleolithic communities, indicating a strong continuity in the local population since the Late Pleistocene[6-10].

Genetic data from present-day populations[11-13] suggests that North African ancestry has contributions from four main sources: 1) an autochthonous Maghrebi component related to a back migration to Africa ~12,000 years ago from Eurasia; 2) a Middle Eastern component probably associated with the Arab conquest; 3) a sub-Saharan component derived from trans-Saharan migrations; and 4) a European component that has been linked to recent historic movements. Paleogenomic studies have begun to provide insights into North African Prehistory[14-16]; however, no research to date has tested whether the Neolithic transition in the Maghreb was driven by local populations who adopted cultural and technological innovations or the migration of people. Here, we perform genome-wide analysis of remains from the Early Neolithic site of Ifri n'Amr or Moussa (IAM; ~5,200 BCE, n=7) and the Late Neolithic site of Kelif el Boroud (KEB; ~3,000 BCE; n=8) (Supplementary Note 1). To test possible migrations through the Strait of Gibraltar, we also analyse human remains from the

southern Iberian Early Neolithic site of El Toro (TOR; ~5,000 BCE; n=12) (Figure 1). This

Iberian Early Neolithic culture bears similarities with early Maghrebi pottery decoration, as

well as bone and lithic tool production traditions which suggest an African influence[17]

(Supplementary Note 1). Including these southern Iberian samples in our analysis enables a

direct test of this hypothesis.

[Figure 1 here]

RESULTS AND DISCUSSION

We sequenced 38 Illumina pair-end libraries from 27 individuals, and selected the best-

conserved libraries for subsequent analyses. Endogenous DNA content was generally low

(2.88% on average) (Supplementary Note 2). Depth of coverage was consistently improved

when enriching using baits targeting specific sites for the Multiethnic Genotyping Array

(MEGA) (~100X), compared to whole-genome capture (~15X) (Supplementary Note 2).

Following enrichment, we generated thirteen low-coverage genomes (five from IAM, four

from KEB and four from TOR), with MEGA coverage ranging from 0.04X to 1.72X depth,

and genome-wide coverage ranging from 0.01X to 0.74X depth (Table 1). All samples

considered in this study met the standard aDNA authentication criteria, including observation

of DNA fragmentation (~46 bp average read length) and damage patterns due to cytosine

deamination toward the 5′ ends of molecules (Supplementary Note 3).

Mitochondrial DNA and Y-chromosome haplogroups obtained for IAM (Moroccan Early

Neolithic) and KEB (Moroccan Late Neolithic) suggest either a population replacement or an

important genetic influx into Morocco between 5,000−3,000 BCE. IAM samples belong to

the mtDNA haplogroups U6a and M1—both of which are associated with the back migration

to Africa from Eurasia in Upper Palaeolithic times[18,19]—while KEB samples belong to

haplogroups K1, T2 and X2, prominently found in Anatolian and European Neolithic

samples[2,20] (Supplementary Note 4). Regarding the paternal lineages, IAM individuals carry

Y chromosomes distantly related to the typically North African E-M81 haplogroup, while

the Y chromosome from KEB belongs to the T-M184 haplogroup; though scarce and broadly

distributed today, this haplogroup has also been observed in European Neolithic individuals[16]

(Supplementary Note 5). Both mtDNA and Y-chromosome lineages (K1, J2 and T2

haplogroups, and G-M201 haplogroup, respectively) for samples from TOR (Iberian Early

Neolithic) are similar to those observed in Europe during Neolithic times[20].

[Table 1 here]

When projected on a Principal Components Analysis (PCA) space built using sub-Saharan

African, North African, European and Middle Eastern population of the Human Genome

Diversity Project (HGDP) dataset genotyped with MEGA, IAM samples are placed close to

Mozabites, while Iberian Neolithic samples fall close to southern European populations

(Supplementary Note 6). As suspected from the mtDNA and Y-chromosome data, KEB

samples do not cluster with IAM and are placed in an intermediate position between IAM

and TOR. We further explored the genetic structure of these samples using the program

ADMIXTURE[21] (Figure 2). At K=5, we observe sub-Saharan African (red), early European

Neolithic (green), North African (yellow), Middle Eastern (violet) and eastern European

components (orange). Congruently with PCA results, TOR is composed of the early

European component, clustering with Sardinian samples, and IAM is composed of the North

African component, clustering with Mozabites. Finally, KEB is placed in an intermediate

position, with ~50% of both early European Neolithic and North African ancestries. It is

worth mentioning that, compared to current North African samples, IAM and KEB do not

show any sub-Saharan African ancestry, suggesting that trans-Saharan migrations occurred

after Neolithic times. This is in agreement with the analysis of present-day genome-wide

data from Morocco, which estimated a migration of western African origin into Morocco

only ~1,200 years ago[11].

West Eurasian populations can be modelled as admixture of four different ancestral

components[2]: Eastern and Western European hunter-gatherers, and Iranian and Levantine

Neolithic. We explored the placement of Moroccan and Southern Iberian Neolithic samples

in this context, and compared their genetic affinities to ancient and present-day West

Eurasian and Levant populations in the Human Origins panel. Interestingly, PCA reveals that

IAM individuals are different from any aDNA sample studied to date (Figure 2;

Supplementary Note 6). When projected, IAM samples are close to modern North Africans,

in the Levantine corner of the PCA space (Figure 2). Southern Iberian Neolithic individuals

from TOR cluster with Sardinians and with other Anatolian and European Neolithic samples.

Moreover, KEB samples are placed halfway between the IAM and Anatolian/European

farmer clusters, in close proximity to Levant aDNA samples and also to Guanche samples[16],

the indigenous population of the Canary Islands known to have a Berber origin[22]. When

compared using ADMIXTURE (See Supplementary Note 7 for details), IAM samples

possess ~100% of a component partially shared by aDNA samples from the Middle East and

Levant at low K values. At K=6, this IAM-like component is observed mainly in modern

North Africa, following a west-to-east cline, and in the Guanches. TOR and other Early

Neolithic samples from Iberia cluster together with farmers from Anatolia, the Aegean area

and Europe. At K=8, the Early Neolithic individuals from Iberia differentiate from the

Anatolian, Aegean and European Early Neolithic samples, and share their main component

(purple) with Middle Neolithic/Chalcolitic samples (Figure 2). Finally, at low K values, KEB

can be explained as having both IAM-like and European Neolithic components, suggesting

an admixture process between IAM-like people and early farmers. Nevertheless, at K=8, the

European component in KEB is predominantly "purple," with some "green" component.

This "green" component is also present, at a low frequency, in Natufians and other ancient

Levantine populations. The substantially larger contribution of the "purple" component,

when compared with the "green", suggests a significant genetic contribution of ancient

Iberians in Morocco (Figure 2). The same admixture profile is observed in Guanches, but the

amount of IAM ancestry is consistently higher in all the samples. Given that the Guanches

could have had originated in a different area of the Maghreb, this result might suggest that

the European Neolithic impact in North Africa was heterogeneous.

[Figure 2 here]

To compare our samples directly to the genomes of ancient and modern populations, we

calculated pair-wise F_{ST} distances, which, unlike PCA and global ancestry analyses, are

insensitive to the inclusion of large numbers of individuals from modern populations. F_{ST}

values indicate that the IAM samples are as differentiated from all other populations as

Yoruba are from non-Africans (Supplementary Note 9), with the sole exception of KEB and,

to a lesser extent, the Guanches and modern North African populations. Given the relatively

low heterozygosity and high identity-by-descent proportions observed in IAM

(Supplementary Note 8), this differentiation could be driven by isolation and genetic drift.

IAM is divergent from the other populations, with the exception of populations that likely

received genetic influx from them. This raises the possibility that IAM was isolated in North

Africa since Palaeolithic times, when a back migration from Eurasia brought mtDNA

haplogroups M1 and U6 to the Maghreb[18]. Although IAM is clearly more similar to KEB

than to any other population, the converse is not true. KEB has lower F_{ST} distances with any

Anatolian, European (excluding European hunter gatherers), Levantine and Iranian

population, rather than with IAM. In the modern DNA reference panel, KEB is similar to

North African, European and Middle Eastern populations. Among the ancient populations,

TOR is more similar to Middle Neolithic/Chalcolithic Europeans, and, among modern

populations, to populations from Spain, North Italy and Sardinia.

To further investigate the genetic affinities of IAM, KEB and TOR samples, we conducted

outgroup f3-statistic analysis[23]. Results indicate that, when KEB and Guanches are excluded,

IAM shares more drift with ancient Levantine populations, such as Natufians (Epipaleolithic)

and Pre-Pottery Neolithic individuals (Figure 3; Supplementary Note 10), than with any

other ancient population. To explore further the connection between IAM and Levantine

populations, we performed an f4-statistic analysis to test whether IAM shares more alleles

with any other population in the Human Origins panel[2,24] than with ancient populations from

the Levant (Supplementary Note 10). Consistently, and also with the exception of KEB and

Guanches, all comparisons indicated higher similarity with Natufians and Levantine farmers.

This suggests that most of IAM ancestry originates from an out-of-Africa source, as IAM

shares more alleles with Levantines than with any sub-Saharan Africans, including the

4,500-year-old genome from Ethiopia[14]. To further test the hypothesis that IAM is more

closely related to out-of-Africa populations, we determined if we could detect Neanderthal

ancestry in IAM, which is typical of non-African populations. A signal of Neanderthal

ancestry has been detected in modern North African populations[25]. A lack of Neanderthal

ancestry in IAM would imply that the signal observed today is a product of more recent

migration into North Africa from the Middle East and Europe in historical times. When

compared to the Neanderthal high coverage genome sequence from Altai[26] and the low-

coverage sequence from Vindija Cave[27], and using the S-statistic[23], we detected a

Neanderthal introgression signal into IAM, suggesting derivation from the same event shared

by non-African populations. All these results together indicate that the origin of IAM was

outside Africa, most probably from the Levant. However, it is important to take into account

that the number of ancient genomes for comparison is low and future sampling can provide

further refinement in the origin of IAM.

Both F_{ST} and outgroup-f3 statistic analyses indicate that KEB shares ancestry with IAM, but

also more genetic drift with Neolithic and Chalcolithic populations from Anatolia and

Europe, with the highest shared genetic drift appearing in Iberian Early Neolithic samples

(Figure 3; Supplementary Note 10). This pattern and the result from ADMIXTURE could be

explained if the KEB population was a mixture between IAM-related and European

Neolithic groups. To formally test this hypothesis, we used an admixture-f3 test[23], using

KEB as the test population, IAM as a reference population and one of the Anatolian and

European Neolithic and Chalcolithic populations as the second reference population. All

comparisons produced negative values of the f3-statistic, which suggests the KEB population

can be modelled as a mixture of IAM and Anatolian/European Neolithic.

TOR has more shared ancestry with Iberian Early Neolithic samples and other Neolithic and

Chalcolithic populations from Europe. Archaeological studies have suggested that there was

an Andalusian Early Neolithic culture with North African influences before the Cardial

expansion into the Western Mediterranean basin[28]. However, we observe that TOR samples

have a similar genetic composition to that of Cardial individuals from Iberia, evidencing a

common origin, and ruling out an Andalusian Early Neolithic distinct from Cardial Culture.

[Figure 3 here]

Finally, although limited by low coverage, phenotypic predictions based on genetic variants

of known effects agree with our estimates of global ancestry. IAM people do not possess any

of the European SNPs associated with light pigmentation, and most likely had dark skin and

eyes. IAM samples present ancestral alleles for pigmentation-associated variants present in

SLC24A5 (rs1426654), SLC45A2 (rs16891982) and OCA2 (rs1800401 and 12913832)

genes. On the other hand, KEB individuals exhibit some European- derived alleles that

predispose individuals to lighter skin and eye colour, including those on genes SLC24A5

(rs1426654) and OCA2 (rs1800401) (Supplementary Note 11).

CONCLUSION

Genetic analyses have revealed that the population history of modern North Africans is quite

complex[11]. Based on our aDNA analysis, we identify an Early Neolithic Moroccan

component that is restricted to North Africa in present-day populations[11], and that is the sole

ancestry in IAM samples. We hypothesize that this component represents the autochthonous

Maghrebi ancestry associated with Berber populations. This Maghrebi component is

different from those of any ancient samples studied so far and is distantly related to that of

Epipaleolithic people from the Levant. Our data suggests that the IAM population was

isolated in the Maghreb since the Upper Palaeolithic back migration, although it is

impossible to be certain without paleogenomic data from North African Palaeolithic samples.

An expansion in Early Neolithic times followed by strong genetic drift might also be

plausible.

Our hypothesis is in agreement with archaeological research pointing to the first stage of the

Neolithic expansion in Morocco as the result of a local population who adopted some

technological innovations, such as pottery production or farming, from neighbouring areas.

By 3,000 BCE, a continuity in the Neolithic spread brought Mediterranean-like ancestry to

the Maghreb, most likely from Iberia. Other archaeological remains, such as African

elephant ivory and ostrich eggs found in Iberian sites, confirm the existence of contacts and

exchange networks through both sides of the Gibraltar strait at this time. Our analyses

strongly support that at least some of the European ancestry observed today in North Africa

is related to prehistoric migrations, and local Berber populations were already admixed with

Europeans before the Roman conquest. Furthermore, additional European/Iberian ancestry

could have reached the Maghreb after KEB people; this scenario is supported by the

presence of Iberian-like Bell-Beaker pottery in more recent stratigraphic layers of IAM and

KEB caves. Future palaeogenomic efforts in North Africa will further disentangle the

complex history of migrations that forged the ancestry of the admixed populations we

observe today.

MATERIAL AND METHODS

Measures to avoid and monitor contamination from modern DNA were applied, at all times,

during sample manipulation. Ancient DNA was extracted from teeth or bone, built into

double-stranded indexed libraries and sequenced on an Illumina NextSeq 500

(Supplementary Note 2). Due to the environmental conditions of the burial sites, we

expected to recover low proportions of endogenous DNA from these ancient remains. To

overcome limitations due to DNA degradation, we applied two different capture methods to

enrich for human reads (Supplementary Note 2): one targeting the whole genome[29] and one

targeting the variants of the MEGA array (Illumina Inc.).

Reads were trimmed and adapters removed using AdapterRemoval[30], and then mapped to the

human reference genome (hg19) using BWA[31]. Low quality (MAPQ<30) and duplicate reads

were removed using SAMtools[32]. MapDamage[33] was used to visualize misincorporation and

fragmentation patterns, and to rescale the quality of bases likely affected by post-mortem

damage. Confidence intervals of sex determination were calculated following Skoglund et

al.[34]. MtDNA haplogroups were determined using HaploGrep[35]. Y-chromosome haplogroup

inference was carried out as in Schroeder et al.[36]. As the reference panel, we used both the

Human Origins panel[2] and the HGDP dataset genotyped with MEGA-ex (Illumina Inc.). For

principal component analysis, we projected the aDNA samples on the PCA space built with

the modern dataset, using smartpca[37] and LASER[38]. Admixture estimations were done using

ADMIXTURE software[21]. F_{ST} distances were calculated using smartpca[37]. Identity-by-

descent proportions were estimated using PLINK[39], and heterozygosity estimations using a

newly developed method for low-coverage genomes (Supplementary Note 8). f-statistics

estimates were calculated using admixtools software[23]. All plots were prepared using R

software[40]. Detailed information about methods is included in the Supplementary Notes.

Acknowledgements C.D.B and R.F. were funded by a grant from the National Science Foundation (1201234).

R.F. was funded by Fundación Canaria Dr. Manuel Morales fellowship. A.E.R.S. was funded by Ciencia sem

Fronteiras fellowship - CAPES, Brazil. B.S. and J.K. were funded by a grant from the Gordon and Betty Moore

Foundation (GBMF-3804).

Author Contributions R.F. and C.D.B conceived the idea for the study, D.M.S, M.D.C.M, J.S., J.M, Y.B.,

F.J.R.S, A.M. and A.T.M assembled skeletal material and provided archaeological background for the samples,

R.F., A.E.R.S., J.K. and A.S. performed work in the wet laboratory, F.M. developed methods for data analysis,

M.R. developed methods for DNA capture, R.F., F.M., M.A.A, P.A.U, G.W., analysed data, C.D.B and B.S.

supervised the study, R.F., F.M. and C.D.B. wrote the manuscript and supplements with input from all co-

authors.

Author Information Sequence data are available through the European Nucleotide Archive (PRJEB22699).

Consensus mtDNA sequences are available at the National Center of Biotechnology Information (Accession

Numbers MF991431-MF991448). The authors declare no competing financial interests. Correspondence and

requests for materials should be addressed to R.F. (e-mail: rfregel@stanford.edu).

REFERENCES

1   Nielsen, R. *et al.* Tracing the peopling of the world through genomics. *Nature* **541**,
302-310, doi:10.1038/nature21347 (2017).

2   Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near
East. *Nature* **536**, 419-424, doi:10.1038/nature19310 (2016).

3   Linstädter, J., Medved, I., Solich, M. & Weniger, G. C. Neolithisation process within
the Alboran territory: Models and possible African impact. *Quaternary International*
274, 219 - 232, doi:10.1016/j.quaint.2012.01.013 (2012).

4   Zilhão, J. Early prehistoric navigation in the Western Mediterranean: Implications for
the Neolithic transition in Iberia and the Maghreb. **Eurasian Prehistory**, 185-200
(2014).

5   Martínez-Sánchez, R. M., Vera-Rodríguez, J. C., Pérez-Jordà, G., Peña-Chocarro, L.
& Bokbot, Y. The beginning of the Neolithic in northwestern Morocco. *Quaternary*
International **In press** (2017).

6   Mulazzani, S. *et al.* The emergence of the Neolithic in North Africa: A new model
for the Eastern Maghreb. *Quaternary International* **410**, 123 - 143,
doi:10.1016/j.quaint.2015.11.089 (2016).

7   Barton, N. *et al.* Human Burial Evidence from Hattab II Cave and the Question of
Continuity in Late Pleistocene–Holocene Mortuary Practices in Northwest Africa.
Cambridge Archaeological Journal **18**, 195–214, doi:10.1017/S0959774308000255
(2008).

8   Lubell, D., Sheppard, P. & Jackes, M. in *Advances in World Archaeology* (eds F.
Wendorf & A. Close) 146-191 (Academic Press, 1984).

9   Linstädter, J. Epipalaeolithic-Neolithic-Transition in the Mediterranean region of
Northwest-Africa. *Quartär* **55**, 41-62 (2008).

10   de Groote, I. & Humphrey, L. T. Characterizing evulsion in the Later Stone Age
Maghreb: Age, sex and effects on mastication. *Quaternary International* **413**, 50 - 61,
doi:https://doi.org/10.1016/j.quaint.2015.08.082 (2016).

11 Henn, B. M. *et al.* Genomic Ancestry of North Africans Supports Back-to-Africa
Migrations. *Plos Genetics* **8**, e1002397-e1002397, doi:10.1371/journal.pgen.1002397
(2012).
12 Arauna, L. R. *et al.* Recent Historical Migrations Have Shaped the Gene Pool of
Arabs and Berbers in North Africa. *Mol Biol Evol* **34**, 318-329,
doi:10.1093/molbev/msw218 (2017).
13 Fadhlaoui-Zid, K. *et al.* Genome-wide and paternal diversity reveal a recent origin of
human populations in North Africa. *PLoS One* **8**, e80293,
doi:10.1371/journal.pone.0080293 (2013).
14 Gallego Llorente, M. *et al.* Ancient Ethiopian genome reveals extensive Eurasian
admixture throughout the African continent. *Science* **350**, 820-822,
doi:10.1126/science.aad2879 (2015).
15 Schuenemann, V. J. *et al.* Ancient Egyptian mummy genomes suggest an increase of
Sub-Saharan African ancestry in post-Roman periods. *Nat Commun* **8**, 15694,
doi:10.1038/ncomms15694 (2017).
16 Rodriguez-Varela, R. *et al.* Genomic Analyses of Pre-European Conquest Human
Remains from the Canary Islands Reveal Close Affinity to Modern North Africans.
Curr Biol **27**, 3396-3402 e3395, doi:10.1016/j.cub.2017.09.059 (2017).
17 Garcia-Borja, P., Aura-Tortosa, J. E., Bernabeu-Auban, J. & Jorda-Pardo, J. F.
Nuevas perspectivas sobre la neolitizacion en la cueva de Nerja (Malaga-España): La
ceramica de la sala del vestibulo. *Zephyrus* **LXVI**, 109-132 (2010).
18 Pennarun, E. *et al.* Divorcing the Late Upper Palaeolithic demographic histories of
mtDNA haplogroups M1 and U6 in Africa. *BMC Evol Biol* **12**, 234,
doi:10.1186/1471-2148-12-234.; eng; ID: 4230 (2012).
19 Secher, B. *et al.* The history of the North African mitochondrial DNA haplogroup U6
gene flow into the African, Eurasian and American continents. *BMC evolutionary*
biology **14**, 109, doi:1471-2148-14-109 [pii] (2014).
20 Haak, W. *et al.* Ancient DNA from European early neolithic farmers reveals their
near eastern affinities. *PLoS Biol* **8**, e1000536 (2010).

21   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry
in unrelated individuals. *Genome research* **19**, 1655-1664,
doi:10.1101/gr.094052.109 [doi] (2009).

22   Maca-Meyer, N. *et al.* Ancient mtDNA analysis and the origin of the Guanches. *Eur
J Hum Genet* **12**, 155-162 (2004).

23   Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* **192**, 1065-+,
doi:10.1534/genetics.112.145037 (2012).

24   Lazaridis, I. *et al.* Ancient human genomes suggest three ancestral populations for
present-day Europeans. *Nature* **513**, 409-413, doi:10.1038/nature13673 (2014).

25   Sanchez-Quinto, F. *et al.* North African populations carry the signature of admixture
with Neandertals. *PLoS One* **7**, e47765, doi:10.1371/journal.pone.0047765 (2012).

26   Prufer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai
Mountains. *Nature* **505**, 43-49, doi:10.1038/nature12886 [doi] (2014).

27   Green, R. E. *et al.* A draft sequence of the Neandertal genome. *Science* **328**, 710-722
(2010).

28   Cortes Sanchez, M. *et al.* The Mesolithic-Neolithic transition in southern Iberia.
Quaternary Research **77**, 221-234 (2012).

29   Carpenter, M. L. *et al.* Pulling out the 1%: Whole-Genome Capture for the Targeted
Enrichment of Ancient DNA Sequencing Libraries. *American Journal of Human
Genetics* **93**, 852-864, doi:10.1016/j.ajhg.2013.10.002 (2013).

30   Lindgreen, S. AdapterRemoval: easy cleaning of next-generation sequencing reads.
BMC research notes **5**, 337, doi:10.1186/1756-0500-5-337 (2012).

31   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

32   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

33   Ginolhac, A., Rasmussen, M., Gilbert, M. T., Willerslev, E. & Orlando, L.
mapDamage: testing for damage patterns in ancient DNA sequences. *Bioinformatics*
27, 2153-2155, doi:10.1093/bioinformatics/btr347 (2011).

34   Skoglund, P., Stora, J., Gotherstrom, A. & Jakobsson, M. Accurate sex identification
of ancient human remains using DNA shotgun sequencing. *Journal of*
Archaeological Science **40**, 4477-4482 (2013).

35   Kloss-Brandstatter, A. *et al.* HaploGrep: a fast and reliable algorithm for automatic
classification of mitochondrial DNA haplogroups. *Hum Mutat* **32**, 25-32,
doi:10.1002/humu.21382 (2011).

36   Schroeder, H. *et al.* Genome-wide ancestry of 17th-century enslaved Africans from
the Caribbean. *Proc Natl Acad Sci U S A* **112**, 3669-3673,
doi:10.1073/pnas.1421784112 (2015).

37   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-
wide association studies. *Nat Genet* **38**, 904-909, doi:10.1038/ng1847 (2006).

38   Wang, C. *et al.* Ancestry estimation and control of population stratification for
sequence-based association studies. *Nat Genet* **46**, 409-415, doi:10.1038/ng.2924
(2014).

39   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
based linkage analyses. *American Journal of Human Genetics* **81**, 559-575,
doi:10.1086/519795 (2007).

40   A language and environment for statistical computing (R Foundation for Statistical
Computing. Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org,
2008).

FIGURES

Figure 1. Geographical location (A) and calibrated radiocarbon date (B) of the

samples included in this study, as well as other ancient DNA samples from the

literature.

Figure 2. Ancestry inference in ancient samples from North Africa and the Iberian

Peninsula. (A) PCA analysis using the Human Origins panel, (B) ADMIXTURE

analysis using the HGDP-MEGA dataset (K=5), (C) ADMIXTURE analysis using

the Human Origins dataset for modern and ancient populations (K=8), and (D)

detailed ADMIXTURE analysis for European Neolithic samples (K=8).

Figure 3. Outgroup f3-statistic for IAM (A), KEB (B) and TOR (C), and admixture
f3-statistic for KEB (D).

Table 1 - Summary statistics for North African and Iberian samples.

Sample	Origin	2 sigma calibrated radiocarbon date (BCE)	MtDNA coverage	MtDNA haplogroup	Contamination estimation	Molecular sex	Y-chromosome haplogroup	SNPs in MEGA dataset	SNPs in Human Origins dataset	MEGA coverage	Genome-wide coverage
IAM.3	Ifri n'Amr or Moussa	5325 - 5210	5X	M1b1	6.42% (21.49% - 1.90%)	XX	-	26,798	5,353	0.04X	0.01X
IAM.4	Ifri n'Amr or Moussa	5215 - 5005	52X	U6a1b	4.66% (6.64% - 3.26%)	XY	E-L19*	62,729	17,093	0.12X	0.02X
IAM.5	Ifri n'Amr or Moussa	5199 - 5176 / 5066 - 4942	82X	U6a1b	3.68% (5.26% - 2.62%)	XY	E-L19*	408,744	208,661	1.72X	0.74X
IAM.6	Ifri n'Amr or Moussa	5290 - 5265 / 5230 - 5195 / 5180 - 5060	21X	U6a7b2	5.28% (8.48% - 3.02%)	XX	-	90,226	18,285	0.17X	0.02X
IAM.7	Ifri n'Amr or Moussa	5000 - 4840	129X	U6a3	2.35% (3.39% - 1.67%)	XY	-	177,041	45,134	0.38X	0.06X
KEB.1	Kelif el Boroud		18X	X2b	3.03% (6.31% - 1.23%)	XX	-	96,946	39,312	0.16X	0.08X
KEB.3	Kelif el Boroud		20X	K1a1b1	3.16% (5.73% - 1.72%)	-	-	-	-	-	-
KEB.4	Kelif el Boroud	3780 - 3650	135X	K1a1b1	2.49% (3.27% - 1.89%)	XX	-	304,607	77,964	0.64X	0.12X
KEB.6	Kelif el Boroud		14X	K1a4a1	3.77% (6.72% - 1.98%)	XY	T-M184	95,152	52,694	0.16X	0.14X
KEB.7	Kelif el Boroud		12X	T2b3	1.44% (4.31% - 0.31%)	XY	-	-	-	-	-
KEB.8	Kelif el Boroud		23X	X2b	3.03% (6.23% - 1.27%)	XX	-	34,385	18,974	0.06X	0.05X
TOR.1	El Toro	5280 - 4780	170X	T2c1d	1.42% (2.07% - 0.98%)	-	-	-	-	-	-
TOR.5	El Toro	5040 - 4850	18X	J2b1a	4.47% (8.76% - 1.96%)	XY	G-M201	-	-	-	-
TOR.6	El Toro		53X	T2b3	4.15% (5.89% - 2.94%)	XX	-	318,110	101,698	0.67X	0.17X
TOR.7	El Toro		126X	T2b3	7.06% (8.36% - 5.88%)	XX	-	115,652	24,719	0.18X	0.03X
TOR.8	El Toro	5280 - 4780	234X	K1a1	3.95% (4.80% - 3.26%)	XX	-	212,370	46,364	0.38X	0.06X
TOR.11	El Toro		124X	K1a2a	3.88% (5.00% - 2.98%)	XX	-	80,687	17,306	0.15X	0.03X
TOR.12	El Toro		174X	J2b1a	1.50% (2.30% - 0.93%)	XY	-	-	-	-	-