Neolithization of North Africa involved the migration of people from both the Levant and Europe

Rosa Fregel[1*], Fernando L. Méndez[1], Youssef Bokbot[2], Dimas Martín-Socas[3], María D. Camalich-Massieu[3], María C. Ávila-Arcos[1,4], Peter A. Underhill[1], Beth Shapiro[5], Genevieve Wojcik[1], Morten Rasmussen[1], Andre E. R. Soares[5§], Joshua Kapp[5], Alexandra Sockell[1], Francisco J. Rodríguez-Santos[6], Abdeslam Mikdad[2], Jonathan Santana[7], Aioze Trujillo-Mederos[3] & Carlos D. Bustamante[1¶]

[1]*Department of Genetics, School of Medicine, Stanford University, Stanford, US.* [2]*Institut National des Sciences de l'Archéologie et du Patrimoine, Rabat, Morocco.* [3]*Department of Prehistory, Faculty of History, Universidad de La Laguna, San Cristóbal de La Laguna, Spain.* [4]*International Laboratory for Human Genome Research, National University of Mexico, Querétaro, Mexico.* [5]*Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, US.* [6]*Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Cantabria, Spain.* [7]*Department of Archaeology, Faculty of Arts and Humanities. Durham University, Durham, UK.*

[¶]Present address: Department of Biomedical Data Science, School of Medicine, Stanford University, Stanford, US.

[§]Present address: Laboratório Nacional de Computação Científica, Petrópolis, Brazil

One of the greatest transitions in the human story was the change from hunter-gatherer to farmer. How farming traditions expanded from their birthplace in the Fertile Crescent has always been a matter of contention. Two models were proposed, one involving the movement of people and the other based on the transmission of ideas. Over the last decade, paleogenomics has been instrumental in settling long-disputed archaeological questions[1], including those surrounding the Neolithic revolution[2]. Compared to the extensive genetic work done on Europe and the Near East, the Neolithic transition in North Africa, including the Maghreb, remains largely uncharacterized. Archaeological evidence suggests this process may have happened through an *in situ* development from Epipaleolithic communities[3,4], or by demic diffusion from the Eastern Mediterranean shores[5] or Iberia[6]. In fact,

Neolithic pottery in North Africa strongly resembles that of European cultures like Cardial and Andalusian Early Neolithic, the southern-most early farmer culture from Iberia. Here, we present the first analysis of individuals' genome sequences from early and late Neolithic sites in Morocco, as well as Andalusian Early Neolithic individuals. We show that Early Neolithic Moroccans are distinct from any other reported ancient individuals and possess an endemic element retained in present-day Maghrebi populations, indicating long-term genetic continuity in the region. Among ancient populations, early Neolithic Moroccans share affinities with Levantine Natufian hunter-gatherers (~9,000 BCE) and Pre-Pottery Neolithic farmers (~6,500 BCE). Late Neolithic (~3,000 BCE) Moroccan remains, in comparison, share an Iberian component of a prominent European-wide demic expansion, supporting theories of trans-Gibraltar gene flow. Finally, the Andalusian Early Neolithic samples share the same genetic composition as the Cardial Mediterranean Neolithic culture that reached Iberia ~5,500 BCE. The cultural and genetic similarities of the Iberian Neolithic cultures with that of North African Neolithic sites further reinforce the model of an Iberian intrusion into the Maghreb.

Present genetic data from modern samples[7-9] suggests that North Africans' ancestry has contributions from four main sources: 1) an autochthonous Maghrebi component related to a back migration to Africa ~12,000 years ago from the Levant; 2) a Middle Eastern component probably associated with the Arab conquest; 3) a sub-Saharan component derived from trans-Saharan migrations; and 4) a European component that has been linked to recent historic movements. Paleogenomic studies have begun to provide insights into North African Prehistory[10,11]; however, no research to date has tested whether the Neolithic transition in the Maghreb was driven by an *in situ* development or the migration of people. Here, we perform genome-wide analysis of remains from the Early Neolithic site of Ifri n'Amr or Moussa (IAM; ~5,000 BCE, n=7) and the Late Neolithic site of Kelif el Boroud (KEB; ~3,000 BCE; n=8) (Supplementary Note 1). To test possible migrations through the Strait of Gibraltar, we also analyse samples from Early Neolithic sites in southern Iberia: El Toro (TOR; ~5,000 BCE; n=12) and Los Botijos (BOT; n=1) (Figure 1). This Andalusian Early Neolithic culture is thought to have arrived prior to Cardial technology, and bears similarities with early Maghrebi

farming traditions[12] (Supplementary Note 1). Including these southern samples in our

analysis enables a direct test of this hypothesis.

[Figure 1]

We sequenced 39 Illumina pair-end libraries from 28 individuals, and selected best

conserved libraries for subsequent analyses. Endogenous DNA content was generally low

(2.06% on average) (Supplementary Note 2). Depth of coverage was consistently

improved when enriching using baits targeting specific sites (~100X), compared to

whole-genome capture (~15X) (Supplementary Note 2). Following enrichment, we

generated fifteen low-coverage genomes (5 from IAM, 4 from KEB and 5 from

TOR/BOT), with coverage ranging from 0.04X to 0.73X depth. All samples considered

in this study met the standard aDNA authentication criteria, including observation of

DNA fragmentation (~46 bp average read length) and damage patterns due to cytosine

deamination toward the 5′ ends of molecules (Supplementary Note 3).

Mitochondrial DNA and Y-chromosome haplogroups obtained for IAM (Moroccan

Early Neolithic) and KEB (Moroccan Late Neolithic) indicate either a population

replacement or an important genetic influx into Morocco between 5,000−3,000 BCE.

IAM samples belong to the mtDNA haplogroups U6a and M1—both of which are

associated with back migration to Africa[13,14]—while KEB samples belong to haplogroups

K1, T2 and X2, prominently found in Anatolian and European Neolithic samples[2,15]

(Supplementary Note 4). Regarding the paternal lineages, IAM individuals carry Y

chromosomes distantly related to the typically North African E-M81 haplogroup, while

the Y chromosome from KEB belongs to the T-M184 haplogroup; though scarce and

broadly distributed today, this haplogroup has also been observed in European Neolithic

individuals[16] (Supplementary Note 5). Both mtDNA and Y-chromosome lineages (K1, J2

and T2 haplogroups, and G-M201 haplogroup, respectively) for samples from TOR/BOT

(Iberian Early Neolithic) are similar to those observed in Europe during Neolithic times[15].

West Eurasian populations can be modelled as admixture of four different ancestral

components[2]: Eastern and Western European hunter-gatherers, Iranian and Levant

Neolithic. We explored the placement of Moroccan and Southern Iberian Neolithic

samples in this context, and compared their genetic affinities to ancient and present-day

West Eurasian and Levant populations. Interestingly, Principal Components Analysis

(PCA) reveals that IAM individuals are different from any aDNA sample studied to date

(Figure 2; Supplementary Note 6). When projected, IAM samples are close to modern

North Africans, in the Levantine corner of the PCA space (Figure 2). Southern Iberian

Neolithic individuals from TOR and BOT cluster with Sardinians and with other

Anatolian and European Neolithic samples. Moreover, KEB samples are placed halfway

between the IAM and Anatolian/European farmer clusters, in close proximity to Levant

aDNA samples. We further explored the genetic structure of these samples using the

program ADMIXTURE[16] (See Supplementary Note 7 for details), with values of K

ranging between 2 and 8. At lower K values, IAM samples possess ~100% of a

component partially shared by aDNA samples from the Middle East and Levant. At K=6,

this IAM-like component is observed mainly in modern North Africa, following a west-

to-east cline. TOR and other Early Neolithic samples from Iberia cluster together with

farmers from Anatolia, the Aegean area and Europe. At K=8 the Early Neolithic

individuals from Iberia differentiate from the Anatolian, Aegean and European Early

Neolithic samples, and share their main component (purple) with Middle

Neolithic/Chalcolitic samples (Figure 2). Finally, at low K values, KEB can be explained

as having both IAM-like and European Neolithic components, suggesting an admixture

process between IAM-like people and early farmers. Nevertheless, at K=8, the European

component in KEB is predominantly "purple," with some "green" component. This

"green" component is also present, at a low frequency, in Natufians and other ancient

Levantine populations. The substantially larger contribution of the "purple" component,

when compared with the "green", suggests a significant genetic contribution of ancient

Iberians in Morocco (Figure 2).

[Figure 2 here]

To compare our samples directly to the genomes of ancient and modern populations, we

calculated pair-wise F_{ST} distances, which, unlike PCA and global ancestry analyses, are

insensitive to the inclusion of large numbers of individuals from modern populations. F_{ST}

values indicate that the IAM samples are as differentiated from all other populations as

Yoruba are from non-Africans (Supplementary Note 9), with the sole exception of KEB

and, to a lesser extent, modern North African populations. Given the relatively low

heterozygosity (~6.6 x 10^{-4}) and high identity-by-descent (~37%) proportions observed in

IAM (Supplementary Note 8), this differentiation could be related to isolation and genetic

drift. Although IAM is clearly more similar to KEB than any other population, the

converse is not true. KEB has lower F_{ST} distances with any Anatolian, European

(excluding European hunter gatherers), Levantine and Iranian population, rather than

with IAM. In the modern DNA reference panel, KEB is similar to North African,

European and Middle Eastern populations. Among the ancient populations, TOR is more

similar to Middle Neolithic/Chalcolithic Europeans, and, among modern populations, to

populations from Spain, North Italy and Sardinia.

To further investigate the genetic affinities of IAM, KEB and TOR samples, we

conducted outgroup f3-statistic analysis[17]. Results indicate that, when KEB is excluded,

IAM shares more drift with ancient Levantine populations, such as Natufians and Levant

Neolithic (Figure 3; Supplementary Note 10), than with any other ancient population.

This result is analogous to preliminary archaeological data pointing to links between

funerary practices in IAM and Pre-Pottery Neolithic B sites in Cyprus (Supplementary

Note 1).

Both F_{ST} and outgroup-f3 statistic analyses indicate that KEB shares ancestry with

IAM, but also more genetic drift with Neolithic and Chalcolithic populations from

Anatolia and Europe, with the highest shared genetic drift appearing in Iberian Early

Neolithic samples (Figure 3; Supplementary Note 10). This pattern and the result from

ADMIXTURE could be explained if the KEB population was a mixture between IAM-

related and European Neolithic groups. To formally test this hypothesis, we used an

admixture-f3 test[17], using KEB as the test population, IAM as a reference population and

one of the Anatolian and European Neolithic and Chalcolithic populations as the second

reference population. All comparisons produced negative values of the f3-statistic, which

suggests the KEB population can be modelled as a mixture of IAM and

Anatolian/European Neolithic. These results also parallel archaeological findings in the

region: Late Neolithic sites in North Africa contain pottery resembling that of the

Andalusian Early Neolithic and Cardial cultures, and ivory tools distinctly associated

with those of Iberian Neolithic sites (Supplementary Note 1).

TOR has more shared ancestry with Iberian Early Neolithic samples and other

Neolithic and Chalcolithic populations from Europe. Archaeological work in southern

Iberia, especially in the Nerja site, has pointed out that the Andalusian Early Neolithic

culture, previous to the Cardial expansion, may have had connections to farmer traditions

in the Maghreb[18]. However, we observe that TOR samples have a similar genetic

composition to that of Cardial individuals from Iberia, evidencing a common origin.

[Figure 3 here]

Finally, phenotypic predictions based on genetic variants of known effects agree with our

estimates of global ancestry. IAM people do not possess any of the European SNPs

associated with light pigmentation, and most likely had dark skin and eyes. IAM samples

present ancestral alleles for pigmentation-associated variants present in SLC24A5

(rs1426654), SLC45A2 (rs16891982) and OCA2 (rs16891982 and 12913832) genes. On

the other hand, KEB individuals exhibit some European- derived alleles that predispose

individuals to lighter skin and eye colour, including those on genes SLC24A5

(rs1426654) and OCA2 (rs16891982) (Supplementary Note 11).

Genetic analyses have revealed that the population history of modern North

Africans is quite complex[7]. Based on our aDNA analysis, we identify an Early Neolithic

Moroccan component that is restricted to North Africa in present-day populations[7], which

is the sole ancestry in IAM samples. We hypothesize that this component represents the

autochthonous Maghrebi ancestry associated with Berber populations. This Maghrebi

component was related to that of Epipaleolithic and Pre-Pottery Neolithic people from

the Levant. By 3,000 BCE, a European Neolithic expansion brought Mediterranean-like

ancestry to the Maghreb, most likely from Iberia. Our analyses demonstrate that at least

some of the European ancestry observed today in North Africa is related to prehistoric

migrations, and local Berber populations were already admixed with Europeans before

the Roman conquest. Furthermore, additional European/Iberian ancestry could have

reached the Maghreb after KEB people; this scenario is supported by the presence of

Iberian-like Bell-Beaker pottery in more recent stratigraphic layers of IAM and KEB

caves. Future palaeogenomic efforts in North Africa will help further disentangle the

complex history of migrations that forged the ancestry of many admixed populations we

observe today.

Methods

Measures to avoid and monitor contamination from modern DNA were applied, at all

times, during sample manipulation. Ancient DNA was extracted from teeth or bone, built

into double-stranded indexed libraries and sequenced on an Illumina NextSeq 500

(Supplementary Note 2). Due to the environmental conditions of the burial sites, we

expected to recover low proportions of endogenous DNA from these ancient remains. To

overcome limitations due to DNA degradation, we applied two different capture methods

to enrich for human reads (Supplementary Note 2): one targeting the whole genome[19] and

one targeting the variants of the Multiethnic Genotyping Array (MEGA) array (Illumina

Inc.).

Reads were trimmed and adapters removed using AdapterRemoval[20], and then

mapped to the human reference genome (hg19) using BWA[21]. Low quality (MAPQ<30)

and duplicate reads were removed using SAMtools[22]. MapDamage[23] was used to

visualize misincorporation and fragmentation patterns, and to rescale the quality of bases

likely affected by post-mortem damage. Confidence intervals of sex determination were

calculated following Skoglund et al.[24]. MtDNA haplogroups were determined using

HaploGrep[25]. Y-chromosome haplogroup inference was carried out as in Schroeder et

al.[26]. As the reference panel, we used both the Human Origins panel[2] and the HGDP

dataset genotyped with MEGA-ex (Illumina Inc.). For principal component analysis, we

projected the aDNA samples on the PCA space built with the modern dataset, using

smartpca[27] and LASER[28]. Admixture estimations were done using ADMIXTURE

software[16]. F_{ST} distances were calculated using smartpca[27]. Identity-by-descent

proportions were estimated using PLINK[29], and heterozygosity estimations using a newly

developed method for low-coverage genomes (Supplementary Note 8). f-statistics

estimates were calculated using admixtools software[17]. All plots were prepared using R

software[30]. Detailed information about methods is included in the Supplementary Notes.

Acknowledgements R.F. and C.D.B were funded by a grant from the National Science Foundation

(1201234). R.F. was funded by Fundación Canaria Dr. Manuel Morales fellowship. A.E.R.S. was funded

by Ciencia sem Fronteiras fellowship - CAPES, Brazil. BS and JK were funded by a grant from the Gordon

and Betty Moore Foundation (GBMF-3804).

Author Contributions R.F. and C.D.B conceived the idea for the study, Y.B., D.M.S, M.D.C.M, F.J.R.S,

A.M., J.S. and A.T.M assembled skeletal material, R.F., A.E.R.S., J.K. and A.S. performed wet laboratory

work, F.M. developed methods for data analysis, M.R. developed methods for DNA capture, R.F., F.M.,

M.A.A, P.A.U, G.W., analysed data, C.D.B and B.S. supervised the study, R.F., F.M. and C.D.B. wrote the

manuscript and supplements with input from all co-authors.

Author Information Sequence data are available through the European Nucleotide Archive

(PRJEB22699). Consensus mtDNA sequences are available at the National Center of Biotechnology

Information (Accession Numbers XXXXXXX-XXXXXXX). The authors declare no competing financial

interests. Correspondence and requests for materials should be addressed to R.F. (e-mail:

rfregel@stanford.edu).

References:

1 Nielsen, R. *et al.* Tracing the peopling of the world through genomics. *Nature*
541, 302-310, doi:10.1038/nature21347 (2017).

2 Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near
East. *Nature* **536**, 419-424, doi:10.1038/nature19310 (2016).

3 Mulazzani, S. *et al.* The emergence of the Neolithic in North Africa: A new
model for the Eastern Maghreb. *Quaternary International* **410**, 123 - 143,
doi:http://dx.doi.org/10.1016/j.quaint.2015.11.089 (2016).

4 Barton, N. *et al.* Human Burial Evidence from Hattab II Cave and the Question of
Continuity in Late Pleistocene–Holocene Mortuary Practices in Northwest Africa.
Cambridge Archaeological Journal **18**, 195–214, doi:
https://doi.org/10.1017/S0959774308000255 (2008).

5 Linstädter, J., Medved, I., Solich, M. & Weniger, G. C. Neolithisation process
within the Alboran territory: Models and possible African impact. *Quaternary*
International **274**, 219 - 232, doi:http://dx.doi.org/10.1016/j.quaint.2012.01.013
(2012).

6 Zilhão, J. Early prehistoric navigation in the Western Mediterranean: Implications
for the Neolithic transition in Iberia and the Maghreb. *Eurasian Prehistory* **11**,
185-200 (2014).

7 Henn, B. M. *et al.* Genomic Ancestry of North Africans Supports Back-to-Africa
Migrations. *Plos Genetics* **8**, e1002397-e1002397,
doi:10.1371/journal.pgen.1002397 (2012).

8 Arauna, L. R. *et al.* Recent Historical Migrations Have Shaped the Gene Pool of
Arabs and Berbers in North Africa. *Mol Biol Evol* **34**, 318-329,
doi:10.1093/molbev/msw218 (2017).

9 Fadhlaoui-Zid, K. *et al.* Genome-wide and paternal diversity reveal a recent
origin of human populations in North Africa. *PLoS One* **8**, e80293,
doi:10.1371/journal.pone.0080293 (2013).

10 Gallego Llorente, M. *et al.* Ancient Ethiopian genome reveals extensive Eurasian
admixture throughout the African continent. *Science* **350**, 820-822,
doi:10.1126/science.aad2879 (2015).

11   Schuenemann, V. J. *et al.* Ancient Egyptian mummy genomes suggest an increase

of Sub-Saharan African ancestry in post-Roman periods. *Nat Commun* **8**, 15694,

doi:10.1038/ncomms15694 (2017).

12   Garcia-Borja, P., Aura-Tortosa, J. E., Bernabeu-Auban, J. & Jorda-Pardo, J. F.

Nuevas perspectivas sobre la neolitizacion en la cueva de Nerja (Malaga-España):

La ceramica de la sala del vestibulo. *Zephyrus* **LXVI**, 109-132 (2010).

13   Pennarun, E. *et al.* Divorcing the Late Upper Palaeolithic demographic histories

of mtDNA haplogroups M1 and U6 in Africa. *BMC Evol Biol* **12**, 234,

doi:10.1186/1471-2148-12-234.; eng; ID: 4230 (2012).

14   Secher, B. *et al.* The history of the North African mitochondrial DNA haplogroup

U6 gene flow into the African, Eurasian and American continents. *BMC*

evolutionary biology **14**, 109, doi:1471-2148-14-109 [pii] (2014).

15   Haak, W. *et al.* Ancient DNA from European early neolithic farmers reveals their

near eastern affinities. *PLoS Biol* **8**, e1000536 (2010).

16   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of

ancestry in unrelated individuals. *Genome research* **19**, 1655-1664,

doi:10.1101/gr.094052.109 [doi] (2009).

17   Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* **192**, 1065-+,

doi:10.1534/genetics.112.145037 (2012).

18   Cortes Sanchez, M. *et al.* The Mesolithic-Neolithic transition in southern Iberia.

Quaternary Research **77**, 221-234 (2012).

19   Carpenter, M. L. *et al.* Pulling out the 1%: Whole-Genome Capture for the

Targeted Enrichment of Ancient DNA Sequencing Libraries. *American Journal of*

Human Genetics **93**, 852-864, doi:10.1016/j.ajhg.2013.10.002 (2013).

20   Lindgreen, S. AdapterRemoval: easy cleaning of next-generation sequencing

reads. *BMC research notes* **5**, 337, doi:10.1186/1756-0500-5-337 (2012).

21   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics* **25**, 1754-1760,

doi:10.1093/bioinformatics/btp324 (2009).

22   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*

25, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

23   Ginolhac, A., Rasmussen, M., Gilbert, M. T., Willerslev, E. & Orlando, L.
mapDamage: testing for damage patterns in ancient DNA sequences.
Bioinformatics **27**, 2153-2155, doi:10.1093/bioinformatics/btr347 (2011).
24   Skoglund, P., Stora, J., Gotherstrom, A. & Jakobsson, M. Accurate sex
identification of ancient human remains using DNA shotgun sequencing. *Journal*
of Archaeological Science **40**, 4477-4482 (2013).
25   Kloss-Brandstatter, A. *et al.* HaploGrep: a fast and reliable algorithm for
automatic classification of mitochondrial DNA haplogroups. *Hum Mutat* **32**, 25-
32, doi:10.1002/humu.21382 (2011).
26   Schroeder, H. *et al.* Genome-wide ancestry of 17th-century enslaved Africans
from the Caribbean. *Proc Natl Acad Sci U S A* **112**, 3669-3673,
doi:10.1073/pnas.1421784112 (2015).
27   Price, A. L. *et al.* Principal components analysis corrects for stratification in
genome-wide association studies. *Nat Genet* **38**, 904-909, doi:10.1038/ng1847
(2006).
28   Wang, C. *et al.* Ancestry estimation and control of population stratification for
sequence-based association studies. *Nat Genet* **46**, 409-415, doi:10.1038/ng.2924
(2014).
29   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
based linkage analyses. *American Journal of Human Genetics* **81**, 559-575,
doi:10.1086/519795 (2007).
30   A language and environment for statistical computing (R Foundation for
Statistical Computing. Vienna, Austria. ISBN 3-900051-07-0, URL
http://www.R-project.org, 2008).
310
311

Figure 1. Geographical location (A) and calibrated radiocarbon date (B) of the samples included in

this study, as well as other ancient DNA samples from the literature. *BOT sample was not

radiocarbon dated, but was assigned to the Early Andalusian period on the basis of the associated

material culture.

Figure 2. Ancestry inference in ancient samples from North Africa and the Iberian Peninsula. (a) PCA

analysis, (b) ADMIXTURE analysis (K=2 - K=8), and (c) detailed ADMIXTURE analysis for European

Neolithic samples (K=8).

Figure 3. Outgroup f3-statistic for IAM (A), KEB (B) and TOR (C), and admixture f3-statistic for KEB

(D).

323