1

Comprehensive analysis of central carbon metabolism reveals multiple connections between nutrients, biosynthetic capacity, and cell morphology in *Escherichia coli*

Short title: Central carbon metabolism, cell growth, and cell morphology

Corey S. Westfall[1] and Petra Anne Levin[1]*

[1] Department of Biology, Washington University, St. Louis, Missouri, U.S.A.

* Corresponding author

E-mail: plevin@wustl.edu (PAL)

Abstract

Bacterial morphology is a complex trait that is highly sensitive to changes in the

environment. For heterotrophic organisms, such as *Escherichia coli*, increases in

nutrient levels are frequently accompanied by several-fold increases in both size

and growth rate. Despite the dramatic nature of these changes, how alterations in

nutrient availability translate into changes in growth and morphology remains a

largely open question. To understand the signaling networks coupling nutrient

availability with size and shape, we examined the impact of deletions in the entirety

of non-essential central carbon metabolic genes on *E. coli* growth rate and cell size.

Our data reveal the presence of multiple metabolic nodes that play important yet

distinctive roles in shaping the cell. Consistent with recent work from our lab and

others, although both are sensitive to nutrient availability, size and growth rate vary

independently. Cell width and length also appear to be independent phenomena,

influenced by different aspects of central carbon metabolism. These findings

highlight the diversity of factors that can impact cell morphology and provide a

foundation for further studies.

Author summary

Often taken for granted, the shape of bacterial cells is a complex trait that is highly

sensitive to environmental perturbations. Nutrients in particular, strongly impact

bacterial morphology together with growth rate. The ubiquitous, rod-shaped

bacteria *Escherichia coli* increases both length and width several fold upon a shift

from nutrient poor to nutrient rich medium, a change accompanied by an equally

dramatic increase in growth rate. Central carbon metabolism is an obvious site for

the integration of nutrient dependent signals that dictate cell size and shape. To

develop a clearer picture of the molecular mechanisms coupling nutrient

assimilation with cell growth and morphology, we screened the entirety of non-

essential carbon metabolic genes for their contribution to growth rate and cell

shape. Our data reveal the presence of multiple regulatory circuits coordinating

different metabolic pathways with specific aspects of cell growth and morphology.

Together, these data firmly establish a role for central carbon metabolism as an

environmentally sensitive sculptor of bacterial cells.

Introduction

The behavior and physiology of single celled organisms is at the mercy of

their environment. Nutrients, in particular, dramatically impact the growth rate, cell

cycle, and morphology of bacteria. *Salmonella typhimurium, Escherichia coli,* and

Bacillus subtilis cells cultured in nutrient rich conditions exhibit mass doubling

times up to 6-fold faster than their counterparts cultured in nutrient poor medium

[1-3]. These increases in growth rate are accompanied by similarly dramatic

increases in cell size. *E. coli* increases length (2-fold) and width (1.5-fold) for a ~3-

fold increase in total square area between nutrient poor and nutrient rich

conditions [3]. *B. subtilis* increases length ~3-fold in nutrient rich conditions while

width remains more or less constant [2,4].

The positive relationship between nutrient availability, growth rate and cell

size is generally referred to as the "Growth Law". Besides cell size, DNA, RNA, and

protein content of the cell also show a positive relationship with growth rate, with

faster growing cells containing more of these macromolecules [1]. Although there is

more total DNA/RNA/protein in larger, faster growing cells, their concentration

remains relatively constant due to the concomitant increase in cell size [1,5].

Despite its name, the relationship between growth rate and cell size is

complicated and there are several, well known exceptions to the Growth Law.

Reducing growth rate by lowering culture temperature does not impact cell length

or width, suggesting that it is nutrients rather than growth rate that determine cell

size [6]. Conversely, mutations that lead to modest delays in division frequently

result in increases in cell length without detectably impacting mass doubling time

[7-9].

Central carbon metabolism: a hub connecting nutrient availability with

growth rate and cell size

As the entry and processing point for the raw materials required for cell

growth and proliferation, central carbon metabolism (CCM) plays a key role in

coordinating nutrient availability with cell growth and size. Recent work indicates

that CCM impacts bacterial morphology in several ways: by its impact on cell cycle

progression [10-12]; by its impact on lipid synthesis [6]; and through its impact on

cell wall biogenesis [13].

CCM-mediated changes in cell cycle progression primarily affect cell length,

while changes in lipid synthesis and cell wall biogenesis impact both length and

width. In an example of the former, UDP-glucose, produced in two steps from

glucose-6-phosphate at the beginning of glycolysis, stimulates interactions between

the moonlighting glucosyltransferase OpgH in *E. coli* and the unrelated enzyme UgtP

in *B. subtilis* and the essential tubulin-like cell division protein FtsZ, delaying

division and increasing cell length during growth in carbon rich medium [4,12]. In

contrast, defects in fatty acid synthesis (FAS) lead to reductions in both length and

width, effectively recapitulating the impact of nutrient limitation on cell morphology

[6,14].

CCM can also effect morphology via its impact on the availability of cellular

building blocks, particularly those required for synthesis of the cell wall. In a recent

study, the Young laboratory determined that defects in *wecE*, a gene encoding an

aminotransferase important for making the *E. coli* outer membrane molecule

enterobacterial common antigen (ECA), led to the formation of swollen, filamentous

cells. ECA is transported to the cell wall via undecaprenyl-phosphate, the lipid

carrier responsible for shuttling precursors for peptidoglycan and other

components of the bacterial cell envelope across the plasma membrane. Careful

genetics revealed the morphological defects associated with defects in *wecE* were

due to sequestration of undecaprenyl-phosphate conjugated to an ECA intermediate,

interfering with peptidoglycan synthesis [15].

More generically, nutrient starvation leads to an accumulation of the small

molecule guanosine tetraphosphate (ppGpp) that interacts directly with RNA

polymerase, to inhibit transcription of a wide array of genes, effectively curtailing

flux through multiple biosynthetic pathways [16]. ppGpp concentration is inversely

proportional to both size and growth rate: ectopic expression of the ppGpp

synthetase RelA during growth in nutrient rich medium reduces both growth rate

and cell size [6,17]. The reduction in size is indirect and dependent on ppGpp-

mediated inhibition of fatty acid synthesis, while the reduction in growth rate is

almost certainly a consequence of ppGpp-dependent inhibition of a host of essential

processes [6].

While these examples highlight several ways in which CCM might impact the

positive relationship between cell growth and cell morphology, they represent only

a small fraction of the pathways that constitute CCM. To identify additional

connections between nutrient levels, metabolism, and cell size in *E. coli*, we

screened non-essential (CCM) gene deletions for defects in cell size and growth rate.

Our data highlight three CCM nodes—the first branch-point of glycolysis, the

pentose-phosphate pathway, and acetyl-CoA metabolism—as critical links between

nutrient availability, cell growth and cell size. While some aspects of CCM

bottlenecks appear to directly impact the relationship between nutrient availability

and biosynthetic flux, other aspects of CCM effect growth and morphology in more

nuanced ways.

Results

Multiple connections exist between CCM, growth rate, and cell size

To identify key steps of CCM that impact the growth and morphology of *E.*

coli, we phenotypically analyzed cells defective in 44 non-essential CCM genes (S1

Table) to. These experiments took advantage of the Keio Collection, an ordered

insertion-deletion library of all non-essential genes in *E. coli* [18] and Coli-Inspector,

an Image J plug in designed for high throughput analysis of bacterial morphology

[19]. Prior to analysis, the Keio insertion-deletions were transduced into the

laboratory wild type strain MG1655. Briefly, single colonies of each mutant were

picked and cultured under nutrient rich conditions (Luria-Bertani (LB) broth

supplemented with 0.2% glucose) to an OD_{600} of ~0.2. Cultures were then back-

diluted to OD_{600} of 0.01 and tracked for 4 generations until they reached a maximum

OD_{600} of 0.2 (the low OD_{600} ensures cells are all actively growing and at

approximately the same growth phase prior to analysis). Cells were sampled, fixed,

and imaged by phase contrast microscopy. Cell size (length and width) was

determined using Coli-Inspector in ImageJ.

Based on size and mass doubling time, we assigned the 44 mutants into six

classes (Table 1). Represented by two mutations, Class I cells were >10% smaller

than wild type cells with mild to undetectable changes in growth rate (within two

SD of wild type mass doubling time). Eight mutations fell into Class II: small (more

than 10% decrease in square area than wild type) with a significant decrease in

doubling time (>20% wild type). Class III was heterogenous, consisting of a

population dominated by smaller than normal cells and punctuated by the presence

of a few (5-10%) very long cells. Three mutants fell into Class IV long (>10% longer

than WT) and two into Class V, heterogenous with a wide distribution of cell lengths

and variable cell widths (Fig 1A). Surprisingly, the majority of mutants (26) were

essentially wild type with regard to growth rate and cell size (Class VI). This finding

suggests the presence of alternative pathways that permit *E. coli* to bypass

particular CCM enzymes and/or that nutrients that allow for bypass are present in

LB. We elected not to investigate the impact of deletions in either *ppsA*, the first step

of gluconeogenesis, or *fumC*, one of the three isoforms involved in the conversion of

fumarate to malate in the TCA cycle, on cell size as we were unable to obtain stable

MG1655 transductants of either mutant. Representative images of each mutant are

shown (Fig 2).

Table 1: CCM deletion mutants exhibiting significant alterations in growth rate

and/or morphology.

Gene Name	Doubling time (per minute)	Area (microns squared)	Pathway
Wild Type	22	5.1	
Class I (Small with doubling times <20% from WT)			
sucC	25	4.6	TCA Cycle
gnd	21	4.6	Pentose-Phosphate
Class II (Small and doubling times >20% from WT)			
crr	27	4.4	Glycolysis
aceE	34	2.9	Glycolysis/Acetate/Fatty acid/TCA
tktA	33	3.7	Pentose-Phosphate
pgl	26	4.5	Pentose-Phosphate
rpiA	28	3.8	Pentose-Phosphate
rpe	37	3.1	Pentose-Phosphate
pta	33	3	Acetate Metabolism
ackA	36	2.7	Acetate Metabolism
Class III (Short and Long)			
acnB	25	5.6	TCA Cycle
zwf	30	4.4	Pentose Phosphate
Class IV (Long)			
yggF	19	6.7	Putative Gluconeogensis
talB	23	5.8	Pentose-Phosphate
eda	22	6	Entner-Douderoff
Class V (heterogenous)			
tpiA	30	ND	Glycolysis
gpmM	21	ND	Glycolysis

Of the different classes, only the eight Class II mutants "obeyed" the Growth

Law, with reductions in growth rate accompanied by predictable reductions in cell

size (Fig 1B). The close correlation between growth rate and size for Class II

mutants suggests that the loss of individual genes and/or their product's catalytic

function, leads to a biosynthetic bottleneck that recapitulates the impact of nutrient

starvation on cellular physiology. In contrast, despite their heterogeneity with

regard to cell length, both Class I and Class IV mutants exhibited more or less

normal growth rates, suggesting a potential defect in the signal transduction

pathways coordinating CCM with cell growth and cell size. The heterogeneity of

Class III and Class V indicate a loss of proper regulation of cell cycle progression

and/or width control, leading to stochastic variations in cell morphology.

Growth in carbon poor medium largely ameliorates CCM mutant

morphological defects

To clarify the impact of nutrient flux on individual CCM mutant phenotypes,

we assessed the growth rate and size of Class I-V mutants in three different media:

LB + 0.2% glucose, the minimal media AB + 0.2% glucose, and AB + 0.2% succinate

representing high nutrient conditions, low nutrient glycolytic conditions, and low

nutrient gluconeogenic conditions respectively. Cells were measured using the

Matlab software package SuperSegger, which facilitates the rapid analysis of large

numbers of clumped cells [20]. In general, SuperSegger results corresponded almost

exactly with the Coli-Inspector data reported above, although cell areas were

smaller over all most likely due to a difference in how the two programs define cell

boundaries and at what point of the cell cycle that they "divide" the parent cell into

two daughter cells (see wild type area in Table 1 and S3 Table).

In light of data indicating that cells defective in *crr*, encoding the

phosphotransferase system subunit EIIA^{glc}, exhibited significant reductions in size

(Table 1), we also examined the nutrient dependent phenotype of cells defective for

adenylate cyclase or the cAMP receptor (Crp) production. Crr is part of a

phosphodependent transferase system that is essential for the import of glucose and

other sugars into *E. coli* [21]. Crr facilitates glucose import via transfer of a

phosphate group from Hpr to EIIB^{glc} and finally to glucose, generating glucose-6-

phosphate at the beginning of glycolysis (reviewed in [22]). During growth on a

non-glycolytic substrate, phosphorylated EIIA^{glc} accumulates and activates the

adenylate cyclase, CyaA, leading to synthesis of cyclic-AMP (cAMP) [23,24]. A co-

activator, cAMP binds to Crp facilitating DNA binding and stimulating transcription

of a broad swathe of genes including those involved in the transport and utilization

of non-glycolytic carbon sources [25,26]. If *crr* impacts cell size via its impact on

cAMP production and Crp activation, defects in both *cyaA* and *crp* should be

phenotypically identical to *crr* mutants with regard to cell size and growth rate. We

found that this is indeed true as deletions in *crr*, *cyaA*, or *crp* led to similar growth

rates (29, 29, and 28 minutes) and cell size (2.1 square micron area for all three

mutants). This implies that the growth rate and size phenotypes of *crr::kan* are

caused not by a lack of glucose import but instead by blocking cAMP synthesis.

217   Generally speaking, most of the CCM mutants retained a positive relationship

218 between nutrient availability and cell size with the exception of the three cAMP

219 mutants, *crr*::*kan*, *cyaA*::*kan*, and *crp*::*kan* (Fig 3). During growth in carbon poor

220 and/or gluconeogenic conditions, the lengths of Class I and Class II mutants

221 approached that of wild type, particularly in AB-succinate medium, and the

222 heterogeneous phenotype exhibited by mutants in Class III and Class IV was largely

223 relieved. Width also decreased, although more modestly. The average width of the

224 mutants ranged from 0.68-0.86 microns during growth in LB-glu and 0.48-0.60

225 microns during growth in AB-suc. Consistent with catabolite repression serving to

226 inhibit division during growth on a gluconeogenic carbon source, the *crp⬚kan*

227 mutants (dotted line in Fig 3B, 3C, and 3D) did not show any change in size when

228 switched from LB-glu to AB-glu. Both *crr*::*kan* and *cyaA*::*kan* (magenta and gray in

229 Fig 3) were similar to *crp*::*kan*, showing only minor, insignificant changes in length,

230 with no change in width. Unfortunately, *crr*, *cyaA*, and *crp* are essential under

231 gluconeogenic conditions, preventing analysis of these mutants in AB-succinate.

232   Although we were able to obtain cell size data for *rpiA*, *crr*, *rpe*, *aceE*, and

233 *sucA* in carbon rich (LB-glu) and carbon poor (AB-glu) conditions, all five mutants

234 were unable to grow in the gluconeogenic condition, AB + 0.2% succinate. At first

235 glance this finding appears to be in conflict with data indicating that *rpiA*, *rpe*, *aceE*,

236 and *sucA* are capable of growth in another carbon poor gluconeogenic condition: M9

237 + 0.1% glycerol [27]. However, this discrepancy likely reflects the ability of glycerol

238 to enter the CCM partway through glycolysis—as dihydroxyacetone phosphate—

while succinate enters through the TCA cycle, leading to dependence on different

pathways through CCM to synthesize all needed carbon metabolites.

Defects in CCM differentially impact assembly of the cell division machinery

To gain insight into the relationship between growth rate, cell cycle

progression and CCM during growth in LB-glucose, we assessed the ability of

different mutants to suppress the heat sensitivity associated with the GTP binding

defective FtsZ mutant, *ftsZ84* (G105S) [28]. While *ftsZ84* supports both FtsZ

assembly and division under permissive conditions (LB at 30°C), under non-

permissive conditions (LB with no salt at 42°C), FtsZ assembly and division are

blocked, leading to filamentation and cell death [7,29]. Based on extensive work

from our group and others, conditions or mutations that enhance FtsZ assembly are

expected to suppress the heat sensitivity of temperature sensitive *ftsZ* alleles

[4,17,30,31]. Changes in the availability of CCM products or components that

normally inhibit FtsZ assembly should restore growth and colony formation to

ftsZ84 mutants under non-permissive conditions while defects in CCM-related

factors that normally promote FtsZ assembly will enhance *ftsZ84* heat sensitivity.

Double mutants in CCM deletions and *ftsZ84* were generated to examine the impact

of these mutants on FtsZ assembly.

Consistent with a positive impact on FtsZ assembly, the majority of Class I-V

mutants promoted growth of *ftsZ84* cells under mildly restrictive conditions (37°C,

LB no salt) (Fig 4A). The strongest suppressors were loss-of-function mutations in

aceE or *ackA* both of which increased *ftsZ84* cell viability 10,000-fold during growth

at 37°C on salt-free LB medium. Mild suppression was observed in loss of function

mutations in *crr*, *tktA*, and *pta* which increased *ftsZ84* cell viability 1,000-fold, 100-

fold, and 100-fold respectively. The *zwf* deletion was alone in leading to

enhancement of *ftsZ84*, i.e. lack of growth at 30° on LB-no salt (Fig 4B). Combined

with the short and long phenotype of *zwf⬚kan*, it appears that loss of *zwf* leads to

impairment of FtsZ activity. Significantly, we did not observe a consistent

correlation between reduced growth rate and suppression of *ftsZ84* heat-sensitivity

among the various CCM mutants.

As part of these experiments we also revisited the impact of a loss of function

mutation in *pgm*, encoding phosphoglucomutase—the enzyme that converts

glucose-6-phosphate to glucose-1-phosphate. In previous work, we had reported

that deletions in *pgm* suppress *ftsZ84* heat sensitivity [12]. However, we were

unable to detect any suppression of *ftsZ84* by *pgm* in this instance. Resequencing

BH173, the strain employed in the original study revealed acquisition of a reversion

mutation that restored *ftsZ84* sequence to wild type.

To gain a better understanding of the mechanisms coordinating central

carbon metabolism with cell morphology, we undertook an in depth analysis of two

distinctive phenotypes: the reduction in length and width observed in cells defective

in catabolite repression and the link between acetate metabolism, cell growth and

cell cycle progression revealed by defects in *aceE, ptaA, and ackA* (Fig 7). The

results of these efforts are described below.

Catabolite-dependent regulation of FtsZ assembly and function

Reasoning that if the product of *crr,* EIIA^{glc}, was impacting division indirectly via its role in stimulation of adenylate cyclase and subsequent activation of the Crp transcription factor, we hypothesized that 1) defects in either the adenylate cyclase (encoded by *cya*) or *crp* should restore growth to *ftsZ84* cells under non-permissive conditions, and 2) the addition of extracellular cAMP (which is able to diffuse through the membrane [32]) to *crr::kan ftsZ84* cells should bypass *crr::kan* mediated suppression of *ftsZ84* heat sensitivity but only in the presence of Crp. Consistent with this model, deletion of either *cya* or *crp* phenocopied *crr▢kan* mediated *ftsZ84* suppression at 37°C on salt free medium (Fig 5). More directly, 5 mM cAMP was sufficient to restore heat sensitivity to *crr::kan ftsZ84* cells as well as cells defective in adenylate cyclase (*cyaA*) but only in the presence of *crp* (Fig 5). Together, these data strongly argue for the presence of at least one inhibitor of FtsZ assembly within the extensive Crp regulon.

cAMP controls cell width through regulation of the morphogene *bolA*

In contrast to wild type cells, which decrease in both length and width upon a shift from glycolytic (low cAMP) to gluconeogenic (high cAMP) conditions, the *crp* deletion showed no significant changes in width when changing media from LB-glu to AB-glu, suggesting a potential role for cAMP in cell width regulation (Fig 3). To explore this idea, we examined the impact of extracellular cAMP on wild type (MG1655) cells cultured in both glycolytic and gluconeogenic conditions.

305    Consistent with a negative impact on cell width, 5 mM exogenous cAMP was

306  sufficient to decrease wild type cell width in both LB and AB + 0.2% glucose (0.96 to

307  0.81 microns and 0.73 to 0.68 microns respectively) (Fig 6A). The addition of cAMP

308  also significantly decreased cell length (3.8 to 3.2 microns for LB and 2.7 to 2.5

309  microns for AB-Glu) and growth rate (2.7 to 2.3 doubling per hour for LB and 0.98 to

310  0.88 doublings per hour for AB-Glu) (Fig 6B). Significantly, cAMP-dependent

311  reductions in size are independent of growth rate. Reducing levels of exogenous

312  cAMP from 5 mM to 1 mM decreased the growth rate of *cyaA*::*kan* (which is unable

313  to synthesize any cAMP) in gluconeogenic conditions (AB + 0.2% succinate) from

314  0.87 to 0.79 doublings per hour, while cell width increased from 0.65 to 0.75

315  microns (Fig 6C). Although width was independent of growth rate, length was not as

316  the *cyaA*::*kan* mutant length decreased with lower cAMP from 2.6 microns in 5 mM

317  cAMP to 2.2 microns in 1 mM cAMP in a manner similar to the growth rate (Fig 6D).

318    cAMP has been implicated in the Crp dependent activation of the

319  transcription factor OmpR , which inhibits expression of a host of genes, including

320  several whose products have morphogenic activity [33]. Of particular interest is

321  *bolA*, encoding a transcription factor that inhibits transcription of *mreB* [34], which

322  encodes a key component of the elongation machinery [35]. Reductions in MreB

323  activity lead to a block in synthesis of lateral cell wall material and the formation of

324  wide and rounded cells [36,37]. *bolA* transcription has previously been shown to be

325  inhibited by addition of extracellular cAMP [38].

326    Together, genetic interactions between *crp, ompR,* and *bolA,* support a model

327  in which cAMP increases cell width indirectly, via down regulation of OmpR and the

concomitant increase in BolA expression leading to repression of MreB mediated

cell wall synthesis. Consistent with this model, cells defective in *ompR* and *bolA*

were refractile to the addition of exogenous cAMP with regard to cell width (Figs 6E

and 6F).

Defects in acetate metabolism support links between fatty acid synthesis and

assembly of the cell division machinery

Cell cycle progression is tightly correlated with growth rate in wild-type

cells. Fast growing cells are not only larger but contain a higher frequency of FtsZ

rings (Z-rings) with approximately 90% of *E. coli* cells containing visible Z-rings in

LB-glucose. Meanwhile, slow growing cells contain fewer Z-rings (~50%). Z-ring

frequency shows a roughly linear relationship to growth rate when grown in

different media (Fig 4C dotted line) [12,39]. Taking advantage of a green fluorescent

protein (GFP)-FtsZ fusion that allows us to label the FtsZ ring without detectably

altering size or growth [6], we assessed the frequency of FtsZ ring formation in the

17 mutants in Classes I through V.

Although the Z-ring frequency and growth rate were correlated in the

majority of CCM mutants, several exhibited a disconnect between these two

parameters, specifically those associated with acetate metabolism (black squares in

Fig 4C). Cells defective in *aceE*, the E1 protein of pyruvate dehydrogenase that

converts pyruvate to acetyl-CoA, exhibited an extremely high (86%) Z-ring

frequency relative to wild type cells cultured in medium of the same growth rate

(~75%) [12], while both *ackA* and *pta*, involved in converting acetyl-CoA to acetate,

exhibited lower Z-ring frequencies relative to growth rate (69% and 67%

respectively). Assembly of the cell division machinery is a multistep process

involving first, recruitment of FtsZ and other so-called early proteins to the nascent

division site followed by a brief delay and assembly of the late proteins, which

include enzymes required for synthesis of the septal cell wall (reviewed in [40]). In

E. coli, division is coincident with recruitment of FtsN, the last protein to be

incorporated into the division machinery and the putative "trigger" for cross wall

synthesis [41,42]. The two-step nature of division machinery assembly suggests two

potential explanations for the elevated Z-ring frequency in *aceE*::*kan* mutants: 1)

premature recruitment of FtsZ to the septum or 2) delays in late protein

recruitment and activation of cytokinesis. Conversely, the lower frequency of FtsZ

rings in *ackA*::*kan* and *pta*::*kan* mutants could be a consequence of either 1) delays

in FtsZ recruitment or 2) premature late protein recruitment and/or activation of

division mediated cell wall synthesis.

To clarify the underlying cause of the elevated Z-ring frequency in *aceE*::*kan*

cells and depressed Z ring frequency in the *ackA*::*kan* and *pta*::*kan* backgrounds we

took advantage of a heterologous, IPTG-inducible FtsN-GFP [43] to assess the

efficiency of FtsN recruitment in all three mutants. Growth in LB-glu showed 32% of

wild-type cells with FtsN-GFP localization at mid-cell (Fig 7A). In contrast, despite

varying Z-ring frequencies, all three mutants exhibit more or less the same

frequency of FtsN ring formation (24%, 23%, and 26% for *aceE*::*kan*, *ackA*::*kan*, and

pta::*kan* respectively) with all three being significantly less than wild type. The

consistency of FtsN ring frequency, despite differences in Z-ring frequency between

the three CCM mutants, suggests that it is delayed recruitment of the late division

proteins and/or activation of cytokinesis that is responsible for the high Z-ring

frequency in *aceE*::*kan* cells while the lower Z-ring frequeny in *ackA*::*kan* and

pta::*kan* mutants would appear to be a consequence of a delay in FtsZ assembly.

Acetyl-CoA metabolism plays a critical role in the ability of cells to synthesize

the wide variety of fatty acids and lipids that constitute the cell envelope [44]. *aceE*,

in particular, encodes a subunit of the pyruvate dehydrogenase complex responsible

for aerobic synthesis of the majority of acetyl-CoA, a necessary precursor in the first

step of FAS [45]. Defects in *aceE* are thus expected to reduce flux through FAS, and

based on previous work, shift the balance in lipid composition [46]. Although not

directly involved in FAS, defects in *pta* and *ackA* may lead to a build-up of acetyl-CoA

and potentially also shift flux through FAS.

Based on previous studies linking lipid composition to cell morphology [47],

we speculated defects in acetate metabolism might result in changes in plasma

membrane composition that interfere with assembly and activity of the cell division

machinery. To test this idea, we measured the Z-ring frequency of wild type and

mutant cells cultured in the presence of sub-inhibitory concentrations of the fatty

acid synthesis inhibitor triclosan (50 ng/mL). This concentration is too low to

impact growth rate but has a modest positive impact on cell size [6]. Triclosan

inhibits FabI, the enzyme that catalyzes the reduction of enoyl-ACP during the

elongation cycle of fatty acid synthesis [48]. Although not a perfect recapitulation of

a block upstream of acetyl-CoA synthesis, as it blocks a downstream step, both

triclosan treatment and the *aceE* mutations should similarly curtail flux through

FAS.

In support of a model in which modest reductions in flux through FAS

negatively impact assembly of the cell division machinery, Z-ring frequency was

significantly increased in wild type cells as well as in *aceE::kan, pta kan*, and

ackA kan cells during growth in sub-MIC (minimum inhibitory concentration), 50

ng/mL triclosan. (Z-ring frequency: wild type 88% +T/95% -T; *pta::kan* 56% -T/

79% + T; *ackA::kan* 60% -T/74% +T; *aceE kan* 79% -T to 87% +T) (Fig 7B). *pta* or

ackA deletion mutants were slightly more sensitive to triclosan with regard to Z-ring

frequency than either *aceE* or wild type cells, likely a reflection of the already high

frequency of FtsZ rings in these latter strains.

Further supporting a positive connection between FAS and assembly of the

cell division machinery, 50 ng/mL triclosan enhanced the heat sensitive phenotype

of conditional mutations in several cell division genes including *ftsA* (*ftsA27*-an

early, actin-like protein), *ftsK* (*ftsK44*—a late division protein involved in

chromosome partitioning), *ftsQ* (*ftsQ1*—a late division protein believed to serve

primarily as a scaffold), and *ftsI* (*ftsI23*—a late division protein with a trans-

peptidase domain) [49-52]. Somewhat surprisingly given its impact on FtsZ ring

frequency (Fig 7B), triclosan had no impact on the heat-sensitivity of *ftsZ84* (Fig 7C).

We speculate that the differential impact of triclosan on *ftsZ84* relative to the other

conditional mutants is a consequence of their subcellular location (cytoplasmic –

FtsZ) versus membrane associated (FtsA) and integrated into the membrane (FtsQ,

FtsK, and FtsI) [40].

Discussion

Our systematic analysis of the relationship between central carbon

metabolism, cell growth, and cell size both reinforces the Growth Law as a general

principal (i.e. growth rate and size are positively correlated) but also calls into

question many of the fundamental assumptions that underlie it. While the largest

class of mutants with observable phenotypes—Class II: small with reduced growth

rates, essentially "obeyed" the law so to speak, growth rate and size varied

independently in the other classes of CCM mutants (Fig 1B). Class II mutants closely

resemble cells cultured in nutrient-poor conditions with regard to growth rate and

size, suggesting these cells are experiencing a near global reduction in biosynthetic

capacity. In contrast, growth rate and morphology are independent of one another

in the other nine CCM mutants (Table 1), consistent with defects in specific

regulatory or structural functions that directly impact cell length and/or width.

These results are in line with accumulating work indicating that growth rate

and cell size are independent phenomena. Modest defects in cell division are well

known to increase bacterial cell size—particularly length—without impacting

growth rate [4,7,50,51,53], while defects in transcription severely curtail growth but

have little detectable impact on morphology [6]. Most importantly, an exhaustive

phenotypic analysis of the entirety of non-essential genes in *E. coli* by the Jacobs-

Wagner group, failed to identify any evidence supporting a direct connection

between growth rate and size [54].

Overflow metabolism and cell size

In a common, yet paradoxical, fermentative mechanism known as "overflow

metabolism", *E. coli* convert up to 50% of available glucose into acetate during

aerobic growth in nutrient rich conditions [55] . Acetate fermentation is inefficient,

resulting in ∼14 fewer ATP per glucose than oxidative phosphorylation and, unlike

other fermentative pathways, does not regenerate NAD^+ from NADH [56]. While it is

unclear why bacteria perform overflow metabolism, it is potentially a more

proteome-efficient means of moving carbon through CCM, ensuring that glycolysis

proceeds at a maximum pace when nutrients are plentiful.

Supporting the idea that overflow metabolism is critical for maximizing

glycolytic flux, defects in *pta*, *ackA*, and *aceE*, encoding the first two enzymes in

acetate fermentation and a subunit of pyruvate dehydrogenase respectively, are

dramatically impaired in both growth rate and cell size. This is consistent with a

global reduction in biosynthetic capacity. Whether this reduction is the result of

lower ATP production or the consequence of a bottleneck in flux through critical

biosynthetic pathways is unclear. Regardless, we suspect that reductions in ATP

levels and/or biosynthesis indirectly result in accumulation of the alarmone ppGpp

[57], providing an explanation for the observed suppression of *ftsZ84* heat-

sensitivity in *pta, ackA,* and *aceE* deletion strains. Increases in ppGpp are well

known to suppress the conditional nature of *ftsZ84,* although the precise mechanism

has yet to be determined [17,58] .

cAMP is a key player in maintaining the correct cellular aspect ratio

A somewhat surprising finding is the identification of cAMP as a contributor

to nutrient dependent reduction in both cell length and width. Accumulating under

gluconeogenic conditions, cAMP binds to the transcriptional activator Crp and

regulates expression of more the than 280 genes [59]. Although the precise

member(s) of the cAMP-Crp regulon responsible remains elusive, our data indicate

that cAMP-Crp function together to reduce width under nutrient poor conditions via

their impact on the morphogenic factor BolA and cell length via its impact on the

assembly of FtsZ (Figs 5 and 6). In this regard, cAMP differs from metabolic

signaling molecule, ppGpp, which genetics indicate is a positive regulator of division

(Fig 8).

CCM is robust but not necessarily efficient

Perhaps the biggest surprise of this study is our finding that the defects in the

majority of non-essential CCM genes had little to no detectable impact on either cell

size or growth rate. While the lack of phenotype in many cases is likely a

consequence of our screening conditions, nutrient-rich, complex medium

supplemented with glucose, in certain cases the robust growth we observed is

difficult to explain. For example, we anticipated that defects in phosphoglucose

isomerase (encoded by *pgi*), the first dedicated step in glycolysis would dramatically

impact growth rate and size by virtue of its impact on not only glycolysis, but also

other associated pathways leading to reductions in ATP synthesis and overall

biosynthetic capacity [60]. Instead, we were unable to distinguish *pgi* mutants from

wild type cells during growth LB-glucose (Table S2). Based on our identification of

seven mutants in the pentose-phosphate pathway, we hypothesize that despite the

ubiquitous nature of glycolysis in nutrient- rich medium, glucose is predominantly

driven through the pentose-phosphate pathway to generate essential building

blocks.

493   Similarly, we observed little in the way of a detectable phenotype associated

with defects in the gene encoding citrate synthase *gltA*. Essential for the complete

oxidation of carbon, GltA synthesizes citrate from oxaloacetate and acetyl-CoA at the

entry point into the TCA cycle [61]. Despite its seeming importance, previous work

indicates that *E. coli* represses *gltA* transcription in the presence of glucose,

suggesting that it can forgo complete carbon oxidation under nutrient rich

conditions [59,62]. It may be that under nutrient rich condition, *E. coli* maximizes

rapid growth over the efficient use of available carbon, a proposition that is in line

with our data indicating that defects in acetate metabolism significantly curtail

growth in nutrient rich medium (Table 1).

Concluding thoughts

505   While this screen demonstrates the ability of systematic phenotypic analysis

to identify new connections between CCM, cell growth and cell size, many questions

remain unanswered. In addition to the large number of insertion mutants with no

observable phenotype, defects in 17 genes resulted in dramatic changes in growth

rate and cell morphology. Of these, we were able to identify the proximal cause in

24% (*crr*, *aceE*, *pta*, *ackA*; Figs 5-7). A better understanding of the remaining 13

mutations will likely require a combination of genetics and metabolomics to tease

apart the root cause of their morphological and growth related defects.

Additionally, although this effort focused specifically on carbon metabolism,

the availability of other nutrients—particularly nitrogen, phosphorous, sulfur and

iron— have also been implicated in growth and morphology. Assessing the

phenotype of the CCM insertion collection under conditions in which these factors

are limiting is thus also worthwhile. Finally, *E. coli* is believed to spend much of its

time under microaerobic or anaerobic conditions. Understanding how oxygen levels

impact growth, cell cycle progression, and cell morphology thus remains a

tantalizingly open question.

Materials and Methods

Strains and media

All chemicals, media components, and antibiotics were purchased from

Sigma-Aldrich (St. Louis, MO). *E. coli* strains were constructed through standard P1

vir transductions [63] moving kanamycin marked insertion deletions from Keio

collection strains [18] purchased from the Coli Genomic Stock Center into MG1655

(F⁻ λ⁻ *ilvG*- *rfb*-50 *rph*-1), our "wild type" parental strain. Cell were cultured in Luria-

Bertani (LB) broth supplemented with 0.2% glucose or AB defined media [64]

supplemented with either 0.2% glucose or 0.2% succinate as the sole carbon source.

Unless otherwise noted (e.g. for heat sensitive conditional alleles of cell division

genes including *ftsZ84*, *ftsA27*, *ftsK44*, *ftsQ1*, and *ftsI23*), cells were cultured at 37° C

and sampled for analysis at OD_{600} between 0.1 and 0.2. For the primary analysis cells

were cultured in 5 mL LB + 0.2% glucose and aerated at 200 rpm. For measuring the

effect of different media, cells were cultured in 200 μL of specified media in a 96-

well plate and aerated at 567 rpm in a BioTek© Eon© plate reader (Fig 3).

Microscopy

Cells were sampled directly from cultures and fixed in 16%

paraformaldehyde (Electron Microscopy Sciences) as described [6]. Fixed cells were

mounted on agarose pads and visualized using a 100x phase-fluor objective on a

Nikon© Ti-E inverted microscope. To facilitate visualization of FtsZ rings, 1 mM IPTG

was added to cultures of cells encoding a chromosomal P_{IPTG}-*ftsZ-GFP* [65].

Expression of a chromosomal P_{IPTG}-*ftsN-GFP* was induced with 50 μM IPTG [43]. GFP

fluorescence was detected with a C-FL GFP filter cube with 470/40nm excitation

and 525/50nm emission (Chroma Technology Corporation) Nikon Elements

software (Nikon Instruments, Inc.) was used for image capture and analysis. >200

cells were counted per experimental replicate and at least three independent

replicates were used for each data point.

Cell Size Measurements

Cell size from phase-contrast images was determined with one of two freely

available segmentation software packages: the ImageJ© plug-in Coli Inspector [19]

(Fig 1, Tables 1 and S2) or SuperSegger, a MatLab© based application [20] (Figs 3-7,

Table S3). As Coli Inspector does not determine area directly, we estimated it as

length x width. SuperSegger directly determines the area of individual cells. We

determined the area of ≥100 cells for each experiment and/or condition. Unless

otherwise noted, average size was determined from 3 independent experiments.

Dot plating efficiency assays

Heat-sensitive cell division alleles, *ftsZ84*, *ftsA27*, *ftsK44*, *ftsQ1*, and *ftsI23*

were cultured in LB at 30°C degrees to an OD_{600} of ~0.5. 1 mL of cells were then

pelleted, washed once with LB-no salt, and re-suspended to an OD_{600} of 0.1 in LB-no

salt. 10 μL of serial dilutions between 10^{-1} to 10^{-6} were plated under permissive

conditions (LB at 30°C) or non-permissive conditions (LB-no salt at 37°C). Due to

the strains extreme sensitivity, LB-no salt at 30°C was non-permissive for *zwf::kan*

ftsZ84 double mutants. For cAMP or triclosan addition experiments, 5 mM cAMP or

50 ng/mL triclosan was added directly to LB agar prior to pouring the plates. Each

plating efficiency experiment was repeated three times with representative images

shown.

Acknowledgements

We would like to thank the Coli Genetic Stock Center (CGSC) for providing

numerous strains quickly. We'd like to thank Norbert Vischer for aid in using Coli

Inspector and Stella Stylianidou and Paul Wiggins for help with SuperSegger. We

would like to thank Elizabeth Mueller for work on the *pgm ftsZ84* double mutant.

We would also like to thank Elizabeth Mueller, Dr. Ashley Sherp, and the Hani Zaher

laboratory for critical feedback on this work.

References

References

1.   Schaechter M, Maaløe O, Kjeldgaard NO. Dependency on medium and
temperature of cell size and chemical composition during balanced grown of
Salmonella typhimurium. J Gen Microbiol. 1958;19: 592–606.

2.   Sargent MG. Control of cell length in Bacillus subtilis. J Bacteriol. 1975;123: 7–
19.

3.   Donachie WD, Begg KJ. Cell length, nucleoid separation, and cell division of
rod-shaped and spherical cells ofEschericia coli. J Bacteriol. 1989;171: 4633–
4639.

4.   Weart RB, Lee AH, Chien A-C, Haeusser DP, Hill NS, Levin PA. A metabolic
sensor governing cell size in bacteria. Cell. 2007;130: 335–347.
doi:10.1016/j.cell.2007.05.043

5.   Donachie WD. Relationship between cell size and time of initiation of DNA
replication. Nature. 1968;219: 1077–9.

6.   Vadia S, Jessica LT, Lucena R, Yang Z, Kellogg DR. Fatty Acid Availability Sets
Cell Envelope Capacity and Dictates Microbial Cell Size. Current Biology. 2017.

7.   Arjes HA, Lai B, Emelue E, Steinbach A, Levin PA. Mutations in the bacterial
cell division protein FtsZ highlight the role of GTP binding and longitudinal
subunit interactions in assembly and function. BMC Microbiol. 2015;15: 209.
doi:10.1186/s12866-015-0544-z

8.   Coltharp C, Buss J, Plumer TM, Xiao J. Defining the rate-limiting processes of
bacterial cytokinesis. Proc Natl Acad Sci USA. 2016;113: E1044–53.
doi:10.1073/pnas.1514296113

9.   Zheng H, Ho P-Y, Jiang M, Bin Tang, Liu W, Li D, et al. Interrogating the
Escherichia coli cell cycle by cell dimension perturbations. P Natl Acad Sci
USA. 2016;113: 15000–15005.

10.   Tymecka-Mulik J, Boss L, Maciąg-Dorszyńska M, Rodrigues JFM, Gaffke L,
Wosinski A, et al. Suppression of the Escherichia coli dnaA46 mutation by
changes in the activities of the pyruvate-acetate node links DNA replication
regulation to central carbon metabolism. Korolev S, editor. PLoS ONE. Public
Library of Science; 2017;12: e0176050. doi:10.1371/journal.pone.0176050

11.   Monahan LG, Hajduk IV, Blaber SP, Charles IG, Harry EJ. Coordinating
bacterial cell division with nutrient availability: a role for glycolysis. MBio.
2014;5. doi:10.1128/mBio.00935-14

12.   Hill NS, Buske PJ, Shi Y, Levin PA. A Moonlighting Enzyme Links Escherichia

coli Cell Size with Central Metabolism. PLoS Genet. Public Library of Science;
2013;9: e1003663. doi:10.1371/journal.pgen.1003663

13.   Irnov I, Wang Z, Jannetty ND, Bustamante JA, Rhee KY, Jacobs-Wagner C.
Crosstalk between the tricarboxylic acid cycle and peptidoglycan synthesis in
Caulobacter crescentus through the homeostatic control of α-ketoglutarate.
Casadesús J, editor. PLoS Genet. Public Library of Science; 2017;13: e1006978.
doi:10.1371/journal.pgen.1006978

14.   Yao Z, Davis RM, Kishony R, Kahne D, Ruiz N. Regulation of cell size in
response to nutrient availability by fatty acid biosynthesis in Escherichia coli.
Proc Natl Acad Sci USA. 2012;109: E2561–8. doi:10.1073/pnas.1209742109

15.   Jorgenson MA, Kannan S, Laubacher ME, Young KD. Dead-end intermediates
in the enterobacterial common antigen pathway induce morphological defects
in Escherichia coli by competing for undecaprenyl phosphate. Mol Microbiol.
2015;100: 1–14. doi:10.1111/mmi.13284

16.   Magnusson LU, Farewell A, Nyström T. ppGpp: a global regulator in
Escherichia coli. Trends in Microbiology. 2005.

17.   Powell BS, Court DL. Control of ftsZ expression, cell division, and glutamine
metabolism in Luria-Bertani medium by the alarmone ppGpp in Escherichia
coli. J Bacteriol. 1998;180: 1053–1062.

18.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio
collection. Mol Syst Biol. EMBO Press; 2006;2: 2006.0008.
doi:10.1038/msb4100050

19.   Vischer NOE, Verheul J, Postma M, van den Berg van Saparoea B, Galli E,
Natale P, et al. Cell age dependent concentration of Escherichia coli divisome
proteins analyzed with ImageJ and ObjectJ. Front Microbiol. 2015;6: 586.
doi:10.3389/fmicb.2015.00586

20.   Stylianidou S, Brennan C, Nissen SB, Kuwada NJ, Wiggins PA. SuperSegger:
robust image segmentation, analysis and lineage tracking of bacterial cells.
Mol Microbiol. 2016;102: 690–700. doi:10.1111/mmi.13486

21.   Gabor E, Göhler A-K, Kosfeld A, Staab A, Kremling A, Jahreis K. The
phosphoenolpyruvate-dependent glucose-phosphotransferase system from
Escherichia coli K-12 as the center of a network regulating carbohydrate flux
in the cell. Eur J Cell Biol. 2011;90: 711–720. doi:10.1016/j.ejcb.2011.04.002

22.   Deutscher J, Francke C, Postma PW. How phosphotransferase system-related
protein phosphorylation regulates carbohydrate metabolism in bacteria.
Microbiol Mol Biol Rev. American Society for Microbiology; 2006;70: 939–

1031. doi:10.1128/MMBR.00024-06

23.   Saier MH, Feucht BU, Hofstadter LJ. Regulation of carbohydrate uptake and
adenylate cyclase activity mediated by the enzymes II of the
phosphoenolpyruvate: sugar phosphotransferase system in Escherichia coli. J
Biol Chem. American Society for Biochemistry and Molecular Biology;
1976;251: 883–892.

24.   Park Y-H, Lee BR, Seok Y-J, Peterkofsky A. In vitro reconstitution of catabolite
repression in Escherichia coli. J Biol Chem. American Society for Biochemistry
and Molecular Biology; 2006;281: 6448–6454. doi:10.1074/jbc.M512672200

25.   Zubay G, Schwartz D, Beckwith J. Mechanism of activation of catabolite-
sensitive genes: a positive control system. P Natl Acad Sci USA. National
Academy of Sciences; 1970;66: 104–110.

26.   Grainger DC, Hurd D, Harrison M, Holdstock J, Busby SJW. Studies of the
distribution of Escherichia coli cAMP-receptor protein and RNA polymerase
along the E. coli chromosome. P Natl Acad Sci USA. National Acad Sciences;
2005;102: 17693–17698. doi:10.1073/pnas.0506687102

27.   Joyce AR, Reed JL, White A, Edwards R, Osterman A, Baba T, et al.
Experimental and computational assessment of conditionally essential genes
in Escherichia coli. J Bacteriol. American Society for Microbiology; 2006;188:
8259–8271. doi:10.1128/JB.00740-06

28.   Ricard M, Hirota Y. Process of cellular division in Escherichia coli:
physiological study on thermosensitive mutants defective in cell division. J
Bacteriol. 1973;116: 314–322.

29.   Addinall SG, Lutkenhaus CJ. Temperature shift experiments with anftsZ84(Ts)
strain reveal rapid dynamics of FtsZ localization and indicate that the Z ring is
required throughout septation and cannot reoccupy division sites once
constriction has initiated. J Bacteriol. 1997;179: 4277–4284.

30.   Levin PA, Kurtser IG, Grossman AD. Identification and characterization of a
negative regulator of FtsZ ring formation in Bacillus subtilis. P Natl Acad Sci
USA. 1999;96: 9642–9647.

31.   Gervais FG, Drapeau GR. Identification, cloning, and characterization of rcsF, a
new regulator gene for exopolysaccharide synthesis that suppresses the
division mutation ftsZ84 in Escherichia coli K-12. J Bacteriol. 1992;174: 8016–
8022.

32.   Jackson DW, Simecka JW, Romeo T. Catabolite repression of Escherichia coli
biofilm formation. J Bacteriol. American Society for Microbiology (ASM);
2002;184: 3406–3410. doi:10.1128/JB.184.12.3406-3410.2002

33.   Huang L, Tsui P, Freundlich M. Positive and negative control of ompB
transcription in Escherichia coli by cyclic AMP and the cyclic AMP receptor
protein. J Bacteriol. American Society for Microbiology (ASM); 1992;174: 664–
670.

34.   Yamamoto K, Nagura R, Tanabe H, Fujita N, Ishihama A, Utsumi R. Negative
regulation of the bolA1p of Escherichia coli K-12 by the transcription factor
OmpR for osmolarity response genes. FEMS Microbiol Lett. 2000;186: 257–
262.

35.   Doi M, Wachi M, Ishino F, Tomioka S, Ito M, Sakagami Y, et al. Determinations
of the DNA sequence of the mreB gene and of the gene products of the mre
region that function in formation of the rod shape of Escherichia coli cells. J
Bacteriol. American Society for Microbiology; 1988;170: 4619–4624.
doi:10.1128/jb.170.10.4619-4624.1988

36.   Kruse T, Bork-Jensen J, Gerdes K. The morphogenetic MreBCD proteins of
Escherichia coli form an essential membrane-bound complex. Mol Microbiol.
Blackwell Publishing Ltd; 2005;55: 78–89. doi:10.1111/j.1365-
2958.2004.04367.x

37.   Divakaruni AV, Baida C, White CL, Gober JW. The cell shape proteins MreB and
MreC control cell morphogenesis by positioning cell wall synthetic complexes.
Mol Microbiol. 2007 ed. 2007;66: 174–188. doi:10.1111/j.1365-
2958.2007.05910.x

38.   Lange R, Hengge-Aronis R. Growth phase-regulated expression of bolA and
morphology of stationary-phase Escherichia coli cells are controlled by the
novel sigma factor sigma S. J Bacteriol. 1991;173: 4474–4481.

39.   Weart RB, Levin PA. Growth rate-dependent regulation of medial FtsZ ring
formation. J Bacteriol. 2003;185: 2826–2834.

40.   Margolin W. FtsZ and the division of prokaryotic cells and organelles. Nat Rev
Mol Cell Biol. 2005;6: 862–871.

41.   Lutkenhaus J. FtsN--trigger for septation. J Bacteriol. 2009;191: 7381–7382.
doi:10.1128/JB.01100-09

42.   Daley DO, Skoglund U, Söderström B. FtsZ does not initiate membrane
constriction at the onset of division. Sci Rep. 2016;6: 33138.
doi:10.1038/srep33138

43.   Gerding MA, Liu B, Bendezú FO, Hale CA, Bernhardt TG, de Boer PAJ. Self-
enhanced accumulation of FtsN at Division Sites and Roles for Other Proteins
with a SPOR domain (DamX, DedD, and RlpA) in Escherichia coli cell
constriction. J Bacteriol. 2009;191: 7383–7401. doi:10.1128/JB.00811-09

44.   Parsons JB, Rock CO. Bacterial lipids: Metabolism and membrane homeostasis.
727   Progress in Lipid Research. 2013;52: 249–276.
728   doi:10.1016/j.plipres.2013.02.002

729   45.   Li M, Ho PY, Yao S, Shimizu K. Effect of lpdA gene knockout on the metabolism
730   in Escherichia coli based on enzyme activities, intracellular metabolite
731   concentrations and metabolic flux analysis by …. Journal of biotechnology.
732   2006.

733   46.   Buttke TM, Ingram L. Inhibition of unsaturated fatty acid synthesis in
734   Escherichia coli by the antibiotic cerulenin. Biochemistry. 1978.

735   47.   Rowlett VW, Mallampalli VKPS, Karlstaedt A, Dowhan W, Taegtmeyer H,
736   Margolin W, et al. Impact of Membrane Phospholipid Alterations in
737   Escherichia coli on Cellular Function and Bacterial Stress Adaptation.
738   jbasmorg.

739   48.   McMurry LM, Oethinger M, Levy SB. Triclosan targets lipid synthesis. Nature.
740   1998;394: 531–532. doi:10.1038/28970

741   49.   Herricks JR, Nguyen D, Margolin W. A thermosensitive defect in the ATP
742   binding pocket of FtsA can be suppressed by allosteric changes in the dimer
743   interface. Mol Microbiol. 2014;94: 713–727. doi:10.1111/mmi.12790

744   50.   Begg KJ, Dewar SJ, Donachie WD. A new Escherichia coli cell division gene,
745   ftsK. J Bacteriol. American Society for Microbiology; 1995;177: 6211–6222.
746   doi:10.1128/jb.177.21.6211-6222.1995

747   51.   Storts DR, Aparicio OM, Schoemaker JM, Markovitz A. Overproduction and
748   identification of the ftsQ gene product, an essential cell division protein in
749   Escherichia coli K-12. J Bacteriol. American Society for Microbiology (ASM);
750   1989;171: 4290–4297.

751   52.   Begg KJ, Donachie WD. Cell shape and division in Escherichia coli:
752   experiments with shape and division mutants. J Bacteriol. 1985;163: 615–
753   622.

754   53.   Labie C, Bouche F, Bouche JP. Isolation and mapping of Escherichia coli
755   mutations conferring resistance to division inhibition protein DicB. J
756   Bacteriol. 1989;171: 4315–4319.

757   54.   Campos M, Dobihal GS, Jacobs-Wagner C. Genome-wide phenotypic analysis of
758   growth, cell morphogenesis and cell cycle events in Escherichia coli. bioRxiv.
759   Cold Spring Harbor Labs Journals; 2017;: 101832. doi:10.1101/101832

760   55.   Aristidou AA, San KY, Bennett GN. Metabolic engineering of Escherichia coli to
761   enhance recombinant protein production through acetate reduction.

Biotechnol Prog. American Chemical Society; 1995;11: 475–478.
doi:10.1021/bp00034a019

56.   Basan M, Hui S, Okano H, Zhang Z, Shen Y. Overflow metabolism in Escherichia
coli results from efficient proteome allocation. Nature. Nature Publishing
Group; 2015;528: 99–104.

57.   Liu K, Bittner AN, Wang JD. Diversity in (p)ppGpp metabolism and effectors.
Curr Opin Microbiol. 2015;24C: 72–79. doi:10.1016/j.mib.2015.01.012

58.   Navarro F, Robin A, D'Ari R, Joseleau-Petit D. Analysis of the effect of ppGpp
on the ftsQAZ operon in Escherichia coli. Mol Microbiol. 1998;29: 815–823.

59.   Zheng D, Constantinidou C, Hobman JL, Minchin SD. Identification of the CRP
regulon using in vitro and in vivo transcriptional profiling. Nucleic Acids Res.
2004;32: 5874–5893. doi:10.1093/nar/gkh908

60.   Adhya S, Schwartz M. Phosphoglucomutase mutants of Escherichia coli K-12. J
Bacteriol. 1971;108: 621–626.

61.   Spencer ME, Guest JR. Molecular cloning of four tricarboxylic acid cyclic genes
of Escherichia coli. J Bacteriol. American Society for Microbiology (ASM);
1982;151: 542–552.

62.   Walsh K, Koshland DE. Characterization of rate-controlling steps in vivo by
use of an adjustable expression vector. P Natl Acad Sci USA. National Acad
Sciences; 1985;82: 3577–3581.

63.   Lennox ES. Transduction of linked genetic characters of the host by
bacteriophage P1. Virology. 1955;1: 190–206.

64.   Clark DJ, Maaløe O. DNA replication and the division cycle in Escherichia coli. J
Mol Biol. 1967.

65.   Hale CA, de Boer PA. Recruitment of ZipA to the septal ring of Escherichia coli
is dependent on FtsZ and independent of FtsA. J Bacteriol. 1999;181: 167–
176.

 Figure Legends

 Figure 1: The effect of central carbon metabolic gene deletions on cell growth

 and size. (A) Mutant phenotypes mapped onto central carbon metabolic pathways.

 Arrows are colored based on the phenotype as follows: black= normal growth and

 size; green = normal growth and small size (<10% smaller than WT); red = slow

 growth (>20% increase in doubling time) and small size; orange = normal growth

 and long cells (>10% larger than WT); blue = short and long cells; purple = highly

 variable size; and dotted lines = essential genes. Brown arrows correspond to

 deletions that obtained suppressor mutations rapidly and were not analyzed. (B)

 Relationship between cell area and growth rate, as doublings per hour. Icons are

 colored as in panel (A).

 Figure 2: Representative images of all CCM mutants exhibiting morphology

 distinct from the wild type parental strain. Merged images of morphologically

 distinct CCM mutant strains expressing an *ftsZ-GFP* fusion to permit visualization of

 the cytokinetic ring. Scale bar = 5 microns and is the same for all images.

 Figure 3: CCM deletions exhibit medium-dependent effects on growth and

 morphology. (A) Cell size versus growth rate in LB-glucose (black diamonds), AB-

 glucose (orange triangles), and AB-succinate (blue circles). Length (B), width (C),

 and area (D) are plotted for each mutant in the three different media. Dotted line =

 crp::*kan* in LB-glucose and AB-glucose.

Figure 4: CCM mutations differentially impact FtsZ localization and activity.

(A) Dot-plating efficiencies of the CCM deletion *ftsZ84* double mutants cultured

under permissive (LB, 30°C) and non-permissive (LB-no salt, 37°C) conditions. (B)

Dot-plating efficiency of *zwf*::*kan ftsZ84* double mutant. Note the reduction in

zwf::*kan ftsZ84* viability under intermediate growth conditions (30°C on LB-no salt).

(C) FtsZ ring frequency versus growth rate for the CCM mutants. For reference we

have included previously reported data on the relationship between nutrient

availability and FtsZ ring frequency (dotted black line) [12]. Mutants are grouped

by metabolic pathway: black squares = acetyl-CoA production and acetate

fermentation; green triangles = pentose-phosphate shunt; purple diamonds = TCA

cycle; orange circle = Entner-Doudoroff pathway; blue cross = cAMP synthesis; and

red plus = gluconeogenesis.

Figure 5: Loss of cAMP signaling suppresses *ftsZ84* heat-sensitivity.

Representative dot-plating efficiencies of *ftsZ84* alone or in combination with

insertion deletions in cAMP signaling genes *crr*, *cyaA*, and *crp*. Cells were plated on

and grown at permissive (LB, 30°C), non-permissive (LB-no salt, 37°C), and non-

permissive containing cAMP (LB-no salt + 5mM cAMP, 37°C) conditions.

Figure 6: cAMP signaling reduces cell width via the morphogenic transcription

factor, *bolA*. Wild-type MG1655 cell size was measured with and without

exogenous 5 mM cAMP in either LB or AB-glucose. Width is shown in (A) and length

in (B). Width (C) and length (D) of *cyaA*::*kan* grown in AB-succinate with different

levels of exogenous cAMP shows that width is not correlated with growth rate but

length is. (E) Deletions in *crp*, *ompR*, or *bolA* abolish the cAMP-dependent cell

reductions in cell width. (F) Proposed genetic pathway linking cAMP to cell width.

Figure 7: Partial inhibition of fatty acid synthesis negatively impacts assembly

and activity of the late division proteins. (A). FtsN septal localization was

measured in MG1655, *aceE*::*kan*, *ackA*::*kan*, and *pta*::*kan* expressing an *ftsN-GFP*

from an IPTG inducible promoter   (B.) FtsZ-ring frequency was measured MG1655,

aceE::*kan*, *ackA*::*kan*, and *pta*::*kan* cells expressing *ftsZ-GFP* from an IPTG-inducible

promoter. Cells were cultured with (gray) or without (black) 50 ng/mL triclosan. (C)

Representative plating efficiencies of wild-type cells and the heat-sensitive cell

division mutants *ftsZ84*, *ftsA27*, *ftsK44*, and *ftsQ1* cultured on LB at 37°C with or

without triclosan. *ftsI23* was plated on LB-no salt at 37°C with or without triclosan.

Figure 8: Proposed pathway linking nutrient levels to cell length and width

through the signaling nucleotides cAMP and ppGpp.

Supporting Information

Table S1. List of bacterial strains used in this study

Table S2. Raw size and growth rate data for all central carbon metabolic gene

deletions.

Table S3. Raw data for figure 3 (CCM mutant growth in different media).

Table S4. Raw data for figure 4C (Z-ring frequency of CCM mutants).