Identification of environmental and genetic factors important for *Cryptococcus*

***neoformans* titan cell formation using new *in vitro* inducing conditions**

Benjamin Hommel,[1,2,8,9] Liliane Mukaremera,[3,*] Radames J. B. Cordero,[4,*] Christopher A.

Desjardins,[5,*] Aude Sturny-Leclère,[1,2,*] John Perfect,[6] James A. Fraser,[7] Arturo Casadevall,[4]

Christina A. Cuomo,[5] Françoise Dromer,[1,2] Kirsten Nielsen,[3] Alexandre Alanio[1,2,8,9]

[*] These authors participated equally to the work

[1] Institut Pasteur, Molecular Mycology Unit, Paris, France

[2] CNRS URA3012, Paris, France

[3] Department of Microbiology and Immunology, University of Minnesota, Minneapolis,

Minnesota, USA

[4] Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School

of Public Health, Baltimore, MD.

[5] Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

[6] Division of Infectious Diseases, Department of Medicine, Duke University Medical Center,

Durham, North Carolina, USA

[7] Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane,

Queensland, Australia

[8] Laboratoire de Parasitologie-Mycologie, Hôpital Saint-Louis, Groupe Hospitalier

Lariboisière, Saint-Louis, Fernand Widal, Assistance Publique-Hôpitaux de Paris (AP-HP),

Paris, France

[9] Université Paris Diderot, Sorbonne Paris Cité, Paris, France

Corresponding Author: Alexandre Alanio, Molecular Mycology unit, Institut Pasteur, 25 rue

du Dr Roux 75724 Paris Cedex 15; email: alexandre.alanio@pasteur.fr; Tel: +33140613255;

Fax: +33145688420

Abstract

A remarkable aspect of the human fungal pathogen *Cryptococcus neoformans* is the morphological changes triggered by the interaction with the host. During infection, a specific morphological change in cell size is observed, particularly in lung tissue, from regular 5 to 7 µm cells (RCs) to titan cells (TCs, > 10 µm and up to 100 µm). However, the stable and reproducible generation of large quantity of TCs was only possible during experimental infection. We implemented here *in vitro* conditions allowing TCs generation with the aim to understand the ancestry of TC, the environmental determinants of TCs formation and perform easily phenotype/genotype analysis to uncover genetic factors underlying TCs formation. This paper reports robust *in vitro* conditions that generates yeasts cells harboring the main characteristics of TCs: large (>10 µm), uninucleate and polyploid cells harboring a single large vacuole with a dense capsule surrounding a thickened melanized cell wall. We observed that TCs formation begins as soon as 8 hours after induction, with a maximal proportion of TCs obtained after 24 hours. TCs derived for the initial oldest cells (mother cells) rather than from daughter cells. Hypoxia, nutrient starvation and low pH stresses were the main factors that trigger TCs formation, while quorum sensing factors like the initial inoculum concentration and the addition of pantothenic acid also impacted TCs formation. Antifungal drugs, resulting in inhibition of ergosterol (fluconazole), or proteins and nucleic acids (flucytosine), altered TCs formation, as did hosts-related products such as serum, phospholipids and anti-capsular antibodies in our settings. We then explored genetic factors important for TCs formation using two approaches. Using strains from H99 lineage strains among which few genetic differences have been described, we showed that TCs formation was dependent on Lmp1, Sgf29 and Gpr4/5-Rim101 proteins. Then, based on a the analysis of natural Pkr1 loss-of-function clinical

isolates and serial clinical isolates, we observed the important role of Pkr1 protein, a negative

regulator of the cyclic adenosine monophosphate / protein kinase A (cAMP/PKA), for TCs

formation. These strains also emphasize the structural and functional relationship between

Pkr1 and Pka.

Our results provide new insights into TCs biogenesis with identification of multiple important

factors/pathways involved. The implementation of such standardized and robust *in vitro*

conditions pave the way for future research focusing on the genetic basis of TCs biogenesis,

biology of TCs and the ontology of morphological changes in *Cryptococcus neoformans*.

Author Summary:

Cryptococcus neoformans is a yeast that is capable of morphological change upon

interaction with host. Particularly, in the lung of infected mice, a subpopulation of yeast

evolves toward gigantism (titan cells size from 10 to 100 μm) that include other characteristics

such as thickened cell wall, dense capsule, polyploidization, large vacuole with peripheral

nucleus and cellular organelles. The generation of large number of such cells outside the lung

of mice have been described but was not reproducible nor standardized. We here report

standardized, reproducible, robust conditions of generation of titan cells and explored the

environmental and genetic factors underlying the genesis of these cells. We showed that Titan

cells were generated upon stresses such as change in the medium, nutrient deprivation,

hypoxia and low pH. Using collection of well characterized reference strains and clinical

isolates, we validated with our model the AMPc/PKA/Rim pathway as the major genetic

determinant of titan cell formation. This study opens the way for a more comprehensive

picture of the ontology of morphological changes in *Cryptococcus neoformans*.

Introduction

The ubiquitous environmental yeast *Cryptococcus neoformans* (Cn) is a basidiomycetous yeast that causes up to 223,000 incident cases of meningoencephalitis and more than 181,000 deaths per year worldwide [1], occurring mostly in immunocompromised individuals with acquired immunodeficiency syndrome (AIDS) [2]. The natural history of most of the cases of this invasive fungal infection proceeds through 3 stages: (i) primary infection via inhalation of desiccated yeasts or basidiospores, with development of sub-clinical pneumonia and spontaneous resolution via granuloma formation; (ii) latency of dormant yeast cells, as demonstrated epidemiologically [3] and biologically [4] and (iii) reactivation and dissemination upon immunosuppression, with meningoencephalitis as the main clinical presentation [5]. From environment to interaction with hosts, the yeasts experience drastic changes that require a capacity to rapidly adapt and survive in host tissues and cause disease [4,6–9]. In hosts, Cn is exposed to various stresses including high temperature, nutrient deprivation, low pH, hypoxia and high levels of free radicals [10].

In response to the host environment, morphological changes are required to survive and cause disease [11]. Specifically, Cn alter its morphology and produce enlarged cells referred to as giant or "titan cells" [12,13]. This phenomenon has been observed in animal and insect models of cryptococcosis, as well as in human lung and brain infections [12–16]. The titan cells (TCs) have increased cell body size, ranging from 10 μm to up to 100 μm in diameter [13,17–21] as compared to the 5-7 μm size of regular cells (RCs). Studies exploring TCs biology have revealed that these are: (i) uninucleate polyploid cells harboring a (ii) large single vacuole surrounded by a (iii) thick cell wall and (iv) a dense and highly crosslinked capsule [17]. TCs also exhibit increased resistance to various stresses including phagocytosis [21], oxidative and

nitrosative stress [20,21], and resistance to the antifungal drug fluconazole [20]. Importantly, TCs production also enhances dissemination, survival and virulence in a mouse model of infection [19]. TCs formation is known to be regulated by the G-protein coupled receptors Gpr5 and Ste3a, that signal through the Gα subunit protein Gpa1 to trigger the cyclic adenosine monophosphate / protein kinase A (cAMP/PKA) signaling pathway [15,20,22–25]. The cAMP/PKA pathway is critical for regulation of other virulence factors in Cn, including capsule formation [22,23], notably through its action on the ubiquitin-proteasome pathway [26]. Pka1 is known to be tuned by a regulatory protein Pkr1, and *pkr1Δ* mutant strains exhibit enlarged capsule [22]. Further studies show TCs formation is increased by high *PKA1* expression or low Pkr1 activity and is decreased by low *PKA1* expression [6]. Downstream of the PKA pathway, Rim101, a major transcription factor that again controls production of many virulence factors, is also necessary for TCs production [18].

To date, studies of TCs formation have been hindered by an inability to consistently and reproducibly generate large quantities of TCs *in vitro*. Although several methods have been reported for inducing large cell *in vitro* there have been persistent problems in easily and consistently implementing these protocols across laboratories [17], presumably because the variables that contribute to inducing conditions are not well understood.

In this study, we developed robust *in vitro* conditions that generates enlarged cells with many of the *in vivo* TCs characteristics and used this protocol to explore environmental and genetic factors involved in TCs formation. The genetic determinants of TCs formation have been investigated through a genotype-phenotype correlation study based on the H99 laboratory reference strains and clinical isolates and validated by adequate KO mutants.

Results

Titan cells generated *in vitro* had similar characteristics to *in vivo* titan cells

While growth in minimal medium (MM) using standard growth conditions has no effect

on cell size, we identified growth conditions that stimulate production of enlarged yeasts cells

and optimized this experimental protocol, referred to as the Titan Cell Generating (or TCG)

protocol, using our H99O reference strain (Fig 1). Observation of these *in vitro*-generated large

cells by microscopy shows many characteristics of TCs including increased cell body size

(diameter >10 μm), thicker refractive cell wall, large central vacuole, and peripheral cell

cytoplasm distribution (Fig 2A). The TCG protocol was reproducible, with H99O reference

strains generating equivalent proportions of TCs in three different laboratories throughout

the world (Fig 2B). Specifically, the proportion of TCs was 39.4% [interquartile range (IQR),

34.1-40.7] in lab 1, 21.0% [12.5-26.8] in lab 2 and 29.9% [23.1-48.5] in lab 3.

TCs differ from regular cells (RCs) in various characteristics including capsule size

melanization, cell wall structure and DNA content. TCs exhibit pronounced melanization,

measured as blackness on the pictures, compared to RCs (Fig 2C, S1 Fig), with a median of the

(max - mean grey intensity per pixel) of 20065 [18785-21887] in TCs vs. 13067 [9660-15998]

in RCs (p<0.0001). TCs exhibit a slight but significant increase in capsule size compared to RCs

(median 3.4 μm in large cells vs. 3.1 μm in RC, p<0.001, Fig 2D). Capsule structure was analyzed

based on binding pattern (fluorescence intensity and distribution) of four monoclonal

antibodies specific for capsular polysaccharides [27] (S2 Fig), and no pronounced difference

between the capsule structures of TCs and RCs was quantified although some differences

could be observed. Chitin content was significantly increased in TCs compared to RCs as

determined by calcofluor white (CFW) fluorescence intensity (fluorescence intensity/pixel/cell

87.9 [71.7-107.7] vs. 66.5 [51.7-79.4], respectively, p<0.0001, Fig2E, S2 Fig). Finally, chitin

levels assessed by N-acetylglucosamine content determined by a biochemical method were

also higher in TCs (156.3 mM/g [153.7-203.7]) than in RCs (97.7 [87.2-119.3]), (p<0.001, Fig2F).

Cell sorting based on CFW staining and fluorescence microscopy allowed us to validate that

cells exhibiting the higher CFW intensity were TCs (FSC/SSC^{high}, S3 Fig).

To analyze DNA content, yeasts obtained at the end of the TCG protocol were stained

with propidium iodide (PI), with fluorescence in TCs (FSC/SSC^{high}) and RCs (FSC/SSC^{low})

compared to haploid (H99O) and diploid (AD7-77) strains grown in Sabouraud medium. DNA

content was higher in TCs (FSC/SSC^{high}) than RCs (FSC/SSS^{low}) cells (Fig 2G), with an increase in

the number of TCs cells with PI fluorescence > 2C (Fig 2H). In contrast, the RCs cells had the

same PI fluorescence pattern as the haploid H99O cells. Similarly, a reverse gating strategy

based on PI intensity shows yeasts with the highest PI intensity were large TCs cells (red

arrows, S4 Fig).

TCs develop from older cells and produce regular daughter cells

To examine temporal changes in cell size with the TCG protocol, we measured yeast

cell sizes at 0, 4, 8, 16, 24 and 120 h using automated analysis with the Icy software. This

automated analysis correlated with manual size measurements (Interclass correlation=0.99)

and TC/RCs classification (Kappa test=0.81±0.07). The median cell size increased during the

first 24 h of incubation, starting at 5.7 μm [5.4-6.0] and increasing to 9.7 μm [8.4-11.0] (Fig

3A). The first TCs cells were observed at 8 h with a progressive increase in the proportion of

TCs overtime, reaching a plateau by 24 h (Fig 3B).

Temporal changes in cell morphology where determined using live cell imaging and

light microscopy. Cells with the large vacuole characteristic of TCs appeared between 4 and 8

hours (Fig 3C, white arrow). Live cell imaging over 18 h (Supplemental material 1) showed that:

(i) TCs swelled from an older (parental) RCs and (ii) TCs produced RCs daughter cells. Calcofluor

white (CFW) staining was transferred to daughter cells upon division [4,28] and was used to

further monitor the ancestry of TCs and RCs cell populations over time using flow cytometry

(Fig 3D). The initial CFW stained population (0 h) consisted of RCs (FSC/SSC^{low}) cells with high

CFW fluorescence intensity. At 24 h, TCs (FSC/SSC^{high}) had high CFW fluorescence intensity,

indicating these cells were in the original CFW stained population. The RCs (FSC/SSC^{low}) cells

were composed of two populations: i) cells with high CFW intensities from the original stained

population and ii) daughter cells with low CFW intensity (Fig 3D left panel) indicating these

were new daughter cells. Direct comparison of cell size (FSC) and CFW intensity at 24 h (Figure

3D, right panel) showed the FSC^{high} cells (TCs) had a higher CFW fluorescence intensity (black

arrow) suggesting that they represent the older progenitor mother cells while the H24 FSC^{low}

cells (RCs) had lower CFW fluorescence intensity suggesting these were newly formed

daughter cells (white arrow). These combined data show that the TCs cells derived from the

initial inoculated cells and produced regular-sized RCs cells upon cell division.

Generation of TCs *in vitro* is influenced by environmental conditions

We tested several parameters affecting steps 2 and 3 of the TCG protocol described in

Fig 1 and identified changes that significantly influence TCs generation as measured by cell

size distribution and TCs proportion (p<0.0001). The first parameter we tested was the growth

medium and transition between different growth mediums (Fig 4A). Initial culture in YPD (step

2) and transfer to MM (step 3) had the highest median cell size at 9.1 µm [6.9-11.1]. Culture

from MM to MM produced fewer TCs, 25.5% (118/463) vs. 39.5%(182/461), although the TCs

tended to be larger with diameters over 20 µm. Light exposure during step 3 also increased

both median cell size and proportion of TCs (Fig 4B). In the light, median cell size was 9.4 µm

[7.3-11.4] vs. 8.4 µm [7.1-9.9] in the dark, with a TCs proportion at 41.6% (983/2362) vs. 24.3%

(1072/4399), respectively (Fig 4B). Incubation temperature at 30°C at step 3 increased cell size

distribution compared to 37°C (9.4 µm [7.3-11.4] vs. 7.0 µm [6.2-8.2]) as well as the proportion

of TCs (41.6% (983/2362) vs. 8.1% (297/3652)) (Fig 4C). The pH of the MM at step 3 also

influenced cell size (Fig 4D), with pH=5.5 producing significantly larger cells that (9.1 µm [6.9-

11.2] and 38.6% of TCs), compared to either lower pH (pH=4: median 5.1 µm [4.4-5.8] (0%

TCs)) or higher pH (pH=7: 8.2 µm [7.2-9.4] (16.4% TCs), or pH=8.5: 6.9µm [5.9-7.9] (0.7% TCs))

(Fig 4D). Finally, hypoxia at step 3 also increased median cell size compared to normoxia (7.5

µm [5.9-9.7]), with chemically induced hypoxia yielding higher median cell sizes compared to

physically induced hypoxia] (10.1 µm [7.8-12.5] vs. 8.9 µm [7.3-10.9, p<0.0001) (Fig 4E). The

proportion of TCs in normoxia (14.5% (732/5050) was lower than in chemically induced

hypoxia or physically induced hypoxia (63.0% (732/1161) and 38.6% (1264/3004),

respectively) (p<0.0001).

Generation of TCs *in vitro* is influenced by fungal and host derived cues

Previous studies of titan cell formation in a murine pulmonary infection model showed

that inoculum concentration can impact titan cell production [12,13]. To explore this

phenomenon, we examined cell size changes in response to different initial concentrations of

cells at step 3 (Fig 5A). Inoculated cell concentration significantly impacted the median cell

size of the yeast population (p<0.0001), with the highest median cell size observed at

10^6cells/mL (9.2 [7.3-11.1]) compared to 10^5 cells/mL (6.3 [5.1-8.3]), 10^4 cells/mL (6.2 [5.1-

7.9]) and 10^7 cells/mL (5.9 [5.2-6.5]) (Fig 5A). Similarly, the proportion of TCs generated was

significantly higher at 10^6 cells/mL (37.6% (896/2382) compared to 10.3% (279/2716) at 10^4

cells/mL, 14.3% (346/2420) at 10^5 cells/mL and 0% at 10^7 cells/mL (0/2177), p<0.0001.

Previous studies reported vitamin B5 (pantothenic acid) is involved in quorum sensing *C.*

neoformans [29]. The addition of vitamin B5 to the TCG protocol had no effect on median cell

size (8.35 µm [6.9-10.2] and 8.3 µm [6.4-10.8], p=0.8011, Fig 5B), but significantly increased

the proportion of TCs (37.9% (1435/3785) vs. 26.9% (983/3650)). Recent studies also implicate

the role of the small Qsp1 peptide by *C. neoformans* cells in quorum sensing [30]. Addition of

Qsp1 peptide significantly decreased median cell size from 9.1 µm [7.1-11.2] to 8.5 µm [6.9-

10.1] (Fig 5C) and TCs proportion from 38.4% (1075/2798) to 26.6% (915/3439), p<0.0001.

We then tested hosts factors that could interact *in vivo* with yeasts cells in the lung such as

anticapsular antibodies, serum and phosphatidylcholine, which both have already been shown

to help producing TCs *in vitro* [13,18,31]. Monoclonal antibodies co-incubated at step 3 that

bind to different epitopes of the capsule inhibited TCs generation, with a decreased cell size

of 7.2 µm [6.1-8.3] for E1 mAb, and 6.8 µm [5.9-7.7] for 18B7 mAb compared to the untreated

control (8.9 µm [7.1-10.6]) (Fig 5D), with a significantly smaller proportion of TCs (3.2%

(44/1360) with 18B7, 5.3% (67/1273) with E1 compared to 33.1% (327/987) for the control).

The addition of fetal calf serum (FCS) significantly decreased median cell size (7.3 µm [6.4-8.1]

vs. (9.1 µm [7.1-11.1], Fig 5E) and the proportion of TCs (2.5% (99/3911) vs. 38.2%

(1234/3228), p<0.0001) compared to control, as did the addition of phosphatidylcholine (PC)

(8.0 µm [7.0-9.0] vs. 9.0 µm [7.1-11.2] for the median cell size, (Fig 5F) and the proportion of

TCs (14.7% (344/2340) vs. 38.2% (1077/2820), p<0.0001).

To understand if the alteration of the metabolism induced by antifungal drugs affected TCs

formation, we tested the effect of co-incubation of fluconazole (inhibitor of ergosterol

synthesis) and flucytosine (inhibitor of nucleic acids formation and transcription) at step 3.

Fluconazole (FLC) exposure during step 3 resulted in significantly smaller median cell sizes

compared to control (7.1 µm [6.2-8.3]) at 1mg/L (data not shown), 6.8 µm [5.9-7.8] at 2 mg/L

(data not shown), and 6.5 µm [5.6-7.2] at 4 mg/L vs. 9.4 µm [7.5-11.3] in the control, Fig 5G)

and a significantly smaller proportion of TCs (5.9% (124/2073) at 1 mg/L, 2.9% (55/1919) at 2

mg/L and 1.0% (19/1877) at 4 mg/L vs. 40.9% (877/2146) in the control, $p < 0.0001$). Flucytosine

exposure also significantly decreased the mean cell size at all concentrations tested (6.5 µm

[5.9-6.9] at 1 mg/L, 6.5 µm [6.1-6.9] at 2.5 mg/L, and 6.4 µm [5.9-6.8] at 5 mg/L compared to

control (8.3 µm [6.8-10.3], Fig 5H) with no TCs observed upon flucytosine exposure. Of note,

the viability of the cells recovered at step 4 after 5 days of drug exposure was unchanged for

fluconazole but reduced for flucytosine ($p < 0.0001$ compared to unexposed, S5 Fig). We then

were interested to test if iterative subcultures with or without the presence of active

molecules (CFW or fluconazole) would affect TCs formation, assuming that the cell wall and

the global metabolism of the sub-cultured progeny would be impaired in the presence of high

concentration of cell wall toxic drug (CFW) or fluconazole, respectively. So, we analyzed the

impact of repeated sub-culture of the cells prior to step 1 on TCs cell production (Fig 5I). Sub-

culture on Sabouraud agar 8 times (8Sub) spanning a one-month period decreased

significantly the median cell size (8.60 µm [7.02-10.07]) compared to the initial culture (0Sub)

(9.17 [6.99-10.90]) ($p < 0.0001$). Addition of CFW to induce cell wall stress during sub-cultures

(8Sub+CFW) had no significant impact on median cell size (8.03 µm [6.82-9.46]) compared to

the 8Sub control. However, iterative exposure to fluconazole during sub-culture increased

significantly median cell size to 10.15 µm [8.04-13.23] (8Sub+FLC) compared to the 8Sub

control ($p < 0.001$, Fig 5I).

In vitro TCs generation is dependent upon genetic background and involves functional

abolition of the _LMP1_ or _SGF29_ genes.

Previous whole genome sequencing studies identified single-nucleotide

polymorphisms (SNPs) and insertions/deletions (indels) between the H99 strains recovered

from various laboratories (Table 1) [32]. To determine whether any of these SNPs or indels

affect TCs generation, we tested the H99S, H99W, H99 CMO18, H99L, KN99α strains in the

TCG assay. H99O produced more TCs than the other H99-derived strains, p<0.0001 (Fig 6A, S6

Fig). These H99 derivative strains were also tested for their titan cell formation in the lungs of

infected mice (Fig 6B). As with *in vitro* TCs production, all the H99 derivative strains showed

lower levels of titan cell formation in the mouse model when compared to H99O (p<0.0001),

with the exception of KN99α that had equivalent titan cell production as H99O (Fig 6B).

Two genes, *LMP1* and *SGF29*, are dramatically affected by SNPs/indels in the H99

derivatives; *LMP1* has a frameshift deletion (H99W and H99 CMO18) and *SGF29* is deleted

(KN99 and H99L). To determine if these genes are involved in TCs production, we analyzed

lmp1Δ and *sgf29Δ* deletion mutants for *in vitro* TCs and *in vivo* titan cell formation (Fig 6C and

Fig 6D, respectively). The *sgf29Δ* mutant in the H99O background manifested halved TCs

generation from 8.1% (49/600) to 4.2% (25/600) (p<0.0001). The same mutant in H99S

manifested no TCs generation. We observed similar results for *lmp1Δ* with no TCs formation

in the *lmp1Δ* mutant. Complementation of *lmp1Δ* using the *lmp1Δ:LMP1* H99S strain restored

TCs generation to that found in H99S (Fig 6C). Importantly, the same trend was observed for

in vivo titan cell formation. The *sgf29Δ* mutant in H99O background reduced titan cell

formation from 18.8% (113/600) to 9.3% (37/400) (p<0.0001). The *lmp1Δ* H99S mutant had

only 3.5% (21/600) titan cell formation compared to 14% for H99S (84/600), p<0.0001, and

this decrease in titan cell formation was restored in the *lmp1Δ:LMP1* H99S strains (9.5%

(57/600)) (Fig 6D).

***In vitro* TCs generation requires signaling through the Gpr/PKA/Rim101 pathway in both**

laboratory reference strains and clinical isolates.

The signal transduction pathway Gpr/PKA/Rim101 is known to regulate TCs formation *in vivo*

[18]. Briefly, the G coupled protein receptor 5 and Ste3a signals through Gpa1 to trigger

cAMP/PKA with downstream the Rim101 transcription factor. This pathway regulates

virulence factors such as capsule or melanin [33,34].

To determine if this same pathway was similarly critical for TCs generation *in vitro*, we

examined cell enlargement in the *gpr4Δ*, *gpr5Δ*, *gpr4Δ/gpr5Δ, rim101Δ* mutants and their

complemented strains in both the H99 and KN99α genetic backgrounds (Fig 7A, S6E Fig).

Rim101 function was similar to that observed *in vivo*, with little TCs generation in the rim101Δ

mutant (1.9% (51/2600)) and full restoration of TCs generation in the complemented strain

(9.2% (239/2600) as compared to the H99O control (11.0% (176/1600) (p<0.0001). Similar to

in vivo titan cell production, the *gpr4Δ/gpr5Δ* had no TCs generation, but surprisingly both of

the single *gpr4Δ* and *gpr5Δ* mutants also lacked TCs generation. Unlike *in vivo* titan cell

production, TCs generation in the *gpr4Δ/gpr5Δ* double mutant was not rescued by GPR5 alone

(Fig 7A). Taken together, these data suggest that signaling through both Gpr4 and Gpr5 via

Rim101 is required for TCs generation *in vitro*.

To determine whether the Gpr/PKA/Rim101 pathway can impact TCs generation in

clinical isolates, we also screened a total of 56 clinical isolates for their ability to produce TCs.

Two isolates (AD2-06a and AD2-02a) produced an increased proportion of TCs relative to

H99O (ratio of clinical strain/H99O of 2.78 and 1.62, respectively). Three additional isolates

(AD4-37a, AD1-95a, AD4-43a) were able to produce TCs but less than H99O (ratio=0.40, 0.23,

0.12, respectively). The proportion of TCs recorded for the other clinical isolates was close to

zero (ratio between 0.1 and 0.01% for five, less than 0.01% for six, and no TCs at all for the remaining 39) (Fig 7B).

The complete genome sequence was obtained for 29 of the 56 clinical isolates and a phylogenetic tree of these strains and the H99O reference strain shows high genetic diversity (Fig 7C, Table 2). Compared to H99O, the high TCs generating strain AD2-06a harbored 31,229 SNPs and was closely related to AD3-55a (31,171 SNPs) and AD3-41a (28,599 SNPs), which were both unable to produce TCs. A total of 19 genes were considered disrupted in AD2-06a and not in AD3-55a and AD3-41a (S2 Table). Duplication of chromosomal regions have been observed in the French sequenced clinical isolates (S3 Table) with AD2-06a harboring a large duplication of chromosome 9 (Chr9, region 465 – 665kb). To assess whether the Chr9 duplicated region could be responsible for TCs formation, we explored a larger collection of *C. neoformans* strains with complete genome sequence [35] and discovered five additional clinical isolates (Ug2459, CCTP20, FFV14, WM-148, WM-626) harboring partial duplications on chr9 duplication (Fig S7). We analyzed TCs generation in these 5 isolates, but only the Ug2459 strain generated TCs *in vitro* and in a low proportion (S7 Fig).

AD2-06a was isolated from the initial cerebrospinal fluid (CSF) sample of an HIV-infected patient at diagnosis of cryptococcal meningitis. Another isolate, AD2-07, was recovered from the CSF of the same patient after 13 days of amphotericin B treatment. Thus, AD2-06a and AD2-07 are closely related with only 137 different SNPs and 40 indels. AD3-55a is in the same clade and also very similar with 370 different SNPs and 64 indels (Fig 7C, 7D). In contrast, these three isolates differ from H99O by 29,000 SNP and 3,386 indels. AD2-07 and AD3-55a were both unable to produce TCs (Fig 7B), allowing a more fine-tuned analysis of SNPs in these closely related strains that could explain the variable TCs generation observed between the strains (Fig 7B).

Comparison of the AD2-06a, AD2-07, and AD3-55a genomes identified four genes with

loss-of-function mutations in AD2-06a but not in AD2-07 or AD3-55a: CNAG_00570 (PKR1) (Fig

7E), CNAG_07475 (hypothetical protein), CNAG_01240 (hypothetical protein) and

CNAG_05335 (hypothetical protein). More precisely, AD2-06a had a frameshift mutation at

glycine 124 in the CNAG_00570 (*PKR1*) leading to a truncated protein of 138 amino acids (Fig

7E).

Pkr1 is the cAMP-dependent protein kinase regulatory subunit that interacts with Pka

to regulate the phosphorylation activity of Pka. We identified that *PKR1* harbored a wild type

sequence in AD2-07, together with a dramatic decrease in TCs generation. In details, the

proportion of TCs generated AD2-07 was 1.0% (31/3047) compared to AD2-06a (39.1%,

(639/1633)) and H99O (15.2% (304/2001)), (p<0.0001). The median cell size of AD2-07 was

significantly decreased compared to AD2-06a and H99O (5.9 μm [5.2-6.6] for AD2-07, 8.5 μm

[7.0-13.0] for AD2-06a, and 7.7 μm [6.5-9.2] for H99O, p<0.0001, S8 Fig).

We then selected additional and already sequenced clinical isolates (S4 Table) that harbored

mutations leading to Pkr1 truncation at different levels (Bt156, Bt58, 8-1, Bt77, Bt117,

Ug2462) [36]. Specifically, a frameshift mutation at amino acid 14 with a stop introduced at

96 for Bt156, and a stop codon introduced at 130 for 2-06a, at 258 for Bt58, at 302 for 8-1, at

439 for Bt77, at 441 for Bt117, and at 445 for Ug2462 was observed (Fig 7E, Table 3). We

hypothesized that the strains with highly impacted/truncated Pkr1 protein would produce

more TCs. These strains were then tested using the TCG protocol together with AD2-06a. The

two largest truncations were observed in AD2-06a and Bt156. These two strains had high level

of TCs generated with a ratio of 2.8±1.0 for AD2-06a and 1.3±0.6 for Bt156 compared to H99O

(Fig7F), and a % of TC of 39.1% (639/1633) (median 8.5μm [7.0-13.0]) and 17.9% (469/2614)

(median 7.7μm [6.4-9.4]), respectively (Fig S8). For the other strains, the ratio and the

proportion of TCs generated and the median size was less than that of H99O: 0.2±0.0, 3.6%

(89/2464) and 6.9µm [6.0-7.8] for 8-1 strain; 0.2±0.1, 2.9% (110/3834) and 6.5µm [5.8-7.1] for

Bt77; 0.1±0.0 0.9% (27/3081) and 6.4µm [5.7-7.2] for Bt117; and 0.1±0.1, 1.5% (54/3628)

5.4µm [4.7-6.1] for Ug2462, p<0.0001) (Fig 7F, S8 Fig).

We then tested the ability of *pkr1Δ* in a KN99α background to generate TCs compared to

KN99α and H99O, and found that the *pkr1Δ* ratio was at 1.1±0.2 compared to the parental

strain KN99 α 0.0±0.0. The *pkr1Δ* median cell size (8.1 µm [6.7-9.5]) exceeded that of KN99α

(6.4 µm [5.5-7.2]) without reaching the level of H99O (8.7 µm [7.1-10.2]), p<0.0001 (data not

shown). And, TCs proportion was at 28.8% (207/719) for H99O, 1.0% (12/1179) for KN99α and

20.4% (191/935) for *pkr1Δ* (p<0.0001, Fig 7G).

Discussion

We identified and validated a new protocol allowing robust generation of TCs *in vitro*. This protocol was discovered serendipitously while testing conditions that could induce dormancy in Cn [4]. We observed yeast cell enlargement under defined growth conditions, then optimized those conditions for TCs generation. It is important to note that the utility of other published protocols to generate TCs *in vitro* were hindered by issues with inter-laboratory reproducibility [13,31]. To establish the inter-laboratory transferability of our protocol it was independently tested in two laboratories (K. Nielsen and A. Casadevall) and the protocol produced equivalent results in all laboratories. Interestingly, TCs generation *in vitro* was also observed by Ballou et al., 2017, using a different set of growth conditions (Ballou et al., companion paper). Exploration of the similarities and differences between these protocols will likely identify the critical environmental conditions that trigger titan cell production *in vivo*.

TCs obtained *in vitro* exhibit many of the characteristics of *in vivo* titan cells recovered from the lungs of infected mice [17]. Similar to previous work on *in vivo* titan cells, we defined TCs as having a cell body size > 10 μm and RCs with a cell body size ≤ 10μm [12,13,18,21]. Melanization was increased in TCs compared to RCs and capsule size was slightly increased in TCs compared to RCs but less than previously shown *in vivo* [12,13]. In addition, immunofluorescence staining with 4 different antibodies showed no major difference between the TCs and RCs capsule structure, indicating a conservation of polysaccharide epitopes consistent with similar structures. The difference in melanization and capsule size between TCs and RCs underscore the ability and the complexity of the regulation of these virulence factors, leading to an adaptation of Cn to stress conditions, including hosts.

The cell wall was clearly thicker in TCs than in RC, with an increased chitin content

(determined both by increased CFW intensity and GlcNAc content). Increased CFW staining

and GlcNAc levels are also increased in *in vivo*-derived titan cells [30]. Finally, TCs generated

with our protocol were also polyplods, as previously shown *in vivo* [10,11].

TCs generation *in vitro* allowed a detailed kinetic analysis that revealed TCs start

forming in as little as 8 h, compared to 24 h for *in vivo* TCs [12,13]. Using our CFW method to

follow cell fate and cell division [4,28], we showed that the TCs were exclusively derived from

cells present in the initial inoculum that evolved progressively toward a TCs phenotype. In

contrast, the RCs were likely to be a mixture of cells from the initial inoculum and progeny of

cell division. TCs cell division produced regular-sized haploid RC, as shown previously for *in*

vivo titan cells [12,13]. We published *in vivo* data that validate this observation [4]. Indeed,

Calcofluor[High] Cn population were associated with yeasts cells harboring high cell size

parameters using multispectral imaging flow cytometry of yeasts recovered from the lung of

infected mice 7 d after inoculation of Cn stained with calcofluor [4].

TCs generation coincided with the appearance of a large vacuole in the yeasts cells at

4 and 8 h of incubation. Recent evidence places vacuoles at the center of networks enabling

nutrient resources to be degraded, sorted and redistributed. As the vacuole volume occupies

much of the total volume of the TCs body size, one can imagine that cell-cycle regulation could

be impacted, ultimately leading to polyploidy [37].

The fact that *in vitro* protocol consistently produced TCs allowed us to test factors that

influenced their generation. In terms of environmental factors, we showed that TCs

generation was impacted by pre-culture medium, initial pH, light exposure, temperature, type

of medium and hypoxia. A metabolic switch between YPD pre-culture and MM incubation was

a key factor to induce TCs generation. This switch is a stress that induces a lot of metabolic

modifications and has been studied extensively in *Saccharomyces cerevisiae* [38]. Hypoxia is

another stress factor in human pathogenic fungi [39]. A major transcriptional regulator of the

fungal hypoxia response is the sterol regulatory element-binding protein (Srebp) [40]. Null

mutants of the Srebp (*sre1Δ*) in Cn display defects in adaptation to hypoxia, ergosterol

synthesis, susceptibility to triazole antifungal drugs and cause a reduction of virulence [41]. As

hypoxia was an important factor for TCs generation, whether Srebp is involved in TCs

formation needs to be determined. Also, oxygen levels in healthy human tissues are 20–70

mmHg (2.5–9% O_2), and can be less than 10 mmHg (~1% O_2) [42]. We know from previous

work on pulmonary aspergillosis that hypoxia have been observed in infected lungs of mice

[43]. Titan cells have been reported in human pulmonary cryptococcosis and well-studied in

murine pulmonary infection following inhalation [13,17–21], although they are also observed

in mouse lungs after intravenous inoculation of animals [4]. These observations lead us to

hypothesize that low oxygen levels in the lungs could be one of signals that trigger titan cell

formation.

Interestingly, quorum sensing is also involved in TCs generation. Indeed, the initial

concentration of yeasts in MM dramatically impacted TCs generation, with 10^6 cells/mL as the

optimal cell concentration (no TCs formation at 10^7 cells/mL). The addition Qsp1 peptide

[30,44] at step 3 to wild type H99O yeasts (already producing Qsp1) decreased TCs

generation, suggesting Qsp1 regulation may be important for sensing the population size.

Finally, pantothenic acid (vitamin B5) [29] had little impact on TCs generation by producing

more TC in proportion. These results show that intercellular communication is important for

the yeast to switch to the TCs phenotype and that this sensing process may involve cell-cell

communication instead of simply sensing nutrient levels. A relationship between stress

sensing and quorum sensing could be hypothesized based on our results and needs to be

further explored.

Host factors such as temperature, addition of lipids, presence of serum, and antibodies

were also tested. The presence of E1 [45] and 18B7 [46] anti-capsular IgG antibodies

decreased TCs generation. This decrease could be related to changes in the yeast metabolism

induced directly by anti-capsular antibodies as shown previously [47] and provide a new

mechanism by which antibodies could alter the course of infection. One could imagine that

specific anti-Cryptococcus anti-capsular antibodies could not diffuse locally in the lung as

compared to other organs. The presence of surfactant protein-D considered as an opsonin in

the lung could impair antibody fixation [48], thus inhibiting the inhibition effect of IgG

antibodies allowing at the end TC formation in the lung. The ability of capsule-binding

antibodies to interfere with TCs formation suggests another mechanism by which humoral

immunity could contribute to protection. Interestingly, addition of serum (FCS 5%) or L-α-

phosphatidylcholine decreased TCs generation, which is different from other protocols for TCs

generation [31] Balou et al. Companion paper]. This difference could be explained by the fact

that we developed a set of inducing conditions based on specific stresses that induce TCs

through parallel or independent pathways to those triggered by serum or lipids. Overall these

results imply the existence of numerous triggers for TCs formation mediated through

independent signaling pathways that can potentially interfere with each other.

The impact of antifungal drugs was also tested. Fluconazole is known to inhibit the 14-alpha-

demethylase (Erg11) involved in ergosterol synthesis, leading to plasma membrane instability

and accumulation of toxic precursors [49]. Flucytosine is a nucleoside analogue leading to

inhibition of DNA replication and protein synthesis [50]. TCs generation was clearly inhibited

by the addition of fluconazole or flucytosine even at concentrations that are below the MIC of

the strain. These data suggest TCs generation may be an active process requiring normal

protein and DNA replication processes, as well as plasma membrane integrity. Conversely,

stressing the yeast by serial passage on fluconazole agar (8 sub-cultures) led to increased TCs

generation, that assess that pre-activation of yeasts in specific conditions (YPD, fluconazole

exposure) are required for TC formation. In other experimental settings, exposure of TCs to

fluconazole selected for aneuploidy in the daughter cells [20]. Of note, exposure to cell-wall

stress induced by serial passage on CFW agar, decreased TCs generation and suggests that

intact cell wall metabolism is required for TCs generation. In these sub-culturing experiments,

we did not investigate subsequent genomic or metabolic changes that arise under these stress

conditions. Serial sub-culture could have induced genetic rearrangements (aneuploidy, SNPs,

indels) or epigenetic variation that could explain the altered TCs generation and can be

explored further in future studies.

Our TCG protocol is practical for studying molecular and genetic mechanisms

underlying TCs generation. The TCG protocol allowed us to identify host, environmental and

yeast factors that impact TCs generation. By taking advantage of strains harboring few genetic

differences (H99 strains or AD2-06a and AD2-07), the TCG protocol is also a powerful tool to

assess genetic factors modulating TCs generation, as shown with *SGF29*, *LMP1*, and *PKR1*.

Sgf29, is a transcription factor that binds H3K4me2/3 and recruits histone deacetylation in *S.*

cerevisae [51]. *LMP1* gene is known to be involved in virulence in mouse model and mating

[32]. It is clear that these genes are required for TCs generation, although their mechanism of

action remains unclear. We also reproduced *in vitro*, the role of the Gpr/PKA/Rim101 pathway

in TCs formation previously observed *in vivo* [18]. Gpr5 signals through Gpa1 to trigger the

PKA pathway that activates the transcription factor Rim101 [18,33]. Altogether, these results

are a proof of concept that the TCG protocol can be used to identify and characterize genes

required for TCs generation. We still only know a handful of genes involved in TCs generation

and it is likely are large number of genes are involved in generation of this complex cell

morphology.

Analysis of differences in TCs generation, combined with complete genome

sequencing, allowed us to identify naturally occurring mutations in the *PKR1* gene that impact

TCs generation. Pkr1 is part of the cAMP/PKA pathway previously shown to modulate titan

cell formation *in vivo* [52]. The Pkr1 protein is known as the regulator of Pka activity [22]. The

pathway is responsible of cell cycle regulation in *S. cerevisae* and in *Xenopus* oocytes [53,54].

We tested the *pkr1Δ* mutant (CNAG_00570) from the Madhani deletion collection and it

produced 20-fold more TCs than the parental KN99α strain, corroborating the regulatory role

of Pkr1 on Pka, and the role of Pka in yeast morphogenesis. Interestingly, a *PKR1* mutation

was identified in an incident clinical isolate (AD2-06a). Surprisingly, the recurrent isolate

recovered following 13 d of amphotericin B treatment (AD2-07) did not have this *PKR1*

mutation. This could be indirect evidence that *PKR1* loss of function impacts the fitness of the

strain under amphotericin B treatment. Interestingly, *PKR1* mutation leading to intron

retention was found in a relapse isolate [55] and shown to be less virulent than the incident

isolate. Whether the virulence observed with the relapse isolates is linked to TCs formation

needs to be further investigated. In *Saccharomyces cerevisiae* the Pka1/Pkr1 complex is a

heterotetramer with 2 catalytic subunits and 2 regulatory units. This complex is dissociated in

the presence of cAMP [56]. Moreover, the architecture of the functional domains of *Pkr1*

include one interaction domain/dimerization at amino acids 2 to 40 and two cAMP binding

domains at amino acids 219 to 351 and 353 to 473, based on INTERPRO data (Fig 7E).

Consequently, PKA1/PKR1 complex dissociation is critical for binding of cAMP on the Pka1

domain. Interestingly, two distinct naturally occurring *PKR1* mutations were identified in the

clinical strains that all impacted TCs formation by reducing the protein length by introducing

a stop codon (Gly125fs for AD2-06 and Asp14fs for Bt58).

Altogether, these results are a proof of concept that the TCG protocol can be used

to identify and characterize genes required for TCs generation. We still only know a handful

of genes involved in TCs generation and it is likely that large number of genes are involved in

generation of this complex cell morphology. Our data provide new insight into the genesis of

TCs and into the environmental, host and genetic factors that influence the generation of TCs.

Our data show that our *in vitro* assay can be used to reproducibly generate TCs that have

similar characteristics to titan cells generated *in vivo*. The conditions identified for TCs

formation provide a robust system that could be invaluable to dissect the molecular

mechanisms that underlie TCs formation and allowed the identification of naturally occurring

mutations that regulate TCs formation.

Material and methods

Ethics statement

Mice (purchased from Jackson Laboratories, Bar Harbor, ME) were handled in accordance with guidelines defined by the University of Minnesota Animal Care and Use Committee (IACUC), under approved protocol numbers 1010A91133 and 130830852. All animal experiments were performed in concordance with the Animal Welfare Act, United States federal law, and NIH guidelines.

Strains and culture medium

The strains and clinical isolates of *C. neoformans* (Cn) used in the study are listed in S1 Table. The study was started with H99 strain called H99O that was kindly provided by J. Heitman (Duke University, Durham, NC) in the late 90's. Strains from the Madhani collection were provided from the Fungal Genetic Stock Center [57].

Cn strains were grown in liquid Yeast Peptone Dextrose (YPD, 1% yeast extract (BD Difco, Le pont de Claix, France) 2% peptone (BD difco), 2% D-glucose (Sigma, Saint Louis, Minnesota, USA)) and in minimal medium (MM, 15mM D-glucose (Sigma), 10 mM $MgSO_4$ (Sigma), 29.4mM KH_2PO_4 (Sigma), 13mM Glycine (Sigma) , 3.0 μM Thiamine (Sigma), [29]). Minimum inhibitory concentration (MIC) of H99O for fluconazole (FLC) and flucytosine (5FC) (both purchased from lsachim, Shimadzu Group Company, Illkirch-Graffenstaden, France) were determined by the EUCAST method and were 8 and 4 mg/L, respectively.

In vitro protocol for TCs generation (TCG protocol)

Cn strain from stock cultures stored in 20% glycerol at -80°C was cultured on Sabouraud agar

plate at room temperature (step 1). After 2 to 5 d of culture, approximately 10^7 cells were

suspended in 10mL YPD in a T25cm^3 flask and cultured 22 h at 30°C, 150 rpm with lateral

shaking until stationary phase (final concentration=2x10^{8}cells/mL) (step 2). Then, one mL of

the suspension was washed twice with MM. The cell concentration was adjusted to 10^6

cells/mL in MM and the suspension was incubated in a 1.5 mL tube (Eppendorf) with the cap

closed, at 30°C, 800 rpm for 5 d using an Eppendorf Thermomixer® (Hambourg, Germany)

(step 3). Cell size was determined as described below. Cells with body size >10 μm were

considered as TCs as described [12]. Results are expressed as median cell size [interquartile

range, IQR] or as median [IQR] of the % of TCs in a given condition for H99 or as a ratio

compared to %TCs obtained with the H99O in experiments involving other strains (clinical

isolates, other H99 strains and mutants).

Capsule size analysis and melanization

Yeasts were observed after India ink staining and capsule size was determined as described

below using the following formula: whole cell size - cell body size)/2.

Capsule immunofluorescence (IF) of TCs was done by incubating approximately 5x10^6 cells/mL

with 10μg/mL of murine-derived monoclonal antibodies to the cryptococcal capsule (IgG1

18B7, IgG1 E1, IgM 12A1, IgM 2D10 [27,45,58,59] in blocking solution (1% bovine serum

albumin in PBS). Cells and mAb mixtures were done in 1.5 mL microcentrifuge tubes at 37 °C

for 1 h under continuous mixing. Next, cells were washed three times with PBS by

centrifugation (5,000 rpm for 5 mins at room temperature) and incubated for 1 h at 37 °C with

5 μg/mL fluorescently labelled secondary-mAbs, goat anti-mouse IgG1-FITCs or IgM-TRITCs

(Southern Biotech) in blocking solution and 1 μg/mL of Uvitex2b (Polysciences, Warrington,

PA) solution to visualize the fungal cell wall. Cells were washed three times with PBS by centrifugation, mounted in glass coverslips and imaged with an Olympus AX 70 microscope equipped with blue, green and red fluorescent filters using 40x and/or oil immersion 100x objectives. Capsule IF of TCs preparations performed in two independent experiments gave consistent results.

TCs melanization was induced following step 3. Cells were washed once with MM, suspended in 1mL of MM supplemented with 1mM of L-DOPA (Sigma D9628), transferred to a 5mL Erlenmeyer flask (for normal oxygenation) and incubated at 30 °C under continuous mixing at 200 rpms for 3 days. Since melanin is resistant to acid hydrolysis, a spherical melanin "ghost" remains following incubation of black cells with 12N HCl for 1 h at 100 °C (a reduced and modified version of the procedure in [60]. Acid-resistant melanin "ghosts" were washed three times in PBS by centrifugation and visualized using light microscopy.

Melanization was measured using imageJ in Icy software by manually circling each cell and measuring the mean gray intensity / pixel /cell. Blackness was calculated as the maximum mean grey intensity minus the mean gray intensity / pixel /cell. Increasing melanin content will result in higher blackness. As melanization darkened the cells, the most darkened cells were the less intense the mean gray was.

Chitin density quantification

Chitin density was determined after calcofluor white staining (CFW, fluorescent brightener 28) adapted from [37]. Briefly, 10^7 Cn cells in 10 mL MM were washed once and 500 µl of 3.7% formaldehyde in PBS was added. Cells were incubated at room temperature for 30-40 minutes, inverting the tube every 5 minutes. Samples were washed twice in PBS, cell concentration was adjusted to 10^6 cell/ml. The supernatant was removed and 1 mL of 0.01

μg/mL CFW in PBS was added, and incubated 5 minutes at 25°C. Cells were then washed twice

in PBS. Results were expressed as median [IQR] of the mean fluorescence intensity/pixel/cell

after picture analysis as described below.

N-acetylglucosamine quantification/ Chitin measurement

H99KN cells were grown in MM for 48 h *in vitro*. Cells were centrifugated 2 min at 14,000 and

were then washed twice with sterile water. These cells were exposed to γ-radiation to remove

layers of the capsule polysaccharide [13]. Cells resuspended in sterile water were transferred

to a 24-well flat-bottom plate and irradiated for 45 min: dose 560 Gy (56,000 rad).

TCs and RCs populations were separated [20]. Washed irradiated cells were filtered using

CellMicroSieves (BioDesign Inc. of New York, Carmel, NY) with a 10 μm pore size. The

CellMicroSieves were rinsed with PBS to remove RCs from the filter. To recover the TCs

population, the CellMicroSieves were inverted and the membrane was washed with PBS. The

TCs population was concentrated by centrifugation at 12,000 g for 1 min. To recover the RCs

population, the filter flow-through was concentrated by centrifugation at 12,000 g for 1 min.

Cellular chitin quantification was adapted from [61]. Purified *in vitro* RCs and TCs were

collected by centrifugation at 14,000 rpm for 2 min and the media were removed. Dry weights

were measured following 2-3 d of evaporation at 37°C. Dried pellets were extracted with 1 ml

6% KOH at 80°C for 90 min. Samples were centrifuged at 14,000 rpm for 20 min. Each pellet

was suspended in 1 ml PBS and spun again. Each pellet was suspended in 0.2 ml of McIlvaine's

Buffer (0.2 M Na_2HPO_4, 0.1 M citric acid, pH 6.0). Five μL of purified *Streptomyces griseus*

chitinase (5 mg/mL in PBS) was added to hydrolyze chitin to GlcNAc and were incubated for 3

d at 37°C. Chitinase-treated samples were spun at 14,000 rpm for 1 min, each 10 μL of sample

supernatant was combined with 10 μL 0.27 M sodium borate, pH 9.0. Samples were heated

to 99.9°C for 10 min. Upon cooling to room temperature, 100 µl of DMAB solution (Ehrlich's

reagent, 10 g p-dimethylaminobenzaldehyde in 12.5 mL concentrated HCl, and 87.5 ml glacial

acetic acid) was added, followed by 20 min incubation at 37°C. Hundred µL was transferred to

96-well plates, and absorbance at 585 nm was recorded. Standard curves were prepared from

stocks of 0.2 to 2.0 mM of GlcNAc (Sigma, Saint Louis, Missouri, USA). The amount of GlcNAc

was calculated as mmol/g cells (dry weight). Results are expressed as median [IQR].

DNA content measurement

A 96-well microtiter plate was filled with 200 µL of a 10^6/mL cell suspension in PBS and

centrifuged 5 minutes at 4000 rpm. The pellet was suspended in 150 µL of ethanol 70% and

incubated in the dark 1 h at 4°C. After discarding the supernatant, a 50 µL mix composed of

44 µL NS (0.01M Tris HCL pH 7.2, 1 mM EDTA, 1mM CaCl2, 0.25 M Sucrose, 2.12 mM $MgCl_2$,

0.1 mM $ZnCl_2$), 5µL RNase A at 10mg/mL and 1.25µL PI at 0.5mg/mL was added in each well

as described [62]. After a 30 minutes' incubation at 30°C in the dark, the plate was sonicated

1 minute and each sample diluted at 1:40 in 50mM Tris HCl. The fluorescence intensity was

measured using the Guava easyCyte 12HT Benchtop Flow Cytometer® (Guava, MERCK,

Kenilworth, New Jersey). Selection of singlets by gating allowed (i) determination of PI

intensity on channel YelB (583/26) in FSC/SSC[high] (TCs) and FSC/SSC[low] (RCs); (ii) determination

of the FSC/SSC distribution in PI[high] and PI[low] population. FlowJo software v.10 was used to

analyze the data. The graphs of the number of yeasts were normalized to the mode to depict

the data in terms of '% of max'. The % of max denotes the number of cells in each bin (the

numerical ranges for the parameter on the *x* axis) divided by the number of cells in the bin

that contains the largest number of cells.

Determination of the ancestry of TCs by flow cytometry

Knowing that CFW staining does not alter Cn viability, and that daughter cells harbored lower CFW signal due to partial cell wall transmission from mother to daughter cells [4,28], we analyzed cell size and CFW fluorescence intensity of the progenies of TCs and RCs following the TCG protocol on 10^6 cells of H99O pre-stained with CFW (channel BluV 448/50 using Guava).

Dynamic imaging

Cell size evolution over time was assessed for strain AD2-06a by dynamic imaging (Nikon Biostation). Briefly, 35 mm sterile culture dish (Hi-Q4, Nikon) were coated for 1 min with E1 antibody at 2 mg/L in order to provide anchor for the capsule. Yeasts (10^5 cells) were added in 1 mL MM and incubated at 32°C for 18 h. Series of 221 images were taken by phase-contrast microscopy every 5 mins at ×100 magnification. Merging was done using ImageJ software in Icy Software.

Impact of various factors on TCs generation

To analyze the various factors that could impact TCs generation, we modified the various steps of the TCG protocol. For step1, stress was produced by 8 subcultures (twice a week for one month) on agar medium or on agar supplemented with CFW (20mg/L) or with fluconazole (32mg/L). For step 2, the pH of MM (normally at 5.5) was set at 4, 7 or 8.5 without buffering. For step 3, initial cell concentration (from 10^4 to 10^7cells/mL) was tested. Hypoxia was generated physically by closing the cap of the Eppendorf® tube during 5 d or chemically upon incubation in MM supplemented with 1 nM $CoCl_2$, cap closed, as already described [63]. The production of TCs was also assessed in the presence of various reagents added at step 3: (1)

Qsp1 peptide (NFGAPGGAYPW, [30]) (Biomatik, Cambridge, Canada) was resuspended at

10mM in water and stored at -80°C until use at 10 µM final ; (2) pantothenic acid purchased

from Sigma (Saint-Louis, Missouri, USA) used at 125 µM; (3) monoclonal anti-capsular

antibodies E1 [45] and 18B7 [46] used at a final concentration of 166 µg/mL in MM [64]; (4)

decomplemented fetal calf serum (FCS, Invitrogen, Carlsbad, CA, USA) at 5 % in MM; (5) L-α-

Phosphatidylcholine from egg yolk (Sigma, Saint-Louis, Missouri, USA) was extemporaneously

reconstituted at 5 mM in MM; (6) antifungal drugs (fluconazole and flucytosine) were tested

at the concentrations close to the MIC (2-fold dilutions) with the diluent (DMSO or water) as

control. Results are expressed as median [IQR]. Growth in the presence or in the absence of

antifungal drugs was evaluated by enumeration of yeasts cells concentration at step 4 of TCG

protocol using the Guava cytometer, starting from 10^6 cells inoculated at step 1.

Production and isolation of TCs from infected mice

Cn strains were cultured overnight at 30°C in YPD broth medium (BD, Hercules, Canada). Cells

were collected by centrifugation, washed with phosphate buffered saline PBS and

resuspended in sterile saline. Groups of 6- to 8-week-old C57BL/6J mice (Jackson Labs, Bar

Harbor, Maine) were anesthetized by intraperitoneal pentobarbital injection and infected

intranasally with 5×10^6 cells in a 50 µl volume. Infected mice were sacrificed by CO_2 inhalation

at 3 days post-infection. The lungs were harvested, homogenized, and then resuspended in

10 ml PBS supplemented with collagenase (1 mg/ml) [13]. Cell homogenates were incubated

for 1hour at 37°C with agitation, and washed several times with double distilled water. The Cn

cells were fixed with 3.7 % formaldehyde for 40 min, washed 3 times with sterile PBS, and

then resuspended in sterile PBS. The proportion of TCs and RCs were determined by

microscopy. Data presented were from 3 mice per strain, except for strains *sgf29Δ* in H99O,

H99S that had 2 mice per strain.

***In vitro* susceptibility testing**

Minimal inhibitory concentrations (MICs) were determined for flucytosine and fluconazole

(Alsachim, Shimadzu Group Company, Illkirch-Graffenstaden, France) according to the

EUCAST method [65].

DNA Sequencing, Variant identification, and Bioinformatic Analysis

Genome DNA was adapted for Illumina sequencing using Nextera reagents. Libraries were

sequenced on an Illumina HiSeq to generate 101 base reads. most data was previously

described [36,66] and one additional isolate was newly sequenced for this study (AD2-07)

(NCBI SRA accession SRR5989089). Reads were aligned to the *C. neoformans* var. *grubii* H99

assembly (GenBank accession GCA_000149245.2 [32] using BWA-MEM version 0.7.12 [67].

Variants were then identified using GATK version 3.4 [68] , where indels were locally realigned,

haplotypeCaller was invoked in GVCF mode with ploidy = 1, and genotypeGVCFs was used to

predict variants in each strain. All VCFs were then combined and sites were filtered using

variantFiltration with QD < 2.0, FS > 60.0, and MQ < 40.0. Individual genotypes were then

filtered if the minimum genotype quality < 50, percent alternate allele < 0.8, or depth < 5.

Variants were then functionally annotated with SnpEff version 4.2 [69]. For phylogenetic

analysis, the 369,574 sites with an unambiguous SNP in at least one isolate and with ambiguity

in at most 10% of isolates were concatenated; insertions or deletions at these sites were

treated as ambiguous to maintain the alignment. Phylogenetic trees were estimated using

RAxML version 8.2.4 [70] under the GTRCAT model in rapid bootstrapping mode. For

determination of PKR1 architecture domains, the INTERPRO tool was used

(http://www.ebi.ac.uk/interpro/protein/J9VH50).

Pictures analysis using Icy software and statistical analysis

To increase the number of events analyzed in each condition tested/each parameter analyzed

(cell size, capsule size and chitin content), pictures of 3-5 fields were taken with an AxioCam

MRm camera (Carl Zeiss, Oberkochen) at x40 on interferential contrast microscope (DMLB2

microscope; Leica, Oberkochen). Image were then analyzed (for cell size and chitin content)

using Icy software v.1.9.2.1. (icy.bioimageanalysis.org) and a specific plugin (HK-Means plugin

(http://icy.bioimageanalysis.org/plugin/HK-Means) that allows analysis of multiple structures

from a bright field. Preliminary experiments were done to compare results obtained with Icy

to "manual" measurements by analyzing about 200 cells on the same pictures for 3

independent experiments. In subsequent experiments, results were pooled for a given

condition from 2 to 3 independent experiments after good reproducibility was assessed.

Statistical analysis was performed with STATA® software (College Station, Texas, v13.0). To

validate the cell size determination using the Icy software, the intraclass correlation

coefficient was calculated. The ability of the automated method to classify the Cn cells as TCs

or RCs compared to visual measurement was evaluated using the Kappa test [71]. To compare

TCs generation in the various conditions, non-parametric tests were performed using the

Kruskal-Wallis test for multiple comparisons or Mann Whitney test as required. GraphPad

Prism software (v.6) was used to generate graphs.

Data access

All sequence data from this study have been submitted to NCBI BioProject

(https://www.ncbi.nlm.nih.gov/bioproject) under accession number PRJNA174567.

The AD2-07 sequence is available in the NCBI SRA under the accession number SRR5989089

(https://www.ncbi.nlm.nih.gov/sra/SRR5989089/)

Acknowledgments

We thank Dr Tihana Bicanic, Shichina Kannmbath, Charles Giamberardino and Jennifer Tenor

who kindly provided specific clinical isolates. The authors want to thank Marie Desnos-Ollivier

for her help with Sanger sequencing, JL Tinevez for technical assistance in Biostation

experiments, Stéphane Dallongeville for help with Icy Software, Stevenn Volant for

biostatistics and Paul-Henri Commere for FACS analysis. We thank Quigly Dragotakes for

helping us determining cell and capsule sizes in specific experiments. The authors

acknowledge Hiten Madhani and member of his laboratory for the knockout collection that

have been made freely available ahead of publication to the scientific community.

References

1.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, et al. Global burden
of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect
Dis. 2017;17: 873. doi:10.1016/s1473-3099(17)30243-8

2.   Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. Estimation
of the current global burden of cryptococcal meningitis among persons living with
HIV/AIDS. AIDS Lond Engl. 2009;23: 525–530. doi:10.1097/QAD.0b013e328322ffac

3.   Garcia-Hermoso D, Janbon G, Dromer F. Epidemiological evidence for dormant
Cryptococcus neoformans infection. J Clin Microbiol. 1999;37: 3204–3209.

4.   Alanio A, Vernel-Pauillac F, Sturny-Leclère A, Dromer F. *Cryptococcus neoformans* host
adaptation: toward biological evidence of dormancy. mBio. 2015;6: e02580-14.
doi:10.1128/mBio.02580-14

5.   Chrétien F, Lortholary O, Kansau I, Neuville S, Gray F, Dromer F. Pathogenesis of cerebral
Cryptococcus neoformans infection after fungemia. J Infect Dis. 2002;186: 522–530.
doi:10.1086/341564

6.   Choi J, Vogl AW, Kronstad JW. Regulated expression of cyclic AMP-dependent protein
kinase A reveals an influence on cell size and the secretion of virulence factors in
Cryptococcus neoformans. Mol Microbiol. 2012;85: 700–715. doi:10.1111/j.1365-
2958.2012.08134.x

7.   Franzot SP, Mukherjee J, Cherniak R, Chen LC, Hamdan JS, Casadevall A. Microevolution of
a standard strain of *Cryptococcus neoformans* resulting in differences in virulence and
other phenotypes. Infect Immun. 1998;66: 89–97.

8.   Hu G, Cheng P-Y, Sham A, Perfect JR, Kronstad JW. Metabolic adaptation in *Cryptococcus*
neoformans during early murine pulmonary infection. Mol Microbiol. 2008;69: 1456–
1475. doi:10.1111/j.1365-2958.2008.06374.x

9.   Gerstein AC, Nielsen K. It's not all about us: evolution and maintenance of *Cryptococcus*
virulence requires selection outside the human host. Yeast. 2017;34: 143–154.
doi:10.1002/yea.3222

10. Brown SM, Campbell LT, Lodge JK. *Cryptococcus neoformans*, a fungus under stress. Curr
Opin Microbiol. 2007;10: 320–325. doi:10.1016/j.mib.2007.05.014

11. Wang L, Lin X. Morphogenesis in fungal pathogenicity: shape, size, and surface. PLOS
Pathog. 2012;8: e1003027. doi:10.1371/journal.ppat.1003027

12. Okagaki LH, Strain AK, Nielsen JN, Charlier C. Cryptococcal cell morphology affects host
cell interactions and pathogenicity. PLoS Pathog. 2010;6: e1000953.
doi:10.1371/journal.ppat.1000953

13. Zaragoza O, García-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodríguez-Tudela JL,
Casadevall A. Fungal cell gigantism during mammalian infection. PLoS Pathog. 2010;6:
e1000945. doi:10.1371/journal.ppat.1000945

14. Love GL, Boyd GD, Greer DL. Large *Cryptococcus neoformans* isolated from brain abscess.
J Clin Microbiol. 1985;22: 1068–1070.

15. Cruickshank JG, Cavill R, Jelbert M. *Cryptococcus neoformans* of unusual morphology. Appl
Microbiol. 1973;25: 309–312.

16. García-Rodas R, Casadevall A, Rodríguez-Tudela JL, Cuenca-Estrella M, Zaragoza O.
Cryptococcus neoformans capsular enlargement and cellular gigantism during *Galleria*
mellonella infection. PloS One. 2011;6: e24485. doi:10.1371/journal.pone.0024485

17. Zaragoza O, Nielsen K. Titan cells in *Cryptococcus neoformans*: cells with a giant impact.
Curr Opin Microbiol. 2013;16: 409–413. doi:10.1016/j.mib.2013.03.006

18. Okagaki LH, Wang Y, Ballou ER, O'Meara TR, Bahn Y-S, Alspaugh JA, et al. Cryptococcal
titan cell formation is regulated by G-protein signaling in response to multiple stimuli.
Eukaryot Cell. 2011;10: 1306–1316. doi:10.1128/EC.05179-11

19. Crabtree JN, Okagaki LH, Wiesner DL, Strain AK, Nielsen JN, Nielsen K. Titan cell production
enhances the virulence of *Cryptococcus neoformans*. Infect Immun. 2012;80: 3776–3785.
doi:10.1128/IAI.00507-12

20. Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, et al. Polyploid titan cells
produce haploid and aneuploid progeny to promote stress adaptation. mBio. 2015;6:
e01340-15. doi:10.1128/mBio.01340-15

21. Okagaki LH, Nielsen K. Titan cells confer protection from phagocytosis in *Cryptococcus*
neoformans infections. Eukaryot Cell. 2012;11: 820–826. doi:10.1128/EC.00121-12

22. D'Souza CA, Alspaugh JA, Yue C, Harashima T, Cox GM, Perfect JR, et al. Cyclic AMP-
dependent protein kinase controls virulence of the fungal pathogen *Cryptococcus*
neoformans. Mol Cell Biol. 2001;21: 3179–3191. doi:10.1128/MCB.21.9.3179-3191.2001

23. Nichols CB, Perfect ZH, Alspaugh JA. A Ras1-Cdc24 signal transduction pathway mediates
thermotolerance in the fungal pathogen *Cryptococcus neoformans*. Mol Microbiol.
2007;63: 1118–1130. doi:10.1111/j.1365-2958.2006.05566.x

24. Bahn Y-S, Jung K-W. Stress Signaling Pathways for the Pathogenicity of *Cryptococcus*.
Eukaryot Cell. 2013;12: 1564–1577. doi:10.1128/EC.00218-13

25. Hu G, Steen BR, Lian T, Sham AP, Tam N, Tangen KL, et al. Transcriptional Regulation by
Protein Kinase A in *Cryptococcus neoformans*. PLOS Pathog. 2007;3: e42.
doi:10.1371/journal.ppat.0030042

26. Geddes JMH, Caza M, Croll D, Stoynov N, Foster LJ, Kronstad JW. Analysis of the protein
kinase A-regulated proteome of *Cryptococcus neoformans* identifies a role for the

ubiquitin-proteasome pathway in capsule formation. mBio. 2016;7: e01862-15.
doi:10.1128/mBio.01862-15

27. Casadevall A, DeShaw M, Fan M, Dromer F, Kozel TR, Pirofski L a. Molecular and idiotypic
analysis of antibodies to *Cryptococcus neoformans* glucuronoxylomannan. Infect Immun.
1994;62: 3864–3872.

28. Alanio A, Desnos-Ollivier M, Dromer F. Dynamics of *Cryptococcus neoformans*-
macrophage interactions reveal that fungal background influences outcome during
cryptococcal meningoencephalitis in humans. mBio. 2011;2: e00158-11.
doi:10.1128/mBio.00158-11

29. Albuquerque P, Nicola AM, Nieves E, Paes HC, Williamson PR, Silva-Pereira I, et al. Quorum
sensing-mediated, cell density-dependent regulation of growth and virulence in
Cryptococcus neoformans. mBio. 2013;5: e00986-13. doi:10.1128/mBio.00986-13

30. Homer CM, Summers DK, Goranov AI, Clarke SC, Wiesner DL, Diedrich JK, et al.
Intracellular action of a secreted peptide required for fungal virulence. Cell Host Microbe.
2016;19: 849–864. doi:10.1016/j.chom.2016.05.001

31. Chrisman CJ, Albuquerque P, Guimarães AJ, Nieves E, Casadevall A. Phospholipids trigger
Cryptococcus neoformans capsular enlargement during interactions with amoebae and
macrophages. PLoS Pathog. 2011;7: e1002047. doi:10.1371/journal.ppat.1002047

32. Janbon G, Ormerod KL, Paulet D, Byrnes EJ, Yadav V, Chatterjee G, et al. Analysis of the
genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex RNA
expression and microevolution leading to virulence attenuation. PLoS Genet. 2014;10:
e1004261. doi:10.1371/journal.pgen.1004261

33. O'Meara TR, Alspaugh JA. The *Cryptococcus neoformans* capsule: a sword and a shield.
Clin Microbiol Rev. 2012;25: 387–408. doi:10.1128/CMR.00001-12

34. Cordero RJB, Casadevall A. Functions of fungal melanin beyond virulence. Fungal Biol Rev.
2017;31: 99. doi:10.1016/j.fbr.2016.12.003

35. Rhodes J, Desjardins CA, Sykes SM, Beale MA, Vanhove M, Sakthikumar S, et al. Tracing
genetic exchange and biogeography of *Cryptococcus neoformans* var. *grubii* at the global
population level. Genetics. 2017; doi:10.1534/genetics.117.203836

36. Desjardins CA, Giamberardino C, Sykes SM, Yu C-H, Tenor JL, Chen Y, et al. Population
genomics and the evolution of virulence in the fungal pathogen *Cryptococcus neoformans*.
Genome Res. 2017;27: 1207–1219. doi:10.1101/gr.218727.116

37. Wiesner DL, Specht CA, Lee CK, Smith KD, Mukaremera L, Lee ST, et al. Chitin recognition
via chitotriosidase promotes pathologic type-2 helper T cell responses to cryptococcal
infection. PLoS Pathog. 2015;11: e1004701. doi:10.1371/journal.ppat.1004701

38. Zhang N, Cao L. Starvation signals in yeast are integrated to coordinate metabolic
reprogramming and stress response to ensure longevity. Curr Genet. 2017;
doi:10.1007/s00294-017-0697-4

39. Nizet V, Johnson RS. Interdependence of hypoxic and innate immune responses. Nat Rev
Immunol. 2009;9: 609–617. doi:10.1038/nri2607

40. Grahl N, Shepardson KM, Chung D, Cramer RA. Hypoxia and fungal pathogenesis: to air or
not to air? Eukaryot Cell. 2012;11: 560–570. doi:10.1128/EC.00031-12

41. Chun CD, Liu OW, Madhani HD. A link between virulence and homeostatic responses to
hypoxia during infection by the human fungal pathogen *Cryptococcus neoformans*. PLoS
Pathog. 2007;3: e22. doi:10.1371/journal.ppat.0030022

42. Lewis JS, Lee JA, Underwood JC, Harris AL, Lewis CE. Macrophage responses to hypoxia:
relevance to disease mechanisms. J Leukoc Biol. 1999;66: 889–900.

43. Grahl N, Puttikamonkul S, Macdonald JM, Gamcsik MP, Ngo LY, Hohl TM, et al. In vivo
hypoxia and a fungal alcohol dehydrogenase influence the pathogenesis of invasive
pulmonary   aspergillosis.   PLoS   Pathog.   2011;7:   e1002145.
doi:10.1371/journal.ppat.1002145

44. Lee H, Chang YC, Nardone G, Kwon-Chung KJ. TUP1 disruption in *Cryptococcus neoformans*
uncovers a peptide-mediated density-dependent growth phenomenon that mimics
quorum sensing. Mol Microbiol. 2007;64: 591. doi:10.1111/j.1365-2958.2007.05666.x

45. Dromer F, Salamero J, Contrepois A, Carbon C, Yeni P. Production, characterization, and
antibody specificity of a mouse monoclonal antibody reactive with *Cryptococcus
neoformans* capsular polysaccharide. Infect Immun. 1987;55: 742–748.

46. Zebedee SL, Koduri RK, Mukherjee J, Mukherjee S, Lee S, Sauer DF, et al. Mouse-human
immunoglobulin G1 chimeric antibodies with activities against *Cryptococcus neoformans*.
Antimicrob Agents Chemother. 1994;38: 1507. doi:10.1128/aac.38.7.1507

47. McClelland EE, Nicola AM, Prados-Rosales R, Casadevall A. Ab binding alters gene
expression in *Cryptococcus neoformans* and directly modulates fungal metabolism. J Clin
Invest. 2010;120: 1355–1361. doi:10.1172/JCI38322

48. Geunes-Boyer S, Beers MF, Perfect JR, Heitman J, Wright JR. Surfactant protein D
facilitates *Cryptococcus neoformans* infection. Infect Immun. 2012;80: 2444–2453.
doi:10.1128/IAI.05613-11

49. Sanglard D, Ischer F, Calabrese D, de Micheli M, Bille J. Multiple resistance mechanisms to
azole antifungals in yeast clinical isolates. Drug Resist Updat. 1998;1: 255.
doi:10.1016/s1368-7646(98)80006-x

50. Loyse A, Dromer F, Day J, Lortholary O, Harrison TS. Flucytosine and cryptococcosis: time
to urgently address the worldwide accessibility of a 50-year-old antifungal. J Antimicrob
Chemother. 2013;68: 2435–2444. doi:10.1093/jac/dkt221

51. Bian C, Xu C, Ruan J, Lee KK, Burke TL, Tempel W, et al. Sgf29 binds histone H3K4me2/3
and is required for SAGA complex recruitment and histone H3 acetylation. EMBO J.
2011;30: 2829–2842. doi:10.1038/emboj.2011.193

52. Kronstad JW, Attarian R, Cadieux B, Choi J, D'Souza CA, Griffiths EJ, et al. Expanding fungal
pathogenesis: *Cryptococcus* breaks out of the opportunistic box. Nat Rev Microbiol.
2011;9: 193–203. doi:10.1038/nrmicro2522

53. Duckworth BC, Weaver JS, Ruderman JV. G2 arrest in *Xenopus* oocytes depends on
phosphorylation of cdc25 by protein kinase A. Proc Natl Acad Sci U S A. 2002;99: 16794–
16799. doi:10.1073/pnas.222661299

54. Mitsuzawa H. Increases in cell size at START caused by hyperactivation of the cAMP
pathway in *Saccharomyces cerevisiae*. Mol Gen Genet MGG. 1994;243: 158–165.

55. Chen Y, Farrer RA, Giamberardino C, Sakthikumar S. Microevolution of serial clinical
isolates of *Cryptococcus neoformans* var. *grubii* and *C. gattii*. mBio. 2017;8: e00166-17.
doi:10.1128/mbio.00166-17

56. Vandamme J, Castermans D, Thevelein JM. Molecular mechanisms of feedback inhibition
of protein kinase A on intracellular cAMP accumulation. Cell Signal. 2012;24: 1610–1618.
doi:10.1016/j.cellsig.2012.04.001

57. McCluskey K, Wiest A, Plamann M. The Fungal Genetics Stock Center: a repository for 50
years of fungal genetics research. J Biosci. 2010;35: 119–126.

58. Casadevall A, Cleare W, Feldmesser M, Glatman-Freedman A, Goldman DL, Kozel TR, et al.
Characterization of a murine monoclonal antibody to *Cryptococcus neoformans*
polysaccharide that is a candidate for human therapeutic studies. Antimicrob Agents
Chemother. 1998;42: 1437–1446.

59. Mukherjee J, Casadevall A, Scharff MD. Molecular characterization of the humoral
responses to *Cryptococcus neoformans* infection and glucuronoxylomannan-tetanus
toxoid conjugate immunization. J Exp Med. 1993;177: 1105–1116.

60. Wang Y, Aisen P, Casadevall A. Melanin, melanin "ghosts," and melanin composition in
Cryptococcus neoformans. Infect Immun. 1996;64: 2420–2424.

61. Banks IR, Specht CA, Donlin MJ, Gerik KJ, Levitz SM, Lodge JK. A chitin synthase and its
regulator protein are critical for chitosan production and growth of the fungal pathogen
Cryptococcus neoformans. Eukaryot Cell. 2005;4: 1902–1912. doi:10.1128/EC.4.11.1902-
1912.2005

62. Tanaka R, Taguchi H, Takeo K, Miyaji M, Nishimura K. Determination of ploidy in
Cryptococcus neoformans by flow cytometry. J Med Vet Mycol Bi-Mon Publ Int Soc Hum
Anim Mycol. 1996;34: 299–301.

63. Ingavale SS, Chang YC, Lee H, McClelland CM, Leong ML, Kwon-Chung KJ. Importance of
mitochondria in survival of *Cryptococcus neoformans* under low oxygen conditions and

tolerance to cobalt chloride. PLoS Pathog. 2008;4: e1000155.
doi:10.1371/journal.ppat.1000155

64. McClelland EE, Ramagopal UA, Rivera J, Cox J, Nakouzi A, Prabu MM, et al. A Small protein
associated with fungal energy metabolism affects the virulence of *Cryptococcus*
neoformans in mammals. PLoS Pathog. 2016;12: e1005849.
doi:10.1371/journal.ppat.1005849

65. Arendrup MC, Cuenca M. Method for the determination of broth dilution minimum
inhibitory concentrations of antifungal agents for conidia forming moulds. 2014.

66. Rhodes J, Beale M, Vanhove M, Jarvis JN, Kannambath S, Simpson JA, et al. A population
genomics approach to assessing the genetic basis of within-host microevolution
underlying recurrent cryptococcal meningitis infection. G3 Genes. 2016;
doi:10.1101/083469

67. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
1303;

68. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The genome
analysis toolkit: a mapReduce framework for analyzing next-generation DNA sequencing
data. Genome Res. 2010;20: 1297–1303. doi:10.1101/gr.107524.110

69. Cingolani P, Platts A, Lily Wang, Coon M, Nguyen T, Wang L, et al. A program for annotating
and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome
of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin). 2012;6: 80–92.
doi:10.4161/fly.19695

70. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinforma Oxf Engl. 2014;30: 1312–1313.
doi:10.1093/bioinformatics/btu033

71. The Kappa Statistic in Reliability Studies: Use, Interpretation, and Sample Size
Requirements. Phys Ther. 2005; doi:10.1093/ptj/85.3.257

 Supporting informations

S1 Fig. TCs melanization. Comparing the blackness of the cell body of TCs (white arrows) and

RCs upon melanization conditions showed that TCs contained more melanin than RCs.

(Bar=10µm).

S2 Fig. Immunofluorescence staining with anti-capsular polysaccharide monoclonal

antibodies (mAbs) 2D10, 12A1, 18B7, E1 does not uncover major differences in capsular

structures between TCs (white arrows) and RCs (black arrows). Each panel correspond to the

same cells observed after staining with (a) India ink; (b) calcofluor white; (c) one of the Mabs;

(d) merge from c and d. (bar=10µm).

S3 Fig. The FSC^{high}/CFW^{high} population of yeasts correspond to TCs. Cells obtained at the end

of the TGC protocol were stained with CFW and sorted by flux cytometry according to size

(FSC) and CFW fluorescence intensity (left panel). Sorted yeasts were observed using bright

field and fluorescence microscopy (right panel) (bar=10µm).

S4 Fig. Among yeasts recovered at the end of the TCG protocol, those with the highest DNA

content have the bigger cell size.

DNA content was analyzed after propidium iodide (PI) staining of yeast cells obtained at the

end of the TCG protocol (H99O TCG), in a control haploid strain (H99O cultured in Sabouraud

agar, H99O Sab) and in a control diploid strain (AD7-77 cultured in Sabouraud agar). Part of

the population of H99O TCG had a higher PI (blue arrow) fluorescence intensity than the

haploid control (upper panel). Gating on the PI intensity showed that the increase in the PI

fluorescence intensity from <20K to >40K corresponded to increase in cell size (FSC) (red

arrows) compared to the diploid (AD7-77) and haploid (H99O Sab) control (lower panel).

S5 Fig. The yeasts growth is maintained in the presence of antifungals after TCG protocol.

10^6 cell/mL were inoculated at step 1 of TCG protocol, then cell growth was evaluated by

enumerating cell concentration obtained at step 4 of TCG protocol using Guava flow

cytometer for cell counting. (**A**) Compared to control, fluconazole did not modify cell growth

in MM whereas (**B**) flucytosine (5FC) reduced it, when used at concentration near the

minimum inhibitory concentration (MIC) for 5 days (*p<0.0001 compared to unexposed

control). The fluconazole and flucytosine MICs for H99O were 8 mg/L and 4 mg/L, respectively.

Experiments were done in triplicates (bars represent mean ± SD).

1000

S6 Fig. **TCs generation is dependent on various genes *in vitro* and *in vivo*.** (**A**) Strains from the

H99 lineage harbored variable abilities to generate TCs with high TCs producer (H99O, S, L)

and low TCs producer (KN99α, H99W, H99 CMO18) *in vitro* and (**B**) *in vivo* (except for KN99α)

compared to H99O. (**C**) The *sgf29Δ* and *lmp1Δ* mutant strains show a decrease in TCs

generation in various H99 backgrounds *in vitro* and (**D**) *in vivo* compared to H99O. (**E**) Rim

pathway is required for TCs generation *in vitro* in H99. Each experiment was done in triplicates.

Results are presented as stacked bar of the % of Titan cells (TCs) and regulars cells (RCs), *

p<0.0001 vs. control H99O.

1009

1010 **S7 Fig. Chr9 ploidy does not influence TCs generation.** A panel of 7 clinical isolates with partial

1011 Chromosome 9 duplication (left panel) was tested for its ability to generate TCs. Only H99O

1012 and AD2-06a exhibited increased cell sizes (middle panel). The % of TC was 67.9 % (431/667)

1013 for 2-06a, 32.1% (429/1339) for H99O, and 4.2% (51/1227), for Ug2459 (Khi2 compared to

1014 H99O, *p<0.0001) with the % TCs ratio to that produced in H99O shown in the right panel.

1015

S8 Fig. *PKR1* mutations influence median cell size

Strains with *PKR1* loss of function mutation showed a variable ability to produce TCs

depending on the resulting truncated proteins. The clinical isolate AD2-07 which did not

harbor the *PKR1* mutation was recovered from the CSF of an HIV-positive patient on day 13 of

amphotericin B treatment while AD2-06a was recovered from its initial CSF. The median cell

size of AD2-07 (5.9 µm [5.2-6.6]) was significantly decreased compared to H99O (7.7 µm [6.6-

9.2]) and 8.5 µm [7.0-13.0] for AD2-06a (p<0.0001). Except Bt156 (median of 7.7 µm [6.4-9.4]),

the others strains had a significantly decreased median size compared to H99O (p<0.0001),

6.9 µm [6.0-7.8] for 8-1 strain; 6.5 µm [5.8-7.1] for Bt77, 6.4 µm [5.7-7.2] for Bt117; and 5.4

µm [4.7-6.1] for Ug2462. Experiments were done in triplicate and pooled.

1026

SI material 1: TCs generation begins between 8 and 12 h and TCs are able to produce yeasts

with regular size.

Dynamic imaging of yeasts from the AD2-06a *C. neoformans* clinical isolate using the Nikon

Biostation IM. Yeasts were seeded in MM on a culture dish that was previously coated with

E1 at 2 mg/L. Images were taken every 5 min for 24 h at ×100 magnification using

transmitted light (white bar=10µm)

1033

1034 **S1 Table: Strains used in this study**

1035

1036 **S2 Table: Gene disrupted in AD2-06a but not in closely related isolated AD3-55a or AD3-**

1037 **41a**

1038

1039 **S3 Table: Clinical isolates with chromosome 9 ploidy variation**

1040

1041 **S4 Table: Strains harboring Pkr1 loss-of-function mutations used in this study**

      Figures and Tables

      Fig 1. *In vitro* protocol of titan cells generation (TCG protocol). The protocol followed several

      steps: (1) *C. neoformans* H99O from a frozen stock culture at -80°C was cultured on Sabouraud

      agar for 2-5 days; (2) Approximately 10^7 yeasts were then suspended in 10mL of liquid Yeast

      Peptone dextrose (YPD) and incubated under agitation (150 rpm) at 30°C for 22 h (stationary

      phase); (3) 1ml of the culture was then washed twice in minimal medium (MM), then 10^6

      yeasts were resuspended in 1mL of MM in a 1.5 mL Eppendorf® tube and incubated at 800

      rpm for up to 120 hours using an Eppendorf thermomixer; (4) A mixture of regular cells and

      of titan cells (TCs) was ready for analysis.

      Fig 2. Titan cells generated *in vitro* harbor the typical phenotype of titan cells produced *in*

      ***vivo*.**

      (**A**) Typical morphology of titan cells (TCs, white arrow) was observed (increased body size (a,

      > 10 µm), capsule (b), thickened cell wall, large central vacuole, and peripheral cell cytoplasm

      distribution) while the size of regular cells (RCs, black arrow) is <10 µm. (**B**) TCs were

      reproducibly generated in lab 1 using H99O and in two independent laboratories (lab 2 and

      lab 3) using their local H99O strain. Cell size was measured manually or by using the Icy

      software on pictures taken in bright field. Each dot represents an independent experiment

      (median [interquartile range, IQR] are presented); (**C**) Melanization was more intense in TCs

      than RCs (*p<0.0001) based on the calculation of the max - mean grey value/pixel of each

      melanin ghost measured (n=19 for TCs and n=531 for RCs) using the ImageJ in Icy software.

      Each dot represented an individual cells and boxes median and IQR (**D**) Capsule size measured

      after India ink staining was significantly larger in TCs than in RCs. Dots represent individual

cells, and boxes median and IQR for 250 cells each (*p<0.001); (**E**) Chitin was denser in TCs

than in RCs according to CFW fluorescence intensity/pixel/cell measured by Icy software after

CFW staining (0.01 µg/ml) at step4 of the TCG protocol (*p<0.0001). Dots represent individual

cells, and boxes median and IQR for 400 cells each (*p<0.001, pooled measurements from 3

independent experiments); (**F**) N-acetylglucosamine (GlcNAc), the monomer component of

chitin, was increased in TCs compared to RCs as measured by a biochemical method after

gamma-irradiation of the yeasts to remove the capsule, allowing a better separation of TCs

and RCs. Each dot represent result from independent experiments (n=7). Results are

presented as median and IQR (p<0.001). (**G**) The yeasts recovered at step4 of the TCG protocol

and analyzed by dot plots (FSC/SSC) using flow cytometry showed two populations FSC/SSC^{high}

and FSC/SSC^{low} representing TCs and RC, respectively; (**H**) DNA content analysis after

propidium iodide (PI) staining showed that the TCs (FSC/SSC^{high}) population harbored

increased PI fluorescence intensity from 2C to >4C as compared to the diploid control (AD7-

77 cultured in Sabouraud agar) while the RCs (FSC/SSC^{low}) population harbored a PI intensity

comparable to the haploid control (H99O cultured in Sabouraud agar).

1081

Fig 3. **Dynamics of TCs generation *in vitro*.** (**A**) Cell body size was measured from samples of

H99O culture (step 3 of the TCG protocol) withdrawn at specific times (H0 to H120) using

pictures taken in bright field and measured with the ICY software (mean number of yeasts

counted ± SD = 219 ± 67, representative of three experiments). Cell size increased starting at

H8 with some cells reaching the threshold of 10 µm (grey dashed line). Each dot represents a

single cell and the bars represent median and IQR; (**B**) TCs generation started between H8 and

H12 and reached a plateau at H24. Each dot represents the proportion of TCs in the

corresponding sample (3 independent experiments); (**C**) Pictures (x400 magnification) taken

overtime showing the progressive increased in cell body size and the appearance of a vacuole

typical of TCs in a large cell at H8 (white arrow) (scale bar 10 μm); (**D**) Cells stained with

calcofluor (CFW) at 0.1 μg/mL prior to incubation using the TCG protocol. CFW fluorescent

intensity is analyzed by flow cytometry in the initial (H0) and the resulting FSC/SSC^{high} (TCs)

and FSC/SSC^{low} (RCs) observed at H24. The initial (H0) (green line) and the H24 FSC/SSC^{high}

(blue line) populations harbored a high CFW fluorescence suggesting that they are mother

cells, with a higher fluorescence for the FSC/SSC^{high}, while two populations of high and low

(black star) CFW fluorescence intensity were observed for the H24 FSC/SSC^{low} cells (Left panel).

The right panel shows the size (FSC) and CFW fluorescence intensity of the yeast populations

at H0 (black density lines) and H24 (yellow density lines). The initial population

(CFW^{high}/FSC^{low}) evolved in two populations, one corresponding to daughter cells (RC,

CFW^{low}/FSC^{low}, white arrow), and the other one corresponding to TCs (CFW^{high}/FSC^{high}, black

arrow).

1103

Fig 4. TCs generation *in vitro* is impacted by various environmental conditions. (A) The

sequence of media used at steps 2 and 3 of the TCG protocol was crucial for TCs generation:

yeasts cultured in YPD and transferred to minimal medium (MM) produced significantly more

TCs (cells >10μm, dotted grey line) than yeasts cultured in MM or YPD and transferred in MM

or YPD, respectively; (B) Exposure to day light at step 3 had a positive impact on TCs generation

compared to incubation in the dark; (C) Raising the incubation temperature to 37°C at step 3

decreased TCs formation compared to 30°C; (D) Modification of the initial pH of the MM used

at step 3 modifies TCs formation with pH 5.5 being optimal while a more acidic (pH=4), a

neutral (pH=7) or an alkaline (pH=8.5) pH inhibited TCs formation; (E) The impact of hypoxia

was tested by physical (closed cap) and by chemical (COCl$_2$ in MM at 1 nM) method and

compared to normoxia (21% oxygen). Physically- and chemically-induced hypoxia enhances

the production of TCs compared to normoxia with a higher proportion of TCs in physically-

compared to chemically-induced hypoxia. All experiments were performed in triplicate and

pooled (mean cell counted ± SD =2305±1438). Median and IQRare shown in black for each

condition (* p<0.0001 vs. reference condition, grey circled dots). The percentages above each

condition represents the % of TCs observed.

1120

Fig 5. Titan cells generation *in vitro* is influenced by quorum sensing factors, host factors

and antifungal drugs. (**A**) The cell concentration at onset of step2 significantly modified TCs

generation with a maximal TCs formation using an initial concentration of 10^6 cells/mL, and an

abolition of TCs formation at 10^7 cells/mL; We then tested several factor at step 3 by adding

(**B**) vitamin B5 (pantothenic acid) at 125 µM which had no impact on cell size distribution; (**C**)

quorum sensing peptide 1 (Qsp1p, 10µM) which significantly decreased cell size; (**D**)

monoclonal anti-cryptococcal capsular polysaccharide antibodies E1 and 18B7 (at 166 µg/mL

in MM) which significantly decreased cell size, as did fetal calf serum (FCS, 5%) (**E**), and

phosphatidylcholine at 5mM (**F**). Fluconazole (**G**) and flucytosine (**H**) at concentration below

the MIC (4 mg/L) drastically impaired TCs formation with no TCs produced upon drug

exposure. (**I**) Step 1 was modified by subculturing H99O 8 times over one month on Sabouraud

agar alone (Sub8), or supplemented with 20mg/L CFW (Sub8+CFW) or with 32mg/L

fluconazole (Sub8+FLC). Compared to initial culture (0Sub), 8 sub-cultures (8Sub) decreased

significantly the cell size (** p<0.0001, vs. 0Sub control, grey circled dots). In addition, iterative

subcultures on CFW and FLC decreased and increased significantly the cell size compared to

the 8Sub control, respectively. Median and IQR are shown in black for each condition (*

p<0.0001 vs. 8Sub control, grey circled dots). The numbers above each condition represents

the % of TCs observed. The experiments were performed in triplicate and pooled (mean cell

counted ± SD = 2455±913).

1140

Fig 6. TCs generation is dependent on various genes *in vitro* and *in vivo*.

The different H99 strains harbored variable abilities to produce TCs compared to H99O (grey

bar) with high TCs producer (H99O, S, L) and low TCs producer (KN99α, H99W, H99 CMO18)

in vitro (**A**) and *in vivo* (**B**). *Sgf29Δ* and *lmp1Δ* deletion mutants show a decrease in TCs

generation in various H99 backgrounds *in vitro* (**C**) and *in vivo* (**D**) compared to H99O. The ratio

to the value obtained for H99O used as a calibrator in each experiment was calculated for

each strain. Bar represent mean ± SD (mean cell counted=600). Khi2 test was performed to

compare the experimental conditions to H99O, they were performed in triplicates and pooled

(*p<0.0001).

1150

Fig 7. Non-synonymous mutation of *PKR1* enhances TCs generation based on clinical isolates

analysis.

(**A**) Rim pathway is required for TCs generation *in vitro* in H99O (grey bar) and KN99a. (**B**) The

screening of 56 *C. neoformans* serotype A MATα clinical isolates identified isolates AD2-06a

(ratio=2.78) and AD2-02a (ratio=1.62) as TCs hyperproducers compared to H99O (grey bar);

(**C**) Phylogenic tree were estimated using RAxML under the GTRCAT model of 29 clinical

isolates. (**D**) Venn diagram representing the number of common or specific SNPs and Indel

between AD2-06a, AD2-07 and AD3-55a. One of the AD2-06a specific SNPs is a non-

synonymous mutation for *PKR1* (mutation Gly125fs); (**E**) Alignment of Pkr1 protein sequences

from selected clinical isolates including H99O and AD2-07 that both harbor a wild type

sequence. The Pkr1 domain architecture is formed by a dimerization domain in green (amino

acids 2 to 40) and two cAMP binding site based on INTERPRO model in red (amino acids 219

to 351 and 353 to 473). (**F**) Strains with *PKR1* loss-of-function mutation showed a variable

ability to produce TC, as compared to H99O (grey bar). The clinical isolate AD2-07, which

produces significantly less TCs was recovered from the CSF of an HIV-positive patient on day

13 of amphotericin B treatment while AD2-06a, which produced the highest proportion of TC,

was recovered from the initial CSF sample of the same patient. (**G**) The *pkr1Δ* mutant strain

increased TCs formation compared to the parental strain KN99α. (*$p < 0.0001$). The ratio to

H99O, used as a calibrator in each experiment, was calculated for each strain (panel A, B, F

and G) and results expressed as mean ±SD. Experiments A, F and G were done in triplicate, B

only once (screening). To compare the experimental conditions to H99O, Khi2 analysis was

performed (*$p < 0.0001$, **$p < 0.05$). Mean cell counted ± SD = 1165±528.

1173

1174   **Tables**

1175

1176   **Table 1: Genes affected by SNPs or Indels in the different H99 strains**

1177

Strains	Genes affected	Gene function	Genetic event
H99O	*CNAG_07634*	hypothetical	Deletion
	CNAG_04078	hypothetical	Insertion
	CNAG_06456	hypothetical	Insertion
H99S	*CNAG_07595*	hypothetical	Deletion/frameshift
	CNAG_12447	miscRNA	Insertion
	CNAG_12900	miscRNA	Insertion
H99L	CNAG_06392, *SGF29*	transcription factor (binds H3K4me2/3 and recruits histone deacetylation)	Deletion
KN99	CNAG_06392, *SGF29*	transcription factor (binds H3K4me2/3 and recruits histone deacetylation)	Deletion
H99W	*CNAG_06765, LMP1*	Hypothetical (involved in mating virulence and melanization)	Deletion/frameshift
H99 CMO18	*CNAG_06765, LMP1*	Hypothetical (involved in mating virulence and melanization)	Deletion/frameshift

1178

1179 **Table 2 – Alignment and SNP statistics of sequenced isolates using H99 as a reference**

1180

1181

Strain	MLST genotype	% of Assembly Covered by aligned reads	Alignment depth (X)	SNPs
AD3-55a	unique	100	133	28,070
AD2-06a	unique	100	129	28,202
AD2-07	unique	99	45	27,797
AD3-41a	unique	100	134	25,378
AD4-92a	A3/M3	100	132	11,906
AD4-47a	A4/M1	100	132	10,018
AD3-58	A4/M1	100	123	9,996
AD3-83a	A1/M1	100	166	172
AD2-99a	A1/M1	100	119	180
AD6-55a	related to A1/M1	100	108	12,440
AD6-54a	related to A1/M1	100	131	12,440
AD5-45a	related to A1/M1	100	142	12,417
AD1-95a	related to A1/M1	100	107	12,521
AD1-90a	related to A1/M1	100	122	12,511
AD5-67a	VNII	100	139	277,381
AD3-9a	VNII	100	111	277,417
AD3-11a	VNII	100	125	268,842
AD4-76a	Th	100	112	43,281
AD3-95a	Th	100	141	43,235
AD5-53a	A5/M5	100	130	42,291
AD2-82a	A5/M5	100	133	42,271
AD1-86a	A5/M5	100	105	43,778
AD5-39a	A4/M4	100	108	43,696
AD4-63a	A4/M4	100	129	43,736
AD2-04a	A4/M4	100	137	43,733
AD1-68a	A4/M4	100	132	43,778

1182

1183 **Table 3 : Clinical strains and mutations**

1184

Strains	Mutations
Ug2462	n.1333C>T|p.Arg445*|1333/1449
Bt117	n.1274_1280dupCTCTCCT|p.Asn428fs|1280/1449
Bt77	n.1293dupA|p.Arg432fs|1293/1449
8-1	n.878_911delCCGAGGGGAGCTCGTTTGGGGAGTTAGCGCTGAT|p.Ser293fs|911/1449
Bt58	n.742G>T|p.Glu248*|742/1449
AD2-06a	n.375_376delAG|p.Gly125fs|376/1449
Bt156	n.41dupA|p.Asp14fs|41/1449

1185

FIG1

1

2

3

4

H99O
Sabouraud

2-5 days
Room temp

10^7 cells
10 mL YPD

22 h
30°C, 150 rpm

10^6 cells
1mL of MM

24 h to 120 h
30°C, 800 rpm

Analysis

Fig2

FIG5

Fig6

Fig7