Natural selection on plasticity of thermal traits in a highly seasonal

environment

Leonardo D. Bacigalupe[1*], Aura M. Barria[1], Avia Gonzalez-Mendez[1], Manuel Ruiz-Aravena[2]

Juan D. Gaitan-Espitia[3,4], Mark Trinder[5] and Barry Sinervo[6*]

[1]Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de

Chile, Casilla 567, Valdivia, Chile

[2] School of Natural Sciences, College of Sciences and Engineering, University of Tasmania,

Hobart, Tasmania, Australia

[3] CSIRO Oceans and Atmosphere, GPO Box 1538, Hobart 7001, TAS, Australia

[4]The Swire Institute of Marine Science and School of Biological Sciences, The University of

Hong Kong, Hong Kong SAR, China

[5] MacArthur Green, 95 South Woodside Road, Glasgow, UK

[6]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA,

95064, USA

*Both authors contributed equally to this work

Correspondence should be addressed to L.D.B (+5663 2293567, lbacigal@gmail.com)

Abstract

For ectothermic species with broad geographical distributions, latitudinal/altitudinal

variation in environmental temperatures (averages and extremes) are expected to shape the

evolution of physiological tolerances and the acclimation capacity (i.e., degree of phenotypic

plasticity) of natural populations. This can create geographical gradients of selection in which

environments with greater thermal variability (e.g., seasonality) tend to favour individuals that

maximize performance across a broader range of temperatures compared to more stable

environments. Although thermal acclimation capacity plays a fundamental role in this context, it

is unknown whether natural selection targets this trait in natural populations. Here we addressed

such an important gap in our knowledge by measuring survival, through mark recapture

integrated into an information-theoretic approach, as a function of the plasticity of critical

thermal limits for activity, behavioural thermal preference and the thermal sensitivity of

metabolism in the northernmost population of the four-eyed frog *Pleurodema thaul*. Overall, our

results indicate that thermal acclimation in this population is not being targeted by directional

selection, although there might be signals of selection on individual traits. According to the most

supported models, survival decreased in individuals with less tolerance to cold when cold-

acclimated (probably because daily low extremes are frequent during the cooler periods of the

year) and increased with body size. However, in both cases, the directional selection estimates

were non-significant.

Keywords: Amphibians, natural selection, physiological plasticity, acclimation, *Pleurodema*

thaul, Atacama Desert

Introduction

It is well known that environmental temperature (T_a) is the abiotic factor with major

incidence in the evolution, ecology and physiology of most of the biodiversity in the planet

(Angilletta 2009 and references therein). The effects of T_a are particularly relevant for

ectotherms as their body temperature depends on T_a and therefore any change in T_a affects their

fitness and performance (e.g. behaviour, growth, reproduction, metabolism). This relationship

between performance and temperature has been described by a thermal performance curve (TPC)

(Huey & Berrigan 2001; Angilletta 2009) which has often been used to describe the thermal

ecology and evolution of ectotherms (Gilchrist 1995; Huey & Kingsolver 1989), their phenotypic

plasticity (Schulte et al. 2011), and to predict their responses to climate change (Clusella-Trullas

et al. 2011; Sinclair et al. 2016). The TPC is best captured by three parameters: a minimum

critical temperature (CT_{Min}), which represents T_a below which performance is minimum; a

maximum critical temperature (CT_{Max}), which represents T_a above which performance is also

minimum and an optimum temperature (T_{Opt}), which represents T_a at which performance is

maximum. Most of these parameters can exhibit geographic variation depending on the

particular environmental context (e.g., local climate) and genetic background of populations

(Gilchrist 1996; Kingsolver et al. 2004; Latimer et al. 2011). Furthermore, this geographic

variation has the potential to create gradients of selection for TPCs across the species distribution

(Kingsolver & Gomulkiewicz 2003) shaping thermal sensitivities, tolerances and thermal

acclimation capacities (i.e., thermal plasticity) of local populations (Seebacher et al. 2012;

Gaitan-Espitia et al. 2014).

Different climate-related hypotheses have been proposed to explain how physiological

tolerances, capacities and their plasticity affect the distributional ranges of species (Bozinovic et

al. 2011). One of them, the climate variability hypothesis (CVH), offers a powerful conceptual

framework to explore the interactions between environmental variability and physiological

performance of ectotherms (e.g., Gaitan-Espitia et al. 2013; 2014). The CVH predicts that

organisms inhabiting more variable environments should have broader ranges of environmental

tolerance and/or greater physiological plasticity that enable them to cope with the fluctuating

environmental conditions (e.g., seasonality) (Ghalambor et al. 2006; Gaitan-Espitia et al. 2017).

In agreement with this hypothesis, other theoretical models have explored the evolutionary

mechanisms underlying local thermal adaptation across heterogeneous environments (e.g.,

Generalist-Specialist models). For instance, the model developed by Lynch and Gabriel (1987),

predicts that temporal environmental heterogeneity selects for more broadly adapted individuals,

whereas in more constant environments the model developed by Gilchrist (1995), predicts that

selection should favor thermal specialists with narrow performance breadth. The mechanistic

understanding of these conceptual frameworks has improved with recent studies showing how in

thermally variable environments directional selection acts on TPC's parameters favoring

organisms that maximize performance across a broader range of temperatures (Logan et al. 2014)

despite the ability of ectotherms to thermoregulate behaviorally (Buckley et al. 2015).

Notwithstanding this progress, whether natural selection targets thermal acclimation capacity

(i.e., plasticity) itself in natural populations remains unknown.

In addition to increasing mean temperatures, it is known that climate change is changing

the frequency and intensity of extreme temperatures and events (Rahmstorf & Coumou 2011;

Wang & Dillon 2014; Vazquez et al. 2016). This, in turn, suggests that both averages and

variances will have an important impact on different performance related traits (e.g. Lardies et al.

2014; Vasseur et al. 2014; Bartheld et al. 2017). Nevertheless, we still do not know whether

selection might also target traits as a function of those extremes.

In this context, populations inhabiting highly seasonal environments characterized also by

daily extreme temperatures, provide a natural laboratory to evaluate the role of natural selection

on the plasticity of critical thermal limits and preferences. We addressed such important gaps in

our knowledge by measuring for the first time survival as a function of the plasticity of thermal

critical temperatures (CT_{Max} and CT_{Min}), preferred temperature (T_{Pref}) and thermal sensitivity of

metabolism (Q_{10}; the magnitude of change in metabolic rate for a 10°C change in T_a) after

acclimation to 10°C and 20°C in the northernmost population of the four-eyed frog *Pleurodema*

thaul. We tested four predictions regarding phenotypic selection and plasticity that built up from

previous findings showing that acclimation to warmer temperatures produces an increase in the

upper but not in the lower limits of the thermal performance curve (Ruiz-Aravena et al. 2014)

(Fig. 1). First, the high seasonality should select for plasticity in TPC parameters and therefore,

the plasticity itself should currently be under directional selection. Second, if daily high extreme

temperatures were frequent, then it would be expected positive directional selection on CT_{max}

when warm as well as cold-acclimated. Third, if daily low extremes were frequent then it would

be expected negative directional selection on CT_{min} during the cooler periods of the year. Finally,

as energy inputs are limited, the energetic definition of fitness indicates that individuals with

higher maintenance costs (i.e. resting metabolic rate) would have less energy available to

allocate to growth, reproduction and/or performance. The main prediction of this principle is that

natural selection should maximize the residual available energy, and therefore, higher

maintenance costs would be associated with lower fitness if no compensations in other functions

were available (Bacigalupe & Bozinovic 2002; Artacho & Nespolo 2009). Thus, our final

prediction is that Q_{10} is not under directional selection.

METHODS

Study organism and laboratory maintenance

Eighty-three adults individuals of *P. thaul* were captured during September 2012 on two

small ponds at Carrera Pinto (27º06'40.2'' S, 69º53'44.3'' W; 2,000 m.a.s.l.), a small oasis in the

Atacama Desert that is known to be the northernmost population of the species (Correa et al.

2007). In both ponds, we performed an exhaustive search across microhabitats (below rocks, in

the vegetation and in the water). All individuals were transported to the laboratory (Universidad

Austral de Chile, Valdivia) within 2 – 3 days of capture. Following capture all animals were

marked by toe clipping and maintained in the laboratory for one month at a temperature of 20º ±

2ºC and with a photoperiod 12D:12L. Animals were housed (N = 5) in terrariums (length x width

x height: 40 x 20 x 20 cm) provided with a cover of moss and vegetation and a small recipient

filled with water. Individuals were fed once a week with mealworms (*Tenebrio sp*. larvae) and

Mazuri® gel diets.

Acclimation and thermal traits

After one month at maintenance conditions, in a split cross design half the frogs were

acclimated to either 10°C or 20°C for two weeks before measuring thermal traits. Frogs were

randomly assigned to the first acclimation temperature using a coin. Next they were acclimated

to the other temperature and again measured thermal traits. We chose these acclimation

temperatures because they are close to the mean minimum temperatures during the breeding

season (August - October, 10ºC) and to the mean temperatures during the active period of the

species (20ºC) at Carrera Pinto (www.cr2.cl). None of the investigators were blinded to the

group allocation during the experiments.

Critical temperatures were determined as the environmental temperature at which an

individual was unable to achieve an upright position within 1 minute (Ruiz-Aravena et al. 2014).

Each individual was placed in a small chamber inside a thermo-regulated bath (WRC-P8,

Daihan, Korea) at 30°C (CT_{Max}) and 5ºC (CT_{Min}) for 15 minutes, after which the bath

temperature was increased (or decreased) at a rate of 0.8ºC per minute (Rezende et al. 2011).

Every minute or at every change in 1ºC, the chamber was turned upside down and we observed if

the animal was able to return to the upright position. When an animal was unable to achieve an

upright position within 1 minute it was allowed to recover at ambient temperature (CT_{Min}) or for

30 minutes in a box with ice packs (CT_{Max}). Body mass (a proxy of body size) was obtained

before each trial using a Shimadzu TX323L electronic balance.

Preferred temperature (T_{Pref}) was determined simultaneously for five individuals in five

open-top terraria (length x width x height: 85 x 12 x 30 cm). Each terrarium had a thermal

gradient between 10ºC and 30ºC produced by an infrared lamp overhead (250 W) on one end,

and ice packs on the other. The organic gardening soil was moisturized at the beginning of each

trial to prevent the desiccation of the frogs. Five individuals were placed at the centre of each one

of the terraria and 45 minutes later we registered T_{Pref} as the dorsal body temperature (T_b) using a

UEi INF155 Scout1 infrared thermometer. Dorsal and cloacal T_b are highly associated (r_P =

0.99) (see Ruiz-Aravena et al. 2014 for details). Body mass was obtained before each trial using

a Shimadzu TX323L electronic balance.

Standard metabolic rate, measured through oxygen consumption at 20°C and 30°C was

measured continuously using an infrared O_2 - CO_2 analyzer (LI-COR LI6262, Lincoln, NV,

USA). The analyzer was calibrated periodically against a precision gas mixture. Although there

was almost no difference between calibrations, baseline measurements were performed before

and after each recording. Flow rates of CO2 – free air was maintained at 100 ml $min^{-1} \pm 1\%$ by a

Sierra mass flow controller (Henderson, NV, USA). We used cylindrical metabolic chambers (60

ml), covered by metal paper. O_2 consumption was recorded during 45 minutes per individual.

Each record was automatically transformed by a macro program recorded in the ExpeData

software (Sable Systems), to (1) transform the measure from % to mlO_2 min^{-1}, taking into

account the flow rate and (2) to eliminate the first 5 min of recordings. For each individual, the

metabolic sensitivity (Q_{10}) was calculated as the ratio between metabolic rate measured at 30ºC

and metabolic rate measured at 20ºC.

Selection on thermal traits

After experiments, all frogs were put back to 20ºC for at least one month before releasing

them. Marked frogs were released at Carrera Pinto in April 2013 and their survival was

monitored on three separate recapture efforts (13th October 2013, 13th June and 9th September

2014). As the desert surrounds these two small ponds dispersal was not a concern.

The relationship between trait plasticity and survival was analyzed using a Cormack-Jolly-Seber

(CJS) framework in Program MARK. An overall goodness of fit test was run using U-Care to

ensure the data were consistent with the assumed structure of the CJS model and to obtain a

value for the over dispersion parameter (c-hat). The time interval between capture occasions (as

a fraction of 1 year and considering also the original capture event) was included in the analysis

to accommodate the unequal intervals. The resulting resighting and survival estimates were

therefore corrected to annual estimates. Survival and resighting parameters were obtained in a

two-stage process. First, the best-fit resighting model was identified from three candidate models

(constant, time dependent and a linear trend). The fit of the three candidate resighting models

was compared using survival modeled as both a constant rate and also as a time-dependent rate,

to ensure that selection of the best-fit resighting model was not influenced by choice of survival

model. Once the best-fit resighting model had been identified (using AICc) this was then

retained for all candidate models. A model selection and an information-theoretic approach

(Burnham & Anderson 2003) was employed to contrast the adequacy of different working

hypotheses (the candidate models) of selection on trait plasticity. The number of candidate

models was kept to a minimum to minimize the likelihood of spurious results (Burnham &

Anderson 2003; Lucaks et al. 2010). Body mass showed a positive relationship with CT_{Max_20} (r_P

= 0.47) and with T_{Pref_10} (r_P = 0.24) and was not associated with any other trait (results not

shown). Therefore, we tested only for a null model (i.e. neither trait under selection), a model

with body mass and models with directional selection for each trait separately and also for

correlational selection (interaction of trait combinations) among the same trait at both

acclimation temperatures, which indicates plasticity. Body mass was included as a covariate in

the case of CT_{Max_20} and T_{Pref_10} (Table 1). All analyses were performed in R version 3.1.3

employing package RMark (Laake 2013). No transformation was required to meet assumptions

of statistical tests. Survival in relation to each covariate was obtained as the model averaged

value across all candidate models weighted by individual model probability (Table 1).

RESULTS

All measured traits including critical thermal limits (CT_{Max}, CT_{Min}), thermal preference

(T_{Pref}) and sensitivity of metabolic rate to temperature (Q_{10}) showed high variance among

individuals (Fig. 2). In addition, for all traits some individuals shifted their thermal traits to

higher values when acclimated to high temperatures, but other individuals showed the reverse

response, that is their traits shifted to lower values after acclimation at higher temperatures (Fig.

3).

Only five out of 28 correlations between physiological traits were statistically significant,

and these involved mostly critical thermal limits. In particular CT_{Max_20} was negatively correlated

with CT_{Min_10} ($r_P = -0.57$) and CT_{Max_10} ($r_P = -0.41$) whilst it was positively correlated with Q_{10_20}

($r_P = 0.26$). Additionally, CT_{Max_10} was positively correlated with CT_{Min_10} ($r_P = 0.31$) and

negatively correlated with CT_{Min_20} ($r_P = -0.25$).

The overall goodness of fit measure for the CJS model indicated a moderate level of

over-dispersion (c-hat $= 2.65$, $P = 0.103$), however with only 3 recapture occasions it was not

possible to identify an alternative starting model and the basic CJS model was adopted as the

basis for subsequent model fitting, with unexplained over-dispersion controlled using the c-hat

adjustment. A constant resighting rate was the best-fit model irrespective of whether survival

was modeled as a constant or time dependent rate (Table 1). Consequently, the constant rate-

resighting model was retained for subsequent modeling of survival. The model selection

procedure indicated that from the 13 candidate models tested, there was not a single best-fit one

(Table 1). In particular, the null model was the most supported (Akaike weight of 0.220), whilst

models including only directional selection on single traits still had some support, with a

cumulative Akaike weight of almost 60% (Table 1). Models including correlational selection

(i.e. plasticity) showed rather weak empirical support (Table 1). Overall, survival decreased as

values of most of the traits increased in both, warm and cold acclimated conditions (Table 2,

Figure 4).

DISCUSSION

To understand how organisms adapt to highly fluctuating environments and whether they will be

able to adaptively respond to current climate change, we need to evaluate whether selection in

nature targets plasticity itself. Populations inhabiting highly seasonal environments that also

experience daily extreme temperatures, provide excellent opportunities to test predictions of the

fitness consequences of such thermal variation on the plasticity of critical thermal limits and

preferences. Here, to the best of our knowledge for the first time, we studied natural selection on

thermal acclimation capacity of performance (CT_{Max} and CT_{Min}), metabolism (Q_{10}) and

behaviour (T_{Pref}). Our results indicate that thermal acclimation in this population is not being

targeted by directional selection, although there might be signals of selection on individual traits.

In part, the relatively weak evidence for natural selection on this system might be a consequence

of the small sample size we used (N = 83), the few recaptures we carried out (n = 3) and the

relatively high value of c-hat in the analyses, which penalizes models on the basis of parameter

number. This prevented us not only from evaluating more complex models (i.e. non linear

selection) but also resulted in estimates of directional selection with rather large SEs and

therefore with 95% confidence intervals that contained the zero in all cases.

Some theoretical models of thermal adaptation across heterogeneous environments (e.g.,

Climate variability hypothesis, generalist-specialist models), suggest that temporal

environmental heterogeneity selects for more broadly adapted individuals (Lynch and Gabriel,

1987; Gilchrist 1995), favoring increased plasticity particularly regarding thermal tolerance traits

(Gunderson & Stillman 2015). Based on these models we predicted that the high seasonality

should select for high plasticity in thermal traits and therefore, the plasticity itself should

currently be under directional selection. Our prediction turned out to be incorrect as models

including plasticity showed relatively weak support.

Frogs of *P. tahul* in the Atacama Desert (the northernmost population of this species) are

exposed to large daily and seasonal oscillations in environmental temperatures. The isothermality

(i.e., ratio between daily and annual thermal ranges; O'Donnell & Ignizio 2012) experienced by

this population (0.65) is ca. 15% higher compared to a population 2,000 km south (0.52), which

experiences narrower daily environmental temperatures at the center of the species' distribution

(Barria & Bacigalupe 2017). This means that the studied population experiences a daily variation

that is almost 65% of its seasonal variation. The high variation on a daily basis together with the

fact that climate change is already changing the frequency and intensity of extreme temperatures

(Rahmstorf & Coumou 2011; Wang & Dillon 2014; Vasquez et al. 2017) made us wonder

whether selection in nature might also target traits as a function of daily extremes. As the cooler

end of the thermal performance curve did not change trough acclimation to warmer temperatures

(Ruiz-Aravena et al. 2014) we predicted negative directional selection on CT_{min} during the cooler

but not the warmer periods of the year. Our results are in agreement with the trend specified by

this prediction, as survival decreased as CT_{min} increased (i.e. less tolerance to cold) when cold-

acclimated (albeit the estimate was non-significant), which was the second most supported

model (Table 1).

Although acclimation produced an increase in the upper limits of the thermal

performance curve in this population (Ruiz-Aravena et al. 2014), we predicted positive

directional selection on CT_{max} when warm as well as cold-acclimated if daily high extreme

temperatures were frequent. Our results do not offer support for this prediction: there was a slight

trend for survival to decrease as CT_{max} increased under warm as well as under cold-acclimated

conditions. However, in both cases estimates were not statistically different from zero.

Nevertheless, this might suggests that selection could be favouring individuals that avoid hot

microhabitats, possibly by means of behaviour (Ruiz-Aravena et al. 2014). Indeed, behavioural

thermoregulation has been proposed as one key factor that prevents an evolutionary response to

selection to raising temperatures (Kearney et al. 2009; Huey et al. 2012; Buckley et al. 2015).

The fact that CT_{Max_20} was negatively correlated with CT_{Min_10} indicates that individuals with

higher cold tolerance might be the ones avoiding hot microhabitats, which opens very interesting

questions for further research.

Regarding the sensitivity of metabolism to temperature (Q_{10}) we predicted that Q_{10} is not

under directional selection. Our results (partially) supported that prediction as the rate at which

survival changed with changes in Q_{10} was very small (Fig. 4, Table 2), although the models with

Q_{10} still showed some support (Table 1). Finally, we also predicted no directional selection on

T_{Pref} as we have previously shown that acclimation to warmer temperatures produced an increase

in this trait (Ruiz-Aravena et al. 2014). Nevertheless, we found a non-significant trend showing

that survival decreased, although at a very low rate, as T_{Pref} increased, which might suggest that

selection favours those individuals that are able to avoid hot microhabitats.

Our results indicate a positive trend of survival with body size (although the directional

selection estimate was non-significant), something that has been previously reported in the

literature (Aubin-Horth et al. 2005; Iida & Fujisaki 2007; Crosby & Latta 2013; Delaney &

Warner 2017). This is somewhat unsurprising, given that body mass is known to be positively

associated with several physiological traits that enhance performance (Castellano et al. 1999;

Madsen & Shine 2000; Hurlbert et al. 2008; Shepherd et al. 2008; Luna et al. 2009) including

plasticity itself (Whitman & Ananthakrishnan 2009). Although this population inhabits two

highly isolated ponds where the presence of competitors (anurans) has not been observed, there

might be a risk of predation by herons (*L.D.B. personal observation*), and thus, a positive

selection for body size might explain it. Nevertheless, further experimental work is needed to

evaluate this possibility.

It is important to mention that we have measured plasticity in just one life stage. It is

likely that other ecological and physiological traits might also be plastic, and their responses to

acclimation might be different and they might even be different between different life stages.

However, we still consider our results show a signal and provide the first evidence that

phenotypic plasticity is not being an actual target of selection in nature and that daily climate

extremes might be selecting for higher tolerance. Nevertheless, further work including multiple

traits and life stages and also in other populations might help unify trends into further generic

hypotheses to clarify the role of plasticity in the viability of ectotherm populations in natural

conditions.

Acknowledgements. We thank Nadia Aubin-Horth, Wolf Blanckenhorn, Dries Bonte, Ray Huey

and Michael Logan for highly valuable comments on a previous version on the manuscript.

Data: data is available for download from the CSIRO Data Access Portal (https://data.csiro.au)

Competing interests. We declare we have no competing interests

Author Contributions. L.D.B conceptualized the study, designed the experimental procedures

and carried out the experiment with A.M.B., A.G.M., M.R.A. and J.D.G.E; M.T., B.S. and L.D.

B. analyzed the data and L.D.B., B.S. and J.D.G. wrote the paper with input from A.M.B and

M.R.A.

Funding. Leonardo Bacigalupe acknowledges funding from FONDECYT grant 1150029. Barry

Sinervo was supported by a Macrosystems grant (EF-1241848) from NSF. Aura Barria and

Manuel Ruiz-Aravena were supported by a CONICYT Doctoral Fellowship.

Ethics. This study did not involve endangered or protected species and was carried out in strict

accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals

of the Comisión Nacional de Investigación Científica y Tecnológica de Chile (CONICYT). All

experiments were conducted according to current Chilean law. The protocol was approved by the

Committee on the Ethics of Animal Experiments of the Universidad Austral de Chile.

332
333

References

Angilletta, M. J. 2009. Thermal Adaptation: a theoretical and empirical synthesis. Oxford, UK: Oxford University press.

Artacho, P. & Nespolo, R. F. 2009 Natural selection reduces energy metabolism in the garden snail, Helix aspersa (*Cornu aspersum*). *Evolution* **63**, 1044–50.

Aubin-Horth, N., Ryan, D. A. J., Good, S. P. & Dodson, J. J. 2005 Balancing selection on size: Effects on the incidence of an alternative reproductive tactic. *Evol. Ecol. Res.* **7**, 1171–1182.

Bacigalupe, L. D. & Bozinovic, F. 2002 Design, limitations and sustained metabolic rate: lessons from small mammals. *J. Exp. Biol.* **205**, 2963–2970.

Barria, A. M. & Bacigalupe, L. D. 2017 Intraspecific geographic variation in thermal limits and acclimatory capacity in a wide distributed endemic frog. *J. Therm. Biol.* **69**, 254–260.

Bartheld, J. L., Artacho, J. P. & Bacigalupe, L. D. 2017. Thermal performance curves under daily thermal fluctuation: A study in helmeted water toad tadpoles. *J. Therm. Biol* 70, 80-85.

Bozinovic, F., Calosi, P. & Spicer, J. I. 2011. Physiological correlates of geographic range in animals. *Annu. Rev. Ecol. Evol. Syst.* **42**, 155-179.

Buckley, L. B., Ehrenberger, J. C. & Angilletta, M. J. 2015. Thermoregulatory behaviour limits local adaptation of thermal niches and confers sensitivity to climate change. *Funct. Ecol.* **29**, 1038–1047.

Burnham, K. P. & Anderson, D. R. 2003 *Model selection and multimodel inference: a practical information-theoretic approach.* Springer Science & Business Media.

Castellano, S., Rosso, A., Doglio, S. & Giacoma, C. 1999. Body size and calling variation in the green toad (*Bufo viridis*). *J. Zool.* **248**, 83–90.

Clusella-Trullas, S., Blackburn, T. M. & Chown, S. L. 2011. Climatic Predictors of Temperature Performance Curve Parameters in Ectotherms Imply Complex Responses to Climate Change. *Am. Nat.* **177**, 738-751.

Correa, C., Sallaberry, M., Gonzalez, B., Soto, E. & Mendez, M. 2007 Amphibia, Anura, Leiuperidae, Pleurodema thaul: Latitudinal and altitudinal distribution extension in Chile. *Check List* **3**, 267–270

Crosby, K. & Latta, R. G. 2013 A test of the reproductive economy hypothesis in plants: More offspring per capita come from large (not small) parents in *Avena barbata. Evol. Ecol.* **27**, 193–203.

380   Delaney, D. M. & Warner, D. A. 2017 Effects of age- and sex-specific density on behaviour and
381   survival in a territorial lizard (*Anolis sagrei*). *Anim. Behav.* **129**, 31–41.
382
383   Gaitan-Espitia, J. D., Arias, M. B., Lardies, M. A. & Nespolo, R. F. 2013. Variation in thermal
384   sensitivity and thermal tolerances in an invasive species across a climatic gradient:
385   lessons from the land snail *Cornu aspersum. PLoS One, 8*(8), e70662
386
387   Gaitan-Espitia, J. D., Bacigalupe, L. D., Opitz, T, Lagos, N. A., Timmermann, T. & Lardies, M.
388   A. 2014. Geographic variation in thermal physiological performance of the intertidal crab
389   *Petrolisthes violaceus* along a latitudinal gradient. *J. Exp. Biol.* **217**, 4379-4386.
390
391   Gaitan-Espitia, J. D., Marshall, D. J., Dupont, S., Bacigalupe, L. D., Bodrossy, L. & Hobday, A.
392   J. 2017. Geographical gradients in selection can reveal genetic constraints for
393   evolutionary responses to ocean acidification. *Biol. Lett.* **13**, 20160784.
394
395   Ghalambor, C. K., Huey, R. B., Martin, P. R., Tewksbury, J. T. & Wang, G. 2006. Are mountain
396   passes higher in the tropics? Janzen's hypothesis revisited. *Integr. Comp. Biol.* **46**, 5–17.
397
398   Gilchrist, G. W. 1995. Specialists and generalists in changing environments. 1. Fitness
399   landscapes of thermal sensitivity. *Am. Nat.***146**:252–270.
400
401   Gilchrist, G. W. 1996. A quantitative genetic analysis of thermal sensitivity in the locomotor
402   performance curve of *Aphidius ervi. Evolution* **50**, 1560–1572.
403
404   Gunderson, A. R. & Stillman, J. H. 2015. Plasticity in thermal tolerance has limited potential to
405   buffer ectotherms from global warming. Proc. *R. Soc. B* **282**, 20150401
406
407   Huey, R. B. & Kingsolver, J. G. 1989. Evolution of thermal sensitivity of ectotherm
408   performance. *Trends Ecol. Evol.* **4**, 131-135.
409
410   Huey, R. B. & Berrigan, D. 1996 Testing evolutionary hypotheses of acclimation. In *Phenotypic*
411   *and evolutionary adaptation to temperature* (eds A. Johnston & A. Bennett), pp. 205–
412   237. Cambridge University Press
413
414   Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M. & Williams, S. E.
415   2012. Predicting organismal vulnerability to climate warming: roles of behaviour,
416   physiology and adaptation. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 1665–1679.
417
418   Hurlbert, A. H., Ballantyne IV, F. & Powell, S. 2008 Shaking a leg and hot to trot: The effects of
419   body size and temperature on running speed in ants. *Ecol. Entomol.* **33**, 144–154.
420
421   Iida, H. & Fujisaki, K. 2007 Seasonal changes in resource allocation within an individual
422   offspring of the wolf spider, Pardosa pseudoannulata (Araneae: Lycosidae). *Physiol.*
423   *Entomol.* **32**, 81–86.
424
425   Kingsolver, J. G. & Gomulkiewicz, R. 2003. Environmental Variation and Selection on

426   Performance Curves. *Integr. Comp. Biol.* **43**, 470–477.
427

428   Kearney, M., Shine, R. & Porter, W. P. 2009 The potential for behavioral thermoregulation to
429   buffer 'cold-blooded' animals against climate warming. *Proc. Natl. Acad. Sci. U. S. A.*
430   **106**, 3835–3840.
431

432   Kingsolver, J. G., Ragland, G. J. & Shlichta, J. G. 2004. Quantitative genetics of continuous
433   reaction norms: thermal sensitivity of caterpillar growth rates. *Evolution* **58,**1521-1529.
434

435   Laake, J. 2013 RMark: an R interface for analysis of capture–recapture data with MARK. In
436   *AFSC Processed Rep 2013-01*, pp. 25. Seattle, Washington, USA.: Alaska Fisheries
437   Science Center, NOAA, National Marine Fisheries Service
438

439   Lardies, M. A., Arias, M. B., Poupin, M. J. & Bacigalupe, L. D. 2014. Heritability of *hsp70*
440   expression in the beetle *Tenebrio molitor*: Ontogenetic and environmental effects. *J.*
441   *Insect Physiol.* **67**, 70-75.
442

443   Latimer, C. A. L., Wilson, R. S. & Chenoweth, S. F. 2011. Quantitative genetic variation for
444   thermal performance curves within and among natural populations of *Drosophila serrata.*
445   *J. Evol. Biol.* **24**, 965–975.
446

447   Logan, M. L., Cox, R. M. & Calsbeek, R. 2014 Natural selection on thermal performance in a
448   novel thermal environment. *Proc. Natl. Acad. Sci. U. S. A.* **39**, 14165-14169.
449

450   Lukacs, P., Burnham, K. & Anderson, D. 2010 Model selection bias and Freedman's paradox.
451   *Ann. Inst. Stat. Math.* **62**, 117–125
452

453   Luna, F., Antenucci, C. D. & Bozinovic, F. 2009 Comparative energetics of the subterranean
454   Ctenomys rodents: breaking patterns. *Physiol. Biochem. Zool.* **82**, 226–35.
455

456   Lynch, M., and W. Gabriel. 1987. Environmental tolerance. *Am. Nat.* **129**:283–303.
457

458   Madsen, T. & Shine, R. 2000 Energy versus risk: costs of reproduction in free-ranging pythons
459   in tropical Australia. *Austral Ecol.* **25**, 670–675.
460

461   O'Donnell, M. S. & Ignizio, D. A. 2012. Bioclimatic predictors for supporting ecological
462   applications in the conterminous United States. Geological Survey Data Series 691.
463

464   Schulte, P. M., Healy, T. M. & Fangue, N. A. 2011.Thermal Performance Curves, Phenotypic
465   Plasticity, and the Time Scales of Temperature Exposure. *Integr. Comp. Biol.* **51**, 691–
466   702.
467

468   Rahmstorf, S. & D. Coumou. 2011. Increase of extreme events in a warming world. *Proc. Natl.*
469   *Acad. Sci. U. S. A.* **108**,17905-17909.
470

471   Rezende, E. L., Tejedo, M. & Santos, M. 2011 Estimating the adaptive potential of critical

472   thermal limits: methodological problems and evolutionary implications. *Funct. Ecol.* **25**,
473   111–121.
474
475   Ruiz-Aravena, M., Gonzalez-Mendez, A., Estay, S. A., Gaitán-Espitia, J. D., Barria-Oyarzo, I.,
476   Bartheld, J. L. & Bacigalupe, L. D. 2014 Impact of global warming at the range margins:
477   phenotypic plasticity and behavioral thermoregulation will buffer an endemic amphibian.
478   *Ecol. Evol.* **4**, 4467–4475.
479
480   Seebacher, F., Holmes, S., Roosen, N. J., Nouvian, M., Wilson, R. S. & Ward, A. J. W. 2012.
481   Capacity for thermal acclimation differs between populations and phylogenetic lineages
482   within a species. *Funct. Ecol.* **26**, 1418–1428.
483
484   Shepherd, B. L., Prange, H. D. & Moczek, A. P. 2008. Some like it hot: Body and weapon size
485   affect thermoregulation in horned beetles. *J. Insect Physiol.* **54**, 604–611.
486
487   Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., Dong,
488   Y., Harley, C. D. G., Marshall, D. J., Helmuth, B. S. & Huey, R. B. 2016. Can we predict
489   ectotherm responses to climate change using thermal performance curves and body
490   temperatures? *Ecol. Lett.* **19**, 1372–1385.
491
492   Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D. G., McCann, K. S.,
493   Savage, V., Tunney, T. D. & O'Connor, M. I. 2014. Increased temperature variation
494   poses a greater risk to species than climate warming. *Proc. R. Soc. B*. **281**, 20132612.
495
496   Vázquez, D. P., Gianoli, E., Morris, W. F. & Bozinovic, F. 2017. Ecological and evolutionary
497   impacts of changing climatic variability. *Biol. Rev.* **92**, 22–42.
498
499   Wang, G. & Dillon, M. E. 2014. Recent geographic convergence in diurnal and annual
500   temperature cycling flattens global thermal profiles. *Nature Climate Change* **4**, 988-992.
501
502   Whitman, D. & Ananthakrishnan, T. 2009 *Phenotypic Plasticity of Insects: Mechanisms and
503   Consequences*. Enfield: Science Publishers.
504

Figure 1. Graphical representation of the predictions tested in this study. (a) Predictions built up
from findings showing that acclimation to warmer temperatures produces an increase in the
upper but not in the lower limits of the thermal performance curve (Ruiz-Aravena et al. 2014).
(b) The high seasonality should have selected for plasticity and therefore, plasticity of all traits
should currently be under directional selection. (c) If daily low extremes are frequent, we expect
negative directional selection on CT_{Min} during the cooler periods of the year (left panel). If daily
high extreme temperatures are frequent, we expect positive directional selection on CT_{Max} during
the warmer periods (right panel) as well as the cooler periods of the year (middle panel). We
predict no directional selection on T_{Pref} and Q_{10} at both acclimation temperatures and on CT_{Min}
when warm acclimated. Cold acclimated is indicated by a _10 subscript while warm acclimated
is indicated by a _20 subscript.

Figure 2. Frequency distribution of CT_{Min}, T_{Pref} and CT_{Max} of the four-eyed frog when
acclimated to 10ºC and 20ºC.

 Figure 3. Individual plasticity in CT_{Min}, T_{Pref}, CT_{Max} and Q_{10} to 10 and 20ºC acclimation
treatments. Each line represents the individual value of the specific traits at each temperature.
For CT_{Min} and CT_{Max} the width of the line is directly proportional to the number of individuals
that showed that specific response.

525
526

Figure 4. Survival estimates of simple directional selection sorted by model probabilities.

w: Akaike weights of the model.

Table 1. Candidate models ordered accordingly to their Akaike weights. Single term models
represent directional (e.g. CT_{Max}) and correlational selection represents plasticity (e.g. CT_{Max_10} *
CT_{Max_20}). CT_{Min} = minimum critical temperature; CT_{Max} = maximum critical temperature; T_{Pref} =
preferred temperature; Q_{10} = thermal sensitivity of metabolism; MB = body mass. Cold
acclimated is indicated by a _10 subscript while warm acclimated is indicated by a _20 subscript.
535

Models		K	AICc	ΔAICc	w_i
1	Null model	2	130.17	0	0.220
2	CT_{Min_10}	3	131.40	1.23	0.119
3	MB	3	131.78	1.61	0.098
4	T_{Pref_20}	3	132.08	1.90	0.085
5	Q_{10_10}	3	132.18	2.01	0.081
6	CT_{Min_20}	3	132.25	2.08	0.078
7	CT_{Max_10}	3	132.26	2.08	0.078
8	Q_{10_20}	3	132.26	2.09	0.077
9	CT_{Min_10} + CT_{Min_20} + CT_{Min_10} * CT_{Min_20}	5	133.38	3.21	0.044
10	MB + CT_{Max_20}	4	133.44	3.27	0.043
11	MB + T_{Pref_10}	4	133.82	3.64	0.036
12	Q_{10_10} + Q_{10_20} + Q_{10_10} * Q_{10_20}	5	134.17	4.00	0.030
13	MB + T_{Pref_10} + T_{Pref_20} + T_{Pref_10} * T_{Pref_20}	6	137.16	6.99	0.007
14	MB + CT_{Max_10} + CT_{Max_20} + CT_{Max_10} * CT_{Max_20}	6	137.62	7.45	0.005

K = number of parameters.
AICc: AIC values corrected for small sample sizes.
w_i: Akaike weights.

Table 2. Directional selection estimates from single terms models with their standard errors (SE)
and 95% confidence intervals (95% CI). CT_{Min} = minimum critical temperature; CT_{Max} =
maximum critical temperature; T_{Pref} = preferred temperature; Q_{10} = thermal sensitivity of
metabolism; MB = body mass. Cold acclimated is indicated by a _10 subscript while warm
acclimated is indicated by a _20 subscript.
545

Trait	Estimate	SE	95% CI
MB	0.209	0.212	-0.206 – 0.625
CT_{Min_10}	-0.248	0.187	-0.616 – 0.119
CT_{Min_20}	-0.030	0.181	-0.384 – 0.324
T_{Pref_10}	-0.025	0.059	-0.140 – 0.090
T_{Pref_20}	-0.026	0.042	-0.109 – 0.056
CT_{Max_10}	0.026	0.257	-0.477 – 0.530
CT_{Max_20}	-0.192	0.195	-0.575 – 0.191
Q_{10_10}	-0.475	1.140	-2.709 – 1.759
Q_{10_20}	-0.048	0.795	-1.607 – 1.510

Figure 1. Graphical representation of the predictions tested in this study. (a) Predictions built up from findings showing that acclimation to warmer temperatures produces an increase in the upper but not in the lower limits of the thermal performance curve (Ruiz-Aravena et al. 2014). (b) The high seasonality should have selected for plasticity and therefore, plasticity of all traits should not currently be under directional selection. (c) If daily low extremes are frequent, we expect negative directional selection on CT_{Min} during the cooler periods of the year (left panel). If daily high extreme temperatures are frequent, we expect positive directional selection on CT_{Max} during the warmer periods (right panel) as well as the cooler periods of the year (middle panel). We predict no directional selection on T_{Pref} and Q_{10} at both acclimation temperatures and on CT_{Min} when warm acclimated. Cold acclimated is indicated by a _10 subscript while warm acclimated is indicated by a _20 subscript.

Figure 2. Frequency distribution of CT_{Min}, T_{Pref} and CT_{Max} of the four-eyed frog when acclimated to 10ºC and 20ºC.

Acclimation 10ºC

Acclimation 20ºC

Figure 3. Individual plasticity in CT_{Min}, T_{Pref}, CT_{Max} and Q_{10} to 10 and 20ºC acclimation treatments. Each line represents the individual value of the specific traits at each temperature. For CT_{Min} and CT_{Max} the width of the line is directly proportional to the number of individuals that showed that specific response.

Figure 4. Survival estimates of simple directional selection sorted by model

probabilities.

w: Akaike weights of the model.