Full title: *Pseudomonas aeruginosa* type IV minor pilins and PilY1 regulate virulence by

modulating FimS-AlgR activity

Short title: Minor pilins regulate virulence via modulation of FimS-AlgR activity

Authors: Victoria A. Marko, Sara L.N. Kilmury, Lesley T. MacNeil, and Lori L. Burrows*.

Affiliations: Department of Biochemistry and Biomedical Sciences and the Michael G.

DeGroote Institute for Infectious Diseases Research, McMaster University, Hamilton, ON,

Canada.

*For correspondence: burrowl@mcmaster.ca

10   Abstract

Type IV pili are expressed by a wide range of prokaryotes, including the opportunistic

pathogen *Pseudomonas aeruginosa*. These flexible fibres mediate twitching motility, biofilm

maturation, surface adhesion, and virulence. The pilus is composed mainly of major pilin

subunits while the low abundance minor pilins FimU-PilVWXE, plus the putative adhesin PilY1,

prime pilus assembly and are proposed to form the pilus tip. The minor pilins and PilY1 are

encoded in an operon that is positively regulated by the FimS-AlgR two-component system.

Independent of pilus assembly, PilY1 is proposed to be a mechanosensory component that - in

conjunction with minor pilins - triggers up-regulation of acute virulence phenotypes upon surface

attachment. Here, we investigated the link between the minor pilins and virulence. *pilW, pilX,*

and *pilY1* mutants had reduced virulence towards *Caenorhabditis elegans* relative to wild type or

a major pilin mutant, implying a role in pathogenicity that is independent of pilus assembly. We

hypothesized that loss of specific minor pilins relieves feedback inhibition on FimS-AlgR,

increasing transcription of the minor pilin operon and other members of the AlgR regulon.

Reporter assays confirmed that FimS-AlgR were required for the increased expression from the

minor pilin operon promoter upon loss of select minor pilins. Overexpression of AlgR or its

hyperactivation via point mutation reduced virulence, and the virulence defects of *pilW*, *pilX*, and

pilY1 mutants were dependent on FimS-AlgR expression. We propose that PilY1 and the minor

pilins inhibit their own expression, and that loss of these proteins leads to FimS-mediated

activation of AlgR, and reduced expression of acute-phase virulence factors. This mechanism

could contribute to adaptation of *P. aeruginosa* in chronic lung infections, as mutations in the

minor pilin operon result in the loss of piliation and increased expression of AlgR-dependent

virulence factors – such as alginate – that are characteristic of such infections.

Author summary

Pseudomonas aeruginosa causes dangerous infections, including chronic lung infections

in cystic fibrosis patients. It uses many strategies to infect its hosts, including the use of

grappling hook-like fibres called type IV pili. Among the components involved in assembly and

function of the pilus are five proteins called minor pilins that - along with a larger protein called

PilY1 - may help the pilus attach to surfaces. In a roundworm infection model, loss of PilY1 and

specific minor pilins delayed killing, while loss of other pilus proteins did not. We traced this

effect to increased activation of the FimS-AlgR regulatory system that inhibits expression of

virulence factors used to initiate infections, while positively regulating chronic infection traits

such as alginate production, a phenotype called mucoidy. A disruption in the appropriate timing

of FimS-AlgR-dependent virulence factor expression when select minor pilins or PilY1 are

missing may explain why those pilus-deficient mutants have reduced virulence compared with

others whose products are not under FimS-AlgR control. Increased FimS-AlgR activity upon

loss of PilY1 and specific minor pilins could help explain the frequent co-occurrence of the non-

piliated and mucoid phenotypes that are hallmarks of chronic *P. aeruginosa* lung infections.

Introduction

Pseudomonas aeruginosa is a Gram-negative opportunistic pathogen, recently listed as

one of the highest priority antimicrobial-resistant threats by the World Health Organization, due

to its high intrinsic antibiotic resistance and recalcitrance to therapy [1]. Among its virulence

factors are filamentous surface appendages called the type IV pili (T4P), sophisticated biological

nanomachines that are broadly distributed among bacteria and archaea [2, 3]. In *P. aeruginosa*,

T4P facilitate surface and host cell adhesion, colonization, biofilm maturation, virulence, and

twitching, a form of surface-associated motility facilitated by cycles of extension, adhesion, and

retraction of T4P fibres [3-11]. T4P are composed of hundreds to thousands of copies of small

proteins called major pilins (PilA in *P. aeruginosa*) along with the low abundance minor pilins

(MPs), FimU-PilVWXE [12-16]. The MPs are encoded in a polycistronic operon with the *pilY1*

gene that codes for a large ~125 kDa non-pilin protein. This operon is positively regulated by the

two-component system (TCS) FimS (AlgZ)-AlgR, where FimS is a predicted histidine sensor

kinase and AlgR is a response regulator that promotes expression of genes important for biofilms

and chronic cystic fibrosis (CF) lung infections [17-19]. The N-termini of immature pilins are

cleaved and methylated at the cytoplasmic face of the inner membrane (IM) by the prepilin

peptidase, PilD, while PilY1 may be processed by signal peptidase 1 [20-23]. Mature pilins are

polymerized into a T4P fibre via an envelope-spanning assembly machinery, where individual

PilA subunits are added or removed at the platform protein, PilC, via action of the ATPases PilB

and PilT, respectively [2, 24].

The MPs and PilY1 are required for T4P function in several bacterial species, including

P. aeruginosa, *Escherichia coli*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, and *Myxococcus*

xanthus [12-15, 25-28]. PilY1 and the MPs were originally proposed to oppose pilus retraction,

as some surface pili remain in *pilY1* and MP mutants when retraction is blocked via deletion of

pilT [21, 26, 27, 29, 30]. We recently showed that deletion of the minor pseudopilins of the Xcp

type II secretion system in a *pilT* background lacking the T4P MPs abolished pilus assembly,

suggesting that when MPs are missing, the minor pseudopilins can prime extension, but cannot

counteract retraction [22]. We also demonstrated that PilY1 and the MPs are present in sheared

pili, and that the loss of PilV, PilW, PilX, or PilY1 excludes the other three components from the

pilus [22]. Thus, PilVWXY1 form a core assembly-initiation subcomplex, while FimU and PilE

connect this complex to PilA. Initiation of assembly with subsequent addition of PilA subunits

beneath would place the MPs at the pilus tip, with PilY1 − the largest component − at the distal

position, supporting the hypothesis that PilY1 is a T4P-associated adhesin [29].

PilY1 and the MPs (and their regulators FimS-AlgR) are required for T4P biogenesis, and

therefore T4P-mediated function [12-15, 17, 19]. However, recent studies hinted at more

enigmatic roles of PilWXY1 in virulence. Bohn et al. [31] showed that in a non-piliated *P.*

aeruginosa background, subsequent loss of *pilY1* reduced virulence in a *Caenorhabditis elegans*

fast killing assay and in a murine airway infection model, and increased resistance to killing by

neutrophils. Thus, PilY1 has a role in virulence that does not require functional pili. Other

studies using *C. elegans* infection models suggested that MP and *pilY1* mutants had attenuated

virulence relative to WT, and in one case, to a non-piliated mutant [32-35]. Recently, Siryaporn

et al. [36] showed that PilWXY1 were required for surface-activated virulence towards amoebae,

while other non-piliated mutants had WT virulence. The N-terminal region of PilY1 has limited

sequence similarity to the eukaryotic von Willebrand factor A (VWFa) domain, which can be

deformed by shear forces [37]. In-frame deletion of this domain from PilY1 allowed normally

avirulent planktonic cells to kill amoebae [36]. PilY1 was therefore proposed to be a

mechanosensor, where deformation of its VWFa domain upon surface interaction induced

expression of virulence factors. One important caveat of that study was that an *algR* mutant

(which also lacks PilY1 and the MPs) had WT virulence towards amoebae [36].

Deformation of PilA subunits by tensile forces acting upon surface attached pili was also

proposed as a possible way to signal surface attachment. Detection of partly unfolded pilins by

the Pil-Chp chemotactic system could lead to increased cyclic adenosine monophosphate

(cAMP) synthesis via the CyaB adenylate cyclase [38, 39]. cAMP is bound by Vfr (virulence

factor regulator), a key transcription factor that promotes expression of virulence factors

involved in motility, attachment, and secretion [38-40]. *fimS-algR* transcription is activated by

Vfr, leading to increased transcription of *fimU-pilVWXY1E* [38]. PilVWXY1 were proposed to

repress their own expression in an AlgR-dependent manner, as the loss of *pilV*, *pilW*, *pilX*, or

pilY1 led to elevated expression of the MP operon and *fimS-algR* [21, 31, 36, 38]. The

mechanism of this putative feedback inhibition is largely uncharacterized, but has been

speculated to involve FimS [38].

Once expression of the MP operon is activated, extracellular PilY1 may sense surface

association via its VWFa domain and transduce this information through the T4P assembly

machinery [36, 38]. This signal is thought to activate an IM-localized diguanylate cyclase, SadC,

and increase levels of c-di-GMP, a second messenger that promotes expression of genes

associated with a biofilm lifestyle while repressing early-phase virulence traits such as swarming

motility [38, 41]. This model was supported by studies demonstrating that loss of *pilW*, *pilX*, or

pilY1 in a high-c-di-GMP background resulted in hyper-swarming, indicative of reduced c-di-

GMP [37, 42]. Rodesney et al. [43] showed that c-di-GMP levels increased in response to shear

forces, and that functional T4P were required for this phenomenon, further supporting this

hypothesis. However, unlike *pilW*, *pilX*, and *pilY1* mutants, a *sadC* mutant had WT virulence

towards amoebae, suggesting the PilWXY1-SadC pathway may be important for surface-

sensing, but not necessarily for surface-activated virulence [36].

Although PilY1 and the MPs influence virulence, the underlying mechanism remains

unclear [31-34, 36]. We hypothesized that a subset of these components represses FimS activity,

such that loss of *pilW*, *pilX*, or *pilY1* activates FimS-AlgR, shifting the bacteria to a less

pathogenic phenotype typically associated with chronic infection. We found that *pilW*, *pilX*, and

pilY1 mutants had attenuated virulence in *C. elegans* slow killing (SK) assays compared to WT

or a *pilA* mutant, and this was was dependent on FimS-AlgR, as double mutants had wild type

(WT) virulence. Hyperactivation (via phospho-mimetic point mutation) or overexpression of

AlgR alone was sufficient to attenuate virulence. Together, these data are consistent with a

model where loss of PilWXY1 relieves feedback inhibition on expression of the AlgR regulon,

resulting in dysregulation of virulence factors that are important for *C. elegans* pathogenesis.

Results

PilWXY1 are important for T4P-independent virulence in PA14 and PAO1

Specific genes in the MP operon were reported to be important for virulence in amoebae,

nematodes, and mouse models, but those studies were done using various strain backgrounds

[31-34, 36]. We first sought to confirm these results in the *C. elegans* SK model, using two well-

studied strains. SK assays were performed using PA14 with deletions of *pilA*, *fimU*, *pilV*, *pilW*,

pilX, *pilY1*, or *pilE* (Fig 1A). *fimU* and *pilE* mutants had WT virulence, while *pilW*, *pilX*, and

pilY1 mutants had reduced virulence compared to WT. *pilW*, *pilX*, and *pilY1* mutants were also

less virulent than a *pilA* mutant, suggesting their reduced virulence was not due to loss of

functional T4P. *pilA* and *pilV* mutants had intermediate virulence, where they were significantly

less pathogenic than WT only in some trials. The twitching and virulence defects of *pilW*, *pilX*,

and *pilY1* mutants could be partially complemented by expression of the relevant gene *in trans*

(Supplementary Fig S1 and S2). The stoichiometry of PilY1 and the MPs is important for

optimal T4P function, which may explain the lack of full complementation [21]. To verify that

these phenotypes were not strain-specific, we tested PAO1 transposon-insertion mutants of *pilA*,

fimU, *pilV*, *pilW*, *pilX*, *pilY1*, and *pilE* in the SK assay (Fig 1B). Similar to the results in PA14,

PilWXY1 were important for T4P-independent virulence. However, the *fimU* and *pilV* mutants

were also less pathogenic than *pilA*; the PA14 and PAO1 MPs are divergent (61-75% amino acid

similarity), so it is possible that FimU and PilV function slightly differently in PAO1 versus

PA14 [44]. To focus on genes that were generally important for virulence of *P. aeruginosa*, we

undertook studies of the mechanism responsible for loss of virulence in the *pilW*, *pilX*, and *pilY1*

mutants.

PilWXY1 promote virulence in a SadC-independent manner

PilWXY1 were previously implicated in promoting c-di-GMP production via SadC, such

that loss of *pilW*, *pilX*, or *pilY1* resulted in a hyper-swarming, biofilm-deficient phenotype,

indicative of low intracellular c-di-GMP [37, 38, 42]. Therefore, we hypothesized that biofilm

defects of *pilW*, *pilX*, and *pilY1* might impede their ability to colonize the *C. elegans* gut, leading

to reduced virulence, and thus tested their ability to form biofilms in liquid SK medium. The

PA14 and PAO1 parent strains and their cognate *pilA*, *fimU*, *pilV*, *pilW*, *pilX*, *pilY1*, and *pilE*

mutants formed negligible levels of biofilm in that medium (Supplementary Fig S3). Next, we

performed swarming motility assays using 0.5% agar SK plates, as swarming is inversely

correlated with biofilm formation [41, 45]. PA14 *pilW*, *pilX*, and *pilY1* mutants had swarming

defects (Fig 2), consistent with their reduced virulence in *C. elegans*, while all PAO1 strains had

low swarming on this media (S4 Fig). These data are inconsistent with reports that PilWXY1

promote c-di-GMP production, and instead suggest that the absence of *pilW*, *pilX*, or *pilY1*

promotes c-di-GMP production, resulting in repression of swarming. We next investigated

whether SadC-mediated biofilm formation affected virulence towards *C. elegans*, as would be

expected if the loss of virulence in *pilW*, *pilX*, and *pilY1* mutants was due to dysregulation of

SadC activity. However, there was no difference in virulence between PA14 and a *sadC* mutant

(Supplementary Fig S5). Further, overexpression of SadC led to a hyper-biofilm phenotype *in*

vitro but a slight reduction in virulence, demonstrating that the amount of biofilm formed does

not correlate with virulence in *C. elegans* (Supplementary Fig S5). Although the exact

mechanisms of *P. aeruginosa* pathogenesis in *C. elegans* are not fully understood, biofilms were

suggested to be important for establishment of infection [46-48]. However, in these assay

conditions, biofilms do not appear to be a major contributor to *P. aeruginosa* pathogenesis.

PilVWXY1 repress expression of the minor pilin operon

After ruling out involvement of the SadC pathway, we next explored the potential role of

FimS-AlgR in PilWXY1-mediated virulence. Informed by previous work in our laboratory,

showing that the sensor kinase PilS of the PilSR TCS interacts directly with PilA in the inner

membrane to regulate major pilin levels [49], we hypothesized that FimS interacts with one or

more MPs, and that loss of that interaction could lead to phosphorylation of AlgR and

subsequent upregulation of the MP operon. Bacterial two-hybrid (BACTH) assays were used to

identify potential interactions between FimS and PilA, FimU, PilV, PilW, PilX, or PilE (Fig 3A).

Positive interactions were identified between FimS and each pilin; however, based on our

experience with PilS, binding of pilins is necessary but not sufficient for regulation.

To decipher which MPs might modulate expression of the operon, we monitored

expression from the *fimU* promoter using a *luxCDABE* reporter. Compared to WT PA14, there

was a ~25-fold increase in luminescence in *pilV*, *pilW*, *pilX*, and *pilY1* mutants (Fig 3B). *fimU*

and *pilA* mutants had ~5-fold increased promoter activity, while a *pilE* mutant was comparable

to WT. *fimS* and *algR* mutants had low baseline luminescence, ~10-fold lower than WT. To

determine whether the increased promoter activity in *pilV*, *pilW*, *pilX*, and *pilY1* mutants was

dependent on FimS-AlgR, either *fimS* or *algR* was deleted in the *pilY1* mutant background. As

previously reported, the *pilY1 algR* double mutant had low luminescence (~10-fold lower than

WT), consistent with AlgR acting as a positive regulator of the MP operon [38]. Loss of *fimS* in

the *pilY1* mutant background also abolished *fimU* promoter activity (~10-fold lower than WT),

supporting the idea that FimS may monitor PilVWXY1 levels and activate AlgR when levels are

low. Based on these data, PilA, FimU, and PilE are unlikely to modulate FimS-AlgR activity

even though they can interact with FimS.

Hyperactivation of AlgR results in attenuated virulence

Thus far, our results supported the hypothesis that loss of PilWXY1 relieves feedback

inhibition on FimS-AlgR, resulting in AlgR activation. Next, we aimed to test whether

hyperactivation of AlgR alone could decrease virulence. We constructed chromosomal $algR_{D54E}$

phospho-mimetic point mutants [50] in both the PA14 and PAO1 backgrounds. We also

constructed $algR_{D54A}$ point mutants, as AlgR phosphorylation is required for transcription of a

subset of genes in its regulon, including the MP operon [17, 50, 51]. SK assays were performed

for PA14 and PAO1 $algR_{D54A}$ and $algR_{D54E}$ mutants, along with PA14 and PAO1 *fimS* and *algR*

deletion mutants. $algR_{D54E}$ mutants of both PA14 and PAO1 were less pathogenic than the

corresponding WT strains, while *algR* deletion and $algR_{D54A}$ mutants had WT virulence (Fig 4).

fimS mutants also had WT virulence. Loss of FimS-AlgR was shown previously to decrease

expression of the minor pilins and PilY1, and prevent pilus assembly [17, 38]. Because our data

show that loss of FimS-AlgR (and thus loss of minor pilin expression) had no impact on

virulence, we conclude that reduced virulence of *pilW*, *pilX*, and *pilY1* mutants is not due to loss

of those gene products directly, but rather to the resulting activation of FimS-AlgR.

Overexpression of AlgR results in attenuated virulence

Increased transcription of *fimS-algR* in a *pilY1* mutant relative to WT has been reported

[36], suggesting that reduced virulence could arise through increased levels of FimS-AlgR

expression as well as activation of the TCS. Therefore, we asked whether increased AlgR levels

would attenuate virulence. When *algR* was expressed *in trans* from a multicopy plasmid in PA14

algR, virulence was reduced compared to the vector-only control (Fig 5A). Because un-

phosphorylated AlgR can also affect transcription of a subset of genes [52, 53], we also tested

the same mutant complemented with $AlgR_{D54A}$. Complementation of the *algR* mutant with

$AlgR_{D54A}$ resulted in a severe virulence defect relative to the vector-only control. Thus, AlgR

hyperactivation and overexpression independently diminish *P. aeruginosa* virulence towards *C.*

elegans. Lastly, as AlgR was reported to be a positive regulator of biofilm formation [54], we

performed biofilm assays for PA14 *algR* complemented with AlgR or AlgR$_{D54A}$. Expression of

both AlgR and AlgR$_{D54A}$ led to hyper-biofilm formation (Fig 5B), further emphasizing that the

ability of a strain to form biofilms in SK medium does not correlate with virulence. Instead, we

suggest that virulence factors that are repressed by FimS-AlgR are important for *C. elegans* slow

killing, and thus an increase in AlgR levels and/or activity attenuates virulence.

The virulence defects of *pilW*, *pilX* and *pilY1* mutants are dependent on FimS-AlgR

To provide further support for this model, we asked whether the virulence defects of

PA14 *pilW*, *pilX*, and *pilY1* mutants were dependent on FimS-AlgR. We deleted *fimS* or *algR* in

the *pilW*, *pilX*, and *pilY1* backgrounds, and tested virulence of the double mutants (Fig 6). We

also deleted *pilW*, *pilX*, and *pilY1* in the *algR*$_{D54A}$ background, to test if AlgR activation was

required for the loss of virulence in *pilW*, *pilX*, and *pilY1* mutants. In all cases, the double

mutants had WT virulence, equivalent to that of the *fimS*, *algR*, or *algR*$_{D54A}$ single mutants.

These results demonstrate that both FimS and AlgR are required for the reduction in virulence

resulting from loss of PilWXY1. Although overexpression of AlgR$_{D54A}$ *in trans* repressed

virulence (Fig 5A), the chromosomal point mutation was sufficient to alleviate the virulence

defect of *pilW*, *pilX*, and *pilY1* mutants, suggesting that AlgR phosphorylation at residue D54 is

important for PilWXY1-mediated virulence.

Loss of FimS-AlgR rescues swarming motility defects of *pilW*, *pilX*, and *pilY1* mutants

After confirming that reduced virulence in PA14 *pilW*, *pilX*, and *pilY1* mutants was

dependent on FimS-AlgR, we compared swarming motility of PA14 *pilW*, *pilX*, *pilY1*, *fimS*,

algR, and *algR*$_{D54A}$ single and double mutants (Fig 7). *pilW*, *pilX*, and *pilY1* mutants had

swarming defects, while swarming of *fimS*, *algR*, and *algR*$_{D54A}$ mutants was similar to WT

PA14. Deletion of *fimS* or *algR* in the *pilW*, *pilX*, and *pilY1* backgrounds restored swarming to

the level of *fimS* and *algR* single mutants. Swarming of the *pilW algR*$_{D54A}$, *pilX algR*$_{D54A}$, and

pilY1 algR$_{D54A}$ mutants resembled that of the *algR*$_{D54A}$ single mutant. These data mirror the SK

assay results, in that the swarming defects of the *pilW*, *pilX*, and *pilY1* mutants are dependent on

FimS and AlgR. Taken together, our data support a model where PilWXY1 repress their own

expression via FimS-AlgR, and loss of PilWXY1 relieves this inhibition, activating AlgR in a

FimS-dependent manner. Activated AlgR then represses genes that are important for swarming

and virulence towards *C. elegans* under SK conditions.

Discussion

P. aeruginosa uses T4P to attach to surfaces and host cells, during biofilm maturation,

and to move across surfaces via twitching motility [2]. The MPs and PilY1 are important players

in T4P biogenesis and function, but also in regulation of swarming motility, surface attachment,

mechanosensation, and virulence [36-38, 42]. The MP operon is positively regulated by FimS-

AlgR, a TCS implicated in regulation of chronic *P. aeruginosa* lung infections [17-19]. Here, we

explored the potential connection between loss of PilWXY1 (and thus, loss of T4P) and AlgR

activation in virulence towards *C. elegans*. We showed that *pilW*, *pilX*, and *pilY1* mutants were

less virulent than WT or a *pilA* mutant – similar to the reduced virulence of these mutants

towards amoebae [36]. These data support the idea that PilWXY1 modulate virulence

independently of their role in T4P assembly. We confirmed previous reports [21, 31, 38] that in

the absence of *pilV*, *pilW*, *pilX*, or *pilY1*, expression of the MP operon is significantly increased,

and that this requires FimS-AlgR. Either hyperactivation or overexpression of AlgR reduced

virulence, while loss of *fimS* or *algR* in *pilW*, *pilX*, or *pilY1* backgrounds reverted it to WT levels.

These data suggest that FimS acts as a molecular thermostat to monitor MP levels, and in

their absence, activates AlgR to upregulate expression of the MP operon (Fig 8). A similar

regulatory mechanism was recently described for the PilSR TCS, where PilS phosphorylates

PilR when PilA levels are low, and dephosphorylates PilR when PilA levels are high [49]. It is

not yet clear whether FimS responds to PilVWXY1 individually or as a group, as these four

proteins are thought to form an inner membrane subcomplex that primes assembly of T4P [22,

55, 56]. When overexpressed individually *in trans*, each of the MPs inhibited twitching motility

in PAO1 [21], but since the others were still expressed from the chromosome, the exact nature of

the signal detected by FimS remains to be determined.

Kuchma et al. [37, 42] reported that loss of *pilW*, *pilX*, or *pilY1* increased swarming

motility and decreased biofilm formation, both indicative of low c-di-GMP levels. As biofilms

were proposed to contribute to *P. aeruginosa* pathogenesis in *C. elegans*, we investigated

whether the reduction in virulence in the absence of PilWXY1 was linked to decreased biofilm

via loss of SadC activation [46-48, 57]. In our hands, levels of *sadC* had no impact on virulence

even though they clearly modulated the amount of biofilm produced in SK media

(Supplementary Fig S5). Irazoqui et al. [48] examined the *C. elegans* gut during *P. aeruginosa*

infection, and described extracellular material that they suggested might indicate presence of a

biofilm. Anti-biofilm compounds reduced *P. aeruginosa* virulence towards *C. elegans*, but a

mechanism of action for those compounds has not been described [47]. Recently, the small RNA

SrbA was shown to modulate both biofilm and virulence towards *C. elegans*; however, deletion

of *srbA* led to altered transcription of at least 26 other genes that may also affect virulence [58].

Rather than using standard biofilm and swarming motility media, we performed these assays in

SK media to more closely mimic the conditions that the bacteria are exposed to in the SK assay.

To our knowledge, using SK media for swarming motility assays has only been attempted once

[33], and it has not been used for biofilm assays. As we found no correlation between biofilm

formation and virulence, we suggest that acute-phase virulence factors may be more important

for *C. elegans* pathogenesis in the SK model.

PilY1 and the MPs have been implicated in surface detection and activation of virulence

[36, 38]. Because loss of PilY1 or any of the MPs prevents T4P assembly and function, it is

important to distinguish phenotypes resulting from lack of those components specifically, rather

than loss of piliation [22]. Luo et al. [38] suggested that association of PilY1 with surfaces

transduces a signal through the T4P system to stimulate c-di-GMP production by SadC, while

Rodesney et al. [43] showed that loss of *pilA*, *pilY1*, or *pilT* prevents surface-activated c-di-GMP

production. Rodesney et al. [43] proposed that both PilY1 and functional T4P are required for

mechanosensation; however, it is not possible to delete *pilY1* without ablating T4P assembly.

Rather than PilY1 signalling through the T4P assembly machinery and SadC [38, 42], T4P-

mediated surface attachment may transiently deplete PilVWXY1 levels in the IM, resulting in

increased FimS-AlgR activity and transition towards a sessile, biofilm lifestyle. Surface-

activated increases in c-di-GMP levels are thought to occur within hours of attachment [38, 43];

this idea is consistent with our swarming data, where activation of FimS-AlgR in response to

PilVWXY1 depletion reduces swarming motility (Fig 2).

Whether the loss of *pilW*, *pilX*, or *pilY1* leads to increased amounts of AlgR, its increased

phosphorylation via FimS, or both remains to be clarified. Okkotsu et al. [50] showed that AlgR

and AlgR$_{D54E}$ levels are comparable, suggesting that the loss of virulence we observed for PA14

$algR_{D54E}$ is attributable to the D54E phospho-mimetic mutation alone. While overexpression of

$AlgR_{D54A}$ *in trans* reduced virulence (Fig 5A), the same mutation on the chromosome reverted

virulence of *pilW*, *pilX,* and *pilY1* mutants to WT levels (Fig 6). Therefore, we suspect that it is

primarily AlgR phosphorylation (or lack of AlgR dephosphorylation) that is responsible for loss

of virulence. However, it is possible that increased AlgR protein levels and phosphorylation both

contribute. Kong et al. [54] showed that AlgR binds *fimS-algR*, suggesting that the TCS could

positively regulate its own transcription in response to reduced PilWXY1 levels.

In addition to being essential for T4P function, FimS and AlgR control alginate

production in the context of chronic CF infections, where *algR* transcription is high [18, 59].

Phosphorylation of AlgR increases binding affinity at some – but not all – of its target sequences

[17, 50, 51, 53]. For example, $AlgR_{D54N}$ failed to support twitching motility, but did not affect

alginate production [17, 51]. FimS is an unorthodox histidine kinase, with 4 transmembrane

domains instead of the more typical 2, and lacks both a periplasmic sensing domain and the

canonical motif involved in ATP coordination, which mediates auto-phosphorylation [19, 60].

Direct interaction and/or phospho-transfer between FimS and AlgR have not been reported.

Rather, the idea that FimS acts as a kinase for AlgR comes from this and other studies

demonstrating similar phenotypes for *fimS, algR,* and $algR_{D54N}$ mutants [17, 18, 61]. FimS and

AlgR promote expression of genes important for production of alginate, biofilms, and c-di-GMP,

and inhibit expression of virulence factors such as the T3SS, pyocyanin, and quorum sensing

[54, 62-65]. Dynamic regulation of AlgR levels are important for pathogenesis, as either deletion

or overexpression of *algR* attenuated virulence in a murine model of acute septicemia [66].

However, an *algR* mutant was also reported to have WT virulence towards amoebae [36]. The

outcomes that result from interaction of *P. aeruginosa* with different hosts will depend on a

variety of factors including host defenses, site of infection, available nutrients, and virulence

repertoire of a particular strain. However, our results suggest that changes in the specific

repertoire of bacterial virulence factors, or the timing of their production, can tip the balance in

the host's favour.

The subset of AlgR-regulated virulence genes important for *C. elegans* pathogenesis is

not clearly defined. Screening of a PA14 transposon library for loss of virulence implicated

several genes encoding regulators rather than individual virulence factors, suggesting that *C.*

elegans pathogenesis is multifactorial [33]. In support of this hypothesis, a study of 18 WT *P.*

aeruginosa strains revealed no correlation between virulence and the presence of any specific

virulence gene [67]. We saw WT virulence for *algR* mutants, consistent with a role for AlgR in

negative regulation of acute phase virulence factors. Factors under positive control of AlgR may

be important during later stages of infection in more complex mammalian infection models [68,

69].

While important for the initial stages of infection, T4P are less critical in chronic CF lung

infections and are often lost when the bacteria grow for extended periods as biofilms [5, 70, 71].

P. aeruginosa CF isolates frequently become mucoid via activation of AlgR, leading to reduced

production of other virulence factors [59, 72, 73]. Although the two outcomes are not necessarily

temporally or mechanistically linked, mutations that achieve both may be advantageous during

chronic CF lung infections. Specifically, loss of PilWXY1 may be adaptive in the context of CF,

leading to AlgR activation and loss of T4P function. It will be interesting to examine the

genotypes of mucoid CF isolates for these types of mutations.

Materials and methods

Bacterial strains and plasmids

Strains and plasmids used in this work are listed in Supplementary Table S1. Bacteria were grown at 37°C for 16 h in 5 ml Luria Bertani Lennox (LB) broth, or on 1.5% agar LB plates, unless otherwise specified. Plasmids were transformed into chemically-competent *E. coli* by heat-shock, and into *P. aeruginosa* by electroporation [74]. Where appropriate, gentamicin (Gm) was added at 15 μg/ml for *E. coli*, and 30 μg/ml for *P. aeruginosa*. Kanamycin (Kan) was added at 50 μg/ml for *E. coli*, and 150 μg/ml for *P. aeruginosa*. Ampicillin (Amp) was added at 100 μg/ml for *E. coli*. L-Arabinose was added at 0.05% where indicated to induce expression from the pBADGr promoter [75].

Cloning procedures

Vectors were constructed using standard cloning procedures, using the primers listed in Supplementary Table S2. Deletion constructs were designed to contain 500-1000 bp homology upstream and downstream the gene to be deleted. Deletion constructs for PA14 *fimU, pilV, pilW, pilX, pilY1,* and *pilE* were synthesized by Genscript in the pUC57Kan vector. pEX18Gm-*sadC* was created by amplifying the *sadC* deletion region from PA14 *sadC roeA* [41], followed by digestion and ligation into pEX18Gm. pEX18Gm-*algR*$_{D54A}$, pEX18Gm-*algR*$_{D54E}$, and pEX18Gm-*fimS* were made by overlap extension PCR [76]. pMS402-P*fimU* was created by amplifying and digesting the promoter region of the PA14 MP operon. Digested pBADGr was treated with alkaline phosphatase prior to ligation to avoid re-circularization of the vector. Constructs were verified by Sanger sequencing (MOBIX lab, McMaster, Hamilton, ON).

Mutant generation by allelic exchange

Allelic exchange was used to remove or alter specific genes [77]. pEX18Gm suicide plasmid derivatives (see Cloning procedures and Table 1) were used to create all mutants in this work. After heat-shock transformation into *E. coli* SM10 cells, pEX18Gm constructs were conjugated into corresponding PA14 or PAO1 parent strains. Cells were then transferred to *Pseudomonas* isolation agar (PIA) Gm100 plates and incubated for 18 h at 37°C, to select for integration of pEX18Gm derivatives into the chromosome. Colonies were streaked onto LB/sucrose and incubated at 30°C for 18 h to select against merodiploids. Resultant colonies were patched onto LB and LB Gm30 to identify gentamicin-sensitive colonies. Regions flanking the desired mutations were amplified and sequenced to confirm success.

Twitching motility assays

Twitching motility assays were performed as previously described [78], with the following modifications. Individual colonies were stab-inoculated in triplicate into 1% agar LB solidified in plasma-treated tissue culture-grade plates (Thermo Fisher) and incubated at 30°C for 48 h. Agar was carefully removed and plates were stained with 1% crystal violet for 5 min. Unbound dye was removed by rinsing with water, then stained twitching areas were measured using ImageJ. Twitching zones were normalized to WT (100%).

Swarming motility assays

SM assays were performed as previously described [79], with the following modifications. The concentration of agar in SK plates (50 mM NaCl, 0.5% agar, 0.35% peptone, 1 mM $CaCl_2$, 1 mM $MgSO_4$, 5 µg/ml cholesterol in 100% ethanol, 20 mM KH_2PO_4, and 5 mM

K_2HPO_4) was reduced to 0.5% (w/v) to promote swarming. Plates were dried individually for 1.5

h at room temperature then 3.5 μl of an overnight culture was spotted onto the center of the

duplicate plates. Plates were individually incubated upright at 37°C for 24 h, followed by

incubation at 22°C for 24 h. Empty swarming motility plates were placed around the perimeter

of arrayed experimental plates to reduce edge effects and promote formation of reproducible

swarming zones. Experiments were repeated three times, 2 technical replicates per trial.

Biofilm assays

Biofilm assays were performed as previously described, with modifications [80]. *P.*

aeruginosa cultures were grown for 16 h at 37°C, diluted 1:200 in fresh LB, and grown to OD_{600}

~ 0.1. Cultures were then diluted 1:500 in liquid SK media (50 mM NaCl, 0.35% peptone, 1 mM

$CaCl_2$, 1 mM $MgSO_4$, 5 μg/ml cholesterol in EtOH, 20 mM KH_2PO_4, and 5 mM K_2HPO_4), then

96-well plates were inoculated with 150 μl each strain, in triplicate. Sterility controls (liquid SK

media) were included throughout the plate to check for contamination. Plates were covered with

peg lids (Nunc) then wrapped in parafilm and incubated at 37°C for 24 h, shaken at 200 rpm.

After incubation, the OD_{600} of the plate was measured to check for uniform growth and lack of

contamination. Peg lids were washed for 10 min in 200 μl/well 1X phosphate-buffered saline

(PBS), then stained with 200 μl/well 0.1% (w/v) crystal violet for 15 min. Unbound crystal violet

was removed by washing lids in 70 ml distilled water 5 times at 10 min intervals. Crystal violet

was solubilized from lids in 200 μl/well 33.3% acetic acid, then the absorbance at 600 nm was

measured. Optical density and absorbance at 600 nm were plotted for growth and biofilm

formation, respectively, then analyzed by one-way ANOVA followed by Dunnett post-test to

compare each mutant to the WT control, p = 0.05. Error bars indicate standard error of the mean.

Representative wells of acetic acid-solubilized crystal violet were imaged.

***Caenorhabditis elegans* slow killing assay**

SK assays were performed as described previously [81]. SK plates (0.35% peptone, 50

mM NaCl, 2% agar, 1 mM CaCl$_2$, 5 μg/ml cholesterol, 1 mM MgSO$_4$, 20 mM KH$_2$PO$_4$, 5 mM

K$_2$HPO$_4$, 100 μM FUDR) were seeded with 100 μl of an overnight culture and incubated

overnight at 37°C. The following day, plates were enriched with 1 ml of an overnight culture

concentrated to 100 μl. Synchronized L4 worms were collected from *E. coli* OP50 plates, washed

twice in M9 buffer, and then >50 worms were seeded onto each bacterial lawn on individual SK

plates. SK plates were incubated at 25°C and scored for dead worms every 24 h. Worms were

considered dead when they did not respond to touch, and were removed from the plate. OP50

was included as a negative control for virulence. Percent survival was plotted as a function of

time. Survival curves were plotted on GraphPad Prism 5.00 for Windows, then compared using

the Gehan-Breslow-Wilcoxon test, p = 0.05. Each assay was performed at least 3 times, and

differences were only considered significant if they were reproducible in all trials.

Representative trials are shown.

Luminescent reporter assay

Luminescent reporter assays were performed as previously described, with minor

modifications [49]. Various strains harbouring the pMS402-P*fimU* plasmid, encoding the

luciferase genes under control of the *fimU* promoter, were grown for 16 h at 37°C in LB Kan150,

then diluted 1:50 in fresh LB Kan150. Subsequently, 100 μl of each culture was added to white-

walled, clear-bottom 96-well plates (Corning) in triplicate, and incubated with shaking at 37°C in

a Synergy 4 microtiter plate reader (BioTek). Luminescence readings were taken every 15 min

for 5 h, and normalized to growth (OD_{600}) at each time point. Readings that exceeded the limit of

detection (>4 000 000 luminescence units) were discarded. At least 3 individual trials were

performed. Error bars indicate standard error of the mean.

Bacterial two-hybrid β-galactosidase activity assay

To test for interactions between FimS and individual pilins, bacterial two-hybrid

(BACTH) assays were performed as previously described [82]. pUT18C and pKT25 derivatives,

encoding the T18 and T25 domains of the *Bordetella pertussis* CyaA adenylate cyclase fused to

the N-terminus of FimS, PilA, FimU, PilV, PilW, PilX, or PilE [22, 49, 83], were co-transformed

into *E. coli* BTH 101 to screen for pairwise interactions. Each interaction was tested in both

orientations, with FimS in pUT18C and each pilin in pKT25, or FimS in pKT25 and each pilin in

pUT18C. Single colonies were inoculated in 5 ml LB Amp100 Kan50 and grown overnight. The

following day, 100 μl was inoculated into 5 ml fresh media and grown toOD_{600} = 0.6, then 5 μl

was spotted onto MacConkey plates (1.5% agar, 100μg/ml ampicillin, 50μg/ml kanamycin, 1%

(w/v) maltose, 0.5mM isopropyl b-D-thiogalactopyranoside) (Difco). Plates were incubated at

30°C for 24 h. An interaction was considered positive when colonies appeared pink, as a result

of fermentation of maltose. BTH 101 expressing pUT18C and pKT25 empty vectors was used as

a negative control, and BTH 101 expressing pUT18C-*fimS* and pKT25-*fimS* was used as a

positive control [49].

Acknowledgements

This work was funded by Canadian Institutes of Health Research (CIHR) grant MOP-86639 to

LLB. VAM held a CIHR Canada Graduate Scholarship. SLNK held an Ontario Graduate

Scholarship.

References

1.   Global priority list of antibiotic-resistant bacteria to guide research, discovery, and

development of new antibiotics [press release]. World Health Organization 2017.

2.   Burrows LL. *Pseudomonas aeruginosa* twitching motility: type IV pili in action. Annu

Rev Microbiol. 2012;66:493-520.

3.   Hospenthal MK, Costa TRD, Waksman G. A comprehensive guide to pilus biogenesis in

Gram-negative bacteria. Nat Rev Micro. 2017;15(6):365-79.

4.   Berry J-L, Pelicic V. Exceptionally widespread nanomachines composed of type IV

pilins: the prokaryotic Swiss Army knives. FEMS Microbiol Rev. 2015;39(1):1-21.

5.   Mahenthiralingam E, Campbell ME, Speert DP. Nonmotility and phagocytic resistance of

Pseudomonas aeruginosa isolates from chronically colonized patients with cystic fibrosis. Infect

Immun. 1994;62(2):596-605.

6.   Chi E, Mehl T, Nunn D, Lory S. Interaction of *Pseudomonas aeruginosa* with A549

pneumocyte cells. Infect Immun. 1991;59(3):822-8.

7.   Doig P, Todd T, Sastry PA, Lee KK, Hodges RS, Paranchych W, et al. Role of pili in

adhesion of *Pseudomonas aeruginosa* to human respiratory epithelial cells. Infect Immun.

1988;56(6):1641-6.

8.   Pier GB, Meluleni G, Neuger E. A murine model of chronic mucosal colonization by

Pseudomonas aeruginosa. Infect Immun. 1992;60(11):4768-76.

9.   Saiman L, Ishimoto K, Lory S, Prince A. The effect of piliation and exoproduct

expression on the adherence of *Pseudomonas aeruginosa* to respiratory epithelial monolayers. J

Infect Dis. 1990;161(3):541-8.

10.   Zoutman DE, Hulbert WC, Pasloske BL, Joffe AM, Volpel K, Trebilcock MK, et al. The

role of polar pili in the adherence of *Pseudomonas aeruginosa* to injured canine tracheal cells: a

semiquantitative morphologic study. Scanning Microsc. 1991;5(1):109-26.

11.   Comolli JC, Hauser AR, Waite L, Whitchurch CB, Mattick JS, Engel JN. *Pseudomonas*

aeruginosa gene products PilT and PilU are required for cytotoxicity in vitro and virulence in a

mouse model of acute pneumonia. Infect Immun. 1999;67(7):3625-30.

12.   Alm RA, Hallinan JP, Watson AA, Mattick JS. Fimbrial biogenesis genes of

Pseudomonas aeruginosa: *pilW* and *pilX* increase the similarity of type 4 fimbriae to the GSP

protein-secretion systems and *pilY1* encodes a gonococcal PilC homologue. Mol Microbiol.

1996;22(1):161-73.

13.   Alm RA, Mattick JS. Identification of a gene, *pilV*, required for type 4 fimbrial

biogenesis in *Pseudomonas aeruginosa*, whose product possesses a pre-pilin-like leader

sequence. Mol Microbiol. 1995;16(3):485-96.

14.   Alm RA, Mattick JS. Identification of two genes with prepilin-like leader sequences

involved in type 4 fimbrial biogenesis in *Pseudomonas aeruginosa*. J Bacteriol.

1996;178(13):3809-17.

15.   Russell MA, Darzins A. The *pilE* gene product of *Pseudomonas aeruginosa*, required for

pilus biogenesis, shares amino acid sequence identity with the N-termini of type 4 prepilin

proteins. Mol Microbiol. 1994;13(6):973-85.

16.   Hobbs M, Collie ES, Free PD, Livingston SP, Mattick JS. PilS and PilR, a two-

component transcriptional regulatory system controlling expression of type 4 fimbriae in

Pseudomonas aeruginosa. Mol Microbiol. 1993;7(5):669-82.

17.   Belete B, Lu H, Wozniak DJ. *Pseudomonas aeruginosa* AlgR regulates type IV pilus

biosynthesis by activating transcription of the *fimU-pilVWXY1Y2E* operon. J Bacteriol.

2008;190(6):2023-30.

18.   Whitchurch CB, Alm RA, Mattick JS. The alginate regulator AlgR and an associated

sensor FimS are required for twitching motility in *Pseudomonas aeruginosa*. Proc Natl Acad Sci

U S A. 1996;93(18):9839-43.

19.   Okkotsu Y, Little AS, Schurr MJ. The *Pseudomonas aeruginosa* AlgZR two-component

system coordinates multiple phenotypes. Front Cell Infect Microbiol. 2014;4:82.

20.   Strom MS, Bergman P, Lory S. Identification of active-site cysteines in the conserved

domain of PilD, the bifunctional type IV pilin leader peptidase/N-methyltransferase of

Pseudomonas aeruginosa. J Biol Chem. 1993;268(21):15788-94.

21.   Giltner CL, Habash M, Burrows LL. *Pseudomonas aeruginosa* minor pilins are

incorporated into type IV pili. J Mol Biol. 2010;398(3):444-61.

22.   Nguyen Y, Sugiman-Marangos S, Harvey H, Bell SD, Charlton CL, Junop MS, et al.

Pseudomonas aeruginosa minor pilins prime type IVa pilus assembly and promote surface

display of the PilY1 adhesin. J Biol Chem. 2015;290(1):601-11.

23.   Sauvonnet N, Vignon G, Pugsley AP, Gounon P. Pilus formation and protein secretion by

the same machinery in *Escherichia coli*. EMBO J. 2000;19(10):2221-8.

24.   McCallum M, Tammam S, Khan A, Burrows LL, Howell PL. The molecular mechanism

of the type IVa pilus motors. Nat Commun. 2017;8:15091.

25.   Ramer SW, Schoolnik GK, Wu C-Y, Hwang J, Schmidt SA, Bieber D. The type IV pilus

assembly complex: biogenic interactions among the bundle-forming pilus proteins of

enteropathogenic *Escherichia coli*. J Bacteriol. 2002;184(13):3457-65.

26.   Winther-Larsen HC, Wolfgang M, Dunham S, Van Putten JPM, Dorward D, Løvold C, et

al. A conserved set of pilin-like molecules controls type IV pilus dynamics and organelle-

associated functions in *Neisseria gonorrhoeae*. Mol Microbiol. 2005;56(4):903-17.

27.   Carbonnelle E, Helaine S, Nassif X, Pelicic V. A systematic genetic analysis in *Neisseria

meningitidis* defines the Pil proteins required for assembly, functionality, stabilization and export

of type IV pili. Mol Microbiol. 2006;61(6):1510-22.

28.   Chang Y-W, Rettberg LA, Treuner-Lange A, Iwasa J, Søgaard-Andersen L, Jensen GJ.

Architecture of the type IVa pilus machine. Science. 2016;351(6278).

29.   Heiniger RW, Winther-Larsen HC, Pickles RJ, Koomey M, Wolfgang MC. Infection of

human mucosal tissue by *Pseudomonas aeruginosa* requires sequential and mutually dependent

virulence factors and a novel pilus-associated adhesin. Cell Microbiol. 2010;12(8):1158-73.

30.   Wolfgang M, Park H-S, Hayes SF, van Putten JPM, Koomey M. Suppression of an

absolute defect in type IV pilus biogenesis by loss-of-function mutations in *pilT*, a twitching

motility gene in *Neisseria gonorrhoeae*. Proc Natl Acad Sci U S A. 1998;95(25):14973-8.

31.   Bohn Y-ST, Brandes G, Rakhimova E, Horatzek S, Salunkhe P, Munder A, et al.

Multiple roles of *Pseudomonas aeruginosa* TBCF10839 PilY1 in motility, transport and

infection. Mol Microbiol. 2009;71(3):730-47.

32.   Garvis S, Munder A, Ball G, de Bentzmann S, Wiehlmann L, Ewbank JJ, et al.

Caenorhabditis elegans semi-automated liquid screen reveals a specialized role for the

chemotaxis gene *cheB2* in *Pseudomonas aeruginosa* virulence. PLoS Pathog.

2009;5(8):e1000540.

33.   Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis A-R, et al.

Genome-wide identification of *Pseudomonas aeruginosa* virulence-related genes using a

Caenorhabditis elegans infection model. PLoS Pathog. 2012;8(7):e1002813.

34.   Lewenza S, Charron-Mazenod L, Giroux L, Zamponi AD. Feeding behaviour of

Caenorhabditis elegans is an indicator of *Pseudomonas aeruginosa* PAO1 virulence. PeerJ.

2014;2:e521.

35.   Jansen G, Crummenerl LL, Gilbert F, Mohr T, Pfefferkorn R, Thänert R, et al.

Evolutionary Transition from Pathogenicity to Commensalism: Global Regulator Mutations

Mediate Fitness Gains through Virulence Attenuation. Molecular Biology and Evolution.

2015;32(11):2883-96.

36.   Siryaporn A, Kuchma SL, O'Toole GA, Gitai Z. Surface attachment induces

Pseudomonas aeruginosa virulence. Proc Natl Acad Sci U S A. 2014;111(47):16860-5.

37.   Kuchma SL, Ballok AE, Merritt JH, Hammond JH, Lu W, Rabinowitz JD, et al. Cyclic-

di-GMP-mediated repression of swarming motility by *Pseudomonas aeruginosa*: the *pilY1* gene

and its impact on surface-associated behaviors. J Bacteriol. 2010;192(12):2950-64.

38.   Luo Y, Zhao K, Baker AE, Kuchma SL, Coggan KA, Wolfgang MC, et al. A hierarchical

cascade of second messengers regulates *Pseudomonas aeruginosa* surface behaviors. mBio.

2015;6(1).

39.   Persat A, Inclan YF, Engel JN, Stone HA, Gitai Z. Type IV pili mechanochemically

regulate virulence factors in *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A.

2015;112(24):7563-8.

40.   Wolfgang MC, Lee VT, Gilmore ME, Lory S. Coordinate regulation of bacterial

virulence genes by a novel adenylate cyclase-dependent signaling pathway. Dev Cell.

2003;4(2):253-63.

41.   Merritt JH, Brothers KM, Kuchma SL, O'Toole GA. SadC reciprocally influences biofilm

formation and swarming motility via modulation of exopolysaccharide production and flagellar

function. J Bacteriol. 2007;189(22):8154-64.

42.   Kuchma SL, Griffin EF, O'Toole GA. Minor pilins of the type IV pilus system participate

in the negative regulation of swarming motility. J Bacteriol. 2012;194(19):5388-403.

43.   Rodesney CA, Roman B, Dhamani N, Cooley BJ, Katira P, Touhami A, et al.

Mechanosensing of shear by *Pseudomonas aeruginosa* leads to increased levels of the cyclic-di-

GMP signal initiating biofilm development. Proc Natl Acad Sci U S A. 2017;114(23):5906-11.

44.   Giltner CL, Rana N, Lunardo MN, Hussain AQ, Burrows LL. Evolutionary and

functional diversity of the *Pseudomonas type* IVa pilin island. Env Microbiol. 2011;13(1):250-

64.

45.   Kuchma SL, Brothers KM, Merritt JH, Liberati NT, Ausubel FM, O'Toole GA. BifA, a

cyclic-di-GMP phosphodiesterase, inversely regulates biofilm formation and swarming motility

by *Pseudomonas aeruginosa* PA14. J Bacteriol. 2007;189(22):8165-78.

46. Zhang L, Fritsch M, Hammond L, Landreville R, Slatculescu C, Colavita A, et al.

Identification of genes involved in *Pseudomonas aeruginosa* biofilm-specific resistance to

antibiotics. PLoS ONE. 2013;8(4):e61625.

47. van Tilburg Bernardes E, Charron-Mazenod L, Reading DJ, Reckseidler-Zenteno SL,

Lewenza S. Exopolysaccharide-repressing small molecules with antibiofilm and antivirulence

activity against *Pseudomonas aeruginosa*. Antimicrob Agents Chemother. 2017.

48. Irazoqui JE, Troemel ER, Feinbaum RL, Luhachack LG, Cezairliyan BO, Ausubel FM.

Distinct pathogenesis and host responses during infection of *C. elegans* by *P. aeruginosa* and *S.*

aureus. PLoS Pathog. 2010;6:e1000982.

49. Kilmury SLN, Burrows LL. Type IV pilins regulate their own expression via direct

intramembrane interactions with the sensor kinase PilS. Proc Natl Acad Sci U S A.

2016;113(21):6017-22.

50. Okkotsu Y, Tieku P, Fitzsimmons LF, Churchill ME, Schurr MJ. *Pseudomonas*

aeruginosa AlgR phosphorylation modulates rhamnolipid production and motility. J Bacteriol.

2013;195(24):5499-515.

51. Whitchurch CB, Erova TE, Emery JA, Sargent JL, Harris JM, Semmler AB, et al.

Phosphorylation of the *Pseudomonas aeruginosa* response regulator AlgR is essential for type IV

fimbria-mediated twitching motility. J Bacteriol. 2002;184(16):4544-54.

52. Stacey SD, Williams DA, Pritchett CL. The *Pseudomonas aeruginosa* two-component

regulator AlgR directly activates *rsmA* expression in a phosphorylation independent manner. J

Bacteriol. 2017.

53.   Ma S, Selvaraj U, Ohman DE, Quarless R, Hassett DJ, Wozniak DJ. Phosphorylation-

independent activity of the response regulators AlgB and AlgR in promoting alginate

biosynthesis in mucoid *Pseudomonas aeruginosa*. J Bacteriol. 1998;180(4):956-68.

54.   Kong W, Zhao J, Kang H, Zhu M, Zhou T, Deng X, et al. ChIP-seq reveals the global

regulator AlgR mediating cyclic di-GMP synthesis in *Pseudomonas aeruginosa*. Nucleic Acids

Res. 2015;43(17):8268-82.

55.   Korotkov KV, Hol WG. Structure of the GspK-GspI-GspJ complex from the

enterotoxigenic *Escherichia coli* type 2 secretion system. Nat Struct Mol Biol. 2008;15(5):462-8.

56.   Cisneros DA, Bond PJ, Pugsley AP, Campos M, Francetic O. Minor pseudopilin self-

assembly primes type II secretion pseudopilus elongation. EMBO J. 2012;31(4):1041-53.

57.   Jenal U, Malone J. Mechanisms of cyclic-di-GMP signaling in bacteria. Annu Rev Genet.

2006;40:385-407.

58.   Taylor PK, Van Kessel ATM, Colavita A, Hancock REW, Mah T-F. A novel small RNA

is important for biofilm formation and pathogenicity in *Pseudomonas aeruginosa*. PLoS ONE.

2017;12(8):e0182582.

59.   Darzins A, Chakrabarty AM. Cloning of genes controlling alginate biosynthesis from a

mucoid cystic fibrosis isolate of *Pseudomonas aeruginosa*. J Bacteriol. 1984;159(1):9-18.

60.   Kim D, Forst S. Genomic analysis of the histidine kinase family in bacteria and archaea.

Microbiology. 2001;147(Pt 5):1197-212.

61.   Cody WL, Pritchett CL, Jones AK, Carterson AJ, Jackson D, Frisk A, et al. *Pseudomonas*

aeruginosa AlgR controls cyanide production in an AlgZ-dependent manner. J Bacteriol.

2009;191(9):2993-3002.

62.   Morici LA, Carterson AJ, Wagner VE, Frisk A, Schurr JR, Honer zu Bentrup K, et al.

Pseudomonas aeruginosa AlgR represses the Rhl quorum-sensing system in a biofilm-specific

manner. J Bacteriol. 2007;189(21):7752-64.

63.   Lizewski SE, Schurr JR, Jackson DW, Frisk A, Carterson AJ, Schurr MJ. Identification

of AlgR-regulated genes in *Pseudomonas aeruginosa* by use of microarray analysis. J Bacteriol.

2004;186(17):5672-84.

64.   Intile PJ, Diaz MR, Urbanowski ML, Wolfgang MC, Yahr TL. The AlgZR two-

component system recalibrates the RsmAYZ posttranscriptional regulatory system to inhibit

expression of the *Pseudomonas aeruginosa* type III secretion system. J Bacteriol.

2014;196(2):357-66.

65.   Jones AK, Fulcher NB, Balzer GJ, Urbanowski ML, Pritchett CL, Schurr MJ, et al.

Activation of the *Pseudomonas aeruginosa* AlgU regulon through *mucA* mutation inhibits cyclic

AMP/Vfr signaling. J Bacteriol. 2010;192(21):5709-17.

66.   Lizewski SE, Lundberg DS, Schurr MJ. The transcriptional regulator AlgR is essential

for *Pseudomonas aeruginosa* pathogenesis. Infect Immun. 2002;70(11):6083-93.

67.   Lee DG, Urbach JM, Wu G, Liberati NT, Feinbaum RL, Miyata S, et al. Genomic

analysis reveals that *Pseudomonas aeruginosa* virulence is combinatorial. Genome Biol.

2006;7(10):R90.

68.   Moradali MF, Ghods S, Rehm BHA. *Pseudomonas aeruginosa* lifestyle: a paradigm for

adaptation, survival, and persistence. Front Cell Infect Microbiol. 2017;7:39.

69.   Lebeaux D, Chauhan A, Rendueles O, Beloin C. From *in vitro* to *in vivo* models of

bacterial biofilm-related infections. Pathogens. 2013;2(2):288-356.

70. Kus JV, Tullis E, Cvitkovitch DG, Burrows LL. Significant differences in type IV pilin

allele distribution among *Pseudomonas aeruginosa* isolates from cystic fibrosis (CF) versus non-

CF patients. Microbiology. 2004;150(5):1315-26.

71. Pasloske BL, Joffe AM, Sun Q, Volpel K, Paranchych W, Eftekhar F, et al. Serial isolates

of *Pseudomonas aeruginosa* from a cystic fibrosis patient have identical pilin sequences. Infect

Immun. 1988;56(3):665-72.

72. Schurr MJ, Yu H, Martinez-Salazar JM, Hibler NS, Deretic V. Biochemical

characterization and posttranslational modification of AlgU, a regulator of stress response in

Pseudomonas aeruginosa. Biochem Biophys Res Commun. 1995;216(3):874-80.

73. Martin DW, Schurr MJ, Mudd MH, Govan JR, Holloway BW, Deretic V. Mechanism of

conversion to mucoidy in *Pseudomonas aeruginosa* infecting cystic fibrosis patients. Proc Natl

Acad Sci U S A. 1993;90(18):8377-81.

74. Dower WJ, Miller JF, Ragsdale CW. High efficiency transformation of *E. coli* by high

voltage electroporation. Nucleic Acids Res. 1988;16(13):6127-45.

75. Asikyan ML, Kus JV, Burrows LL. Novel proteins that modulate type IV pilus retraction

dynamics in *Pseudomonas aeruginosa*. J Bacteriol. 2008;190(21):7022-34.

76. Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. Site-directed mutagenesis by

overlap extension using the polymerase chain reaction. Gene. 1989;77(1):51-9.

77. Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE, Tseng BS, et al. Precision-

engineering the *Pseudomonas aeruginosa* genome with two-step allelic exchange. Nat Protoc.

2015;10(11):1820-41.

78. Gallant CV, Daniels C, Leung JM, Ghosh AS, Young KD, Kotra LP, et al. Common β-

lactamases inhibit bacterial biofilm formation. Molec Microbiol. 2005;58(4):1012-24.

79.   Köhler T, Curty LK, Barja F, van Delden C, Pechère J-C. Swarming of *Pseudomonas*

aeruginosa is dependent on cell-to-cell signaling and requires flagella and pili. J Bacteriol.

2000;182(21):5990-6.

80.   Wenderska IB, Chong M, McNulty J, Wright GD, Burrows LL. Palmitoyl-DL-carnitine

is a multitarget inhibitor of *Pseudomonas aeruginosa* biofilm development. ChemBioChem.

2011;12(18):2759-66.

81.   Tan MW, Mahajan-Miklos S, Ausubel FM. Killing of *Caenorhabditis elegans* by

Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. Proc Natl Acad Sci

U S A. 1999;96(2):715-20.

82.   Buensuceso RNC, Nguyen Y, Zhang K, Daniel-Ivad M, Sugiman-Marangos SN,

Fleetwood AD, et al. The conserved tetratricopeptide repeat-containing C-terminal domain of

Pseudomonas aeruginosa FimV is required for its cyclic AMP-dependent and -independent

functions. J Bacteriol. 2016;198(16):2263-74.

83.   Nguyen Y, Harvey H, Sugiman-Marangos S, Bell SD, Buensuceso RNC, Junop MS, et

al. Structural and functional studies of the *Pseudomonas aeruginosa* minor pilin, PilE. J Biol

Chem. 2015;290(44):26856-65.

Figure captions

Fig 1. **PilWXY1 contribute to T4P-independent virulence.**

(A) SK assays for PA14 *pilA*, *fimU*, *pilV*, *pilW*, *pilX*, *pilY1*, and *pilE* mutants. Synchronized L4

worms were seeded onto SK plates and scored for death every 24 h, then plotted as "percent

survival" over the course of the assay. "Day" represents the number of days after L4 on which

the plates were scored. PA14 *fimU* and *pilE* mutants had similar virulence to WT, *pilA* and *pilV*

mutants were slightly less virulent than WT in some trials, and *pilW, pilX,* and *pilY1* mutants

were less virulent than all other strains tested. (B) (A) SK assays for PAO1 *pilA*, *fimU*, *pilV*,

pilW, *pilX*, *pilY1*, and *pilE* mutants. The PAO1 *pilE* mutant had similar virulence to WT, the *pilA*

mutant was slightly less virulent in some trials, and *fimU*, *pilV*, *pilW*, *pilX*, and *pilY1* mutants

were much less virulent. In (A) and (B), asterisks indicate strains that were consistently less

virulent than a *pilA* mutant in all trials by Gehan-Breslow-Wilcoxon test at p = 0.05, n = 3 trials.

Fig 2. PA14 *pilW*, *pilX*, and *pilY1* mutants have swarming motility defects.

SM assays for PA14 *pilA, fimU, pilV, pilW, pilX, pilY1,* and *pilE* mutants. SM assays were

performed in duplicate on 0.5% agar SK plates, and imaged after 24 h incubation at 37°C and 24

h at 22°C. PA14 *pilW*, *pilX*, and *pilY1* mutants had reduced SM relative to WT. Representative

images are shown, n = 3 trials.

Fig 3. PilVWXY1 repress their expression via FimS-AlgR.

(A) BACTH assays for FimS, PilA, and MPs. Protein fusions with T18 and T25 fragments of the

CyaA adenylate cyclase were screened for interactions on MacConkey plates. FimS interacted

with PilA, FimU, PilV, PilW, PilX, and PilE in at least one orientation. Positive (+) or negative

(-) interactions are indicated below each image, n = 3. (B) *fimU* promoter activity in PA14 *pilA*,

fimU, *pilV*, *pilW*, *pilX*, *pilY1*, *pilE*, *fimS*, *algR*, *pilY1 fimS*, or *pilY1 algR* mutants. pMS402-

P*fimU*, containing the *fimU* promoter upstream of the *lux* genes, was introduced into strains of

interest. Assays were set up in technical triplicate, and measurements were taken every 15 min

over 5 h. Loss of *pilV*, *pilW*, *pilX*, or *pilY1* led to highly elevated *fimU* promoter activity. *pilA*

and *fimU* mutants had slightly increased promoter activity relative to WT. Loss of *fimS* or *algR*

reverted *fimU* promoter activity in the *pilY1* mutant to baseline. n = 3 trials.

Fig 4. **AlgR hyperactivation reduces virulence.**

(A) SK assays for PA14 *fimS*, *algR*, *algR*$_{D54A}$, and *algR*$_{D54E}$ mutants. The *fimS*, *algR,* and

algR$_{D54A}$ mutants had WT virulence, while the *algR*$_{D54E}$ mutant was less virulent than WT. (B)

SK assays for PAO1 *algR*, *algR*$_{D54A}$, and *algR*$_{D54E}$ mutants. The *algR* and *algR*$_{D54A}$ mutants had

WT virulence, while the *algR*$_{D54E}$ mutant was less virulent than WT. For (A) and (B), asterisks

indicate strains that were less virulent than WT in all trials, by Gehan-Breslow-Wilcoxon test at

p = 0.05, n = 3 trials.

Fig 5. AlgR promotes biofilm formation and represses virulence.

(A) SK assays for *algR* deletion and overexpression strains. Loss of *algR* had no effect on SK,

while overexpression of pBADGr-*algR* or pBADGr-*algR*$_{D54A}$ reduced virulence at 0.05%

arabinose. Asterisks indicate strains that were consistently less virulent than PA14 + pBADGr in

all trials by Gehan-Breslow-Wilcoxon test at p = 0.05, n = 3 trials. (B) Biofilm assays for *algR*

deletion and overexpression strains. Microtiter plate biofilm assays were performed in liquid SK

media over 24 h, in triplicate. Biofilms were stained with 1% crystal violet then solubilized in

acetic acid. Loss of *algR* had no effect on biofilm formation. When grown at 0.05% arabinose,

overexpression of pBADGr-*algR* or pBADGr-*algR*$_{D54A}$ increased biofilm formation, p < 0.001.

Significance was determined by one-way ANOVA followed by Dunnett post-test relative to WT,

n = 3 trials.

Fig 6. The virulence defect of *pilW*, *pilX*, and *pilY1* mutants is dependent on FimS-AlgR.

SK assays for *pilW*, *pilX*, *pilY1*, *fimS*, *algR*, and *algR*$_{D54A}$ single and double mutants. *fimS*, *algR*,

and *algR*$_{D54A}$ mutants have WT virulence. *pilW*, *pilX*, and *pilY1* have reduced virulence relative

to WT, *fimS*, *algR*, and *algR*$_{D54A}$ mutants. Combination of *pilW*, *pilX*, or *pilY1* mutations with

fimS, *algR*, or *algR*$_{D54A}$ mutations results in virulence equivalent to *fimS*, *algR*, and *algR*$_{D54A}$

single mutants, respectively. All graphs represent 1 trial, separated into 3 graphs for clarity.

Asterisks indicate strains that were consistently less virulent than PA14 by Gehan-Breslow-

Wilcoxon test at $p = 0.05$, $n = 3$.

Fig 7. The swarming motility defect of *pilW*, *pilX*, and *pilY1* mutants is dependent on FimS-

AlgR.

SM assays for *pilW*, *pilX*, *pilY1*, *fimS*, *algR*, and *algR*$_{D54A}$ single and double mutants. The

reduction in SM of *pilW*, *pilX*, and *pilY1* mutants was dependent on FimS-AlgR. *pilW*, *pilX*, and

pilY1 had reduced SM relative to WT. *fimS*, *algR*, and *algR*$_{D54A}$ mutants had WT swarming.

Double mutants swarmed similar to *fimS*, *algR*, and *algR*$_{D54A}$ single mutants. Representative

images are shown, $n = 3$.

Fig 8. Model for regulation of the MP operon via FimS-AlgR.

When PilVWXY1 are absent, FimS promotes phosphorylation of AlgR (phosphate indicated by

yellow star). Phospho-AlgR binds the *fimU* promoter to promote expression of *pilY1* and the MP.

Phospho-AlgR promotes expression of genes associated with chronic infections, and represses

those associated with acute infections. As PilVWXY1 accumulate in the IM, they are detected by

FimS, leading to reduced AlgR phosphorylation. Abbreviations: PilV, V (orange); PilW, W

(cyan); PilX, X (pink); PilY1, Y1 (dark purple); IM, inner membrane.

Supporting information

S1 Fig. Twitching motility of *pilW*, *pilX*, and *pilY1* mutants can be complemented *in trans*.

Colonies were stab-inoculated into 1% agar LB plates, in triplicate. Plates were stained with

crystal violet after 48 h at 30°C. Complementation of PA14 *pilW*, *pilX*, and *pilY1* mutants with

pBADGr-*pilW*, pBADGr-*pilX*, or pBADGr-*pilY1*, respectively, led to increased TM relative to

complementation with pBADGr alone. Numbers indicate percent twitching area relative to WT,

n = 3.

S2 Fig. Virulence of *pilW*, *pilX*, and *pilY1* mutants can be complemented *in trans*.

Complementation of PA14 *pilW*, *pilX*, and *pilY1* mutants with pBADGr-*pilW*, pBADGr-*pilX*, or

pBADGr-*pilY1*, respectively, restored virulence to near-WT levels. Asterisks indicate strains that

were consistently less virulent than WT by Gehan-Breslow-Wilcoxon test at p = 0.05, n = 3.

Individual graphs represent separate trials.

S3 Fig. PA14 and PAO1 produce low levels of biofilm in liquid slow killing media.

Biofilm assays for (A) PA14 and (B) PAO1 *pilA, fimU, pilV, pilW, pilX, pilY1*, and *pilE* mutants.

Very little biofilm formation was detectable in liquid SK media for any strains. There were no

differences in biofilm formation as determined by one-way ANOVA followed by Dunnett post-

test relative to WT at p = 0.05, n = 3.

S4 Fig. PAO1 does not swarm optimally on 0.5% agar slow killing plates.

SM assays for PAO1 *pilA, fimU*, *pilV*, *pilW*, *pilX*, *pilY1*, and *pilE* mutants. SM assays were

performed in duplicate on 0.5% agar SK plates, and imaged after 24 h incubation at 37°C and 24

h at 22°C. All strains formed very small smarming zones, and the patterns were not reproducible

between strains. Representative images are shown, n = 3.

S5 Fig. SadC promotes biofilm formation but is not required for virulence.

(A) Biofilm assays for *sadC* deletion and overexpression strains. PA14 *sadC* biofilm levels were

similar to WT. Expression of SadC *in trans* from a multicopy plasmid led to increased biofilm

formation relative to WT at 0% (due to leaky promoter) and 0.05% arabinose, p < 0.001.

Significance was determined by one-way ANOVA followed by Dunnett post-test relative to

PA14 + pBADGr, n = 3. (B) SK assays for *sadC* deletion and overexpression strains.

Overexpression of SadC led to a subtle but reproducible loss of virulence relative to WT at 0%

and 0.05% arabinose. A *sadC* mutant had WT virulence. Asterisks indicate strains that were

consistently less virulent than PA14 + pBADGr by Gehan-Breslow-Wilcoxon test at p = 0.05, n

= 3.

WT pilA fimU pilV

pilW pilX pilY1 pilE

3 cm

A

T18-T25   FimS-FimS

−   +

+T25 FimS

T18C →   PilA   FimU   PilV   PilW   PilX   PilE

+   +   +   +   +   +

+T18C FimS

T25 →   PilA   FimU   PilV   PilW   PilX   PilE

+   +   +   −   +   +

B

Legend (right side):
- PA14 + P*fimU*
- *pilA* + P*fimU*
- *fimU* + P*fimU*
- *pilV* + P*fimU*
- *pilW* + P*fimU*
- *pilX* + P*fimU*
- *pilY1* + P*fimU*
- *pilE* + P*fimU*
- *fimS* + P*fimU*
- *algR* + P*fimU*
- *pilY1 fimS* + P*fimU*
- *pilY1 algR* + P*fimU*

Y-axis: Relative Luminescence (RLU), 1.0×10^{0} to 1.0×10^{8}

X-axis: Time (min), 0 to 300

A

Percent survival vs Day

No arabinose
- PA14 + pB
- *algR* + pB
- *algR* + *algR*
- *algR* + D54A

0.05% arabinose
- PA14 + pB
- *algR* + pB
- *algR* + *algR* *
- *algR* + D54A *

B

Growth (OD$_{600}$) Biofilm (Absorbance$_{600}$)

Planktonic Growth
Biofilm

***   ***

Arabinose: − − − − + + + +

PA14 + pB
algR + pB
algR + algR
algR + D54A
PA14 + pB
algR + pB
algR + algR
algR + D54A

PA14   *pilW*   *pilX*   *pilY1*

fimS   *pilW fimS*   *pilX fimS*   *pilY1 fimS*

algR   *pilW algR*   *pilX algR*   *pilY1 algR*

$algR_{D54A}$ $pilW\ algR_{D54A}$ $pilX\ algR_{D54A}$ $pilY1\ algR_{D54A}$

3 cm

PA14 + pB	*pilW* + pB	*pilW* + *pilW*	*pilX* + pB	*pilX* + *pilX*	*pilY1* + pB	*pilY1* + *pilY1*
100 ± 13	2 ± 2	95 ± 4	2 ± 2	68 ± 3	5 ± 1	38 ± 4

3 cm