Manuscript submitted to *Applications in Plant Sciences* – Genomic Resources Article

Thomson et al.*, - Lecythidaceae Plastome markers*

Complete plastome sequence from *Bertholletia excelsa* and 23 related

species yield informative markers for Lecythidaceae[1]

Ashley M. Thomson*[2,3], Oscar M. Vargas*[2], Christopher W. Dick[2,4]

[2]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,
MI 48109
[3]Faculty of Natural Resources Management. Lakehead University, Thunder Bay, Ontario,
Canada, P7B 5E1
[4]Smithsonian Tropical Research Institute, Republic of Panama
* These authors contributed equally to this work
14
15 Email addresses: AMT: athomson@lakeheadu.ca
16   OMV: oscarvargash@gmail.com
17   CWD: cwdick@umich.edu
18 Number of words: 3247
19
20 [1] Manuscript received ______; revision accepted _______.
21 [4] Author for correspondence: oscarvargash@gmail.com

ABSTRACT

- *Premise of the study:* The tropical tree family Lecythidaceae has enormous ecological and economic importance in the Amazon basin. Lecythidaceae species can be difficult to identify without molecular data, however, and phylogenetic relationships within and among the most diverse genera are poorly resolved.
- *Methods:* To develop informative genetic markers for Lecythidaceae, we used genome skimming to assemble *de novo* the full plastome of the Brazil nut tree (*Bertholletia excelsa*) and 23 other Lecythidaceae species. Indices of nucleotide diversity and phylogenetic signal were used to identify regions suitable for genetic marker development.
- *Results:* The *B. excelsa* plastome contained 160,472 bp and was arranged in a quadripartite structure. Using the 24 plastome alignments, we developed primers for 10 coding and non-coding DNA regions containing exceptional nucleotide diversity and phylogenetic signal. We additionally developed 19 chloroplast simple sequence repeats (cpSSRs) for population-level studies.
- *Discussion:* The coding region *ycf1* and the spacer *rpl16-rps3* outperformed plastid DNA markers previously used for barcoding and phylogenetics. Used in a phylogenetic analysis, the matrix of 24 plastomes showed with 100% bootstrap support that *Lecythis* and *Eschweilera* are polyphyletic. The plastomes and primers presented in this study will facilitate a broad array of ecological and evolutionary studies in Lecythidaceae.

Keywords: Amazon; *Bertholletia excelsa*; DNA barcoding; genetic markers; Lecythidaceae; plastome; tropical trees.

INTRODUCTION

Lecythidaceae (*sensu latu*) is a pantropical family of trees with three subfamilies:

Foetidioideae, which is restricted to Madagascar; Planchonioideae, found in the tropical forests of

Asia and Africa; and the Neotropical clade Lecythidoideae (Mori et al.*,* 2007), which contains ca.

234 of the ca. 278 known species in the broader family (Mori et al.*,* 2007; Huang et al.*,* 2015;

Mori et al.*,* 2017; Mori, 2017). Neotropical Lecythidaceae are understory, canopy, or emergent

trees with distinctive floral morphology and woody fruit capsules. Among Lecythidaceae species

are the iconic Brazil nut tree, *Bertholletia excelsa* Bonpl.; the oldest documented angiosperm

tree, *Cariniana micrantha* Ducke (dated at >1400 years old in Manaus, Brazil; Chambers et al.*,*

1998); the cauliflorous cannonball tree commonly grown in botanical gardens, *Couroupita*

guianensis Aubl.; and important timber species (e.g. *Carinaria legalis* (Mart.) Kuntze).

Lecythidaceae is the third most abundant family of trees in the Amazon forest, following

Fabaceae and Sapotaceae (ter Steege et al.*,* 2013). The most species-rich genus, *Eschweilera,*

with ca. 99 species (Mori, 2017), is also the most abundant tree genus in the Amazon basin (ter

Steege et al.*,* 2013), and *Eschweilera coriacea* (DC.) S.A.Mori is the most common tree species

in much of Amazonia (ter Steege et al.*,* 2013). Lecythidaceae provide important ecological

services such as carbon sequestration, and food resources for pollinators (bats and large bees) and

seed dispersers (monkeys and agouties) (Prance and Mori, 1979; Mori and Prance, 1990).

64   Tools for species-level identification and phylogenetic analyses of Lecythidaceae could

significantly advance research on Amazon tree diversity. However, despite their ease of

identification at the family level, species-level identification of many Lecythidaceae (especially

Eschweilera) is notoriously difficult when based on sterile (i.e. without fruit or floral material)

herbarium specimens, and flowering of individual trees often occurs only at multi-year intervals

(Mori and Prance, 1987). As a complement to other approaches, DNA barcoding (Dick and

Kress, 2009; Dexter et al.*,* 2010) may help to identify species or clades of Lecythidaceae.

71   A combination of two protein-coding plastid regions (*matK* and *rbcL*) have been proposed

as core plant DNA barcodes (Hollingsworth et al.*,* 2009), although other coding and non-coding

plastome regions (*psbA-trnH, rpoB, rpoC1, trnL, ycf5*) and the internal transcribed spacer (ITS)

of nuclear ribosomal genes, have been recommended as supplemental barcodes for vascular

plants (Kress et al., 2005; Lahaye et al.*,* 2008; Li et al.*,* 2011). However, an evaluation of a subset

of these markers (ITS, *psbA-trnH, matK, rbcL, rpoB, rpoC1, trnL*) on Lecythidaceae in French

Guiana (Gonzales et al.*,* 2009) showed poor performance for species identification. Furthermore,

the use of traditional markers (plastid *ndhF, trnL-F,* and *trnH-psbA*, and nuclear ITS) for

phylogenetic analysis has produced weakly supported trees (Mori et al., 2007; Huang et al., 2015)

indicating a need to develop more informative markers and/or increase molecular sampling.

The main objectives of this study were to (1) assemble, annotate, and characterize the first

complete plastome sequence of Lecythidaceae, from the iconic Brazil nut tree *Bertholletia*

excelsa; (2) obtain a robust backbone phylogeny for the Neotropical clade using newly-

assembled draft plastome sequences for an additional 23 species; and (3) develop a novel set of

informative molecular markers for DNA barcoding and broader evolutionary studies.

86   **METHODS**

Plant material and DNA library preparation—We performed genomic skimming on

24 Lecythidaceae species, including 23 Lecythidoideae and one outgroup species (*Barringtonia*

edulis Seem.) from the Planchonioideae. The sampling included all 10 Lecythidoideae genera

(Appendix 1). Silica-dried leaf tissue from herbarium-vouchered collections was collected by

Scott Mori and colleagues and loaned by the New York Botanical Garden. Total genomic DNA

was extracted from 20 milligrams of dried leaf tissue using the NucleoSpin Plant II extraction kit

(Machery-Nagel, Bethlehem, PA, USA) with SDS lysis buffer. Prior to DNA library preparation,

5 micrograms of total DNA were fragmented using a Covaris S-series sonicator (Covaris, Inc.

Woburn, MA, USA), following the manufacturer's protocol, to obtain ca. 300 bp insert-sizes. We

prepared the sequencing library using the NEBNext DNA library Prep Master Mix and Multiplex

Oligos for Illumina Sets (New England BioLabs Inc. Ipswich, MA, USA) according to the

manufacturer's protocol. Size-selection was carried out prior to PCR using Pippin Prep (Sage

Science, Beverly, MA, USA). Molecular mass of the finished paired-end library was quantified

using an Agilent 2100 Bioanalyzer (Agilent Technologies Inc., Santa Clara, CA, USA) and by

101 qPCR using an ABI PRISM 7900HT (ThermoFisher Scientific, Waltham, MA, USA) at the

102 University of Michigan DNA Sequencing Core (Ann Arbor, MI, USA). We sequenced the

103 libraries on one lane of the Illumina HiSeq 2000 (Illumina Inc., San Diego, CA, USA) with a

104 paired-read length of 100 bp.

105   *Plastome assembly*—Illumina adaptors and barcodes were excised from raw reads using

106 Cutadapt v.1.4.2 (Martin, 2011). Reads were then quality-filtered using Prinseq v. 0.20.4

107 (Schmieder and Edwards, 2011), which trimmed 5' and 3' sequence ends with Phred quality

108 score <20 and removed all trimmed sequences <50 bp in length, with >5% ambiguous bases, or

109 with mean Phred quality score <20. A combination of *de novo* and reference-guided approaches

110 were used to assemble the plastomes. First, chloroplast reads were separated from the raw read

111 pool by Blast-searching all raw reads against a database consisting of all complete angiosperm

112 plastome sequences available on GenBank (accessed in 2014). Any aligned reads with an e-value

113 $<1^{-5}$ were retained for subsequent analysis. The filtered chloroplast reads were *de novo*

114 assembled using Velvet v.7.0.4 (Zerbino and Birney, 2008) with kmer values of 71, 81, and 91

115 using a low-coverage cutoff of 5 and minimum contig length of 300. The assembled contigs were

116 then mapped to a reference genome (see below) using Geneious v. R8 (Kearse et al*.,* 2012) to

117 determine their order and direction using the reference-guided assembly tool with medium

118 sensitivity and iterative fine-tuning options. Finally, raw reads were iteratively mapped onto the

119 draft genome assembly to extend contigs and fill gaps using low-sensitivity reference-guided

120 assembly in Geneious. We first assembled the draft genome of *Bertholletia excelsa*; the

121 plastomes of the remaining 23 species were assembled subsequently using the plastome of *B.*

122 *excelsa* as a reference. The *B. excelsa* plastome was annotated using DOGMA (Wyman et al*.,*

123 2004) with the default settings for chloroplast genomes. Codon start and stop positions were

124 determined using the open reading frame finder in Geneious and by comparison with the

plastome sequence of *Camellia sinensis* var. *pubilimba* Hung T. Chang (Genbank ID:

KJ806280). A circular representation of the *B. excelsa* plastome was made using OGDraw V1.2

(Lohse et al., 2007). The complete annotated plastome of *B. excelsa* and the draft plastomes of

the remaining 23 Lecythidaceae species sampled were deposited in GenBank (Appendix 1).

Identification of molecular markers—Chloroplast simple sequence repeats (cpSSRs) in

B. excelsa were identified using the Phobos Tandem Repeat Finder v.3.3.12 (Mayer, 2010) by

searching for uninterrupted repeats of nucleotide units of 1 to 6 bp in length, with thresholds of

$\geq$12 mononucleotide, $\geq$6 dinucleotide, and $\geq$4 trinucleotide, and $\geq$3 tetra-, penta-, and

hexanucleotide repeats (Sablok et al., 2015). We developed primers to amplify the cpSSRs using

Primer 3 v.2.3.4 (Untergasser et al., 2012) with the default options and setting the PCR product

size range between 100 and 300 bp.

The 24 plastomes were aligned with MAFFT v.7.017 (Katoh et al., 2002) and scanned for

regions of high nucleotide diversity, π (Nei 1987), using a sliding window analysis implemented

in DNAsp v.5.10.1 (Librado and Rozas, 2009) with a window and a step size of 600 bp. Levels of

nucleotide diversity were plotted using the generic function "plot" in R (R core development

group), and windows with values over the 95[th] percentile were considered of high π.

Taking into account that DNA barcodes can also be used in phylogenetic analyses and

because regions with high π do not necessarily have high phylogenetic signal (e.g. unalignable

hypervariable regions), we employed a log-likelihood approach modified from Walker et al.

(2017) to identify phylogenetically influential regions. First, we inferred a phylogenetic tree with

the plastome alignment (including only one inverted repeat) by performing 100 independent

maximum likelihood (ML) searches using a GTRGAMMA model with RAxML v. 8.2.9

(Stamatakis, 2014). Those searches resulted in the same topology that was subsequently

annotated with the summary from 100 bootstraps using "sumtrees.py" v.4.10 (Sukumaran and

Holder, 2010). Then, we calculated the site-specific log-likelihood in the alignment over the

plastome phylogeny and calculated their differences site-wise to the averaged log-likelihood per

site of 1000 randomly permuted trees (tips were randomly shuffled). Log-likelihood scores were

calculated with RAxML under using a GTRGAMMA model. The site-wise log-likelihood

differences (LD) were calculated using 600 bp non-overlapping windows with a custom R script

(see below). We interpreted greater LD as an indication of greater phylogenetic signal, and

windows with LD above the 95[th] percentile were considered to have exceptional phylogenetic

signal.

157   Primers flanking the top ten regions with high π were designed using Primer 3 with

default program options. We employed a maximum product size of 1300 bp because lower

cutoffs values (i.g. 600 bp) made the primer design extremely challenging due to the lack of

conserved regions. Primers were designed to amplify across all 23 Neotropical species without

the use of degenerate bases. However, primers with a small number of degenerate bases were

permitted for some regions where primer development otherwise would not have been possible

due to high sequence variability in the priming sites. We investigated the potential of our markers

to produce robust phylogenies by calculating individual gene trees in RAxML v.8.2.9 in an ML

search with 100 rapid bootstraps (option "-f a") using the GTRGAMMA model. To evaluate the

number of markers needed to obtain a resolved tree with an average of ~90 bootstrap support

(BS), we first concatenated the two markers with the highest π and inferred a tree; subsequently

we added the marker with the next highest π score. We iterated this process until we obtained a

matrix with each of the 10 markers developed. For every tree obtained, we calculated its average

BS and its Robinson-Foulds distance (RF) (Robinson and Foulds, 1981) from the plastome

phylogeny, using a custom R script employing the packages APE (Paradis et al., 2004) and

Phangor (Schliep 2011). Scripts and alignments used for this study can be found at

https://bitbucket.org/oscarvargash/lecythidaceae_plastomes.

174   **RESULTS**

175   ***Lecythidaceae plastome features***—The sequenced plastome of *Bertholletia excelsa*

contained 160,472 base pairs and 115 genes, of which 4 were rRNAs and 30 were tRNAs (Fig. 1,

Table 1). The arrangement of the *B. excelsa* plastome had a typical angiosperm quadripartite

structure with a single copy region of 85,830 bp, a small single copy region of 16,670 bp, and

two inverted of repeats of 27,481 bp each. Relative to *Camellia sinensis* var. *pubilimba*, we found

no gene gain/losses in *B. excelsa*. The only structural difference found is that *B. excelsa* contains

the sequential genes *trnH-GUG*, *rps3*, *rpl22*, and *rps19* in the inverted repeat while *C. sinensis*

var. *pubilimba* contains these genes in the large single copy region. Similarly, no gene gain/losses

are found when *B. excelsa* is compared to other neotropical Lecythidaceae plastomes assembled

herein (Table 2). In addition to *B. excelsa,* the plastome of *Eschweilera alata* was also completely

assembled; the coverage for the remaining plastomes ranged between 85% and 99.60%

(Appendix 1).

187   ***Identification of molecular markers***—Within the plastome of *Bertholletia excelsa* we

found 23 cpSSRs, 22 of which were in non-coding regions and one in the *ndhD* coding region.

We designed 19 primer pairs with an acceptable product length, annealing temperature, and GC

content for cpSSRs located in noncoding regions (Table 3). π exceeded the 95[th] percentile for

nine 600 bp windows (Fig. 2A, Table 4, 5). Similarly, 13 windows were over the 95[th] percentile

for LD (Fig. 2B, Table 4, 5) indicating high phylogenetic signal. While most of the informative

windows were in non-coding regions, two consecutive regions were positioned in the *ycf1* gene.

Six windows contained both high π and LD. As expected, high π and greater LD largely agreed.

Based on the rank of the windows obtained for π, we developed primers for the following regions

(ordered from high to low π): *ycf1, rpl16-rps3, psbM-trnD, ccsA-ndhD, trnG-psaB, petD-rpoA,*

psbZ-trnfM, trnE-trnT, and *trnT-psbD* (Table 6).

198   ***Phylogenetics of the plastomes and the developed markers***—The ML analysis of the

plastome alignment for Lecythidaceae (145,487 sites) yielded a fully resolved phylogeny with

high BS for all clades (Fig. 3). Of the genera in which the sampling included multiples species,

Eschweilera and *Lecythis* were polyphyletic, while *Allantoma, Corythophora, Couratari,* and

Gustavia were monophyletic (*Bertholletia* is monospecific, and only one species of *Couroupita,*

Cariniana, and *Grias* were included in the analysis). The trees obtained from individual markers

with high π had an average BS of 73 throughout their nodes, while the trees obtained from two or

more concatenated regions had an average BS of 89 (Fig. 4A, Fig. S1). None of the gene trees,

single or combined (Fig. S1), recovered the topology obtained using the complete plastome

matrix (none of the gene trees obtained a RF=0, Fig. 4B). In general, matrices with concatenated

markers (mean RF=6) outperformed single markers (mean RF = 13.8) (Fig. 4B).

DISCUSSION

210   ***Genetic markers from the Lecythidaceae plastome***—We are publishing the first full

plastome for Lecythidaceae, including high-depth coverage of the Brazil nut tree, *Bertholletia*

excelsa, and 23 draft genomes representing all Lecythoideae genera and a Paleotropical outgroup

taxon. We found no significant gene losses or major rearrangements when the plastome of

Bertholletia excelsa was compared with that of *Camellia sinensis* var. *pubilimba*, a closely

related plastome (Theaceae).

We inferred a robust backbone phylogeny for Lecythoideae using the 24 aligned

plastomes. All nodes in our topology had 100% bootstrap support except for a node that connects

three closely related species of *Eschweilera* (Fig. 3). The topology agreed with previous but

weakly supported (<50% BS) Lecythidaceae phylogenies, based on chloroplast and nuclear

internal transcribed spacer (ITS) sequences (Mori et al., 2007, Huang et al., 2015), indicating that

Eschweilera and *Lecythis* are polyphyletic. Although the polyphyly of these two genera is well

supported with all available data, some inferred species-level relationships may change with

increased taxonomic sampling and the inclusion of nuclear genomic data.

We measured nucleotide diversity (π) and a proxy for phylogenetic signal using a log-

likelihood approach (LD) modified from Walker et al. (2017). These calculations helped us to

evaluate the performance of specific chloroplast regions as potential phylogenetic markers. The

core plant DNA barcodes *matK* and *rbcL* did not exhibit high π or LD in our analysis (Table 5).

Of the secondary plastome barcodes mentioned in the literature (*rpoC1, rpoB, trnL, psbA-trnH;*

Kress et al, 2005, Lahaye et al., 2008, Hollingsworth 2009, Li et al., 2011) only *psbA-trnH*

showed high LD (Table 5) although it did not exhibit exceptionally high values of π. In contrast,

the regions *ycf1, rpl16-rps3, psbM-trnD, ccsA-ndhD, trnG-psaB, petD-rpoA, psbZ-trnfM, trnE-*

trnT, and *trnT-psbD* displayed the highest values of π and LD and therefore outperformed all the

previously proposed plant DNA barcodes.

Phylogenetic trees calculated from concatenated marker sets (based on the π rank)

outperformed single regions in terms of support (BS) and accuracy (RF) (Fig. 4). In fact, tree

topologies using single markers deviated from the complete plastome tree (mean RF= 13.8). The

best performing concatenated matrix contained all 10 regions for which we developed primers.

However, the combination of *ycf1* and *rpl16−rps3* produced an average BS ~90 (Fig. 4A) with

reasonable accuracy (RF = 4, Fig. 4B); we conclude that these two regions, amplified in three

PCRs (Table 6), are promising markers for DNA barcoding, phylogeny, and phylogeography in

Lecythidaceae. Although barcoding efficiency in species-rich clades (i.e. *Eschweilera/Lecythis)*

might decline with the addition of more samples, *ycf1* and *rpl16–rps3* effectively distinguished

between three closely-related species within the *E. parvifolia* clade (see branch lengths in Fig.

S1), suggesting that these markers might effectively distinguish between many other closely

related species. Our results and conclusions agree with those of Dong et al. (2015) who proposed

ycf1 as a universal barcode for land plants.

The 19 cpSSR markers developed for noncoding portions of the *B. excelsa* plastome

provide a useful resource for population genetic studies. Because of their fast stepwise mutation

rate relative to SNPs, cpSSRs can also be used for finer grain phylogeographic analyses (e.g.

Lemes et al., 2010; Twyford et al., 2013). This may be especially useful for species that exhibit

little geographic structuring across parts of their ranges. Because they are maternally transmitted

and can be variable within populations, the cpSSRs may also be used to track dispersal of seeds

and seedlings relative to the maternal source trees.

Because of their high level of polymorphism and phylogenetic signal content, we

anticipate using the cpDNA markers presented here to study the phylogeography of widespread

Lecythidaceae species, such as *Couratari guianensis* and *Eschweilera coriacea,* which range

from the Amazon basin into Central America.

Barcoding of tropical trees—DNA barcoding of tropical trees has been useful for several

applications (Dick and Kress, 2009), including community phylogenetic analyses (Kress et al.,

2009), inferring the species identity of the gut content (diet) of herbivores (García-Robledo et al.,

2013), and for species identification of seedlings (Gonzalez et al., 2009). The power of DNA

barcodes to discriminate among species should be high if the studied species are distantly related;

for example, Kress et al. (2009) were able to discriminate 281 of 296 tree and shrub species from

Barro Colorado Island (BCI) using standard DNA barcodes, but they were not able to

discriminate among some congeneric species in the species-rich genera *Inga* (Fabaceae), *Ficus*

(Moraceae), and *Piper* (Piperaceae). Gonzales et al. (2009) encountered similar challenges with

Eschweilera species in their study of trees and seedlings in Paracou, French Guiana. The latter

study tested a wide range of putative DNA barcode regions (*rbcLa*, *rpoC1*, *rpoB*, *matK*, *trnL*,

psbA-trnH, ITS) but did not include the markers presented in this paper.

Limitations of plastome markers for phylogeny and species ID—These newly-identified

plastome markers are not free of limitations. First, plastome-based phylogenies should be

interpreted with caution, as they can disagree with nuclear markers and species trees due to

introgression and or lineage sorting issues (Rieseberg and Soltis, 1997; Sun et al., 2015; Vargas

et al., 2017). These same processes limit the cpDNA for species identification. For example,

cpDNA haplotypes of *Nothofagus*, *Eucalyptus*, *Quercus*, *Betula*, and *Acer* were more strongly

determined by geographic location than by species-identity due to the occurrence of localized

introgression within these groups (Petit et al., 1993; Palme et al., 2004; Saeki et al., 2011;

Premoli et al., 2012; Nevill et al., 2014; Thomson et al., 2015). The occurrence of haplotype

sharing in closely-related Lecythidacae species has, to date, not been examined at a large scale

and it is therefore not possible to conclude to what extent introgression or incomplete lineage

sorting might affect this group. We suggest that future studies utilizing cpDNA barcodes for

Neotropical Lecythidaceae test species from several shared geographic localities to examine to

what extent haplotypes tend to be shared among species at the same localities. Nuclear DNA

markers may also be used to examine phylogenetic incongruence and identify cases where

introgression might have occurred.

ACKNOWLEDGEMENTS

289   National Science Foundation (DEB 1240869 and FESD Type I 1338694 to CD) and the

290 University of Michigan (Associate Professor Award to CWD) provided financial support for this

291 work. We would like to thank Scott Mori, Gregory Stull, Caroline Parins-Fukuchi, Joseph

292 Walker, and three anonymous reviewers for their useful comments; and Scott Mori and the New

293 York Botanical Garden for providing access to curated DNA samples of Lecythidaceae.

LITERATURE CITED:

CHAMBERS, J.Q., N. HIGUCHI, AND J.P. SCHIMEL. 1998. Ancient trees in Amazonia. *Nature* 391:

135–136. http://dx.doi.org/10.1038/34325.

DEXTER, K.G., T.D. PENNINGTON, AND C.W. CUNNINGHAM. 2010. Using DNA to assess errors in

tropical tree identifications: How often are ecologists wrong and when does it matter?

Ecological Monographs 80: 267–286.

DICK, C.W., AND W.J. KRESS. 2009. Dissecting tropical plant diversity with forest plots and a

molecular toolkit. *BioScience* 59: 745–755. doi:10.1525/bio.2009.59.9.6

DONG, W., C. XU, C. LI, J. SUN, Y. ZUO, S. SHI, T. CHENG, ET AL. 2015. *ycf1*, the most promising

plastid DNA barcode of land plants. *Scientific Reports* 5: 8348. doi:10.1038/srep08348

GARCÍA-ROBLEDO, C., D.L. ERICKSON, C.L. STAINES, T.L. ERWIN, AND W.J. KRESS. 2013.

Tropical Plant-Herbivore Networks: Reconstructing Species Interactions Using DNA

Barcodes. *PLoS ONE* 8: e52967. doi:10.1371/journal.pone.0052967

GONZALEZ, M.A., C. BARALOTO, J. ENGEL, S.A. MORI, P. PÉTRONELLI, B. RIÉRA, A. ROGER, ET

AL. 2009. Identification of Amazonian trees with DNA barcodes. *PLoS ONE* 4: e7483.

doi:10.1371/journal.pone.0007483

HOLLINGSWORTH, P.M., L.L. FORREST, J.L. SPOUGE, M. HAJIBABAEI, S. RATNASINGHAM, M. VAN

DER BANK, M.W. CHASE, ET AL. 2009. A DNA barcode for land plants. *Proceedings of the*

National Academy of Sciences of the United States of America 106: 12794–129797.

doi:10.1073/pnas.0905845106

HUANG, Y.Y., S.A. MORI, AND L.M. KELLY. 2015. Toward a phylogenetic-based generic

classification of Neotropical Lecythidaceae–I. Status of *Bertholletia*, *Corythophora*,

Eschweilera and *Lecythis*. *Phytotaxa* 203: 85–121. doi:10.11646/phytotaxa.203.2.1

KATOH, K., K. MISAWA, K. KUMA, AND T. MIYATA. 2002. MAFFT: a novel method for rapid

multiple sequence alignment based on fast Fourier transform. *Nucleic acids research* 30:

3059–3066. doi:10.1093/nar/gkf436

KEARSE, M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG, S. STURROCK, S. BUXTON, ET

AL. 2012. Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649.

doi:10.1093/bioinformatics/bts199

KRESS, W.J., D.L. ERICKSON, F.A. JONES, N.G. SWENSON, R. PEREZ, O. SANJUR, AND E.

BERMINGHAM. 2009. Plant DNA barcodes and a community phylogeny of a tropical forest

dynamics plot in Panama. *Proceedings of the National Academy of Sciences of the United*

States of America 106: 18621–6. doi:10.1073/pnas.0909820106

KRESS, W.J., K.J. WURDACK, E.A. ZIMMER, L.A. WEIGT, AND D.H. JANZEN. 2005. Use of DNA

barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences of*

the United States of America 102: 8369–8374. doi:10.1073/pnas.0503123102

LAHAYE, R., M. VAN DER BANK, D. BOGARIN, J. WARNER, F. PUPULIN, G. GIGOT, O. MAURIN, ET

AL. 2008. DNA barcoding the floras of biodiversity hotspots. *Proceedings of the National*

Academy of Sciences of the United States of America 105: 2923–2928.

doi:10.1073/pnas.0709936105

LEMES, M.R., C.W. DICK, C. NAVARRO, A.J. LOWE, S. CAVERS, AND R. GRIBEL. 2010.

Chloroplast DNA microsatellites reveal contrasting phylogeographic structure in mahogany

(*Swietenia macrophylla* King, Meliaceae) from Amazonia and Central America. *Tropical*

Plant Biology 3: 40–49. doi:10.1007/s12042-010-9042-5

LI, D.-Z., L.-M. GAO, H.-T. LI, H. WANG, X.-J. GE, J.-Q. LIU, Z.-D. CHEN, ET AL. 2011.

Comparative analysis of a large dataset indicates that internal transcribed spacer (ITS)

should be incorporated into the core barcode for seed plants. *Proceedings of the National*

Academy of Sciences of the United States of America 108: 19641–19646.

doi:10.1073/pnas.1104551108

LIBRADO, P., and J. ROZAS. 2009. DnaSP v5: A software for comprehensive analysis of DNA

polymorphism data. *Bioinformatics* 25: 1451–1452. doi:10.1093/bioinformatics/btp187

LOHSE, M., O. DRECHSEL, and R. BOCK. 2007. OrganellarGenomeDRAW (OGDRAW): A tool

for the easy generation of high-quality custom graphical maps of plastid and mitochondrial

genomes. *Current Genetics* 52: 267–274. doi:10.1007/s00294-007-0161-y

MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17: 10–12. doi:10.14806/ej.17.1.200

MAYER, C. 2010. Phobos. Available from: http://www.rub.de/spezzoo/cm/cm_phobos.htm

MORI, S.A., E.A. KIERNAN, N.P. SMITH, L.M. KELLEY, Y.-Y. HUANG, G.T. PRANCE, and B.

THIERS. 2017. Observations on the phytogeography of the Lecythidaceae clade (Brazil nut

family). *Phytoneuron* 30: 1–85.

MORI, S.A., AND G.T. PRANCE. 1987. A Guide to Collecting Lecythidaceae. *Annals of the*

Missouri Botanical Garden 74: 321–330.

MORI, S.A., AND G.T. PRANCE. 1990. Lecythidaceae. Part II. The zygomorphic-flowered New

World genera (*Couroupita*, *Corythophora*, *Bertholletia*, *Couratari*, *Eschweilera*, &

Lecythis). *Flora Neotropica* 21: 1–376.

MORI, S.A., C.H. TSOU, C.C. WU, B. CRONHOLM, AND A.A. ANDERBERG. 2007. Evolution of

Lecythidaceae with an emphasis on the circumscription of Neotropical genera: Information

from combined *ndhF* and *trnL-F* sequence data. *American Journal of Botany* 94: 289–301.

doi:10.3732/ajb.94.3.289

MORI, S.A. 2017. The Lecythidaceae pages. http://sweetgum.nybg.org/science/projects/lp/

Nei, M. 1987. *Molecular Evolutionary Genetics*. New York: Columbia University Press.

NEVILL, P.G., T. DESPRÉS, M.J. BAYLY, G. BOSSINGER, AND P.K. ADES. 2014. Shared

phylogeographic patterns and widespread chloroplast haplotype sharing in *Eucalyptus*

368 species with different ecological tolerances. *Tree Genetics and Genomes* 10: 1079–1092.

369 doi:10.1007/s11295-014-0744-y

370 PALME, A.E., Q. SU, S. PALSSON, AND M. LASCOUX. 2004. Extensive sharing of chloroplast

371 haplotypes among European birches indicates hybridization among *Betula pendula*, *B.*

372 *pubescens* and *B. nana*. *Molecular Ecology* 13: 167–178. doi:10.1046/j.1365-

373 294X.2003.02034.x

374 PARADIS, E., J. CLAUDE, AND K. STRIMMER. 2004. APE: Analyses of phylogenetics and evolution

375 in R language. *Bioinformatics* 20: 289–290. doi:10.1093/bioinformatics/btg412

376 PETIT, R.J., A. KREMER, AND D.B. WAGNER. 1993. Geographic structure of chloroplast DNA

377 polymorphisms in European oaks. *Theoretical and Applied Genetics: International Journal*

378 *of Plant Breeding Research* 87: 122–128. doi:10.1007/BF00223755

379 PRANCE, G.T., AND S.A. MORI. 1979. Lecythidaceae–Part I. The actinomorphic-flowered New

380 World Lecythidaceae (*Asteranthos, Gustavia, Grias, Allantoma, & Cariniana*). *Flora*

381 *Neotropica Monograph* 21: 1–270.

382 PREMOLI, A.C., P. MATHIASEN, M.C. ACOSTA, and V.A. RAMOS. 2012. Phylogeographically

383 concordant chloroplast DNA divergence in sympatric *Nothofagus* s.s. How deep can it be?

384 *New Phytologist* 193: 261–275. doi:10.1111/j.1469-8137.2011.03861.x

385 R CORE TEAM. 2017. R: A Language and Environment for Statistical Computing. Available from:

386 https://www.r-project.org/

RIESEBERG, L.H., AND D.E. SOLTIS. 1991. Phylogenetic consequences of cytoplasmic gene flow

in plants. *Evolutionary Trends in Plants* 5: 64–84.

ROBINSON, D.F., AND L.R. FOULDS. 1981. Comparison of phylogenetic trees. *Mathematical*

Biosciences 53: 131–147. doi:10.1016/0025-5564(81)90043-2

SABLOK, G., G. V. PADMA RAJU, S.B. MUDUNURI, R. PRABHA, D.P. SINGH, V. BAEV, G.

YAHUBYAN, ET AL. 2015. ChloroMitoSSRDB 2.00: More genomes, more repeats, unifying

SSRs search patterns and on-the-fly repeat detection. *Database* 2015: 1–10.

doi:10.1093/database/bav084

SAEKI, I., C.W. DICK, B. V. BARNES, AND N. MURAKAMI. 2011. Comparative phylogeography of

red maple (*Acer rubrum* L.) and silver maple (*Acer saccharinum* L.): Impacts of habitat

specialization, hybridization and glacial history. *Journal of Biogeography* 38: 992–1005.

doi:10.1111/j.1365-2699.2010.02462.x

SCHLIEP, K.P. 2011. phangorn: Phylogenetic analysis in R. *Bioinformatics* 27: 592–593.

doi:10.1093/bioinformatics/btq706

SCHMIEDER, R., AND R. EDWARDS. 2011. Quality control and preprocessing of metagenomic

datasets. *Bioinformatics* 27: 863–864. doi:10.1093/bioinformatics/btr026

STAMATAKIS, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* 30: 1312–1313. doi:10.1093/bioinformatics/btu033

SUKUMARAN, J., AND M.T. HOLDER. 2010. DendroPy: A Python library for phylogenetic

computing. *Bioinformatics* 26: 1569–1571. doi:10.1093/bioinformatics/btq228

SUN, M., D.E. SOLTIS, P.S. SOLTIS, X. ZHU, J.G. BURLEIGH, AND Z. CHEN. 2015. Deep

phylogenetic incongruence in the angiosperm Rosidae clade. *Molecular Phylogenetics and*

Evolution 83: 156–166. doi:10.1016/j.ympev.2014.11.003

TER STEEGE, H., N.C.A. PITMAN, D. SABATIER, C. BARALOTO, R.P. SALOMÃO, J.E. GUEVARA,

O.L. PHILLIPS, ET AL. 2013. Hyperdominance in the Amazonian tree flora. *Science* 342: 325–

342. doi:10.1126/science.1243092

THOMSON, A.M., C.W. DICK, AND S. DAYANANDAN. 2015. A similar phylogeographical structure

among sympatric North American birches (*Betula*) is better explained by introgression than

by shared biogeographical history. *Journal of Biogeography* 42: 339–350.

doi:10.1111/jbi.12394

TWYFORD, A.D., C.A. KIDNER, N. HARRISON, AND R.A. ENNOS. 2013. Population history and

seed dispersal in widespread Central American *Begonia* species (Begoniaceae) inferred from

plastome-derived microsatellite markers. *Botanical Journal of the Linnean Society* 171:

260–276.

UNTERGASSER, A., I. CUTCUTACHE, T. KORESSAAR, J. YE, B.C. FAIRCLOTH, M. REMM, AND S.G.

ROZEN. 2012. Primer3-new capabilities and interfaces. *Nucleic Acids Research* 40: 1–12.

doi:10.1093/nar/gks596

VARGAS, O.M., E.M. ORTIZ, and B.B. SIMPSON. 2017. Conflicting phylogenomic signals reveal a

pattern of reticulate evolution in a recent high-Andean diversification (Asteraceae: Astereae:

Diplostephium). *New Phytologist* 214: 1736–1750. doi:10.1111/nph.14530

WALKER, J.F., J.W. BROWN, AND S.A. SMITH. 2017. Analyzing contentious relationships and

outlier genes in phylogenomics. *bioRxiv*. http://dx.doi.org/10.1101/115774

WYMAN, S.K., R.K. JANSEN, AND J.L. BOORE. 2004. Automatic annotation of organellar genomes

with DOGMA. *Bioinformatics* 20: 3252–3255. doi:10.1093/bioinformatics/bth352

ZERBINO, D.R., AND E. BIRNEY. 2008. Velvet: algorithms for de novo short read assembly using

de Bruijn graphs. *Genome Research* 18: 821–829. doi:10.1101/gr.074492.107

DATA ACCESSIBILITY:

DNA sequences: Genbank accessions MF359935–MF359958, BioProject SUB2740669

Plastome alignment, gene alignments, trees, and R code:

https://bitbucket.org/oscarvargash/lecythidaceae_plastomes

TABLES

TABLE 1. Genes contained within the chloroplast genome of *Bertholletia excelsa*.

Function	Gene group	Gene name
Self-replication	Ribosomal proteins (LSU)	*rpl2, rpl14, rpl16, rpl20, rpl22, rpl23, rpl32, rpl33, rpl36*
	Ribosomal proteins (SSU)	*rps2, rps3, rps4, rps7, rps8, rps11, rps12, rps14, rps15, rps16, rps18, rps19*
	RNA polymerase subunits	*rpoA, rpoB, rpoC1, rpoC2*
	Ribosomal RNAs	*rrn4.5, rrn5 rrn16, rrn23*
	Transfer RNAs	*trnA-UGC, trnC-GCA, trnD-GUC, trnE-UUC, trnF-GAA, trnG-GCC, trnG-UCC, trnH-GUG, trnI-CAU, trnI-GAU, trnK-UUU, trnL-CAA, trnL-UAA, trnL-UAG, trnfM-CAU, trnM-CAU, trnN-GUU, trnP-UGG, trnQ-UUG, trnR-AGC, trnR-UCU, trnS-GCU, trnS-GGA, trnS-UGA, trnT-GGU, trnT-UGU, trnV-GAC, trnV-UAC, trnW-CCA, trnY-GUA*
Photosynthesis	Photosystem I	*psaA, psaB, psaC, psaI, psaJ*
	Photosystem II	*psbA, psbB, psbC, psbD, psbE, psbF, psbH, psbI, psbJ, psbK, psbL, psbM, psbN, psbT, psbZ*
	NADH dehydrogenase	*ndhA, ndhB,ndhC, ndhD, ndhE, ndF, ndhG, ndhH, ndhI, ndhJ, ndhK*
	Cytochrome b/f complex	*petA, petB, petD, petG, petL, petN*
	ATP synthase	*atpA, atpB, atpE, atpF, atpH, atpI*
Other genes	RubisCO large subunit	*rbcL*
	Subunit of acetyl-CoA-carboxylase	*accD*
	Envelope membrane protein	*cemA*
	Protease	*clpP*
	c-type cytochrome synthase	*ccsA*
	Translational initiation factor	*infA*
	Maturase	*matK*
Unknown function	Hypothetical chloroplast reading frames	*ycf1, ycf2, ycf3, ycf4, yc15*

441   TABLE 2. Comparison for plastome subunits for the samples for which the inverted repeats (IR)

442   were completely assembled. Length and GC content of the large single copy (LSC) and small

443   single copy (SSC) regions in partial plastomes are estimates only.

Species	LSC length (bp)	SSC Length (bp)	IR Length (bp)	GC content (%)	Protein-coding genes	rRNAs	tRNAs
Allantoma decandra	85,269	18,738	27,618	36.9	81	4	30
Allantoma lineata	85,119	18,756	27,635	36.9	81	4	30
Bertholletia excelsa	85,840	18,950	27,841	36.4	81	4	30
Corythophora amapaensis	85,861	18,778	27,638	36.7	81	4	30
Corythophora labriculata	85,673	18,759	27,594	36.7	81	4	30
Couratari macrosperma	83,785	18,728	27,614	37.0	81	4	30
Couratari stellata	85,547	18,491	27,576	36.9	81	4	30
Eschweilera alata	85,056	18,721	27,635	36.6	81	4	30
Eschweilera caudiculata	84,713	18,759	27,638	37.0	81	4	30
Eschweilera congestiflora	84,815	18,167	27,715	37.1	81	4	30
Eschweilera integrifolia	84,688	18,796	27,592	36.9	81	4	30
Eschweilera micrantha	85,286	18,719	27,668	36.8	81	4	30
Eschweilera wachenheimii	85,378	18,815	27,603	36.8	81	4	30
Lecythis pneumatophora	85,506	18,845	27,622	36.7	81	4	30

444
445

446   TABLE 3. Primers for the amplification of simple sequence repeats in the plastome of *Bertholletia excelsa*. All primers pairs amplify

447   non-coding sequences with the exception of *ndhD*.

Repeat Unit	location	No. Repeats	Region	Primer F (5' - 3')	Primer R (5' - 3')	Product Size (bp)
A	396-409	14	*trnH-psbA*	CCAAAATCATGAACTAACCCCCA	ACCAAGAGGGCGTTATTGCT	226
C	3686-3702	17	*trnK intron*	TGAAGTCGTGTTGCTGAGATCT	CTGTTGATAAGTTTGCCGAGGT	197
A	5680-5691	12	*rps16 intron*	GAGGTTTTCTCCTCGGACGG	ACCACTCATTAAACGAAATGCCT	244
AAAG	9396-9407	3	*trnS-trnG*	GTCCACTCAGCCATCTCTCC	AGCCCGGCCATAGGAATAAA	297
A	9769-9780	12	*trnS-trnG*	TTTATTCCTATGGCCGGGCT	TGCATTGTTTAAGAATCCATAGTTTCA	246
A	17925-17938	14	*rps2-rpoC2*	TTTTCCCCACACTTCCCCTC	TGTCCGGTCATTTGATTTGGT	192
A	29392-29403	12	*rpoB-trnC*	AAGAGAGGAGAAGTTTTTAGGCA	CCTTACCACTCGGCCATGTC	232
AAAG	34775-34786	3	*trnT-psbD*	GGGATGCGAGAAAGAGACTT	CAAAAGTATATCTTTCTACGGGTCG	250
AAAAT	38160-38174	3	*trnS-psbZ*	TACCGGTTTTCAAGACCGGG	TCACAAATGGGCATGCTGGA	201
AAAG	47627-47638	3	*ycf3-trnS*	ACCCATCAATCATTCGATTCGT	GAAAGATCTTTCCTTGGGGGA	168
AAAT	49610-49625	4	*trnT-trnL*	NO SUITABLE PRIMERS FOUND		NA
AATT	50016-50027	3	*trnT-trnL*	NO SUITABLE PRIMERS FOUND		NA
AAAATT	75475-75492	3	*clpP-psbB*	CCACTGAACAAGGGAGAGCC	ACCAAGGCAAACCCATGGAA	128
AAAAT	77155-77169	3	*psbB-psbT*	TGAATCACTGCTTTTCTTTGACTCT	AGGCGGTTCTCGAAAGAAGA	183
A	85073-85085	13	*rpl16-trnH*	TTCAATCTCGGGATTCTTTGAGA	TCGCCTGCGAAAACTTAACT	246
AAAT	102172-102183	3	*rps12-trnV*	TCGATCAATCCCTTTGCCCT	CGTACTCCTCGCTCAATGAGA	248
A	106208-106219	12	*trnL intron*	TGGAGCACCTAACAACGCAT	AGACCTCCGGGAAAAGCATG	119
AACTT	117345-117359	3	*rpl32-trnL*	AGAGTAAACACAAGATACAAGGGT	GTGGGTTAGGTCAATCGGGA	194
AT	117609-117622	7	*rpl32-trnL*	AGTCAACGTCAAAATTAATGAATGGT	AGGTTGAACGCGAGCGATAT	177
AAAT	119729-119740	3	*ndhD*	AAATAACTCCCGCGGTCCAG	GCTTCTCTTGCATTACCGGG	240
AAT	122820-122831	4	*ndhG-ndhI*	NO SUITABLE PRIMERS FOUND		NA

AAT	122843-122854	4	*ndhG-ndhI*	NO SUITABLE PRIMERS FOUND		NA
AATC	125271-125282	3	*ndhA intron*	AACCCGCTTCAAGCCATGAT	AAACGGCTTATAAATTCGCAGT	130

448
449

TABLE 4. Regions of the chloroplast binned in windows of 600 sites with high (above the 95[th]

percentile) nucleotide diversity (π) and/or site-wise log-likelihood score differences (LD). LSC:

large single copy. SSC: small single copy (see main text). Coding regions are indicated in

windows that have the same 5' and 3' expressed flanking regions in column 3. Notice that no

regions are reported for the inverted repeat (IR). Coordinates are given on the alignment and the

Bertholletia excelsa plastome that are assembled with the standard LSC-SSC-IR structure.

Location in the alignment	*Bertholletia* plastome location	Closest flanking expressed region		Region	π	LD
		5'	3'			
1–600	1–490	*trnH*	*psbA*	LSC		*
5401–6000	4885–5373	*trnK-UUU*	*rps16*	LSC		*
34801–35400	30925–31450	*petN*	*trnD-GUC*	LSC		*
35401–36000	31451–31967	*psbM*	*trnD-GUC*	LSC	*	*
37201–37800	33027–33573	*trnE-UUC*	*trnT-GGU*	LSC	*	*
39601–40200	34893–35433	*trnT-GGU*	*psbD*	LSC		*
43801–44400	38798–39254	*psbZ*	*trnfM-CAU*	LSC	*	*
44401–45000	39255–39744	*trnfM-CAU*	*psaB*	LSC	*	*
61201–61800	54771–55275	*trnV-UAC*	*atpE*	LSC		*
78601–79200	70230–70771	*psaJ*	*rps18*	LSC		*
89401–90000	80536–81103	*petD*	*rpoA*	LSC	*	
95401–96000	85455–85906	*rpl16*	*rps3*	LSC	*	
131401–132000	119237–119759	*ccsA*	*ndhD*	SSC	*	
140401–141000	127827–128402	*rps15*	*ycf1*	SSC		*
144001–144600	131283–131868	*ycf1*	*ycf1*	SSC	*	*
144601–145200	131869–132446	*ycf1*	*ycf1*	SSC	*	*

456
457

458   TABLE 5. Nucleotide diversity (π) and differences in log-likelihood (LD) scores of the informative

459   windows identified in this study (bold) and previously proposed barcode markers. High values,

460   above the 95th percentile, are indicated in bold.

Region	Pi	LD
ccsA–ndhD	**0.0258**	247.12
matK	0.0153	136.92
petD–rpoA	**0.0246**	260.79
petN–trnD	0.0228	**361.07**
psaJ–rps18	0.0176	**309.10**
psbM–trnD	**0.0292**	**330.41**
psbZ–trnfM	**0.0246**	**373.97**
rbcL	0.0105	95.03
rpl16–rps3	**0.0345**	275.89
rpoB	0.0097	120.53
rpoC1	0.0103	178.60
rps15–ycf1	0.0212	**284.57**
trnE–trnT	**0.0241**	**522.51**
trnfM–psaB	**0.0254**	**375.76**
trnH–psbA	0.0126	**310.47**
trnK–rps16	0.0164	**350.44**
trnL	0.0106	192.27
trnT–psbD	0.0239	**291.15**
trnV–atpE	0.0128	**379.77**
ycf1 **(1)**	**0.0273**	**462.53**
ycf1 **(2)**	**0.0469**	**313.12**

462   TABLE 6. Primer sequences designed to amplify the ten most polymorphic Lecythidaceae

463   plastome regions, as sorted by decreasing nucleotide diversity (π). The product size (length)

464   references the *Bertholletia excelsa* plastome.

Window in the alignment	π	Region	Forward primer Sequence (5' - 3')	Reverse primer Sequence (5' - 3')	Length (bp)
144103-145487	0.04691	*ycf1*(1)	AGAACCTTTGATTATGTCTCGACG	AGAGACATGCTATAAAAATAGCCCA	1186
95034-95741	0.03446	*rpl16-rps3*	AGAGTTTCTTCTCATCCAGCTCC	GCTTAGTGTGTGACTCGTTGG	1014
35585-36413	0.02920	*psbM-trnD*	CCGTTCTTTCTTTTCTATAACCTACCC	ACGCTGGTTCAAATCCAGCT	1093
143235-144102	0.02733	*ycf1*(2)	TGATTCGAATCTTTTAGCATTAKAACT	KCGTCGAGACATAATCAAAGGT	1189
131180-132054	0.02576	*ccsA-ndhD*	CCGAGTGGTTAATAATGCACGT	GCTTCTCTTGCATTACCGGG	1180
44398-45132	0.02537	*trnG-psaB*	TCGATYCCCGCTATCCGCC	GCCAATTTGATTCGATGGAGAGA	883
89032-89688	0.02464	*petD-rpoA*	TGGGAGTGTGTGACTTGAACT	TGACCCATCCCTTTAGCCAA	824
43412-44397	0.02456	*psbZ-trnfM*	TCCAATTGRCTGTTTTTGCATTAATTG	CCTTGAGGTCACGGGTTCAA	706
37444-38345	0.02409	*trnE-trnT*	AGACGATGGGGGCATACTTG	CCACTTACTTTTTCTTTTGTTTGTTGA	1324
38346-40085	0.02391	*trnT-psbD*	GGCGTAAGTCATCGGTTCAA	CCCAAAGCGAAATAGGCACA	1717

465
466

467 **FIGURES**

469   Fig. 1. Plastome map of the Brazil-nut tree *Bertholletia excelsa*. Genes outside the circle are

470   transcribed clockwise, genes inside the circle are transcribed counter-clockwise. Gray bars in the

471   inner ring show the GC content percentage.

Fig. 2. A) Sliding window plot of nucleotide diversity (π) across the alignment of 24 sequenced

Lecythidaceae plastomes. B) Alignment site-wise differences in log-likelihood (LD) calculated

from the chloroplast topology vs. the average scores of 1000 random trees using a 600-site

window. Regions with π and LD above the 95th percentiles are indicated with dashed lines.

Continuous vertical lines indicate the boundaries, from left to right, among the large single copy,

the inverted repeat, and the small single copy.

480   Fig. 3. Maximum likelihood phylogeny inferred from plastomes of 23 Neotropical Lecythidaceae.

481   Numbers at nodes indicate bootstrap support.

Fig. 4. A) Average bootstrap support for trees inferred from matrices of concatenated regions with relatively high nucleotide diversity sorted in ascending order; and B) Robinson-Foulds distance (RF) sorted in descending order. Lower RF distances, which measures the number of different bipartitions from the complete plastome topology, indicate better accuracy.

SUPPORTING INFORMATION

Fig. S1. Phylograms obtained from single and combined markers with high nucleotide diversity.

Numbers at nodes indicate bootstrap support.

APPENDIX 1. Lecythidaceae species sequenced with their voucher, assembly information, and

GenBank accession number. All voucher specimens are deposited at the New York Botanical

Garden Herbarium (NY).