TITLE

Incorporating trait-based mechanisms, alternative interaction modes and forbidden links

increases the minimum dimensionality of ecological networks

RUNNING TITLE

Networks' minimum mechanistic dimensionality

AUTHORS' DETAILS

Diogenis A. Kiziridis (correspondence); Departments of Mathematics and Biosciences,

Swansea University, Swansea, UK; danis.k@zoho.com

Lynne Boddy; Cardiff School of Biosciences, Cardiff University, Cardiff, UK;

boddyl@cardiff.ac.uk

Daniel C. Eastwood; Department of Biosciences, Swansea University, Swansea, UK;

d.c.eastwood@swansea.ac.uk

Chenggui Yuan; Department of Mathematics, Swansea University, Swansea, UK;

c.yuan@swansea.ac.uk

Mike S. Fowler; Department of Biosciences, Swansea University, Swansea, UK;

m.s.fowler@swansea.ac.uk

ABSTRACT

1.   Multiple traits are involved in intra- and interspecific interactions, influencing

interaction outcomes and community structure. Knowing the minimum number of traits

required to explain observed network structures, i.e. the 'minimum dimensionality', can help

us to focus on key biological questions. Existing theoretical methods for estimating minimum

dimensionality are often simplified in three ways: (*i*) they are phenomenological, (*ii*) they lack

the ubiquitous feature of alternative interaction (e.g. feeding) modes, (*iii*) they do not consider

phenological and/or phenotypic mismatches, known as 'forbidden links'. Here we develop a

method to estimate the 'minimum mechanistic dimensionality' of ecological networks, which

can assume alternative interaction modes and account for forbidden links.

2.   Our method compares the performance of traits, e.g. predator mouth gape versus prey

body size. A system of linear inequalities—making assumptions of how traits are involved in

alternative interaction modes—is constructed to explain the observed interaction outcomes.

We attempt to solve these inequalities with mixed integer linear programming. Minimum

mechanistic dimensionality is the minimum number of traits required for a feasible system of

inequalities. We illustrate how the inclusion of alternative interaction modes alters the

explanation of an intransitive 'rock–paper–scissors' system of competing marine

invertebrates. We apply our method to 658 published empirical ecological networks—

including animal dominance, predator–prey, primary consumption, pollination, parasitism and

seed dispersal networks—to determine whether minimum dimensionality estimates change

following the inclusion of concepts emerging from our framework, i.e., alternative interaction

modes, trait-mediated 'forbidden links' and a mechanistic approach to describing trait-based

interactions.

3.   The minimum mechanistic dimensionality was typically higher in ecological

networks when including alternative interaction modes (54% of empirical networks),

'forbidden interactions' as trait-mediated interaction outcomes (92%), or a mechanistic

perspective (81%), compared to dimensionality estimates in the absence of these features.

4.   Our method can inform us about how many traits are mechanistically involved in

pairwise interactions, to generate theoretical networks of more accurate structure at the

interaction outcome level. In general, it can reduce the risk of omitting essential traits, for

improved understanding, explanation, and prediction of community structures, and structure-

dependent community, ecosystem, and evolutionary processes.

KEYWORDS

Species interactions, trait-based mechanisms, niche space, phenotype space, food web

intervality, multilayer ecological networks, mutualism, trophic interactions, social networks,

cyclic rock–paper–scissors intransitive game.

1   INTRODUCTION

Understanding community structure can improve our predictions of structure-dependent

population dynamics (Berlow et al., 2009), evolution (Becerra, 2003), ecosystem processes

(Woodward, Papantoniou, Edwards, & Lauridsen, 2008), and conservation (Mello et al.,

2015). Communities are structured by different forces, including evolution; spatio-temporal

distribution of species due to neutral, historical, dispersal, or habitat filtering processes;

interaction neutrality; and traits (Bartomeus et al., 2016; Poisot, Stouffer, & Gravel, 2015;

Vázquez, Chacoff, & Cagnolo, 2009). A useful representation of the structure of interactions

is through network approaches (Delmas et al., 2019; Newman, 2003). Focusing on the

contribution of traits in interaction networks, i.e. of how individuals interact via a subset of

their phenotypic characters, allows us to ask what the minimum number of traits is that must

be involved given the interaction outcomes we observe, hereafter called 'minimum

dimensionality'. The minimum dimensionality of a system can concentrate our efforts on

investigating which traits underpin community structure (Eklöf et al., 2013).

A mechanistic method for estimating the minimum dimensionality of an interaction network

must make assumptions about how interactions are expected to occur. Interactions in different

systems appear to share four common features: (1) interactions can be of various types within

a system, from cooperative and mutualistic, to antagonistic or victim–exploiter interactions,

within or between species; (2) alternative modes can lead to a successful outcome in a given

type of interaction, such as the different signals which plants use to attract pollinators

(Schiestl & Johnson, 2013), novel pathways to bacterial infection (Meyer et al., 2012),

alternative combative mechanisms employed in fungal competition for space (Boddy, 2000),

and different feeding modes (Kiørboe, 2011); (3) multiple tasks may need to be completed for

success via a given interaction mode, e.g., the encounter–detection–identification–approach–

subjugation–consumption steps of a typical predation sequence (Endler, 1991); and (4) for a

specific task, the traits involved are measurable, competing features of the interacting players

(McGill, Enquist, Weiher, & Westoby, 2006): physiological (e.g. plant–herbivore toxin versus

detoxification enzyme concentrations), morphological (e.g. animal–plant proboscis length

versus depth of nectar in floral tube), behavioural (e.g. cheating versus punishment in

cleaner–client fish), and phenological (e.g. predator–prey temporal presence versus absence).

Here, we attempt to combine all four of these features in a single framework for the first time.

Despite the accumulated empirical knowledge about how interactions occur, the two available

methods to estimate minimum dimensionality lack a mechanistic perspective, or an explicit

incorporation of the widespread feature of alternative interaction modes. Eklöf et al. (2013)

developed a method for estimating minimum dimensionality in different interaction types,

rooted in the theory of food web intervality in niche space (Cohen, 1977). In this method,

dimensions originate phenomenologically, each dimension potentially accounting for multiple

traits. Dalla Riva and Stouffer's (2016) minimum dimensionality method avoids this approach

by adopting a simple trait space representation for trophic interactions. Network structure is

explained mechanistically here by explicitly modelling exploiter–resource interactions via

trait matching. However, Dalla Riva and Stouffer (2016) model interactions via a single

interaction mode, since the contributions act additively from each exploiter–resource trait

pair. The same focus on a single interaction mode appears in Eklöf et al. (2013), because

niche dimensions act in conjunction to determine the niche of the exploiter (resource).

Here, we aimed to develop a new method for estimating the minimum number of traits

required to explain the interaction outcomes across a broad range of ecological networks. This

'minimum mechanistic dimensionality' method encompasses all four features of interactions

outlined by empirical studies, within a simple phenotypic trait space representation. We ask if

our minimum dimensionality estimate is higher than other approaches across a range of

different empirical ecological network types: under the assumption of alternative interaction

modes compared to a single mode; with failure outcomes taken into account instead of

omitted; and under its mechanistic perspective compared to the minimum dimensionality

under the phenomenological, niche approach of Eklöf et al. (2013). We show that essential

traits can be missed from ecological networks because the lower bound of the number of traits

involved in 658 empirical systems we interrogated was frequently underestimated by omitting

the presence of alternative interaction modes, the trait-mediated failure outcomes, and the

mechanistic involvement of traits competing in pairs.

2   METHODS

Here, we outline our general approach using empirical examples based on a nectarivory

network and an intransitive system of competing marine invertebrates. We describe the

interactions among species in the context of exploiter and resource roles, before going on to

define and calculate the minimum mechanistic dimensionality of these networks using a linear

inequalities method. We then describe how we compared competing minimum dimensionality

estimates across 658 empirical examples, ranging from social hierarchies, to mutualistic

networks, to food webs.

2.1 A generalised mechanistic description of how interactions occur

We illustrate our mechanistic description with an empirical plant–animal system of two

interaction types, reported in Kennedy (1978): nectarivory pursued by four animal species,

and biotic pollination pursued by a plant species (Fig. 1). For nectarivory, the animals are the

exploiters, and the plant the resource; for pollination, the plant is now considered the

exploiter, and the animals the resources. This system can be represented by two bipartite

exploiter–resource graphs, regarded as two layers of a multilayer network (Kivelä et al.,

2014). There are four possible pairwise interaction outcomes: the plant is pollinated and an

animal exploits the plant's nectar; neither the plant nor an animal succeed in exploiting the

other; the plant is not pollinated, but its nectar is exploited; or the plant is pollinated, and its

nectar is not exploited. Note that nectarivory and pollination failures are also considered

interaction outcomes in our description.

FIGURE 1 The mechanistic description of interactions applied to an empirical system (lowland wet forest near

Puerto Viejo, Heredia, Costa Rica, described in Kennedy, 1978). From left to right, the animals reported are the

three Euglossine bees *Eulaema speciosa*, *E. seabrai*, and *E. meriana*, and the hummingbird is *Phaethornis*

longirostris. The plant species is the closed-flowered *Calathea marantifolia* (a closed, white flower is shown in the

species token). The interaction modes via which animals succeed are given in parentheses next to the outcomes

indicated by arrows. We show only some indicative tasks in the modes, for the sake of illustration. Species token

size indicates the relative size of the representative individual.

Successful nectarivory or pollination outcomes could occur via alternative modes. For

instance, the animals adopt two alternative modes of achieving nectarivory: accessing nectar

either by opening petals or puncturing sepals (Fig. 1). Success in a given mode may require

success in multiple tasks. For example, the petal-opening nectarivory mode requires success

in at least five tasks (Fig. 1). For a success in a task, an exploiter's performance in a trait,

termed 'power', must be sufficiently higher than the resource's performance in a

corresponding trait called 'toughness' (taken from the creature combat rules of the card game

Magic: The Gathering® in Garfield, 2017). We consider that exploiter and resource are

challenged in one trait 'dimension' of their phenotype space, where the performance of the

corresponding power–toughness traits compete to determine who succeeds in that task. Using

Boolean logic terms, modes can be represented as OR-associated clauses of AND-associated

tasks. Any structure of logical statements can be equivalently expressed in this 'disjunctive

normal form' (Cohn, 2003), herein termed 'interaction form'.

2.2 The minimum mechanistic dimensionality of ecological networks

In the nectarivory example above, the animals succeeded via two alternative modes, one five-

dimensional (one spatial and four morphological traits for the bees), the other two-

dimensional (one spatial and one morphological trait for the hummingbird) (Fig. 1).

Minimally, we could observe these four animal successes in nectarivory via a one-

dimensional mode: all animals could 'easily' consume nectar from a plant with open, wide

and short flowers just because of spatial matching with the flowered plant. In other words, if

this system was mechanistically minimal, the same outcomes would have occurred only

because of interactions in one exploiter–resource dimension. Thus, the idea behind our

method is to find a minimal interaction form which is sufficient for the mechanistic

explanation of all the observed interaction outcomes of a system. By comparing a

theoretically minimal interaction form with the empirically observed one, we can gain insight

into the extra strategies, measures, or defences employed by the players. In the example

illustrated in Fig. 1, the plant imposed six exploitation barriers (*sensu* Santamaría &

Rodríguez-Gironés, 2015), challenging the animals in six dimensions instead of the

theoretically required one dimension.

2.2.1 Rock–paper–scissors: an intransitive network with marine invertebrates

The success of the animals at the empirical plant–animal system in Kennedy (1978) could

minimally arise by interactions in one pair of exploiter–resource trait dimensions. However,

one dimension is not sufficient to mechanistically explain the outcomes of an intransitive

network of outcomes between three (or more) species, such as the cyclic spatial replacement

of marine invertebrates studied by Jackson and Buss (1975): an ectoproct species *Stylopoma*

spongites (player *A*) replaces a sponge species *Tenaciella* sp. (player *B*), which in turn

replaces sponge species *Toxemna* sp. (player *C*), and which in turn replaces the ectoproct

species player A. We consider all three species both exploiters and resources in spatial

replacement (Fig. 2).

FIGURE 2 The mechanistic explanation of outcomes of an intransitive three-player interaction in a rock–paper–

scissors empirical system requires at least two exploiter–resource trait pairs, i.e. two dimensions. The empirical

system is reported in Jackson and Buss (1975), with three encrusting invertebrate species: the ectoproct *Stylopoma*

spongites (player *A*), and the sponges *Tenaciella* sp. and *Toxemna* sp. (players *B* and *C*, respectively). We show

three attempts to minimally explain the observed outcomes, with the first attempt being mechanistically

impossible. We indicate hypothetical tasks, and power | toughness trait scores in arbitrary units of performance.

It is impossible to explain the observed outcomes in this unipartite graph with interactions via

a one-dimensional interaction mode. The single pair of exploiter–resource power–toughness

traits is involved in a system of six inequalities, to satisfy the observed successes and failures.

For the successful replacements, the power of a winning exploiter must be greater than the

toughness of a defeated resource. For the failures, the power of a losing exploiter must be less

than or equal to the toughness of an undefeated resource. This system of six linear inequalities

creates a cyclic sequence of ever-increasing power–toughness scores (the impossible '1 one-

dimensional mode' in Fig. 2).

Our framework provides two alternative minimal mechanistic explanations for the emergence

of a rock–paper–scissors system. First, we can find solutions if we add a second pair of

power–toughness traits in the same mode (Fig. 2). We explain the failure of players A and B

as failure in the first dimension (e.g. overgrowth), and the failure of C as failure in the second

dimension (e.g. destruction of rival, even if C can overgrow B). Second, we can find solutions

if we add a second pair of power–toughness traits in a new one-dimensional mode (Fig. 2). In

that case, we explain the success of A and B as success via the first mode (e.g. destructive

overgrowth), and the success of C as success via a second mode (e.g. allelochemical

elimination). Minimal explanation II is described by Jackson and Buss (1975) for this cryptic

reef system: player A replaces B via overgrowth, player B replaces C via overgrowth, but

player A is replaced by C due to toxic effects. Appendix S1 in the Supporting Information

includes the complete systems of linear inequalities for rock–paper–scissors under minimal

explanations I and II.

2.2.2 The linear inequalities method for calculating the minimum mechanistic dimensionality

As illustrated with the rock–paper–scissors example (Fig. 2), systems might require more than

one pair of exploiter–resource competing traits for their outcomes to be explained

mechanistically. One method to find this minimum number of dimensions is by attempting to

solve a system of linear inequalities. If the system of linear inequalities is impossible, a

simple strategy is to increase the number of dimensions (d) by one, and retry. The minimum d

≥ 1 for a feasible system of inequalities is our minimum mechanistic dimensionality. In the

marine invertebrates example of Fig. 2, there were two minimal explanations: an additional

trait pair belonging to the same mode (minimal explanation I); or belonging to a new one-

dimensional mode (minimal explanation II). We will focus on these two extreme

explanations, although there could be intermediate minimal interaction forms for more than

two dimensions.

When a new task is added to a single mode, that permits feasibility of the system of linear

equalities, the d exploiter–resource trait pairs (dimensions) must be involved in the same

mode (minimal explanation I, Fig. 2). On one hand, an observed success of exploiter A

against resource B must be the result of success in all tasks (e.g. a successful parasite has

overcome all the host defences). Specifically, the power of exploiter A in any trait pair i, $P_{A,i} \geq$

0, must be greater than the toughness of resource B in that trait pair, $T_{B,i} \geq 0$: $P_{A,i} > T_{B,i}$. Since

success might require more than the marginal superiority of the exploiter's power, we can add

a superiority threshold, $t_{A,B,i} > 0$, making the task success requirement $P_{A,i} \geq T_{B,i} + t_{A,B,i}$. On the

other hand, an observed failure of exploiter A against resource B must be the result of failure

in at least one task (e.g. a parasite fails against at least one host defence). We can use a binary

variable as an indicator of failure in trait pair i, $f_{A,B,i}$ (Williams, 2013). If $f_{A,B,i} = 1$, then

exploiter A fails against resource B in trait pair i; otherwise, $f_{A,B,i} = 0$, representing exploiter

success in the task. The demand for a failure in at least one trait pair i can be formulated with

the linear inequality $\sum_{i=1}^{d} f_{A,B,i} \geq 1$. Finally, we include bounds for the power–toughness

differences for computational efficiency (Williams, 2013): a sufficiently negative lower

bound of the exploiter's power inferiority in case of task failure, m; and a sufficiently positive

upper bound of the exploiter's power superiority in case of task success, M. Thus, for an

observed failure, the following pair of inequalities must be satisfied in any trait pair i:

$$P_{A,i} + M f_{A,B,i} \leq T_{B,i} + M, \tag{1}$$

$$P_{A,i} - m f_{A,B,i} \geq T_{B,i} + t_{A,B,i}. \tag{2}$$

The extra inequality $\sum_{i=1}^{d} f_{A,B,i} \geq 1$ forces at least one of the binary indicator variables to equal

one, i.e. failure in at least one task. In the case of a task failure in trait pair i, $f_{A,B,i} = 1$,

inequality (1) is the task failure requirement, and inequality (2) is the lower bound for the

exploiter's power inferiority. In case of a task success, $f_{A,B,i} = 0$, inequality (1) gives the upper

bound for the exploiter's power superiority, and inequality (2) becomes a success

requirement.

When a new one-dimensional mode is added, permitting a feasible system (minimal

explanation II, Fig. 2), each of the d pairs of competing exploiter–resource traits must be

involved in a different mode. On one hand, the observed failure of any exploiter A against any

resource B must be the result of failure in all d modes, $P_{A,i} \leq T_{B,i}$ in any mode i (e.g. a parasite

cannot invade via any of the alternative host entrances). On the other hand, the observed

success of exploiter A against resource B must come from success via at least one mode (e.g.

a parasite invades via at least one host entrance). We now use a binary variable, $s_{A,B,i}$, to

indicate success via mode i. Given the same bounds as in minimal explanation I, the

following pair of inequalities must be satisfied to indicate exploiter success in any mode i:

$$P_{A,i} + m\, s_{A,B,i} \geq T_{B,i} + t_{A,B,i} + m, \qquad (3)$$

$$P_{A,i} - M\, s_{A,B,i} \leq T_{B,i}. \qquad (4)$$

With the extra inequality $\sum_{i=1}^{d} s_{A,B,i} \geq 1$, we force at least one of the indicator variables to

equal one, i.e. exploiter success occurs via at least one mode.

A complete system of linear inequalities incorporates all observed successes and failures of

all possible exploiter–resource pairs (see Appendix S1 in the Supporting Information for

examples of complete systems of linear inequalities under minimal explanations I and II).

Such systems of linear inequalities, with continuous trait values and integer indicator

variables, can be formulated and attempted to be solved as mixed integer linear programming

problems (Williams, 2013). In both minimal explanations (I and II), minimum mechanistic

dimensionality is the minimum d leading to a feasible system of inequalities.

2.3 Comparing the minimum dimensionality of empirical networks

We applied our method to 658 empirical systems, covering six different types of ecological

networks: animal social dominance networks, food webs with basal species excluded, basal–

consumer, plant–pollinator, host–parasite, and seed dispersal networks (see Appendix S1 for

details). Considering a single interaction type in each system, and assuming adequate

sampling effort (e.g. no observed failures due to rarity), we computed five dimensionality

measures (Appendix S1). Four minimum mechanistic dimensionality measures were based on

our framework: (*a*) a single, potentially multidimensional mode; (*b*) potentially multiple, one-

dimensional modes; (*c*) as measure (*b*), but excluding observed failures; (*d*) as measure (*c*)

but with players interacting via a common trait per dimension, rather than a power against

toughness trait. The fifth dimensionality measure we considered (*e*) was the

phenomenological measure of Eklöf et al. (2013). We used these five dimensionality

measures to answer three questions about minimum mechanistic dimensionality (MMD): (1)

Is MMD higher under the assumption of alternative one-dimensional modes (dimensionality

measure *b*), compared to the assumption of a single multidimensional mode (dimensionality

measure *a*)? (2) Is MMD higher with observed failures taken into account (dimensionality *b*),

or excluded (dimensionality *c* or *d*)? (3) Is MMD higher than the phenomenological

dimensionality measure developed by Eklöf et al. (2013) (dimensionality *a* versus *e*)?

The systems of linear inequalities for our four minimum mechanistic dimensionality measures

were formulated and solved with the Gurobi Optimizer (Gurobi Optimization and Inc., 2017)

as mixed integer linear programming problems (Williams, 2013). R and Python code for

formulating and solving for these four minimum mechanistic dimensionalities are provided

(DATA ACCESSIBILITY). We computed the fifth dimensionality with code provided in the

Supporting Information of Eklöf et al. (2013). The references we provide for the empirical

networks (DATA ACCESSIBILITY) were retrieved from five data sources (Cohen, 2010;

Ortega, Fortuna, & Bascompte, 2017; Shizuka & McDonald, 2015; Stanko & Miklisova,

2014; Thompson & Townsend, 2004). Additionally, we provide characteristics, and raw data

from the five computed dimensionality measures, for each of the 658 empirical systems

(DATA ACCESSIBILITY).

3   RESULTS

For the five dimensionality measures we considered, the inclusion of alternative interaction

modes, forbidden links and a mechanistic approach to describing interactions consistently

increased the minimum dimensionality estimate across a wide range of empirical ecological

networks.

3.1 Multimodal versus unimodal minimum mechanistic dimensionality

Our minimum mechanistic dimensionality was frequently higher under the alternative modes

explanation than under the single mode explanation (Fig. 3). The dimensionality assuming

alternative modes (horizontal axes in Fig. 3) increased faster than the dimensionality

assuming a single interaction mode (vertical axes in Fig. 3), especially in the systems of non-

basal consumption, biotic pollination, ectoparasitism, and seed dispersal (Fig. 3b,d–f). 54% of

the empirical systems had higher dimensionality if alternative modes were assumed, with only

7% of the systems having higher unimodal dimensionality (Fig. 4a).

FIGURE 3 Minimum mechanistic dimensionality estimates from 658 empirical systems. Cell shading shows how many of the n systems had the specific pair of values in our two minimum dimensionality measures, i.e. number of exploiter–resource trait pairs in: alternative one-dimensional modes (x-axis; minimal explanation II); a single mode (y-axis; minimal explanation I). We focused on one interaction type in each empirical system: (a) animal dominance in unipartite graphs (6–31 individuals); (b) consumption of non-basal species in unipartite food web graphs (6–57 species, basal species excluded from the original food webs); (c) consumption of basal species by consumers exclusively feeding on them in bipartite graphs (11–91 species; from the same food webs used in (b)); (d) plant biotic pollination in bipartite graphs (8–114 species); (e) ectoparasitism of small mammals in bipartite graphs (8–92 species); and (f) plant seed dispersal in bipartite graphs (6–86 species). Parameter values in the linear inequalities method: $m = -200$; $M = 200$; $t_{A,B,i} = 1$, for all pairs of exploiter A with resource B, in any trait pair i.

FIGURE 4 Comparisons of three minimum dimensionality measures estimated from 658 empirical systems. For

each empirical network, we calculated the ratio of: (a) our minimum mechanistic dimensionality under minimal

explanation II (alternative one-dimensional modes), to our minimum mechanistic dimensionality under minimal

explanation I (a single mode); (b) our minimum mechanistic dimensionality under minimal explanation II, to the

same dimensionality with the failures excluded from the system of linear inequalities; and (c) our minimum

mechanistic dimensionality under minimal explanation I, to the comparable phenomenological dimensionality of

Eklöf et al. (2013). Violin plots show the normalised distributions of the dimensionality ratios. The dotted

horizontal lines give the ratio of one.

3.2 Minimum mechanistic dimensionality with failure outcomes taken into account versus

excluded

Minimum mechanistic dimensionality was higher in 92% of the empirical systems when

failure outcomes were included, instead of excluded (Fig. 4b). In the rest of the empirical

systems, the two dimensionality estimates were equal. Here, we compared our minimum

multimodal dimensionality with the same dimensionality but with any failure inequalities

excluded from the inequalities system. The minimum dimensionality with failures excluded

always equals one because all exploiters can have a single power trait with value greater than

the value of the single toughness trait of any resource, in the absence of any inequalities

constraining the power scores, hence explaining any structure of only observed successes

unimodally. We further required that exploiters and resources possess the same trait for power

and toughness in the unipartite systems of animal dominance and non-basal consumption,

instead of the default power–toughness trait pair. In that way, the unipartite systems could

require more than one dimension with failures excluded. Even in this case of modelling trait

competition with a single, common trait per dimension, 79% of the unipartite systems had

higher minimum dimensionality with failures included rather than excluded.

3.3 Mechanistic versus phenomenological minimum dimensionality

In 81% of the empirical systems, our minimum mechanistic dimensionality was higher than

the phenomenological dimensionality estimate of Eklöf et al. (2013). We assumed a single

mode (minimal explanation I), which is comparable to the niche approach of Eklöf et al.

(2013). Only 2% of the networks had higher minimum dimensionality under the

phenomenological approach, with no bipartite networks among them (Fig. 4c).The minimum

number of trait pairs for the explanation of all outcomes in our approach, was 1.5 times larger

(median) than with the phenomenological dimensionality across all networks. Note that since

our dimensionality refers to exploiter–resource trait pairs, the actual number of necessary

traits is double our dimensionality, i.e. the present approach suggested a median of 3 times

more traits required for the explanation of the observed outcomes than the Eklöf et al. (2013)

approach.

4   DISCUSSION

We introduced a method for calculating the minimum number of traits required to explain all

observed interaction outcomes of an ecological network mechanistically, using a general

framework which is applicable to different interaction types, modes, tasks, and types of traits.

Applying our method to 658 empirical systems, we showed that the minimum number of

traits involved is typically underestimated when ignoring any of the three framework features

we combined here for the first time: (1) alternative interaction modes; (2) trait-mediated

failure outcomes; and (3) a mechanistic description of interactions.

With our generalised framework, minimum mechanistic dimensionality can explicitly

incorporate the alternative interaction modes observed empirically, e.g. alternative feeding

modes. In previous theoretical trait-based works, an exploiter has to overcome all the barriers

or defences of a potential resource to consume or parasitise the resource (Gilman, Nuismer, &

Jhwueng, 2012; Santamaría & Rodríguez-Gironés, 2007); similarly, in other theoretical works

adopting a niche approach, a niche arises from the intersection of all niche dimension

intervals (Eklöf et al., 2013; Stouffer, Camacho, & Amaral, 2006). The interaction mode in

our framework is equivalent to these two approaches, since an exploiter's performance must

be sufficiently high in all the mode's tasks. By additionally generalising to alternative

interaction modes, we showed that minimum mechanistic dimensionality can differ,

frequently being larger under alternative interaction modes than in a single mode (Fig. 4a).

Additionally, different assumptions about the interaction form can lead to alternative minimal

explanations of the outcomes, as in rock–paper–scissors systems (Fig. 2), offering a new

mechanistic perspective in the study of intransitive networks (Szolnoki et al., 2014).

We adopted a phenotype rather than a niche space representation of traits. Studies of

interactions commonly use the 'resource-utilization' approach to representing the 'ecological

niche' concept (MacArthur & Levins, 1967, Schoener, 1989). Despite its operational

advantage (Schoener, 1989), dimensions usually arise phenomenologically, as in the

minimum dimensionality of Eklöf et al. (2013). For example, body size is a trait with high

explanatory power in food webs (Stouffer, Rezende, & Amaral, 2011). However, more traits

allometrically scaling with body size are mechanistically involved in trophic interactions

(Woodward et al., 2005). Even if taken mechanistically, realised niches commonly span a

range of the resource gradient (MacArthur & Levins, 1967), implying two traits per niche

dimension. For instance, with the maximum prey size range limited by a predator's mouth

gape, the niche range minimum must be delimited due to a second trait, like the predator's

inability to handle smaller prey. Another problem is that exploiters are excluded from the

niche space because it is created by trait dimensions of the resources (MacArthur & Levins,

1967; Schoener, 1989). Our framework accounts for the traits of both interacting players

simultaneously, and a dimension is simply a challenged trait-axis in the phenotype space of

exploiters and resources, as in the approach of Dalla Riva and Stouffer (2016). In that way,

we found that our minimum mechanistic dimensionality assuming a single interaction mode

was frequently higher than the comparable phenomenological dimensionality of Eklöf et al's

(2013) niche-based approach (Fig. 4c).

We regarded failures as trait-mediated outcomes of an interaction, meaning more traits are

expected to be involved in the interactions (Fig. 4b). The 'success' outcomes in our

framework are the 'interactions' typically described in the literature (Bartomeus et al., 2016);

and our 'failure' outcomes are synonymous to the 'forbidden links' or 'forbidden interactions'

of some authors (Jordano, Bascompte, & Olesen, 2003; Morales-Castilla, Matias, Gravel, &

Araújo, 2015). Under our framework, players interact given their mere inclusion in the study

system, even if they never actually meet, e.g. through spatio-temporal mismatch. Behavioural

studies commonly employ a few predictor traits to explain only the observed dominance

events in a system (Chase & Seitz, 2011), i.e. only the successes. In contrast, we found that

three to six pairs of competing traits must be involved in several dominance systems (Fig. 3a).

For example, in the elephant family named 'AA' in Archie et al. (2006), almost all observed

dominance events are directed towards younger elephants, and the authors conclude the

system is an age-ordered dominance hierarchy based only on the successes, in agreement with

our one dimension with failures-excluded analysis (Fig. 4b). Our minimum mechanistic

dimensionality though, explicitly incorporating failures, suggests three trait pairs here under

both minimal explanations, because there are several older–younger pairs where no

dominance or aggression was observed, i.e. failures unaccounted for by Archie et al. (2006).

In other words, our framework predicts mechanisms preventing the occurrence of dominance

between these older–younger pairs. Most elephants dominated only younger members of their

matriline, and of two specific matrilines (Archie et al., 2006). These two preferences are

candidates for the two extra dimensions that we expect under our framework, which are

overlooked when ignoring failure outcomes.

Our framework is applicable to various types of traits. Continuous-valued quantitative traits

can be modelled directly (e.g. for animal reaching nectar in Fig. 1). Comparison of traits with

ordered levels (binary, semiquantitative, and quantitative but discontinuous; Legendre &

Legendre, 1998) can be modelled with appropriate scaling (e.g. for animals separating petals

in Fig. 1). Categorical qualitative traits can be redefined as binary traits (e.g. a qualitative trait

for inhabiting a forest can be redefined to a binary presence–absence trait as in Fig. 1).

Moreover, the inequality rule can model both cases of competing traits (Abrams, 2000;

Santamaría & Rodríguez-Gironés, 2007): difference traits (also called barrier traits, or

unidirectional axes of vulnerability), and matching traits (complementarity traits, or

bidirectional axes of vulnerability). The natural case in our framework is the difference traits,

since a larger power–toughness difference contributes to success. However, if we state the

tasks appropriately, matching traits can be reformulated as difference traits (e.g. the difference

traits formulation for the animal–plant spatial matching in Fig. 1).

The proposed method for calculating the minimum dimensionality is mechanistic, based on

biologically plausible features of interactions; yet, it is simple with its phenotype space

representation, and easy to adjust and extend. In this first account, we assumed two simple

and extreme minimal interaction forms (minimal explanations I and II), but the user can input

any minimum number of traits and trait values, in any interaction form. Additionally, we

presented a deterministic version, but future versions could be probabilistic (Dalla Riva &

Stouffer, 2016; Poisot et al., 2016), e.g. with more probable outcomes explained by larger

power–toughness differences. Assuming adequate sampling effort, our mechanistic

description has not considered the effects of phylogenetic relationship (Rohr & Bascompte,

2014), and abundance (Vázquez & Aizen, 2003), which could also be incorporated in future

extensions.

In conclusion, we have outlined a novel method of estimating the minimum dimensionality of

ecological networks, which is relatively straightforward to adopt and calculate. Studies can

rely on network models reproducing community structure more accurately at the interaction

outcome level (Olito & Fox, 2015), and can reduce the risk of missing important traits that are

involved mechanistically, in alternative interaction modes, or only in failure outcomes. In that

way, our method could contribute to a better understanding, explanation, and prediction of

community structure and structure-dependent processes.

ACKNOWLEDGEMENTS

The authors would like to thank David Gilljam for comments which improved the manuscript.

Swansea University supported DAK with a student scholarship.

AUTHORS' CONTRIBUTIONS

DAK developed the method and the required code, compiled the empirical networks dataset,

computed the dimensionalities of the networks, analysed the data, and wrote the first draft of

the manuscript. All authors contributed to the development of the method, data analysis, and

to the writing of the manuscript.

DATA ACCESSIBILITY

Data and code will be made available at the Dryad Digital Repository.

REFERENCES

Abrams, P. A. (2000). The evolution of predator–prey interactions: theory and evidence.

Annual Review of Ecology and Systematics, *31*, 79–105.

doi:10.1146/annurev.ecolsys.31.1.79

Archie, E. A., Morrison, T. A., Foley, C. A. H., Moss, C. J., & Alberts, S. C. (2006).

Dominance rank relationships among wild female African elephants, *Loxodonta*

africana. *Animal Behaviour*, *71*(1), 117–127. doi:10.1016/j.anbehav.2005.03.023

Bartomeus, I., Gravel, D., Tylianakis, J. M., Aizen, M. A., Dickie, I. A., & Bernard-Verdier,

M. (2016). A common framework for identifying linkage rules across different types

of interactions. *Functional Ecology*, *30*(12), 1894–1903. doi:10.1111/1365-

2435.12666

Becerra, J. X. (2003). Synchronous coadaptation in an ancient case of herbivory. *Proceedings*

of the National Academy of Sciences, *100*(22), 12804–12807.

doi:10.1073/pnas.2133013100

Berlow, E. L., Dunne, J. A., Martinez, N. D., Stark, P. B., Williams, R. J., & Brose, U.

(2009). Simple prediction of interaction strengths in complex food webs. *Proceedings*

of the National Academy of Sciences, *106*(1), 187–191. doi:10.1073/pnas.0806823106

Boddy, L. (2000). Interspecific combative interactions between wood-decaying

basidiomycetes. *FEMS Microbiology Ecology*, *31*(3), 185–194. doi:10.1111/j.1574-

6941.2000.tb00683.x

Chase, I. D., & Seitz, K. (2011). Self-structuring properties of dominance hierarchies: a new

perspective. *Advances in Genetics*, *75*, 51–81. doi:10.1016/B978-0-12-380858-

5.00001-0

Cohen, J. E. (1977). Food webs and the dimensionality of trophic niche space. *Proceedings of*

the National Academy of Sciences, *74*(10), 4533–4536. doi:10.1073/pnas.74.10.4533

Cohen, J. E. (2010). Ecologists' Co-Operative Web Bank. Version 1.1. Machine-readable

database of food webs. New York: The Rockefeller University. Retrieved 2

September 2017, from

http://digitalcommons.rockefeller.edu/cohen_joel_laboratory/1/

Cohn, P. M. (2003) Basic algebra: groups, rings and fields. Springer-Verlag, London, UK

Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food

webs' backbones using functional traits. *Oikos*, *125*(4), 446–456.

doi:10.1111/oik.02305

Débarre, F., Nuismer, S. L., & Doebeli, M. (2014). Multidimensional (co)evolutionary

stability. *The American Naturalist*, *184*(2), 158–171. doi:10.1086/677137

Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Dalla Riva, G. V., Fortin, M.-J., …

Poisot, T. (2019). Analysing ecological networks of species interactions. *Biological*

Reviews, *94*(1), 16–36. doi:10.1111/brv.12433

Dy, R. L., Richter, C., Salmond, G. P. C., & Fineran, P. C. (2014). Remarkable mechanisms

in microbes to resist phage infections. *Annual Review of Virology*, *1*, 307–331.

doi:10.1146/annurev-virology-031413-085500

Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N. P., … Allesina, S.

(2013). The dimensionality of ecological networks. *Ecology Letters*, *16*(5), 577–583.

doi:10.1111/ele.12081

Eklöf, A., & Stouffer, D. B. (2016). The phylogenetic component of food web structure and

intervality. *Theoretical Ecology*, *9*(1), 107–115. doi:10.1007/s12080-015-0273-9

Endler, J. A. (1991). Interactions between predators and prey. In J. R. Krebs & N. B. Davies

(Eds.), *Behavioural ecology: an evolutionary approach* (3rd ed., pp. 169–196).

Oxford: Blackwell Scientific Publications.

Fontaine, C., Guimarães Jr., P. R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten, W. H.,

… Thébault, E. (2011). The ecological and evolutionary implications of merging

512 different types of networks. *Ecology Letters*, *14*(11), 1170–1181. doi:10.1111/j.1461-

513 0248.2011.01688.x

514 Garfield, R. (2017). *Magic: The Gathering*® *comprehensive rules*. Renton, WA: Wizards of

515 the Coast LLC. Retrieved 2 September 2017, from

516 http://magic.wizards.com/en/game-info/gameplay/rules-and-formats/rules

517 Gilman, R. T., Nuismer, S. L., & Jhwueng, D.-C. (2012). Coevolution in multidimensional

518 trait space favours escape from parasites and pathogens. *Nature*, *483*(7389), 328–330.

519 doi:10.1038/nature10853

520 Gurobi Optimization and Inc. (2017). Gurobi Optimizer, version 7.5. Retrieved 2 September

521 2017, from http://www.gurobi.com

522 Jackson, J. B. C., & Buss, L. (1975). Allelopathy and spatial competition among coral reef

523 invertebrates. *Proceedings of the National Academy of Sciences*, *72*(12), 5160–5163.

524 doi:10.1073/pnas.72.12.5160

525 Jordano, P., Bascompte, J., & Olesen, J. M. (2003). Invariant properties in coevolutionary

526 networks of plant–animal interactions. *Ecology Letters*, *6*(1), 69–81.

527 doi:10.1046/j.1461-0248.2003.00403.x

528 Kennedy, H. (1978). Systematics and pollination of the "closed-flowered" species of

529 *Calathea* (Marantaceae). *University of California Publications in Botany*, *71*, 1–90.

530 Kiørboe, T. (2011). How zooplankton feed: mechanisms, traits and trade-offs. *Biological*

531 *Reviews*, *86*(2), 311–339. doi:10.1111/j.1469-185X.2010.00148.x

532 Kivelä, M., Arenas, A., Barthelemy, M., Gleeson, J. P., Moreno, Y., & Porter, M. A. (2014).

533 Multilayer networks. *Journal of Complex Networks*, *2*(3), 203–271.

534 doi:10.1093/comnet/cnu016

535 Legendre, P., & Legendre, L. (1998). *Numerical ecology* (2nd ed.). Amsterdam: Elsevier

536 Science.

537 MacArthur, R., & Levins, R. (1967). The limiting similarity, convergence, and divergence of

538 coexisting species. *The American Naturalist*, *101*(921), 377–385. doi:10.1086/282505

McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community

ecology from functional traits. *Trends in Ecology & Evolution*, *21*(4), 178–185.

doi:10.1016/j.tree.2006.02.002

Mello, M. A. R., Rodrigues, F. A., Costa, L. da F., Kissling, D. W., Şekercioğlu, Ç. H.,

Marquitti, F. M. D., & Kalko, V. E. K. (2015). Keystone species in seed dispersal

networks are mainly determined by dietary specialization. Oikos, 124(8), 1031–1039.

doi:10.1111/oik.01613

Meyer, J. R., Dobias, D. T., Weitz, J. S., Barrick, J. E., Quick, R. T., & Lenski, R. E. (2012).

Repeatability and contingency in the evolution of a key innovation in phage lambda.

Science, *335*(6067), 428–432. doi:10.1126/science.1214449

Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic

interactions from proxies. *Trends in Ecology & Evolution*, *30*(6), 347–356.

doi:10.1016/j.tree.2015.03.014

Newman, M. E. J. (2003). The structure and function of complex networks. *SIAM Review*,

45(2), 167–256. doi:10.1137/S003614450342480

Olito, C., & Fox, J. W. (2015). Species traits and abundances predict metrics of plant–

pollinator network structure, but not pairwise interactions. *Oikos*, *124*(4), 428–436.

doi:10.1111/oik.01439

Ortega, R., Fortuna, M. A., & Bascompte, J. (2017). Web of Life dataset. Retrieved 2

September 2017, from http://www.web-of-life.es/

Poisot, T., Cirtwill, A. R., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. B. (2016).

The structure of probabilistic networks. *Methods in Ecology and Evolution*, *7*(3),

303–312. doi:10.1111/2041-210X.12468

Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: why ecological interaction

networks vary through space and time. *Oikos*, *124*(3), 243–251.

doi:10.1111/oik.01719

Rohr, R. P., & Bascompte, J. (2014). Components of phylogenetic signal in antagonistic and

mutualistic networks. *The American Naturalist*, *184*(5), 556–564.

doi:10.1086/678234

Santamaría, L., & Rodríguez-Gironés, M. A. (2007). Linkage rules for plant–pollinator

networks: trait complementarity or exploitation barriers? *PLOS Biology*, *5*(2), e31.

doi:10.1371/journal.pbio.0050031

Santamaría, L., & Rodríguez-Gironés, M. A. (2015). Are flowers red in teeth and claw?

Exploitation barriers and the antagonist nature of mutualisms. *Evolutionary Ecology*,

29(3), 311–322. doi:10.1007/s10682-015-9757-8

Schiestl, F. P., & Johnson, S. D. (2013). Pollinator-mediated evolution of floral signals.

Trends in Ecology & Evolution, *28*(5), 307–315. doi:10.1016/j.tree.2013.01.019

Schoener, T. W. (1989). The ecological niche. In J. M. Cherrett (Ed.), *Ecological concepts:*

the contribution of ecology to an understanding of the natural world (pp. 79–114).

Oxford: Blackwell Scientific Publications.

Shizuka, D., & McDonald, D. B. (2015). Data from: The network motif architecture of

dominance hierarchies. *Dryad Digital Repository*. Retrieved 16 August 2017, from

http://dx.doi.org/10.5061/dryad.f76f2

Stanko, M., & Miklisova, D. (2014). Data from: Empirical evaluation of neutral interactions

in host-parasite networks. *Dryad Digital Depository*. Retrieved 17 August 2017, from

http://datadryad.org/resource/doi:10.5061/dryad.70sj1

Stouffer, D. B., Camacho, J., & Amaral, L. A. N. (2006). A robust measure of food web

intervality. *Proceedings of the National Academy of Sciences*, *103*(50), 19015–19020.

doi:10.1073/pnas.0603844103

Stouffer, D. B., Rezende, E. L., & Amaral, L. A. N. (2011). The role of body mass in diet

contiguity and food-web structure. *Journal of Animal Ecology*, *80*(3), 632–639.

doi:10.1111/j.1365-2656.2011.01812.x

Szolnoki, A., Mobilia, M., Jiang, L.-L., Szczesny, B., Rucklidge, A. M., & Perc, M. (2014).

Cyclic dominance in evolutionary games: a review. *Journal of The Royal Society*

Interface, *11*(100), 20140735. doi:10.1098/rsif.2014.0735

Thompson, R. M., & Townsend, C. R. (2004). Interaction Web DataBase. Retrieved 17

August 2017, from

https://www.nceas.ucsb.edu/interactionweb/html/thomps_towns.html

Vázquez, D. P., & Aizen, M. A. (2003). Null model analyses of specialization in plant–

pollinator interactions. *Ecology*, *84*(9), 2493–2501. doi:10.1890/02-0587

Vázquez, D. P., Chacoff, N. P., & Cagnolo, L. (2009). Evaluating multiple determinants of

the structure of plant–animal mutualistic networks. *Ecology*, *90*(8), 2039–2046.

doi:10.1890/08-1837.1

Williams, H. P. (2013). *Model building in mathematical programming* (5th ed.). West Sussex:

Wiley-Blackwell.

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J. M., Olesen, J. M., Valido, A., &

Warren, P. H. (2005). Body size in ecological networks. *Trends in Ecology &*

Evolution, *20*(7), 402–409. doi:10.1016/j.tree.2005.04.005

Woodward, G., Papantoniou, G., Edwards, F., & Lauridsen, R. B. (2008). Trophic trickles and

cascades in a complex food web: impacts of a keystone predator on stream

community structure and ecosystem processes. *Oikos*, *117*(5), 683–692.

doi:10.1111/j.2008.0030-1299.16500.x