A near complete haplotype-phased genome of the dikaryotic wheat stripe rust

fungus *Puccinia striiformis* f. sp. *tritici* reveals high inter-haplome diversity

Authors:

Benjamin Schwessinger[a,*,+], Jana Sperschneider[b], William S. Cuddy[c,d,+,%], Diana P.

Garnica[a,$], Marisa E. Miller[f], Jennifer M. Taylor[h], Peter N. Dodds[h], Melania Figueroa[f,g],

Robert F. Park[c], John P. Rathjen[a,+]

Research School of Biology, The Australian National University, Acton 2601, ACT,

Australia[a]; Centre for Environment and Life Sciences, CSIRO Agriculture and Food,

Perth, WA, Australia[b]; Plant Breeding Institute, Faculty of Agriculture and Environment,

The University of Sydney, Narellan, NSW, Australia[c]; Elizabeth Macarthur Agricultural

Institute, NSW Department of Primary Industries, Woodbridge Road, Menangle, NSW,

2568, Australia[d]; Department of Plant Pathology, University of Minnesota, St. Paul, MN,

USA[f]; Stakman-Borlaug Center for Sustainable Plant Health, University of Minnesota,

St. Paul, MN, USA[g]; [h]Black Mountain Laboratories, CSIRO Agriculture and Food,

Canberra, ACT, Australia; [$]Current address: Department of Agriculture and Water

Resources, Canberra City ACT 2601 Australia

Running head: Haplotype phasing of the dikaryotic wheat stripe rust fungus *Puccinia*

striiformis f. sp. *tritici*

[+] Co-corresponding authors

*Address correspondence about data analysis to Benjamin Schwessinger

(benjamin.schwessinger@anu.edu.au, +61 405919737)

[%]Address correspondence about biological material to William S. Cuddy

(will.cuddy@dpi.nsw.gov.au, +61 2 4640 6515)

Abstract:

A long-standing biological question is how evolution has shaped the genomic

architecture of dikaryotic fungi. To answer this, high quality genomic resources that

enable haplotype comparisons are essential. Short-read genome assemblies for

dikaryotic fungi are highly fragmented and lack haplotype-specific information due to the

high heterozygosity and repeat content of these genomes. Here we present a diploid-

aware assembly of the wheat stripe rust fungus *Puccinia striiformis* f. sp. *tritici* based on

long-reads using the FALCON-Unzip assembler. RNA-seq datasets were used to infer

high quality gene models and identify virulence genes involved in plant infection referred

to as effectors. This represents the most complete *Puccinia striiformis* f. sp. *tritici*

genome assembly to date (83 Mb, 156 contigs, N50 1.5 Mb) and provides phased

haplotype information for over 92% of the genome. Comparisons of the phase blocks

revealed high inter-haploid diversity of over 6%. More than 25% of all genes lack a clear

allelic counterpart. When investigating genome features that potentially promote the

rapid evolution of virulence, we found that candidate effector genes are spatially

associated with conserved genes commonly found in basidiomycetes. Yet candidate

effectors that lack an allelic counterpart are more distant from conserved genes than

allelic candidate effectors, and are less likely to be evolutionarily conserved within the *P.*

striiformis species complex and *Pucciniales*. In summary, this haplotype-phased

assembly enabled us to discover novel genome features of a dikaryotic plant

pathogenic fungus previously hidden in collapsed and fragmented genome assemblies.

Importance:

Current representations of eukaryotic microbial genomes are haploid, hiding the

genomic diversity intrinsic to diploid and polyploid life forms. This hidden diversity

contributes to the organism's evolutionary potential and ability to adapt to stress

conditions. Yet it is challenging to provide haplotype-specific information at a whole-

genome level. Here, we take advantage of long-read DNA sequencing technology and a

tailored-assembly algorithm to disentangle the two haploid genomes of a dikaryotic

pathogenic wheat rust fungus. The two genomes display high levels of nucleotide and

structural variations, which leads to allelic variation and the presence of genes lacking

allelic counterparts. Non-allelic candidate effector genes, which likely encode important

pathogenicity factors, display distinct genome localization patterns and are less likely to

be evolutionary conserved than those which are present as allelic pairs. This genomic

diversity may promote rapid host adaptation and/or be related to the age of the

sequenced isolate since last meiosis.

Introduction

The Basidiomycota and the Ascomycota constitute the two largest fungal phyla and

contain many of the most damaging crop pathogens (1). The dominant life phase for

most basidiomycete species is dikaryotic, where two haploid nuclei coexist within one

cell (2). To date, about 475 basidiomycete fungal genome sequences representing

some 245 species are available in the public domain (September 2017,

https://www.ncbi.nlm.nih.gov/genome/). These genome references are either

representations of the haploid life stage of a species (3), or collapsed and mosaic

assemblies of the dikaryotic state (4–7). Hence, the information about the inter-haploid

genome variation in dikaryotic basidiomycota beyond single nucleotide polymorphisms

(SNPs) and small insertions and deletions (INDELs) is very limited. The absence of

haplotype-phased information limits the studies of genome architecture and evolution,

particularly for the rust fungi of the order *Pucciniales*, many of which are extremely

destructive pathogens of economically important crops including cereals, coffee, and

soybean (8–13).

Stripe, stem and leaf rusts are the three rust diseases affecting wheat production, one

of the most important staples in human diets (14). Of these, stripe rust caused by

Puccinia striiformis f. sp. *tritici* (*Pst*) is currently the most damaging disease with

estimated annual losses of $USD 1 billion (15, 16). As a biotrophic pathogen, *Pst*

colonizes living hosts and extracts large amounts of nutrients from plant cells through

specialized structures called haustoria. The large tax on host energy reserves caused

by *Pst* infection results in yield losses mostly associated with poor grain filling (17).

The full life cycle of *Pst* involves asexual and sexual reproductive phases associated

with the production of specific spore types (17, 18). The damage to wheat occurs during

the asexual cycle and results from the repeated infections throughout the growing

season, which cause exponential amplification of dikaryotic urediniospores. *Pst* infects

more than 30 varieties of *Berberis* spp. and *Mahonia* spp. to complete its full sexual life-

cycle involving four additional spore stages and sexual recombination during meiosis

(19–21). The sexual stages are restricted geographically to the Himalayan region

(Nepal, Pakistan and China), where it leads to high levels of genetic diversity that are

largely absent from other parts of the world. This makes the extended Himalaya region

the center of diversity of *Pst* and the main source for new highly virulent *Pst* isolates

(12, 22).

Genetic resistance in the host plant, particularly race-specific resistance, is often used

in the field to reduce damage by pathogenic rust fungi (23, 24). Race-specific resistance

is generally conferred by dominant resistance (*R*) genes in the host which recognize

specific avirulence (*Avr*) alleles within the pathogen. Mechanistically, *Avr* alleles encode

variants of virulence effector proteins, and the *R* gene typically encodes a nucleotide-

binding leucine rich repeat (NB-LRR) protein that detects the Avr protein within the

infected plant cell. In case of *Pst*, more than 75 *y*ellow *r*ust resistance genes (*Yr*) have

been catalogued to date. A given *Pst* isolate has a characteristic spectrum of *Avr* alleles

that can be distinguished on a set of wheat tester lines containing these *Yr* genes (25).

The collective virulence phenotypes on the differential set defines the *Pst* pathotype.

Wheat stripe rust epidemics are associated with the appearance of genetically novel

pathotypes which are not recognized by currently employed *R* genes and hence grow

on commercial wheat cultivars. As such, incursions of exotic stripe rust isolates with

new virulences can play a role in disease outbreaks, for instance the Warrior *Pst* isolate

which invaded Europe in 2011 was highly successful as it was virulent on the wheat

cultivars cultivated at that time (26, 27). In addition to this novel exotic incursion, it is

well documented that *Pst* rapidly evolved new virulence traits on a continental scale in

Australia following its introduction in 1979 (28). However, the mechanisms underlying

the evolution of these new pathotypes remains understudied as no genetic locus

contributing to the evolution of virulence has yet been identified in *Pst*. While new

combinations of alleles generated during sexual recombination can lead to the

emergence of new pathotypes, the contribution of other genetic and molecular events to

pathogen evolution during asexual reproduction is unclear. Presumably, the occurrence

of mutation explains the loss of *Avr* specificities and the adaptation to otherwise

resistant wheat varieties (13, 28).

Most agriculturally important fungi are haploid with small genomes (29). Rusts on the

other hand are dikaryotic in the asexual phase and have expanded genomes with large

amounts of repetitive sequence (6, 7). It is likely that the separation of rust genomes

into two haploid copies contributes to their rapid evolution. Existing *Pst* genome

sequences suffer from the use of short-read sequencing technologies, which prevents

characterization of individual haploid genomes (referred to as haplome), while the high

percentage of repetitive DNA reduces the size of contigs that can be assembled (4, 5,

30). The overall similar gene content of each genome causes the reads from allelic

variants to collapse upon assembly, producing a consensus sequence that loses

haplotype (phasing) information. Read mapping to the consensus reference reveals that

the two genomes are highly heterozygous for SNPs (5, 7), but differences in effector

and gene content are undetectable. These problems can be addressed to some extent

by using traditional Sanger long-sequence reads or strategies such as fosmid-to-fosmid

sequencing (6, 7), however, these approaches are expensive. Opportunities to resolve

the questions at higher resolution have arisen from new technologies that generate very

long sequencing reads (31, 32).

Here, we use long-read sequencing to provide a near-complete haplotype phased

genome assembly for an isolate of the first pathotype of *Pst* detected in Australia in

1979 (28). It is the most complete *Pst* genome to date with over 97% of all

basidiomycete benchmarking universal single copy orthologs (BUSCOs) captured (33).

In addition, phased haplotype information for over 92% of the genome enabled us to

detect high inter-haplome diversity at the nucleotide and structural level, which leads to

allelic variation and 25% of all genes lacking a clear allelic counterpart. We identified

over 1,700 candidate effector genes, which are more often spatially associated with

each other and conserved BUSCOs rather than a clear preferentially spatial association

with repetitive elements. Non-allelic candidate effectors that lack counterparts in the

other haplome are less likely to be evolutionarily conserved in other rust fungi. Thus, the

highly contiguous haplotype assembly has allowed discovery of novel genome features

that may be linked to the rapid evolution of this devastating pathogen.

Results and Discussion

Haplotype-aware genome assembly of an Australian *Puccinia striiformis* f. sp.

***tritici* isolate**

The main aim of this study was to generate a high quality reference genome for *Pst*. For

this purpose we sequenced a single pustule isolate of the Australian founder pathotype

Pst 104E137A-, collected in 1982 (abbreviated as *Pst*-104E). We sequenced 13 PacBio

SMRT cells obtaining a total of 13.7 Gb of data with an average read length of 10,710

bases, and a read length N50 of 15,196 bases (Table S1). We assembled these data

using the diploid-aware assembler FALCON-Unzip (31) to obtain a synthetic haplotype-

phased reference genome. The FALCON-Unzip assembler is designed to phase

structural variations and associated single nucleotide polymorphisms (SNPs) into

distinct haplotype blocks. This gives rise to a primary assembly (primary contigs) and

linked haplotype blocks (haplotigs). The haplotigs represent the alternative genome

structure with respect to primary contigs. FALCON-Unzip does not always link

physically connected phase blocks, and primary contigs can represent sequences from

either of the two haploid genomes (31).

Previous unphased *Pst* genome assemblies ranged in size between 53 and 115 Mb (4,

5, 7, 30). In an attempt to reconcile the difference in the reported genome sizes, we

used GenomeScope to estimate the haploid genome size using *k*-mer frequencies (30-

mer) in two Illumina short-read datasets of *Pst*-104E (34). Based on this analysis we

estimated a haploid genome size of 68-71 Mb, with a heterozygosity (SNPs and

INDELs) rate of approximately 1.2%. We assembled our long-read data into 156

primary contigs with a total length of 83 Mb after manual curation. The corresponding

phased haplotype blocks were contained in 475 haplotigs with a total size of 73 Mb

(Table 1).

These assembly statistics are a vast improvement over previous assemblies in terms of

connectivity and number of contigs (Figure 1A). The primary assembly has a contig N50

of 1.3 Mb compared to a scaffold N50 of 0.5 Mb for *Pst*-78 or contig N50 of 5.1 Kb for

Pst-130, often referred to as the reference genome (4, 27, 30). In addition we identified

1,302 (97.5%) out of the 1,335 benchmarking genes (BUSCO v2;

http://busco.ezlab.org/) (33) that are highly conserved in basidiomycetes, with only 10

(0.7%) missing in our combined assembly before filtering for genes related to

transposable elements (TE). Our final assembly has 1,292 (96.8%) complete BUSCOs

with 19 (1.4%) missing. This compares to wide variation in BUSCOs that can be

identified from previous assemblies, ranging from 35.7% for *Pst*-887 to 95.6% for *Pst*-78

(Figure 1B). In summary, our assembly is currently the most complete *Pst* genome in

terms of contiguity, haplotype-phased information, and gene content. This advance

provides a new resource to investigate genome architecture and inter-haplotype

variation within this dikaryotic plant pathogen.

High levels of inter-haplotype block variation

The *Pst*-104E primary assembly covers 83 Mb in a total of 156 primary contigs. Within

this assembly, 99 primary contigs (~80 Mb) are associated with 475 haplotigs (~73 Mb),

representing phased information for 92% of the primary contigs. These primary contigs

are referred to as primary contigs with haplotigs. The remaining 57 primary contigs

(~3.6 Mb) have no associated haplotig (Table 1). Of these, 51 are likely single-

haplotype-specific sequences, because these display similar average coverage to

phased regions of the genome when mapping independently generated Illumina short-

read data to the PacBio long-read assembly. This high level of phasing enabled us to

investigate the inter-haplotype variation at a whole-genome scale. Previous studies

using Illumina short-reads mapped against the consensus merged haplotype

assemblies estimated the inter-haplome variations based on heterozygous SNPs at

between 0.5% - 1% (5, 7, 30). Taking a similar approach, we identified approximately

0.5% (416,460 heterozygous SNPs) of the genome as variable when mapping Illumina

short reads against primary contigs only. However, we estimated a dramatically higher

level of inter-haplome variation when using this phased assembly. For this analysis, we

aligned all haplotigs with their corresponding primary contigs and estimated variations

using Assemblytics (35, 36). Assemblytics defines six major categories of (structural)

variations including insertions and deletions, tandem repeats identified by overlapping

alignments and other types of repeats suggested by gapped non-unique contig

alignments (see Figure 2A for illustration of the six different variant categories) and

divides these according to size in bins (Figure 2A). This analysis revealed that structural

variation comprised 6.4% (~5.10/79.77 Mb) of the primary assembly space when

compared to their respective haplotigs (Figure 2A) (35). The variation of two primary

contigs and their respective haplotigs is illustrated in the dot plots shown in Figure 2B

and C, which visualizes large-scale inversions, deletions and insertions in haplotigs

associated with two primary contigs. It is plausible that the actual difference between

the two haplotypes is higher than the estimated 6%, because calculations were

restricted to a maximal variant size of 10 kb and do not include primary contigs without

haplotigs. Overall, the dramatic difference in estimated inter-haplome variation between

previous assemblies (5, 7, 30) and short-read based prediction programs (34) is likely

caused by the fact that most of the observed variations are contained in size bins

greater than 500 bases, which is not detectable with Illumina short-read data and by

highly fragmented assemblies.

Over half of the Pst-104E genome is covered by repetitive

We annotated primary contigs and haplotigs independently based on our observations

of high levels of heterozygosity between the two (Figure 2 and S1). We first identified

and classified transposable elements (TEs) using the REPET pipeline (37) to the order

level based on the Wicker classification (38). We further transferred superfamily

annotation from the underlying BLAST (39) hits if they agreed with the REPET

annotations and with each other. There was no major difference between TE coverage

of primary contigs 54% (~45 Mb) and haplotigs with 53% (~39 Mb) (Figure S1).

However, primary contigs that lacked haplotigs had a larger proportion of TEs with a

total coverage of 67%, which might explain their increased fragmentation and reduced

contig length (Table 1). The composition of TE families on primary contigs versus

haplotigs was very similar (Figure S1). Both retrotransposons (Class I) and DNA

transposons (Class II) cover 30% of the genome each (note that distinct TE classes can

overlap). For Class I transposons, the long terminal repeat (LTR) order was the most

prominent with ~27% coverage, and within this order elements from the Gypsy and

Copia superfamilies were most prominent. The only other Class I orders with greater

than 1% genome coverage were LARD and DIRS elements. Class II elements were

dominated by TIR elements with a genome coverage of ~20%, with significant

contributions of elements belonging to the hAT, MuDR, PIF-Harbinger, Tc1-Mariner,

and CATCA superfamily. More than 6% of the genome was covered by Class II

elements that could not be classified below the class level and showed no homology to

previously identified TEs. This is in contrast to the minimal coverage by unclassifiable

Class I elements (0.05%). Overall, this is the highest number of identified transposable

elements detected in any *Pst* genome assembly so far as previous reports varied from

17% to 50% (4, 7, 30). Such increased content of identified transposable elements is

likely caused by the increased contiguity and the absence of any unidentified bases

(Ns) in our assembly (Figure 1).

High levels of inter-haplome structural variation lead to variable gene content

between primary contigs and haplotigs

We also annotated gene models on primary contigs and haplotigs independently using

an extensive set of newly generated and publicly available RNA-seq data (40). This is in

contrast to the published *Pst* genomes that are nearly exclusively annotated using *ab*

initio gene finding approaches without specific gene expression data (4, 5, 7, 30). The

newly generated RNA-seq datasets were obtained from dormant and germinated

urediniospores, wheat leaf tissue six and nine days post infection (dpi), and haustoria-

enriched fractions. These datasets were complemented by publicly available RNAseq

data from germinated spores, and infected wheat tissue sampled at 13 different time

points x plant genotype combinations (40). We used this extensive expression data in a

comprehensive genome annotation pipeline (41–45) and identified 15,928 and 14,321

gene models on primary contigs and haplotigs respectively, after filtering for genes

related to TE function (Tables 1 and S2) (46, 47). The protein sequences of these

genes were functionally annotated using an extensive set of tools (Table S2 and File

S1) (33, 48–52). We obtained very similar annotation levels for primary contigs and

haplotigs with about 52% of all proteins having at least one functional annotation in the

following categories; GO terms, InterPro match, Pfam domain, EggNog term, KEGG

pathway annotation, MEROPs catalytic domain, or carbohydrate hydrolyzing enzymatic

domains (CAZy) (33, 48–52). The level of functional annotation for *Pst* proteins

identified as BUSCO orthologs was near complete with only two or one protein (< 0.1%)

lacking any functionally recognizable domain (Table S2). This pattern was reversed

when characterizing candidate effectors (see identification below) as approximately

83% of all proteins lacked a conserved functional domain.

Overall, the haplotype-phased assembly did not show biased distribution of any

particular gene annotation group (Table S2); this is consistent with the high level of

haplotype phasing. This encouraged us to investigate the relationship between the two

haplotype phase block assemblies (primary contigs compared to haplotigs) in terms of

gene content. One must keep in mind that these two assemblies do not actually

represent the true haploid genomes, because of potential haplotype switching between

primary contigs and haplotigs, and the inability to assert independent contigs to one

specific haplome (31). However, we argue that a relational comparison between the two

assemblies is still valuable in order to investigate the approximate inter-haplome gene

diversity. Therefore, to simplify the analysis we treated primary contigs and haplotigs as

two representative genetic units. We used Proteinortho in the synteny mode to identify

syntenic allele pairs between the primary contigs and haplotigs (53). We identified a

total of 10,921 potential syntenic allele pairings including 10,785 primary proteins and

10,860 haplotig proteins (Table S3, File S2). Of these, 9,756 were properly paired

where the haplotig gene models were located on an associated haplotig that overlapped

with the primary gene model when performing targeted whole-genome alignments

(Figure S2A, Table S2). These correspond to 'classic' alleles in a diploid organism.

Another 450 pairs were not directly linked as the haplotig containing the allelic ortholog

did not overlap with the primary gene model although it was associated with the primary

contig (Figure S2B and File S2). These may be simple rearrangements linked to

inversions or repeat duplications. A further 715 pairs were completely unlinked as the

allele-containing haplotig was not associated with the respective primary contig in our

assembly (Figure S2C and File S3). We randomly selected 176 of these loci and

investigated them manually by whole genome alignment of haplotigs to primary contigs,

followed by micro-synteny analysis of the identified gene loci (36, 54, 55). An example

of this analysis is illustrated in Figure 3. In this case, a ~40 kb region present both in

primary contig 014 and haplotig 027_006 showed micro-synteny for three genes each,

namely *Pst104E_05635-05637* and *Pst_104E_24450-24452*, respectively (Figure 3D),

while the overall macro-synteny was not conserved (Figure 3A-C). This may have been

caused by genetic transposition of the identified region from the chromosomal region

corresponding to a haplotig that fully aligned with primary contig 014 into the sequence

of the chromosomal region corresponding to haplotig 027_006. We found support for

such allele transposition, either via cut and paste or copy and paste mechanisms, in

71/176 cases. The remaining cases could not be categorized confidently and may

represent complex genomic regions, genetically linked contigs that were broken up

during the assembly process, gene duplication events, or miss-assemblies. Based on

this manual inspection we estimate that approximately 280 loci (71/176*715 total pairs)

containing alleles might be rearranged in one of the two haploid genomes. We identified

a further 912 loci that clustered at the protein level yet their genomic location was not

syntenic between the two haplotype phase block assemblies (File S3). We refer to

these genes as non-allelic inter-haplome paralogs. In summary, this suggests that over

3% (~1,192/30,249) of all genes are closely related at the protein level but do not reside

in regions displaying macro-synteny.

We identified 4,761 primary and 2,931 haplotig genes that did not cluster at the

protein level using Proteinortho and hence may represent singletons, with singletons

being defined as genes of a diploid organism that lack inter-haplome alleles and

paralogs (Table S3). Of the 4,761 primary genes, 663 were located in regions where the

assembly was not haplotype-phased based on coverage analysis using Illumina short-

read data (File S4). Hence we identified 7,029 true singletons (File S5) when comparing

both haplotype phase block assemblies and 1,506 of these singletons are referred to as

single haplotype genes (File S5), because they lacked any BLAST hit (blastn, e-value <

0.01) when using the gene sequence as a query against the other haplotype-phase

block genome sequence. These single haplotype genes are often linked in clusters,

because for 1,164 single haplotype genes at least one of their nearest neighbors is also

a haplotype-specific gene, compared to 212 of an equally sized random subsample of

all genes (Fisher's exact test, p-value ~ $2.3*10^{-109}$). Similarly, 1,492 haplotype-specific

genes are located in regions where primary contigs and associated haplotigs do not

align, indicating haplotype specific regions. Single haplotype genes are highly enriched

in these regions as only 251 of an equally sized random subsample of all genes

displayed a similar location (Fisher's exact test, p-value ~ $4.5*10^{-265}$). Taken together,

these findings suggest that there are numerous large presence-absence structural

polymorphisms between the two haploid genomes that can span multiple adjacent

genes, and therefore represent many of the haplotype-specific genes. To study the

overall conservation of these single haplotype genes we queried them against the

EnsemblFungi cDNA and NCBI nr databases (blastn, e-value < 0.01) (56, 57). Out of

1506 genes, 1424 had at least one significant hit in either database, with the top hits in

all cases being fungal sequences. The remaining 82 genes lacked any sequence

homology to known fungal genes. These genes were significantly shorter compared to

all genes (mean length 538 bases versus 1538, two-sided Student's t-test, p-value ~

$2.38e^{-07}$). We identified expression evidence for 27/82 of these genes including 7 of 10

predicted candidate effectors. This is consistent with observations in other fungi for

which isolate-specific genes tend to be shorter and lower expressed than genes that are

conserved between isolate (58). Overall, the high levels of non-allelic genes (~25%) and

single haplotype genes (~5%) illustrates that the large inter-haplome polymorphism at

the nucleotide and structural level (Figure 2 and 3B, C) results in significant differences

in gene content between the two.

Complementary candidate effector gene prediction using machine learning and *in*

***planta* expression data**

The diversity of plant pathogen effectors makes them impossible to identify based on

protein sequences alone (59). Only a small number of effectors have thus far been

confirmed in rust fungi, namely AvrP123, AvrP4, AvrL567, AvrM, RTP1, and

PGTAUSPE-10-1 (60), AvrL2 and AvrM14 (61), PstSCR1 (62) and PEC6 (63). At the

sequence level, effectors do not share common domains or motifs, apart from the

presence of a signal peptide. To predict candidate effectors in *Pst*-104E, we utilized a

combination of gene expression analysis and machine learning methods. First, we

predicted fungal rust secretomes based on a protocol optimized for recovering fungal

candidate effectors (64). We observed large differences in secretome sizes across rust

proteomes, e.g. the stripe rust isolate *Pst*-887 had a small secretome compared to *Pst*-

104E (Table S4). Overall the number of secreted proteins appeared to correlate with

completeness of *Pst* genome assemblies based on BUSCO analysis (Figure 1B and

Table S4). This implies that it is difficult to perform comprehensive orthology analyses

between current *Pst* assemblies given that many appear to be incomplete in terms of

BUSCOs and therefore are likely incomplete for other gene families also, including

secreted proteins.

To predict candidate effectors, we used the machine learning approach EffectorP on all

secreted proteins without predicted transmembrane domains (64). Overall we identified

1,069 and 969 candidate effectors from primary contigs and haplotigs respectively (File

S6). We complemented this *in silico* approach with a detailed expression analysis of

Pst-104E genes that encode secreted proteins. We used gene expression data and *k*-

means clustering to predict clusters in the secretome that are differentially expressed

during infection and exhibit similar expression profiles (Figure 4, File S7). For the

primary contigs of *Pst*-104E, this resulted in eight predicted clusters. The expression

profiles of three clusters (clusters 2, 3 and 8) resembled the expected expression

patterns of haustorially-delivered cytoplasmic rust effectors, namely a high expression in

haustorial tissue and at the infection time points (6 and 9 dpi) as well as low expression

in spores (Figure 4A). In total there are 809 genes in clusters 2, 3, and 8, of which 306

(~38%) were also identified by EffectorP as candidate effectors (Table S5). Upon closer

inspection of primary contig expression patterns, cluster 8 in particular exhibits the

highest overall haustorial expression and overall lowest expression in spores, indicating

it is likely to contain cytoplasmic effectors. Interestingly, whilst cluster 8 shows the

lowest percentage of EffectorP predicted candidate effectors (26%), it has the highest

percentage of proteins with a predicted nuclear localization signal (NLS) (Table S5)

(65). We also observed that proteins in cluster 8 are mostly larger (average length of

410 aa) than other known rust effectors (the largest is AvrM, at 314 aa), which might

indicate that *Pst* utilizes a class of larger effector proteins that target host nuclei.

Similarly, oomycete pathogens secrete a class of cytoplasmic effectors called Crinklers

that carry NLSs (66, 67) and these are not predicted as candidate effectors by

EffectorP, possibly due to their larger size. Therefore, we included both *in planta* up-

regulated secreted proteins as well as EffectorP predicted proteins as candidate

effectors. In total, we identified 1,572 candidate effectors on primary contigs when

combining predictions based on *in planta* expression analysis and EffectorP. We

identified similar expression patterns for secreted proteins on haplotigs. Clusters 11, 13,

14 and 15 shared a similar expression profile to clusters 2, 3, and 8 and contained 673

genes (Table S6 and S7). Of these, 234 (~37%) were also identified by EffectorP

amounting to a total of 1,388 candidate effectors on haplotigs. Overall, we identified a

set of 1,725 non-redundant candidate effectors when combining all candidate effectors

on primary contigs and haplotigs (File S8).

Candidate effector genes are spatially associated with conserved genes and with

each other

For many filamentous plant pathogens a 'two speed' genome has been suggested to

contribute to rapid evolution in terms of candidate effector variability (68). For example,

in fungal plant pathogens such as *Fusarium oxysporum* spp. and *Verticillium dahliae,*

lineage specific genomic regions and/or dispensable chromosomes are enriched for

TEs and candidate effector genes (69–74). In several *Phytophthora* spp., candidate

effectors have been reported to localize in gene sparse, TE-rich regions, which show

signs of accelerated evolution (68, 69, 74). It is not known if rust genomes have a

comparable genome architecture that facilitates rapid evolution of candidate effector

genes. Hence we investigated the genomic location of candidate effectors in relation to

several genomic features including TEs, neighboring genes, BUSCOs, other candidate

effectors, and AT content (Figures 5, 6, S3, and S4). We focused mostly on candidate

effectors on primary contigs, because the primary assembly is far more contiguous

compared to its haplotigs thereby empowering our analysis (Figure 1). In addition, we

made use of our haplotype phased assembly and investigated if allelic candidate

effector variants show distinct features compared to haplotype-singletons. In all cases

we used a random subset of genes and BUSCO gene sets as control groups. We

envisioned BUSCO genes as a particularly well-suited control group as these are

conserved within the phylum of basidiomycetes (75) and can therefore be considered as

part of the *Pst* core genome. On the contrary, candidate effector genes are reported to

be more specific to the class, species, or isolate level (6, 76). This observation also

holds true for *Pst*-104E, because we only observed 40 BLAST hits outside the class of

Pucciniomycetes for 1,725 non-redundant candidate effectors using EnsemblFungi

cDNA as reference (blastn, e-value $1*e^{-5}$).

We first tested if candidate effectors are located in gene sparse regions when compared

to all genes or BUSCOs. For this analysis, we generated density plots with the 5' and 3'

prime distance to its closest neighbor for each gene (68). When comparing gene

distance density hexplots we observed very similar distributions between candidate

effectors and all genes. Candidate effectors in general did not appear to be located in

gene sparse regions and neither did BUSCOs (Figure 5A). Similar effects have been

reported for other rust species, such as the oat crown rust pathogen *Puccinia coronata*

f. sp. *avenae* (77). Next, we tested if candidate effectors are linked to TEs as observed

for other plant pathogenic fungi (74). We compared the minimum distance of all genes,

BUSCOs and candidate effectors to TEs. Candidate effectors globally did not display a

preferential association with TEs when compared with genes in general (Figure 5B).

However, on close examination of the relative spatial distribution of TEs, candidate

effectors and BUSCOs on the 30 largest contigs we could identify some regions where

candidate effectors are closely associated with TEs (Figure S3). The observation that

candidate effectors are not associated globally to TEs is consistent with reports of other

rust fungi including *P. coronata* f. sp. *avenae*, *P. graminis tritici*, and *Melampsora larici-*

populina (6, 77). In case of *Pst*, we aim to answer the question of TEs involvement in

the evolution of novel virulences on a continental scale by re-sequencing *Pst*-104E

mutant progeny with distinct virulence profiles, collected in Australia between 1980 and

2003 (28).

The observation that candidate effectors and BUSCOs show similar localization

patterns relative to all genes and TEs led us to investigate if these two gene groups are

spatially associated, and if each group clusters with itself. We first compared the

minimum distance between genes of the same group when subsampling to an equal

number of genes in each group. Indeed, when comparing the minimum distances

between candidate effectors we found that these were less than the minimum distances

between a random subset of genes (Figure 5C). Also, BUSCOs were more closely

associated with each other than a random subset of genes. Consistently, when we

investigated the number of candidate effectors clustered within a minimum given

distance we found that candidate effectors are more clustered than BUSCOs or an

equal sized random subset of all genes (Figure S4). A similar trend yet to a lesser

degree was observed for BUSCOs. Clustering of candidate effectors was also identified

as a feature of several smut fungi including *Ustilago maydis* and *Sporisorium*

scitamineum (3, 78). In these related basidomycete plant pathogens, candidate effector

gene clusters are born via tandem duplication and hypothesized to contribute to the

rapid evolution of these genes linked to TEs.

The observed spatial association of both BUSCOs and candidate effectors with

themselves led us investigate if these two gene groups are spatially associated with

each other. Indeed, candidate effectors were located more closely to BUSCOs and vice

versa when compared to a random subsample of genes (Figure 5D). This is a surprising

observation because BUSCOs are defined by their overall conservation while candidate

effectors are far less conserved. In obligate biotrophic fungi a subset of effectors might

be essential, because host colonization is an absolute requirement for survival. This is

in contrast to plant pathogens that are also able to grow saprophytically such as *Z.*

tritici, *V. dahliae, U. maydis* and *P. infestans* (3, 72, 73, 79). Hence it might be

advantageous for obligate biotrophs like *Pst* to link these essential effectors to core

genome components to ensure their inheritance and conservation within the species

complex.

The candidate effector allele status influences association with conserved genes

and evolutionary conservation

We next investigated if the distance of candidate effectors to BUSCOs is correlated with

their allelic variation. We calculated the normalized Levenshtein distance of cDNA and

amino acid alignments for all allele pairs. The normalized Levenshtein distance

measures the required single-character edits (insertions, deletions or substitutions) to

convert two strings into each other, e.g. an alignment of two allele sequences, while

accounting for difference in sequence length. It can therefore be used as a proxy for

sequence variation between two alleles (80). We did not observe any significant

difference between the Levenshtein distance at the cDNA level when comparing

BUSCOs and candidate effectors, whereas alleles of all other genes were more variable

than candidate effectors (Table 2). This was in contrast to the variation seen at the

protein level where candidate effectors were more variable than BUSCOs (Table 2).

This suggests that changes at the DNA level are more likely to result in changes to the

protein sequence in case of candidate effectors. We therefore also calculated the ratio

of nonsynonymous to synonymous mutations for all alleles (dN/dS ratio) whenever

possible (81). Indeed, analysis of the dN/dS ratios supported our previous observation

that for candidate effectors, changes in the DNA sequence were more likely to alter the

protein sequence (Table 2). This suggests that candidate effectors evolve faster than

BUSCOs and most other allele pairs even though they are spatially associated with

BUSCOs. The sequence variation in candidate effector allele pairs was not correlated

with distance to the closest BUSCO, using either Levenshtein distances on the protein

level or dN/dS as a proxy (Spearman, correlation < |0.06|, p > 0.15). Subsequently, we

investigated if candidate effector singletons were more distant to BUSCOs than their

allelic counterparts. These singletons have either diverged sufficiently from their

ancestral allele counterparts, were lost due to structural rearrangements and mutations,

or encode *de novo* evolved candidate effectors. The candidate effector singletons were

found to be located more distantly from BUSCOs than allelic candidate effectors (Figure

6A), but were not more distant from other genes in general (Figure 6B). Nonetheless,

we reasoned that these candidate effector singletons might be more likely to be isolate-

or species-specific given their distinct genomic locations when compared to allelic

candidate effectors. We tested if candidate effector singletons are more likely to lack

orthologs in publicly available *Pst* genomes or other genomes of *Pucciniales* species

(82). Out of a total of 453 candidate effector singletons, 116 lacked an ortholog in five

other *Pst* genomes, compared to 118 out of 1,272 allelic candidate effectors. Singletons

are therefore more likely to be isolate-specific than are allelic candidate effectors

(Fischer's exact test, p ~ $1.36e^{-16}$). We observed a similar trend when comparing *Pst*-

104E with the six publically available *Pucciniales* genomes. Of 985 candidate effectors

lacking orthologs in other rust fungi, 313 were singletons and 672 allelic, also showing a

enrichment for candidate effector singletons (Fischer's exact test, p ~ $4.45e^{-26}$).

Conclusion

Using long-read sequencing technology we are now starting to uncover the genomic

diversity of dikaryotic fungi that was previously hidden by a reliance on short-read

sequence assemblies. We used this approach to generate a highly contiguous

haplotype-phased assembly of the Australian founder *Pst* pathotype*.* We are now able

to describe the levels of inter-haplome diversity both on the structural and gene levels. It

is difficult to fully evaluate the significance of observed levels of variations without

additional experiments and in the absence of similar studies. With over 6% variation, the

inter-haplome diversity of *Pst*-104E is higher than that reported for *P. coronata* f. sp.

avenae which ranges between 2.1 and 2.7% (77). It is also higher than the variation

observed between two isolates of *Z. tritici* (3D7 vs. MG2, 4.9%), an ascomycete

pathogen of wheat that undergoes frequent sexual cycles (58, 72), and two isolates of

V. dahliea (JR2 vs. VdLs17, 1.7%), a ascomycete pathogen of tomato that almost

exclusively propagates asexually (73). These comparisons suggest that the observed

inter-haplome diversity of *Pst* is relatively high. *Pst*-104E belongs to the 'North Western

European' (NW European) lineage of *Pst,* which has undergone long-term asexual

reproduction. The NW European *Pst* lineage can be traced back to its first sampling in

mid-1950 in the Netherlands, and has not shown any signs of sexual recombination

since (22, 83, 84). Consistently, the two *Pca* isolates that show much less inter-haplome

variation than *Pst*-104E are from populations that reproduce both sexually and

asexually on common buckthorn and oat, respectively (77). Frequent sexual

recombination is likely to reduce inter-haplome diversity and to purge mutations that are

deleterious in the monokaryon stage (85). On the other hand, long term clonal lineages

might accumulate polymorphisms that clear unwanted *Avr* genes but also contribute to

genomic decay. With respect to this, it is an interesting point if *Pst*-104E is still viable as

a monokaryon. The accumulation of large scale polymorphisms and potentially

deleterious mutations in each haplome of *Pst*-104E might have been buffered in the

dikaryon stage, but it is likely that it represents a terminal lineage of *Pst* in agreement

with the Muller's rachet hypothesis (85). Isolates from the NW European lineage show a

reduction in teliospore production on wheat, the entry point into the *Pst* sexual cycle,

when compared to isolates from the Himalayan region where sexual reproduction is

common (86). Also, successful sexual reproduction under laboratory conditions has

been reported only for *Pst* isolates that emerged recently from the center of diversity in

the Himalayan region (87), but not for isolates that have undergone long term clonal

reproduction such as the NW European lineage (personal communication J. Rodriguez-

Algaba). Furthermore, *Pst* populations of the NW European lineage have been

complete replaced by more recent *Pst* incursions in Europe and Australia (15, 26).

In future it will be important to generate high quality genomes for more *Pst* isolates,

including from sexual populations in the Himalayan regions (88). This will enable us to

investigate what role sexual vs. asexual reproduction play in terms of genome evolution

of a dikaryon in wild vs. agricultural settings. For now, the near-complete haplotype-

phased genome of *Pst*-104E provides a first insight into the genetic architecture of a

dikaryotic rust fungus pathogenic on wheat. In itself it is a high quality reference

genome enabling investigation of the rapid and devastating evolution of the fungus to

virulence during its asexual reproduction cycle in all wheat growing areas today.

Material and Methods

Data availability

The following are the NCBI accession numbers for data generated in course of this

manuscript which is registered as BioProject PRJNA396589.

Short read archive accession numbers are as follows:

SRX311905-14 and SRX311918-20: PacBio 10-20kb Blue Pippen kit, RSII, 13 SMRT

cells.

SRX311916 and 17: genomic DNA TruSeq library, HiSeq 2000, 100 bases paired end

library.

SRX311915: genomic DNA TruSeq PCR free, MiSeq, 250 bases paired end library.

SRX3191029-43: TruSeq v2 RNAseq samples , HiSeq 2000 100 bases paired end

library.

Bioinformatic scripts, supplemental files and genome annotation can be found on this

manuscripts github page https://github.com/BenjaminSchwessinger/Pst_104_E137_A-

_genome.

The genome will also be available at MycoCosm.

***Puccinia stiriformis* f. sp. *tritic*i pathotype, growth conditions and spore**

amplification

The isolate of pathotype 104E137A- was collected from the field in 1982 (Plant

Breeding Institute accession 821559=415), tested and propagated as described

previously (28). This pathotype is virulent on Heines VII (*Yr2,* Yr25), Vilmorin 23 (*Yr3*),

Hybrid 46 (*Yr4*), Stubes dickkopf, Nord Deprez, Suwon92/Omar and Avocet S (28). The

rust propagated for PacBio sequencing was produced by selecting a single pustule of

the original isolate (increase 0415Ga) on wheat plants of the susceptible variety

'Morocco'. The initial inoculation involved rubbing leaves of the susceptible host with a

spores from a sterile cotton tip. Plants were incubated under plastic in the dark at 9.5 ^{o}C

for 18 h before being transferred to a greenhouse microclimate set at 22 ^{o}C ±2 ^{o}C. After

6 d, plants were observed and all leaves were removed except for one leaf which

showed signs of infection by a single fleck indicating a rust pustule was soon to erupt

from the location. After pustule eruption, the single pustule selection was repeated to

ensure that the starter material for propagation was a single genotype. Multiplication of

rust occurred using *Triticum aestivum* 'Morocco'. For multiplication, 20 seeds of

'Morocco' were placed as a single layer into four inch pots filled with pasteurised soil

and watered with a half-strength solution of liquid fertiliser (Aquasol, Yates). At full

coleoptile emergence, each pot was treated with 50 mL maleic hydrazide solution (2mL

L^{-1} Slow Grow 270, Kendron). At full leaf emergence plants were inoculated by rubbing

with the pustules formed by the previous step and incubated as described previously.

Once four pots of 'Morocco' were heavily infected then rust was collected and

inoculated onto 64 four inch pots and a differential set to check pathotype identity and

purity. Rust was collected from the 64 pots using a GRA-101 large spore cyclone

(Tallgrass Solutions) attached to a domestic vacuum cleaner. Rust spores were dried

over silica gel for 7 days before being sieved with a 50 μm sieve and being stored at -80

^{o}C until DNA extraction.

DNA extraction and genome sequencing

DNA was extracted from dried dormant *Pst* urediniospores as we described it in great

detail elsewhere (89, 90). PacBio sequencing was performed at the Ramaciotti Centre

(Sydney, Australia). For library preparation the 20kb Blue Pippen kit (PacBio) was used.

DNA libraries were sequenced on a PacBio RSII instrument using P6-C4 chemistry. In

total we sequenced 13 SMRT cells (Table S1). DNA samples from the same *Pst*

pathotype were also sequenced with Illumina short read technology. We sequenced one

TruSeq library on a HiSeq 2000 instrument as a 100 bases paired end library at the

University of Western Sydney (Sydney, Australia). We sequenced one TruSeq PCR free

250 bases paired end library on an Illumina MiSeq instrument at the Ramaciotti Center

(Sydney, Australia).

Genome assembly and manual curation

For genome assembly we used FALCON-Unzip github tag 1.7.4 with the parameters

described in File S9 and S10 (31). We checked the resulting contigs for eukaryotic

contamination by blastn searches against the NCBI nucleotide reference database

(downloaded 04052016) (39). None of the contigs had predominant non-eukaryotic

sequences as best BLAST hits at a given position. We performed two manual curation

steps. In the first step we reasoned that some of the primary contigs without haplotigs

might actually represent haplotigs that could not be connected to their respective

primary contigs in the assembly graph because of too large a difference between the

two haplotypes. We aligned all primary contigs without haplotigs to primary contigs with

haplotigs using mummer version 3 (36). We screened the best alignments of each

primary contig without haplotig for percentage alignment, length of alignments, and if

they align to regions in the primary contigs that previously had not been covered by a

haplotig alignment. Using this approach we re-assigned 55 primary contigs without

haplotigs (~6Mb) to haplotigs (Table S8). In the second step of manual curation we

removed all contigs with an average coverage of greater 2000x when using Illumina

short read data. In total we removed 18 primary contigs (~0.6Mb) and 7 haplotigs

(~0.2Mb) of which most were mitochondrial contigs based on blastn analysis. The final

assembly contains 156 primary contigs (~83Mb) and 475 haplotigs (~73Mb) (Table S8).

Coverage analysis and identification of unphased regions in primary contigs

We aimed to assess the coverage within contigs and between contigs when mapping

Illumina short read data on primary contigs (p) and primary contigs and haplotigs (ph) at

the same time. We reasoned that unphased region of primary contigs should have

about twice the coverage of phased regions when mapping against ph and similar

coverage comparing mapping against p vs. ph. We trimmed Illumina short reads using

Trimmomatic v0.35 (91) (ILLUMINACLIP:adapter.fa:2:30:10 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:25 MINLEN:35) and assessed read quality with FastQC v0.11.4

(92). Reads were mapped against p or ph using BWA-MEM v0.7.15-r1142-dirty using

the standard parameters (93). The coverage for each position was calculated with

samtools v1.3.1 using depth with the '-aa' flag (94). Unphased regions on primary

contigs were defined as outlined above and converted to bed format. See jupyter

notebook Pst_104E_v12_coverage_analysis_ submission_21092017 in the github repo.

Repeat annotation

Repeat regions of the primary contigs and haplotigs were predicted independently. We

used the REPET pipeline v2.5 (37, 95) for repeat annotation in combination with

Repbase v21.05 (46). First, we used TEdenovo to predicted novel repetitive elements

following the instructions (96) using the parameters given in File S11. The set of TEs

provided by Tedenovo were used to annotate all repetitive elements using Teanno

following the instructions including the methodological advice (97) using the parameters

given in File S12. Annotation was performed on genome version 0.4 and subsequently

filtered for version 1.0 (Table S7). We transferred the superfamily annotation according

to Wicker (38) for all elements from the underlying database hits if these agreed with

each other and the REPET annotation. See jupyter notebooks

Pst_104E_v12_TE_filtering_and_summary_p_contigs_ submission_21092017 and

Pst_104E_v12_TE_filtering_and_summary_h_contigs_ submission_21092017 in the

github repo for full analysis details.

Gene model annotation

We annotated genes on primary contigs and haplotigs independently. We combined

RNAseq guided *ab initio* prediction using CodingQuarry v2.0 (42) and BRAKER v1.9

(43) with *de novo* transcriptome assembly approaches using Trinity v2.2.0 (98) and

PASA v2.0.1 (41). Gene models were unified using EvidenceModeler v1.1.1 (41) using

the weights given in File S13.

We mapped the trimmed RNAseq reads described in this study (see below) and

previously (40) against primary contigs and haplotigs using hisat2 v2.1.0 (--max-

intronlen 10000 --min-intronlen 20 --dta-cufflinks) (45). For *ab initio* predictions we

reconstructed transcripts using stringtie v1.2.3 (-f 0.2) (99). We ran CodingQuarry (-d) in

the pathogen mode using SignalP4 (100) for secretome prediction on the soft-masked

genome using RepeatMasker v4.0.5 (-xsmall -s -GC 43) . Similarly, we used the

stringtie reconstructed transcripts as training set for the *ab initio* prediction pipeline

BRAKER 1 v1.9 (43) and used the non-repeat masked genome as reference.

We used Trinity v2.2.0 to obtain *Pst* transcripts both in the *de novo* mode and the

genome guided mode (98). Several RNAseq samples contained host and pathogen

RNA as they were prepared from infected wheat tissue. We first mapped all reads to

primary contigs and haplotigs using hisat2 (see above). We extracted mapped RNAseq

reads using piccard tools SamToFastq (101). Only these reads mapping against *Pst*

contigs were used in the *de novo* pipeline of Trinity (--seqType fq). For genome guided

assembly we used bam files generated with hisat2 as starting point for Trinity (--

jacard_clip, --genome_gudied_max_intron 10000). We used the PASA pipeline v2.0.2

to align both sets of Trinity transcripts against *Pst* contigs with BLAT and GMAP using

the parameters given in File S14 (102).

The different gene models were combined using EvidenceModeler v.1.1.1 to get the

initial gene sets for primary contigs and haplotigs (41). These were filtered for homology

with proteins encoded in transposable elements. We used blastp to search for

homology in the Repbase v21.07 peptides database with an e-value cut-off of $1*e^{-10}$. In

addition, we used transposonPSI (103, 104). We used the outer union of both

approaches to remove genes coding for proteins associated with transposable elements

from our list of gene models.

Protein annotation

For initial protein annotation we used the fungal centric annotation pipeline funannotate

v0.3.10 (105). This included annotation for proteins with homology to swissprot

(uniref90, downloaded 22/09/2016) (50), to carbohydrate-active enzyme (dbCAN,

downloaded 22/09/2016) (49), to peptidases (MEROPS v10.0) (52, 106), for proteins

with eggnog terms (eggnog v4.5) (107) and SignalP4 (100). This annotation was

complemented by interproscan v5.21-60 (-iprlookup -goterms -pa) (48), eggnog-mapper

v0.99.2 (-m diamond and –d euk) (51), SignalP 3 (108), and EffectorP v1.01 (64, 109).

Biological material and molecular biology methods for *Pst* gene expression

analysis

We investigated gene expression of *Pst* at five different developmental stages or tissue

types. We extracted total RNA from dormant spores, germinated spores after 16 hours,

6 and 9 days post infection (dpi) of wheat and from haustoria isolated from wheat leaves

at 9 dpi.

In the case of dormant spores, spores were harvest from infected wheat at 14-18 dpi of

wheat, dried under vacuum for 1 hour and stored at -80^{o}C until use. For germination,

fresh spores were heat treated for 5 minutes at 42^{o}C and sprinkled on top of sterile Milli-

Q (MQ) water. The container was covered with Clingfilm and spores were incubated at

100% humidity at 10^{o}C in the dark for 16 hours before harvesting. For infection assays,

dormant spores were heat treated for 5 minutes at 42^{o}C, mixed with talcum powder (1:7

w/w) and sprayed homogenously with a manual air pump onto seven-day old wheat

seedlings wetted with water using a spray bottle. Plants were maintained in a container

at 100% humidity in the dark at 10^{o}C for 24 hours. At this point plants were transferred

to a constant temperature growth cabinet at 17^{o}C with a 16:8 light cycle. We collected

infected wheat leaf samples 6 and 9 dpi. Haustoria were purified from wheat leaves at 9

dpi (110). Infected wheat leaves (~20 g) were surface sterilised with 70% ethanol,

washed and blended in 250 ml of 1x isolation buffer (0.2 M sucrose, 20 mM MOPS pH

7.2, 1x IB). The homogenate was passed consecutively through 100 μm and 20 μm

meshes to remove cell debris. The filtrate was centrifuged at 1080 g for 15 min at 4^{o}C

and the resulting pellets resuspended in 80 mL 1x IB containing 30% Percol (v/v). The

suspension was centrifuged at 25,000 g for 30 min at 4^{o}C. The upper 10 mL of each

tube was recovered, diluted 10 times with 1x IB and centrifuged at 1080 g for 15 min at

4^{o}C. The pellets were resuspended in 20 mL of 1x IB containing 25% Percoll (v/v) and

centrifuged at 25,000 g for 30 min at 4^{o}C. The upper 10 mL of each tube was

recovered, diluted 10 times in 1x IB and centrifuged at 1080 g for 15 min at 4^{o}C. Pellets

were stained with ConA-488 to visualize haustoria under the fluorescence microscope.

The final pellets were frozen in liquid nitrogen and stored at -80^{o}C prior to RNA

isolation.

RNA for all samples was isolated as follows. Total RNA was isolated using the QIAGEN

Plant RNeasy kit following the manufacturer's instructions. Initial RNA quality and purity

checks were performed on a NanoDrop ND-1000 UV-Vis Spectrophotometer. Samples

were treated with DNase I (New England Biolabs) following the manufacturer's

instructions. Samples were purified using the QIAGEN Plant RNeasy kit following the

cleanup protocol, and RNA was eluted from columns in 50 µl of RNase-free water. The

concentration and integrity of all final RNA samples were verified on the Agilent 2100

bioanalyzer, using the RNA 6000 nano and pico kits. Three biological replicates were

processed.

RNA samples were sequenced at the Ramaciotti Centre (Sydney, Australia) on an

Illumina HiSeq 2000 instrument as 100-bp paired end reads. Approximately 10 µg of

total RNA per biological sample was processed with the TruSeq RNA Sample

Preparation Kit v2.

Differential expression analysis

We trimmed Illumina RNAseq reads using Trimmomatic v0.35 (91)

(ILLUMINACLIP:adapter.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:25

MINLEN:35) and assessed read quality with FastQC v0.11.4 (92). We mapped reads

using gene models as a guide using STAR v020201 (111). We first generated a

genome reference in the genomeGenerate mode using our gff for gene models (--

runMode genomeGenerate –sjdbGTFfile --sjdbGTFtagExonParentTranscript Parent).

We mapped our RNAseq reads against this reference using STAR in the alignReads

mode (--runMode alignReads

--readFilesCommand gunzip –c --outFilterType BySJout --outFilterMultimapNmax

20 --alignSJoverhangMin 8 --alignSJDBoverhangMin 1 --outFilterMismatchNmax 999 --

alignIntronMin 20 --alignIntronMax 10000 --alignMatesGapMax 1000000 --outSAMtype

BAM SortedByCoordinate --outSAMstrandField intronMotif --outFilterIntronMotifs

RemoveNoncanonical --quantMode GeneCounts). We used featureCounts v1.5.3 and

our gene annotation to quantify the overlaps of mapped reads with each gene model (-t

exon –g Parent) (112). We identified differentially expressed genes in either haustoria or

infected leaves relative to germinated spores (|log fold change| > 1.5 and an adjusted p-

value < 0.1) using the DESeq2 R package (113). *k*-means clustering was performed on

average rlog transformed values for each gene and condition. The optimal number of

clusters was defined using the elbow plot method and circular heatmaps drawn using

Circos (114). Scripts regarding the gene expression analysis can be found in the

gene_expression folder of the github repo.

We compared the expression pattern of alleles in different clusters (Table S6 and S7) in

jupyter notebook Pst_104E_v12_secretome_expression_cluster_analysis_

submission_21092017 in the github repo.

BUSCO analysis

We used BUSCO2 v2.0 4 beta to identify core conserved genes and to assess genome

completeness (33). In all cases we ran BUSCO2 in the protein mode using the

basidiomycota reference database downloaded 01/09/2016 (-l basidiomycota_odb9 -m

protein). We combined BUSCO identification on primary contigs and haplotigs non-

redundantly to asses completeness of the combined assembly. For detail see jupyter

notebook Pst_104E_v12_BUSCO_summary_submission_21092017 in the github repo.

Inter-haplome variation analysis

We mapped trimmed reads against primary contigs using BWA-MEM v0.7.15-r1142-

dirty using the standard parameters (93). We called SNPs with freebayes default

parameters (115) and filtered the output with vcffilter v1.0.0-rc1 (-f "DP >10" -f "QUAL >

20") (116). SNP calls were summarized with real time genomic vcfstats v3.8.4 (117).

We aligned all haplotigs to their corresponding primary contigs using nucmer of the

mummer package (-maxmatch -l 100 -c 500) (36). We fed these alignments into

Assemblytics to estimate the inter-haplom variation for each primary contig – haplotig

pairing (35). For this analysis, we used a unique anchor length of 8kb based on the

length of identified TEs in our *Pst* assembly and a maximum feature length of 10kb. For

consistency, we used nucmer alignments filtered by Assemblytics for the allele status

analysis (see below). Analysis and summary of variations is shown in jupyter notebook

Pst_104E_v12_assemblytics_analysis_ submission_21092017 and

Pst_104E_v12_nucmer_and_assemblytics submission_21092017 in the github repo.

Allele status analysis

We used proteinortho v5.16 in the synteny mode with default parameters (-synteny) to identify alleles between the primary assembly and haplotigs (53). We parsed the results and defined three major allele status categories as follows. Allele pairs were parsed from the 'poff-graph' output file. Non-allelic inter-haplome paralogues were parsed from the 'proteinortho' output file and checked for absence in the 'poff-graph' output file. Potential singletons were defined as gene models being absent from both of these two output files. Alleles were further subdivided into alleles for which the primary and associated haplotig gene models were located on contigs that aligned with each other at the position of the primary gene model (Figure S2A), alleles for which the primary and associated haplotig gene models were located on contigs that did not align with each other at the position of the primary gene model (Figure S2B), and alleles for which the allele of a primary gene model was not located on a haplotig associated with the respective primary contig (Figure S2C). Potential singletons were screened for being located in regions of the primary assembly that were unphased based on Illumina coverage analysis (see above). Genes located in these regions were defined as unphased and removed from the initial list. All other gene models constitute haplome singletons. Analysis details can be found in the jupyter notebooks Pst_104E_v12_defining_alleles_ submission_21092017 and Pst_104E_v12_missing_allele_QC_ submission_21092017.

Allele variation analysis

We assessed the variation of allele pairs using three approaches. We calculated the

Levenshtein distance (80) on the CDS alignments of two alleles, on the codon based

protein alignments and we calculated the dN/dS ratios using these two alignments sets

with yn00 paml version 4.9 (81). The CDS of two alleles were aligned using muscle

v3.8.31 (118) and codon based alignments were generated using PAL2NAL v14 (119).

The Levenshtein distance was calculated in python using the distance module v 0.1.3

(120). Analysis details can be found in the jupyter notebook

Pst_104E_v12_post_allele_analysis_ submission_21092017.

Genome architecture analysis

We used bedtools v2.25.0 (121) and the python module pybedtools (122) to perform

various genome analysis tasks. This included the calculation of nearest neighbours

using the closest function. Details of the analysis can be found in the jupyter notebooks

Pst_104E_v12_post_allele_analysis submission_21092017 and

Pst_104E_v12_effectors submission_21092017.

Orthology analysis of candidate effector analysis

We performed orthology analysis with proteinortho v5.16 (-singles) (53) of all non-

redundant candidate effectors with publicly available *Pst* genomes. *Pst*-130 (4) and

Pst-78 (30) protein sets were downloaded from MycoCosm (05/09/2017) (82). *Pst*-0821,

Pst-21, *Pst*-43 and *Pst*-887 were downloaded from yellowrust.com (30/03/2017) (5,

123). We performed a similar analysis searching for candidate effector orthologs in

Pucciniales excluding *Pst* genomes. *Puccinia triticina* 1-1 BBBD Race 1 (30), *Puccinia*

graminis f. sp. *tritici* v2.0 (6), *Puccinia coronata-avenae* 12SD80 and 12NC29 (124), and

Melampsora lini CH5 (125) genomes were downloaded from MycoCosm (05/09/2017).

The *Puccinia sorghi* genome (126) (ASM126337v1) was downloaded from NCBI

(05/09/2017).

Data and statistical analysis

We used the python programming language (127) in the jupyter notebook environment

for data analysis (128). In particular, we used pandas (129), numpy (130), matplotlib

(131) and seaborn (132) for data crunching and plotting. Statistical analysis was

performed using the Scipy (130) and statsmodel (133) toolkits.

Acknowledgment

We thank the following colleagues for technical advice; Ying Zhang, Sylvain Forêt,

Marcin Adamski, Adam Taranto, and Megan McDonald. We thank Ashlea Grewar for

technical assistance with rust multiplication. We thank the following colleagues for

feedback on the manuscript; Adam Taranto, Megan McDonald, Sajid Ali, Annemarie

Fejer Justesen, and Sambasivam Periyannan.

The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE

Office of Science User Facility, is supported by the Office of Science of the U.S.

Department of Energy under Contract No. DE-AC02-05CH11231.

Funding information

BS was supported by a Human Frontiers Science Program long-term postdoctoral

fellowship (LT000674/2012) and a Discovery Early Career Research Award

(DE150101897). BS and JPR were supported by a sequencing grant from Bioplatforms

Australia. JS is supported by a CSIRO OCE Postdoctoral Fellowship. We thank Ashlea

Grewar for technical assistance with rust multiplication. RFP acknowledges the

generous support of Judith and David Coffey and family, and of the Australian Grains

Research and Development Corporation. MF is supported by the University of

Minnesota Experimental Station USDA-NIFA Hatch/Figueroa project MIN-22-058,

M.E.M is supported by a USDA-NIFA Postdoctoral Fellowship Award (2017-67012-

26117).

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Authors' contributions

BS amplified rust spores, extracted high molecular weight DNA, performed assembly

using FALCON-Unzip, performed manual curation of assembly, annotated the genome

and proteome, performed all additional bioinformatic analysis except for differential

expression analysis of RNA-seq data, conceived study, and wrote the manuscript. JS

performed differential expression analysis, contributed ideas to data analysis and wrote

manuscript. WC performed pathotyping of the rust strain, amplified rust spores and

commented on the manuscript. DG performed infection assays, RNAseq assay,

extracted haustoria for RNAseq assay, and commented on the manuscript. MEM

contributed ideas to data analysis and commented on the manuscript. JMT and PND

contributed ideas to the differential expression analysis and commented on the

manuscript. MF contributed ideas to data analysis and wrote the manuscript. RFP

provided *Pst* urediniospores and commented on manuscript. JPR contributed ideas to

methodological development, data analysis and commented on the manuscript.

References

1.   2017. Stop neglecting fungi. Nat Microbiol 2:nmicrobiol2017120.

2.   Spatafora JW, Aime MC, Grigoriev IV, Martin F, Stajich JE, Blackwell M. 2017.

The Fungal Tree of Life: from Molecular Systematics to Genome-Scale

Phylogenies. Microbiol Spectr 5.

3.   Kämper J, Kahmann R, Bölker M, Ma L-J, Brefort T, Saville BJ, Banuett F,

Kronstad JW, Gold SE, Müller O, Perlin MH, Wösten HAB, de Vries R, Ruiz-

Herrera J, Reynaga-Peña CG, Snetselaar K, McCann M, Pérez-Martín J,

Feldbrügge M, Basse CW, Steinberg G, Ibeas JI, Holloman W, Guzman P,

Farman M, Stajich JE, Sentandreu R, González-Prieto JM, Kennell JC, Molina L,

Schirawski J, Mendoza-Mendoza A, Greilinger D, Münch K, Rössel N, Scherer M,

Vraneš M, Ladendorf O, Vincon V, Fuchs U, Sandrock B, Meng S, Ho ECH, Cahill

MJ, Boyce KJ, Klose J, Klosterman SJ, Deelstra HJ, Ortiz-Castellanos L, Li W,

Sanchez-Alonso P, Schreier PH, Häuser-Hahn I, Vaupel M, Koopmann E,

Friedrich G, Voss H, Schlüter T, Margolis J, Platt D, Swimmer C, Gnirke A, Chen

F, Vysotskaia V, Mannhaupt G, Güldener U, Münsterkötter M, Haase D,

Oesterheld M, Mewes H-W, Mauceli EW, DeCaprio D, Wade CM, Butler J, Young

S, Jaffe DB, Calvo S, Nusbaum C, Galagan J, Birren BW. 2006. Insights from the

genome of the biotrophic fungal plant pathogen Ustilago maydis. Nature 444:97–

101.

4.   Cantu D, Govindarajulu M, Kozik A, Wang M, Chen X, Kojima KK, Jurka J,

Michelmore RW, Dubcovsky J. 2011. Next generation sequencing provides rapid

access to the genome of Puccinia striiformis f. sp. tritici, the causal agent of wheat

stripe rust. PloS One 6:e24230.

5.   Cantu D, Segovia V, MacLean D, Bayles R, Chen X, Kamoun S, Dubcovsky J,

Saunders DG, Uauy C. 2013. Genome analyses of the wheat yellow (stripe) rust

pathogen Puccinia striiformis f. sp. tritici reveal polymorphic and haustorial

expressed secreted proteins as candidate effectors. BMC Genomics 14:270.

6.   Duplessis S, Cuomo CA, Lin Y-C, Aerts A, Tisserant E, Veneault-Fourrey C, Joly

DL, Hacquard S, Amselem J, Cantarel BL, Chiu R, Coutinho PM, Feau N, Field M,

Frey P, Gelhaye E, Goldberg J, Grabherr MG, Kodira CD, Kohler A, Kües U,

Lindquist EA, Lucas SM, Mago R, Mauceli E, Morin E, Murat C, Pangilinan JL,

Park R, Pearson M, Quesneville H, Rouhier N, Sakthikumar S, Salamov AA,

Schmutz J, Selles B, Shapiro H, Tanguay P, Tuskan GA, Henrissat B, Peer YV

de, Rouzé P, Ellis JG, Dodds PN, Schein JE, Zhong S, Hamelin RC, Grigoriev IV,

Szabo LJ, Martin F. 2011. Obligate biotrophy features unraveled by the genomic

analysis of rust fungi. Proc Natl Acad Sci 108:9166–9171.

7.   Zheng W, Huang L, Huang J, Wang X, Chen X, Zhao J, Guo J, Zhuang H, Qiu C,

Liu J, Liu H, Huang X, Pei G, Zhan G, Tang C, Cheng Y, Liu M, Zhang J, Zhao Z,

Zhang S, Han Q, Han D, Zhang H, Zhao J, Gao X, Wang J, Ni P, Dong W, Yang

L, Yang H, Xu J-R, Zhang G, Kang Z. 2013. High genome heterozygosity and

endemic genetic recombination in the wheat stripe rust fungus. Nat Commun

4:2673.

8.   Goellner K, Loehrer M, Langenbach C, Conrath U, Koch E, Schaffrath U. 2010.

Phakopsora pachyrhizi, the causal agent of Asian soybean rust. Mol Plant Pathol

11:169–177.

9.   Nazareno ES, Li F, Smith M, Park RF, Kianian SF, Figueroa M. Puccinia coronata

f. sp. avenae: a threat to global oat production. Mol Plant Pathol n/a-n/a.

10. Talhinhas P, Batista D, Diniz I, Vieira A, Silva DN, Loureiro A, Tavares S, Pereira

AP, Azinheira HG, Guerra-Guimarães L, Várzea V, Silva M do C. 2017. The

coffee leaf rust pathogen Hemileia vastatrix: one and a half centuries around the

tropics. Mol Plant Pathol 18:1039–1051.

11. Park RF, Golegaonkar PG, Derevnina L, Sandhu KS, Karaoglu H, Elmansour HM,

Dracatos PM, Singh D. 2015. Leaf Rust of Cultivated Barley: Pathology and

Control. Annu Rev Phytopathol.

12. Hovmøller MS, Sørensen CK, Walter S, Justesen AF. 2011. Diversity of Puccinia

striiformis on Cereals and Grasses. Annu Rev Phytopathol 49:197–217.

13. Schwessinger B. 2016. Fundamental wheat stripe rust research in the 21st

century. New Phytol n/a-n/a.

14.   Bread Wheat - Improvement and Production.

15. Wellings CR. 2011. Global status of stripe rust: a review of historical and current

threats. Euphytica 179:129–141.

16. Beddow JM, Pardey PG, Chai Y, Hurley TM, Kriticos DJ, Braun H-J, Park RF,

Cuddy WS, Yonow T. 2015. Research investment implications of shifts in the

global geography of wheat stripe rust. Nat Plants 15132.

17. Chen W, Wellings C, Chen X, Kang Z, Liu T. 2014. Wheat stripe (yellow) rust

caused by Puccinia striiformis f. sp. tritici. Mol Plant Pathol 15:433–446.

18. Schwessinger B. 2017. Fundamental wheat stripe rust research in the 21(st)

century. New Phytol 213:1625–1631.

19. Zhao J, Wang L, Wang Z, Chen X, Zhang H, Yao J, Zhan G, Chen W, Huang L,

Kang Z. 2013. Identification of Eighteen Berberis Species as Alternate Hosts of

Puccinia striiformis f. sp. tritici and Virulence Variation in the Pathogen Isolates

from Natural Infection of Barberry Plants in China. Phytopathology 103:927–934.

20. Zhao J, Wang M, Chen X, Kang Z. 2016. Role of Alternate Hosts in Epidemiology

and Pathogen Variation of Cereal Rusts. Annu Rev Phytopathol 54:207–228.

981 21. Jin Y, Szabo LJ, Carson M. 2010. Century-Old Mystery of Puccinia striiformis Life

982   History Solved with the Identification of Berberis as an Alternate Host.

983   Phytopathology 100:432–435.

984 22. Ali S, Gladieux P, Leconte M, Gautier A, Justesen AF, Hovmøller MS, Enjalbert J,

985   de Vallavieille-Pope C. 2014. Origin, Migration Routes and Worldwide Population

986   Genetic Structure of the Wheat Yellow Rust Pathogen Puccinia striiformis f.sp.

987   tritici. PLoS Pathog 10:e1003903.

988 23. Periyannan S, Milne RJ, Figueroa M, Lagudah ES, Dodds PN. 2017. An overview

989   of genetic rust resistance: From broad to specific mechanisms. PLOS Pathog

990   13:e1006380.

991 24. Ellis JG, Lagudah ES, Spielmeyer W, Dodds PN. 2014. The past, present and

992   future of breeding rust resistant wheat. Plant-Microbe Interact 5:641.

993 25. Park RF. 2008. Breeding cereals for rust resistance in Australia. Plant Pathol

994   57:591–602.

995 26. Hovmøller MS, Walter S, Bayles RA, Hubbard A, Flath K, Sommerfeldt N, Leconte

996   M, Czembor P, Rodriguez-Algaba J, Thach T, Hansen JG, Lassen P, Justesen

997   AF, Ali S, de Vallavieille-Pope C. 2016. Replacement of the European wheat

998   yellow rust population by new races from the centre of diversity in the near-

999   Himalayan region. Plant Pathol 65:402–411.

1000 27. Hubbard A, Lewis CM, Yoshida K, Ramirez-Gonzalez RH, Vallavieille-Pope C de,

1001   Thomas J, Kamoun S, Bayles R, Uauy C, Saunders DG. 2015. Field

pathogenomics reveals the emergence of a diverse wheat yellow rust population.

Genome Biol 16:23.

28.   Wellings CR. 2007. Puccinia striiformis in Australia: a review of the incursion,

evolution, and adaptation of stripe rust in the period 1979–2006. Aust J Agric Res

58:567–575.

29.   Dean R, Van Kan J a. L, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu

PD, Rudd JJ, Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The Top 10

fungal pathogens in molecular plant pathology. Mol Plant Pathol 13:414–430.

30.   Cuomo CA, Bakkeren G, Khalil HB, Panwar V, Joly D, Linning R, Sakthikumar S,

Song X, Adiconis X, Fan L, Goldberg JM, Levin JZ, Young S, Zeng Q, Anikster Y,

Bruce M, Wang M, Yin C, McCallum B, Szabo LJ, Hulbert S, Chen X, Fellers JP.

2016. Comparative Analysis Highlights Variable Genome Content of Wheat Rusts

and Divergence of the Mating Loci. G3 Genes Genomes Genet g3.116.032797.

31.   Chin C-S, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, Dunn

C, O'Malley R, Figueroa-Balderas R, Morales-Cruz A, Cramer GR, Delledonne M,

Luo C, Ecker JR, Cantu D, Rank DR, Schatz MC. 2016. Phased diploid genome

assembly with single-molecule real-time sequencing. Nat Methods 13:1050–1054.

32.   Wang M, Beck CR, English AC, Meng Q, Buhay C, Han Y, Doddapaneni HV, Yu

F, Boerwinkle E, Lupski JR, Muzny DM, Gibbs RA. 2015. PacBio-LITS: a large-

insert targeted sequencing method for characterization of human disease-

associated chromosomal structural variations. BMC Genomics 16:214.

33.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015.

1024   BUSCO: assessing genome assembly and annotation completeness with single-

1025   copy orthologs. Bioinformatics 31:3210–3212.

1026   34.   Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J,

1027   Schatz MC. GenomeScope: fast reference-free genome profiling from short reads.

1028   Bioinformatics.

1029   35.   Nattestad M, Schatz MC. 2016. Assemblytics: a web analytics tool for the

1030   detection of variants from an assembly. Bioinformatics 32:3021–3023.

1031   36.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg

1032   SL. 2004. Versatile and open software for comparing large genomes. Genome

1033   Biol 5:R12.

1034   37.   Quesneville H, Bergman CM, Andrieu O, Autard D, Nouaud D, Ashburner M,

1035   Anxolabehere D. 2005. Combined Evidence Annotation of Transposable

1036   Elements in Genome Sequences. PLOS Comput Biol 1:e22.

1037   38.   Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A,

1038   Leroy P, Morgante M, Panaud O, Paux E, SanMiguel P, Schulman AH. 2007. A

1039   unified classification system for eukaryotic transposable elements. Nat Rev Genet

1040   8:973–982.

1041   39.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

1042   search tool. J Mol Biol 215:403–410.

1043   40.   Dobon A, Bunting DCE, Cabrera-Quio LE, Uauy C, Saunders DGO. 2016. The

1044   host-pathogen interaction between wheat and yellow rust induces temporally

1045   coordinated waves of gene expression. BMC Genomics 17:380.

1046   41.   Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR,

1047   Wortman JR. 2008. Automated eukaryotic gene structure annotation using

1048   EVidenceModeler and the Program to Assemble Spliced Alignments. Genome

1049   Biol 9:R7.

1050   42.   Testa AC, Hane JK, Ellwood SR, Oliver RP. 2015. CodingQuarry: highly accurate

1051   hidden Markov model gene prediction in fungal genomes using RNA-seq

1052   transcripts. BMC Genomics 16:170.

1053   43.   Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1:

1054   Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and

1055   AUGUSTUS. Bioinformatics 32:767–769.

1056   44.   Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL.

1057   2015. StringTie enables improved reconstruction of a transcriptome from RNA-

1058   seq reads. Nat Biotechnol 33:290–295.

1059   45.   Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low

1060   memory requirements. Nat Methods 12:357–360.

1061   46.   Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive

1062   elements in eukaryotic genomes. Mob DNA 6:11.

1063   47.   Haas BJ, Zeng Q, Pearson MD, Cuomo CA, Wortman JR. 2011. Approaches to

1064   Fungal Genome Annotation. Mycology 2:118–141.

1065   48.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen

1066   J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew

1067   M, Yong S-Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein

1068   function classification. Bioinforma Oxf Engl 30:1236–1240.

1069   49.   Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. dbCAN: a web resource for

1070   automated carbohydrate-active enzyme annotation. Nucleic Acids Res 40:W445-

1071   451.

1072   50.   Bairoch A, Apweiler R. 2000. The SWISS-PROT protein sequence database and

1073   its supplement TrEMBL in 2000. Nucleic Acids Res 28:45–48.

1074   51.   Huerta-Cepas J, Forslund K, Pedro Coelho L, Szklarczyk D, Juhl Jensen L, von

1075   Mering C, Bork P. Fast genome-wide functional annotation through orthology

1076   assignment by eggNOG-mapper. Mol Biol Evol.

1077   52.   Rawlings ND, Barrett AJ, Finn R. 2016. Twenty years of the MEROPS database

1078   of proteolytic enzymes, their substrates and inhibitors. Nucleic Acids Res

1079   44:D343–D350.

1080   53.   Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. 2011.

1081   Proteinortho: Detection of (Co-)orthologs in large-scale analysis. BMC

1082   Bioinformatics 12:124.

1083   54.   Veltri D, Wight MM, Crouch JA. 2016. SimpleSynteny: a web-based tool for

1084   visualization of microsynteny across multiple species. Nucleic Acids Res 44:W41-

1085   45.

1086   55.   Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison

1087   visualizer. Bioinforma Oxf Engl 27:1009–1010.

1088   56.   2016. Database resources of the National Center for Biotechnology Information.

1089   Nucleic Acids Res 44:D7–D19.

1090   57.   Kersey PJ, Allen JE, Armean I, Boddu S, Bolt BJ, Carvalho-Silva D, Christensen

1091   M, Davis P, Falin LJ, Grabmueller C, Humphrey J, Kerhornou A, Khobova J,

1092   Aranganathan NK, Langridge N, Lowy E, McDowall MD, Maheswari U, Nuhn M,

1093   Ong CK, Overduin B, Paulini M, Pedro H, Perry E, Spudich G, Tapanari E, Walts

1094   B, Williams G, Tello–Ruiz M, Stein J, Wei S, Ware D, Bolser DM, Howe KL,

1095   Kulesha E, Lawson D, Maslen G, Staines DM. 2016. Ensembl Genomes 2016:

1096   more genomes, more complexity. Nucleic Acids Res 44:D574–D580.

1097   58.   Plissonneau C, Stürchler A, Croll D. 2016. The Evolution of Orphan Regions in

1098   Genomes of a Fungal Pathogen of Wheat. mBio 7:e01231-16.

1099   59.   Sperschneider J, Dodds PN, Gardiner DM, Manners JM, Singh KB, Taylor JM.

1100   2015. Advances and Challenges in Computational Prediction of Effectors from

1101   Plant Pathogenic Fungi. PLoS Pathog 11:e1004806.

1102   60.   Petre B, Joly DL, Duplessis S. 2014. Effector proteins of rust fungi. Front Plant Sci

1103   5.

1104   61.   Anderson C, Khan MA, Catanzariti A-M, Jack CA, Nemri A, Lawrence GJ,

1105   Upadhyaya NM, Hardham AR, Ellis JG, Dodds PN, Jones DA. 2016. Genome

1106   analysis and avirulence gene cloning using a high-density RADseq linkage map of

1107   the flax rust fungus, Melampsora lini. BMC Genomics 17:667.

1108   62.   Dagvadorj B, Ozketen AC, Andac A, Duggan C, Bozkurt TO, Akkaya MS. 2017. A

1109   Puccinia striiformis f. sp. tritici secreted protein activates plant immunity at the cell

1110   surface. Sci Rep 7:1141.

1111   63.   Liu C, Pedersen C, Schultz-Larsen T, Aguilar GB, Madriz-Ordeñana K, Hovmøller

1112   MS, Thordal-Christensen H. 2016. The stripe rust fungal effector PEC6

1113   suppresses pattern-triggered immunity in a host species-independent manner and

1114   interacts with adenosine kinases. New Phytol n/a-n/a.

1115   64.   Sperschneider J, Gardiner DM, Dodds PN, Tini F, Covarelli L, Singh KB, Manners

1116   JM, Taylor JM. 2015. EffectorP: predicting fungal effector proteins from

1117   secretomes using machine learning. New Phytol n/a-n/a.

1118   65.   Sperschneider J, Catanzariti A-M, DeBoer K, Petre B, Gardiner DM, Singh KB,

1119   Dodds PN, Taylor JM. 2017. LOCALIZER: subcellular localization prediction of

1120   both plant and effector proteins in the plant cell. Sci Rep 7:srep44598.

1121   66.   van Damme M, Bozkurt TO, Cakir C, Schornack S, Sklenar J, Jones AME,

1122   Kamoun S. 2012. The Irish Potato Famine Pathogen Phytophthora infestans

1123   Translocates the CRN8 Kinase into Host Plant Cells. PLoS Pathog 8:e1002875.

67. Ramirez-Garcés D, Camborde L, Pel MJC, Jauneau A, Martinez Y, Néant I,

Leclerc C, Moreau M, Dumas B, Gaulin E. 2016. CRN13 candidate effectors from

plant and animal eukaryotic pathogens are DNA-binding proteins which trigger

host DNA damage response. New Phytol 210:602–617.

68. Dong S, Raffaele S, Kamoun S. 2015. The two-speed genomes of filamentous

pathogens: waltz with plants. Curr Opin Genet Dev 35:57–65.

69. Faino L, Seidl MF, Shi-Kunne X, Pauper M, Berg GCM van den, Wittenberg AHJ,

Thomma BPHJ. 2016. Transposons passively and actively contribute to evolution

of the two-speed genome of a fungal pathogen. Genome Res 26:1091–1100.

70. Shi X, Faino L, Berg G van den, Thomma B, Seidl M. 2017. Evolution within the

fungal genus Verticillium is characterized by chromosomal rearrangement and

gene loss. bioRxiv 164665.

71. Ma L-J, van der Does HC, Borkovich KA, Coleman JJ, Daboussi M-J, Di Pietro A,

Dufresne M, Freitag M, Grabherr M, Henrissat B, Houterman PM, Kang S, Shim

W-B, Woloshuk C, Xie X, Xu J-R, Antoniw J, Baker SE, Bluhm BH, Breakspear A,

Brown DW, Butchko RAE, Chapman S, Coulson R, Coutinho PM, Danchin EGJ,

Diener A, Gale LR, Gardiner DM, Goff S, Hammond-Kosack KE, Hilburn K, Hua-

Van A, Jonkers W, Kazan K, Kodira CD, Koehrsen M, Kumar L, Lee Y-H, Li L,

Manners JM, Miranda-Saavedra D, Mukherjee M, Park G, Park J, Park S-Y,

Proctor RH, Regev A, Ruiz-Roldan MC, Sain D, Sakthikumar S, Sykes S,

Schwartz DC, Turgeon BG, Wapinski I, Yoder O, Young S, Zeng Q, Zhou S,

Galagan J, Cuomo CA, Kistler HC, Rep M. 2010. Comparative genomics reveals

mobile pathogenicity chromosomes in Fusarium. Nature 464:367–373.

72.   Goodwin SB, M'Barek SB, Dhillon B, Wittenberg AHJ, Crane CF, Hane JK, Foster

AJ, Lee TAJV der, Grimwood J, Aerts A, Antoniw J, Bailey A, Bluhm B, Bowler J,

Bristow J, Burgt A van der, Canto-Canché B, Churchill ACL, Conde-Ferràez L,

Cools HJ, Coutinho PM, Csukai M, Dehal P, Wit PD, Donzelli B, Geest HC van

de, Ham RCHJ van, Hammond-Kosack KE, Henrissat B, Kilian A, Kobayashi AK,

Koopmann E, Kourmpetis Y, Kuzniar A, Lindquist E, Lombard V, Maliepaard C,

Martins N, Mehrabi R, Nap JPH, Ponomarenko A, Rudd JJ, Salamov A, Schmutz

J, Schouten HJ, Shapiro H, Stergiopoulos I, Torriani SFF, Tu H, Vries RP de,

Waalwijk C, Ware SB, Wiebenga A, Zwiers L-H, Oliver RP, Grigoriev IV, Kema

GHJ. 2011. Finished Genome of the Fungal Wheat Pathogen Mycosphaerella

graminicola Reveals Dispensome Structure, Chromosome Plasticity, and Stealth

Pathogenesis. PLOS Genet 7:e1002070.

73.   Faino L, Seidl MF, Datema E, Berg GCM van den, Janssen A, Wittenberg AHJ,

Thomma BPHJ. 2015. Single-Molecule Real-Time Sequencing Combined with

Optical Mapping Yields Completely Finished Fungal Genome. mBio 6:e00936-15.

74.   Möller M, Stukenbrock EH. 2017. Evolution and genome architecture in fungal

plant pathogens. Nat Rev Microbiol.

75.   Tuskan G, DiFazio S, Jansson S, Bohlmann J, Grigoriev I, Hellsten U, Putnam N,

Ralph S, Rombauts S, Salamov A, Schein J, Sterck L, Aerts A, Bhalerao R,

Bhalerao R, Blaudez D, Boerjan W, Brun A, Brunner A, Busov V, Campbell M,

Carlson J, Chalot M, Chapman J, Chen G, Cooper D, Coutinho P, Couturier J,

Covert S, Cronk Q. 2006. The genome of black cottonwood, Populus trichocarpa

(Torr. & Gray). Science 313:1596–1604.

76.   Saunders DGO, Win J, Cano LM, Szabo LJ, Kamoun S, Raffaele S. 2012. Using

Hierarchical Clustering of Secreted Protein Families to Classify and Rank

Candidate Effectors of Rust Fungi. PLoS ONE 7:e29847.

77.   Miller ME, Zhang Y, Omidvar V, Sperschneider J, Schwessinger B, Raley C,

Palmer JM, Garnica D, Upadhyaya N, Rathjen J, Taylor JM, Park RF, Dodds PN,

Hirsch CD, Kianian SF, Figueroa M. 2017. De novo assembly and phasing of

dikaryotic genomes from two isolates of Puccinia coronata f. sp. avenae, the

causal agent of oat crown rust. bioRxiv 179226.

78.   Dutheil JY, Mannhaupt G, Schweizer G, M. K. Sieber C, Münsterkötter M,

Güldener U, Schirawski J, Kahmann R. 2016. A Tale of Genome

Compartmentalization: The Evolution of Virulence Clusters in Smut Fungi.

Genome Biol Evol 8:681–704.

79.   Haas BJ, Kamoun S, Zody MC, Jiang RH, Handsaker RE, Cano LM, Grabherr M,

Kodira CD, Raffaele S, Torto-Alalibo T, Bozkurt TO, Ah-Fong AM, Alvarado L,

Anderson VL, Armstrong MR, Avrova A, Baxter L, Beynon J, Boevink PC,

Bollmann SR, Bos JI, Bulone V, Cai G, Cakir C, Carrington JC, Chawner M, Conti

L, Costanzo S, Ewan R, Fahlgren N, Fischbach MA, Fugelstad J, Gilroy EM,

Gnerre S, Green PJ, Grenville-Briggs LJ, Griffith J, Grunwald NJ, Horn K, Horner

NR, Hu CH, Huitema E, Jeong DH, Jones AM, Jones JD, Jones RW, Karlsson

EK, Kunjeti SG, Lamour K, Liu Z, Ma L, Maclean D, Chibucos MC, McDonald H,

McWalters J, Meijer HJ, Morgan W, Morris PF, Munro CA, O'Neill K, Ospina-

Giraldo M, Pinzon A, Pritchard L, Ramsahoye B, Ren Q, Restrepo S, Roy S,

Sadanandom A, Savidor A, Schornack S, Schwartz DC, Schumann UD,

Schwessinger B, Seyer L, Sharpe T, Silvar C, Song J, Studholme DJ, Sykes S,

Thines M, van de Vondervoort PJ, Phuntumart V, Wawra S, Weide R, Win J,

Young C, Zhou S, Fry W, Meyers BC, van West P, Ristaino J, Govers F, Birch

PR, Whisson SC, Judelson HS, Nusbaum C. 2009. Genome sequence and

analysis of the Irish potato famine pathogen Phytophthora infestans. Nature

461:393–8.

80.   Levenshtein VI. 1966. Binary Codes Capable of Correcting Deletions, Insertions

and Reversals. Sov Phys Dokl 10:707.

81.   Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous

substitution rates under realistic evolutionary models. Mol Biol Evol 17:32–43.

82.   Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A,

Zhao X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014.

MycoCosm portal: gearing up for 1000 fungal genomes. Nucleic Acids Res

42:D699-704.

83.   Ali S, Rodriguez-Algaba J, Thach T, Sørensen CK, Hansen JG, Lassen P, Nazari

K, Hodson DP, Justesen AF, Hovmøller MS. 2017. Yellow Rust Epidemics

Worldwide Were Caused by Pathogen Races from Divergent Genetic Lineages.

Front Plant Sci 8.

84.   Thach T, Ali S, de Vallavieille-Pope C, Justesen AF, Hovmøller MS. 2016.

Worldwide population structure of the wheat rust fungus Puccinia striiformis in the

past. Fungal Genet Biol 87:1–8.

85.   Muller HJ. 1964. THE RELATION OF RECOMBINATION TO MUTATIONAL

ADVANCE. Mutat Res 106:2–9.

86.   Ali S, Leconte M, Walker A-S, Enjalbert J, de Vallavieille-Pope C. 2010. Reduction

in the sex ability of worldwide clonal populations of Puccinia striiformis f.sp. tritici.

Fungal Genet Biol 47:828–838.

87.   Rodriguez-Algaba J, Walter S, Sørensen CK, Hovmøller MS, Justesen AF. 2014.

Sexual structures and recombination of the wheat rust fungus Puccinia striiformis

on Berberis vulgaris. Fungal Genet Biol 70:77–85.

88.   Ali S, Gladieux P, Rahman H, Saqib MS, Fiaz M, Ahmad H, Leconte M, Gautier A,

Justesen AF, Hovmøller MS, Enjalbert J, de Vallavieille-Pope C. 2014. Inferring

the contribution of sexual reproduction, migration and off-season survival to the

temporal maintenance of microbial populations: a case study on the wheat fungal

pathogen Puccinia striiformis f.sp. tritici. Mol Ecol 23:603–617.

89.   Schwessinger B, Rathjen JP. 2017. Extraction of High Molecular Weight DNA

from Fungal Rust Spores for Long Read Sequencing. Methods Mol Biol Clifton NJ

1659:49–57.

90.   High quality DNA from Fungi for long read sequencing e.g. PacBio.

91.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
1232   sequence data. Bioinforma Oxf Engl 30:2114–2120.

1233   92.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput
1234   Sequence Data.

1235   93.   Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with
1236   BWA-MEM. ArXiv13033997 Q-Bio.

1237   94.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis
1238   G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The
1239   Sequence Alignment/Map format and SAMtools. Bioinforma Oxf Engl 25:2078–
1240   2079.

1241   95.   Flutre T, Duprat E, Feuillet C, Quesneville H. 2011. Considering Transposable
1242   Element Diversification in De Novo Annotation Approaches. PLOS ONE
1243   6:e16526.

1244   96.   TEdenovo tuto - URGI.

1245   97.   TEannot tuto - URGI.

1246   98.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X,
1247   Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A,
1248   Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N,
1249   Regev A. 2011. Trinity: reconstructing a full-length transcriptome without a
1250   genome from RNA-Seq data. Nat Biotechnol 29:644–652.

99.   Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.

Nat Protoc 11:1650–1667.

100.   Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0:

discriminating signal peptides from transmembrane regions. Nat Methods 8:785–

786.

101.   Picard Tools - By Broad Institute.

102.   Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR,

Wortman JR. 2008. Automated eukaryotic gene structure annotation using

EVidenceModeler and the Program to Assemble Spliced Alignments. Genome

Biol 9:R7.

103.   TransposonPSI: An Application of PSI-Blast to Mine (Retro-)Transposon ORF

Homologies.

104.   Haas BJ, Zeng Q, Pearson MD, Cuomo CA, Wortman JR. 2011. Approaches to

Fungal Genome Annotation. Mycology 2:118–141.

105.   Palmer J. 2017. funannotate: Fungal genome annotation scripts. Python.

106.   MEROPS - the Peptidase Database.

107.   Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T,

Mende DR, Sunagawa S, Kuhn M, Jensen LJ, von Mering C, Bork P. 2016.

eggNOG 4.5: a hierarchical orthology framework with improved functional

annotations for eukaryotic, prokaryotic and viral sequences. Nucleic Acids Res

44:D286–D293.

108. Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of

signal peptides: SignalP 3.0. J Mol Biol 340:783–795.

109. Sperschneider J, Dodds PN, Taylor JM, Duplessis S. 2017. Computational

Methods for Predicting Effectors in Rust Pathogens. Methods Mol Biol Clifton NJ

1659:73–83.

110. Garnica DP, Upadhyaya NM, Dodds PN, Rathjen JP. 2013. Strategies for Wheat

Stripe Rust Pathogenicity Identified by Transcriptome Sequencing. PLoS ONE

8:e67150.

111. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P,

Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner.

Bioinforma Oxf Engl 29:15–21.

112. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. Bioinforma Oxf Engl

30:923–930.

113. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

114. Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, Jones SJ,

Marra MA. 2009. Circos: An information aesthetic for comparative genomics.

Genome Res 19:1639–1645.

115. Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read

sequencing. ArXiv12073907 Q-Bio.

116. 2017. vcflib: a simple C++ library for parsing and manipulating VCF files, + many

command-line utilities. C++, vcflib.

117. 2017. rtg-tools: RTG Tools: Utilities for accurate VCF comparison and

manipulation. Java, Real Time Genomics.

118. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res 32:1792–1797.

119. Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein

sequence alignments into the corresponding codon alignments. Nucleic Acids Res

34:W609–W612.

120. Meyer M. Distance: Utilities for comparing sequences. C, Python.

121. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics 26:841–842.

122. Dale RK, Pedersen BS, Quinlan AR. 2011. Pybedtools: a flexible Python library

for manipulating genomic datasets and annotations. Bioinformatics 27:3423–

3424.

123.   Wheat Yellow Rust Genomics.

124.   Miller ME, Zhang Y, Omidvar V, Sperschneider J, Schwessinger B, Raley C,

1311   Palmer JM, Garnica D, Upadhyaya N, Rathjen J, Taylor JM, Park RF, Dodds PN,

1312   Hirsch CD, Kianian SF, Figueroa M. 2017. De novo assembly and phasing of

1313   dikaryotic genomes from two isolates of Puccinia coronata f. sp. avenae, the

1314   causal agent of oat crown rust. bioRxiv 179226.

1315   125.   Nemri A, Saunders DGO, Anderson C, Upadhyaya NM, Win J, Lawrence G,

1316   Jones D, Kamoun S, Ellis J, Dodds P. 2014. The genome sequence and effector

1317   complement of the flax rust pathogen Melampsora lini. Plant-Microbe Interact

1318   5:98.

1319   126.   Rochi L, Diéguez MJ, Burguener G, Darino MA, Pergolesi MF, Ingala LR, Cuyeu

1320   AR, Turjanski A, Kreff ED, Sacco F. 2016. Characterization and comparative

1321   analysis of the genome of Puccinia sorghi Schwein, the causal agent of maize

1322   common rust. Fungal Genet Biol FG B.

1323   127.   Oliphant TE. 2007. Python for Scientific Computing. Comput Sci Eng 9:10–20.

1324   128.   Perez F, Granger BE. 2007. IPython: A System for Interactive Scientific

1325   Computing. Comput Sci Eng 9:21–29.

1326   129.   McKinney W. 2010. Data Structures for Statistical Computing in Python, p. 51–56.

1327   *In* .

1328 130. Walt S van der, Colbert SC, Varoquaux G. 2011. The NumPy Array: A Structure

1329   for Efficient Numerical Computation. Comput Sci Eng 13:22–30.

1330 131. Hunter JD. 2007. Matplotlib: A 2D Graphics Environment. Comput Sci Eng 9:90–

1331   95.

1332 132. seaborn: statistical data visualization — seaborn 0.8.1 documentation.

1333

1334

1335 **Figure Legends**

1336

Figure 1: The *Pst*-104E genome is the most contiguous and complete assembly

for *Puccinia striiformis* f. sp. *tritici*

A) Assembly comparison of the *Pst*-104E primary and haplotig assembly with the two

best publicly available *Pst* genome assemblies, *Pst*-78 and *Pst*-130. The histograms

and the left y-axis show log10 counts of contigs within each size bin. The dots and the

right y-axis show the cumulative size of small to large sorted contig lengths. Each dot

represents a single contig of given size shown on the x-axis. Each plot also shows the

number of contigs or scaffolds, total assembly size, N50 of the assembly and NG50

assuming a genome size of 85 Mb. NG50 is the N50 of an assembly considering the

estimated genome size instead of the actual assembly size. This enables comparisons

between different sized assemblies.

B) Genome completeness assessed using benchmarking universal single-copy

orthologs (BUSCOs) for basidiomycota (odb9) as proxy. The graph shows BUSCO

results for *Pst*-104E primary (p), haplotig (h) and non-redundantly combined (ph)

assemblies in comparison to all publicly available *Pst* genome assemblies with gene

models including *Pst*-78, *Pst*-130, *Pst*-21, *Pst*-43, *Pst*-0821, and *Pst*-887. The analysis

was performed on the protein level using publicly available gene models. The *

indicates the actual number of identified BUSCOs for the complete *Pst*-104E ph

assembly before filtering gene models for similarity with genes encoded by transposable

elements.

1357

Figure 2: The *Pst*-104E genome is characterized by high levels of inter-haplome

variation

A) Summary of inter-haplome variation between primary contigs and their respective

haplotigs using Assemblytics. Each plot indicates the number of bases that are spanned

by the specific variation category, which is illustrated by a cartoon. The number labelling

each histogram represents the % of the total size of primary contigs with haplotigs that

are contained within this variation type and size bin. B) and C) Two representative

whole genome alignments of primary contigs 019 and 028 with their respective

haplotigs. This illustrates the large-scale variations summarized in A).

1367

Figure 3: Allele transposition in the *Pst*-104E genome

A-C) Dot plots of whole genome alignments generated using the mummer toolset where

the x-axis represents primary contig and the y-axis the haplotig sequence. A) shows the

whole genome alignments of haplotigs_027_xxx to primary contig 014. B) shows the

whole genome alignment of haplotigs_027_xxx to primary contig 027. C) shows the

whole genome alignment of haplotigs_014_xxx to primary contig 014. Black lines

indicate alignments in the forward direction and red lines in the reverse direction in the

haplotig sequence. The black rectangles highlight a ~40 kb region in haplotig_027_006

that does not align to primary contig 027 yet aligns to a region in primary contig 014,

which is not covered by an associated haplotig of 014. In D) we show micro-synteny

analysis of this extended region with primary contig 014 on top and haplotig_027_006

on the bottom. Gene models identified as alleles are labeled with their locus tag and

shaded by a light blue background. Vertical grey shading illustrates the blastn identity

between sequences on both contigs according to the scale shown in the right bottom

corner next to the sequence scale bar. Start and stop positions for each contig

sequence are given at the start and the end of each contig.

1384

Figure 4: Identification of candidate effectors based on detailed expression

analysis of secreted proteins of both *Pst*-104E assemblies

A) Clustering of *Pst*-104E secretome expression profiles for genes located on primary

contigs. Blue color intensity indicates the relative expression level using rlog

transformed read counts in spores, germinated spores, haustoria, and in wheat tissue at

6 and 9 days post infection. For example, cluster 8 shows the lowest relative expression

in spores and the highest in haustoria compared to the other clusters. B) Clustering of

Pst-104E secretome expression profiles for genes located on haplotigs.

1393

Figure 5: Candidate effector genes are spatially associated with conserved genes

and with each other

A) Nearest neighbor gene distance density hexplots for three gene categories,

including all genes, BUSCOs and candidate effectors. Each subplot represents a

distance density hexplot with the log10 3'-flanking and 5'-flanking distance to its nearest

neighboring gene plotted along the x-axis and y-axis, respectively. B) Violin plots for the

distance to the most proximal transposable element for genes in each category. C)

Violin plots for the distance to the most proximal gene in the same category for

subsamples of each category equivalent to the smallest category size (n=1444). D)

Violin plots for the minimum distance of candidate effectors and BUSCOs to each other

or a random subset of genes (n = 1444). The p-values in B, C, and D are calculated

using the Wilcoxon rank-sum test corrected for multiple testing (Bonferroni, alpha=0.05).

Violin plots shown in C and D have outliers removed and show the interquartile range

for illustration purposes

1408

Figure 6: The candidate effector allele status influences association with

conserved genes

A) Violin plots for the distance to the most proximal BUSCO for candidate effectors in

each category. The Kruskal–Wallis one-way analysis of variance of all three categories

shows a significant difference between the three samples (p ~ $2.36e^{-06}$). B) Violin plots

for the distance to the most proximal gene for candidate effectors in each category. The

Kruskal–Wallis one-way analysis of variance of all three categories shows no significant

difference between the three samples (p ~ 0.08). The p-values in A and B are calculated

using the Wilcoxon rank-sum test corrected for multiple testing (Bonferroni, alpha=0.05).

*Wilcoxon rank-sum test comparisons with non-allelic inter-haplome paralogs lack

statistical power due to the small samples size of n=28.

1420

1421   **Legends for supplemental Tables, Figures and Files**

1422

1423   **Supplemental Table 1: Summary table of PacBio genome sequencing using 13**

1424   **SMRT cells**

1425

1426   **Supplemental Table 2: Annotation of gene classes on primary contigs and**

1427   **haplotigs**

1428   For each category the number of proteins and the percentage of proteins having a hit

1429   within each category is given. First number in each column indicates the total number of

1430   proteins and the second number the percentage within each category.

1431

Supplemental Table 3: Summary of inter-haplome allele analysis

Summary table describing the allele and conservation state of *Pst*-104E genes. Primary

contigs and haplotigs were treated as two representative units for orthology analysis in

Proteinortho with the synteny flag. The three major categories are highlighted in bold.

In case of alleles they can be subdivided in three categories as illustrated in

Supplemental Figure 2. Superscript A, B, C, correspond to Supplemental Figure 2 A, B,

C.

The number in brackets for haplotype singletons is the number of singletons given by

Proteinortho without filtering for unphased gene models based on genome coverage

analysis. True haplotype singletons are located in phased regions of the genome.

1442

1443 **Supplemental Table 4: Number of total proteins and predicted secreted proteins**

1444 **in publicly available cereal rust genomes**

1445 Numbers in brackets indicates the percentage of protein that are predicted to be

1446 secreted in each proteome using SignalP3.

1447

1448 **Supplemental Table 5: Summary table of genes located in indicated expression**

1449 **clusters in regards to EffectorP, nuclear localization and apoplastic localization**

1450 **prediction**

1451 We used EffectorP (Ref.), Localizer (ref), and ApoplastP (ref) for predictions.

1452

Supplemental Table 6: Summary of the allele state and orthologous expression

pattern of secreted protein coding genes located on primary contigs

The top half of the table shows how many genes are within each cluster and of those

how many are allelic, non-allelic protein 'orthologs' or singletons.

The bottom half of the table shows how the expression of alleles of primary genes

cluster in the haplotig gene expression analysis.

1459

Supplemental Table 7: Summary of the allele state and orthologous expression

pattern of secreted protein coding genes located on haplotigs

The top half of the table shows how many genes are within each cluster and of those

how many are allelic, non-allelic protein 'orthologs' or singletons.

The bottom half of the table shows how the expression of alleles of haplotig genes

cluster in the primary gene expression analysis.

1466

1467 **Supplemental Table 8: Changes in genome size and contig number at different**

1468 **steps in the assembly process**

1469

Supplemental Figure 1: Over half of the *Pst*-104E genome is covered by repetitive

elements

A) Shows repetitive element annotation on primary contigs and B) on haplotigs. Top

panels show percentage genome coverage for all repetitive elements and different sub-

categories. These include transposable elements (TEs) of the Class I (RNA

retrotransposons) and Class II (DNA transposons), simple sequence repeats (SSR),

and unclassifiable repeats (no Cat). Middle and bottom panels show percent genome

coverage of Class I and Class II TEs categorized to class, order, and superfamily levels

wherever possible. Repetitive elements were identified using the REPET pipeline and

classifications inferred from the closest BLAST hit (see method section).

1480

1481

Supplemental Figure 2: Illustration of the three observed allele configuration

categories in the *Pst*-104E haplotype-phased assembly

A) 'Classic' alleles identified by Proteinortho for which the gene models of allele pairs

are located in regions of primary contigs and associated haplotigs that overlap in whole

genome alignments. B) Alleles detected on non-overlapping associated haplotigs.

These are alleles identified by Proteinortho for which the gene models of allele pairs are

located in regions of primary contigs and associated haplotigs that do not overlap in

whole genome alignments. C) Alleles detected on non-associated haplotigs and primary

contigs that might represent long range transpositions. These are alleles identified by

Proteinortho for which the gene models of allele pairs are located on haplotigs that are

not associated with the primary contig. See also Figure 4 for detailed illustration of one

example.

1494

Supplemental Figure 3: Spatial correlation of AT richness, repeat coverage,

candidate effectors and BUSCOs

Circa plot of the 30 largest primary contigs ordered by size. The total length shown is

greater than 50 Mb. See legend for details of each track. Lines with reverse arrowheads

illustrate regions in the *Pst*-104E genome where candidate effectors cluster in repeat

poor regions. Lines with round arrowheads illustrate regions in the *Pst*-104E genome

where candidate effectors appear to cluster with each other close to repeat dense

regions. RE is the abbreviation for repetitive element.

1503

Supplemental Figure 4: Candidate effector genes are clustered

A) Illustration of the clustering approach. In this example genes 1-3 in the black

category are clustered together as each is separated by less than 12 kb to their closest

neighbour of the same category. Gene 4 is not part of the cluster as it is more than 12

kb away from gene 3. The clustering algorithm is oblivious to the fact that gene I from

the red category is interspersed between genes 2 and 3 of the black category. This

cluster thus contain 3 black genes. B) Clustering of equal numbers of genes (n=1444)

using a maximum distance cut off of 12 kb. The y-axis represents different gene

categories and the x-axis represents the number of genes linked by a minimum distance

of 12 kb. The numbers next to the circles indicate the total number of genes in each

cluster size bin. The circles are drawn in relative size reflecting this number.

1515

1516 **Supplemental File 1: Functional annotation of *Pst*-104E protein coding genes**

1517 Tab separated list of gene IDs followed by annotation term or accession number

1518

Supplemental File 2: Allele pairs identified between primary contigs and haplotigs

of the *Pst*-104E assembly

Tab separated list of gene IDs with one pairing per line followed by one of the three

classifications: "Classic" alleles, alleles on non-overlapping associated haplotigs, alleles

on non-associated haplotigs. See Figures S2 for illustration of these allele

classifications.

1525

1526 **Supplemental File 3: Non-allelic inter-haplome paralogs of the *Pst*-104E assembly**

1527 List of gene IDs that we identified as non-allelic inter-haplome paralogs.

1528

1529 **Supplemental File 4: Unphased gene loci on primary contigs of the *Pst*-104E**

1530 **assembly**

1531 List of gene IDs that we identified as being unphased based on primary contigs after

1532 coverage analysis of Illumina short read mapping data.

1533

1534 **Supplemental File 5: Singletons of the *Pst*-104E assembly**

1535 List of gene IDs that we identified as singletons as these lack inter-haplome alleles and

1536 paralogs.

1537

1538 **Supplemental File 6: EffectorP predicted effector candidates of the *Pst*-104E**

1539 **assembly**

1540 List of gene IDs that we identified as candidate effectors using the machine learning

1541 approach EffectorP.

1542

1543 **Supplemental File 7: Expression pattern of the *Pst*-104E secretome**

1544 Tab separated list of gene IDs followed by their expression cluster as shown in Figure 4.

1545

1546 **Supplemental File 8: Non-redundant list of candidate effectors of the *Pst*-104E**

1547 **assembly**

1548 List of gene IDs of all candidate on primary contigs and haplotig effector candidate

1549 singletons.

1550

1551 **Supplemental File 9: Parameters of the FALCON assembly pipeline used for the**

1552 ***Pst*-104E assembly**

1553

1554   **Supplemental File 10: Parameters of the FALCON-Unzip processing of the *Pst*-**

1555   **104E assembly**

1556

1557   **Supplemental File 11: Parameters of the REPET Tedenovo repeat prediction of**

1558   **the *Pst*-104E assembly**

1559

1560   **Supplemental File 12: Parameters of the REPET Teanno repeat prediction of the**

1561   ***Pst*-104E assembly**

1562

1563   **Supplemental File 13: Parameters of EvidenceModeler to unify gene predictions**

1564   **of the *Pst*-104E assembly**

1565

1566   **Supplemental File 14: Parameters of PASA pipeline to map *de novo* transcripts on**

1567   **the *Pst*-104E assembly**

1568

1569

1570

1571

1572

Tables:

Table 1: Summary of *Pst*-104E genome assembly and annotation
Summary statistics for the genome assembly according to the three different contig
categories as described in the main text.
[a]Candidate effectors have been predicted based on the machine learning algorithm

EffectorP and transcriptional upregulation during infection of wheat as described in the

main text.

1580

	Primary assembly		Haplotype assembly
	Primary contigs with haplotigs	Primary contigs without haplotigs	Haplotigs
Number of contigs	99	57	475
Bases	79,770,604	3,585,012	73,478,481
TEs Coverage	53.72	67.17	52.82
Number of genes	15303	625	14321
Average gene length	1210	1290	1189
Average introns per gene	3.45	2.70	3.42
Number of genes per 10kb	1.92	1.74	1.95
BUSCOs	1395	49	1293

Number of BUSCOs per 10kb	0.17	0.14	0.18
Number of candidate effectors[a]	1523	49	1390
Number of candidate effectors[a] per 10kb	0.19	0.14	0.19

1581

1582

Table 2: Candidate effector alleles are more variable than BUSCOs alleles on the
protein level
Summary of normalized Levenshtein distances and dN/dS ratios calculated for CDS

alignments and codon based amino acid sequence alignments.

A Percentage of genes or proteins for which the normalized Levenshtein distance is

greater 0.

B are calculated using the Wilcoxon rank-sum test corrected for multiple testing

(Bonferroni, alpha=0.05).

C Number of loci for which dN/dS ratios could be calculated using yn (ref.).

D Percentage of loci for which dN/dS was not 0.

1593

	BUSCOs	Candidate effectors	Other Genes

#of loci with Levenshtein distance CDS	1198	1214	8509
% of genes showing variation[A]	81	66	79
Median	0.0069	0.0044	0.0074
Mean	0.0288	0.0409	0.0579
Wilcoxon-rank Sum test vs. candidate effectors[B]	~9.47*e-02	N.A.	~1.21*e-10
#of loci with Levenshtein distance protein	1198	1214	8509
% of proteins showing variation[A]	65	60	70
Median	0.0028	0.0060	0.0075
Mean	0.0264	0.0474	0.0637
Wilcoxon-rank Sum test vs. candidate effectors[B]	~9.46*e-05	N.A.	~1.86*e-10
#of loci with dN/dS ratio[C]	859	619	5403
%of loci showing variation[D]	75	87	85
Median	0.0802	0.3972	0.2840
Mean	0.2012	0.4797	0.3432

| Wilcoxon-rank Sum test vs. candidate effectors[B] | ~2.42*e-54 | N.A. | ~1.91*e-06 |

1594

1595

1596

1597

1598

1599

A Minimum distance of candidate effectors in given category to closest BUSCO

B Minimum distance of candidate effectors in given category to closest gene