Unliganded progesterone receptor governs estrogen receptor gene expression by regulating DNA methylation in breast cancer cells

Gaetano Verde [1, 2,*], Lara I De Llobet [1], Roni H.G. Wright [1], Javier Quilez [1], Sandra Peiró [3], François Le Dily [1], and Miguel Beato [1,*]

[1] Centre de Regulació Genòmica (CRG), Barcelona Institute for Science and Technology; Barcelona, Spain. Universitat Pompeu Fabra, Barcelona, Spain. gverde@uic.es; lara.llobet@crg.eu; roni.wright@crg.eu; javier.quilez@crg.eu; francois.ledily@crg.eu; miguel.beato@crg.eu

[2] Present address: Faculty of Medicine and Health Sciences, Universitat Internacional de Catalunya, Barcelona, Spain; gverde@uic.es

[3] Vall d'Hebron Institute of Oncology, Barcelona, Spain; speiro@vhio.net

* Correspondence: gverde@uic.es; miguel.beato@crg.eu

Abstract: Breast cancer prognosis and response to endocrine therapy strongly depends on the expression of the estrogen and progesterone receptors (ER and PR, respectively). Although much is known about ERα gene (*ESR1*) regulation after hormonal stimulation, how it is regulated in hormone-free condition is not fully understood. We used ER+/PR+ positive breast cancer cells to investigate the role of PR in *ESR1* regulation in the absence of hormones. We show that PR binds to the low-methylated *ESR1* promoter and maintains both gene expression and DNA methylation of the ESR1 locus in hormone-deprived breast cancer cells. Depletion of PR reduces *ESR1* expression, with a concomitant increase in gene promoter methylation. The high amount of methylation in the *ESR1* promoter of PR-depleted cells persists after the stable re-expression of PR and inhibits PR binding to this genomic region. As consequence, the rescue of PR expression in PR-depleted cells is insufficient to restore *ESR1* expression. Consistently, DNA methylation impedes PR binding to consensus progesterone responsive elements. These findings contribute to understand the complex crosstalk between PR and ER, and suggest that the analysis of *ESR1* promoter methylation in breast cancer cells can help to design the appropriate targeted therapies for breast cancer patients

Keywords: progesterone receptor; estrogen receptor; gene expression; DNA methylation; breast cancer, endocrine therapy.

1. Introduction

Estrogen and progesterone are main players in the progression and outcome of breast cancers [1]. Both hormones act through their cognate receptors, estrogen receptor (ER) and progesterone receptor (PR) [1]. Upon hormone exposure, PR and ER exhibit enhanced binding to specific DNA sequences known as hormone-responsive elements, which are generally located within target gene enhancers or promoters [1,2]. The DNA-bound receptors orchestrate the assembly of large cofactor-containing protein complexes that can either positively or negatively affect gene transcription [1]. In addition, hormone-activated ER and PR attached to the cell membrane can trigger rapid signaling by interacting with several kinases, which also participate in hormonal gene regulation [1,3].

PR and ER profoundly affect the breast cancer cell biology. Approximately one-third of breast cancers lack both hormone receptors (ER–/PR–) and generally shows poor histological differentiation with higher growth rates [4]. These cancers rarely respond to hormone therapies and exhibit a poor clinical outcome compared to breast cancers that express both hormone receptors (ER+/PR+) [4;5]. In addition, ER-positive tumors lacking PR (ER+/PR-) are less likely to respond to endocrine therapy compared to ER+/PR+ cancers [4;5]. Consistent with this data, hormone-free PR

enhances breast cancer cell response to estrogen and to selective ER modulator used for the
endocrine therapy, including tamoxifen and others antiestrogens [6;7]. However, how the presence
of hormone-free PR affects the breast cancer sensitivity to these external stimuli is not completely
understood.
About two-third of breast cancers are resistant to steroid hormones at the time of diagnosis but
often retain functional hormone receptors and remain highly sensitive to growth factors. Growth
factors induce breast cancer cell proliferation by enhancing the transcriptional activity of the
hormone-free PR via the protein kinases signaling pathways [8]. The phosphorylated and
under-sumoylated hormone-free PR positively regulates the expression of growth promoting genes
by recruiting steroid receptor coactivator 1 (SRC1) to gene promoters [9]. In contrast, sumoylated PR
down-regulates the expression of the same class of genes by recruiting histone deacetylase 3
(HDAC3) to promoters [9]. Thus, the hormone-free PR affects breast cancer progression by
regulating gene expression at chromatin level, even if the molecular mechanisms put into play are
not completely clearly.
The key role of hormone-free PR for the responsiveness of breast cancers to endocrine therapy
and growth factors [7,8] as well as the high correlation between PR and ER levels in breast cancers
[10] led us to explore whether and how PR regulates the ERα gene (*ESR1*) expression in
hormone-free breast cancer cells. Here we report that PR binds to the low-methylated *ESR1*
promoter to maintain its basal expression and its low level of DNA methylation in hormone-free
breast cancer cells. Consistent with these data, we show that DNA methylation hinders PR binding
to hormone-responsive elements

2. Results

2.1. PR is required to maintain ESR1 gene basal expression in hormone-free breast cancer cells

Comparing the *ESR1* gene expression in T47D breast cancer cells and a derived clone selected
for its low PR expression (T47D-Y) [11], as previously observed [12], we confirmed that low PR
expression is accompanied by a low expression of *ESR1* at both the transcript and protein levels
(Figure 1, a and b, *left panels*). Although the PR inhibits *ESR1* gene expression in T47D cells upon
progestin exposure [13], the decrease of *ESR1* expression in T47D-Y cells led us to hypothesize that
the hormone-free PR could be involved in maintaining basal *ESR1* gene expression. To test this
possibility, we knocked down PR in T47D breast cancer cells using short-hairpin RNA approach
(shPR) and analyzed *PR* and *ESR1* gene expression by RT-qPCR and Western blotting assays.
Concomitant with the decrease of PR levels, RNA and protein amount of *ESR1* decreased in shPR
cells compared to control cells (shC) (Figure 1, a and b, *right panels*). Interestingly, the strongest PR
knockdown cells showed the lowest *ESR1* levels (Figure S1, a and b), demonstrating a quantitative
relationship between the levels of the two hormone receptors.
To further confirm the decrease of PR and ER levels in PR-depleted cells (shPR) and
PR-deficient cells (T47D-Y), we analyzed the transcription of PR-target or ER-target genes in control
cells, T47D-Y and shPR cells treated with progestin (R5020), estradiol (E2) or vehicle (ethanol). The
results showed that the progestin-mediated gene expression was strongly affected in T47D-Y and
shPR cells compared to control cells, confirming that the low PR levels of T47D-Y and shPR cells
affected the transcription of PR-target genes (Figure S2). In the same manner, the estradiol-mediated
induction of *TFF1* gene transcription in control cells was strongly reduced in shPR and T47D-Y cells
(Figure 1c), consistent with the reduced ERα level upon PR loss.
Finally, to test the role of PR in the maintenance of *ESR1* gene expression in different cellular
backgrounds, we depleted the PR levels in two additional ER/PR positive cell lines, MCF7 and
BT474, using the same short-hairpin RNA approach. Despite a moderate decrease of the PR levels
in these cell lines, this slight decrease of PR was accompanied by a concomitant decrease of *ESR1*
expression compared to control cells (Figure S1, c and d), confirming the importance of PR levels in
maintaining *ESR1* gene expression in different breast cancer cell lines.

Figure 1. Loss of PR reduces the *ESR1* expression in hormone-deprived T47D breast cancer cells. (**a**) Gene-specific mRNA expression measured by quantitative RT-PCR in T47D or PR-deficient cells (T47D-Y) (*left panel*) and T47D cells transduced with shRNA against PR (shPR, clone trcn0000010776) or scramble shRNA (shC) (*right panel*). The gene-specific expression levels were normalized to *GAPDH* expression and are represented as relative values in the T47D cells. *RMND1* was used as a PR-independent control. *PGR*, PR-encoding gene; *ESR1*, ER-encoding gene. Error bars represent the SEM of three independent experiments. **P less or equal than 0.01, ***P less or equal than 0.005, unpaired two-tailed Student's *t*-test. (**b**) PR and ERα protein levels measured by Western blot in T47D and T47D-Y cells (left panel) and in T47D transduced with shRNA against PR (shPR; clone trcn0000010776) or scramble shRNA (shC) (right panel). α-tubulin protein was used as loading control. The vertical white line depicts a removed lane between the two samples. Blots are representative of three independent experiments. (**c**) PR depletion impairs estrogen-mediated gene transcription. T47D cells, PR-deficient (T47D-Y) cells, short hairpin control (shC) and PR-depleted cells (shPR, clone trcn0000010776) were treated with estradiol (E2, 10 nM) or ethanol (vehicle) for 6 hours, at which point *TFF1* mRNA expression was measured by quantitative RT-PCR. The *TFF1* gene expression was normalized to *GAPDH* expression and is represented as fold change relative to the vehicle (E2/vehicle). Error bars represent the SEM of three independent experiments. *P less or equal than 0.05, **P less or equal than 0.01, unpaired two-tailed Student's *t*-test.

2.2 PR binds to the ESR1 locus in hormone-free breast cancer cells

To test whether PR directly regulates *ESR1* gene expression prior to hormone stimulation, we analyzed ChIP-seq data obtained with an antibody to PR in serum-starved T47D cells [14]. In the absence of hormones, PR appears to bind to two genomic regions within the *ESR1* locus, one located within the gene promoter (chromosome 6: 152,128,500–152,129,000 hg19) and another within the third intron (chromosome 6: 152,222,250–152,222,650 hg19) (Figure 2a). The specificity of these two hormone-free PR binding events was confirmed by ChIP-qPCR using T47D cells and PR-deficient cells (T47D-Y) or PR-depleted cells (shPR). PR bound to the promoter and the intronic regions in T47D cells but not in T47D-Y cells (Figure 2b) or in shPR cells (Figure 2c). Strikingly, our analysis of previously published ChIP-seq experiments performed in the same conditions [2,15] revealed that the intronic sequence bound by PR in hormone-deprived T47D cells exhibited marks of active enhancers, including histone H3 mono-methylation on lysine 4 (H3K4me1) and DNase hypersensitivity (Figure 2a) [16].

We also tested PR binding at *ESR1* locus in MCF7 cells by ChIP-qPCR and found that in this
case the hormone-free PR also bound at *ESR1* promoter but not at the enhancer-like site in the third
intron of *ESR1* gene (Figure S3).

Figure 2. PR binds to the promoter and to an enhancer-like intron of the *ESR1* gene in
hormone-deprived T47D breast cancer cells. (**a**) Screen shot from the UCSC genome browser
showing the *ESR1* gene, the RNA reads, and the ChIP-seq results from PR binding, with a peak at the
gene promoter marked by polymerase 2 binding (Pol2), histone 3 trimethylated at lysine 4
(H3K4me3), low DNA methylation (5mC), and a CpG island at the bottom. Another PR peak is
found in an intronic region containing the classical enhancer epigenetic marks of DNase
hypersensitive site (DNase), histone 3 monomethylated at lysine 4 (H3K4me1), and low DNA
methylation signal (5mC). The negative control immunoprecipitation is indicated by the IgG
antibody. (**b, c**) ChIP assay was performed with a specific antibody against PR or total rabbit IgG in
T47D cells and PR-deficient cells (T47D-Y) (**b**) or PR-depleted cells (shPR, clone trcn0000010776) and
control cells expressing a scrambled shRNA (shC) (**c**). Specific binding was assessed by quantitative
PCR amplification of the *ESR1* gene promoter, an enhancer-like intronic sequence, and a genomic
region localized at 3'-end of the enhancer-like intron (negative control region). Error bars represent
the SEM of three independent experiments. **P* less or equal than 0.05, ***P* less or equal than 0.01,
***P less or equal than 0.005, unpaired two-tailed Student's *t*-test.

2.3 Rescue of PR does not restore ESR1 gene expression in PR-deficient cells

To explore whether stable expression of PR restores *ESR1* gene expression, we stably expressed
PR in PR-deficient cells (T47D-Y+PR) and analyzed *ESR1* expression. Unexpectedly, *ESR1*
expression remained significantly reduced at both transcript and protein levels after re-establishing
PR levels (Figure 3, a and b). Similarly, the estrogen-mediated induction of the *TFF1* gene remained
reduced after PR rescue (Figure 3c). In contrast, the progestin-mediated gene transcription was
significantly restored in PR-rescue cells (Figure S2), demonstrating the capability of re-expressed PR
to mediate the PR-target gene transcription.
Together, these data demonstrate that PR re-expression alone is insufficient to restore *ESR1*
gene expression to a level comparable to wild-type cells, suggesting that the *ESR1* gene is stably
repressed through another mechanism once PR is absent in breast cancer cells.

Figure 3. PR rescue of PR-deficient cells does not restore *ESR1* gene expression. Gene-specific mRNA (**a**) and protein (**b**) expression measured by quantitative RT-PCR and Western blot, respectively, in T47D control cells, PR-deficient cells (T47D-Y) and PR-rescue cells (T47D-Y+PR) (lanes T47D and T47D-Y of this image are the same as in Figure 1b and are shown here for comparison with T47DY+PR). The gene-specific mRNA expression levels were normalized to *GAPDH* expression and are represented as values relative to the T47D cells. *PGR*, PR-encoding gene; *ESR1*, ER-encoding gene. Error bars represent the SEM of three independent experiments. *P less or equal than 0.05, **P less or equal than 0.01, ***P less or equal than 0.005, unpaired two-tailed Student's *t*-test. Blots are representative of three independent experiments. (**c**) PR rescue of PR-depleted cells does not restore the estrogen-mediated gene expression. T47D, PR-deficient cells (T47D-Y), and PR-rescue (T47D-Y+PR) cells were treated with estradiol (E2, 10 nM) or ethanol (vehicle) for 6 hours, at which point TFF1 mRNA expression levels were measured by quantitative RT-PCR. Gene-specific expression levels were normalized to *GAPDH* expression and are represented as values relative to the vehicle (E2/vehicle). Error bars represent the SEM of three independent experiments. **P less or equal than 0.01, unpaired two-tailed Student's *t*-test.

2.4 Lack of PR affects DNA methylation at the ESR1 promoter

DNA methylation at the *ESR1* promoter represents one of the main epigenetic mechanisms for stably repressing *ESR1* expression in breast cancers [17]. To explore whether PR loss affects the DNA methylation profile of the *ESR1* locus, we compared the DNA methylation pattern at the *ESR1* promoter and intronic PR-binding sites between T47D control cells, PR deficient cells (T47D-Y) (Figure 4a), and PR-depleted cells (shPR) (Figure 4b) before and after PR rescue. Methylation of the *ESR1* promoter was strongly increased in PR-deficient and PR-depleted cells, and high DNA methylation amount of this genomic region persisted after PR rescue (T47D-Y+PR cells) (Figure 4a). In contrast, the DNA methylation profile of the intronic-PR binding site was not increased, neither in PR deficient nor PR depleted cells, and did not significantly change after PR rescue (Figure 4a).

Consistent with these data, as previously demonstrated [18], the analysis of the TGCA breast invasive carcinoma (BRCA) dataset showed a clear difference in *ESR1* gene methylation levels when the patients were segregated based on the PR status. PR negative breast carcinoma patients present higher methylation levels of the *ESR1* gene than PR-positive breast carcinomas. The gain of *ESR1* methylation in PR-negative breast carcinomas was stronger at gene promoter rather than within the gene body (Figure S3).

Figure 4. The loss of PR increases the DNA methylation level at the *ESR1* promoter. DNA methylation of the *ESR1* promoter and the enhancer-like intron was assessed by MeDIP-qPCR in T47D cells, T47D-Y cells and T47D-Y cells with stable PR transfection (T47D-Y+PR) (**a**), or in T47D cells transduced with shRNA against *PR* (shPR; clone trcn0000010776) or scrambled shRNA (shC) (**b**). The results are represented as values relative to the control (T47D cell in a or shC cells in b). IgG, negative control for immunoprecipitation. Error bars represent the SEM of three independent experiments. **P less or equal than 0.01, unpaired two-tailed Student's *t*-test.

2.5 DNA methylation impedes PR binding to hormone responsive elements

DNA methylation can directly affect the affinity of transcription factors towards their binding sites [19]. To check whether higher *ESR1* promoter methylation levels affect PR binding to this genomic region, we compared PR binding levels at the *ESR1* locus between T47D control cells, PR-deficient cells (T47D-Y), and PR-rescue cells (T47D-Y+PR) by ChIP-qPCR assay. As described above, PR bound to the *ESR1* promoter and to an enhancer-like intronic sequence in control cells, whereas, this binding was completely impaired in PR-deficient cells (Figure 2b). Rescue of PR in PR-deficient cells (T47D-Y+PR) completely restored PR binding at the low-methylated intronic sequence; however, it only partially restored PR binding at the highly-methylated promoter site, suggesting that hyper-methylation at the *ESR1* promoter impedes PR binding to this genomic region (Figure 5a). To test this hypothesis, we treated the PR-rescue cells (T47D-Y+PR) with the demethylating agent 5-azacytidine (5-azaC) or vehicle (control) and then compared the PR binding at the *ESR1* locus between control and 5-azaC-treated cells. The results showed that 5-azaC treatments significantly reduced DNA methylation levels at *ESR1* promoter and had a tendency to increase PR binding levels at this genomic site (Figure 5, b and c). In contrast, 5-azaC treatments did not affect the DNA methylation and the PR binding at the *ESR1* intronic region (Figure 5, b and c).

The CpG island at the *ESR1* promoter contains a canonical progesterone-responsive elements (PRE) encompassing a CpG as well as several half palindromic PRE sites with one or two neighboring CpGs (Figure 6a). To determine whether methylation of PRE affects the PR binding in vitro, we tested whether PR can bind to methylated or unmethylated forms of PRE oligonucleotides by electrophoresis mobility shift assay (EMSA). We observed that PR bound more efficiently to the unmethylated probes than to their methylated counterparts, especially when the PRE contained two CpGs rather than one (Figure 6, b and c). Further, an unmethylated PRE, but not its methylated counterpart, was a high-affinity competitor in EMSA for an oligonucleotide probe without CpG, which was previously shown to be a strong PR binding site (8) (Figure 6d). In addition, we observed that the CpG-containing PRE were bound by PR less efficiently than the PRE without CpGs (Figure 6, b and c), suggesting that the presence of CpG, even not methylated, reduces PR binding affinity. Finally, we analyzed the methylation of all genomic regions bound by hormone-free PR [14]. Despite having a higher CpG content than their flanking genomic regions, all the hormone-free PR binding sites had an overall lower level of methylation than their surrounding areas (Figure S4), suggesting that not only the *ESR1* locus but also the other PR binding sites require low levels of methylation for PR binding.

Figure 5. DNA methylation affects PR binding to the *ESR1* promoter. (**a**) The high-methylated *ESR1* promoter, in contrast to the low-methylated intronic sequence, was only partially bound by PR in PR-rescued cells (T47D-Y+PR). ChIP assays were performed with a specific antibody against PR. Specific binding was assessed by quantitative PCR amplification of the *ESR1* gene promoter and an enhancer-like intronic sequence in T47D, PR-deficient cells (T47D-Y) and PR-rescue cells (T47D-Y+PR). Error bars represent the SEM of three independent experiments. *P less or equal than 0.05, **P less or equal than 0.01, ***P less or equal than 0.005, unpaired two-tailed Student's *t*-test. (**b**) The 5-azacytidine (5-azaC) demethylated *ESR1* promoter. DNA methylation analysis was performed using T47D-Y+PR cells treated for 112 hours with the demethylating agent 5-azaC (5 μM) or vehicle (Control). The results are represented as fold change relative to the control. Error bars represent the SEM of three independent experiments. **P less or equal than 0.01. (**c**) The demethylating agent 5-azaC increases PR binding at the *ESR1* promoter in PR-rescued cells (T47D-Y+PR). ChIP was performed as in figure 5b using T47D-Y+PR cells treated for 112 hours with the demethylating agent 5-azaC (5 μM) or vehicle (Control). Error bars represent the SEM of three independent experiments. $P = 0.07$, unpaired two-tailed Student's *t*-test.

Figure 6. DNA methylation impedes PR binding to progesterone-responsive elements (PREs). (**a**) Screen shot from the UCSC genome browser showing the CpG island (CpG 89) at the *ESR1* promoter and the positions of PREs containing one or two CpG dinucleotides. (**b**, **c**) Electrophoretic-mobility shift assay using the indicated amount of purified human PR to capture the PRE with no CpG (ACAGTTTGT; No CpG), one methylated (MetCpG) or unmethylated CpG (UnmetCpG) (ACGGTTTGT) (**b**), two methylated (MetCpGs) or two unmethylated CpGs (UnmetCpGs) (ACGGTTCGT); (**c**). Quantification of the percentage of PR binding to different probes is shown in the lower part of the gel images. **P* less or equal than 0.05, ***P* less or equal than 0.01, unpaired two-tailed Student's *t*-test. (**d**) A double-stranded oligonucleotide probe with no CpG (ACAGTTTGT) was incubated with 2.4 μg of purified human PR and analyzed by PAGE either in the absence (−) or presence (+) of 100-fold excess of unlabelled oligonucleotides containing two unmethylated (UNMET) or two methylated (MET) CpGs (ACGGTTCGT). Error bars represent the SEM of three independent experiments. ****P* less or equal than 0.005, unpaired two-tailed Student's *t*-test. The dashed grey line indicates that a lane between the two samples was removed.

3. Discussion

The study of the role of PR in hormone-free breast cancer cells helps to clarify how these cells respond to external stimuli, including growth factors and ER modulators used for the endocrine therapy. We show here that PR binds the *ESR1* locus and is required to maintain the *ESR1* gene expression in hormone-free breast cancer cells. When the levels of PR are reduced, *ESR1* gene expression decreases in parallel with an increase of the DNA methylation level at the *ESR1* promoter, suggesting that hormone-free PR maintains *ESR1* expression by preserving a low DNA methylation at the *ESR1* promoter. In line with these results, as previously demonstrated [18], our in silico analysis confirmed that PR-negative breast cancer patients present higher *ESR1* promoter methylation than PR-positive breast carcinomas. Consistently, demethylation of the *ESR1* promoter reactivates *ESR1* expression in ER-negative breast cancer cells [20].

The loss of PR did not affect DNA methylation of the *ESR1* intronic site, which in contrast to *ESR1* promoter lacks a CpG island, suggesting that PR selectively prevents methylation around CpG islands. However, whether the PR regulates *ESR1* promoter methylation in a direct or an indirect manner is not clear. Different molecular mechanisms activated by PR down regulation could be responsible for *ESR1* promoter hypermethylation in PR-depleted cells. In this context, it was suggested that DNA methylation can be a secondary event following gene silencing, rather than

being a major event preceding it [21]. Thus, we cannot exclude that the reduced levels of *ESR1*
transcription induced by PR depletion may drive the gain of *ESR1* promoter methylation in
PR-depleted cells.
In cells stably lacking PR expression, the stable re-expression of PR did not affect the
hyper-methylation found at the *ESR1* promoter, and was insufficient to reactivate *ESR1* gene
expression. Moreover, rescue with PR completely restored the PR binding at low-methylated
intronic sequence but only partially restored it at the highly methylated *ESR1* promoter site,
suggesting that DNA methylation affects the PR binding to DNA. Consistently, the treatment of
PR-rescued cells with the demethylating agent 5-azaC showed a tendency to increase PR binding at
ESR1 promoter but not at the low-methylated intronic site. Moreover, *in vitro* DNA binding
experiments showed that PR preferentially bound unmethylated PRE oligonucleotides rather than
their methylated counterpart, demonstrating that PR is a methylation-sensitive DNA binding
protein. Consistent with these data, we also observed that all the PR binding sites in hormone-free
breast cancer cells show low methylation levels. Taken together, these data demonstrate that the
gain of DNA methylation at the *ESR1* promoter observed upon PR loss increases DNA methylation
that impedes PR binding to the *ESR1* promoter and maintains reduced *ESR1* transcription after PR
rescue (Figure 7).
In many cases, the transcriptional regulation of steroid target genes requires the action of
regulatory sequences located far away from the promoters [22]. A significant fraction of these distal
sequences engages in physical interactions with promoters, suggesting that they act as enhancers
[22]. In this study, we showed that the PR binding site within the *ESR1* intronic sequence in T47D
breast cancer cells exhibits the classical epigenetic marks found at active enhancer regions, including
the monomethylation of lysine 4 of histone 3 (H3K4me1), a low DNA methylation, and a DNase
hypersensitivity [2,15]. This suggests that PR through its binding to the *ESR1* promoter and the
enhancer-like intronic sequence could facilitate the interaction between these two genomic regions
to enhance the *ESR1* transcription in T47D breast cancer cells. However, further analyses are
required to investigate the enhancer activity of the *ESR1* intronic PR binding site and its possible
interaction with the gene promoter.
The expression of ER and PR strongly affects breast cancer prognosis and response to endocrine
therapy, with double-negative ER–/PR– breast cancers having worse clinical outcome than ER+/PR+
breast cancers [4;5]. Although ER is expressed in a large proportion of PR-positive breast cancers
(ER+/PR+), a smaller percentage of patients express ER without PR (ER+/PR-) [4], suggesting that
PR-independent molecular mechanisms can still maintain a lower ER expression. In agreement, the
ER+/PR+ breast cancers are much more likely to respond to estrogen and selective ER modulator
(SERM) therapy than breast cancers that lack PR protein (ER+/PR−) [5]. Since the SERM treatments
target ER, how the absence of PR affects this kind of therapy was unclear. Our finding that
PR-deficient cells show lower ER levels compared to control cells, suggests that the ER+/PR+ breast
cancers are more likely to respond to SERM therapy because they could maintain the *ESR1* levels
higher than breast cancers without PR protein.
In conclusion, in this study, we demonstrate that PR binds to the low-methylated *ESR1*
promoter and maintains both gene expression and the DNA methylation profile of the *ESR1* locus in
hormone-free breast cancer cells. These data expand our understanding of the complex crosstalk
between PR and ER, and suggest that the analysis of DNA methylation of *ESR1* promoter in breast
cancer cells can help to design more appropriate targeted therapies for different types of breast
cancer patients.

Figure 7. Model of regulation of the *ESR1* gene expression and DNA methylation by PR in
hormone-free breast cancer cells. In hormone-free ER+/PR+ breast cancer cells, PR binds to
low-methylated gene promoter as well as to an enhancer-like intronic sequence of *ESR1*. PR binding
at the gene promoter is required for maintaining *ESR1* transcription. In the absence of PR, DNA
methylation (mC) increases at the ESR1 promoter, and ESR1 transcription is reduced. Re-expression
of PR in PR-depleted cells leads to PR binding to the low-methylated enhancer-like intronic
sequence, but the high level of DNA methylation (mC) at the ESR1 promoter impedes PR binding to
this genomic region. Consequently, re-expression of PR in PR-depleted cells is insufficient to restore
ESR1 expression.

4. Materials and Methods

4.1 Cell culture and hormone treatments

The T47D-MTVL (T47D) breast cancer cells used in this study have a stable integrated copy of
the luciferase reporter gene driven by the MMTV promoter [14]. The T47D, T47D-Y [11], T47D-Y+PR
[12], were routinely grown in RPMI 1640 medium, MCF7 and BT474 cells in DMEM medium. The
mediums were supplemented with 10% FBS, glutamine and standard antibiotics. For the
experiments in the absence of hormones, cells were grown 48 hours in medium without phenol red
supplemented with 10% dextran-coated charcoal treated FBS (DCC/FBS). In contrast to normal FBS,
the DCC/FBS does not contain hormone-similar compounds that can activate PR activity. Moreover,
since the cell cycle phase can influence PR activity [23], cells were then synchronized in G0/G1 by
serum starvation for 16 hours to avoid cell cycle phase variability. For the hormone treatments, cells
were then induced with E2 (10 nM), R5020 (10 nM), or vehicle (ethanol) for 6 hours. For
5-azacytidine treatments, T47D-Y+PR cells were grown for 96 hours (48 hours using DMEM medium
supplemented with 10% FBS, and 48 hours with DMEM without phenol red supplemented with 10%
DCC/FBS) with 5 μM of 5-azacytidine (A3656, Sigma-Aldrich) or vehicle (1:1 acetic acid to water).
Cells were finally synchronized in G0/G1 by 16 hours of serum starvation before performing PR
binding and DNA methylation analysis.

4.2 Lentivirus preparation and infection

HEK-293 cells were transfected with pVSV-G [24] and pCMVΔR8.91 [25], together with the
pLKO.1-puro non-targeting vector (SHC001; Sigma-Aldrich) or pLKO.1-shRNA against PR using

CaCl₂ to permeabilize the cell membrane. Two different clones of PLKO.1-shRNA against PR have
been used: clone trcn0000010776 and clone trcn0000003321 (SHCLND-NM_000926, Sigma-Aldrich).
The viral particles containing the shRNA were collected 72 hours after the transfection and used to
stably infect breast cancer cells. Cell populations were finally selected with puromycin (1 µg/ml) and
processed to quantify mRNA and protein expression.

4.3 Reverse transcription and quantitative PCR

Total RNA was isolated with the RNeasy extraction kit (Qiagen). Complementary DNA
(cDNA) was generated from 100ng of total RNA with the First Strand cDNA Superscript II Synthesis
kit (Invitrogen; #11904018) and analyzed by quantitative PCR. Gene-specific expression was
regularly normalized to *GAPDH* expression. Primers sequences are listed in Table S1.

4.4 Western blotting

Cell lysates were resolved on SDS-polyacrylamide gels, and the proteins were transferred to
Hybond-ECL nitrocellulose membranes (Amersham). Membranes were blocked with TBS-0.1%
Tween 20 (TBS-T) with 5% of skimmed milk, incubated for 1 hour at room temperature with a
primary antibody (antibody against PR, sc-7208 from Santa Cruz Biotechnology; antibody against
ERα, sc-543 from Santa Cruz Biotechnology; antibody against α-tubuline, T9026 from Sigma),
diluted in TBS-T with 2.5% skimmed milk. After three washes with TBS-T, membranes were
incubated for 45 min at room temperature with horseradish peroxidase–conjugated secondary
antibodies (GE Healthcare). Antibody binding was detected by chemiluminescence on a LAS-3000
image analyzer (Fuji PhotoFilm).

4.5 Chromatin immunoprecipitation

Chromatin immunoprecipitation (ChIP) assays were performed as described previously [26],
with minor modifications. Cells were cross-linked in medium containing 1% formaldehyde for
10 min at 37°C, and crosslinking was quenched with 125 mM glycine for 5 min at room temperature.
After cells were lysed in hypotonic buffer, the nuclei were lysed with SDS-lysis buffer. Chromatin
was sheared by sonication and incubated 16 hours with 5µg of antibody against progesterone
receptor (PR, Santa Cruz Biotechnology, sc-7208) or rabbit IgG (Cell Signaling, #2729s).
Immunocomplexes were recovered with protein A agarose bead slurry (Diagenode, #C03020002) for
2 hours with rotation and washed with ChIP buffer (Diagenode, #K0249001) and Tris-EDTA buffer.
For reversing the crosslinking, samples were incubated with proteinase K (10 mg/ml) at 65°C for 16
hours. DNA was purified and analyzed by quantitative PCR. Primer sequences are listed in table S1.
For PR ChIP-seq analysis, the reads of the previously published PR ChIP-seq [14] were trimmed
using Trimmomatic (version 0.33) with the parameters values recommended by Bolger *et al*. [27].
The trimmed reads were aligned to the hg19 assembly version of the human genome using BWA
(version: 0.7.12-r1039) [28]. The FASTA file containing the genome reference sequence of hg19 was
downloaded from the UCSC Genome Browser discarding the random scaffolds and the alternative
haplotypes from the reference sequence for the alignment [29]. BWA-MEM algorithm and SAMtools
(version: 1.2, using htslib 1.2.1) [30] were used to convert SAM files to BAM files and to sort them to
retain only uniquely aligned reads. The PR binding sites were identified with the MACS2 tool
(version 2.1.0.20150420) [31]. Peaks were additionally filtered until those remaining had a false
discovery rate (FDR) q-value $< 10^{-6}$ and a 4-fold enrichment over the control sample (input), leaving
476 peaks for subsequent analyses.

4.6 DNA methylation

The DNA methylation analyses were performed by methylated DNA immunoprecipitation
assay coupled with quantitative-PCR (MeDIP-qPCR) or high-throughput sequencing (MeDIP-seq).
For MeDIP-qPCR, genomic DNA was randomly sheared by sonication to generate fragments
between 300 and 700 bp. Sonicated DNA was denatured and then immunoprecipitated using an

413 antibody against 5mC (Eurogentec; #BI-MECY-1000) or mouse IgG antibody as previously described
414 [32]. The immunocomplexes were recovered using 8 μl Dynabeads (M-280; Life Technologies), and
415 the pull-down products were detected by quantitative-PCR. Primers sequences are listed in Table S1.
416  For MeDIP-seq, adaptors from the NEBNext Ultra DNA Library Prep Kit from Illumina were
417 added to the fragmented DNA. Fragmented DNA was immunoprecipitated with antibody against
418 5mC as described above, and the amplified library was prepared using NEBNext Ultra DNA Library
419 Prep Kit for Illumina (E7370L) following manufacturer's instructions. Amplified libraries were
420 sequenced, and reads were aligned with BowTie v1.1.2 using the reference human genome version
421 19 (hg19) as previously described [33]. The mapped reads were filtered for duplicates and used to
422 compute genome-wide reads-per-million (RPM) normalized signal tracks. The 5mC and CpG heat
423 maps were generated using DeepTools (version 2.2.0) [34] and BEDtools (version v2.24.0) [35] and
424 the matrix underlying the heatmaps was used to generate the 5mC and CpG average profiles. To test
425 the significance of the overall reduction of 5mC methylation observed in the progesterone-receptor
426 binding sites (PRBs), we calculated the average 5mC normalized read counts signal over each PRBs
427 and random regions resulting from shuffling the genomic coordinates of the PRBs, while keeping
428 their sizes as in the true set of regions (this second step was repeated 1,000 times to generate an
429 empirical null distribution of 5mC methylation averaged values). The Mann-Whitney U Test was
430 applied using the stats module of the Python's SciPy library [36].
431  The DNA methylation data obtained by MeDIP-seq method are available in the GEO
432 repository. Accession number: GSE107966. For the reviewers: to access to this data, use the following
433 link: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE107966 with the secure token:
434 efwfueewdfizjov

435 *4.7 Electrophoresis mobility-shift assay*

436  Recombinant human PR (isoform B; PRB) was expressed in baculovirus and purified as
437 previously described [37]. Radioactive double-stranded oligonucleotides containing the
438 progesterone-responsive elements (PRE) were incubated with the indicated amounts of PR-B for 20
439 min at room temperature and analyzed in a 5% acrylamide-bisacrylamide electrophoresis gel. The
440 radioactivity of the DNA-protein complex was then quantified by using the PhosphorImager and
441 ImageQuant software (Molecular Dynamics). For the EMSA competition assay, a radioactive
442 oligonucleotide without CpGs was first mixed with 100-fold of non-radioactive unmethylated or
443 methylated probe containing two CpGs and then incubated with 2.4 μg of PR-B for 20 min at room
444 temperature. DNA-protein complexes either in absence or presence of unlabelled oligonucleotides
445 were then analyzed as described above. Oligonucleotides sequences are listed in Table S1.

446 **Supplementary Materials:** Figure S1: Depletion of PR reduces the *ESR1* expression in hormone-free breast
447 cancer cells; Figure S2: PR depletion and PR rescue affect progestin-mediated gene transcription; Figure S3: PR
448 binds at *ESR1* gene promoter in MCF7 breast cancer cells; Figure S4: DNA methylation at the *ESR1* locus is
449 higher in PR-negative than in PR-positive breast invasive cancers; Figure S5: The PR binding sites are low
450 methylated in hormone-deprived breast cancer cells. Table S1: Oligonucleotide sequences.

451 **Author Contributions:** G.V., F.L.D., and M.B designed the study and strategy of the project; G.V. and L.I.D.L.
452 performed the experimental work; J.Q. provided the bioinformatics analysis of high throughput data. R.H.G.W
453 performed the analysis of the TGCA dataset. G.V. and F.L.D. discussed the results and wrote the paper. S.P.
454 helps with discussion of the results. All the authors contributed to editing the manuscript.

455 **Funding:** We received funding from Spanish Ministry of Economy and Competitiveness, Plan Nacional project
456 SAF 2013-42497-P; Centro de Excelencia Severo Ochoa 2013-2017; the CERCA Programme/ Generalitat de
457 Catalunya; G.V. has received funding from the Spanish Ministry of Economy and Competitiveness, "Juan de la
458 Cierva Incorporation" fellowship (ref. IJCI-2014-20723), the European Union Seventh Framework Programme
459 (FP7/2007-2013) under grant agreement number 299429 and the European Molecular Biology Organization
460 (EMBO long-term fellowship ALTF 1106-2011, cofunded with the European Commission EMBOCOFUND2010,
461 GA-2010-267146).

Acknowledgments: We acknowledge support from Daniel Soronellas and Giancarlo Castellano for the DNA methylation mapping of MeDIP-sequencing analysis, Jofre Font for the PR purification, Enrique Vidal for revision of the statistical analysis. Roberto Ferrari and Aldo Verde for their support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Hilton, HN; Clarke, CL; Graham, JD. Estrogen and progesterone signaling in the normal breast and its implications for cancer development. *Mol Cell Endocrinol*. **2017**, pii: S0303-7207(17), 30433-1, 10.1016/j.mce.2017.08.011.

2. Ballaré, C; Castellano, G; Gaveglia, L; Althammer, S; González-Vallinas, J; Eyras, E; Le Dily, F; Zaurin, R; Soronellas, D; Vicent, GP; Beato, M. Nucleosome-Driven Transcription Factor Binding and Gene Regulation. *Mol. Cell*. **2013**, 49: 67–79, 10.1016/j.molcel.2012.10.019.

3. Wierer, M; Verde, G; Pisano, P; Molina, H; Font-Mateu, J; Di Croce, L; Beato, M. PLK1 signaling in breast cancer cells cooperates with estrogen receptor-dependent gene transcription. *Cell Rep.* **2013**, 3(6):2021-32, 10.1016/j.celrep.2013.05.024.

4. Dunnwald, LK; Rossing, MA; Li, CI. Hormone receptor status, tumor characteristics, and prognosis: a prospective cohort of breast cancer patients. *Breast Cancer Res*. **2007**, 9(1):R6, 10.1186/bcr/bcr1639.

5. Cui, X; Schiff, R; Arpino, G; Osborne, CK; Lee, AV. Biology of progesterone receptor loss in breast cancer and its implications for endocrine therapy. *J Clin Oncol*. **2005**, 23(30):7721-35, 10.1200/JCO.2005.09.004.

6. Daniel, AR; Gaviglio, AL; Knutson, TP; Ostrander, JH; D'Assoro, AB; Ravindranathan, P; Peng, Y; Raj, GV; Yee, D; Lange, CA. Progesterone receptor-B enhances estrogen responsivenss of breast cancer cells via scaffolding PELP1- and estrogen receptor-containing transcription complexes. *Oncogene*. **2015**, 34: 506–515, 10.1038/onc.2013.579.

7. Encarnacion, CA; Ciocca, SR; McGuire, WL; Clark, GM; Fuqua, SAW; Osborne, CK. Measurement of steroid hormone receptors in breast cancer patients on tamoxifen. *Breast Cancer Res Treat.* **1993**, 26(3):237-46.

8. Qiu, M; Lange, CA. MAP kinases couple multiple functions of human progesterone receptors: degradation, transcriptional synergy, and nuclear association. *J Steroid Biochem Mol Biol*. **2003**, 85(2-5):147-57, 10.1016/S0960-0760(03)00221-8.

9. Daniel, AR; Lange, CA. Protein kinases mediate ligand-independent derepression of sumoylated progesterone receptors in breast cancer cells. *Proc Natl Acad Sci USA*. **2009**, 106(34):14287-92, 10.1073/pnas.0905118106.

10. Bendrat, K; Fritz, P; Müller, S; Brockmöller, S; Debus, A; Friedrichs, K; Lindner, C; Brinkmann, F; Heidemann, E; Niendorf, A. Improved Risk Stratification for Breast Cancer Samples Based on the Expression Ratio of the Estrogen and Progesterone Receptor. *Anticancer Res*. **2016**, 36(8):3855-63.

11. Sartorius, CA; Groshong, SD; Miller, LA; Powell, RL; Tung, L; Takimoto, GS; Horwitz, KB. New T47D breast cancer cell lines for the independent study of progesterone B- and A-receptors: Only antiprogestin-occupied B-receptors are switched to transcriptional agonists by cAMP. *Cancer Res.* **1994**, 54: 3868-77.

12. Quiles, I; Millán-Ariño, L; Subtil-Rodríguez, A; Miñana, B; Spinedi, N; Ballaré, C; Beato, M; Jordan, A. Mutational analysis of progesterone receptor functional domains in stable cell lines delineates sets of genes regulated by different mechanisms. *Mol. Endocrinol*. **2009**, 23: 809–826, 10.1210/me.2008-0454.

13. Alexander, IE; Shine, J; Sutherland, RL. Progestin regulation of estrogen receptor messenger RNA in human breast cancer cells. *Mol. Endocrinol*. **1990**, 6: 821-828, 10.1210/mend-4-6-821.

14. Vicent, GP; Nacht, AS; Zaurin, R; Font-Mateu, J; Soronellas, D; Le Dily F; Reyes, D; Beato, M. Unliganded progesterone receptor mediated targeting of an RNA-containing repressive complex silences a subset of hormone-inducible genes. *Genes Dev.* **2013**, 27: 1179–1197, 10.1101/gad.215293.113.

15. Le Dily, F; Baù, D; Pohl, A; Vicent, GP; Serra, F; Soronellas, D; Castellano, G; Wright, RH; Ballaré, C; Filion, G; Marti-Renom, MA; Beato, M. Distinct structural transitions of chromatin topological domains correlate with coordinated hormone-induced gene regulation. *Genes Dev*. **2014**, 28: 2151–2162, 10.1101/gad.241422.114.

16. Calo, E; Wysocka, J. Modification of enhancer chromatin: What, How, and Why? *Molecular Cell*. **2013**, 49: 825–837, 10.1016/j.molcel.2013.01.038.

17. Martínez-Galán, J; Torres-Torres, B; Núñez, MI; López-Peñalver, J; Del Moral, R; Ruiz De Almodóvar, JM;
Menjón, S; Concha, A; Chamorro, C; Ríos, S; Delgado, JR. ESR1 gene promoter region methylation in free
circulating DNA and its correlation with estrogen receptor protein expression in tumor tissue in breast
cancer patients. *BMC Cancer*. **2014**, 14: 59, 10.1186/1471-2407-14-59.

18. Widschwendter, M; Siegmund, KD; Müller, HM; Fiegl, H; Marth, C; Müller-Holzner, E; Jones, PA; Laird,
PW. Association of breast cancer DNA methylation profiles with hormone receptor status and response to
tamoxifen. *Cancer Res*. **2004**, 64: 3807–3813, 10.1158/0008-5472.CAN-03-3852.

19. Yin, Y; Morgunova, E; Jolma, A; Kaasinen, E; Sahu, B; Khund-Sayeed, S; Das, PK; Kivioja, T; Dave, K;
Zhong, F; Nitta, KR; Taipale, M; Popov, A; Ginno, PA; Domcke, S; Yan, J; Schübeler, D; Vinson, C; Taipale,
J. Impact of cytosine methylation on DNA binding specificities of human transcription factors. *Science*.
2017, 356(6337), pii: eaaj2239, 10.1126/science.aaj2239.

20. Ferguson, AT; Lapidus, RG; Baylin, SB; Davidson, NE. Demethylation of the estrogen receptor gene in
estrogen receptor-negative breast cancer cells can reactivate estrogen receptor gene expression. *Cancer Res*.
1995, 55: 2279–2283.

21. Jones, PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat Rev Genet*.
2012, 13(7):484-92, 10.1038/nrg3230.

22. Carroll, JS; Liu, XS; Brodsky; AS, Li, W; Meyer, CA; Szary, AJ; Eeckhoute, J; Shao, W; Hestermann, EV;
Geistlinger, TR; Fox, EA; Silver, PA; Brown, M. Chromosome-wide mapping of estrogen receptor binding
reveals long-range regulation requiring the forkhead protein FoxA1. *Cell*. **2005**, 122: 33–43,
10.1016/j.cell.2005.05.008.

23. Dressing, GE; Knutson, TP; Schiewer, MJ; Daniel, AR; Hagan, CR; Diep, CH; Knudsen, KE; Lange, CA.
Progesterone receptor-cyclin D1 complexes induce cell cycle-dependent transcriptional programs in breast
cancer cells. *Mol Endocrinol*. **2014**, 28(4):442-57, 10.1210/me.2013-1196.

24. Stewart, SA; Dykxhoorn, DM; Palliser, D; Mizuno, H; Yu, EY; An, DS; Sabatini, DM; Chen, IS; Hahn, WC;
Sharp, PA; Weinberg, RA; Novina, CD. Lentivirus-delivered stable gene silencing by RNAi in primary
cells. *RNA*. **2003**, 9: 493–501.

25. Zufferey, R; Nagy, D; Mandel, RJ; Naldini, L; Trono, D. Multiply attenuated lentiviral vector achieves
efficient gene delivery in vivo. *Nat Biotechnol*. **1997**, 15: 871–875, 10.1038/nbt0997-871.

26. Strutt, H; Paro, R. Mapping DNA target sites of chromatin proteins in vivo by formaldehyde crosslinking.
Methods Mol Biol. **1999**, 119: 455–67, 10.1385/1-59259-681-9:455.

27. Bolger, AM; Lohse, M; Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics. **2014**, 30: 2114–2120, 10.1093/bioinformatics/btu170.

28. Li, H; Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*.
2009, 25: 1754–60, 10.1093/bioinformatics/btp324

29. Karolchik, D; Barber, GP; Casper, J; Clawson, H; Cline, MS; Diekhans, M; Dreszer, TR; Fujita, PA;
Guruvadoo, L; Haeussler, M; Harte, RA; Heitner, S; Hinrichs, AS; Learned, K; Lee, BT; Li, CH; Raney, BJ;
Rhead, B; Rosenbloom, KR; Sloan, CA; Speir, ML; Zweig, AS; Haussler, D; Kuhn, RM; Kent, WJ. The UCSC
Genome Browser database: 2014 update. *Nucleic Acids Res*. **2014**, 42: D764-70, 10.1093/nar/gkt1168.

30. Li, H; Handsaker, B; Wysoker, A; Fennell, T; Ruan, J; Homer, N; Marth, G; Abecasis, G; Durbin, R; 1000
Genome Project Data Processing Subgroup. The sequence Alignment/Map format and SAMtools.
Bioinformatics. **2009**, 25, 2078–9, 10.1093/bioinformatics/btp352.

31. Zhang, Y; Liu, T; Meyer, CA; Eeckhoute, J; Johnson, DS; Bernstein, BE; Nusbaum, C; Myers, RM; Brown,
M; Li, W; Liu, XS. Model-based analysis of ChIP-Seq (MACS). *Genome Biol*. **2008**, 9: R137,
10.1186/gb-2008-9-9-r137.

32. Mohn, F; Weber, M; Schübeler, D; Roloff, TC. Methylated DNA immunoprecipitation (MeDIP). *Methods*
Mol Biol. **2009**, 507:55-64, 10.1007/978-1-59745-522-0_5

33. Langmead, B; Trapnell, C; Pop, M; Salzberg, SL. Ultrafast and memory-efficient alignment of short DNA
sequences to the human genome. *Genome Biol*. **2009**, 10: R25, 10.1186/gb-2009-10-3-r25.

34. Ramírez, F; Ryan, DP; Grüning, B; Bhardwaj, V; Kilpert, F; Richter, AS; Heyne, S; Dündar, F; Manke, T.
deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res*. **2016**, 44:
160–165, 10.1093/nar/gkw257.

35. Quinlan, AR. BEDTools: The swiss-army tool for genome feature analysis. *Curr. Protoc. Bioinformatics*.
2014, 47: 11.12.1-11.12.34, 10.1002/0471250953.bi1112s47.

36. Oliphant, TE. Python for scientific computing. *Comput. Sci. Eng*. **2007**, 9(3): 10–20, 10.1109/MCSE.2007.58

37. Di Croce, L; Koop, R; Venditti, P; Westphal, HM; Nightingale, KP; Corona, DF; Becker, PB; Beato, M.
Two-step synergism between the progesterone receptor and the DNA- binding domain of nuclear factor 1
on MMTV minichromosomes. *Mol. Cell*. **1999**, 4(1): 45–54, 10.1016/S1097-2765(00)80186-0.