1   A model species for agricultural pest genomics: the genome of the Colorado potato beetle,
2   *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae)
3

Sean D. Schoville[1]*, Yolanda H. Chen[2], Martin N. Andersson[3], Joshua B. Benoit[4], Anita
Bhandari[5], Julia H. Bowsher[6], Kristian Brevik[2], Kaat Cappelle[7], Mei-Ju M. Chen[8], Anna K.
Childers[9,10], Christopher Childers[8], Olivier Christiaens[7], Justin Clements[1], Elena N. Elpidina[11],
Patamarerk Engsontia[12], Markus Friedrich[13], Inmaculada García-Robles[14], Richard A. Gibbs[15],
Chandan Goswami[16], Alessandro Grapputo[17], Kristina Gruden[18], Marcin Grynberg[19], Bernard
Henrissat[20,21,22], Emily C. Jennings[4], Jeffery W. Jones[13], Megha Kalsi[23], Sher A. Khan[24],
Abhishek Kumar[25,26], Fei Li[27], Vincent Lombard[20,21], Xingzhou Ma[27], Alexander Martynov[28],
Nicholas J. Miller[29], Robert F. Mitchell[30], Monica Munoz-Torres[31], Anna Muszewska[19], Brenda
Oppert[32], Subba Reddy Palli[23], Kristen A. Panfilio[33,34], Yannick Pauchet[35], Lindsey C. Perkin[32],
Marko Petek[18], Monica F. Poelchau[8], Éric Record[36], Joseph P. Rinehart[10], Hugh M. Robertson[37],
Andrew J. Rosendale[4], Victor M. Ruiz-Arroyo[14], Guy Smagghe[7], Zsofia Szendrei[38], Elise M.
Szuter[4], Gregg W.C. Thomas[39], Alex S. Torson[6], Iris M. Vargas Jentzsch[33], Matthew T.
Weirauch[40,41], Ashley D. Yates[42,43], George D. Yocum[10], June-Sun Yoon[23], Stephen Richards[15]

1   Department of Entomology, University of Wisconsin-Madison
2   Department of Plant and Soil Sciences, University of Vermont
3   Department of Biology, Lund University
4   Department of Biological Sciences, University of Cincinnati
5   Department of Molecular Physiology, Christian-Albrechts-University at Kiel
6   Department of Biological Sciences, North Dakota State University
7   Department of Crop Protection, Ghent University
8   USDA-ARS National Agricultural Library, Beltsville, MD USA
9   USDA-ARS Bee Research Lab, Beltsville, MD USA
10   USDA-ARS Insect Genetics and Biochemistry Research Unit, Fargo, ND USA
11   A.N. Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University
12   Department of Biology, Faculty of Science, Prince of Songkla University, Thailand
13   Department of Biological Sciences, Wayne State University
14   Department of Genetics, University of Valencia
15   Department of Molecular and Human Genetics, Baylor College of Medicine
16   National Institute of Science Education and Research, Bhubaneswar, India
17   Department of Biology, University of Padova
18   Department of Biotechnology and Systems Biology, National Institute of Biology, Slovenia
19   Institute of Biochemistry and Biophysics, Polish Academy of Sciences
20   Architecture et Fonction des Macromolécules Biologiques, CNRS, Aix-Marseille Université, 13288
37   Marseille, France
21   INRA, USC 1408 AFMB, F-13288 Marseille, France
22   Department of Biological Sciences, King Abdulaziz University, Saudi Arabia
23   Department of Entomology, University of Kentucky
24   Department of Entomology, Texas A&M University, College Station
25   Department of Genetics & Molecular Biology in Botany, Christian-Albrechts-University at Kiel
26   Division of Molecular Genetic Epidemiology, German Cancer Research Center
27   Department of Entomology, Nanjing Agricultural University
28   Center for Data-Intensive Biomedicine and Biotechnology, Skolkovo Institute of Science and Technology
29   Department of Biology, Illinois Institute of Technology
30   Department of Biology, University of Wisconsin-Oshkosh
31   Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory
32   USDA-ARS Center for Grain and Animal Health Research
33   Institute for Developmental Biology, University of Cologne
34   School of Life Sciences, University of Warwick, Gibbet Hill Campus

35 Department of Entomology, Max Planck Institute for Chemical Ecology, Jena, Germany
36 INRA, Aix-Marseille Université, UMR1163, Biodiversité et Biotechnologie Fongiques, Marseille, France
37 Department of Entomology, University of Illinois at Urbana-Champaign
38 Department of Entomology, Michigan State University
39 Department of Biology and School of Informatics and Computing, Indiana University
40 Center for Autoimmune Genomics and Etiology, Division of Biomedical Informatics and Division of
Developmental Biology, Cincinnati Children's Hospital Medical Center.
41 Department of Pediatrics, University of Cincinnati College of Medicine.
42 Department of Entomology, The Ohio State University
43 Center for Applied Plant Sciences, The Ohio State University
62
63 *Corresponding Author: Sean Schoville, University of Wisconsin-Madison, Department of
64 Entomology, 1630 Linden Drive, Madison, WI 53706; sean.schoville@wisc.edu; +1-608-262-
65 2956
66

Abstract

Background: The Colorado potato beetle, *Leptinotarsa decemlineata* Say, is one of the most challenging agricultural pests to manage. It has shown a spectacular ability to not only rapidly adapt to a broad range of solanaceaeous plants and variable climates during its global invasion, but, most notably, to rapidly evolve resistance to insecticides (over 50 different compounds in all major classes, in some cases within the first year of use). To examine evidence of rapid evolutionary change, and to understand the genetic basis of herbivory and insecticide resistance, we tested for structural and functional genomic changes relative to other arthropod species, using whole-genome sequencing, transcriptome sequencing, and a large community-driven annotation effort.

Results: We present a 140x coverage whole genome sequence from a single female *L. decemlineata*, with a reference gene set of 24,740 genes. Transposable elements comprise at least 17% of the genome, and are heavily represented in an analysis of rapidly evolving gene families compared to other Coleoptera. Population genetic analyses provide evidence of high levels of nucleotide diversity, local geographic structure, and recent population growth in pest populations, pointing to the availability of considerable standing genetic variation. These factors may play an important role in rapid evolutionary change. Adaptations to plant feeding are evident in gene expansions and differential expression of digestive enzymes (e.g. cysteine peptidase genes) in gut tissues, as well as expansions of the gustatory receptors for bitter tasting plants in the nightshade family, Solanaceae. Despite its notoriety for adapting to insecticides, *L. decemlineata* has a similar suite of genes involved in resistance (metabolic detoxification and cuticle penetration) compared to other beetles, although expansions in specific cytochrome P450 subfamilies are known to be associated with insecticide resistance. Finally, this beetle has

interesting duplications in RNAi genes that might be linked to its high sensitivity to RNAi and

could be important in the future development of gene targeted pesticides.

Conclusions: As a representative of one of the most evolutionarily diverse lineages, the *L.*

decemlineata genome will undoubtedly provide new opportunities for deeper understanding on

the ecology, evolution, and management of this species, as well as new opportunities to leverage

genomic technologies to understand the basis of a broad range of phenotypes and to develop

sustainable methods to control this widely successful pest.

Keywords: insects; evolution; pesticide resistance; herbivory; plant-insect interactions; pest

management; whole-genome sequence

Background

The Colorado potato beetle, *Leptinotarsa decemlineata* Say 1824 (Coleoptera: Chrysomelidae), is widely considered one of the world's most successful globally-invasive insect herbivores, with costs of ongoing management reaching tens of millions of dollars annually [1] and projected costs if unmanaged reaching billions of dollars [2]. This beetle was first identified as a pest in 1859 in the Midwestern United States, after it expanded from its native host plant, *Solanum rostratum* (Solanaceae), onto potato (*S. tuberosum*) [3]. As testimony to the difficulty in controlling *L. decemlineata*, the species has the dubious honor of starting the pesticide industry, when Paris Green (copper (II) acetoarsenite) was first applied to control it in the United States in 1864 [4]. *Leptinotarsa decemlineata* is now widely recognized for its ability to rapidly evolve resistance to insecticides, as well as a wide range of abiotic and biotic stresses [5], and for its global expansion across 16 million km^2 to cover the entire Northern Hemisphere within the 20th century [6]. Over the course of 150 years of research, *L. decemlineata* has been the subject in more than 9,700 publications (according to the Web of Science[TM] Core Collection of databases) ranging from molecular to organismal biology from the fields of agriculture, entomology, molecular biology, ecology, and evolution.

In order to be successful, *L. decemlineata* evolved to exploit novel host plants, to inhabit colder climates at higher latitudes [7–9], and to cope with a wide range of novel environmental conditions in agricultural landscapes [10,11]. Genetic data suggest the potato-feeding pest lineage directly descended from populations that feed on *S. rostratum* in the U.S. Great Plains [12]. This beetle subsequently expanded its range northwards, shifting its life history strategies to exploit even colder climates [7,8,13], and steadily colonized potato crops despite substantial geographical barriers [14]. *Leptinotarsa decemlineata* is an excellent model for understanding

pest evolution in agroecosystems because, despite its global spread, individuals disperse over

short distances and populations often exhibit strong genetic differentiation [15–17], providing an

opportunity to track the spread of populations and the emergence of novel phenotypes. The

development of genomic resources in *L. decemlineata* will provide an unparalleled opportunity

to investigate the molecular basis of traits such as climate adaptation, herbivory and host

expansion, and chemical detoxification. Perhaps most significantly, understanding its ability to

evolve rapidly would be a major step towards developing sustainable methods to control this

widely successful pest in agricultural settings.

Given that climate is thought to be the major factor in structuring the range limits of

species [18], the latitudinal expansion of *L. decemlineata,* spanning more than 40° latitude from

Mexico to northern potato-producing countries such as Canada and Russia [6], warrants further

investigation. Harsh winter climates are thought to present a major barrier for insect range

expansions, especially near the limits of a species' range [7,19]. To successfully overwinter in

temperate climates, beetles need to build up body mass, develop greater amounts of lipid storage,

have a low resting metabolism, and respond to photoperiodic keys by initiating diapause [20,21].

Although the beetle has been in Europe for less than 100 years, local populations have

demonstrating remarkably rapid evolution in life history traits linked to growth, diapause and

metabolism [8,13,20]. Understanding the genetic basis of these traits, particularly the role of

specific genes associated with metabolism, fatty acid synthesis, and diapause induction, could

provide important information about the mechanism of climate adaptation.

Although *Leptinotarsa decemlineata* has long-served as a model for the study of host

expansion and herbivory due to its rapid ability to host switch [17,22], a major outstanding

question is what genes and biological pathways are associated with herbivory in this species?

While >35,000 species of Chrysomelidae are well-known herbivores, most species feed on one

or a few host species within the same plant family [23]. Within *Leptinotarsa*, the majority of

species feed on plants within Solanaceae and Asteraceae, while *L. decemlineata* feeds

exclusively on solanaceous species [24]. It has achieved the broadest host range amongst its

congeners (including, but not limited to: buffalobur (*S. rostratum*), potato (*S. tuberosum*),

eggplant (*S. melongena),* silverleaf nightshade (*S. elaeagnifolium*), horsenettle (*S. carolinense*),

bittersweet nightshade (*S. dulcamara),* tomato *(S. lycopersicum),* and tobacco (*Nicotiana*

tabacum)) [17,22,25], and exhibits geographical variation in the use of locally abundant *Solanum*

species [26].

Another major question is what are the genes that underlie the beetle's remarkable

capacity to detoxify plant secondary compounds and are these the same biological pathways used

to detoxify insecticidal compounds [27]? Solanaceous plants are considered highly toxic to a

wide range of insect herbivore species [28], because they contain steroidal alkaloids and

glycoalkaloids, nitrogen-containing compounds that are toxic to a wide range of organisms,

including bacteria, fungi, humans, and insects [29], as well as glandular trichomes that contain

additional toxic compounds [30]. In response to beetle feeding, potato plants upregulate

pathways associated with terpenoid, alkaloid, and phenylpropanoid biosynthesis, as well as a

range of protease inhibitors [31]. A complex of digestive cysteine proteases is known to underlie

L. decemlineata's ability to respond to potato-induced defenses [32,33]. There is evidence that

larvae excrete [34] and perhaps even sequester toxic plant-based compounds in the hemolymph

[35,36]. Physiological mechanisms involved in detoxifying plant compounds, as well as other

xenobiotics, have been proposed to underlie pesticide resistance [27]. To date, while cornerstone

of *L. decemlineata* management has been the use of insecticides, the beetle has evolved

resistance to over 50 compounds and all of the major classes of insecticides. Some of these

chemicals have even failed to control *L. decemlineata* within the first year of release [10], and

notably, regional populations of *L. decemlineata* have demonstrated the ability to independently

evolve resistance to pesticides and to do so at different rates [37]. Previous studies have

identified target site mutations in resistance phenotypes and a wide range of genes involved in

metabolic detoxification, including carboxylesterase genes, cytochrome P450s, and glutathione

S-transferase genes [38–42].

To examine evidence of rapid evolutionary change underlying *L. decemlineata*'s

extraordinary success utilizing novel host plants, climates, and detoxifying insecticides, we

evaluated structural and functional genomic changes relative to other beetle species, using

whole-genome sequencing, transcriptome sequencing, and a large community-driven biocuration

effort to improve predicted gene annotations. We compared the size of gene families associated

with particular traits against existing available genomes from related species, particularly those

sequenced by the i5k project (http://i5k.github.io), an initiative to sequence 5,000 species of

Arthropods. While efforts have been made to understand the genetic basis of phenotypes in *L.*

decemlineata (for example, pesticide resistance) [32,43,44], previous work has been limited to

candidate gene approaches rather than comparative genomics. Genomic data can not only

illuminate the genetic architecture of a number of phenotypic traits that enable *L. decemlineata*

to continue to be an agricultural pest, but can also be used to identify new gene targets for

control measures. For example, recent efforts have been made to develop RNAi-based pesticides

targeting critical metabolic pathways in *L. decemlineata* [41,45,46]. With the extensive wealth of

biological knowledge and a newly-released genome, this beetle is well-positioned to be a model

system for agricultural pest genomics and the study of rapid evolution.

Results and Discussion

Genome Assembly, Annotation and Assessment

A single female *L. decemlineata* from Long Island, NY, USA, a population known to be

resistant to a wide range of insecticides [47,48], was sequenced at ~140x coverage and

assembled with ALLPATHS [49] followed by assembly improvement with ATLAS

(https://www.hgsc.bcm.edu/software/). The average coleopteran genome size is 760 Mb (ranging

from 160-5,020 Mb [50]), while most of the beetle genome assemblies have been smaller (mean

assembly size 286 Mb, range 160-710 Mb) [51-55]. The draft genome assembly of *L.*

decemlineata is 1.17 Gb and consists of 24,393 scaffolds, with a N50 of 414 kb and a contig N50

of 4.9 kb. This assembly is more than twice the estimated genome size of 460 Mb [56], with the

presence of gaps comprising 492 Mb, or 42%, of the assembly. As this size might be driven by

underlying heterozygosity, we also performed scaffolding with REDUNDANS [57], which

reduced the assembly size to 642 Mb, with gaps reduced to 1.3% of the assembly. However, the

REDUNDANS assembly increased the contig N50 to 47.4 kb, the number of scaffolds increased

to 90,205 and the N50 declined to 139 kb. For all downstream analyses, the ALLPATHS

assembly was used due to its increased scaffold length and reduced number of scaffolds.

The number of genes in the *L. decemlineata* genome predicted based on automated

annotation using MAKER was 24,671 gene transcripts, with 93,782 predicted exons, which

surpasses the 13,526-22,253 gene models reported in other beetle genome projects (**Figure 1**)

[51-55]. This may be in part due to fragmentation of the genome, which is known to inflate gene

number estimates [58]. To improve our gene models, we manually annotated genes using expert

opinion and additional mRNA resources (see **Supplementary Methods** in **Additional File 1** for

more details). A total of 1,364 genes were manually curated and merged with the unedited

MAKER annotations to produce an official gene set (OGS v1.0) of 24,850 transcripts, comprised

of 94,859 exons. A total of 12 models were curated as pseudogenes. A total of 1,237 putative

transcription factors (TFs) were identified in the *L. decemlineata* predicted proteome (**Figure 2**).

The predicted number of TFs is similar to some beetles, such as *Anoplophora glabripennis*

(1,397) and *Hypothenemus hampei* (1,148), but substantially greater than others, such as

Tribolium castaneum (788), *Nicrophorus vespilloides* (744), and *Dendroctonus ponderosae*

(683) [51-55].

We assessed the completeness of the assembly and OGS computationally using

benchmarking sets of universal single-copy orthologs (BUSCOs) based on 35 holometabolous

insect genomes [59], as well as manually by assessing the completeness and co-localization of

the homeodomain transcription factor gene clusters. Using the reference set of 2,442 BUSCOs,

the genome and OGS were 93.0% and 71.8% complete, respectively. We found an additional

4.1% and 17.9% of the BUSCOs present but fragmented in the genome and OGS, respectively.

For the highly conserved Hox and Iroquois Complex (Iro-C) clusters, we located and annotated

complete gene models for all 12 expected orthologs, but these were split across six different

scaffolds (**Additional File 1 Supplementary Figure 1S** and **Table 1S**). All linked Hox genes

occurred in the expected order and with the expected, shared transcriptional orientation,

suggesting that the current draft assembly was correct but incomplete (see also **Supplementary**

Figures 2S and **3S**). Assuming direct concatenation of scaffolds, the Hox cluster would span a

region of 3.7 Mb, similar to the estimated 3.5 Mb Hox cluster of *A. glabripennis* [54]. While

otherwise highly conserved with *A. glabripennis*, we found a tandem duplication for *Hox3/zen*

and an Antennapedia-class (ANTP-class) homeobox gene with no clear ortholog in other

arthropods.

Gene Annotation, Gene Family Evolution and Differential Expression

We estimated a phylogeny among six coleopteran genomes (*A. glabripennis*, *Agrilus*

planipennis, *D. ponderosae*, *L. decemlineata*, *Onthophagus taurus*, and *T. castaneum*;

unpublished genomes available at http://i5k.github.io/) [51,52,54] using a conserved set of single

copy orthologs and compared the official gene set of each species to understand how gene

families evolved the branch representing Chysomelidae. *Leptinotarsa decemlineata* and *A.*

glabripennis (Cerambycidae) are sister taxa (**Figure 1**), as expected for members of the same

superfamily Chrysomeloidea. We found 166 rapidly evolving gene families along the *L.*

decemlineata lineage (1.4% of 11,598), 142 of which are rapid expansions and the remaining 24

rapid contractions **(Table 1)**. Among all branches of our coleopteran phylogeny, *L. decemlineata*

has the highest average expansion rate (0.203 genes per million years), the highest number of

genes gained, and the greatest number of rapidly evolving gene families.

Examination of the functional classification of rapidly evolving families in *L.*

decemlineata (**Supplementary Tables 2S** and **3S**) indicates that a subset of families are clearly

associated with herbivory. The peptidases, comprising several gene families that play a major

role in plant digestion [32,60], displayed a significant expansion in genes (OrthoDB family Id:

EOG8JDKNM, EOG8GTNV8, EOG8DRCSN, EOG8GTNT0, EOG8K0T47, EOG88973C,

EOG854CDR, EOG8Z91BB, EOG80306V, EOG8CZF0X, EOG80P6ND, ,EOG8BCH22,

EOG8ZKN28, EOG8F4VSG, EOG80306V, EOG8BCH22, EOG8ZKN28, EOG8F4VSG).

While olfactory receptor gene families have rapidly contracted (EOG8Q5C4Z, EOG8RZ1DX),

subfamilies of odorant binding proteins and gustatory receptors have grown (see Sensory

Ecology section below). The expansion of gustatory receptor subfamilies are associated with

bitter receptors, likely reflecting host plant detection of nightshades (Solanaceae). Some gene

families associated with plant detoxification and insecticide resistance have rapidly expanded

(Glutathione S-transferases: EOG85TG3K, EOG85F05D, EOG8BCH22; UDP-

glycosyltransferase: EOG8BCH22; cuticle proteins: EOG8QJV4S, EOG8DNHJQ; ABC

transporters: EOG83N9TJ), whereas others have contracted (Cytochrome P450s: EOG83N9TJ).

Finally, gene families associated with immune defense (fibrinogen: EOG8DNHJQ;

Immunoglobulin: EOG8Q87B6, EOG8S7N55, EOG854CDX, EOG8KSRZK, EOG8CNT5M)

exhibit expansions that may be linked to defense against pathogens and parasitoids that

commonly attack exposed herbivores [61]. A substantial proportion of the rapidly evolving gene

families include proteins with transposable element domains (25.3%), while other important

functional groups include DNA and protein binding (including many transcription factors),

nuclease activity, protein processing, and cellular transport.

Diversification of transcription factor (TF) families potentially signals greater complexity

of gene regulation, including enhanced cell specificity and refined spatiotemporal signaling [62].

Notably, several TF families are substantially expanded in *L. decemlineata*, including HTH_psq

(194 genes vs. a mean of only 24 across the insects shown in **Figure 2**), MADF (152 vs. 54), and

THAP (65 vs. 41). Two of these TF families, HTH_psq and THAP, are DNA binding domains in

transposons. Of the 1,237 *L. decemlineata* TFs, we could infer DNA binding motifs for 189

(15%) (**Supplementary Table 4S**), mostly based on DNA binding specificity data from

Drosophila melanogaster (124 TFs), but also from species as distant as human (45 TFs) and

mouse (11 TFs). Motifs were inferred for a substantial proportion of the TFs in the largest TF

families, including Homeodomain (59 of 90, 66%), bHLH (34 of 46, 74%), and Forkhead box

(14 of 16, 88%). We could only infer a small number of C2H2 zinc finger motifs (21 of 439,

~5%), which is expected as these sequences evolve quickly by shuffling zinc finger arrays,

resulting in largely dissimilar DNA-binding domains across metazoans [63]. Collectively, the

almost 200 inferred DNA binding motifs for *L. decemlineata* TFs provide a unique resource to

begin unraveling gene regulatory networks in this organism.

To identify genes active in mid-gut tissues, life-stages, and sex differences, we examined

differential transcript expression levels using RNA sequencing data. Comparison of significantly

differentially expressed genes with >100-fold change, after Bonferroni correction, indicated

higher expression of digestive enzymes (proteases, peptidases, dehydrogenases and transporters)

in mid-gut versus whole larval tissues, while cuticular proteins were largely expressed at lower

levels (**Figure 3A, Supplementary Table 5S**). Comparison of an adult male and female showed

higher expression of testes and sperm related genes in males, while genes involved in egg

production and sterol biosynthesis are more highly expressed in females (**Figure 3B,**

Supplementary Table 6S). Comparisons of larvae to both an adult male (**Figure 3C,**

Supplementary Table 7S) and an adult female (**Figure 3D, Supplementary Table 8S**) showed

higher expression of larval-specific cuticle proteins, and lower expression of odorant binding and

chemosensory proteins. The adults, both drawn from a pesticide resistant population, showed

higher constitutive expression of cytochrome p450 genes compared to the larval population,

which is consistent with the results from previous studies of neonicotinoid resistance in this

population [48].

Transposable Elements

Transposable elements (TEs) are ubiquitous mobile elements within most eukaryotic

genomes and play critical roles in both genome architecture and the generation of genetic

variation [64]. Through insertional mutagenesis and recombination, TEs are a major contributor

to the generation of novel mutations within the genome [65]. TEs can generate a number of

different types of mutations, including mutations resulting in exonization, modulating alternative

splicing, disrupting or silencing genes, or altering cis-acting sequences such as transcription

factor binding sites [66]. Thus increasingly, TE activities are thought to generate much of the

genetic diversity that contributes to rapid evolution [67‑69]. For instance, an estimated 50-80%

of all mutation events in *D. melanogaster* are caused by TEs [70]. We found that at least 17% of

the *L. decemlineata* genome consists of TEs (**Supplementary Table 9S**), which is greater than

the 6% found in *T. castaneum* [51], but less than some Lepidoptera (35% in *Bombyx mori* and

25% in *Heliconius melpomene*) [71,72]. LINEs were the largest TE class, comprising ~10% of

the genome, while SINEs were not detected. Curation of the TE models with intact protein

domains resulted in 334 current models of potentially active TEs, meaning that these TEs are

capable of transposition and excision. Within the group of active TEs, we found 191 LINEs, 99

DNA elements, 38 LTRs, and 5 Helitrons. Given that TEs have been associated with the ability

of species to rapidly adapt to novel selection pressures [73‑75], particularly via alterations of

gene expression patterns in neighboring genomic regions, we scanned gene rich regions (1 kb

neighborhood size) for active TE elements. Genes with active neighboring TEs have functions

that include transport, protein digestion, diapause, and metabolic detoxification (**Supplementary**

Table 10S). Because TE elements have been implicated in conferring insecticide resistance in

other insects [76], future work should investigate the role of these TE insertions on rapid

evolutionary changes within pest populations of *L. decemlineata*.

Population Genetic Variation and Invasion History

To understand the propensity for *L. decemlineata* pest populations to rapidly evolve across a range of environmental conditions, we examined geographical patterns of genomic variability and the evolutionary history of *L. decemlineata*. High levels of standing variation are one mechanism for rapid evolutionary change [77]. We identified 1.34 million biallelic single nucleotide polymorphisms (SNPs) from pooled RNAseq datasets, or roughly 1 variable site for every 22 base pairs of coding DNA. This rate of polymorphism is exceptionally high when compared to vertebrates (e.g. ~1 per kb in humans, or ~1 per 500 bp in chickens) [78,79], and is 8-fold higher than other beetles (1 in 168 for *D. ponderosae* and 1 in 176 bp for *O. taurus*) [52,80] and 2 to 5-fold higher than other dipterans (1 in 54 bp for *D. melanogaster* and 1 in 125 bp for *Anopheles gambiae*) [78,81]. It is likely that these values simply scale with effective population size, although the dipterans, with the largest known population sizes, have reduced variation due to widespread selective sweeps and genetic bottlenecks [82].

Evolutionary relationships and the amount of genetic drift among Midwestern USA, Northeastern USA, and European *L. decemlineata* populations were estimated based on genome-wide allele frequency differences using a population graph. A substantial amount of local genetic structure and high genetic drift is evident among all populations, although both the reference lab strain from New Jersey and European populations appear to have undergone more substantial drift, suggestive of strong inbreeding (**Figure 4**). The allele frequency spectrum was calculated for populations in Wisconsin, Michigan, and Europe to estimate the population genetic parameter θ, or the product of the mutation rate and the ancestral effective population size, and the ratio of contemporary to ancestral population size in models that allowed for single or multiple episodes

of population size change. Estimates of ancestral θ are much higher for Wisconsin ($\theta = 12595$)

and Michigan ($\theta = 93956$) than Europe ($\theta = 3.07$; **Supplementary Table 11S**), providing

support for a single introduction into Europe following a large genetic bottleneck [15]. In all

three populations, a model of population size growth is supported, in agreement with historical

accounts of the beetles expanding from the Great Plains into the Midwestern U.S. and Europe

[3,15], but the dynamics of each population appear independent, with the population from

Michigan apparently undergoing a very recent decline in contemporary population size (the ratio

of contemporary to ancient population size is 0.066, compared to 3.3 and 2.1 in Wisconsin and Europe,

respectively).

Sensory Ecology and Host Plant Detection

To interact with their environment, insects have evolved neurosensory organs to detect

environmental signals, including tactile, auditory, chemical and visual cues [83]. We examined

neural receptors, olfactory genes, and light sensory (opsin) genes to understand the sensory

ecology and host-plant specializations of *L. decemlineata*.

We found high sequence similarity in the neuroreceptors of *L. decemlineata* compared to

other Coleoptera. The transient receptor potential (TRP) channels are permeable transmembrane

proteins that respond to temperature, touch, pain, osmolarity, pheromones, taste, hearing, smell

and visual cues of insects [84,85]. In most insect genomes, there are typically 13-14 TRP genes

located in insect stretch receptor cells and several are targeted by commercial insecticides [86].

We found 12 TRP genes present in the *L. decemlineata* genome, including the two TRPs

(Nanchung and Inactive) that are targeted by commercial insecticides, representing a complete

set of one-to-one orthologs with *T. castaneum.* Similarly, the 20 known amine neurotransmitter

receptors in *T. castaneum* are present as one-to-one orthologs in *L. decemlineata* [87,88]. Amine

receptors are G-protein-coupled receptors that interact with biogenic amines, such as

octopamine, dopamine and serotonin. These neuroactive substances regulate behavioral and

physiological traits in animals by acting as neurotransmitters, neuromodulators and

neurohormones in the central and peripheral nervous systems [89].

The majority of phytophagous insects are restricted to feeding on several plant species

within a genus, or at least restricted to a particular plant family [90]. Thus, to find their host

plants within heterogeneous landscapes, insect herbivores detect volatile organic compounds

through olfaction, which utilizes several families of chemosensory gene families, such as the

odorant binding proteins (OBPs), odorant receptors (ORs), gustatory receptors (GRs), and

ionotropic receptors (IRs) [91]. OBPs directly bind with volatile organic compounds emitted

from host plants and transport the ligands across the sensillar lymph to activate the membrane-

bound ORs in the dendrites of the olfactory sensory neurons [92]. The ORs and GRs are 7-

transmembrane proteins related within a superfamily [93] of ligand-gated ion channels [94]. The

ionotropic receptors are related to ionotropic glutamate receptors and function in both smell and

taste [95]. These four gene families are commonly large in insect genomes, consisting of tens to

hundreds of members.

We compared the number of genes found in *L. decemlineata* in the four chemosensory

gene families to *T. castaneum* and *A. glabripennis* (**Supplementary Table 12S**, as well as

Tables 13S-15S for details). While the OBP family is slightly enlarged, the three receptor

families are considerably smaller in *L. decemlineata* than in either *A. glabripennis* or *T.*

castaneum. However, each beetle species exhibits species-specific gene subfamily expansions

(**Supplementary Figures 4S-7S**); in particular, members of the GR family related to bitter taste

are expanded in *L. decemlineata* relative to *A. glabripennis* and other beetles. Among the OBPs,

we identified a major *L. decemlineata*-specific expansion of proteins belonging to the Minus-C

class (OBPs that have lost two of their six conserved cysteine residues) that appear unrelated to

the 'traditional' Minus-C subfamily in Coleoptera, indicating that coleopteran OBPs have lost

cysteines on at least two occasions.

To understand the visual acuity of *L. decemlineata*, we examined the G-protein-coupled

transmembrane receptor opsin gene family. We found five opsins, three of which are members of

rhabdomeric opsin (R-opsin) subfamilies expressed in the retina of insects [51,96]. Specifically,

the *L. decemlineata* genome contains one member of the long wavelength-sensitive R-opsin and

two short wavelength UV-sensitive R-opsins. The latter were found to be closely linked in a

range of less than 20,000 bp, suggestive of recent tandem gene duplication. Overall, the

recovered repertoire of retinally-expressed opsins in *L. decemlineata* [97] is consistent with the

beetle's attraction to yellow light and to the yellow flowers of its ancestral host plant, *S.*

rostratum [98,99], and is consistent with the beetle's sensitivity in the UV- and LW-range [100].

In addition, we found a member of the Rh7 R-opsin subfamily, which is broadly conserved in

insects including other beetle species (*A. glabripennis*), although it is missing from *T.*

castaneum. Finally, *L. decemlineata* has a single ortholog of the c-opsin subfamily shared with *T.*

castaneum, which is absent in *A. glabripennis* and has an unclear role in photoreception [101].

Host Plant Utilization

Protein digestion

Insect herbivores are fundamentally limited by nitrogen availability [102], and thus need

to efficiently break down plant proteins in order to survive and develop on host plants [103].

Leptinotarsa decemlineata has serine and cysteine digestive peptidases (coined "intestains")

[32,104,105], as well as aspartic and metallo peptidases [33], for protein digestion. For the vast

majority of plant-eating beetles (the infraorder Cucujiformia, which includes Chrysomelidae),

cysteine peptidases contribute most strongly to proteolytic activity in the gut [105,106]. In

response to herbivory, plants produce a wide range of proteinase inhibitors to prevent insect

herbivores from digesting plant proteins [103,105,107]. Coleopteran peptidases are differentially

susceptible to plant peptidase inhibitors, and our annotation results suggest that gene duplication

and selection for inhibitor insensitive genotypes may have contributed to the success of leaf-

feeding beetles (Chrysomelidae) on different plants. We found that gene expansion of cysteine

cathepsins from the C1 family in *L. decemlineata* correlates with the acquisition of greater

digestive function by this group of peptidases, which is supported by gene expression activity of

these genes in mid-gut tissue (**Figure 3A, Figure 5**). The gene expansion may be explained by

an evolutionary arms race between insects and plants that favors insects with a variety of

digestive peptidases in order to resist plant peptidase inhibitors [108,109] and allows for

functional specialization [110].

Cysteine peptidases of the C1 family were represented by more than 50 genes separated

into four groups with different structure and functional characteristics (**Supplementary Table**

16S): cathepsin L subfamily, cathepsin B subfamily, TINAL-like genes, and cysteine peptidase

inhibitor domains (CPIDs). Cathepsin L subfamily cysteine peptidases are endopeptidases [111]

that can be distinguished by the cathepsin propeptide inhibitor domain I29 (pfam08246)

[112,113]. Within the cathepsin L subfamily, we found sequences that were similar to classical

cathepsin L, cathepsin F, and cathepsin O. However, there were 28 additional predicted

peptidases of this subfamily that could not be assigned to any of the "classical" cathepsin types,

and most of these were grouped into two gene expansions (uL1 and uL2) according to their

phylogenetic and structural characteristics. Cathepsin B subfamily cysteine peptidases are

distinguished by the specific peptidase family C1 propeptide domain (pfam08127). Within the

cathepsin B subfamily, there was one gene corresponding to typical cathepsin B peptidases and

14 cathepsin B-like genes. According to the structure of the occluding loop, only the typical

cathepsin B may have typical endo- and exopeptidase activities, while a large proportion of

cathepsin B-like peptidases presumably possesses only endopeptidase activity due to the absence

of a His-His active subsite in the occluding loop, which is responsible for exopeptidase activity

[111]. Only one gene corresponding to a TINAL-like-protein was present, which has a domain

similar to cathepsin B in the C-terminus, but lacks peptidase activity due to the replacement of

the active site Cys residue with Ser [114]. Cysteine peptidase inhibitor domain (CPID) genes

encode the I29 domain of cysteine peptidases without the mature peptidase domain. Within the

CPID group, there were seven short inhibitor genes that lack the enzymatic portion of the

protein. A similar trend of "stand-alone inhibitors" has been observed in other insects, such as *B.*

mori [115]. These CPID genes may be involved in the regulation of cysteine peptidases. We note

that we found multiple fragments of cysteine peptidase genes, suggesting that the current list of

L. decemlineata genes may be incomplete. Comparison of these findings with previous data on

L. decemlineata cysteine peptidases [116] demonstrates that intestains correspond to several

peptidase genes from the uL1 and uL2 groups (**Supplementary Table 16S**). These data, as well

as literature for Tenebrionidae beetles [117], suggest that intensive gene expansion is typical for

peptidases that are involved in digestion.

We also found a high number of digestion-related serine peptidase genes in the *L.*

decemlineata genome (**Supplementary Table 17S**), but they contribute only a small proportion

of the beetle's total gut proteolytic activity [32]. Of the 31 identified serine peptidase genes and

fragments, we annotated 16 as trypsin-like peptidases and 15 as chymotrypsin-like peptidases.

For four chymotrypsin-like and one trypsin-like peptidase, we identified only short fragments.

All complete (and near-complete) sequences have distinctive S1A peptidase subfamily motifs, a

conserved catalytic triad, conserved sequence residues such as the "CWC" sequence and

cysteines that form disulfide bonds in the chymotrypsin protease fold. The number of serine

peptidases was higher than expected based upon the number of previously identified EST clones

[32], but lower than the number of chymotrypsin and trypsin genes in the *T. castaneum* genome.

Carbohydrate digestion

Carbohydrates are the other category of essential nutrients for *L. decemlineata*. The

enzymes that assemble and degrade oligo- and polysaccharides, collectively termed

Carbohydrate active enzymes (CAZy), are categorized into five major classes: glycoside

hydrolases (GH), polysaccharide lyases, carbohydrate esterases (CE), glycosyltransferases (GT)

and various auxiliary oxidative enzymes [118]. Due to the many different roles of carbohydrates,

the CAZy family profile of an organism can provide insight into "glycobiological potential" and,

in particular, mechanisms of carbon acquisition [119]. We identified 182 GHs assigned to 25

families, 181 GTs assigned to 41 families, and two CEs assigned to two families in *L.*

decemlineata; additionally, 99 carbohydrate-binding modules (which are non-catalytic modules

associated with the above enzyme classes) were present and assigned to 9 families

(**Supplementary Table 18S**; the list of CAZy genes is presented in **Supplementary Table 19S**).

We found that *L. decemlineata* has three families of genes associated with plant cell wall

carbohydrate digestion (GH28, GH45 and GH48) that commonly contain enzymes that target

pectin (GH28) and cellulose (GH45 and GH48), the major structural components of leaves [120].

We found evidence of massive gene duplications in the GH28 family (14 genes) and GH45

family (11 genes, plus one additional splicing variant), whereas GH48 is represented by only

three genes in the genome [121]. Overall, the genome of *L. decemlineata* shows a CAZy profile

adapted to metabolize pectin and cellulose contained in leaf cell walls. The absence of specific

members of the families GH43 (α-L-arabinofuranosidases) and GH78 (α-L-rhamnosidases)

suggests that *L. decemlineata* can break down homogalacturonan, but not substituted

galacturonans such as rhamnogalacturonan I or II [120]. The acquisition of these plant cell wall

degrading enzymes has been linked to horizontal transfer in the leaf beetles and other

phytophagous beetles [120], with strong phylogenetic evidence supporting the transfer of GH28

genes from a fungal donor (Pezizomycotina) in *L. decemlineata*, as well as in the beetles *D.*

ponderosae and *Hypothenemus hampei*, but a novel fungal donor in the more closely related

cerambycid beetle *A. glabripennis* [54] and a bacterial donor in the weevil *Callosobruchus*

maculatus [120].

Insecticide Resistance

To understand the functional genomic properties of insecticide resistance, we examined

genes important to neuromuscular target site sensitivity, tissue penetration, and prominent gene

families involved in Phase I, II, and III metabolic detoxification of xenobiotics [122]. These

include the cation-gated nicotinic acetylcholine receptors (nAChRs), the γ-amino butyric acid

(GABA)-gated anion channels and the histamine-gated chloride channels (HisCls), cuticular

proteins, cytochrome P450 monooxygenases (CYPs), and the Glutathione S-transferases (GSTs).

Many of the major classes of insecticides (organochlorides, organophosphates,

carbamates, pyrethroids, neonicotinoids and ryanoids) disrupt the nervous system (particularly

ion channel signaling), causing paralysis and death [123]. Resistance to insecticides can come

from point mutations that reduce the affinity of insecticidal toxins to ligand-gated ion

superfamily genes [124]. The cys-loop ligand-gated ion channel gene superfamily is comprised

of receptors involved in mediating synaptic ion flow during neurotransmission [125]. A total of

22 cys-loop ligand-gated ion channels were identified in the *L. decemlineata* genome in numbers

similar to those observed in other insects [126], including 12 nAChRs, three GABA receptors,

and two HisCls (**Supplementary Figure 8S**). The GABA-gated chloride channel homolog of the

Resistance to dieldrin (*Rdl*) gene of *D. melanogaster* was examined due to its role in resistance

to dieldrin and other cyclodienes in Diptera [127]. The coding sequence is organized into 10

exons (compared to nine in *D. melanogaster*) on a single scaffold, with duplications of the third

and sixth exon (**Supplementary Figure 9S**). Alternative splicing of these two exons encodes for

four different polypeptides in *D. melanogaster* [128,129], and as the splice junctions are present

in *L. decemlineata*, we expect the same diversity of *Rdl*. The point mutations in the

transmembrane regions TM2 and TM3 of *Rdl* are known to cause insecticide resistance in

Diptera [124,127], but were not observed in *L. decemlineata*.

Cuticle genes have been implicated in imidacloprid resistant *L. decemlineata* [130] and at

least one has been shown to have phenotypic effects on resistance traits following RNAi

knockdown [131]. A total of 163 putative cuticle protein genes were identified and assigned to

one of seven families (CPR, CPAP1, CPAP3, CPF, CPCFC, CPLCG, and TWDL)

(**Supplemental Figure 10S** and **Table 20S**). Similar to other insects, the CPR family, with the

RR-1 (soft cuticle), RR-2 (hard cuticle), and unclassifiable types, constituted the largest group of

cuticle protein genes (132) in the *L. decemlineata* genome. While the number of genes in *L.*

decemlineata is slightly higher than in *T. castaneum* (110), it is similar to *D. melanogaster* (137)

[132]. Numbers in the CPAP1, CPAP3, CPF, and TWDL families were similar to other insects,

and notably no genes with the conserved sequences for CPLCA were detected in *L.*

decemlineata, although they are found in other Coleoptera.

A total of 89 CYP (P450) genes were identified in the *L. decemlineata* genome, an

overall decrease relative to *T. castaneum* (143 genes). Due to their role in insecticide resistance

in *L. decemlineata* and other insects [38,48,133], we examined the CYP6 and CYP12 families in

particular. Relative to *T. castaneum*, we observed reductions in the CYP6BQ, CYP4BN, and

CYP4Q subfamilies. However, five new subfamilies (CYP6BJ, CYP6BU, CYP6E, CYP6F and

CYP6K) were identified in *L. decemlineata* that were absent in *T. castaneum*, and the CYP12

family contains three genes as opposed to one gene in *T. castaneum* (CYP12h1). We found

several additional CYP genes not present in *T. castaneum*, including CYP413A1, CYP421A1,

CYP4V2, CYP12J and CYP12J4. Genes in CYP4, CYP6, and CYP9 are known to be involved

in detoxification of plant allelochemicals as well as resistance to pesticides through their

constitutive overexpression and/or inducible expression in imidacloprid resistant *L. decemlineata*

[48,130].

GSTs have been implicated in resistance to organophosphate, organochlorine and

pyrethroid insecticides [134] and are responsive to insecticide treatments in *L. decemlineata*

[130,135]. A total of 27 GSTs were present in the *L. decemlineata* genome, and while they

represent an expansion relative to *A. glabripennis*, all have corresponding homologs in *T.*

castaneum. The cytosolic GSTs include the epsilon (11 genes), delta (5 genes), omega (4 genes),

theta (2 genes) and sigma (3 genes) families, while two GSTs are microsomal (**Supplementary**

Figure 11S). Several GST-like genes present in the *L. decemlineata* genome represent the Z

class previously identified using transcriptome data [135].

Pest Control via the RNAi Pathway

RNA interference (RNAi) is the process by which small non-coding RNAs trigger

sequence-specific gene silencing [136]. RNAi plays a role in various cellular processes,

including the protection against viruses and mobile genetic elements such as transposons, gene

regulation and cellular development [137]. There are several classes of interfering RNAs, with

three important varieties including small interfering RNAs (siRNAs), micro RNAs (miRNAs)

and piwi-interacting RNAs (piRNAs). The application of exogenous double stranded RNA

(dsRNA) has been exploited as a tool to suppress gene expression for functional genetic studies

[138,139] and for pest control [41,140].

We annotated a total of 49 genes associated with RNA interference, most of them were

found on a single scaffold. All genes from the core RNAi machinery (from all three major RNAi

classes) were present in *L. decemlineata*, including fifteen genes encoding components of the

RNA Induced Silencing Complex (RISC) and genes known to be involved in double-stranded

RNA uptake, transport, and degradation **(Supplementary Table 21S).** A complete gene model

was annotated for *R2D2*, an essential component of the siRNA pathway that interacts with *dicer-*

2 to load siRNAs into the RISC, and not previously detected in the transcriptome of the *L.*

decemlineata mid-gut [141]. The core components of the small interfering RNA (siRNA)

pathway were duplicated, including *dicer-2*, an RNase III enzyme that cleaves dsRNAs and pre-

miRNAs into siRNAs and miRNAs respectively [142,143]. The *dicer-2a* and *dicer-2b* CDS have

60% nucleotide identity to each other, and 56% and 54% identity to the *T. castaneum dicer-2*

homolog, respectively. The *argonaute-2* gene, which plays a key role in RISC by binding small

non-coding RNAs, was also duplicated. A detailed analysis of these genes will be necessary to

determine if the duplications provide functional redundancy. The duplication of genes in the

siRNA pathway may play a role in the high sensitivity of *L. decemlineata* to RNAi knockdown

[144] and could benefit future efforts to develop RNAi as a pest management technology.

Conclusion

The whole-genome sequence of *L. decemlineata*, provides novel insights into one of the

most diverse animal taxa, Chrysomelidae. It is amongst the largest beetle genomes sequenced to

date, with a minimum assembly size of 640 Mb (ranging up to 1.17 Gb) and 24,740 genes. The

genome size is driven in part by a large number of transposable element families, which

comprise at least 17% of the genome and appear to be rapidly expanding relative to other beetles.

Population genetic analyses suggest high levels of nucleotide diversity, local geographic

structure, and evidence of recent population growth, which helps to explain how *L. decemlineata*

rapidly evolves to exploit novel host plants, climate space, and overcome a range of pest

management practices (including a large and diverse number of insecticides). Digestive

enzymes, in particular the cysteine peptidases and carbohydrate-active enzymes, show evidence

of gene expansion and elevated expression in gut tissues, suggesting the diversity of the genes is

a key trait in the beetle's phytophagous lifestyle. Additionally, expansions of the gustatory

receptor subfamily for bitter tasting might be a key adaptation to exploiting hosts in the

nightshade family, Solanaceae, while expansions of novel subfamilies of CYP and GST proteins

are consistent with rapid, lineage-specific turnover of genes implicated in *L. decemlineata*'s

capacity for insecticide resistance. Finally, *L. decemlineata* has interesting duplications in RNAi

genes that might increase its sensitivity to RNAi and provide a promising new avenue for

pesticide development. The *L. decemlineata* genome promises new opportunities to investigate

the ecology, evolution, and management of this species, and to leverage genomic technologies in

developing sustainable methods of pest control.

Materials and Methods

Genome Characteristics and Sampling of DNA and RNA

Previous cytological work determined that *L. decemlineata* is diploid and consists of 34

autosomes plus an XO system in males, or an XX system in females [145]. Twelve

chromosomes are submetacentric, while three are acrocentric and two are metacentric, although

one chromosome is heteromorphic (acrocentric and/or metacentric) in pest populations [146].

The genome size has been estimated with Feulgen densitometry at 0.46 pg, or approximately 460

Mb [47]. To generate a reference genome sequence, DNA was obtained from a single adult

female, sampled from an imidacloprid resistant strain developed from insects collected from a

potato field in Long Island, NY. Additionally, whole-body RNA was extracted for one male and

one female from the same imidacloprid resistant strain. Raw RNAseq reads for 8 different

populations were obtained from previous experiments: two Wisconsin populations

(PRJNA297027) [130], a Michigan population (PRJNA400685), a lab strain originating from a

New Jersey field population (PRJNA275431), and three samples from European populations

(PRJNA79581 and PRJNA236637) [43,121]. All RNAseq data came from pooled populations or

were combined into a population sample from individual reads. In addition, RNA samples of an

adult male and female from the same New Jersey population were sequenced separately using

Illumina HiSeq 2000 as 100 bp paired end reads (deposited in the GenBank/EMBL/DDBJ

database, PRJNA275662), and three samples from the mid-gut of 4th-instar larvae were

sequenced using SOLiD 5500 Genetic Analyzer as 50 bp single end reads (PRJNA400633).

Genome Sequencing, Assembly, Annotation and Assessment

Four Illumina sequencing libraries were prepared, with insert sizes of 180 bp, 500 bp, 3

kb, and 8 kb, and sequenced with 100 bp paired-end reads on the Illumina HiSeq 2000 platform

at estimated 40x coverage, except for the 8kb library, which was sequenced at estimated 20x

coverage. ALLPATHS-LG v35218 [49] was used to assemble reads. Two approaches were used

to scaffold contigs and close gaps in the genome assembly. The reference genome used in

downstream analyses was generated with ATLAS-LINK v1.0 and ATLAS GAP-FILL v2.2

(https://www.hgsc.bcm.edu/software/). In the second approach, REDUNDANS was used [57], as

it is optimized to deal with heterozygous samples. The raw sequence data and *L. decemlineata*

genome have been deposited in the GenBank/EMBL/DDBJ database (Bioproject accession

PRJNA171749, Genome assembly GCA_000500325.1, Sequence Read Archive accessions:

SRX396450- SRX396453). This data can be visualized, along with gene models and supporting

data, at the i5k Workspace@NAL: https://i5k.nal.usda.gov/Leptinotarsa_decemlineata and

https://apollo.nal.usda.gov/lepdec/jbrowse/ [147].

Automated gene prediction and annotation were performed using MAKER v2.0 [148],

using RNAseq evidence and arthropod protein databases. The MAKER predictions formed the

basis of the first official gene set (OGS v0.5.3). To improve the structural and functional

annotation of genes, these gene predictions were manually and collaboratively edited using the

interactive curation software Apollo [149]. For a given gene family, known insect genes were

obtained from model species, especially *T. castaneum* [51] and *D. melanogaster* [150], and the

nucleotide or amino acid sequences were used in *blastx* or *tblastn* [151,152] to search the *L.*

decemlineata OGS v0.5.3 or genome assembly, respectively, on the i5k Workspace@NAL. All

available evidence (AUGUSTUS, SNAP, RNA data, etc.), including additional RNAseq data not

used in the MAKER predictions, were used to inspect and modify gene predictions. Changes

were tracked to ensure quality control. Gene models were inspected for quality, incorrect splits

and merges, internal stop codons, and gff3 formatting errors, and finally merged with the

MAKER-predicted gene set to produce the official gene set (OGS v1.0; merging scripts are

available upon request). For focal gene families (e.g. peptidase genes, odorant and gustatory

receptors, RNAi genes, etc.), details on how genes were identified and assigned names based on

functional predictions or evolutionary relationship to known reference genes are provided in the

Supplementary Material (**Additional File 1**).

To assess the quality of our genome assembly, we first used BUSCO v2.0 [59] to

determine the completeness of the genome assembly and the official gene set (OGS v1.0),

separately. We benchmarked our data against 35 insect species in the Endopterygota obd9

database, which consists of 2,442 single-copy orthologs (BUSCOs). Secondly, we annotated and

examined the genomic architecture of the Hox and Iroquois Complex gene clusters. For this,

tBLASTn searches were performed against the genome using orthologous Hox gene protein

sequences from *T. castaneum* (Tcas3.0) and *A. glabripennis*. Provisional *L. decemlineata* models

were refined, and potential gene duplications were identified, via iterative and reciprocal BLAST

and by manual inspection and correction of protein alignments generated with ClustalW2 [153],

using RNAseq expression evidence when available.

Gene Family Evolution

In order to identify rapidly evolving gene families along the *L. decemlineata* lineage, we

obtained ~38,000 ortho-groups from 72 Arthropod species as part of the i5k pilot project

(Thomas et al. *in prep*) from OrthoDB version 8 [154]. For each ortho-group, we took only those

genes present in the order Coleoptera, which was represented by the following six species: *A.*

glabripennis, *A. planipennis*, *D. ponderosae*, *L. decemlineata*, *O. taurus*, and *T. castaneum*

(http://i5k.github.io/) [51,52,54]. Finally, in order to make accurate inferences of ancestral

states, families that were present in only one of the six species were removed. This resulted in a

final count of 11,598 gene families that, among these six species, form the comparative

framework that allowed us to examine rapidly evolving gene families in the *L. decemlineata*

lineage.

Aside from the gene family count data, an ultrametric phylogeny is also required to

estimate gene gain and loss rates. To make the tree, we considered only gene families that were

single copy in all six species and that had another arthropod species also represented with a

single copy as an outgroup. Outgroup species were ranked based on the number of families in

which they were also single copy along with the coleopteran species, and the highest ranking

outgroup available was chosen for each family. For instance, *Pediculus humanus* was the most

common outgroup species. For any gene family, we chose *P. humanus* as the outgroup if it was

also single copy. If it was not, we chose the next highest ranking species as the outgroup for that

family. This process resulted in 3,932 single copy orthologs that we subsequently aligned with

PASTA [155]. We used RAxML [156] with the PROTGAMMAJTTF model to make gene trees

from the alignments and ASTRAL [157] to make the species tree. ASTRAL does not give

branch lengths on its trees, a necessity for gene family analysis, so the species tree was again

given to RAxML along with a concatenated alignment of all one-to-one orthologs for branch

length estimation. Finally, to generate an ultrametric species tree with branch lengths in millions

of years (my) we used the software r8s [158], with a calibration range based on age estimates of

a crown Coleopteran fossil at 208.5-411 my [159]. This calibration point itself was estimated in a

similar fashion in a larger phylogenetic analysis of all 72 Arthropod species (Thomas et al. *in*

prep).

With the gene family data and ultrametric phylogeny as the input data, gene gain and loss

rates (λ) were estimated with CAFE v3.0 [160]. CAFE is able to estimate the amount of

assembly and annotation error (ε) present in the input data using a distribution across the

observed gene family counts and a pseudo-likelihood search, and then is able to correct for this

error and obtain a more accurate estimate of λ. Our analysis had an ε value of about 0.02, which

implies that 3% of gene families have observed counts that are not equal to their true counts.

After correcting for this error rate, $\lambda = 0.0010$ is on par with those previously those found for

other Arthropod orders (Thomas et al. *in prep*). Using the estimated λ value, CAFE infers

ancestral gene counts and calculates p-values across the tree for each family to assess the

significance of any gene family changes along a given branch. Those branches with low p-values

are considered rapidly evolving.

Gene Expression Analysis

RNAseq analyses were conducted to 1) establish male, female, and larva-enriched gene

sets and 2) identify specific genes that are enriched within the digestive tract (mid-gut) compared

to entire larva. RNAseq datasets were trimmed with CLC Genomics v.9 (Qiagen) and quality

was assessed with FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Each

dataset was mapped to the predicted gene set (OGS v1.0) using CLC Genomics. Reads were

mapped with >90% similarity over 60% of length, with two mismatches allowed. Transcripts per

million (TPM) was used as a proxy for gene expression and fold changes were determined as the

TPM in one sample relative to the TPM of another dataset [161]. The Baggerly's test (t-type test

statistic) followed by Bonferroni correction at 0.01 was used to identify genes with significant

enrichment in a specific sample [162]. This stringent statistical analysis was used as only a single

replicate was available for each treatment. Enriched genes were removed, and mapping and

expression analyses were repeated to ensure low expressed genes were not missed. Genes were

identified by BLASTx searching against the NCBI non-redundant protein databases for

arthropods with an expectation value (E-value) < 0.001.

Transposable Elements

We investigated the identities (family membership), diversity, and genomic distribution

of active transposable elements within *L. decemlineata* in order to understand their contribution

to genome structure and to determine their potential positional effect on genes of interest

(focusing on a TE neighborhood size of 1 kb). To identify TEs and analyze their distribution

within the genome, we developed three repeat databases using: 1) RepeatMasker [163], which

uses the library repeats within Repbase (http://www.girinst.org/repbase/), 2) the program

RepeatModeler [164], which identifies de-novo repeat elements, and 3) literature searches to

identify beetle transposons that were not found within Repbase. The three databases were used

within RepeatMasker to determine the overall TE content in the genome.

To eliminate false positives and examine the genome neighborhood surrounding active

TEs, all TE candidate models were translated in 6 frames and scanned for protein domains from

the Pfam and CDD database (using the software transeq from Emboss, hmmer3 and rps-blast).

The protein domain annotations were manually curated in order to remove: a) clear false

positives, b) old highly degraded copies of TEs without identifiable coding potential, and c) the

correct annotation when improper labels were given. The TE models that contained protein

domains were mapped onto the genome and used for our neighborhood analysis: we extracted

the 1 kb flanking regions for each gene and scanned these regions for TEs with intact protein

coding domains.

Population Genetic Variation and Demographic Analysis

Population genetic diversity of pooled RNAseq samples was used to examine genetic

structure of pest populations and past population demography. For Wisconsin, Michigan and the

lab strains from New Jersey, we aligned the RNAseq data to the genomic scaffolds, using

Bowtie2 version 2.1.0 [165] to index the genome and generate aligned SAM files. We used

Burrows-Wheeler Aligner (BWA) version 0.7.5a [166] to align the RNAseq from the three

populations from Europe. SAMtools/BCFtools version 0.1.19 [167] was used to produce BAM

and VCF files. All calls were filtered with VCFtools version 0.1.11 [168] using a minimum

quality score of 30 and minimum depth of 10. All indels were removed from this study.

Population specific VCF files were sorted and merged using VCFtools, and the allele counts

were extracted for each SNP. These allele frequency data were then used to infer population

splits and relative rates of genetic drift using Treemix version 1.12 [169]. We ran Treemix with

SNPs in groups of 1000, choosing to root the tree with the Wisconsin population.

To infer patterns of demographic change in the Midwestern USA (Wisconsin and

Michigan) and European populations, the genome-wide allele frequency spectrum was used in

dadi version 1.6.3 [170] to infer demographic parameters under several alternative models of

population history. The history of *L. decemlineata* as a pest is relatively well-documented. The

introduction of *L. decemlineata* into Europe in 1914 [171] is thought to have involved a strong

bottleneck [15] followed by rapid expansion. Similarly, an outbreak of *L. decemlineata* in

Nebraska in 1859 is thought to have preceded population expansion into the Midwest reaching

Wisconsin in 1865 [3]. For each population, a constant-size model, a two-epoch model of

instantaneous population size change at a time point τ, a bottle-growth model of instantaneous

size change followed by exponential growth, and a three-epoch model with a population size

change of fixed duration followed by exponential growth, was fit to infer θ, the product of the

ancestral effective population size and mutation rate, and relative population size changes.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and material: All data generated or analyzed during this study have been

made publicly available (see Methods for NCBI accession numbers), or included in this

published article and its supplementary information files. The genome assembly and official

gene set can also be accessed at: https://data.nal.usda.gov/dataset/leptinotarsa-decemlineata-

genome-assembly-10_5667 and https://data.nal.usda.gov/dataset/leptinotarsa-decemlineata-

genome-annotations-v053_5668

Competing interests: The authors declare that they have no competing interests.

Funding: We would like to acknowledge the following funding sources: sequencing, assembly

and automated annotation was supported by NIH grant NHGRI U54 HG003273 to RAG; the

UVM Agricultural Experiment Station Hatch grant to YHC (VT-H02010); the NIH postdoctoral

training grant to RFM (K12 GM000708); MMT's work with Apollo was supported by NIH

grants (5R01GM080203 from NIGMS, and 5R01HG004483 from NHGRI) and by the Director,

793 *Authors' contributions*: All authors contributed to the manual annotation effort, data analysis,

794 and data interpretation, in addition to reading and approving the final manuscript. SDS and YHC

795 coordinated the project and drafted the manuscript. SR coordinated genome sequencing,

796 assembly and automated annotation at the Baylor College of Medicine Human Genome

797 Sequencing Center. MFP, CC, MMT, and M-JMC coordinated the biocuration of the genome.

798 GWCT generated the phylogeny and conducted the gene family analysis. MTW conducted the

799 transcription factor analysis. JBB conducted the RNAseq analysis. KB, AM, and YHC conducted

800 the transposable element analysis. SDS and JC conducted the population genetics analysis.

801 *Acknowledgements*: We sincerely thank the sequencing, assembly and annotation teams at

802 the Baylor College of Medicine Human Genome Sequencing Center for their efforts. Mention of

803 trade names or commercial products in this publication is solely for the purpose of providing

804 specific information and does not imply recommendation or endorsement by the U.S.

805 Department of Agriculture. USDA is an equal opportunity provider and employer.

References

1. Grafius E. Economic impact of insecticide resistance in the Colorado potato beetle

(Coleoptera: Chrysomelidae) on the Michigan potato industry. J Econ Entomol. 1997;90:1144–

51.

2. Skryabin K. Do Russia and Eastern Europe need GM plants? Nat Biotechnol. 2010;27:593–5.

3. Walsh BD. The new potato bug and its natural history. Practical Entomol. 1865;1:1–4.

4. Gauthier NL, Hofmaster RN, and Semel M. History of Colorado potato beetle control. In:

Lashomb JH and Casagrande R, editors. Advances in Potato Pest Management. Stroudsburg:

Hutchinson Ross Publishing Co.; 1981. p. 13–33.

5. Hare JD. Ecology and management of the Colorado potato beetle. Annu Rev Entomol.

1990;41:81–100.

6. Weber DC. Colorado Potato Beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera:

Chrysomelidae). In: Capinera JL, editor. Encyclopedia of Entomology. Dordrecht: Springer

Netherlands; 2008. p. 1008–13.

7. Lyytinen A, Boman S, Grapputo A, Lindström L, Mappes J. Cold tolerance during larval

development: effects on the thermal distribution limits of *Leptinotarsa decemlineata*. Entomol

Exp Appl. 2009;133:92–9.

8. Hiiesaar K, Jõgar K, Williams IH, Kruus E, Metspalu L, Luik A, et al. Factors affecting

development and overwintering of second generation Colorado Potato Beetle (Coleoptera:

Chrysomelidae) in Estonia in 2010. Acta Agric Scand Sect B - Soil Plant Sci. 2013;63:506–15.

9. Hiiesaar K, Jõgar K, Williams IH, Luik A, Kruus E, Metspalu L, et al. Phenology and

overwintering of the Colorado potato beetle *Leptinotarsa decemlineata* Say in 2008–2015 in

Estonia. Acta Agric Scand Sect. B — Soil Plant Sci. 2016;1–8.

10. Alyokhin A, Baker M, Mota-Sanchez D, Dively G, Grafius E. Colorado potato beetle

resistance to insecticides. Am J Potato Res. 2008;85:395–413.

11. Sparks TC, Nauen R. IRAC: Mode of action classification and insecticide resistance

management. Pestic Biochem Physiol. 2015;121:122–8.

12. Izzo V, Chen YH, Schoville SD, Wang C, Hawthorne DJ. Origin of pest lineages of the

Colorado potato beetle, *Leptinotarsa decemlineata*. bioRxiv. 2017;doi: 10.1101/156612.

13. Lehmann P, Lyytinen A, Piiroinen S, Lindström L. Northward range expansion requires

synchronization of both overwintering behaviour and physiology with photoperiod in the

invasive Colorado potato beetle (*Leptinotarsa decemlineata*). Oecologia. 2014;176:57–68.

14. Liu N, Li Y, Zhang R. Invasion of Colorado potato beetle, *Leptinotarsa decemlineata*, in

China: dispersal, occurrence, and economic impact. Entomol Exp Appl. 2012;143:207–17.

15. Grapputo A, Boman S, Lindström L, Lyytinen A, Mappes J. The voyage of an invasive

species across continents: genetic diversity of North American and European Colorado potato

beetle populations. Mol Ecol. 2005;14:4207–19.

16. Azeredo-Espin A, Schroder R, Huettel M, Sheppard W. Mitochondrial DNA variation in

geographic populations of Colorado potato beetle, *Leptinotarsa decemlineata* (Coleoptera;

Chrysomelidae). Experientia. 1991;47:483–5.

17. Lu W, Kennedy GG, Gould F. Genetic analysis of larval survival and larval growth of two

populations of *Leptinotarsa decemlineata* on tomato. Entomol. Exp. Appl. 2001;99:143–55.

18. Gaston KJ. The how and why of biodiversity. Nature. 2003;421:900–1.

19. Turnock WJ, Fields PG. Winter climates and cold hardiness in terrestrial insects. Eur J

Entomol. 2005;102:561–576.

20. Piiroinen S, Ketola T, Lyytinen A, Lindström L. Energy use, diapause behaviour and

northern range expansion potential in the invasive Colorado potato beetle. Funct Ecol.

2011;25:527–36.

21. Lehmann P, Lyytinen A, Piiroinen S, Lindström L. Latitudinal differences in diapause related

photoperiodic responses of European Colorado potato beetles (*Leptinotarsa decemlineata*).

Evol Ecol. 2015;29:269–82.

22. Hsiao TH. Host plant adaptations among geographic populations of the Colorado potato

beetle. Entomol Exp Appl. 1978;24:237–47.

23. Jolivet P. Food habits and food selection of Chrysomelidae. Bionomic and evolutionary

perspectives. In: Jolivet P, Petitpierre E, Hsiao TH, editors. Biology of the Chrysomelidae.

Dordrechf: Kluwer Acad. Press; 1988. p. 1–24.

24. Hsiao TH. Host specificity, seasonality and bionomics of *Leptinotarsa* beetles. Biology of

the Chrysomelidae. Springer; 1988. p. 581–99.

25. Hare JD, Kennedy GG. Genetic variation in plant-insect associations: survival of

Leptinotarsa decemlineata populations on *Solanum carolinense*. Evolution. 1986;40:1031–43.

26. Jacques Jr. RL, Fasulo TR. Colorado Potato Beetle, *Leptinotarsa decemlineata* (Say) and

False Potato Beetle, *Leptinotarsa juncta* (Germar) (Insecta: Coleoptera: Chrysomelidae). Univ.

Florida IFAS Ext. 2000;EENY146:1–5.

27. Alyokhin A, and Chen YH. Adaptation to toxic hosts as a factor in the evolution of

insecticide resistance. Curr Opin Insect Sci. 2017;21:33–38.

28. Cárdenas PD, Sonawane PD, Heinig U, Bocobza SE, Burdman S, Aharoni A. The bitter side

of the nightshades: Genomics drives discovery in Solanaceae steroidal alkaloid metabolism.

Phytochemistry. 2015;113:24–32.

29. Milner SE, Brunton NP, Jones PW, Brien NMO, Collins SG, Maguire AR. Bioactivities of

glycoalkaloids and their aglycones from *Solanum* species. J. Agric. Food Chem.

2011;59:3454–84.

30. Dimock MB, Tingey WM. Host acceptance behaviour of Colorado potato beetle larvae

influenced by potato glandular trichomes. Physiol Entomol. 1988;13:399–406.

31. Lawrence SD, Novak NG, Ju CJ-T, Cooke JEK. Examining the molecular interaction

between potato (*Solanum tuberosum*) and Colorado potato beetle *Leptinotarsa decemlineata*.

Botany. 2008;86:1080–91.

32. Petek M, Turnšek N, Gašparič MB, Novak MP, Gruden K, Slapar N, et al. A complex of

genes involved in adaptation of *Leptinotarsa decemlineata* larvae to induced potato defense.

Arch Insect Biochem Physiol. 2012;79:153–81.

33. Novillo C, Castañera P, Ortego F. Characterization and distribution of chymotrypsin-like and

other digestive proteases in Colorado potato beetle larvae. Arch Insect Biochem Physiol.

1997;36:181–201.

34. Armer CA. Colorado potato beetle toxins revisited: Evidence the beetle does not sequester

host plant glycoalkaloids. J Chem. Ecol. 2004;30:883–8.

35. Deroe C, Pasteels JM. Defensive mechanisms against predation in the Colorado beetle

(*Leptinotarsa decemlineata*, Say). Arch Biol Sci. 1977;88:289–304.

36. Hsiao TH, Fraenkel G. Properties of Leptinotarsin: a toxic hemolymph protein from the

Colorado potato beetle. Toxicon. 1969;7:119–30.

37. Forgash AJ. Insecticide resistance in the Colorado potato beetle. In: Ferro DN, Voss RH,

editors. Proc Symp Colorado Potato Beetle, XVII Int Congr Entomol. August 1984. Amherst,

MA: Massachusetts Agricultural Experimental Station; 1985. p. 33–52.

38. Wan P-J, Shi X-Q, Kong Y, Zhou L-T, Guo W-C, Ahmat T, et al. Identification of

cytochrome P450 monooxygenase genes and their expression profiles in cyhalothrin-treated

Colorado potato beetle, *Leptinotarsa decemlineata*. Pestic Biochem Physiol. 2013;107:360–8.

39. Zhang J, Pelletier Y, Goyer C. Identification of potential detoxification enzyme genes in

Leptinotarsa decemlineata (Say) and study of their expression in insects reared on different

plants. J Plant Sci. 2008;88:621–9.

40. Lü F-G, Fu K-Y, Li Q, Guo W-C, Ahmat T, Li G-Q. Identification of carboxylesterase genes

and their expression profiles in the Colorado potato beetle *Leptinotarsa decemlineata* treated

with fipronil and cyhalothrin. Pestic Biochem Physiol. 2015;122:86–95.

41. Zhu F, Xu J, Palli R, Ferguson J, Palli SR. Ingested RNA interference for managing the

populations of the Colorado potato beetle, *Leptinotarsa decemlineata*. Pest Manag Sci.

2011;67:175–82.

42. Clements J, Schoville S, Peterson N, Lan Q, Groves RL. Characterizing molecular

mechanisms of imidacloprid resistance in select populations of *Leptinotarsa decemlineata* in

the Central Sands region of Wisconsin. PLoS One. 2016;11:e0147844.

43. Kumar A, Congiu L, Lindström L, Piiroinen S, Vidotto M, Grapputo A. Sequencing, de novo

assembly and annotation of the Colorado potato beetle, *Leptinotarsa decemlineata*,

transcriptome. PLoS One. 2014;9:e86012.

44. Rinkevich FD, Su C, Lazo TA, Hawthorne DJ, Tingey WM, Naimov S, et al. Multiple

evolutionary origins of knockdown resistance (kdr) in pyrethroid-resistant Colorado potato

beetle, *Leptinotarsa decemlineata*. Pestic Biochem Physiol. 2012;104:192–200.

45. Zhou LT, Jia S, Wan PJ, Kong Y, Guo WC, Ahmat T, et al. RNA interference of a putative

S-adenosyl-L-homocysteine hydrolase gene affects larval performance in *Leptinotarsa*

decemlineata (Say). J Insect Physiol. 2013;59:1049–56.

46. Zhang J, Khan SA, Hasse C, Ruf S, Heckel DG, Bock R. Full crop protection from an insect

pest by expression of long double-stranded RNAs in plastids. Science 2015;347:991-994.

47. Mota-Sanchez D, Hollingworth RM, Grafius EJ, Moyer DD. Resistance and cross-resistance

to neonicotinoid insecticides and spinosad in the Colorado potato beetle, *Leptinotarsa*

decemlineata (Say) (Coleoptera: Chrysomelidae). Pest Manag Sci. 2006;62:30–7.

48. Zhu F, Moural TW, Nelson DR, Palli SR. A specialist herbivore pest adaptation to

xenobiotics through up-regulation of multiple Cytochrome P450s. Sci. Rep. 2016;6:20421.

49. Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, et al.

ALLPATHS: de novo assembly of whole-genome shotgun microreads. Genome Res.

2008;18:810–20.

50. Gregory TR. Animal genome size database [Internet]. 2017. Available from:

http://www.genomesize.com

51. Tribolium Genome Sequencing Consortium. The genome of the model beetle and pest

Tribolium castaneum. Nature. 2008;452:949–55.

52. Keeling CI, Yuen MM, Liao NY, Docking TR, Chan SK, Taylor GA, et al. Draft genome of

the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, a major forest pest. Genome

Biol. 2013;14(3):R27.

53. Cunningham CB, Ji L, Wiberg RAW, Shelton J, McKinney EC, Parker DJ, et al. The genome

and methylome of a beetle with complex social behavior, *Nicrophorus vespilloides*

(Coleoptera: Silphidae). Genome Biol Evol 2015;7(12):3383-3396.

54. McKenna DD, Scully ED, Pauchet Y, Hoover K, Kirsch R, Geib SM, et al. Genome of the

Asian longhorned beetle (*Anoplophora glabripennis*), a globally significant invasive species,

reveals key functional and evolutionary innovations at the beetle–plant interface. Genome Biol.

2016;17:227.

55. Vega FE, Brown SM, Chen H, Shen E, Nair MB, Ceja-Navarro JA, et al. Draft genome of

the most devastating insect pest of coffee worldwide: the coffee berry borer, *Hypothenemus*

hampei. Sci Rep. 2015;5:12525.

56. Petitpierre E, Segarra C, Juan C. Genome size and chromosomal evolution in leaf beetles

(Coleoptera , Chrysomelidae). Hereditas. 1993;6:1–6.

57. Pryszcz LP, Gabaldón T. Redundans: an assembly pipeline for highly heterozygous genomes.

Nucleic Acids Res. 2016;44:e113–e113.

58. Denton JF, Lugo-Martinez J, Tucker AE, Schrider DR, Warren WC, Hahn MW. Extensive

error in the number of genes inferred from draft genome assemblies. PLoS Comput Biol.

2014;10(12):e1003998.

59. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 2015;31:3210–2.

60. Gruden K, Štrukelj B, Popovič T, Lenarčič B, Bevec T, Brzin J, et al. The cysteine protease

activity of Colorado potato beetle (*Leptinotarsa decemlineata* Say) guts, which is insensitive to

potato protease inhibitors, is inhibited by thyroglobulin type-1 domain inhibitors. Insect

Biochem Mol Biol. 1998;28:549–60.

61. Smilanich AM, Dyer LA, Chambers JQ, Bowers MD. Immunological cost of chemical

defence and the evolution of herbivore diet breadth. Ecol Lett. 2009;12:612–21.

62. Vidal NM, Grazziotin AL, Iyer LM, Aravind L, Venancio TM. Transcription factors,

chromatin proteins and the diversification of Hemiptera. Insect Biochem Mol Biol. 2016;69:1–

13.

63. Najafabadi HS, Mnaimneh S, Schmitges FW, Garton M, Lam KN, Yang A, et al. C2H2 zinc

finger proteins greatly expand the human regulatory lexicon. Nat Biotech. 2015;33:555–62.

64. Chénais B, Caruso A, Hiard S, Casse N. The impact of transposable elements on eukaryotic

genomes: from genome size increase to genetic adaptation to stressful environments. Gene.

2012;509:7–15.

65. Tu Z. Insect Transposable Elements. In: Gilbert LI, editor. Insect Molecular Biology and

Biochemistry. London, U. K.: Academic Press; 2012.

66. Casacuberta E, González J. The impact of transposable elements in environmental

adaptation. Mol Ecol. 2013;22:1503–17.

67. Bourque G, Leong B, Vega VB, Chen X, Lee YL, Srinivasan KG, et al. Evolution of the

mammalian transcription factor binding repertoire via transposable elements. Genome Res.

2008;18:1752–62.

68. Cordaux R, Batzer MA. The impact of retrotransposons on human genome evolution. Nat

Rev Genet. 2009;10:691–703.

69. González J, Petrov DA. The adaptive role of transposable elements in the *Drosophila*

genome. Gene. 2009;448:124–33.

70. Biémont C, Vieira C. Genetics: junk DNA as an evolutionary force. Nature. 2006;443:521–4.

71. Lavoie CA, Platt RN, Novick PA, Counterman BA, Ray DA. Transposable element

evolution in *Heliconius* suggests genome diversity within Lepidoptera. Mob DNA. 2013;4:21.

72. Osanai-Futahashi M, Suetsugu Y, Mita K, Fujiwara H. Genome-wide screening and

characterization of transposable elements and their distribution analysis in the silkworm,

Bombyx mori. Insect Biochem Mol Biol. 2008;38:1046–57.

73. Schrader L, Kim JW, Ence D, Zimin A, Klein A, Wyschetzki K, et al. Transposable element

islands facilitate adaptation to novel environments in an invasive species. Nat Commun.

2014;5.

74. González J, Karasov TL, Messer PW, Petrov DA. Genome-wide patterns of adaptation to

temperate environments associated with transposable elements in *Drosophila*. PLoS Genet.

2010;6:e1000905.

75. Cridland JM, Thornton KR, Long AD. Gene expression variation in *Drosophila*

melanogaster due to rare transposable element insertion alleles of large effect. Genetics.

2015;199:85–93.

76. Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, et al. A single P450 allele

associated with insecticide resistance in *Drosophila*. Science. 2002;297:2253–6.

77. Messer PW, Petrov DA. Population genomics of rapid adaptation by soft selective sweeps.

Trends Ecol Evol 2013;28(11):659-669.

78. Aquadro CF, Bauer DuMont V, Reed FA. Genome-wide variation in the human and fruitfly:

a comparison. Curr Opin Genet Dev. 2001;11:627–34.

79. International Chicken Polymorphism Map Consortium. A genetic variation map for chicken

with 2.8 million single-nucleotide polymorphisms. Nature. 2004;432:717–22.

80. Choi J-H, Kijimoto T, Snell-Rood E, Tae H, Yang Y, Moczek AP, et al. Gene discovery in

the horned beetle *Onthophagus taurus*. BMC Genomics 2010;11(1):703.

81. Morlais I, Ponçon N, Simard F, Cohuet A, Fontenille D. Intraspecific nucleotide variation in

Anopheles gambiae: new insights into the biology of malaria vectors. Am J Trop Med Hyg.

2004;71:795–802.

82. Campos JL, Zhao L, Charlesworth B. Estimating the parameters of background selection and

selective sweeps in *Drosophila* in the presence of gene conversion. Proc Natl Acad Sci U S A.

2017;114:E4762–71.

83. Dangles O, Irschick D, Chittka L, Casas J. Variability in sensory ecology: expanding the

bridge between physiology and evolutionary biology. Q Rev Biol. 2009;84:51–74.

84. Matsuura H, Sokabe T, Kohno K, Tominaga M, Kadowaki T. Evolutionary conservation and

changes in insect TRP channels. BMC Evol Biol. 2009;9:228.

85. Venkatachalam K, Montell C. TRP channels. Annu. Rev. Biochem. 2007;76:387–417.

86. Nesterov A, Spalthoff C, Kandasamy R, Katana R, Rankl NB, Andrés M, et al. TRP channels

in insect stretch receptors as insecticide targets. Neuron. 2015;86:665–71.

87. Hauser F, Cazzamali G, Williamson M, Park Y, Li B, Tanaka Y, et al. A genome-wide

inventory of neurohormone GPCRs in the red flour beetle *Tribolium castaneum*. Front

Neuroendocrinol. 2008;29:142–65.

88. Nishi Y, Sasaki K, Miyatake T. Biogenic amines, caffeine and tonic immobility in *Tribolium*

castaneum. J Insect Physiol. 2010;56:622–8.

89. Roeder T. Biochemistry and molecular biology of receptors for biogenic amines in locusts.

Microsc Res Tech. 2002;56:237–47.

90. Schoonhoven L, Van Loon JJA, Dicke M. Insect–Plant Biology. 2nd Ed. Oxford, U. K.:

Oxford University Press; 2005.

91. Leal WS. Odorant reception in insects: roles of receptors, binding proteins, and degrading

enzymes. Annu Rev Entomol. 2013;58:373–91.

92. Pelosi P, Zhou J-J, Ban LP, Calvello M. Soluble proteins in insect chemical communication.

Cell Mol Life Sci. 2006;63:1658–76.

93. Robertson HM, Warr CG, Carlson JR. Molecular evolution of the insect chemoreceptor gene

superfamily in *Drosophila melanogaster*. Proc Natl Acad Sci U S A. 2003;14537–42.

94. Sato K, Tanaka K, Touhara K. Sugar-regulated cation channel formed by an insect gustatory

receptor. Proc Natl Acad Sci U S A. 2011;108:11680–5.

95. Rytz R, Croset V, Benton R. Ionotropic Receptors (IRs): Chemosensory ionotropic glutamate

receptors in *Drosophila* and beyond. Insect Biochem Mol Biol. 2013;43:888–97.

96. Jackowska M, Bao R, Liu Z, McDonald EC., Cook TA, Friedrich M. Genomic and gene

regulatory signatures of cryptozoic adaptation: Loss of blue sensitive photoreceptors through

expansion of long wavelength-opsin expression in the red flour beetle *Tribolium castaneum*.

Front Zool. 2007;4:24.

97. Sharkey CR, Fujimoto MS, Lord NP, Shin S, McKenna DD, Suvorov A, et al. Overcoming

the loss of blue sensitivity through opsin duplication in the largest animal group, beetles. Sci

Rep. 2017;7:8.

98. Izzo VM, Mercer N, Armstrong J, Chen YH. Variation in host usage among geographic

populations of *Leptinotarsa decemlineata*, the Colorado potato beetle. J Pest Sci. 2014;87:597–

608.

99. Otálora-Luna F, Dickens JC. Multimodal stimulation of Colorado potato beetle reveals

modulation of pheromone response by yellow light. PLoS One. 2011;6:e20990.

100. Briscoe AD, Chittka L. The evolution of color vision in insects. Annu Rev Entomol.

2001;46:471–510.

101. Leung NY, Montell C. Unconventional roles of opsins. Annu Rev Cell Dev Biol. 2017;33.

102. Price PW, Denno RF, Eubanks MD, Finke DL, Kaplan I. Insect Ecology: Behavior,

Populations and Communities. New York, NY: Cambridge University Press.; 2011.

103. Martinez M, Santamaria M, Diaz-Mendoza M, Arnaiz A, Carrillo L, Ortego F, et al.

Phytocystatins: defense proteins against phytophagous insects and Acari. Int J Mol Sci.

2016;17:1747.

104. Wolfson JL, Murdock LL. Suppression of larval Colorado potato beetle growth and

development by digestive proteinase inhibitors. Entomol Exp Appl. 1987;44:235–40.

105. Bolter CJ, Jongsma MA. Colorado potato beetles (*Leptinotarsa decemlineata*) adapt to

proteinase inhibitors induced in potato leaves by methyl jasmonate. J Insect Physiol.

1995;41:1071–8.

106. Murdock LL, Brookhart G, Dunn PE, Foard DE, Kelley S, Kitch L, et al. Cysteine digestive

proteinases in Coleoptera. Biochem Physiol. 1987;87:783–7.

107. Chye M-L, Sin S-F, Xu Z-F, Yeung EC. Serine proteinase inhibitor proteins: Exogenous

and endogenous functions. Vitr Cell Dev Biol-Plant. 2006;42:100–8.

108. Gruden K, Popovič T, Cimerman N, Križaj I, Štrukelj B. Diverse enzymatic specificities of

digestive proteases, "intestains", enable Colorado potato beetle larvae to counteract the potato

defence mechanism. Biol Chem. 2003;384:305–10.

109. Gruden K, Kuipers AG., Gunčar G, Slapar N, Štrukelj B, Jongsma MA. Molecular basis of

Colorado potato beetle adaptation to potato plant defence at the level of digestive cysteine

proteinases. Insect Biochem Mol Biol. 2004;34:365–75.

110. Goptar IA, Semashko TA, Danilenko SA, Lysogorskaya EN, Oksenoit ES, Zhuzhikov DP,

et al. Cysteine digestive peptidases function as post-glutamine cleaving enzymes in tenebrionid

stored-product pests. Comp Biochem Physiol Part B 2012;161:148–54.

111. Turk V, Stoka V, Vasiljeva O, Renko M, Sun T, Turk B, et al. Cysteine cathepsins: From

structure, function and regulation to new frontiers. Biochim Biophys Acta - Proteins

Proteomics. 2012;1824:68–88.

112. Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, et al. CDD:

NCBI's conserved domain database. Nucleic Acids Res. 2014;43:D222-D226.

113. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the

protein families database. Nucleic Acids Res. 2013;42: D222-D230.

114. Wex T, Lipyansky A, Brömme NC, Wex H, Guan XQ, Brömme D. TIN-ag-RP, a novel

catalytically inactive cathepsin B-related protein with EGF domains, is predominantly

expressed in vascular smooth muscle cells†. Biochemistry. 2001;40:1350–7.

115. Yamamoto Y, Kurata M, Watabe S, Murakami R, Takahashi SY. Novel cysteine proteinase

inhibitors homologous to the proregions of cysteine proteinases. Curr Protein Pept Sci.

2002;3:231–8.

116. Sainsbury F, Rhéaume A-J, Goulet M-C, Vorster J, Michaud D. Discrimination of

differentially inhibited cysteine proteases by activity-based profiling using cystatin variants

with tailored specificities. J Proteome Res. 2012;11:5983–93.

117. Martynov AG, Elpidina EN, Perkin L, Oppert B. Functional analysis of C1 family cysteine

peptidases in the larval gut of *Tenebrio molitor* and *Tribolium castaneum*. BMC Genomics.

2015;16:75.

118. Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B. Expansion of the enzymatic

repertoire of the CAZy database to integrate auxiliary redox enzymes. Biotechnol Biofuels

2013;6(1):41.

119. Scully ED, Hoover K, Carlson JE, Tien M, Geib SM. Mid-gut transcriptome profiling of

Anoplophora glabripennis, a lignocellulose degrading cerambycid beetle. BMC Genomics.

2013;14:850.

120. Kirsch R, Gramzow L, Theißen G, Siegfried BD, Heckel DG, Pauchet Y. Horizontal gene

transfer and functional diversification of plant cell wall degrading polygalacturonases: key

events in the evolution of herbivory in beetles. Insect Biochem Mol Biol. 2014;52:33–50.

121. Pauchet Y, Wilkinson P, Chauhan R, Ffrench-Constant RH. Diversity of beetle genes

encoding novel plant cell wall degrading enzymes. PloS One 2010;5:e15635.

122. Meyer UA. Overview of enzymes of drug metabolism. J Pharmacokinet Pharmacodyn

1996;24(5):449-459.

123. Bloomquist JR. Chloride channels as tools for developing selective insecticides. Arch Insect

Biochem Physiol. 2003;54:145–56.

124. Le Goff G, Hamon A, Bergé J-B, Amichot M. Resistance to fipronil in *Drosophila*

simulans: influence of two point mutations in the RDL GABA receptor subunit. J Neurochem.

2005;92:1295–305.

125. Jones AK, Sattelle DB. The cys-loop ligand-gated ion channel gene superfamily of the

nematode, *Caenorhabditis elegans*. Invert Neurosci. 2008;8:41–7.

126. Jones AK, Sattelle DB. Diversity of insect nicotinic acetylcholine receptor subunits. In:

Thany SH, editor. Insect Nicotinic Acetylcholine Receptors. New York, NY: Springer New

York; 2010. p. 25–43.

127. Ffrench-Constant RH, Mortlock DP, Shaffer CD, MacIntyre RJ, Roush RT. Molecular

cloning and transformation of cyclodiene resistance in *Drosophila*: an invertebrate gamma-

aminobutyric acid subtype A receptor locus. Proc Natl Acad Sci U S A. 1991;88:7209–13.

128. Ffrench-Constant RH, Rocheleau TA. *Drosophila* γ-aminobutyric acid receptor gene Rdl

shows extensive alternative splicing. J Neurochem. 1993;60:2323–6.

129. Buckingham SD, Biggin PC, Sattelle BM, Brown LA, Sattelle DB. Insect GABA receptors:

splicing, editing, and targeting by antiparasitics and insecticides. Mol Pharmacol.

2005;68:942–51.

130. Clements J, Schoville S, Peterson N, Lan Q, Groves RL. Characterizing molecular

mechanisms of Imidacloprid resistance in select populations of *Leptinotarsa decemlineata* in

the Central Sands Region of Wisconsin. PLoS One. 2016;11:e0147844.

131. Clements J, Schoville S, Peterson N, Huseth AS, Lan Q, Groves RL. RNA interference of

three up-regulated transcripts associated with insecticide resistance in an imidacloprid resistant

population of *Leptinotarsa decemlineata*. Pestic Biochem Physiol. 2017;135:35–40.

132. Willis JH. Structural cuticular proteins from arthropods: Annotation, nomenclature, and

sequence characteristics in the genomics era. Insect Biochem Mol Biol. 2010;40:189–204.

133. McDonnell CM, King D, Comeron JM, Li H, Sun W, Berenbaum MR, et al. Evolutionary

toxicogenomics: diversification of the Cyp12d1 and Cyp12d3 Genes in *Drosophila* species. J

Mol Evol. 2012;74:281–96.

134. Hu F, Dou W, Wang J-J, Jia F-X, Wang J-J. Multiple glutathione S-transferase genes:

identification and expression in oriental fruit fly, *Bactrocera dorsalis*. Pest Manag Sci.

2014;70:295–303.

135. Han J-B, Li G-Q, Wan P-J, Zhu T-T, Meng Q-W. Identification of glutathione S-transferase

genes in *Leptinotarsa decemlineata* and their expression patterns under stress of three

insecticides. Pestic Biochem Physiol. 2016;133:26–34.

136. Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific

genetic interference by double-stranded RNA in *Caenorhabditis elegans*. Nature.

1998;391:806–11.

137. Siomi H, Siomi MC. On the road to reading the RNA-interference code. Nature.

2009;457:396–404.

138. Revuelta L, Ortego F, Díaz-Ruíz JR, Castañera P, Tenllado F, Hernández-Crespo P.

Contribution of ldace1 gene to acetylcholinesterase activity in Colorado potato beetle. Insect

Biochem Mol Biol. 2011;

139. Wan P-J, Fu K-Y, Lü F-G, Guo W-C, Li G-Q. Knockdown of a putative alanine

aminotransferase gene affects amino acid content and flight capacity in the Colorado potato

beetle *Leptinotarsa decemlineata*. Amino Acids. 2015;47:1445–54.

140. Baum JA, Bogaert T, Clinton W, Heck GR, Feldmann P, Ilagan O, et al. Control of

coleopteran insect pests through RNA interference. Nat Biotechnol. 2007;25:1322.

141. Swevers L, Huvenne H, Menschaert G, Kontogiannatos D, Kourti a., Pauchet Y, et al.

Colorado potato beetle (Coleoptera) gut transcriptome analysis: expression of RNA

interference-related genes. Insect Mol Biol. 2013;22:668–84.

142. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in

the initiation step of RNA interference. Nature. 2001;409:363–6.

143. Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, et al. Genes and mechanisms

related to RNA interference regulate expression of the small temporal RNAs that control *C.*

elegans developmental timing. Cell. 2001;106:23–34.

144. Yoon J-S, Shukla JN, Gong ZJ, Mogilicherla K., and Palli SR. RNA interference in the

Colorado potato beetle, *Leptinotarsa decemlineata*: Identification of key contributors. Insect

Biochem Mol Biol. 2016;78:78-88.

145. Guénin H-A, Scherler M. La formule chromosomiale du doryphore *Leptinotarsa*

decemlineata Stål. Rev Suisse Zool. 1951;58:359–70.

146. Hsiao TH, Hsiao C. Chromosomal analysis of *Leptinotarsa* and *Labidomera* species.

Genetica. 1983;60:139–50.

147. Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee C-Y, et al. The i5k

Workspace@NAL—enabling genomic data access, visualization and curation of arthropod

genomes. Nucleic Acids Res 2015;43(D1):D714-D719.

148. Holt C, Yandell M. MAKER2: an annotation pipeline and genome-database management

tool for second-generation genome projects. BMC Bioinformatics. 2011;12:491.

149. Lee E, Helt GA, Reese JT, Munoz-Torres MC, Childers CP, Buels RM, et al. Web Apollo:

a web-based genomic annotation editing platform. Genome Biol. 2013;14:R93.

150. Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, et al. The

genome sequence of *Drosophila melanogaster*. Science 2000;287(5461):2185-2195.

151. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10:421.

152. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.

J Mol Biol. 1990;215:403–10.

153. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.

Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23:2947–8.

154. Kriventseva E V, Tegenfeldt F, Petty TJ, Waterhouse RM, Simão FA, Pozdnyakov IA, et al.

OrthoDB v8: update of the hierarchical catalog of orthologs and the underlying free software.

Nucleic Acids Res. 2015;43:D250-D256.

155. Mirarab S, Nguyen N, Guo S, Wang L-S, Kim J, Warnow T. PASTA: ultra-large multiple

sequence alignment for nucleotide and amino-acid sequences. J Comput Biol. 2015;22:377–86.

156. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014;2010–1.

157. Mirarab S, Warnow T. ASTRAL-II: coalescent-based species tree estimation with many

hundreds of taxa and thousands of genes. Bioinformatics. 2015;31:i44--i52.

158. Sanderson MJ. r8s: inferring absolute rates of molecular evolution and divergence times in

the absence of a molecular clock. Bioinformatics. 2003;19:301–2.

159. Wolfe JM, Daley AC, Legg DA, Edgecombe GD. Fossil calibrations for the arthropod Tree

of Life. Earth-Sci Rev 2016;160(Supp. C):43-110.

160. Han M V, Thomas GWC, Lugo-Martinez J, Hahn MW. Estimating gene gain and loss rates

in the presence of error in genome assembly and annotation using CAFE 3. Mol Biol Evol.

2013;30:1987–97.

161. Rosendale AJ, Romick-Rosendale LE, Watanabe M, Dunlevy ME, Benoit JB. Mechanistic

underpinnings of dehydration stress in the American dog tick revealed through RNA-seq and

metabolomics. J Exp Biol. 2016;219:1808–19.

162. Baggerly KA, Deng L, Morris JS, Aldaz CM. Differential expression in SAGE: accounting

for normal between-library variation. Bioinformatics. 2003;19:1477–83.

163. Smit, AFA, Hubley, R & Green P. RepeatMasker Open-4.0.

164. Smit A, Hubley R. RepeatModeler Open-1.0.

165. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat. Methods.

2012;9:357.

166. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25:1754–60.

167. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. Bioinformatics. 2009;25:2078–9.

168. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call

format and VCFtools. Bioinformatics. 2011;27:2156–8.

169. Pickrell JK, Pritchard JK. Inference of population splits and mixtures from genome-wide

allele frequency data. PLoS Genet. 2012;8:e1002967.

170. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint

demographic history of multiple populations from multidimensional SNP frequency data.

PLoS Genet. 2009;5:e1000695.

171. Jacques Jr RL. The Potato Beetles. Flora Fauna Handb. No. 3. Taylor & Francis; 1988

1226

1227 **Table 1:** Summary of gene gain and loss events inferred after correcting for annotation and

1228 assembly error across all 6 species. The number of rapidly evolving families is shown in

1229 parentheses for each type of change and the rate is genes per million years.

	Expansions			Contractions			No Change	Average Expansion
	Families	Genes gained	Rate	Families	Genes lost	Rate		
Anoplophora glabripennis	865 (13)	1231	1.42	1988 (107)	3125	1.57	5850	-0.182341
Agrilus plannipennis	739 (100)	1991	2.69	707 (8)	769	1.09	7257	0.119108
Dendroctonus ponderosae	933 (72)	1982	2.12	1606 (21)	1887	1.17	6164	0.006438
Leptinotarsa decemlineata	426 (40)	855	2.01	1556 (48)	2116	1.36	6721	-0.127501
Onthophagus taurus	1299 (142)	2952	2.27	767 (24)	895	1.17	6637	0.203380
Tribolium castaneum	786 (51)	1428	1.82	516 (27)	909	1.76	7401	0.055645

1230

Figure Legend

Figure 1. Ultrametric tree with branch lengths in millions of years for *Leptinotarsa decemlineata*

relative to five other Coleoptera genomes. The *L. decemlineata* lineage is shown in orange.

Branches are labeled with their length in years (top) and with the number of gene family

expansions (blue) and contractions (purple) that occurred on that lineage. Rapid changes for both

types are in parentheses.

Figure 2. Heatmap distribution of the abundance of transcription factor families in *Leptinotarsa*

decemlineata compared to other insects. Each entry indicates the number of TF genes for the

given family per genome, based on presence of predicted DNA binding domains. Color scale is

log (base 2) and the key is depicted at the top (light blue means the TF family is completely

absent). Families discussed in the main text are indicated by arrows.

Figure 3. Volcano plots showing statistically significant gene expression differences after

Bonferroni correction in *Leptinotarsa decemlineata*. A) Mid -gut tissue versus whole larvae, B)

an adult male versus an adult female, C) an adult male versus whole larvae, and D) an adult

female versus whole larvae. Points outside the gray area indicate >100-fold-differences in

expression. Blue points indicate down-regulated genes and red points indicate up-regulated genes

in each contrast.

Figure 4. Population genetic relationships and relative rates of genetic drift among *Leptinotarsa*

decemlineata pest populations based on single nucleotide polymorphism data. Population codes:

NJ- New Jersey lab strain, WIs- imidacloprid susceptible population from Arlington, Wisconsin,

WIr- imidacloprid resistant population from Hancock, Wisconsin, MI- imidacloprid resistant

population from Michigan, and EU- European samples combined from Italy and Russia.

Figure 5. Phylogenetic relationships of the cysteine peptidase gene family in *Leptinotarsa*

decemlineata compared to model insects. Species abbreviations are: *L. decemlineata* (Ld, green

color), *Drosophila melanogaster* (Dm, blue color), *Apis mellifera* (Am, purple color), and

Tribolium castaneum (Tc, red color). Mid-gut gene expression (TPM) of highly expressed *L.*

decemlineata cysteine peptidases is shown as bar graphs across three replicate treatments.

1258

1259

1260
1261 **Figure 1.** Ultrametric tree with branch lengths in millions of years for *Leptinotarsa decemlineata*
1262 relative to five other Coleoptera genomes. The *L. decemlineata* lineage is shown in orange.
1263 Branches are labeled with their length in years (top) and with the number of gene family
1264 expansions (blue) and contractions (purple) that occurred on that lineage. Rapid changes for both
1265 types are in parentheses.

1266
1267 **Figure 2.** Heatmap distribution of the abundance of transcription factor families in *Leptinotarsa*
1268 *decemlineata* compared to other insects. Each entry indicates the number of TF genes for the
1269 given family per genome, based on presence of predicted DNA binding domains. Color scale is
1270 log (base 2) and the key is depicted at the top (light blue means the TF family is completely
1271 absent). Families discussed in the main text are indicated by arrows.

1272
Figure 3. Volcano plots showing statistically significant gene expression differences after
Bonferroni correction in *Leptinotarsa decemlineata*. A) Mid-gut tissue versus whole larvae, B)
an adult male versus an adult female, C) an adult male versus whole larvae, and D) an adult
female versus whole larvae. Points outside the gray area indicate >100-fold-differences in
expression. Blue points indicate down-regulated genes and red points indicate up-regulated genes
in each contrast.
1279

1280

1281
1282 **Figure 4.** Population genetic relationships and relative rates of genetic drift among *Leptinotarsa*
1283 *decemlineata* pest populations based on single nucleotide polymorphism data. Population codes:
1284 NJ- New Jersey lab strain, WIs- imidacloprid susceptible population from Arlington, Wisconsin,
1285 WIr- imidacloprid resistant population from Hancock, Wisconsin, MI- imidacloprid resistant
1286 population from Michigan, and EU- European samples combined from Italy and Russia.
1287

1289
1290 **Figure 5**. Phylogenetic relationships of the cysteine peptidase gene family in *Leptinotarsa*
1291 *decemlineata* compared to model insects. Species abbreviations are: *L. decemlineata* (Ld, green
1292 color), *Drosophila melanogaster* (Dm, blue color), *Apis mellifera* (Am, purple color), and
1293 *Tribolium castaneum* (Tc, red color). Mid-gut gene expression (TPM) of highly expressed *L.*
1294 *decemlineata* cysteine peptidases is shown as bar graphs across three replicate treatments.
1295

Additional Files

Additional File 1.pdf Supplementary Materials. Contains additional methods, results, figures and

tables.

Additional File 2.fasta Precursor miRNA nucleotide sequences.

Additional File 3.fasta Peptide sequences of annotated olfactory genes. Includes the Odorant

Binding Proteins (OBPs), Odorant Receptors (ORs), Gustatory Receptors (GRs), and Ionotropic

Receptors (IRs). The IRs from *Tribolium castaneum* are also included.