Overlooked biodiversity loss in Amazonian smallholder agriculture

Jacob B. Socolar[1†*]

Elvis H. Valderrama Sandoval[2]

David S. Wilcove[1,3]

1: Department of Ecology and Evolutionary Biology, Princeton University, Princeton NJ

2: Universidad Nacional de la Amazonía Peruana, Iquitos Perú

3: Woodrow Wilson School of Public Policy, Princeton University, Princeton NJ

†: jacob.socolar@gmail.com

*Current address, Department of Ecology and Evolutionary Biology, University of Connecticut,

Storrs CT

ABSTRACT

Diversified smallholder agriculture is the main human land-use affecting the western Amazon, home

to the world's richest terrestrial biota, but the scant available data to date have suggested that the

biodiversity impacts of this land-use are small. Here, we present comprehensive surveys of birds and

trees in primary forest and smallholder agricultural mosaics in northern Peru. These surveys reveal

substantial biodiversity losses that have been overlooked by other studies. Avian biodiversity losses

arise primarily from biotic homogenization across infrequently surveyed forest habitats (a loss of

beta-diversity). Furthermore, tree species richness declines much more steeply than bird richness.

Statistical modeling of local habitat features that allow forest-associated species to persist in the

smallholder mosaic strongly suggests that our results represent a best-case scenario for Amazonian

agricultural biodiversity. We conclude that previous assessments of the biodiversity value of

Amazonian smallholder agriculture have been overly optimistic because they are restricted to upland

habitat, thereby missing losses in beta diversity; do not evaluate trees; and/or rely on generalizations

from less speciose areas of the Neotropics, where habitat specialization amongst species is less

prevalent. Smallholder agriculture will likely expand in western Amazonia due to infrastructure

development, and it must be seen as a serious threat to the region's biodiversity.

Key words: Amazon, beta diversity, birds, Peru, trees

INTRODUCTION

The western Amazon is the global epicenter of terrestrial biodiversity (Jenkins et al. 2013)

and the largest remaining tropical forest wilderness (Tyukavina et al. 2015), but it is nevertheless

threatened by human activities. In contrast to the mechanized agriculture and ranching in

southeastern Amazonia, the principal driver of forest loss in the western Amazon is smallholder

slash-and-burn agriculture (Finer & Novoa 2016; Ravikumar et al. 2017). This practice creates

mosaics of cultivations and secondary forest surrounding human settlements. The prospect of

increased smallholder settlement in western Amazonia in the wake of roadbuilding and

hydrocarbons development has raised alarm for this bastion of tropical biodiversity. For example,

most of western Amazonia is covered in hydrocarbons concessions, the development of which

would provide road access for settlers (Finer & Orta-Martínez 2010; Laurance et al. 2014).

Existing data on biodiversity in western Amazonian agriculture are extremely scant,

consisting of two small-scale studies of birds and dung beetles, respectively (Andrade & Torgler

1994; Korasaki et al. 2013). These studies document comparable avian richness in slash-and-burn

mosaic and primary forest (Andrade & Torgler 1994), and comparable dung beetle richness in young

secondary forest and primary forest (Korasaki et al. 2013). These isolated results contrast with

results obtained from more intensive land-use change in the eastern Amazon (e.g. large-scale

agriculture, fragmentation, silviculture, or fire; Ferraz et al. 2007; Barlow et al. 2007; Berry et al.

2008; Mahood et al. 2011; Gardner et al. 2013; Lees et al. 2015; Moura et al. 2016). However, they

agree with numerous Mesoamerican studies that have documented high levels of biodiversity in

smallholder mosaics (Daily et al. 2001; Sekercioglu et al. 2007; Ranganathan et al. 2008; Karp et al.

2011; Mendenhall et al. 2011) and have generated sustained debates over the relative conservation

benefits of protecting primary-forest versus preventing agricultural intensification/industrialization

of smallholder mosaics, especially given limited funding for conservation (Phalan et al. 2011; Gibson

et al. 2011). This debate has a special urgency in the western Amazon, where forests remain largely

intact but under increasing threat from smallholder agriculture, including inside protected areas

(Finer & Novoa 2016).

Despite concordant results from Andrade and Torgler (1994) and Korasaki et al.

(2013) that would downplay the significance of smallholder agriculture and conversion of primary

forest to secondary forest, there are strong reasons to think that the impacts of smallholder activities

on Amazonian biodiversity might be more severe than previously recognized. Thus, the biodiversity

impacts of the main land-use affecting the world's richest terrestrial biota remain unknown. First,

the few Amazonian studies that have examined smallholder agriculture either have included

relatively few smallholder sites (e.g. 7 out of 361 sites in smallholder habitats in Moura et al. 2013) or

are limited in their total sampling (Andrade & Torgler 1994; Korasaki et al. 2013). For example,

Andrade and Torgler (1994) found bird diversity comparable to primary forest in Colombian slash-

and-burn mosaics, but this conclusion rests on only understory birds sampled over a relatively small

area.

Second, Amazonia is more species-rich than other areas of the Neotropics, so data from

Mesoamerica might not generalize to the Amazon. Ecological theory predicts that habitat

specialization among species should be more frequent in hyperdiverse communities (MacArthur &

Levins 1967), and this might predispose Amazonian communities to be more sensitive to habitat

alteration. Consistent with this idea, modelling work suggests that a given deforestation scenario

would impact Amazonian bird communities more heavily than their Mesoamerican counterparts

(Newbold et al. 2014).

Third, previous studies focused on the upland (*terra firme*) forest of uplifted clay terraces. Yet

Amazonia contains additional forest types that are critical for biodiversity and are also impacted by

slash-and-burn. These include floodplain habitats, bamboo forests, and forests on white-sand soils,

all of which harbor specialist species that do not occur in terra firme forests (Remsen & Parker 1983;

Wittmann et al. 2006; Fine et al. 2010; Alvarez Alonso et al. 2013; Socolar et al. 2013). Because biotic

homogenization can drive landscape-scale biodiversity loss in tropical forests (Karp et al. 2012; Solar

et al. 2015; Alroy 2017; Giam 2017), effective conservation planning requires an extensive

comparison of biodiversity in intact and degraded landscapes across multiple forest types (Socolar et

al. 2016). However, we are unaware of any data from the western Amazon that evaluate the

biodiversity consequences of land-use change across multiple forest-types simultaneously.

Here, we quantify the biodiversity consequences of Amazonian slash-and-burn agriculture

based on extensive field surveys of bird and tree diversity in Loreto department, Peru. In Loreto,

upland, floodplain, and white-sand forests collectively harbor the richest avifauna and tree flora on

Earth (ter Steege et al. 2013). Although the area remains largely roadless, the city of Iquitos is the

world's largest city without an outside road link (circa 0.5 million inhabitants), and slash-and-burn

mosaics are ubiquitous along rivers and local roads (Mäki et al. 2001). Furthermore, slash-and-burn

is practiced to varying degrees, often legally, inside the region's protected areas (Pulgar Vidal &

Gamboa Moquillaza 2013). Therefore, our results stand to inform conservation practices and

priorities across this hyperdiverse yet understudied region in the face of ongoing smallholder

expansion.

Despite previous work suggesting that the biodiversity value of tropical smallholder

landscapes is very high (Sekercioglu et al. 2007; Karp et al. 2011; Mendenhall et al. 2011), we

hypothesized that habitat specialists would fare poorly at disturbed sites, driving landscape-scale

biodiversity declines via a reduction in beta-diversity (Karp et al. 2012; Socolar et al. 2016).

METHODS

STUDY SITES

We conducted fieldwork in the Amazonian lowlands of Loreto Department, Peru within 230

km of the city of Iquitos. Natural habitats in the region are varied and interdigitate at fine spatial

scales. We focused on four terrestrial habitats that harbor distinctive biological communities.

Quintessential *upland forest* grows on uplifted clay-soil terraces of (Higgins et al. 2011). These uplands

are the most spatially extensive habitat in the region and the richest in bird and tree species.

Floodplain forest along major rivers, subject to protracted flooding during January-June (Espinoza et al.

2013), differs from the uplands in tree and bird species composition (Remsen & Parker 1983;

Wittmann et al. 2004). *White-sand forest* occurs on deposits of pure white-sand soil (arenosols) and

supports a characteristic avifauna and flora that is absent from other habitats (Fine et al. 2010;

Alvarez Alonso et al. 2013). Lastly, *river islands* harbor *Cecropia* (Urticaceae)-dominated woodland

with a characteristic suite of specialist birds (Rosenberg 1990). Slash-and-burn agriculture affects all

of these habitats, removing primary forest vegetation and replacing it with a heterogeneous mosaic

of clearings, hedgerows, and secondary forests (Figure 1). Typical crops include manioc, corn, camu-

camu, and watermelon on floodplains; manioc, plantain, rice, small buffalo pastures, and small

aquaculture ponds in uplands; manioc and pineapple on white sands; and rice, watermelon, and

manioc on islands.

We sampled bird and tree communities at intact sites (primary forest) and disturbed sites

(slash-and-burn mosaics of active cultivation and fallow secondary forest). We selected twenty intact

sites within 230 km of Iquitos harboring accessible habitat that is largely undisturbed by humans for

as long as records are available, except for light selective logging at floodplain sites and widespread

hunting of game animals (see supplementary material). These sites spanned the major forest habitats

of the region: ten in uplands spanning both banks of the Amazon River, six on floodplains, and four

in white-sand forest. We were unable to find intact examples of river islands large enough to

accommodate our sampling scheme. We then selected twenty disturbed sites in slash-and-burn

mosaic, each paired with an intact site for forest type, soil texture, and geographic proximity. At each

study site, we established six sampling points spaced by at least 210 meters to avoid double-counting

during avian point counts.

During subsequent vegetation assessment, we determined that six sampling points on

different transects were unsuitable for analysis due to their inadvertent location in transitional

habitat at the edge of the forest-type of interest. Flooding and time constraints prevented us from

sampling trees at two study sites (one in intact floodplain and another in intact uplands), and we

removed their paired disturbed sites from the tree dataset. Thus, the final dataset contained 234 bird

sampling points and 209 tree sampling points. See supplementary material for details of site

selection, site spacing, and site characteristics.

BIODIVERSITY DATA

We surveyed birds and trees at each sampling point. For birds, a single observer

(REDACTED) conducted four ten-minute 100-meter-radius point-counts at each sampling point

during July-December 2013-2014. Surveys ran from first light until mid-morning, and were not

conducted in rain or windy conditions. We visited most points in both years and rotated the visit

order to ensure that each point received early-morning coverage. To assemble our final dataset for

analysis, we aggregated data across the four visits to each point by taking the maximum count for

each species from any visit.

We made two modifications to standard point-count protocols, tailored to the challenges of

detecting skittish species and birds in mixed-species flocks (see supplementary material). First, we

included detections of species that flushed during our approach to and departure from each point

(within 100 m). Second, when mixed flocks detected during the point count lingered within 100

meters after the count period, we proceeded to follow the flock until we identified all of its

participants or until it moved >100 m from the point. We separately recorded individuals detected

via these modifications, permitting us to include them or exclude them from analysis (see

SENSITIVITY ANALYSIS, below).

To survey trees, we established a 50x2 m^2 tree plot at a fully randomized location within 100

m of each sampling point (equivalent to 0.6 Gentry transects per site; (Gentry 1988). Within these

plots, we identified every tree greater than 2.5 cm diameter at breast height. We collected a voucher

for each species (except for palms with very large leaves), deposited in the herbarium at the

Universidad Nacional de la Amazonía Peruana (UNAP). One botanist (REDACTED) conducted all

sampling and made all species determinations with reference to the UNAP herbarium collections.

See supplementary information for detailed bird and tree survey protocols.

BIODIVERSITY COMPARISONS

We used sample-based rarefaction to compare bird and tree richness in intact and disturbed

landscapes on a per-area basis (Chao et al. 2014). For trees, this revealed dramatic diversity loss due

to massively reduced densities of individuals at disturbed sites (i.e., cleared areas have fewer trees).

Therefore, we used individual-based rarefaction to test for a second-order effect of slash-and-burn

on tree diversity, controlling for the number of individuals sampled. For both birds and trees, we

performed rarefaction analysis on each forest type separately (upland, floodplain, white-sand) and

for all forest types combined. We also visualized patterns of community change using non-metric

multidimensional scaling.

Some bird species that we did not record at intact sites are well known to be common on

intact river-islands (Rosenberg 1990). As noted above, we were unable to sample intact river island

habitat because in our study area all accessible river islands large enough to accommodate our

sampling scheme have been settled, cleared, or otherwise disturbed by people. Therefore, we

conducted a follow-up analysis to account for bias related to the intact river-island avifauna. We

obtained a comprehensive list of bird species that were common on intact river-islands within the

study area thirty years ago (Rosenberg 1990). We then repeated our analysis while excluding these

species from all datasets, thereby removing their influence on our conclusions. We stress that we

selected these species not because they are prevalent in disturbed samples, but because they are

known to be prevalent in an intact river-island habitat that we were unable to sample. By removing

only common river-island species, we are confident that we removed very few species that would

not have appeared in the dataset for intact forest types, had we been able to sample river islands.

Therefore, this analysis mitigates bias in the comparison between intact and disturbed habitats.

POPULATION COMPARISONS

For every species of bird and tree in the dataset, we calculated Bayesian point-estimates and

95% credible intervals for the multiplicative change (fold-change) in abundance between intact and

disturbed sites. To do so, we assumed that the number of individuals detected at intact and

disturbed sites were realizations of Poisson processes. This implies that the total count at disturbed

sites is a binomial draw from the summed count at intact and disturbed sites, and furthermore that

the logarithm of the fold-change between the Poisson means is equal to the logit of the binomial

proportion p (Przyborowski & Wilenski 1940). We computed the posterior density of p using the

Jeffreys prior, and we used the posterior density of $p/(1-p)$ for inference on the fold-change (Brown

et al. 2001).

DISTRIBUTION OF DISTURBANCE-SENSITIVE SPECIES

To understand what features of disturbed points allow them to support species characteristic

of intact forests, we defined *disturbance-sensitive species* as those that are more abundant in intact forest

than disturbed forest, and *disturbance-sensitive counts* as the total number of individuals belonging to

disturbance-sensitive species detected at each point. We then fit generalized linear mixed models for

birds and trees to assess the relationship between disturbance-sensitive counts and local habitat data

(see below) across the disturbed points.

Local habitat data

At every sampling point, we recorded the number and size of streams and estimates of

percent cover of 10 vegetation formations within 100 m of the point (see supplementary material).

Using Landsat 8 imagery downloaded from the Global Forest Change Data website (Hansen et al

2013), we built a random-forest classifier of the study landscape as intact, disturbed, or open water at

30 m resolution. We validated our classification against the central coordinates of our 240 sampling

points, and then we extracted the area classified as intact within 200, 500, and 5000 m of each

disturbed sampling point. We also measured the distance from each disturbed point to the nearest

primary forest (continuously forested since 1985, before the acceleration of forest clearance in the

region; (Mäki et al. 2001) and to the nearest river (channel width > 30 m) based on visual

examination of Landsat imagery in the USGS Landsat Look viewer, supplemented with aerial

imagery in Google Earth.

Mixed models

Initially, we assumed that any species recorded in higher numbers in intact than disturbed

habitat is disturbance-sensitive. For birds and trees, we fit ordinary and zero-inflated Poisson and

negative binomial mixed models (treating study-site as a random effect) for the disturbance-sensitive

counts using a variety of predictors describing local vegetation cover at the 100 m scale, forest cover

at 0.2 − 5 km spatial scales, and proximity to major rivers (see supplementary material). We used the

small-sample corrected Akaike information criterion (AICc) to select covariates and error structure

that yielded parsimonious models, and we base inference on broad agreement across all top-

performing models.

To verify that our conclusions were robust to uncertainty in which species are disturbance-

sensitive, we re-analyzed the model with the lowest AICc score as follows. Using the binomial

likelihood described above, we computed the probability that each species in the dataset is

disturbance-sensitive by integrating the posterior distribution for the binomial proportion (based on

a uniform prior) from 0 to 0.5. We then randomly assigned each species to be disturbance-sensitive

or not based on these probabilities, re-computed the disturbance-sensitive counts, and fit the

regression model to these counts under a Bayesian mode of inference using Markov-chain Monte

Carlo sampling implemented in JAGS (Plummer 2003). We repeated this process 500 times,

combined the posterior chains for inference, and compared the resulting parameter estimates to the

corresponding frequentist estimates.

SENSITIVITY AND DETECTABILITY ANALYSIS FOR BIRDS

To ensure that our non-standard point-count methodology did not bias avian sampling, we

repeated our analyses using only detections obtained via standard protocols. We used an N-mixture

model to determined that avian detectability is likely to be at least as high in disturbed habitats as

intact habitats (see supplementary material). Therefore, if anything, our results overestimate the

biodiversity value of smallholder landscapes.

RESULTS

Across pristine and disturbed habitats combined we recorded 455 bird species and 751 tree

species; the bird dataset is among the richest single-observer point-count datasets ever assembled.

We found very high avian richness in slash-and-burn mosaics. In fact, in each habitat studied

(uplands, floodplain, white-sands), sample-based rarefaction revealed that bird richness at disturbed

sites was comparable to intact sites (Figure 2). However, tree richness declined severely. This decline

partly resulted from dramatic reductions in the number of individuals at disturbed sites (i.e. cleared

areas have fewer trees) but was exacerbated by changes in the species-abundance distribution (Figure

2).

Importantly, considering each habitat in isolation substantially underestimated the difference

in bird richness between intact and disturbed landscapes. Across habitats, reductions in beta-

diversity caused modest but significant declines in bird richness. Moreover, the apparent biodiversity

value of smallholder landscapes was substantially inflated by the spurious absence of river-island

species from our intact sites (an artifact of our inability to sample intact river islands). When the

influence of these poorly sampled river-island species is removed from both the intact and disturbed

points, it becomes apparent that intact landscapes have dramatically higher avian richness than

disturbed landscapes in our study region (Figure 2). This occurs because the river-island avifauna

overlaps more with disturbed habitats than with other intact habitats in the study area. We did not

observe a similar pattern in trees, though non-metric multidimensional scaling suggests that some

homogenization might have occurred (Figure 3). Instead, uplands dominated the tree species

richness of all intact sites combined, minimizing the opportunity for specialists in other habitats to

contribute to richness patterns (Figure 2).

Disturbed sites consistently clustered separately from intact sites in terms of their species

composition, and non-metric multidimensional scaling of community composition revealed that the

difference between intact and disturbed sites corresponded to the first axis of variation (Figure 3).

The second axis of variation, corresponding to an edaphic gradient from floodplains through

uplands to white sands, was collapsed at disturbed sites, reflecting the loss of beta-diversity among

forest types. These patterns are consistent for birds and trees and for a variety of incidence- and

abundance-based dissimilarity metrics (Figure S5). Thus, disturbance in addition to driving species

loss, smallholder agriculture drives the disassembly and re-arrangement of primary forest bird

communities.

Furthermore, large numbers of disturbance-sensitive species showed dramatically reduced

abundance at disturbed sites (Figure 4). For example, we detected the Screaming Piha (*Lipaugus*

vociferans) 137 times at intact sites, and only once at disturbed sites. Similarly, we detected the tree

Eschweilera coriacea (Lecythidaceae) thirty-one times at intact sites and only once at disturbed sites. In

the rarefaction analysis, such species contribute to the disturbed-site total, but in fact they are

severely harmed by slash-and-burn practices. Among the 249 bird and 221 tree species for which we

detected a significant change in abundance, 57% and 86% declined, respectively. Of the birds that

significantly increased in abundance in our dataset, fully 39% are common on intact river islands

(Rosenberg 1990). Failure to detect significant abundance changes was generally a consequence of

low sample size (and probably not a consequence of small effect size). The median sample size

among species without a significant effect was two for birds and one for trees.

Mixed models revealed a major positive influence of local forest cover and nearby primary

forest on the abundance of disturbance-sensitive birds and trees that was consistent across all well-

performing models (Table 1). For birds, the most important components of this effect were primary

forest cover at a radius of 5 km and secondary forest cover at a radius of 100 m. For trees, the key

components were secondary forest cover at a radius of 100 m and primary forest cover at a radius of

200 m. These effects were robust despite uncertainty in which species are disturbance-sensitive.

DISCUSSION

Our results constitute the first large-scale biodiversity assessment of slash-and-burn

agriculture in western Amazonia, and one of the first biodiversity assessments in degraded

Amazonian landscapes to explicitly consider multiple natural habitat types. These features define a

key knowledge gap for conservation science, because western Amazonia is the epicenter of terrestrial

biodiversity on Earth (Jenkins et al. 2013), harbors multiple types of forest, is heavily affected by

slash-and-burn agriculture (Finer & Novoa 2016), and features extensive species turnover (beta-

diversity) between natural habitats (Tuomisto et al. 1995; Pomara et al. 2012).

Our results are sobering. Diversity loss, community turnover, and large numbers of

declining, disturbance-sensitive species characterize the transition from intact forest to slash-and-

burn mosaic. Slash-and-burn agriculture collapses avian beta-diversity across forest-types, and this

process drives substantial reductions in gamma-diversity for birds. Because slash-and-burn mosaics

are as diverse as primary forest within any single forest type (e.g. within upland foreest), previous

work was unable to detect this decline (Andrade & Torgler 1994; Korasaki et al. 2013). Previous

studies have treated an environmental domain equivalent to the first panel of our Figure 2.

Moreover, within the slash-and-burn mosaic, secondary forest cover and proximity to

primary forest were consistent, strong predictors of the occurrence of disturbance-sensitive species.

According to our vegetation classifier, the median disturbed point in our dataset was surrounded by

over 19% primary forest at a radius of 0.2 km, 28% at 0.5 km, and 57 % at 5 km. The proximity of

intact habitat, coupled with the high heterogeneity and low land-use intensity of the slash-and-burn

mosaic (the median disturbed point contained 30% closed-canopy secondary-forest cover within a

100 m radius), strongly suggests that our results are a best-case scenario for biodiversity in

Amazonian smallholder agriculture. The conservation value of slash-and-burn mosaics in our study

area depends on extensive fallow areas (i.e. secondary forests) and spillover from primary forest.

Recent work from elsewhere in Amazonia suggests that the biodiversity impacts of

smallholder agriculture might be even more severe than our methods can detect. Space-for-time

substitutions might underestimate the severity of impacts in Amazonia (França et al. 2016), perhaps

due to inadequate primary-forest baseline data. Furthermore, the negative impacts of agricultural

disturbance can spill across into adjacent primary forest, leading to substantial additional losses of

conservation value (Barlow et al. 2016). In our study area, a few species (e.g. curassows in the genus

Mitu) are so heavily hunted that they are absent even at intact sites, and our analyses cannot shed

light on their disturbance-sensitivity.

Some implications of our results extend beyond the western Amazon. In particular, we note

that many previous comparisons of biodiversity value at intact and degraded tropical sites have been

restricted to a single natural habitat (or have analyzed multiple habitats separately), with variable

results (e.g Daily et al. 2003; Peh et al. 2006; Ranganathan et al. 2008; Phalan et al. 2011; Kurz et al.

2014). Our results show that the large-scale pattern across multiple habitats is gloomier than single-

habitat results suggest, at least for birds. This conclusion is consistent with the observation that

smallholder agriculture reduces pairwise avian compositional dissimilarities across biogeographic

regions of Costa Rica (Karp *et al.* 2012). We expand on this result by showing that the

homogenization produced by smallholder agriculture drives substantial losses of regional gamma-

diversity (this is not a forgone conclusion; see Socolar *et al.* 2016). Moreover, we show that

homogenizing effects are important not only across widely spaced biogeographic regions, but also

across fine-scale habitat formations that structure Amazonian communities. The vast majority of

biodiversity assessments of Neotropical agriculture have focused on uplands and therefore missed

the additional biodiversity losses driven by homogenization across forest-types. Habitat differences

within biogeographic regions are globally ubiquitous (e.g. due to variation in elevation, soils,

hydrology, climate, etc), and revealing the full impacts of disturbance requires sampling that is both

spatially extensive and locally comprehensive with respect to habitat variation (Gardner et al. 2013;

Solar et al. 2015). Habitat specialization and spatial turnover are characteristic of hyperdiverse

species communities, suggesting that habitat degradation might have its worst effects precisely where

biodiversity is highest.

We also note that previous studies of biodiversity in Neotropical agricultural landscapes have

broadly neglected trees. A recent meta-analysis of the biodiversity value of degraded tropical

landscapes was unable to include a single study of tree diversity in Neotropical agriculture (Gibson

et al. 2011). This situation might arise because lower tree diversity in cleared areas is perceived as a

forgone conclusion (more attention has been paid to shrubs and forbs, e.g. (Mayfield & Daily 2005).

Nevertheless, trees make up a critical component of tropical biodiversity, and maintaining tropical

tree diversity is likely essential for the long-term conservation a variety of coevolved species (Koh et

al. 2004). Moreover, the impacts of agriculture on tree diversity are even more severe than could be

predicted by declines in abundance alone; agricultural landscapes are species-poor even after

controlling for the number of individual trees sampled. Thus, field inventories of tree communities

are crucial for accurately assessing the biodiversity consequences of slash-and-burn agriculture, and

our results paint a bleak picture.

We do not mean to dismiss innovative efforts, including efforts inside protected areas, to

harmonize conservation objectives with the livelihoods of local people (Pulgar Vidal & Gamboa

Moquillaza 2013). There is a clear humanitarian mandate for such efforts, and they can prevent the

even greater losses of biodiversity that result from the conversion of disturbed forests and

agricultural mosaics to soy monocultures or tree plantations. However, we do mean to sound the

alarm over the potential consequences of ongoing smallholder expansion. There will be severe

biodiversity losses if settlers gain access to the last remaining tropical wildernesses in western

Amazonia, no matter how lightly they tread.

ACKNOWLEDGMENTS

REFERENCES

Alroy J. 2017. Effects of habitat disturbance on tropical forest biodiversity. Proceedings of the
National Academy of Sciences **114**:6056–6061.
Alvarez Alonso J, Metz MR, Fine PV. 2013. Habitat specialization by birds in western Amazonian
white-sand forests. Biotropica **45**:365–372. Wiley Online Library.
Andrade GI, Torgler HR. 1994. Sustainable use of the tropical rain forest: evidence from the

avifauna in a shifting-cultivation habitat mosaic in the Colombian Amazon. Conservation
Biology **8**:545–554.
Barlow J et al. 2007. Quantifying the biodiversity value of tropical primary, secondary, and plantation
forests. Proceedings of the National Academy of Sciences of the United States of America
104:18555–18560.
Barlow J et al. 2016. Anthropogenic disturbance in tropical forests can double biodiversity loss from
deforestation. Nature **535**:144–147. Nature Publishing Group.
Berry NJ, Phillips OL, Ong RC, Hamer KC. 2008. Impacts of selective logging on tree diversity
across a rainforest landscape: the importance of spatial scale. Landscape Ecology **23**:915–929.
Brown LD, Cai TT, DasGupta A. 2001. Interval estimation for a binomial proportion. Statistical
science **16**:101–117.
Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014. Rarefaction and
extrapolation with Hill numbers: a framework for sampling and estimation in species diversity
studies. Ecological Monographs **84**:45–67.
Daily GC, Ceballos G, Pacheco J, Suzán G, Sánchez-Azofeifa A. 2003. Countryside biogeography of
neotropical mammals: conservation opportunities in agricultural landscapes of Costa Rica.
Conservation Biology **17**:1814–1826. Wiley Online Library.
Daily GC, Ehrlich PR, Sanchez-Azofeifa A. 2001. Countryside biogeography: use of human-
dominated habitats by the avifauna of southern Costa Rica. Ecological Applications **11**:1–13.
Espinoza JC, Ronchail J, Frappart F, Lavado W, Santini W, Guyot JL. 2013. The major floods in the
Amazonas river and tributaries (western Amazon basin) during the 1970–2012 period: a focus
on the 2012 flood. Journal of Hydrometeorology **14**:1000–1008.
Ferraz G, Nichols JD, Hines JE, Stouffer PC, Bierregaard RO, Lovejoy TE. 2007. A large-scale
deforestation experiment: effects of patch area and isolation on Amazon birds. Science **315**:238–
241.
Fine PVA, García-Villacorta R, Pitman NCA, Mesones I, Kembel SW. 2010. A floristic study of the
white-sand forests of Peru. Annals of the Missouri Botanical Garden **97**:283–305.
Finer M, Novoa S. 2016. Large-scale vs. small-scale deforestation in the Peruvian Amazon. MAAP
32.
Finer M, Orta-Martínez M. 2010. A second hydrocarbon boom threatens the Peruvian Amazon:
trends, projections, and policy implications. Environmental Research Letters **5**:014012.
França F, Louzada J, Korasaki V, Griffiths H, Silveira JM, Barlow J. 2016. Do space-for-time
assessments underestimate the impacts of logging on tropical biodiversity? An Amazonian case
study using dung beetles. Journal of Applied Ecology **53**:1098–1105.
Gardner TA et al. 2013. A social and ecological assessment of tropical land uses at multiple scales:
the Sustainable Amazon Network. Philosophical Transactions of the Royal Society B: Biological
Sciences **368**:20120166–20120166.
Gentry AH. 1988. Tree species richness of upper Amazonian forests. Proceedings of the National
Academy of Sciences **85**:156–159.
Giam X. 2017. Global biodiversity loss from tropical deforestation. Proceedings of the National
Academy of Sciences of the United States of America **114**:5775–5777. National Acad Sciences.
Gibson L et al. 2011. Primary forests are irreplaceable for sustaining tropical biodiversity. Nature
478:378–381. Nature Publishing Group.
Higgins MA, Ruokolainen K, Tuomisto H, Llerena N, Cardenas G, Phillips OL, Vásquez R,
Räsänen M. 2011. Geological control of floristic composition in Amazonian forests. Journal of
Biogeography **38**:2136–2149.
Jenkins CN, Pimm SL, Joppa LN. 2013. Global patterns of terrestrial vertebrate diversity and

conservation. Proceedings of the National Academy of Sciences **110**:E2602–E2610. National
Acad Sciences.
Karp DS, Rominger AJ, Zook J, Ranganathan J, Ehrlich PR, Daily GC. 2012. Intensive agriculture
erodes β-diversity at large scales. Ecology Letters **15**:963–970.
Karp DS, Ziv G, Zook J, Ehrlich PR, Daily GC. 2011. Resilience and stability in bird guilds across
tropical countryside. Proceedings of the National Academy of Sciences **108**:21134–21139.
National Acad Sciences.
Koh LP, Dunn RR, Sodhi NS, Colwell RK, Proctor HC, Smith VS. 2004. Species coextinctions and
the biodiversity crisis. Science **305**:1632–1634.
Korasaki V, Braga RF, Zanetti R, Moreira FMS, Vaz-de-Mello FZ, Louzada J. 2013. Conservation
value of alternative land-use systems for dung beetles in Amazon: valuing traditional farming
practices. Biodiversity and Conservation **22**:1485–1499.
Kurz DJ, Nowakowski AJ, Tingley MW, Donnelly MA, Wilcove DS. 2014. Biological Conservation.
Biological Conservation **170**:246–255. Elsevier Ltd.
Laurance WF et al. 2014. A global strategy for road building. Nature:1–7. Nature Publishing Group.
Lees AC, Moura NG, de Almeida AS, Vieira ICG. 2015. Poor Prospects for Avian Biodiversity in
Amazonian Oil Palm. PloS one **10**:e0122432.
MacArthur R, Levins R. 1967. The limiting similarity, convergence, and divergence of coexisting
species. American Naturalist **101**:377–385. JSTOR.
Mahood SP, Lees AC, Peres CA. 2011. Amazonian countryside habitats provide limited avian
conservation value. Biodiversity and Conservation **21**:385–405.
Mayfield MM, Daily GC. 2005. Countryside biogeography of neotropical herbaceous and shrubby
plants. Ecological Applications **15**:423–439. Eco Soc America.
Mäki S, Kalliola R, Vuorinen K. 2001. Road construction in the Peruvian Amazon: process, causes
and consequences. Environmental Conservation **28**:199–214. Cambridge Univ Press.
Mendenhall CD, Sekercioglu CH, Brenes FO, Ehrlich PR, Daily GC. 2011. Predictive model for
sustaining biodiversity in tropical countryside. Proceedings of the National Academy of Sciences
108:16313–16316. National Acad Sciences.
Moura N, Lees AC, Andretti CB, Davis BJW, Solar RRC, Aleixo A, Barlow J, Ferriera J, Gardner
TA. 2013. Avian biodiversity in multiple-use landscapes of the Brazilian Amazon. Biological
Conservation **167**:339–348. Elsevier Ltd.
Moura NG, Lees AC, Aleixo A, Barlow J, Berenguer E, Ferreira J, Mac Nally R, Thomson JR,
Gardner TA. 2016. Idiosyncratic responses of Amazonian birds to primary forest disturbance.
Oecologia **180**:903–916. Springer Berlin Heidelberg.
Newbold T, Scharlemann JPW, Butchart SHM, Sekercioglu CH, Joppa L, Alkemade R, Purves DW.
2014. Functional traits, land-use change and the structure of present and future bird
communities in tropical forests. Global Ecology and Biogeography **23**:1073–1084.
Peh KSH, Sodhi NS, de Jong J, Sekercioglu C, Yap CAM, Lim SLH. 2006. Conservation value of
degraded habitats for forest birds in southern Peninsular Malaysia. Diversity and Distributions
12:572–581.
Phalan B, Onial M, Balmford A, Green RE. 2011. Reconciling Food Production and Biodiversity
Conservation: Land Sharing and Land Sparing Compared. Science **333**:1289–1291.
Plummer M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs
sampling. in K. Hornik, F. Leisch, and A. Zeileis, editors. Vienna, Austria.
Pomara LY, Ruokolainen K, Tuomisto H, Young KR. 2012. Avian composition co-varies with
floristic composition and soil nutrient concentration in Amazonian upland forests. Biotropica
44:545–553.
Przyborowski J, Wilenski H. 1940. Homogeneity of results in testing samples from Poisson series:

With an application to testing clover seed for dodder. Biometrika **31**:313–323.

Pulgar Vidal M, Gamboa Moquillaza P. 2013. Reserva Nacional Allpahuayo Mishana Plan Maestro
2013 - 2018. Pages 1–24. Lima, Peru.

Ranganathan J, Daniels RJR, Chandran MDS, Ehrlich PR, Daily GC. 2008. Sustaining biodiversity in
ancient tropical countryside. Proceedings of the National Academy of Sciences **105**:17852–
17854. National Acad Sciences.

Ravikumar A, Sears RR, Cronkleton P, Menton M, Pérez-Ojeda del Arco M. 2017. Is small-scale
agriculture really the main driver of deforestation in the Peruvian Amazon? Moving beyond the
prevailing narrative. Conservation Letters **10**:170–177.

Remsen JV Jr, Parker TA III. 1983. Contribution of river-created habitats to bird species richness in
Amazonia. Biotropica:223–231. JSTOR.

Rosenberg GH. 1990. Habitat specialization and foraging behavior by birds of Amazonian river
islands in northeastern Peru. The Condor:427–443. JSTOR.

Sekercioglu C, Loarie SR, Oviedo Brenes F, Ehrlich PR, Daily GC. 2007. Persistence of forest birds
in the Costa Rican agricultural countryside. Conservation Biology **21**:482–494.

Socolar J, Robinson SK, Terborgh JW. 2013. Bird Diversity and Occurrence of Bamboo Specialists
in Two Bamboo Die-Offs in Southeastern Peru. The Condor **115**:253–262.

Socolar JB, Gilroy JJ, Kunin WE, Edwards DP. 2016. How should beta-diversity inform biodiversity
conservation? Trends in Ecology & Evolution **31**:67–80. Elsevier Ltd.

Solar RR de C et al. 2015. How pervasive is biotic homogenization in human-modified tropical
forest landscapes? Ecology Letters **18**:1108–1118.

Steege ter H et al. 2013. Hyperdominance in the Amazonian tree flora. Science **342**:1243092–
1243092.

Tuomisto H, Ruokolainen K, Kalliola R, Linna A, Danjoy W, Rodriguez Z. 1995. Dissecting
Amazonian biodiversity. Science **269**:63–66.

Tyukavina A, Hansen MC, Potapov PV, Krylov AM, Goetz SJ. 2015. Pan-tropical hinterland forests:
mapping minimally disturbed forests. Global Ecology and Biogeography **25**:151–163.

Wittmann F, Junk WJ, Piedade MTF. 2004. The várzea forests in Amazonia: flooding and the highly
dynamic geomorphology interact with natural forest succession. Forest Ecology and
Management **196**:199–212.

Wittmann F, Schongart J, Montero JC, Motzer T, Junk WJ, Piedade MTF, Queiroz HL, Worbes M.
2006. Tree species composition and diversity gradients in white-water forests across the
Amazon Basin. Journal of Biogeography **33**:1334–1347.

FIGURE AND TABLE CAPTIONS

Figure 1: In western Amazonia, slash-and-burn agriculture converts primary forest (**a**) to various

disturbed habitats across uplands, floodplains, and white-sands, resulting in a heterogeneous mosaic

of secondary habitats. Shown here are tangles following abandonment of a floodplain agricultural

plot (**b**), barren ground and scrub following agricultural abandonment on white-sands (**c**), and a

mosaic of upland secondary forest and active agricultural plots (**d**). (**b-d**) represent the range of

slash-and-burn habitats in a highly diversified mosaic, not typical differences between soil types.

Figure 2: Sample-based rarefaction (mean and 95% confidence interval) for birds (top row) shows

that within each forest type, disturbed sites are as species-rich as their intact counterparts. However,

when forest-types are aggregated, intact forest is more diverse, especially after accounting for the

distribution of river-island species. For trees (bottom row), individual-based rarefaction shows that

richness plummets in disturbed forests, due to low individual abundance and a second-order effect

of altered species-abundance distributions after controlling for the number of individuals sampled.

Aggregated forest-types do not show greater tree-richness differences than individual forest-types;

instead, the uplands dominate the species pool and are effectively as diverse as all habitats combined.

Figure 3: Non-metric multidimensional scaling (NMDS) based on Raup-Crick dissimilarities for

point-scale bird data (top; stress = 0.21) and site-scale tree data (bottom; stress = 0.22; point-scale

tree data were too sparse for NMDS). In both cases, the first NMDS axis captures the difference

between intact and disturbed sites, while the second axis captures the gradient from nutrient-rich

floodplains to nutrient-poor white-sands. Intact and disturbed sites segregate almost completely.

Heterogeneity between forest types at intact sites is collapsed at disturbed sites.

Figure 4: Multiplicative changes in abundance for birds (top) and trees (bottom). Species

significantly different from one (i.e. abundance differs by land-use class) are given by dark points

with 95% credible intervals. Most tree species plummet in abundance. Bird communities include

species that fare well following disturbance, but 57% of species with significant abundance-changes

declined, often dramatically (note the logarithmic y-axis).

Table 1: Results of models for counts of disturbance-sensitive birds and trees, summarizing results

for the top performing model (credible intervals for the effect size) and for all models within 2 AICc

units of the top performing model (frequency of inclusion and sign of effect). Forest type was

included in all models as a control. All of the best-performing bird models used a NB1 negative

binomial error structure without zero-inflation. Five of the best-performing tree models (including

the top model) used a zero-inflated NB2 negative binomial error structure, one used NB2 error

without zero-inflation.

Predictor/Interpretation	Birds			Trees		
	freq†	sign‡	95% CI*	freq†	sign‡	95% CI*
ForestType: Upland		-	-0.706 — 0.695		-	-3.285 — 0.002
ForestType: Floodplain		-	-1.268 — 0.366		-	-3.939 — -0.026
Secondary forest (% cover within 100 m)	11/11	+	0.002 — 0.015	6/6	+	0.004 — 0.032
Tall secondary forest: canopy > 20 m (% cover within 100 m)	11/11	+	0.010 — 0.022	1/6	+	
Scrub/gap lacking closed canopy > 5 m tall (% cover within 100 m)	5/11	-		0/6		
Non-habitat: water, grass, dirt, buildings (% cover within 100 m)	6/11	-	-0.020 — -0.001	0/6		
Streams: width (m) of incised channel within 100 m	4/11	+		1/6	+	
intact200: % cover of primary forest within 200 m	3/11	+		6/6	+	1.491 — 7.464
intact500: % cover of primary forest within 500 m	0/11			0/6		
intact5000: % cover of primary forest within 5000 m	11/11	+	0.417 — 2.529	0/6		
PrimaryDist: distance (km) to nearest primary forest	0/11			1/6	-	
RiverDist: distance (km) to nearest river > 30 m wide	11/11	-	-0.222 — -0.022	1/6	+	

551 † Frequency of predictor's inclusion among models within two AICc units of the top-performing model.

552 ‡ No predictor for either birds or trees entered multiple models within two AICc units of the top-performing model with opposite signs.

553 * 95% Bayesian credible interval from top-performing model, accounting uncertainty in which species are disturbance-sensitive.