Histone deacetylase inhibition reduces deleterious cytokine release induced by ingenol stimulation

Erin T. Larragoite[*1], Racheal A. Nell[*2], Laura J. Martins[1], Louis R. Barrows[3], Vicente Planelles[1], Adam M. Spivak[2§]

[*]Contributed equally

Authors' Affiliation: [1]Department of Pathology, University of Utah, Salt Lake City, United States; [2]Department of Medicine, University of Utah School of Medicine, Salt Lake City, United States, [3]Department of Pharmacology and Toxicology, University of Utah, Salt Lake City, United States

[§] Corresponding Author:

Adam M. Spivak

50 North Medical Drive, Division of Infectious Diseases, Room 4B319, Department of Medicine, University of Utah School of Medicine, Salt Lake City, UT 84132

Phone: 801-587-1964

Fax: 801-585-3377

adam.spivak@hsc.utah.edu

ETL erin.larragoite@path.utah.edu

RAN racheal.nell@outlook.com

LJM laura.martins@path.utah.edu

LRB lbarrows@pharm.utah.edu

VP vicente.planelles@path.utah.edu

AMS adam.spivak@hsc.utah.edu

Abstract

Introduction: Latency reversal agents (LRAs), such as protein kinase C (PKC) agonists, constitute a promising strategy for exposing and eliminating the HIV-1 latent reservoir. PKC agonists activate NF-κB and, in turn, induce deleterious pro-inflammatory cytokine production. Adjuvant pharmacological agents, such as ruxolitinib, a JAK inhibitor, and rapamycin, an mTOR inhibitor, have previously been combined with LRAs to reduce deleterious pro-inflammatory cytokine secretion without inhibiting HIV-1 viral reactivation in vitro. Histone deacetylase inhibitors (HDACi) are known to dampen pro-inflammatory cytokine secretion in the context of other diseases and can synergize with other LRAs to bring dormant proviruses out of latency. In this study we investigated whether a broad panel of epigenetic modifiers, including HDACi, could effectively dampen PKC-induced pro-inflammatory cytokine secretion during latency reversal.

Methods: We screened an epigenetic modifier library to identify compounds that reduced intracellular IL-6 production induced by the PKC agonist Ingenol-3,20-dibenzoate. We further tested the most promising epigenetic inhibitor class, histone deacetylase inhibitors, for their ability to reduce a broad panel of pro-inflammatory cytokines *ex vivo*.

Results: We identified nine epigenetic modulators that reduced PKC-induced intracellular IL-6. In cells from aviremic individuals living with HIV-1, the HDAC1-3 inhibitor, suberohydroxamic acid (SBHA), reduced secretion of pro-inflammatory cytokines TNF-α, IL-5, and IL-17 when used in combination with Ingenol-3,20-dibenzoate.

Conclusion: The addition of SBHA to Ingenol-3,20-dibenzoate reduces deleterious cytokine production during latency reversal in aviremic donor PBMCs. The ability of SBHA to reduce

PKC-induced pro-inflammatory cytokines when used in combination with Ingenol-3,20-dibenzoate makes the combination an attractive latency reversal therapy.

Keywords: HIV-1; Latency Reversing Agent; Ingenol-3,20-dibenzoate; suberohydroxamic acid; cytokine; HDACi

1. Introduction

Antiretroviral therapy (ART) has improved the lives of individuals living with HIV-1 by durably blocking viral replication, reducing plasma viremia levels below the limit of detection (~20-50 copies/ml) [1-3], and allowing for reconstitution of the adaptive immune system. However, ART does not result in HIV-1 eradication due to the presence of a transcriptionally silent, long-lived latent viral reservoir [4-6]. The presence of this latent reservoir makes ART a lifelong necessity [7]. Additionally, not all patients have access to ART or are compliant with ART regimens due to high costs [8], side effects [9], and social stigma [10, 11]. Aviremic individuals taking ART are prone to low-level chronic immune activation and exhaustion [12], cardiovascular disease [13, 14], non-AIDS related cancers [15], and neurologic complications [16], underscoring the importance of strategies targeting the latent HIV-1 reservoir.

The "shock and kill" strategy [17] proposes that a latency reversing agent (LRA) be used to transcriptionally reactivate the latent reservoir followed by targeting infected cells with immune-mediated mechanisms [17, 18]. Immunological strategies include the use of broadly neutralizing antibodies (bNAbs), cytokines, vaccines, chimeric antigen receptors, and other immune-mediated mechanisms to enhance recognition of HIV-1 infected cells [19, 20]. While all

strategies hold merit, enhanced targeting of latently infected cells via immune-mediated mechanisms would still be dependent on unmasking the latent reservoir.

Several mechanistic classes have emerged as potential LRAs including positive transcription elongation factor b (PTEFb) activators [21, 22], cytokines [23], histone deacetylase inhibitors (HDACi) [24-28], and protein kinase C (PKC) agonists [18, 28-30]. Spina *et al.* [29] demonstrated that PKC agonists are uniquely efficacious LRAs across multiple in vitro models of HIV-1 latency, whereas other well characterized LRAs such as HDACi had functionality only in select in vitro models. PKC agonists are thought to reactivate latent HIV-1 though activation of NF-κB and have been shown to additionally activate AP-1 and NFAT [31]. On the other hand, PKC agonists have undesired effects, including induction of T cell activation [32] and pro-inflammatory cytokine secretion. We previously demonstrated that latency reversal with the PKC agonist Ingenol-3,20-dibenzoate, used in combination with a second pharmacological agent, ruxolitinib (an FDA approved Janus Kinase [JAK] inhibitor), led to potent reactivation of latent HIV-1 in the absence of pro-inflammatory cytokine secretion[33]. Additionally, rapamycin, an inhibitor of mTOR, has been shown to reduce pro-inflammatory cytokine secretion induced upon reactivation of latent HIV-1 with CD3 and CD28 antibodies [34]. These studies indicate that latency reversal and pro-inflammatory cytokine secretion can be uncoupled. In the present study we expanded our search for epigenetic modifiers that could be used as an adjuvant therapy to reduce cytokine secretion induced by PKC agonists.

While ruxolitinib and rapamycin have been shown to reduce in vivo cytokine secretion in the context of HIV-1, HDACi have also been observed to inhibit pro-inflammatory cytokine secretion in vivo in the context of graft-versus-host disease [35] and rheumatoid arthritis [36] as well as in vitro, in the context of LPS stimulation of human peripheral blood mononuclear cells

(PBMCs) [37]. Additionally, HDACi are well-known HIV LRAs and it has been shown that they can have additive or synergistic activity with PKC agonists when reactivating latent HIV-1 [27, 38, 39]. Therefore, we hypothesized that HDACi and potentially other epigenetic modifiers could decrease PKC-induced pro-inflammatory cytokines secretion and be used as a combination therapy with PKC agonists during latency reversal. In this study, we performed a screen of epigenetic modifiers and report that suberohydroxamic acid (SBHA), an HDAC1- and 3-inhibitor, suppresses PKC-induced pro-inflammatory cytokines.

2. Materials and Methods

Participants

Participants living with and without HIV-1 infection were recruited in accordance with active University of Utah Institutional Review Board (IRB) protocols 58246 and 67637 respectively. Participants living with HIV-1 infection were aviremic (plasma viral loads less than 50 HIV-1 RNA copies/mL) for a minimum of 6 months and prescribed an ART regimen initiated during chronic HIV-1 infection for a minimum of 12 months.

In vitro epigenetic inhibitor screening

PBMCs were isolated from healthy donors using a Lymphoprep density gradient (Cat# 07861, StemCell Technologies) prior to being cultured in RPMI medium supplemented with 10% FBS and 5 U/mL penicillin/streptomycin overnight to remove monocytes via adherence. Non-adherent PBMCs were then cultured at a density of 1×10^5 cells/100µL in the presence of 100nM compounds from a Cayman Chemical Epigenetics Screening Library obtained from the

University of Utah Drug Discovery Core Facility (Item No. 11076) or 100nM ruxolitinib (Cayman Chemical Co., Item No. 11609), a control for cytokine inhibition, for 1.5 h at 37°C. Cells were then incubated in medium alone or the presence of 100nM IDB for 40 hours post-stimulant exposure. At 40 hours post-stimulant exposure, 0.067 µL/100µL BD GolgiStop™ Protein Transport Inhibitor (Cat# 554724) was added to each sample to inhibit cytokine secretion. Cells were then fixed and stained at 48 hours prior to flow cytometry analysis.

Intracellular cytokine staining and flow cytometry

Cells were washed with 1x phosphate buffered saline (PBS) prior to staining the cells with 0.1µL/100 µL Fixable Viability Dye eFluor® 450 (Cat# 65-0863-14, Affymetrix eBioscience) for 30 min at 4°C. Cells were then washed with 1x PBS prior to fixing with 100 µL BD Cytofix/Cytoperm™ for 30 min at 4°C. Once cells were fixed, they were washed with a perm/wash solution (1x PBS, 3% FBS, 0.1% Saponin, 0.05% Sodium Azide) prior to staining cells in 100 µL perm/wash with 0.5 µL APC anti-human IL-6 antibody (Cat# 501112, Biolegend®) overnight at 4°C. Cells were finally washed in 1x PBS and re-suspended in PBS prior to flow cytometry [BD FACSCanto™ flow cytometer with FACSDiva™ acquisition software (Becton-Dickinson, Mountain View, CA)] and analysis with FlowJo (TreeStar Inc, Ashland, OR).

Selection of compounds

The percentage of IL-6-positive cells in Ingenol-3,20-dibenzoate-alone-treated cells was compared to cells treated with epigenetic modifiers and IDB in order to calculate intracellular IL-6 fold-change. The top epigenetic modifiers that reduced intracellular IL-6 by four-fold or

greater ('hits') were selected for testing in aviremic patient cells to determine whether these compounds would diminish PKC induced pro-inflammatory cytokines.

Ex vivo cell culture, qPCR, and cytokine measurements

PBMCs were isolated from donors living with HIV-1 infection prior to the isolation of resting CD4$^+$ T cells (rCD4s) (EasySep^{TM} Human Resting CD4$^+$ T Cell Isolation Kit, Cat # 17962, StemCell Technologies). rCD4s were cultured at a density of 5 x 10^6/mL in RPMI supplemented with 10% FBS and 5U/mL penicillin/streptomycin. Epigenetic modifiers (Chidamide, Apicidin, Entinostat, Suberohydroxamic Acid (SBHA), Pyroxamide, Panobinostat, WDR5-0103, Iniparib and CCG-100602) that reduced intracellular IL-6 were added at 50nM to rCD4s in the presence of medium or IDB (45nM) for 48 hours. Medium or DMSO and Dynabead^{TM} Human T-Activator αCD3/αCD28 (αCD3/αCD28) (Cat# 11132D, Gibco^{TM}) stimulated cells were used as controls. At 48 hours post-stimulation, supernatant was collected and the rapid ex vivo evaluation of anti-latency assay (REVEAL) assay was performed as previously described [28] to quantify viral release. Additionally, PBMCs from HIV-1 positive aviremic donors were cultured at a density of 3 x 10^6/ml in RPMI supplemented with 10% FBS and 5U/mL penicillin/streptomycin. The epigenetic modifiers that induced the highest level of reactivation when combined with IDB (SBHA, Panobinostat, and Pyroxamide) were added at 75nM or 100nM concentrations in the presence of medium or IDB for 48 hours. Cells cultered in medium alone were used as controls. Supernatant was collected and sent to ARUP Laboratories to measure ex vivo cytokine secretion via a commercially available quantitative multiplex bead assay to measure the following cytokines: interferon gamma (IFN-γ), tumor necrosis factor alpha

(TNF-α), interleukin (IL) 1 beta (IL-1β), IL-2, soluble IL-2 receptor (IL-2r), IL-4, IL-5, IL-6, IL-8, IL-10, IL-12, and IL-13.

Statistical Analysis

Significant changes in reactivation or the secretion of pro-inflammatory cytokines in cells treated with epigenetic modifiers in combination with IDB compared to IDB alone was calculated with GraphPad Prism Version 5.0f (GraphPad Software, Inc., San Diego CA). Statistical significance was calculated using a non-parametric Wilcoxon matched-pairs signed rank test.

3. Results

HDACi have been previously described to inhibit cytokine secretion [35-37]. Therefore, we investigated the ability of HDACi and other epigenetic modifiers to reduce pro-inflammatory cytokine secretion upon latency reversal with a PKC agonist. We conducted a screen of 96 epigenetic modifiers (Cayman Chemical Co.) in primary CD4$^+$ T cells to identify those that more efficiently quelled the induction of pro-inflammatory cytokines during latency reversal with ingenol-3,20-dibenzoate (IDB). We used IL-6 as a prototypical cytokine for the screen. Nine epigenetic modifiers were identified which reduced intracellular IL-6 levels by four-fold or greater in the context of IDB stimulation (**Figure 1**). These 'hits' include six HDACi (Chemical Abstracts Service (CAS) # 743420-02-2 (chidamide), CAS # 183506-66-3 (apicidin), CAS # 209783-80-2 (entinostat), CAS # 38937-66-5 (suberohydroxamic acid (SBHA)), CAS # 382180-17-8 (pyroxamide), CAS # 404950-80-7 (panobinostat)), one mixed lineage leukemia (MLL) inhibitor Cas # 890190-22-4 (WDR5-0103), and two compounds categorized as 'miscellaneous'

as they did not have the same function as other epigenetic modifiers in the library. Out of the two compounds categorized as miscellaneous, compound Cas # 160003-66-7 (Iniparib), has an unknown function and compound Cas # 1207113-88-9 (CCG-100602) is an inhibitor of Rho pathway-mediated signaling [40]. The positive control, ruxolitinib, also reduced intracellular IL-6 by greater than fourfold as previously shown [33]. As HDACi accounted for six out of nine 'hits', HDACi were selected for further validation ex vivo on cells from aviremic individuals living with HIV. The top three HDACi SBHA, Panobinostat, and Pyroxamide were determined to be the most promising epigenetic modifiers to reduce pro-inflammatory cytokine production in the context of IDB stimulation. We examined SBHA and Panobinostat for their ability to reduce secretion of a broad panel of pro-inflammatory cytokines ex vivo (**Figure 2**).

Treatment of aviremic patient PBMCs with IDB significantly increased TNF-α (p = 0.0156), IFN-γ (p = 0.0312), IL-1β (p =0.0156), IL-5 (p = 0.0156), IL-8 (p = 0.0156), IL-13 (p = 0.0156), and IL-17 (p = 0.0156) (**Figure 2A**). The addition of 75nM or 100nM SBHA to 100nM IDB was found to significantly reduce secreted levels of TNF-α (p = 0.0078), IL-5 (p = 0.0156), IL-2r (p = 0.0078), and IL-17 (p = 0.0156) (n = 8) (**Figure 2A**) ex vivo. For IL-13 statistical significance was not achieved (p = 0.0547) upon treatment of cells with IDB and 75nM SBHA. However, the addition of SBHA to IDB stimulation did not diminish the induction of IFN-γ and IL-8, which are significantly induced by IDB treatment. The effect of SBHA on IDB -induced IL-6 and IL-1β levels appeared inconsistent between donors ex vivo, with donors clustered into two distinct populations. Three donors (H027, H052, and H058) appeared to cluster above the mean pg/mL IL-6 when donor PBMCs were treated with Ingenol-3,30-dibenzoate and SBHA at 75nM or 100nM. In the remaining 5 donors (H037, H055, H063, H065 and H068), IL-6 appeared to decrease to near the limit of detection, though this decrease was not statistically

significant when treated with either 75nM (p =0.0625) or 100nM (p = 0.1250) SBHA. Secreted IL-2, IL-4, IL-10, and IL-12 levels remaining unchanged upon treatment with IDB and the combination therapy. The addition of 100nM panobinostat to 100nM IDB did not significantly reduce secreted levels of any cytokines measured (**Figure 2B**; TNF-α (p = 0.2500), IFN-γ (p = 0.5000), IL-1β (p = 0.2500), IL-2 (p > 0.9999), IL-2r (p = 0.2500), IL-4 (p > 0.9999), IL-5 (p = 0.5000), IL-6 (0.5000), IL-8 (p > 0.9999), IL-10 (0.5000), IL-12 (no difference), IL-13 (p = 0.2500)) (Figure 3B). Secreted IL-17 was not measured.

4. Discussion

In order to eliminate the latent HIV-1 reservoir, LRA therapies must be evaluated based on both their potency and deleterious side effects. PKC agonists are among the most potent LRAs that have emerged across multiple models of HIV-1 latency [29, 30]. However, PKC agonists are also known to induce T cell activation [32] and secretion of pro-inflammatory cytokines [33, 41]. Previous reports have demonstrated that LRAs can be combined with a second pharmacological agent to suppress deleterious pro-inflammatory cytokine secretion induced upon treatment with PKC agonists [33] and TCR stimulation [34]. While HDAC inhibitors such as SAHA [35, 37], Trichostatin A [36], and Nicotinamide [36] have been shown to reduce pro-inflammatory cytokine secretion in other disease contexts [35, 36], we here for the first time have examined the ability of these compounds to do so.

HDAC1-3 inhibition is a known mechanism of inhibiting the production of pro-inflammatory cytokines [42-45]. HDAC1- and HDAC3-independent knock-down have been previously shown to reduced LPS-induced IL-6 and TNF-α in murine microglia [42] and IL-1

pro-inflammatory cytokine induction in HEK293IL-1R and mouse embryonic fibroblasts [45]. Additionally, other HDAC1-3 inhibitors such as entinostat have been shown to reduce the production of pro-inflammatory cytokines IL-1β and IL-6 in a rat collagen-induced arthritis model [43] and inflammatory gene mRNA levels for IL-1β, IL-17, and IFN-γ in an experimental autoimmune neuritis rat model [44]. In this study we showed that Entinostat reduced intracellular IL-6 production but to a lower degree than SBHA, Panobinostat, and Pyroxamide and was therefore not studies further. However, it does strengthen the case that HDAC1-3 inhibition is a candidate pathway for inhibiting pro-inflammatory cytokines induced by PKC stimulation during HIV-1 latency reversal, as we demonstrated by combining known HDAC1-3 inhibitor SBHA with IDB to reduce the production of PKC-induced TNF-α, IL-17, and IL-5 ex vivo. While SBHA reduced PKC-induced pro-inflammatory cytokines, Panobinostat, a pan-HDAC inhibitor, did not significantly reduce pro-inflammatory cytokines. Panobinostat has been previously shown to significantly reduce IL-6 in a Hodgkin's Lymphoma phase II clinical trial [46] and IFN-γ secretion in a lymphocyte-target cell killing assay [47]. Other pan-HDACi such as TSA have been reported to reduce IL-6 [36] and TNF-α [36, 48] secretion in LPS-stimulated macrophages, although other studies have shown that TSA has no effect on other pro-inflammatory genes such as IL-1β and IL-10 [48]. It is unclear why Panobinostat did not reduce PKC-induced pro-inflammatory cytokine production.

The combination therapy of SBHA and IDB significantly reduced PKC-induced pro-inflammatory cytokines suggesting this combination therapy may be an effective strategy to decouple latency reversal and pro-inflammatory cytokine induction. Future experiments will examine the effect of one of our unexplored 'hits', Pyroxamide, on the production of PKC-induced pro-inflammatory cytokines.

5. Conclusions

SBHA, a known HDAC1-3 inhibitor, substantially decreases Ingenol-3,20-dibenzoate-induced TNF-α, IL-2r, IL-5, and IL-17. These results suggest that IDB and SBHA may be a potential LRA combination therapy to reduce PKC-induced pro-inflammatory cytokines during latency reversal.

Acknowledgements

The authors express their sincere gratitude to the study participants for their continued participation in this work and ongoing translational research. This work was supported in part by the National Institute of Allergy and Infectious Diseases (NIAID) grants AI122377-01 and AI143567-02 (VP), NIAID Ruth L. Kirschstein National Research Service Award T32AI055434-11 (ETL), NIAID grant R21AI124823-02 (LRB), and the Doris Duke Charitable Foundation Clinical Scientist Development Award CSDA201612 (AMS). The research reported in this publication was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number UL1TR002538. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing interests

The authors declare no competing interests.

Ethics approval

Heathy donors and HIV-1 positive aviremic individuals on ART were recruited for phlebotomy according to Institutional Review Board (IRB) protocols 67637 (approved May 31, 2017) and 58246 (approved Jan 4, 2017) at the University of Utah.

Figure Legends

Figure 1. Epigenetic modifiers screening identifying compounds that dampen cytokine production induced by Ingenol-3,20-dibenzoate. Screening of 96 epigenetic modifiers for compounds that reduced intracellular IL-6 induced by IDB in healthy donor PBMCs (n=1) revealed nine epigenetic modifier 'hits' that reduce intracellular IL-6 by fourfold or greater (at or below dotted line) when compared to IDB treatment alone. Ruxolitinib was included as a positive control for reduction of intracellular IL-6 when combined with Ingenol-3,20-dibenzoate. Compounds are listed by CAS number and color coded according to categorized function. All nine 'hits' were selected for further testing.

Figure 2A) SBHA reduces pro-inflammatory cytokine secretion induced by IDB ex vivo. Solid lines represent the mean change in pro-inflammatory cytokine concentrations in the supernatant of PBMCs isolated from HIV-1 positive aviremic individuals (n = 8) and treated with SBHA (75nM or 100nM) and IDB (100nM) for 72 hours compared to IDB alone. The dotted line represents the baseline for the assay. *P value <0.05; **P value <0.01. 2**B)** Panobinostat has no significant effect on IDB induced pro-inflammatory cytokine production ex vivo. PBMCs from aviremic individual was treated with Panobinostat (100nM) and IDB (100nM). The mean change in pro-inflammatory cytokines is indicated by a solid line and dotted lines indicate the baseline for the assay.

References

1.	Gulick RM, Mellors JW, Havlir D, Eron JJ, Gonzalez C, McMahon D, et al. Treatment with indinavir, zidovudine, and lamivudine in adults with human immunodeficiency virus infection and prior antiretroviral therapy. N Engl J Med. 1997;337(11):734-9.

2.	Hammer SM, Squires KE, Hughes MD, Grimes JM, Demeter LM, Currier JS, et al. A controlled trial of two nucleoside analogues plus indinavir in persons with human immunodeficiency virus infection and CD4 cell counts of 200 per cubic millimeter or less. AIDS Clinical Trials Group 320 Study Team. N Engl J Med. 1997;337(11):725-33.

3.	Perelson AS, Essunger P, Cao Y, Vesanen M, Hurley A, Saksela K, et al. Decay characteristics of HIV-1-infected compartments during combination therapy. Nature. 1997;387(6629):188-91.

4.	Chun TW, Stuyver L, Mizell SB, Ehler LA, Mican JA, Baseler M, et al. Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy. Proceedings of the National Academy of Sciences. 1997;94(24):13193-7.

5.	Finzi D, Hermankova M, Pierson T, Carruth LM, Buck C, Chaisson RE, et al. Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. Science (New York, NY). 1997;278(5341):1295-300.

6.	Wong JK, Hezareh M, Günthard HF, Havlir DV, Ignacio CC, Spina CA, et al. Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. Science (New York, NY). 1997;278(5341):1291-5.

7.   Torbett BE, Goodsell DS, Richman DD, editors. The Future of HIV-1 Therapeutics-Resistance is Futile. Cham: Springer International Publishing; 2015.

8.   Long L, Fox M, Sanne I, Rosen S. The high cost of second-line antiretroviral therapy for HIV/AIDS in South Africa. AIDS. 2010;24(6):915-9.

9.   Saberi P, Neilands TB, Vittinghoff E, Johnson MO, Chesney M, Cohn SE. Barriers to antiretroviral therapy adherence and plasma HIV RNA suppression among AIDS clinical trials group study participants. AIDS Patient Care STDS. 2015;29(3):111-6.

10.   Rintamaki LS, Davis TC, Skripkauskas S, Bennett CL, Wolf MS. Social stigma concerns and HIV medication adherence. AIDS Patient Care STDS. 2006;20(5):359-68.

11.   Katz IT, Ryu AE, Onuegbu AG, Psaros C, Weiser SD, Bangsberg DR, et al. Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. J Int AIDS Soc. 2013;16(3 Suppl 2):18640.

12.   International ASSWGoHIVC, Deeks SG, Autran B, Berkhout B, Benkirane M, Cairns S, et al. Towards an HIV cure: a global scientific strategy. Nat Rev Immunol. 2012;12(8):607-14.

13.   Borges AH, Dubrow R, Silverberg MJ. Factors contributing to risk for cancer among HIV-infected individuals, and evidence that earlier combination antiretroviral therapy will alter this risk. Curr Opin HIV AIDS. 2014;9(1):34-40.

14.   High KP, Brennan-Ing M, Clifford DB, Cohen MH, Currier J, Deeks SG, et al. HIV and aging: state of knowledge and areas of critical need for research. A report to the NIH Office of AIDS Research by the HIV and Aging Working Group. J Acquir Immune Defic Syndr. 2012;60 Suppl 1:S1-18.

15. Gopal S, Achenbach CJ, Yanik EL, Dittmer DP, Eron JJ, Engels EA. Moving forward in HIV-associated cancer. J Clin Oncol. 2014;32(9):876-80.

16. Elbirt D, Mahlab-Guri K, Bezalel-Rosenberg S, Gill H, Attali M, Asher I. HIV-associated neurocognitive disorders (HAND). Isr Med Assoc J. 2015;17(1):54-9.

17. Deeks SG. HIV: Shock and kill. Nature. 2012;487(7408):439-40.

18. Spivak AM, Planelles V. HIV-1 Eradication: Early Trials (and Tribulations). Trends Mol Med. 2016;22(1):10-27.

19. Burton DR, Hangartner L. Broadly Neutralizing Antibodies to HIV and Their Role in Vaccine Design. Annu Rev Immunol. 2016;34:635-59.

20. Vandergeeten C, Fromentin R, Chomont N. The role of cytokines in the establishment, persistence and eradication of the HIV reservoir. Cytokine Growth Factor Rev. 2012;23(4-5):143-9.

21. Bartholomeeusen K, Xiang Y, Fujinaga K, Peterlin BM. Bromodomain and extra-terminal (BET) bromodomain inhibition activate transcription via transient release of positive transcription elongation factor b (P-TEFb) from 7SK small nuclear ribonucleoprotein. J Biol Chem. 2012;287(43):36609-16.

22. Peng W, Hong Z, Chen X, Gao H, Dai Z, Zhao J, et al. Thiostrepton Reactivates Latent HIV-1 through the p-TEFb and NF-kappaB Pathways Mediated by Heat Shock Response. Antimicrob Agents Chemother. 2020;64(5).

23. Chun TW, Engel D, Mizell SB, Ehler LA, Fauci AS. Induction of HIV-1 replication in latently infected CD4+ T cells using a combination of cytokines. J Exp Med. 1998;188(1):83-91.

24.   Archin NM, Kirchherr JL, Sung JA, Clutton G, Sholtis K, Xu Y, et al. Interval dosing with the HDAC inhibitor vorinostat effectively reverses HIV latency. J Clin Invest. 2017;127(8):3126-35.

25.   Wu G, Swanson M, Talla A, Graham D, Strizki J, Gorman D, et al. HDAC inhibition induces HIV-1 protein and enables immune-based clearance following latency reversal. JCI Insight. 2017;2(16).

26.   Divsalar DN, Simoben CV, Schonhofer C, Richard K, Sippl W, Ntie-Kang F, et al. Novel Histone Deacetylase Inhibitors and HIV-1 Latency-Reversing Agents Identified by Large-Scale Virtual Screening. Front Pharmacol. 2020;11:905.

27.   Laird GM, Bullen CK, Rosenbloom DI, Martin AR, Hill AL, Durand CM, et al. Ex vivo analysis identifies effective HIV-1 latency-reversing drug combinations. J Clin Invest. 2015;125(5):1901-12.

28.   Spivak AM, Bosque A, Balch AH, Smyth D, Martins L, Planelles V. Ex Vivo Bioactivity and HIV-1 Latency Reversal by Ingenol Dibenzoate and Panobinostat in Resting CD4(+) T Cells from Aviremic Patients. Antimicrob Agents Chemother. 2015;59(10):5984-91.

29.   Spina CA, Anderson J, Archin NM, Bosque A, Chan J, Famiglietti M, et al. An in-depth comparison of latent HIV-1 reactivation in multiple cell model systems and resting CD4+ T cells from aviremic patients. PLoS Pathog. 2013;9(12):e1003834.

30.   Darcis G, Kula A, Bouchat S, Fujinaga K, Corazza F, Ait-Ammar A, et al. An In-Depth Comparison of Latency-Reversing Agent Combinations in Various In Vitro and Ex Vivo HIV-1 Latency Models Identified Bryostatin-1+JQ1 and Ingenol-B+JQ1 to Potently Reactivate Viral Gene Expression. PLoS Pathog. 2015;11(7):e1005063.

31.   Genot EM, Parker PJ, Cantrell DA. Analysis of the role of protein kinase C-alpha, -epsilon, and -zeta in T cell activation. J Biol Chem. 1995;270(17):9833-9.

32.   Korin YD, Brooks DG, Brown S, Korotzer A, Zack JA. Effects of prostratin on T-cell activation and human immunodeficiency virus latency. J Virol. 2002;76(16):8118-23.

33.   Spivak AM, Larragoite ET, Coletti ML, Macedo AB, Martins LJ, Bosque A, et al. Janus kinase inhibition suppresses PKC-induced cytokine release without affecting HIV-1 latency reversal ex vivo. Retrovirology. 2016;13(1):88.

34.   Martin AR, Pollack RA, Capoferri A, Ambinder RF, Durand CM, Siliciano RF. Rapamycin-mediated mTOR inhibition uncouples HIV-1 latency reversal from cytokine-associated toxicity. J Clin Invest. 2017;127(2):651-6.

35.   Reddy P, Maeda Y, Hotary K, Liu C, Reznikov LL, Dinarello CA, et al. Histone deacetylase inhibitor suberoylanilide hydroxamic acid reduces acute graft-versus-host disease and preserves graft-versus-leukemia effect. Proc Natl Acad Sci U S A. 2004;101(11):3921-6.

36.   Grabiec AM, Krausz S, de Jager W, Burakowski T, Groot D, Sanders ME, et al. Histone deacetylase inhibitors suppress inflammatory activation of rheumatoid arthritis patient synovial macrophages and tissue. J Immunol. 2010;184(5):2718-28.

37.   Leoni F, Zaliani A, Bertolini G, Porro G, Pagani P, Pozzi P, et al. The antitumor histone deacetylase inhibitor suberoylanilide hydroxamic acid exhibits antiinflammatory properties via suppression of cytokines. Proc Natl Acad Sci U S A. 2002;99(5):2995-3000.

38.   Perez M, de Vinuesa AG, Sanchez-Duffhues G, Marquez N, Bellido ML, Munoz-Fernandez MA, et al. Bryostatin-1 synergizes with histone deacetylase inhibitors to reactivate HIV-1 from latency. Curr HIV Res. 2010;8(6):418-29.

39.   Lu HK, Gray LR, Wightman F, Ellenberg P, Khoury G, Cheng WJ, et al. Ex vivo response to histone deacetylase (HDAC) inhibitors of the HIV long terminal repeat (LTR) derived from HIV-infected patients on antiretroviral therapy. PLoS One. 2014;9(11):e113341.

40.   Evelyn CR, Wade SM, Wang Q, Wu M, Iniguez-Lluhi JA, Merajver SD, et al. CCG-1423: a small-molecule inhibitor of RhoA transcriptional signaling. Mol Cancer Ther. 2007;6(8):2249-60.

41.   Kulkosky J, Culnan DM, Roman J, Dornadula G, Schnell M, Boyd MR, et al. Prostratin: activation of latent HIV-1 expression suggests a potential inductive adjuvant therapy for HAART. Blood. 2001;98(10):3006-15.

42.   Durham BS, Grigg R, Wood IC. Inhibition of histone deacetylase 1 or 2 reduces induced cytokine expression in microglia through a protein synthesis independent mechanism. J Neurochem. 2017;143(2):214-24.

43.   Lin HS, Hu CY, Chan HY, Liew YY, Huang HP, Lepescheux L, et al. Anti-rheumatic activities of histone deacetylase (HDAC) inhibitors in vivo in collagen-induced arthritis in rodents. Br J Pharmacol. 2007;150(7):862-72.

44.   Zhang ZY, Zhang Z, Schluesener HJ. MS-275, an histone deacetylase inhibitor, reduces the inflammatory reaction in rat experimental autoimmune neuritis. Neuroscience. 2010;169(1):370-7.

45.   Ziesche E, Kettner-Buhrow D, Weber A, Wittwer T, Jurida L, Soelch J, et al. The coactivator role of histone deacetylase 3 in IL-1-signaling involves deacetylation of p65 NF-kappaB. Nucleic Acids Res. 2013;41(1):90-109.

46.   Oki Y, Buglio D, Zhang J, Ying Y, Zhou S, Sureda A, et al. Immune regulatory effects of panobinostat in patients with Hodgkin lymphoma through modulation of serum cytokine levels and T-cell PD1 expression. Blood Cancer J. 2014;4:e236.

47.   Klein JM, Henke A, Sauer M, Bessler M, Reiners KS, Engert A, et al. The histone deacetylase inhibitor LBH589 (panobinostat) modulates the crosstalk of lymphocytes with Hodgkin lymphoma cell lines. PLoS One. 2013;8(11):e79502.

48.   Weiss U, Moller M, Husseini SA, Manderscheid C, Hausler J, Geisslinger G, et al. Inhibition of HDAC Enzymes Contributes to Differential Expression of Pro-Inflammatory Proteins in the TLR-4 Signaling Cascade. Int J Mol Sci. 2020;21(23).

Figure 1

Figure 2A

Figure 2B