Impaired competence in flagellar mutants of *Bacillus subtilis* is connected to the regulatory

network governed by DegU

Theresa Hölscher[1],*, Tina Schiklang[1],*, Anna Dragoš[1], Anne-Kathrin Dietel[2,3], Christian Kost[2,3], Ákos T.

Kovács[1,4#]

[1] Terrestrial Biofilms Group, Institute of Microbiology, Friedrich Schiller University Jena, Jena,

Germany

[2] Experimental Ecology and Evolution Group, Department of Bioorganic Chemistry, Max Planck

Institute for Chemical Ecology, Jena, Germany

[3] Department of Ecology, School of Biology/Chemistry, University of Osnabrück, Osnabrück,

Germany

[4] Bacterial Interactions and Evolution Group, Department of Biotechnology and Biomedicine,

Technical University of Denmark, Kgs Lyngby, Denmark

* Contributed equally

[#] **For correspondence. Email** atkovacs@dtu.dk

Running title: Hindered competence by lack of motility

Keywords: flagella, *Bacillus subtilis*, competence, phosphorylation, viscosity

Summary

The competent state is a developmentally distinct phase, in which bacteria are able to take up and

integrate exogenous DNA into their genome. *Bacillus subtilis* is one of the naturally competent

bacterial species and the domesticated laboratory strain 168 is easily transformable. In this study,

we report a reduced transformation frequency of *B. subtilis* mutants lacking functional and

structural flagellar components. This includes *hag*, the gene encoding the flagellin protein forming

the filament of the flagellum. We confirm that the observed decrease of the transformation

frequency is due to reduced expression of competence genes, particularly of the main competence

regulator *comK*. The impaired competence is due to an increase in the phosphorylated form of the

response regulator DegU, which is involved in regulation of both flagellar motility and competence.

Altogether, our study identified a close link between motility and natural competence in *B. subtilis*

suggesting that hindrance in motility has great impact on differentiation of this bacterium not

restricted only to the transition towards sessile growth stage.

Originality-Significance statement

Understanding how versatile bacterial phenotypes influence each other is important for our basic

understanding of microbial ecology. Our research highlights the novel intertwinement of bacterial

differentiation and reveal how lack of single cell motility adjusts DNA exchange among bacterial

strains.

Introduction

When facing stressful environmental conditions, bacteria can respond with a variety of post-exponential modifications including secretion of degradative enzymes, sporulation, or genetic competence. *Bacillus subtilis* is one of the bacterial species that are able to take up free DNA from the environment and incorporate it into its own genome, a phenomenon referred to as natural competence (Dubnau, 1991). To import extracellular DNA into *B. subtilis* cells, a pseudopilus formed by proteins encoded by the *comG* operon facilitates binding to the receptor protein ComEA, which is located in the bacterial cell membrane (Inamine and Dubnau, 1995; Chen *et al.*, 2005). As only single stranded DNA is imported, the membrane-associated nuclease NucA catalyzes cleavage of the DNA after successful binding (Provvedi *et al.*, 2001). Subsequent transport of the DNA through a membrane channel formed by the protein ComEC is mediated by the ATPase ComFA that probably requires the transmembrane proton motive force (Maier *et al.*, 2004).

To take up DNA, cells have to be in a developmental state, in which a specific set of genes and regulators are expressed (Dubnau, 1991; Berka *et al.*, 2002). Regulation of the whole apparatus required for competence development is complex. Briefly, entry into the competence state occurs in a bistable manner during the early stationary phase, where a minority of cells produces high level of the competence master regulator ComK above a certain threshold that is required to switch on competence development, the so called 'K-state' (van Sinderen *et al.*, 1995; Maamar and Dubnau, 2005; Smits *et al.*, 2005; Dubnau and Losick, 2006). It was demonstrated that noise in the expression of *comK* determines the competent subpopulation and allows a dynamic stress response regarding competence development (Maamar *et al.*, 2007; Mugler *et al.*, 2016). Eventually, ComK activates the expression of late competence operons encoding the DNA-binding and -uptake machinery as well as genes, whose products are responsible for DNA integration (Berka *et al.*, 2002; Ogura *et al.*, 2002; Hamoen *et al.*, 2003). Increase of the ComK level is linked to a quorum sensing-mediated accumulation of the small ComS protein, which interferes with ComK degradation during the exponential phase (Turgay *et al.*, 1998). ComK is able to bind the *comK* promoter, triggering its own transcription, thus creating an auto-stimulatory loop (van Sinderen and Venema, 1994). This binding is further stabilized by the non-phosphorylated form of the regulator DegU to increase the level of ComK above the threshold sufficient for competence development (Hamoen *et al.*, 2000).

However, DegU is not only crucial for competence initiation, but also involved in the regulation of many other processes including protease production, biofilm development, and, particularly, flagellar motility (Murray *et al.*, 2009; Mukherjee and Kearns, 2014). The main components of the hook and basal body of the flagellum are encoded by the large *fla/che* operon. Transcription of this operon is activated by a complex formed by the regulator SwrA and phosphorylated DegU (DegU~P),

which binds to one of the *fla/che* promoters (Mordini *et al.*, 2013; Mukherjee and Kearns, 2014).

Amongst other genes, the operon contains the gene encoding the sigma factor σ^D that activates

transcription of motility genes outside the *fla/che* operon like the *hag* gene (encoding flagellin),

motA and *motB* (encoding flagellar stator proteins), as well as transcription of *lytF*, which is

necessary for separation of motile cells after cell division (Serizawa *et al.*, 2004; Chen *et al.*, 2009).

The level of σ^D and its position in the *fla/che* operon determines the cell fate, i.e. subpopulations of

motile single cells or non-motile chains (Cozy and Kearns, 2010). The function of DegU~P changes in

the absence of SwrA. In this case, DegU~P seems to inhibit motility via the same promoter of the

fla/che operon (Amati *et al.*, 2004). Additionally, DegU~P can activate the anti-sigma factor FlgM by

binding to its promoter region in the absence of SwrA (Hsueh *et al.*, 2011) allowing FlgM to

antagonize σ^D (Caramori *et al.*, 1996). Consequently, DegU~P indirectly suppresses transcription of

σ^D-dependent genes (Hsueh *et al.*, 2011). It was suggested that a completion of flagellum assembly

can be sensed by the DegSU two component system: FlgM, which is activated by DegU~P, causes

inhibition of σ^D-dependent genes, when the assembly of the flagellum is impeded (Cozy and Kearns,

2010; Hsueh *et al.*, 2011).

In addition to its role on modulating the expression of flagellum-related genes in *B. subtilis*, the

phosphorylation and therefore the activity of DegU, has been shown to be influenced by a

mechanical signal transmitted by the flagellum (Cairns *et al.*, 2013). Inhibition of flagellar rotation by

the flagellar clutch or by tethering the flagella results in an increased DegU~P level in the cell.

In this study, we report a correlation between motility function and competence development,

which in *B. subtilis* is connected by the multifunctional response regulator DegU. We show that

mutants lacking a functional flagellum such as Δ*hag*, Δ*motA*, and Δ*flgE* exhibited a reduced

transformation frequency. This was due to a decrease in competence gene expression, particularly

reduced levels of the competence master regulator ComK, which can be reverted by overexpressing

comK in the *hag* mutant. Finally, we suggest that the reduced transformation frequency was likely

due to an imbalance in the phosphorylation level of DegU.

Results

Lack of active flagella impairs competence for DNA uptake in B. subtilis.

While genetically modifying various *B. subtilis* strains, a striking difference in transformation frequency was observed between the wild type and a non-motile mutant lacking the gene encoding flagellin, *hag*. To explore this phenomenon, we tested the transformability of wild type (strain 168) and *hag*-mutant in competence medium (see Experimental Procedures), where the *hag*-mutant showed a more than 100-fold reduced transformation frequency relative to the wild type (Fig. 1A, B): while the transformation frequency of the wild type ranged between $3\cdot10^{-5}$ and $5\cdot10^{-5}$, that of the hag-mutant was reduced to values below $3\cdot10^{-7}$. Similarly, the undomesticated *B. subtilis* strains DK1042 (transformable derivative of NCIB 3610) and PS216 showed reduced transformation efficiency when the *hag* gene of these strains was disrupted (Fig. S1). To investigate whether this difference in transformation frequency between the two strains resulted from a lower growth rate of the *hag*-mutant, the growth behavior of wild type and *hag*-mutant grown in competence medium was evaluated over time. As depicted in Fig. 1B, the *hag*-mutant showed a clear growth advantage and reached a higher OD compared to the wild type (unpaired two-sample t-test with Welch Correction: P = 0.001, n = 5), thus supporting our previous observations (Hölscher *et al.*, 2015). Further, it was tested whether the addition of DNA at different time points would increase the transformation frequency of the *hag*-mutant. However, the mutant showed a consistently low transformation frequency over the course of several hours, indicating that a shifted timing of the initiation of the competence state is unlikely to be the reason for the observed decrease in transformation frequency (Fig. 1C). To test whether this phenomenon is restricted to the *hag*-mutant or connected to the lack of an active motility apparatus in general, mutants lacking other functional flagellum-related genes were investigated. The transformation frequencies of mutants lacking the gene encoding one of the flagellar motor units, *motA*, and the gene encoding the hook protein, *flgE*, were decreased in both cases compared to the wild type (Fig. 2; unpaired two-sample t-test with Welch Correction: P = 0.01 for WT - Δ*motA*, P = 0.039 for WT - Δ*flgE*, n = 9 for both). Although the wild type transformation frequency was slightly different, the transformation frequencies of both Δ*motA* and Δ*flgE* were around 10-times lower than that of the wild type (Fig. 2). In contrast, a *cheA*-mutant lacking the main chemotaxis sensor kinase showed a similar transformation frequency than the wild type (Fig. S2; unpaired two-sample t-test with Welch Correction: P = 0.232, n = 3), suggesting that the presence of an active flagellum, but not directed motility *per se* is required for full competence development. In sum, these results demonstrate that the observed impaired competence is linked to a loss of flagellar function.

Lack of competence in flagellar mutants is due to the reduced expression of competence genes.

To determine if the detected diminished transformation frequency of flagellar mutants was due to

altered competence gene expression, the fluorescent reporter P*comG*-*gfp* was introduced into these

strains. This reporter allows the detection of cells expressing the *comG* operon-encoding genes

required for pseudopilus formation and DNA uptake. In addition, this reporter provides a proxy on

the activity of the ComK protein, the master regulator of competence. Qualitative microscopy

analyses of cultures harboring the reporter, and which were grown in competence medium for 5 h,

showed indeed a decreased number of fluorescent (i.e. *comG* expressing) cells in the *hag* mutant

compared to the wild type, whereas a control strain lacking *comK* showed no fluorescence (Fig. 3A).

For quantitative determination of competence gene expression within the population, flow

cytometric measurements were performed that revealed 24.7% of fluorescent cells in wild type

cultures (mean value), but only 4.6% of fluorescent cells for the *hag*-mutant (Fig. 3B, C; unpaired

two-sample t-test with Welch Correction: P = 0.004, n = 3), thus confirming the microscopy results.

Similarly, the *motA* and *flgE* mutants were analyzed microscopically as well as by using flow

cytometry. Both methods revealed fewer cells activated transcription of competence genes in these

mutants compared to the wild type (Fig. 4; unpaired two-sample t-test with Welch Correction: P =

0.017 for WT - Δ*motA*, P = $1.3 \cdot 10^{-9}$ for WT - Δ*flgE*, n = 3 for both; mean percentage of fluorescent

cells: 16.7% for wild type, 4.5% for Δ*motA*, 4.2% for Δ*flgE*). Flow cytometry measurements at

different time points during growth in competence medium confirmed a similarly reduced fraction of

competent cells in the *hag* mutant compared to the wild type strain (Fig. S3).

Reduced competence in hag *mutant can be rescued by overexpression of* comK.

The reduced competence gene expression in the tested flagellar mutants suggested a regulatory link

between flagellar motility and competence. To investigate if regulatory elements upstream of *comK*

were responsible for our observations and if a bypass of those could therefore rescue

transformation frequency in the flagellar mutant, we examined a strain with an additional copy of

comK under the control of a xylose-inducible promoter (P*xyl*-*comK*). Indeed, in combination with P*xyl*-

comK, the transformation level of the *hag* mutant increased back to a level that was statistically

indistinguishable from wild type levels (mean transformation frequency of $5.3 \cdot 10^{-6}$ for the wild type

and $8.5 \cdot 10^{-6}$ for Δ*hag* P*xyl*-*comK*; Kruskal-Wallis test: P = 0.453, n = 9, Fig. 5A). Despite this observed

increase in the *hag* strain upon *comK* overexpression, the wild type strain, which contained an

inducible copy of *comK* showed a higher transformation frequency (Fig. 5A, Kruskal-Wallis test: P =

$3.4 \cdot 10^{-4}$ for WT – WT P*xyl*-*comK*, P = $3.4 \cdot 10^{-4}$ for WT P*xyl*-*comK* - Δ*hag* P*xyl*-*comK*, n = 9 for both), which

was probably due to higher levels of *comK* transcription at the native locus as previously observed
(Hahn *et al.*, 1996).
171
Reduced competence in flagellar mutants is likely connected to unbalanced DegU phosphorylation.
As the above results suggested that regulatory elements in response to impaired flagellar motility
are responsible for the decreased *comK* expression, we investigated DegU as a likely candidate
causing the reduced competence in flagellar mutants. As non-phosphorylated DegU was implicated
to be required for *comK* transcription (Dahl *et al.*, 1992; Hamoen *et al.*, 2000), two variants of *degU*
were tested: *degU*32, which harbors a mutation resulting in an extended half-life and thus higher
stability of the phosphorylated form of the DegU protein (DegU~P), and *degU*146, which is cannot be
phosphorylated (Dahl *et al.*, 1991; Dahl *et al.*, 1992; Kunst *et al.*, 1994). Both variants were tested in
wild type as well as the Δ*hag* background to observe differences in transformability compared to the
wild type strain. The results of this experiment indicated that the transformation frequency of the
*degU*32 strain was slightly decreased (Figure 5B), which is consistent with previous publications,
suggesting that non-phosphorylated DegU is required for priming *comK* transcription. The observed
difference, however, was only marginally significant in our experimental setup (Figure 5B; Kruskal-
Wallis test: $P = 0.078$, $n = 6$). Surprisingly, when combined with the Δ*hag* mutation, the
transformability of *degU*32 increased significantly far above wild type levels, despite presumably
possessing low levels of non-phosphorylated DegU to induce the ComK auto-stimulatory loop (Fig.
5B; Kruskal-Wallis test: $P = 0.004$, $n = 6$). Furthermore, we observed a tendency towards a reduced
albeit non-significant transformation frequency in the *degU*146 strain compared to the wild type
(Fig. 5B; Kruskal-Wallis test: $P>0.05$, $n = 6$). This result was similar to the one observed for *degU*32,
although no negative impact on transformability was expected in strain *degU*146 due to the
abolished phosphorylation of DegU. Interestingly, the *degU*146 strain combined with the Δ*hag*
mutation exhibited transformation frequencies at the same level than the wild type strain (Fig. 5B;
Kruskal-Wallis test: $P>0.05$, $n = 6$) that was significantly higher than the transformation frequency of
the single *degU*146 mutant (Fig. 5B; Kruskal-Wallis test, $P = 0.007$, $n = 6$). These results suggest that
altering the phosphorylation state of DegU in flagellar mutants can revert the negative impact on
competence, which was caused by a lack of motility.
198
Increased viscosity enhances competence in B. subtilis
A recent study showed that restricting the flagellar rotation by viscous medium results in induction
of flagellar gene transcription and activation of the DegSU two-component system in *Paenibacillus*
sp. NAIST15-1 (Kobayashi *et al.*, 2017). Accordingly, we tested whether an increased viscosity of the

medium changes the transformability in *B. subtilis*. Indeed, the average transformation frequency of

the wild type strain was three-fold higher in a medium of increased viscosity. The corresponding

statistical test, however, indicated only a trend towards a statistically significant difference (Fig. 6;

unpaired two-sample t-test: $P = 0.095$, $n = 4$).

Discussion

Many cellular processes in *B. subtilis* are tightly connected through their underlying regulatory

networks. Examples include motility and biofilm formation or biofilm formation and sporulation

(e.g. Vlamakis *et al.*, 2013; Marlow *et al.*, 2014; Hölscher *et al.*, 2015). Here, we report an additional

connection between flagellar motility and competence development. We could show that mutants

with impaired flagellar function have defects in competence development. Such mutants displayed a

considerably lower transformation frequency and expression of late competence genes, suggesting

that it is due to an altered expression of the competence master regulator gene *comK*. The rescue

experiment with an inducible *comK* confirmed that indeed competence could be rescued in the Δ*hag*

strain, since Δ*hag* P$_{xyl}$-*comK* exhibited a wild type transformation level.

In a recently published study, similar effects were observed even though different methods have

been used: Diethmaier and colleagues found that the expression of *comK* is lower in deletion

mutants of the *fla/che* operon, *hag*, and the second stator gene *motB* (Diethmaier *et al.*, 2017).

While in their study the expression of *comK* was primarily monitored using a *comK* promoter fusion,

our experiments predominantly assayed transformation frequency. Both studies, however, report a

negative effect of the deletion of flagellar components on competence development. The extent to

which wild type and mutant differ in competence development is in the same order of magnitude

between the studies: for example Diethmaier *et al*. observe a 10-fold reduced number of *comG*

expressing-cells in Δ*hag* (Diethmaier *et al.*, 2017), whereas our flow cytometry experiments showed

a slightly lower, 5-fold reduction. Additionally, by investigating the transformation frequency in a

cheA mutant, we could also show that the chemotactic response does not seem to have an influence

on competence development.

Investigating modified variants of the response regulator DegU, we found that the transformation

frequency of the *hag*-mutant could be restored to wild type level when the mutant carried a non-

phosphorylatable DegU variant (*degU*146). This result suggests that a high level of DegU~P in the

flagellar mutants was the reason for the decreased expression of the competence genes and *comK*,

which could be counteracted by introducing a non-phosphorylatable variant of DegU. By additionally

investigating a strain harboring a *degU-yfp* fusion, Diethmaier *et al*. also suggested an increased level

of DegU~P to be present in the *hag*-mutant (Diethmaier *et al.*, 2017), which is consistent with our

conclusions. In addition, the authors detected a reduced expression of *comK* in a strain with the

*degU*32 variant, which produces a form of DegU~P with higher stability (Diethmaier *et al.*, 2017).

Comparable results were obtained by Msadek *et al*., who found that high levels of DegU~P inhibit

competence (Msadek *et al.*, 1990). We observed a similar, although weak statistical trend towards a

reduced transformation frequency in *degU*32 strain. Miras and Dubnau (2016) have recently

highlighted that differences in the DegU phosphorylation pathway among diverse *B. subtilis* isolates
were likely responsible for variance in DNA transformation efficiency among certain domesticated
and undomesticated strains. Moreover, slight differences in competence induction levels could also
be affected by strain-specific characteristics. For example, *B. subtilis* 168 strains derived from
different laboratories can exhibit striking variations in biofilm robustness (Gallegos-Monterrosa *et
al.*, 2016). As suggested by Diethmaier and colleagues, the reduced transformation frequency in
*degU*32 might be caused by the high DegU~P levels of this strain. However, the *degU*32 strain
exhibits a non-motile phenotype and in the undomesticated strains DegU32 is not able to interact
with SwrA at the P_A promoter of the *fla/che* operon (*swrA* is inactive in domesticated strains),
leading to repression of P_A (*fla/che*) (Amati *et al.*, 2004; Mordini *et al.*, 2013). Due to low or no
expression of the basic flagellar genes, this phenotype could mimic the situation observed in the
flagellar mutants. In addition, we observed an increased transformation frequency when the *hag*
gene was deleted from the *degU*32 background. This is in contrast to the model assuming that
increased levels of phosphorylated DegU in the cells lowers competence. Therefore, it is possible
that yet unidentified factors are also involved in connecting motility and competence development
that might be independent of DegU~P. At this point however, we cannot provide a reasonable
explanation for the increased transformation frequency of Δ*hag degU*32.

Interestingly, induction of competence state has negative impact on motility in *B. subtilis*. ComK
negatively controls *hag* gene expression by stimulating the transcription of *comFA-C* operon and the
downstream located anti-sigmaD factor coding gene, *flgM* (Liu and Zuber, 1998). This feedback loop
presents another intriguing connection between these two cellular processes.

Diethmaier *et al*. proposed that increased DegU~P and lower *comK* expression in the flagellar
mutants and in a strain with straight flagella was caused by a lower viscous load (Diethmaier *et al.*,
2017). In line with this report, we also observed that higher viscosity in the medium resulted in an
increased transformation frequency. Nevertheless, a possible role of the DegSU two-component
system in sensing incomplete assembly of flagella and dysfunction as suggested previously (Hsueh *et
al.*, 2011; Cairns *et al.*, 2013) could also explain the increased DegU~P levels in the flagellar mutants.

Together, our results identify a connection between two major physiological processes, providing
another example of the complexity of intracellular regulatory networks and the vast amount of tasks
a single regulator can cover.

Experimental Procedures

Strains and cultivating conditions

The strains used in this study and their mutant derivatives are listed in Table S1. Mutants

constructed in this study were obtained by natural transformation of a *B. subtilis* receptor strain with

genomic DNA from a donor strain. Strain TB831 was created by transformation of strain 168 P$_{xyl}$-

comK with genomic DNA of strain GP902 (J. Stülke lab collection). To obtain strains TB926 and

TB925, genomic DNA of strain 168 P$_{comG}$-*gfp* was used to transform strain TB710 and TB689,

respectively. Strain TB928 was obtained by transforming strain 168 P$_{xyl}$-*comK* with genomic DNA of

GP901 (J. Stülke lab collection). To create strain TB935 and TB936, strain 168 was transformed with

genomic DNA obtained from strain QB4371 (Kunst *et al.*, 1994) and QB4458 (Dahl *et al.*, 1991),

respectively. Their derivatives harbouring also a mutation of *hag* (TB923 and TB924) were created by

transformation with genomic DNA, which was obtained from GP901. In-frame deletions of *motA*,

flgE, and *cheA* were created using plasmids pEC1, pDP306, and pDP338, respectively, as previously

described (Courtney *et al.*, 2012; Chan *et al.*, 2014; Calvo and Kearns, 2015). Strains were verified by

fluorescence microscopy (P$_{comG}$-*gfp* reporter), PCR (*hag* mutants), or sequencing (*degU* variants),

using the oligonucleotides listed in Table S2. For experiments with strains harboring the inducible

construct P$_{xyl}$-*comK*, 1% of xylose (final concentration) was added for induction (see van den Esker *et*

al., 2017). To increase medium viscosity, 10% Ficoll400 (Carl Roth) was added to the medium before

culture inoculation and the mix was vortexed vigorously for ca 20 s.

Transformation frequency assay

To assess the transformation frequency of different strains, a modified version of the transformation

protocol from Konkol *et al*. (2013) was used. 1 ml of each culture grown in 3 ml Lysogeny broth (LB)

medium (LB-Lennox, Carl Roth; 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, and 5 g L^{-1} NaCl) for 16 h was

centrifuged for 2 min at 11,000 x g. The pellet was washed twice in de-ionized water and was re-

suspended in 100 µl de-ionized water. The re-suspended culture was diluted (1:80) in complete

competence medium (MC: 1,8 ml de-ionized water, 6.7 µl 1M MgSO$_4$, 50 µl 0.2% L-Tryptophan, 200

µl 10xMC; per 100 ml 10xMC: 14,036 g K$_2$HPO$_4$ [x3H$_2$O], 5,239 g KH$_2$PO$_4$, 20 g glucose, 10 ml 300 mM

tri-sodium citrate, 1 ml 83.97 mM ammonium iron (III) citrate, 1 g casein hydrolysate, 2 g potassium

glutamate [H$_2$O]) and incubated at 37°C, 225 rpm. For experiments with strains harboring P$_{xyl}$-*comK*,

10x MC with fructose instead of glucose was used. After 6 h incubation time, 5 µl DNA with an

antibiotic marker (PY79 *safA*::Tet gDNA, 60 ng/µl) was added to 500 µl culture. Any alteration in

incubation time before addition of the DNA is indicated in the results section. Each culture was

incubated for 30 min, then 500 µl fresh LB medium was added and the culture was incubated for

another 1 h under the conditions mentioned above. Serial dilutions of cultures supplemented with

DNA were prepared and plated on LB medium supplemented with 1.5 % agar to determine the

number of colony forming units (cfu). Additionally, 50 μl and 100 μl undiluted cultures

supplemented with gDNA as well as controls were plated on tetracycline (Tet) containing LB-agar

plates (10 μg ml^{-1} Tet) to determine the number of transformant colonies. The transformation

frequency was calculated by dividing the number of transformants per ml by cfu per ml.

Growth curve experiments

To examine growth properties, cultures were inoculated in LB medium from frozen glycerol stocks

and incubated for ca. 16 h at 37°C shaking at 225 rpm. Cultures were diluted 1:100 in 200 μl fresh

completed MC medium (see above) and the OD_{590nm} was recorded for 16 h using a TECAN Infinite

F200 PRO microplate reader. The cultures were incubated with orbital shaking with a duration of

800 s and an amplitude of 3 mm at 37°C and the OD_{590} was measured every 15 min.

Fluorescence microscopy

Strains were investigated using a confocal laser scanning microscope (LSM 780, Carl Zeiss) equipped

with an argon laser and a Plan-Apochromat/1.4 Oil DIC M27 63× objective. Cultures were grown

prior microscopy for 5 h (if not indicated otherwise) in competence medium under the same

conditions as described above (see section transformation frequency assay). Excitation of the

fluorescent reporter (GFP) was performed at 488nm and the emitted fluorescence was recorded at

493-598nm. For image visualization, Zen 2012 software (Carl Zeiss) was used, brightness and

contrast were adjusted equally in all images.

Flow cytometry

Flow cytometric measurements were performed using a Partec CyFlow® Space (Sysmex Partec

GmbH, Germany), which was equipped with a solid-state laser for excitation of green/yellow

fluorescent proteins at 488 nm. Single cells were detected in forward and sideward scatter channels

as well as in one fluorescent channel. A minimum of 40,000 cells were analyzed for the experiments.

To define the background fluorescence signal, non-labelled *B. subtilis* cultures were analyzed as

control. Cultures used for measurements were grown for 5 h (if not indicated otherwise) in

competence medium under the same conditions as described above (see section transformation

frequency assay). For evaluation of the data, the FlowJo® software (FlowJo LLC, Ashland, USA) was

used and a gate was set at 3 fluorescence units for all samples to isolate the fluorescent population

and determine the percentage of fluorescent cells.

Statistics

Statistical analyses were performed using OriginPro 2016 (V93E, OriginLab Northampton, USA).

Unpaired two-sample t-test with Welch Correction or a Kruskal-Wallis test was used to test for

significant differences.

Acknowledgement

We thank Daniel Kearns for providing plasmids, Anne Richter for constructing the in-frame deletion mutants, and Tadeja Marin for initial observations. We thank Wiep Klaas Smits for his comment on our bioRxiv submission. This project was funded by grant KO4741/3-1 from the German Research Foundation (DFG). T.H. and A.D. were supported by International Max Planck Research School and Alexander von Humboldt foundation fellowships, respectively. CK was supported by the Volkswagen Foundation (I/85 290) as well as the German Research Foundation (SFB 944/2-2016, KO 3909/2-1). Á.T.K. was supported by a Startup fund from the Technical University of Denmark.

Conflict of interest. None declared

References

Amati, G., Bisicchia, P., and Galizzi, A. (2004) DegU-P represses expression of the motility *fla-che*

operon in *Bacillus subtilis*. *J Bacteriol* **186**: 6003–6014.

Berka, R.M., Hahn, J., Albano, M., Draskovic, I., Persuh, M., Cui, X., *et al.* (2002) Microarray analysis

of the *Bacillus subtilis* K-state: Genome-wide expression changes dependent on ComK. *Mol Microbiol*

43: 1331–1345.

Cairns, L.S., Marlow, V.L., Bissett, E., Ostrowski, A., and Stanley-Wall, N.R. (2013) A mechanical signal

transmitted by the flagellum controls signalling in *Bacillus subtilis*. *Mol Microbiol* **90**: 6–21.

Calvo, R.A., and Kearns, D.B. (2015) FlgM is secreted by the flagellar export apparatus in *Bacillus*

subtilis. *J Bacteriol* **197**: 81–91.

Caramori, T., Barilla, D., Nessi, C., Sacchi, L., and Galizzi, A. (1996) Role of FlgM in sigma(D)-

dependent gene expression in *Bacillus subtilis*. *J Bacteriol* **178**: 3113–3118.

Chan, J.M., Guttenplan, S.B., and Kearns, D.B. (2014) Defects in the flagellar motor increase synthesis

of poly-γ-glutamate in *Bacillus subtilis*. *J Bacteriol* **196**: 740–753.

Chen, I., Christie, P.J., and Dubnau, D. (2005) The ins and outs of DNA transfer in bacteria. *Science*

310: 1456–1460.

Chen, R., Guttenplan, S.B., Blair, K.M., and Kearns, D.B. (2009) Role of the σD-dependent autolysins

in *Bacillus subtilis* population heterogeneity. *J Bacteriol* **191**: 5775–5784.

Courtney, C.R., Cozy, L.M., and Kearns, D.B. (2012) Molecular characterization of the flagellar hook in

Bacillus subtilis. *J Bacteriol* **194**: 4619–4629.

Cozy, L.M., and Kearns, D.B. (2010) Gene position in a long operon governs motility development in

Bacillus subtilis. *Mol Microbiol* **76**: 273–285.

Dahl, M.K., Msadek, T., Kunst, F., and Rapoport, G. (1991) Mutational analysis of the *Bacillus subtilis*

DegU regulator and its phosphorylation by the DegS protein kinase. *J Bacteriol* **173**: 2539–2547.

Dahl, M.K., Msadek, T., Kunst, F., and Rapoport, G. (1992) The phosphorylation state of the DegU

response regulator acts as a molecular switch allowing either degradative enzyme synthesis or

expression of genetic competence in *Bacillus subtilis*. *J Biol Chem* **267**: 14509–14514.

Diethmaier, C., Chawla, R., Canzoneri, A., Kearns, D.B., and Dubnau, D. (2017) Viscous drag on the

flagellum activates *Bacillus subtilis* entry into the. *Mol Microbiol* **106:** 367-380.

Dubnau, D. (1991) Genetic competence in *Bacillus subtilis*. *Microbiol Rev* **55**: 395–424.

Dubnau, D., and Losick, R. (2006) Bistability in bacteria. *Mol Microbiol* **61**: 564–572.

van den Esker, M.H., Kovács, Á.T., and Kuipers, O.P. (2017) YsbA and LytST are essential for pyruvate

utilization in *Bacillus subtilis*. *Environ Microbiol* **19**: 83–94.

Gallegos-Monterrosa, R., Mhatre, E., and Kovács, Á.T. (2016) Specific *Bacillus subtilis* 168 variants

form biofilms on nutrient-rich medium. *Microbiol* **162**: 1922–1932.

Hamoen, L.W., Venema, G., and Kuipers, O.P. (2003) Controlling competence in *Bacillus subtilis*:

shared use of regulators. *Microbiol* **149**: 9–17.

Hamoen, L.W., Werkhoven, a F. Van, Venema, G., and Dubnau, D. (2000) The pleiotropic response

regulator DegU functions as a priming protein in competence development in *Bacillus subtilis. Proc*

Natl Acad Sci U S A **97**: 9246–9251.

Hölscher, T., Bartels, B., Lin, Y.-C., Gallegos-Monterrosa, R., Price-Whelan, A., Kolter, R., *et al.* (2015)

Motility, chemotaxis and aerotaxis contribute to competitiveness during bacterial pellicle biofilm

development. *J Mol Biol* **427**: 3695–3708.

Hsueh, Y.H., Cozy, L.M., Sham, L.T., Calvo, R.A., Gutu, A.D., Winkler, M.E., and Kearns, D.B. (2011)

DegU-phosphate activates expression of the anti-sigma factor FlgM in *Bacillus subtilis*. *Mol Microbiol*

81: 1092–1108.

Inamine, G.S., and Dubnau, D. (1995) ComEA, a *Bacillus subtilis* integral membrane protein required

for genetic transformation, is needed for both DNA binding and transport. *J Bacteriol* **177**: 3045–

3051.

Kobayashi, K., Kanesaki, Y., and Yoshikawa, H. (2017) Surface sensing for flagellar gene expression on

solid media in *Paenibacillus sp*. NAIST15-1. *Appl Environ Microbiol* **83:** e00585-17.

Konkol, M.A., Blair, K.M., and Kearns, D.B. (2013) Plasmid-encoded comi inhibits competence in the

ancestral 3610 strain of *Bacillus subtilis*. *J Bacteriol* **195**: 4085–4093.

Kunst, F., Msadek, T., Bignon, J., and Rapoport, G. (1994) The DegS/DegU and ComP/ComA two-

component systems are part of a network controlling degradative enzyme synthesis and

competence in *Bacillus subtilis*. *Res Microbiol* **145**: 393–402.

Liu, J., and Zuber, P. (1998) A molecular switch controlling competence and motility: competence

regulatory factors ComS, MecA, and ComK control sigmaD-dependent gene expression in *Bacillus*

subtilis. *J Bacteriol* **180:** 4243-4251.

Maamar, H., and Dubnau, D. (2005) Bistability in the *Bacillus subtilis* K-state (competence) system

requires a positive feedback loop. *Mol Microbiol* **56**: 615–24.

Maamar, H., Raj, A., and Dubnau, D. (2007) Noise in Gene Expression Determines. *Science* 526–529.

Maier, B., Chen, I., Dubnau, D., and Sheetz, M.P. (2004) DNA transport into *Bacillus subtilis* requires

proton motive force to generate large molecular forces. *Nat Struct Mol Biol* **11**: 643–649.

Marlow, V.L., Porter, M., Hobley, L., Kiley, T.B., Swedlow, J.R., Davidson, F. a, and Stanley-Wall, N.R.

(2014) Phosphorylated DegU manipulates cell fate differentiation in the *Bacillus subtilis* biofilm. *J*

Bacteriol **196**: 16–27.

Miras, M., and Dubnau, D. (2016) A DegU-P and DegQ-Dependent Regulatory Pathway for the K-

state in *Bacillus subtilis*. *Front Microbiol* **7**: 1–14.

Mordini, S., Osera, C., Marini, S., Scavone, F., Bellazzi, R., Galizzi, A., and Calvio, C. (2013) The Role of

SwrA, DegU and PD3 in *fla/che* Expression in *B. subtilis*. *PLoS One* **8**: e85065.

Msadek, T., Kunst, F., Henner, D., Klier, A., Rapoport, G., and Dedonder, R. (1990) Signal transduction

pathway controlling synthesis of a class of degradative enzymes in *Bacillus subtilis*: expression of the

regulatory genes and analysis of mutations in *degS* and *degU*. *J Bacteriol* **172**: 824–834.

Mugler, A., Kittisopikul, M., Hayden, L., Liu, J., Wiggins, C.H., Süel, G.M., and Walczak, A.M. (2016)

Noise expands the response range of the *Bacillus subtilis* competence circuit. *PLoS Comput Biol* **12**:

1–21.

Mukherjee, S., and Kearns, D.B. (2014) The structure and regulation of flagella in *Bacillus subtilis*.

Annu Rev Genet **48**: 319–40.

Murray, E.J., Kiley, T.B., and Stanley-Wall, N.R. (2009) A pivotal role for the response regulator DegU

in controlling multicellular behaviour. *Microbiol* **155**: 1–8.

Ogura, M., Yamaguchi, H., Kobayashi, K., Ogasawara, N., and Fujita, Y. (2002) Whole-genome analysis

of genes regulated by the *Bacillus subtilis* competence transcription factor ComK. *J Appl Microbiol*

184: 2344–2351.

Provvedi, R., Chen, I., and Dubnau, D. (2001) NucA is required for DNA cleavage during

transformation of *Bacillus subtilis*. *Mol Microbiol* **40**: 634–644.

Serizawa, M., Yamamoto, H., Yamaguchi, H., Fujita, Y., Kobayashi, K., Ogasawara, N., and Sekiguchi, J.

(2004) Systematic analysis of SigD-regulated genes in *Bacillus subtilis* by DNA microarray and

Northern blotting analyses. *Gene* **329**: 125–136.

van Sinderen, D., Luttinger, A., Kong, L., Dubnau, D., Venema, G., and Hamoen, L. (1995) *comK*

encodes the competence transcription factor, the key regulatory protein for competence

development in *Bacillus subtilis*. *Mol Microbiol* **15**: 455–462.

van Sinderen, D., and Venema, G. (1994) ComK acts as an autoregulatory control switch in the signal

transduction route to competence in *Bacillus subtilis*. *J Bacteriol* **176**: 5762–5770.

Smits, W.K., Eschevins, C.C., Susanna, K.A., Bron, S., Kuipers, O.P., and Hamoen, L.W. (2005) Stripping

Bacillus: ComK auto-stimulation is responsible for the bistable response in competence

development. *Mol Microbiol* **56**: 604–614.

Turgay, K., Hahn, J., Burghoorn, J., and Dubnau, D. (1998) Competence in *Bacillus subtilis* is

controlled by regulated proteolysis of a transcription factor. *EMBO J* **17**: 6730–6738.

Vlamakis, H., Chai, Y., Beauregard, P., Losick, R., and Kolter, R. (2013) Sticking together: building a

biofilm the *Bacillus subtilis* way. *Nat Rev Microbiol* **11**: 157–68.

Figures

Fig. 1. Transformation frequency is reduced in a mutant lacking flagellin protein. (A) Transformation frequency of *B. subtilis* wild type and *hag*-mutant after 6 h incubation in competence medium (unpaired two-sample t-test with Welch Correction: P = 3.1·10^{-5}, n = 9). The inset shows a zoom-in of the *hag*-mutant data. (B) Growth dynamics of wild type and hag-mutant during 12 h incubation in competence medium. Standard deviations for the measurements are depicted in light grey (unpaired two-sample t-test with Welch Correction: P = 0.001, n = 5). Arrows indicate the time points of DNA addition to investigate the transformation frequency over time, which is shown as box-and-whisker plot in (C). The line in the boxes represents the median, the box indicates the 25[th]-75[th] percentile. Asterisks indicate statistically significant differences between wild type and *hag*-mutant (unpaired two-sample t-test with Welch Correction for WT - Δ*hag*:, P = 0.125 for 3 h, P < 0.01 for 4 h, 5 h, 6 h; n = 6).

Fig. 2. Mutants impaired in flagellar function exhibit lower transformation frequencies. Deletion of

the gene encoding a flagellar stator (*motA*; A) or the gene encoding the hook protein (*flgE*; B) results

in significantly lower transformation frequency of the respective strain compared to the wild type

after incubation in competence medium for 6 h. The line in the boxes represents the median, the

box indicates 25th-75th percentile. Asterisks indicate statistically significant differences (unpaired

two-sample t-test with Welch Correction: P = 0.01 for Δ*motA*; P = 0.039 for Δ*flgE*; n = 9 for all).

Fig. 3. Fewer cells of the *hag*-mutant express competence genes compared to the wild type. (A)

Representative microscopy images of strains harboring the P*comG-gfp* reporter in wild type, Δ*hag* or

Δ*comK* genetic background. Images were recorded after incubation in competence medium for 5 h.

The scale bar represents 50 μm. (B) Histograms of flow cytometric measurements showing the cell

count and the fluorescence in arbitrary units for wild type, Δ*hag*, and Δ*comK* including background

fluorescence. Representative images are shown for each strain. (C) Percentage of fluorescent cells

determined from the data in (B) for wild type and *hag*-mutant by isolating the fluorescent

population with fluorescence intensities above 3 A.U. Asterisks indicate significant differences

(unpaired two-sample t-test with Welch Correction: P = 0.036, n = 3).

Fig. 4. Competence gene expression is reduced in mutants lacking a functional flagellum. (A)

Representative microscopy images of strains harboring the P$_{comG}$-*gfp* reporter in wild type, Δ*motA*, or

Δ*flgE* genetic background. Images were recorded after 5 h incubation in competence medium. The

scale bar represents 50 µm. (B) Histograms of flow cytometric measurements showing the cell count

and the fluorescence in arbitrary units for wild type, Δ*motA* or Δ*flgE*. Representative images are

shown for each strain. (C) Percentage of fluorescent cells determined from the data in (B) by

isolating the fluorescent population with fluorescence intensities above 3 A.U. showing a significant

difference (asterisks) between wild type and Δ*motA* (P = 0.017) as well as wild type and Δ*flgE* (P <

0.001) with n = 3 for both (unpaired two-sample t-test with Welch Correction).

Fig. 5. Synthetically induced *comK* and *degU*146 increase competence of Δ*hag*. (A) Transformation frequencies of wild type compared to strains harboring a xylose-inducible copy of *comK* (P$_{xyl}$-*comK*) with wild type or Δ*hag* genetic background (Kruskal-Wallis test for WT - WT P$_{xyl}$-comK: P = 3.4·10^{-4}; for WT P$_{xyl}$-*comK* - Δ*hag* P$_{xyl}$-*comK*: P = 3.4·10^{-4}, n = 9 for both). (B) Transformation frequencies of WT compared to strains harboring either a phosphorylated DegU variant (*degU*32) or a non-phosphorylatable DegU variant (*degU*146) in wild type or Δ*hag* background. Strain Δ*hag degU*32 is significantly different from all other strains (Kruskal-Wallis test: P < 0.05 for all, n=6). The line in the boxes represents the median, the box indicates 25th-75th percentile. Asterisks indicate statistically significant differences (Kruskal-Wallis test: P = 0.007, n=6).

523 **Fig. 6.** Competence is improved in viscous medium. Transformation frequency of the wild type strain

524 grown in normal competence medium and in medium with increased viscosity. The line in the boxes

525 represents the median, the box indicates 25th-75th percentile, # indicates marginally significant

526 differences (unpaired two-sample t-test: P = 0.095, n = 4).