Hippocampal atrophy and intrinsic brain network alterations relate to impaired capacity for mind wandering in neurodegeneration

Authors

Claire O'Callaghan,[1,2] James M. Shine,[2] John R. Hodges,[2,3,4] Jessica R. Andrews-Hanna,[5] and Muireann Irish[2,3,6]

Affiliations

[1] Department of Psychiatry and Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge, UK

[2] Brain and Mind Centre, University of Sydney, Sydney, Australia

[3] Australian Research Council Centre of Excellence in Cognition and its Disorders, Sydney, Australia

[4] Sydney Medical School, University of Sydney, Sydney, Australia

[5] Institute of Cognitive Science, University of Colorado Boulder, Boulder, CO

[6] School of Psychology, University of Sydney, Sydney, Australia

Corresponding author

Associate Professor Muireann Irish, Brain and Mind Centre and School of Psychology, the University of Sydney, 100 Mallet Street, Camperdown, NSW 2050, Australia.
+61 2 9114 4165; muireann.irish@sydney.edu.au

Figures: 5   **Tables:** 2

Supplementary figures: 4   **Supplementary tables:** 1

Abstract

Mind wandering represents the human capacity for internally focussed thought, and relies upon dynamic interactions between default and frontoparietal networks. The majority of studies in the field have characterised mind wandering in healthy people, yet there is limited understanding of how this capacity is affected in clinical populations. The present study used a validated thought sampling task, to probe the capacity for mind wandering in two neurodegenerative disorders; the behavioural variant of frontotemporal dementia (n=28) and Alzheimer's disease (n=22), compared to healthy older controls (n=28). These disorders were selected due to their canonical profiles of neural dysfunction across key sites of the default and frontoparietal networks. Behaviourally, mind wandering frequency was found to be reduced in the patient groups, leading to an increase in stimulus-bound thoughts. These behavioural profiles were associated with distinct regions of grey matter loss, as revealed by voxel-based morphometry, predominantly in the hippocampal complex and striatum. Resting state functional connectivity further revealed associations between impaired mind wandering performance and altered connectivity within and between regions of the frontoparietal and default networks. Together, these findings are the first to describe altered mind wandering in neurodegenerative disorders, which was associated with hippocampal atrophy and aberrations in the functional integrity of the default and frontoparietal networks. These results corroborate current theoretical frameworks emphasising that cooperation between default and frontoparietal regions is critical for producing and sustaining internally focussed thought. Notably this study reveals a new dimension of cognitive dysfunction not previously documented in neurodegenerative disorders.

Mind wandering – the ability to disengage from the external environment and direct attention inwardly to a stream of thought – is fundamental to the human experience. Dynamic interactions within and between large-scale brain networks govern the initiation and maintenance of mind wandering (1, 2). Of particular interest in this context are interactions between the default network and the frontoparietal control network (3-7).

In a recently proposed framework, spontaneous and unconstrained internally oriented thought is generated by fluctuations in the medial temporal lobe system of the default network, with weak influence from frontoparietal regions (2). More deliberative thought, for example planning or thinking through a problem, corresponds to reduced variability in the medial temporal system, and increased coupling between the frontoparietal network and default network core (2). The medial temporal lobe system therefore emerges as influential in the origin of spontaneous thoughts, with frontoparietal control regions becoming increasingly important for subsequent elaboration and metacognitive processing (8).

Exploring mind wandering in clinical populations can provide unique information about its cognitive and neural substrates. Altered mind wandering is documented in many clinical populations, and may constitute an important neurocognitive endophenotype across disorders. Perseverative mind wandering that is more frequent or salient, with negative content, has been reported in depressive rumination, neuroticism and dysphoria (9-11). These perseverative and ruminative styles of mind wandering are suggested to reflect an overly constrained mode of function in the default network, leading to excessive stability of thoughts (2). In contrast, higher rates of unintentional, spontaneous mind wandering are associated with increased obsessive-compulsive and attention-deficit/hyperactivity symptomatology in non-clinical samples (12, 13). Similarly, higher frequencies of mind wandering have been noted in schizophrenia, which correlate with the severity of positive symptoms (14). An unconstrained default network, due to local hyperactivity or relaxed influence from frontoparietal regions, may underpin such excessive variation and incoherence of thoughts, as seen in psychosis (2). Crucially, however, no study to date has directly investigated the relationship between mind wandering capacity and network alterations in neuropsychiatric populations. Instead, putative neural correlates of mind wandering in clinical disorders have been inferred based on the known network alterations in those diseases. As such, it remains unclear how pathological brain states impact the frequency and phenomenology of mind wandering.

The present study directly tests whether regional changes in the default and frontoparietal networks are associated with alterations in mind wandering capacity, by focusing on neurodegenerative disorders. These syndromes afford a unique opportunity to study the impact of network level dysfunction on mind wandering, given well established pathology primarily targeting, but not restricted to, key nodes of the default and frontoparietal networks (15-17). This approach is an important extension to recent work confirming that focal lesions to nodes of the default network, in the hippocampus and medial prefrontal cortex, impact mind wandering (18, 19). Moreover, on the cognitive level, mind wandering is supported by distinct component processes, many of which are disrupted in neurodegenerative populations with dementia. These underlying processes include flexible allocation of internally vs. externally focussed attention. Also, episodic memory, prospection, mental imagery, and socio-emotional processing, which form the content of mind wandering; and, working memory processes, which sustain a stream of thought, buffering it from distraction (20-22). Dementia syndromes are associated with marked disruption across many of these component processes, for example autobiographical memory retrieval (23, 24), mental simulation (25-27), working memory and attentional allocation (28, 29). Given this well-established dysfunction in key brain regions, and the disruption of core cognitive processes fundamental to constructing and maintaining spontaneous thought, it follows that distinct changes in the frequency and phenomenology of mind wandering should be present in dementia (30). No study to date, however, has empirically investigated the capacity for mind wandering in dementia, yet this line of enquiry is crucial to establish how damage to functional brain networks impacts internally generated thought processes, whilst yielding important new insights into the cognitive symptomatology of these syndromes.

To that end, we explored mind wandering capacity in two dementia subtypes: Alzheimer's disease (AD) and behavioural variant frontotemporal dementia (bvFTD). AD, characterised by prominent memory deficits, is associated with pathological changes in the default and frontoparietal networks, particularly the hippocampus, medial temporal lobe subsystem and posterior cingulate cortex, extending into prefrontal and parietal regions with disease progression (17, 31-35). In contrast, bvFTD is distinguished by behavioural dysfunction, including disinhibition, apathy, loss of empathy, emotional blunting, stereotypical behaviours, and loss of insight (36, 37). Early pathological changes in

bvFTD are most prominent in regions comprising the salience and default networks, including the dorsomedial and ventromedial prefrontal cortices, as well as widespread changes across the insula, amygdalae, thalamus and striatum with disease progression (17, 32, 38-40).

Quantifying the nature and content of mind wandering in clinical disorders is inherently challenging, presumably reflecting the dearth of studies in this field. Dominant experimental approaches to mind wandering typically require subjects to monitor or self-identify extraneous thoughts during performance of an ongoing cognitive task (3, 41-43). Such approaches, however, rely on dual-tasking and metacognitive capacities that are diminished in dementia, limiting the extent to which reliable conclusions can be drawn from existing measures. To circumvent these methodological constraints, we employed the "Shape Expectations" task – a paradigm developed to measure mind wandering under conditions of low cognitive demand, which was validated in non-pathological ageing and performance was shown to correlate with resting state variations in default network connectivity (44). In keeping with existing frameworks, the task quantifies mind wandering as thoughts unrelated to the current environment or to the task at hand, and it can be used to quantify mind wandering propensity along a conceptual continuum (21, 45).

The objectives of this study were twofold. First, we aimed to quantify the capacity for mind wandering in dementia syndromes during conditions of low cognitive demand using a validated experimental task. Second, we sought to characterise how disease-related alterations in (i) regional grey matter, and (ii) functional connectivity in the default and frontoparietal networks, relate to changes in mind wandering performance. In doing so, we aimed to validate current frameworks of mind wandering in a clinical model, by showing that the integrity of the default and frontoparietal networks is essential to support mind wandering capacity.

Results

Overall mind wandering performance

The Shape Expectations task probes mind wandering capacity under conditions of low cognitive demand using an experience-sampling procedure across 9 experimental trials (see (44). The scoring system conceptualises mind wandering along a continuum, ranging from 1

to 4; level 1 represents stimulus-bound thought and level 4 represents mind wandering. To calculate overall mind wandering performances, total percentages of each level across the task were calculated (i.e., total counts of 1s, 2s, 3s or 4s divided by 9 trials).

Figure 1 displays the percentage of responses at each scoring level of the mind wandering continuum. A repeated measures ANOVA revealed no main effect of response level ($F(3, 225) = 1.59$, $p = .193$), however, a significant main effect of group ($F(2, 75) = 4.67$, $p < .05$) was present. Between group Sidak post-hoc comparisons revealed significantly higher levels of stimulus-bound responses (Level 1) in bvFTD compared to control ($p < .0001$) and AD ($p < .05$) participants. Numerically, controls had less stimulus bound instances than AD patients although this difference was not statistically significant ($p = .30$). Further, bvFTD patients showed attenuated Level 2 responses relative to the AD and control groups (all p values < .05).

A significant group by response level interaction was observed ($F(6, 225) = 4.60$, $p < .0001$). Within-group post-hoc tests revealed significantly fewer stimulus-bound responses (Level 1) in controls relative to all other response categories (all p values < .05). In contrast, stimulus-bound responses were significantly elevated in the bvFTD group relative to Level 2 responses ($p < .05$). Finally, AD responses did not differ significantly across the four levels (p values > .20).

Figure 1 – Overall proportion of mind wandering scores across groups

Effect of trial duration on mind wandering frequency

As the experimental trials were of varying durations (Short: ≤20 seconds, Medium: 30-60 seconds, Long: ≥90 seconds), we explored whether trial length was related to the average score obtained across each of the three trial durations. A significant main effect was observed for trial duration ($F(2, 150) = 15.33$, $p < .0001$) with higher average scores achieved on long trials relative to all short and medium durations (p values $< .0001$), irrespective of group. A significant main effect of group was also evident ($F(2, 75) = 4.24$, $p = .018$) indicative of the bvFTD group achieving lower average scores relative to controls across all durations ($p=.014$). No other significant differences were evident at the group level (all p values $> .3$). Further, the group x duration interaction was not significant ($F(4, 150) = 1.66$, $p =.163$) (See Supplementary Figure 1).

Mind wandering index score

To directly examine stimulus bound (level 1) vs. mind wandering (level 4) responses, an index score was created by subtracting the % level 1 responses from % level 4. A larger, positive index score reflects a tendency to engage in mind wandering as opposed to stimulus bound thought, with negative scores reflecting the reverse profile.

Figure 2 shows the average mind wandering index score across participant groups, providing a measure of the propensity to engage in mind wandering versus stimulus bound thought (i.e., % Level 4 – % Level 1 responses). Significant group differences were observed ($F(2, 75) = 7.12$, $p < .01$), driven exclusively by the bvFTD group. Relative to controls, bvFTD patients displayed an increased propensity for stimulus-bound thought at the expense of pure mind wandering ($p < .01$). In contrast, no such differences were evident in the AD group ($p = .453$), although their mind wandering index was numerically lower than that of controls.

Figure 2 – Average mind wandering index score across participant groups

Legend – Mind wandering index score (i.e., % difference in Level 4 minus Level 1 responses), reflecting the propensity to engage in mind wandering relative to stimulus bound thought, averaged across groups. Higher scores reflect an increased propensity to engage in mind wandering as opposed to stimulus bound thought; with lower scores reflecting a tendency toward stimulus bound thought. Error bars show the standard error of the mean. **$p < .01$.

Grey matter correlates of mind wandering performance

Voxel-based morphometry was used to determine the relationship between scores on the mind wandering index and regional grey matter loss. Figure 3 displays the common grey matter regions that covaried with the mind wandering index score in AD and bvFTD patient groups. Three distinct grey matter clusters were identified (Table 2) including: (i) a right striatal cluster encompassing the caudate head, extending to the nucleus accumbens; (ii) a left striatal cluster including the dorsal caudate and extending posteriorly to the thalamus; and (iii) a medial temporal lobe cluster involving the left hippocampus and parahippocampal gyrus.

Table 2. Voxel-based morphometry (VBM) overlap analysis showing common grey matter correlates of mind wandering performance in bvFTD and AD patients.

Regions	Hemisphere (L/R/B)	MNI coordinates for voxel of maximal intensity			Number of voxels	*p*
		X	*Y*	*Z*		
Caudate head; nucleus accumbens	R	12	0	16	316	.001
Dorsal caudate, thalamus	L	-14	-4	14	263	.001
Posterior hippocampus; parahippocampal gyrus	L	-24	-38	-6	114	.001

Results derived from an overlap masking procedure, using voxelwise approach and reported at *p* < .005 uncorrected with a cluster extent threshold of 100 contiguous voxels. L = Left, R = Right; MNI = Montreal Neurological Institute

Figure 3 – Common grey matter regions associated with the mind wandering index score across patient groups

Overlap analysis showing grey matter regions significantly associated with the mind wandering index score in both bvFTD and AD patients. a) Right striatum cluster: right caudate head extending to nucleus accumbens; b) Left striatum cluster: left dorsal caudate extending to thalamus; c) Left hippocampal/parahippocampal gyrus cluster. Results reported at *p* < .005 uncorrected with a cluster extent threshold of 100 contiguous voxels.

For all participants, mean grey matter volumes were then extracted from the regions identified in the overlap analysis. These regional volumes were plotted to show the directionality of the relationship between mind wandering index and grey matter. As illustrated in Figure 4, grey matter volume in each of the three key regions was positively correlated with the mind wandering index score: right striatum: *r* = .56, *p* < .0001; left striatum: *r* = .67, *p* < .0001; hippocampus/parahippocampus: *r* = .50, *p* < .0001. This is

consistent with decreased grey matter in these regions being associated with a reduced mind wandering frequency and concomitant increase in stimulus-bound thought.

Figure 4 – Association between grey matter volume of the three clusters of interest and mind wandering index scores for all participants

Relationship between grey matter volumes in the three significant clusters and mind wandering index score, plotted for all participant groups: controls (blue), bvFTD (green), AD (red). a) Right striatal cluster; b) Left striatal cluster; c) Left hippocampal/parahippocampal cluster. Grey matter volumes in mm^3

Seed region connectivity and mind wandering performance

The relationship between seed region connectivity and the mind wandering index score was then examined. Regions where the relationship between mind wandering index and connectivity differed significantly between the groups are shown in Figure 5. Attenuated mind wandering performance in bvFTD relative to controls, as captured by the mind wandering index, was associated with a pattern of reduced core and frontal connectivity, but increased intra-hippocampal connectivity. Specifically, bvFTD patients' mind wandering index score was associated with reduced connectivity between the left DLPFC and left PCC ($Z = -2.63$, *p* < .05), and between the left DLPFC and right amPFC (Z = -2.36, p < .05), in the context of stronger connectivity between the left anterior and posterior hippocampus ($Z = 2.19$, p < .05).

Relative to controls, AD patients' mind wandering index was associated with reduced core and dorsal PFC connectivity between the left PCC and right DLPFC (Z = -2.19, p < .05), in the context of increased connectivity between the hippocampus and medial PFC (right anterior hippocampus to right amPFC: Z = 2.58, p < .05; left posterior hippocampus to left amPFC: Z = 2.04, p < .05).

Comparing the two patient groups, the mind wandering index in bvFTD patients was associated with weaker inter-hemispheric dorsolateral connectivity relative to AD (left DLPFC to right DLPFC: Z = -2.10, p < .05) and stronger inter-hemispheric hippocampal connectivity (left anterior to right posterior hippocampi: Z = 2.09, p < .05). No other between group comparisons emerged as significant.

Figure 5 – Seed region connectivity and mind wandering performance

Seed regions where the relationship between mind wandering index and connectivity differed significantly between the groups. FPN = frontoparietal network; DN = default network; HC = hippocampus. Results shown in neurological convention (i.e., left and right side of brain shown on left and right side of image).

Discussion

Our results provide an empirical demonstration of distinct changes in mind wandering capacity in two dementia syndromes, associated with discrete grey matter correlates and functional connectivity changes in the default and frontoparietal networks. To our knowledge, this is the first investigation of mind wandering in dementia, and its associated neural correlates. These findings corroborate current theoretical frameworks that emphasise the essential role of the medial temporal lobe, and interactions between the default and frontoparietal networks, in the generation and evolution of mind wandering (2).

Our most striking behavioural finding was a clear bias toward stimulus-bound thought, at the expense of fully-fledged instances of mind wandering, in individuals with bvFTD. This manifested as a negative mind wandering index, suggesting an inability to shift or progress thoughts beyond stimuli in the immediate environment. By contrast, healthy older controls displayed a positive mind wandering index, reflecting preferential engagement in mind wandering under periods of low cognitive demand, in line with previous findings using this task (44). This tendency to mind wander in healthy individuals is to be expected given the monotonous nature of the stimuli, and minimal task requirements, designed to provoke off-task thought. While a decreased propensity to engage in mind wandering versus stimulus bound thought (as evidenced by the mind wandering index) was observed in AD relative to controls, these differences did not emerge as statistically significant. Whether the mind wandering index would continue to decrease in AD with disease progression remains an important question that longitudinal studies would be well placed to address. Nevertheless, taken together, the behavioural data suggests distinct changes in the distribution of Level 1 to Level 4 responses on the Shape Expectations task in dementia syndromes, with a shift towards stimulus-bound forms of thought at the expense of mind wandering.

Existing evidence suggests that component processes supporting mind wandering are compromised in dementia, most notably those processes involving memory-based

constructive simulation (30, 46). AD and bvFTD patients display comparable episodic memory dysfunction (47-50). Both groups also display marked impairments in future-oriented forms of thinking, including prospective memory (51, 52) and constructive simulation of future episodes (26, 53, 54). Accordingly, reductions in mind wandering performance commensurate with the magnitude of these memory-based constructive impairments might be predicted. Our results, however, suggest that mind wandering capacity is more vulnerable in bvFTD, as it was significantly impaired relative to controls. In contrast to AD, bvFTD is associated with marked behavioural rigidity, manifesting as preservative behaviour and adherence to stereotypical routines (55, 56). This has been quantified via cognitive tasks where bvFTD patients show limited verbal generativity (29), and reduced capacity for divergent and flexible thinking (57) and problem solving (58). An intriguing possibility exists whereby this narrowed behavioural and cognitive repertoire in bvFTD may reflect, in part, the reduced ability to generate spontaneous thoughts and to transition flexibly between them.

Our neuroimaging findings underscore the role of key regions of the default and frontoparietal networks in supporting internally generated thought, corroborating previous reports in healthy individuals (3, 6, 7). Irrespective of dementia subtype, mind wandering index scores were associated with decreased connectivity between the default network core (PCC) and the frontoparietal network (dlPFC). Cooperation between the frontoparietal and default network is thought to support processes relevant to mind wandering, including autobiographical planning, recollection, mental simulation, and creativity (59-62). Indeed, higher trait levels of deliberative mind wandering in healthy individuals are associated with increased cortical thickness, and increased functional connectivity, in regions that bridge the default and frontoparietal networks and promote their integration (63). Limited evidence from task-based fMRI of AD patients also supports this cooperative network interaction, whereby performance on memory tasks is not only mediated by default network regions, but also depends on recruitment of prefrontal executive regions, including the dlPFC (64, 65). The posterior cingulate cortex may play a key role in default-frontoparietal network cooperation, by maintaining connectivity with the default network and simultaneously facilitating integration between other default network and frontoparietal regions in a dynamic and context-dependent manner (62). PCC dysfunction in dementia might therefore impair the maintenance of deliberative thought processes. Given that changes in the PCC are evident from the mild cognitive impairment stages that precede Alzheimer's disease (66), the impact

of relatively focal PCC dysfunction on mind wandering style in prodromal AD warrants further attention.

The mind wandering index was also associated with hyper-connectivity within the hippocampus in bvFTD, and from the hippocampus to the amPFC in AD. Furthermore, in both patient groups, decreased grey matter in the left hippocampus and parahippocampus correlated with reduced mind wandering. Convergent measures have shown that neural activity in the hippocampus and parahippocampus (8), entorhinal cortex (67) and temporal cortex (68) precedes spontaneous free recall of episodic memories. This accords with a large body of rodent work implicating hippocampal sharp wave ripple events (SWRs) in the replay (and preplay) of previously learnt or future behavioural sequences (69-74). In combined intracranial recordings with resting state fMRI in anesthetised monkeys, hippocampal SWRs precede increased activation of the default network (75). Together, these findings link spontaneous activation in the hippocampus and surrounding regions with both recall and prospection, which may then engage the default network more broadly to support the elaboration of memories and simulations. Consistent with this proposal, higher trait levels of spontaneous mind wandering are correlated with increased functional connectivity in medial temporal regions of the default network, and with increased cortical thickness in the retrosplenial cortex (63) – a region adjacent to the posterior hippocampal and parahippocampal regions identified in our VBM analysis. A critical role for the hippocampus in mind wandering was also recently confirmed, as individuals with selective bilateral hippocampal damage exhibited reduced diversity in their mind wandering content (18). Our findings complement and extend these results by confirming that functional and structural alterations in the hippocampus and parahippocampus mediate alterations in mind wandering in dementia.

Taken together, our results suggest that disrupted capacity for mind wandering in both AD and bvFTD reflects a combination of decreased frontoparietal-default network connectivity and increased hippocampal connectivity. Direct comparison of the patient groups revealed this pattern to be amplified in bvFTD, as their mind wandering index was associated with reduced bilateral dlPFC connectivity and increased bilateral hippocampal connectivity, relative to AD. This resonates with our behavioural finding of disproportionately impaired mind wandering in the bvFTD group. Abnormalities in the default and frontoparietal

networks are well established in AD and bvFTD; however, comparatively greater salience network dysfunction is observed in bvFTD, encompassing the anterior cingulate cortex and anterior insula (17, 76). Reduced causal influence from the salience network over the default network has also been shown during a reasoning task in bvFTD (77). The salience network is proposed to mediate dynamic shifts between default and executive control networks (78), facilitating transitions between external and internal focus. Greater salience network dysfunction and its reduced causal influence on the default network may impair bvFTD patients' ability to direct attention away from the external environment. This may disrupt the capacity for internally oriented thought as attention remains focussed on external information, consistent with the high levels of stimulus-bound thought bvFTD patients exhibited on our task.

Finally, we found an association between striatal grey matter loss and reduced mind wandering, irrespective of dementia subtype. The basal ganglia represents a network hub vulnerable to degeneration across neurodegenerative disorders (79), and abnormalities across the striatum are suggested to contribute to an array of cognitive and neuropsychiatric features in these conditions (80). We speculate that the involvement of the basal ganglia in supporting large scale network communication (81) may explain its association with attenuated mind wandering in dementia. This is consistent with known functional connectivity between the striatum and large-scale cortical networks, including default and frontoparietal (82), and the convergence of these functional networks in distinct zones of the striatum (83). Striatal degeneration may impair the integration of information from disparate brain networks, which is necessary to support more abstract forms of cognition, for example mind wandering. As the dynamic integration and segregation of network communication is increasingly recognised as important for mind wandering (1, 84), further investigation of the striatal influence over large scale network topology in this context is warranted.

In summary, to our knowledge, this is the first study to empirically measure mind wandering capacity in dementia syndromes, and reveals a definitive shift towards stimulus-bound thought at the expense of mind wandering. This narrowed range of internal mentation relates to hippocampal and striatal grey matter loss, and functional changes in the default and frontoparietal networks. Our findings corroborate current theoretical frameworks which emphasise the cooperation between default and frontoparietal regions in producing and sustaining internally focussed thought (2). Future work is needed to identify the trait level

and phenomenological characteristics of altered mind wandering in dementia, and to determine how reductions in this adaptive and enriching cognitive capacity relates to broader cognitive and behavioural symptoms in these syndromes. Given the ubiquity of mind wandering in everyday life, we stress the importance of understanding how loss of this fundamental human capacity impacts wellbeing and sense of self in individuals living with dementia.

Methods and Materials

Case selection

See Table 1 for demographic and clinical details. Twenty-eight individuals meeting current diagnostic criteria for bvFTD (85) and 22 individuals with a clinically probable diagnosis of AD (86) were recruited from the FRONTIER research group in Sydney, Australia. Briefly, bvFTD patients presented with marked changes in behaviour and personality, social and emotional dysregulation and executive dysfunction. AD patients were characterised by significant episodic memory dysfunction, temporal disorientation, and visuospatial deficits, with relatively preserved socio-emotional function. Patient groups showed characteristic patterns of brain atrophy on structural magnetic resonance imaging (MRI). Global cognitive function was assessed using the Addenbrooke's Cognitive Examination-Revised (ACE-R; 87), covering domains of attention and orientation, memory, fluency, language, and visuospatial function. The Cambridge Behavioural Inventory-Revised (CBI-R; 88) was included as an informant-rated scale to assess functional status and behavioural disturbance. Twenty-eight age-matched control participants were recruited from a volunteer research panel and screened for cognitive impairment using an established cut-off of 88 or above on the ACE-R. Exclusion criteria for all participants included prior history of mental illness, significant head injury, movement disorders, cerebrovascular disease, alcohol and other drug abuse, and limited English proficiency. The study was approved by the South Eastern Sydney Local Area Health and University of New South Wales ethics committees and all participants provided informed consent in accordance with the Declaration of Helsinki.

Demographics and clinical characteristics

Demographic details and clinical characteristics of study participants are presented in Table 1. The groups were matched for gender distribution (χ^2 = 1.7, p =.419), however age differences were present (F(2,75) = 4.8, p = .011) reflecting that bvFTD patients were

significantly younger than controls ($p = .016$). Global cognitive function differed across the groups (ACE-R: $F(2,75) = 47.1$, $p < .0001$), with both patient groups performing significantly worse than controls (p values $< .0001$). While AD patients displayed greater cognitive impairment relative to bvFTD (ACE-R: $p =. 036$), greater functional and behavioural decline was evident in the bvFTD relative to AD group (CBI-R: ($t(48) = -2.7$, $p = .009$). Patient groups were matched for disease duration (years elapsed since onset of symptoms: $t(48) = .698$, $p = .488$)

Table 1. Demographics and clinical characteristics of study participants.

	bvFTD	AD	Controls	bvFTD vs Control	AD vs Control	bvFTD vs AD
N	28	22	28	-	-	-
Sex (M:F)	17:11	15:7	14:14	n.s.	n.s.	n.s.
Age	61.5 (8.9)	67.0 (9.6)	67.9 (6.4)	*	n.s.	n.s.
Disease duration§	4.4 (3.3)	5.1 (3.9)	-	-	-	n.s.
CBI-R (% endorsed)	35.5 (11.1)	25.4 (13.8)	-	-	-	**
ACE-R total (max 100)	75.9 (14.1)	67.7 (12.9)	96.2 (2.6)	***	***	*

Standard deviations presented in parentheses. n.s. = non significant; * = $p < .05$; ** = $p < .01$; *** = $p < .001$
§Years since symptom onset; bvFTD = behavioural variant frontotemporal dementia; AD = Alzheimer's disease; CBI-R = Cambridge Behavioural Inventory-Revised (higher scores indicating greater levels of functional decline and behavioural disturbance); ACE-R = Addenbroke's Cognitive Examination-Revised.

Mind wandering experimental task

Participants viewed static, two-dimensional coloured geometric shapes presented individually on a computer screen, following which they were prompted to report on their thoughts that arose during stimulus presentation. The task consisted of 9 trials, each presenting a commonplace shape (e.g., blue square, yellow circle) for varying durations (Short: ≤20 seconds, Medium: 30-60 seconds, Long: ≥90 seconds). Total testing time was approximately 15 minutes. Responses were digitally recorded and transcribed for scoring.

The scoring system represents a continuum ranging from levels 1-4. Level 1 represents stimulus-bound thoughts, for example if a participant reports thinking about only what was presented on the screen, i.e., "a blue square". Level 4 responses, by contrast, are fully-fledged instances of mind wandering, where the participant's thoughts are completely unrelated to the stimulus or the task at hand. Levels 2 and 3 represent intermediary responses. The scoring protocol and representative participant responses are included in Supplementary Information.

The final score awarded for each trial is the highest level achieved on that trial, ranging from 1-4. Total percentages of each level across the task were calculated (i.e., total counts of 1s, 2s, 3s or 4s divided by 9). A mind wandering index was also derived to compare the extent of level 1 (stimulus bound) with level 4 (mind wandering) responses. This was achieved by subtracting the % level 1 responses from % level 4. The mind wandering index score was used as a covariate in the neuroimaging analysis, rather than the percentage of mind wandering (level 4) as many patients scored 0 for this, leading to reduced variance in the sample.

Statistical analysis

Behavioural analyses were conducted using the open-source statistical environment R (R Core Team, 2013). Demographic variables were compared via one-way ANOVA or independent samples t-tests. *A priori*, variables were plotted and checked for normality of distribution using Kolmogorov–Smirnov tests. Performance on the Shape Expectations task was compared via one-way ANOVA and repeated measures ANOVA, with Sidak post hoc comparisons.

Imaging acquisition

Participants underwent whole-brain T1 weighted imaging using a 3T Philips MRI scanner with standard quadrature head coil (eight channels) using the following sequences: coronal orientation, matrix 256 x 256, 200 slices, 1 mm^2 in-plane resolution, slice thickness 1 mm, echo time/repetition time = 2.6/5.8 ms, flip angle $\alpha = 8°$. Structural scans were not available for three bvFTD patients, one AD patient, and three controls due to imaging contraindications (e.g., pacemaker; metallic stent).

A subset of 18 bvFTD, 15 AD, and 15 controls underwent task-free resting state imaging. T2*-weighted echo planar functional images were acquired in sequential order with repetition

time (TR) = 2 s, echo time (TE) = 30 ms, flip angle = 90°, 32 axial slices covering the whole brain, field of view (FOV) = 240 mm, inter-slice gap = 0 mm, and raw voxel size = 2.1 mm by 2.6 mm by 4.5 mm thick. T1-weighted images were used for co-registration with functional images. A single 7-min run was performed in the scanner, which consisted of the participant lying still with their eyes open. Participants were questioned immediately afterwards to ensure they had not fallen asleep during the sequence.

Voxel-based morphometry (VBM) pre-processing and analysis

Voxel-based morphometry (VBM) was performed on the three dimensional T1-weighted scans, using the FSL-VBM toolbox in the FMRIB software library package FSL (http://www.fmrib.ox.ac.uk/fsl/). Scans were skull-stripped using the BET algorithm in FSL (89) and tissue segmentation was completed using FMRIB's Automatic Segmentation Tool (FAST v4.0) (90). A study-specific grey matter template was created by canvassing the maximum equal amounts of scans across the three groups and registered to the Montreal Neurological Institute Standard space (MNI 152) using a non-linear b-spline representation of the registration warp field. Grey matter partial volume maps were non-linearly registered to the study template and modulated by dividing by the Jacobian of the warp field, to correct for any contraction/enlargement caused by the non-linear component of the transformation (91). After normalisation and modulation, the grey matter maps were smoothed using an isotropic Gaussian kernel (standard deviation = 3 mm; full width half maximum = 8 mm).

VBM group comparisons

Group differences in grey matter intensity across patients and controls were compared using a voxelwise general linear model (GLM). Significant clusters were formed using the threshold-free cluster enhancement (TFCE) method (92) via permutation-based nonparametric testing with 5000 permutations per contrast (93). Group differences in grey matter intensity were assessed using t-tests, tested for significance at $p < .01$, corrected for multiple comparisons via family-wise error (FWE) correction across space. Age was added as a covariate in the analysis given the significant age differences between the control and bvFTD groups. Results of group grey matter comparisons are reported in Supplementary information. Relative to controls, both patient groups showed characteristic patterns of widespread grey matter atrophy, as shown in Supplementary Table 1 and Supplementary Figure 2.

VBM analysis of mind wandering performance

To examine the relationship between grey matter intensity and mind wandering performance, separate voxelwise GLMs were conducted within the bvFTD and the AD patient groups (not combined with controls) using the mind wandering index score as a covariate in the design matrix. An overlap analysis was conducted using the results from each covariate analysis to identify grey matter regions that were commonly implicated across the patient groups. The contrasts of the statistical maps generated from the bvFTD and AD covariate analyses were scaled to a common threshold ($p < .005$, uncorrected) and multiplied to create an inclusive, or overlap, mask across patient groups. A conservative cluster extent threshold of 100 contiguous voxels was employed to guard against false positive findings.

Resting state fMRI (rsfMRI) pre-processing and analysis

rsfMRI scans were pre-processed using FEAT (FMRI Expert Analysis Tool), from the FSL FMRIB software library. Scans were motion corrected using MCFLIRT (94); six motion parameters (estimated by MCFLIRT) and cerebrospinal fluid and white matter signals were regressed out using the CompCor strategy (95). Mean framewise displacement (FD) values were calculated from MCFLIRT. Mean FD values were: controls = 0.149 ± 0.06; AD = 0.117 ± 0.33; bvFTD = 0.247 ± 0.19. There were no differences in FD values between the groups: AD vs. control $t = 1.449$; $p = 0.159$; bvFTD vs control $t = 1.861$; $p = 0.148$; AD vs. bvFTD $t = 1.598$; $p = 0.117$. Mean FD was also shown not to correlate significantly with the mind wandering index scored used as a covariate in the imaging analysis ($r = -0.147$; $p = 0.324$). Following these pre-statistics, linear registrations were performed using FLIRT: participants' rsfMRI scans were registered to their T1 scans that had been skull-stripped using BET; and the T1 scans were registered to standard space (MNI 152). These transformations were concatenated and applied to the raw EPI scans, registering the rsfMRI scans to standard space.

Seed regions of interest of a 4mm diameter were created in standard space, using the fslmaths function in FSL. The location of the seeds was selected based on previously published co-ordinates of key hubs within the default network (5), frontoparietal network (96), and hippocampus (97). The default network seeds included the hippocampal formation (HF; ±22 -20 -26), a midline core, which included the posterior cingulate cortex (PCC; ±8 -56 26) and the anteromedial prefrontal cortex (amPFC; ±6 52 -2); and the ventromedial prefrontal cortex (vmPFC; 0 26 -18) (5). Frontoparietal network seeds included the dorsolateral prefrontal cortex (right DLPFC; 43 22 34, left DLPFC; -43 22 34) (96). The hippocampus was seeded

posteriorly (left -23 -26 -15, right 25 -26 -15) and anteriorly (left -18 -14 -18, right 18 -14 -18) (97). Using FLIRT, the ROIs were registered to individuals' rsfMRI scans using the transformations derived during pre-processing.

Group comparisons of seed region connectivity

Patterns of seed region connectivity were initially examined. Within the groups, one sample t-tests were conducted between each pairwise edge for the 13 regions of interest, corrected for multiple comparisons using a false discovery rate of $q \leq .05$. Group differences in seed region connectivity are shown in Supplementary Figures 3 and 4.

Seed region connectivity and mind wandering performance

Group differences in the relationship between seed region connectivity and the mind wandering index score were then examined. To establish the relationship between connectivity and mind wandering performance, within each of the three groups, participants' mind wandering index scores were correlated with each pairwise edge for the 13 regions of interest. Dunn and Clark statistics were calculated to compare these correlations between the groups at a significance level of $p < .05$ (i.e. $Z \geq |2|$). Seed region results were visualised with BrainNet viewer (98).

Acknowledgments:

We thank Nadene Dermody and Jody Kamminga for assistance with data collection and scoring. CO is supported by a National Health and Medical Research Council Neil Hamilton Fairley Fellowship GNT1091310. JMS is supported by a National Health and Medical Research Council CJ Martin Fellowship GNT1072403. JAH is supported by the University of Arizona and a grant from the John Templeton Foundation, "Prospective Psychology Stage 2: A Research Competition" to Martin Seligman. The opinions expressed in this publication are those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation. MI is supported by an Australian Research Council Future Fellowship (FT160100096).

This work is supported by funding to ForeFront, a collaborative research group dedicated to the study of frontotemporal dementia and motor neuron disease, from the National Health and Medical Research Council (NHMRC) (APP1037746) and the Australian

Research Council (ARC) Centre of Excellence in Cognition and its Disorders (CE11000102).

Conflict of interest disclosure: The authors report no conflicts of interest.

References

1.   Kucyi A (2017) Just a thought: How mind-wandering is represented in dynamic brain connectivity. *NeuroImage*.

2.   Christoff K, Irving ZC, Fox KC, Spreng RN, & Andrews-Hanna JR (2016) Mind-wandering as spontaneous thought: a dynamic framework. *Nature Reviews Neuroscience* 17(11):718-731.

3.   Christoff K, Gordon AM, Smallwood J, Smith R, & Schooler JW (2009) Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proceedings of the National Academy of Sciences* 106(21):8719-8724.

4.   Buckner RL, Andrews-Hanna JR, & Schacter DL (2008) The brain's default network. *Annals of the New York Academy of Sciences* 1124(1):1-38.

5.   Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, & Buckner RL (2010) Functional-Anatomic Fractionation of the Brain's Default Network. *Neuron* 65(4):550-562.

6.   Fox KC, Spreng RN, Ellamil M, Andrews-Hanna JR, & Christoff K (2015) The wandering brain: Meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *NeuroImage* 111:611-621.

7.   Mason MF*, et al.* (2007) Wandering minds: the default network and stimulus-independent thought. *Science* 315(5810):393-395.

8.   Ellamil M*, et al.* (2016) Dynamics of neural recruitment surrounding the spontaneous arising of thoughts in experienced mindfulness practitioners. *NeuroImage* 136:186-196.

9.   Smallwood J, O'Connor RC, Sudbery MV, & Obonsawin M (2007) Mind-wandering and dysphoria. *Cognition and Emotion* 21(4):816-842.

10.   Perkins AM, Arnone D, Smallwood J, & Mobbs D (2015) Thinking too much: self-generated thought as the engine of neuroticism. *Trends in cognitive sciences* 19(9):492-498.

11.   Smallwood J & Andrews-Hanna J (2013) Not all minds that wander are lost: The importance of a balanced perspective on the mind-wandering state Towards a balanced perspective of the mind-wandering state. *Frontiers in Psychology* 4:441.

12.   Seli P, Smallwood J, Cheyne JA, & Smilek D (2015) On the relation of mind wandering and ADHD symptomatology. *Psychonomic bulletin & review* 22(3):629-636.

13.   Seli P, Risko EF, Purdon C, & Smilek D (2016) Intrusive thoughts: linking spontaneous mind wandering and OCD symptomatology. *Psychological research*:1-7.

14.   Shin D-J, *et al.* (2015) Away from home: the brain of the wandering mind as a model for schizophrenia. *Schizophrenia research* 165(1):83-89.

15.   Pievani M, de Haan W, Wu T, Seeley WW, & Frisoni GB (2011) Functional network disruption in the degenerative dementias. *The Lancet Neurology* 10(9):829-843.

16.   Irish M, Piguet O, & Hodges JR (2012) Self-projection and the default network in frontotemporal dementia. *Nature Reviews Neurology* 8(3):152-161.

17.   Zhou J, *et al.* (2010) Divergent network connectivity changes in behavioural variant frontotemporal dementia and Alzheimer's disease. *Brain* 133(5):1352-1367.

18.   McCormick C, Rosenthal CR, Miller TD, & Maguire E (2017) Mind-wandering in people with hippocampal amnesia. *bioRxiv*.

19.   Bertossi E & Ciaramelli E (2016) Ventromedial prefrontal damage reduces mind-wandering and biases its temporal focus. *Social Cognitive and Affective Neuroscience* 11(11):1783-1791.

20.   Poerio GL, *et al.* (2017) The role of the default mode network in component processes underlying the wandering mind. *Social cognitive and affective neuroscience*.

21.   Smallwood J & Schooler JW (2015) The science of mind wandering: empirically navigating the stream of consciousness. *Annual review of psychology* 66:487-518.

22.   Smallwood J (2013) Distinguishing how from why the mind wanders: a process–occurrence framework for self-generated mental activity. *Psychological bulletin* 139(3):519.

23.   Piolino P, *et al.* (2003) Autobiographical memory and autonoetic consciousness: triple dissociation in neurodegenerative diseases. *Brain* 126(10):2203-2219.

24.   Irish M, *et al.* (2011) Profiles of recent autobiographical memory retrieval in semantic dementia, behavioural-variant frontotemporal dementia, and Alzheimer's disease. *Neuropsychologia* 49(9):2694-2702.

25.   Duval C, *et al.* (2012) What happens to personal identity when semantic knowledge degrades? A study of the self and autobiographical memory in semantic dementia. *Neuropsychologia* 50(2):254-265.

26.   Irish M, *et al.* (2015) Scene construction impairments in Alzheimer's disease–A unique role for the posterior cingulate cortex. *cortex* 73:10-23.

27.   Irish M, Addis DR, Hodges JR, & Piguet O (2012) Considering the role of semantic memory in episodic future thinking: evidence from semantic dementia. *Brain* 135(7):2178-2191.

28.   Stopford CL, Thompson JC, Neary D, Richardson AMT, & Snowden JS (2012) Working memory, attention, and executive function in Alzheimer's disease and frontotemporal dementia. *Cortex* 48(4):429-446.

29.   Possin KL, *et al.* (2013) Dissociable executive functions in behavioral variant frontotemporal and Alzheimer dementias. *Neurology* 80(24):2180-2185.

30.   O'Callaghan C & Irish M (2017) Candidate mechanisms of spontaneous cognition as revealed by dementia syndromes. *The Oxford Handbook of Spontaneous Thought*, eds Fox KCR & Christoff K (Oxford University Press, New York).

31.   Buckner RL*, et al.* (2005) Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *The Journal of Neuroscience* 25(34):7709-7717.

32.   Seeley WW, Crawford RK, Zhou J, Miller BL, & Greicius MD (2009) Neurodegenerative diseases target large-scale human brain networks. *Neuron* 62(1):42.

33.   Damoiseaux JS, Prater KE, Miller BL, & Greicius MD (2012) Functional connectivity tracks clinical deterioration in Alzheimer's disease. *Neurobiology of Aging* 33(4):828. e819-828. e830.

34.   Jones DT*, et al.* (2016) Cascading network failure across the Alzheimer's disease spectrum. *Brain* 139(2):547-562.

35.   Braak H & Braak E (1991) Neuropathological stageing of Alzheimer-related changes. *Acta neuropathologica* 82(4):239-259.

36.   Piguet O, Hornberger M, Mioshi E, & Hodges JR (2011) Behavioural-variant frontotemporal dementia: diagnosis, clinical staging, and management. *The Lancet Neurology* 10(2):162-172.

37.   Rascovsky K*, et al.* (2007) Diagnostic Criteria for the Behavioral Variant of Frontotemporal Dementia (bvFTD): Current Limitations and Future Directions. *Alzheimer Disease & Associated Disorders* 21(4):S14-S18.

38.   Broe M*, et al.* (2003) Staging disease severity in pathologically confirmed cases of frontotemporal dementia. *Neurology* 60(6):1005-1011.

39.   Seeley WW*, et al.* (2008) Frontal paralimbic network atrophy in very mild behavioural variant frontotemporal dementia. *Archives of Neurology* 65:2249-2255.

40.   Landin-Romero R*, et al.* (2017) Disease-specific patterns of cortical and subcortical degeneration in a longitudinal study of Alzheimer's disease and behavioural-variant frontotemporal dementia. *Neuroimage* 151:72-80.

41.   Stawarczyk D, Majerus S, Maquet P, & D'Argembeau A (2011) Neural correlates of ongoing conscious experience: both task-unrelatedness and stimulus-independence are related to default network activity. *PLoS ONE* 6(2):e16997.

42.   Schooler JW, Reichle ED, & Halpern DV (2004) Zoning Out while Reading: Evidence for Dissociations between Experience and Metaconsciousness. *Thinking and seeing: Visual metacognition in adults and children*, ed Levin DT (MIT Press, Cambridge, Massachusetts), pp 203-226.

43.   Maillet D & Schacter DL (2016) When the mind wanders: Distinguishing stimulus-dependent from stimulus-independent thoughts during incidental encoding in young and older adults. *Psychology and aging* 31(4):370.

44.   O'Callaghan C, Shine JM, Lewis SJG, Andrews-Hanna JR, & Irish M (2015) Shaped by our thoughts – A new task to assess spontaneous cognition and its associated neural correlates in the default network. *Brain and Cognition* 93(0):1-10.

45.   Stawarczyk D, Majerus S, Maj M, Van der Linden M, & D'Argembeau A (2011) Mind-wandering: Phenomenology and function as assessed with a novel experience sampling method. *Acta Psychologica* 136(3):370-381.

46.   Irish M & Piolino P (2016) Impaired capacity for prospection in the dementias– Theoretical and clinical implications. *British Journal of Clinical Psychology* 55(1):49-68.

47.   Bertoux M*, et al.* (2016) Social cognition deficits: the key to discriminate behavioral variant frontotemporal dementia from Alzheimer's disease regardless of amnesia? *Journal of Alzheimer's Disease* 49(4):1065-1074.

48.   Hornberger M, Piguet O, Graham A, J., Nestor PJ, & Hodges JR (2010) How preserved is episodic memory in behavioral variant frontotemporal dementia? *Neurology* 74:472-479.

49.   Irish M, Piguet O, Hodges JR, & Hornberger M (2014) Common and unique gray matter correlates of episodic memory dysfunction in frontotemporal dementia and alzheimer's disease. *Human Brain Mapping* 35(4):1422-1435.

50.   Frisch S*, et al.* (2013) Dissociating Memory Networks in Early Alzheimer's Disease and Frontotemporal Lobar Degeneration - A Combined Study of Hypometabolism and Atrophy. *PLOS ONE* 8(2):e55251.

51.   Kamminga J, O'Callaghan C, Hodges JR, & Irish M (2014) Differential Prospective Memory Profiles in Frontotemporal Dementia Syndromes. *Journal of Alzheimer's Disease* 38(3):669-679.

52.   Dermody N, Hornberger M, Piguet O, Hodges JR, & Irish M (2016) Prospective memory impairments in Alzheimer's disease and behavioral variant frontotemporal dementia: Clinical and neural correlates. *Journal of Alzheimer's Disease* 50(2):425-441.

53.   Addis DR, Sacchetti DC, Ally BA, Budson AE, & Schacter DL (2009) Episodic simulation of future events is impaired in mild Alzheimer's disease. *Neuropsychologia* 47(12):2660-2671.

54.   Irish M, Hodges JR, & Piguet O (2013) Episodic future thinking is impaired in the behavioural variant of frontotemporal dementia. *Cortex* 49(9):2377-2388.

55.   Nyatsanza S*, et al.* (2003) A study of stereotypic behaviours in Alzheimer's disease and frontal and temporal variant frontotemporal dementia. *Journal of Neurology, Neurosurgery & Psychiatry* 74(10):1398-1402.

56.   Bozeat S, Gregory CA, Lambon Ralph MA, & Hodges JR (2000) Which neuropsychiatric and behavioural features distinguish frontal and temporal variants of frontotemporal dementia from Alzheimer's disease? *Journal of Neurology, Neurosurgery & Psychiatry* 69:178-196.

57.   de Souza LC*, et al.* (2010) Poor creativity in frontotemporal dementia: A window into the neural bases of the creative mind. *Neuropsychologia* 48(13):3733-3742.

58.   Eslinger PJ*, et al.* (2007) Oops! Resolving social dilemmas in frontotemporal dementia. *Journal of Neurology, Neurosurgery & Psychiatry* 78(5):457-460.

59.   Spreng RN, Stevens WD, Chamberlain JP, Gilmore AW, & Schacter DL (2010) Default network activity, coupled with the frontoparietal control network, supports goal-directed cognition. *Neuroimage* 53(1):303-317.

60.   Gerlach KD, Spreng RN, Gilmore AW, & Schacter DL (2011) Solving future problems: default network and executive activity associated with goal-directed mental simulations. *Neuroimage* 55(4):1816-1824.

61.   Beaty RE, Benedek M, Silvia PJ, & Schacter DL (2015) Creative Cognition and Brain Network Dynamics. *Trends in Cognitive Sciences*.

62.   Fornito A, Harrison BJ, Zalesky A, & Simons JS (2012) Competitive and cooperative dynamics of large-scale brain functional networks supporting recollection. *Proceedings of the National Academy of Sciences* 109(31):12788-12793.

63.   Golchert J*, et al.* (2017) Individual variation in intentionality in the mind-wandering state is reflected in the integration of the default-mode, fronto-parietal, and limbic networks. *NeuroImage* 146:226-235.

64.   Dhanjal NS & Wise RJ (2014) Frontoparietal cognitive control of verbal memory recall in Alzheimer's disease. *Annals of neurology* 76(2):241-251.

65.   Grady CL*, et al.* (2003) Evidence from functional neuroimaging of a compensatory prefrontal network in Alzheimer's disease. *The Journal of Neuroscience* 23(3):986-993.

66.   Rombouts SARB, Barkhof F, Goekoop R, Stam CJ, & Scheltens P (2005) Altered resting state networks in mild cognitive impairment and mild Alzheimer's disease: An fMRI study. *Human Brain Mapping* 26(4):231-239.

67.   Gelbard-Sagiv H, Mukamel R, Harel M, Malach R, & Fried I (2008) Internally generated reactivation of single neurons in human hippocampus during free recall. *Science* 322(5898):96-101.

68.   Burke JF*, et al.* (2014) Theta and high-frequency activity mark spontaneous recall of episodic memories. *Journal of Neuroscience* 34(34):11355-11365.

69.   Dragoi G & Tonegawa S (2011) Preplay of future place cell sequences by hippocampal cellular assemblies. *Nature* 469(7330):397-401.

70.   Pfeiffer BE & Foster DJ (2013) Hippocampal place-cell sequences depict future paths to remembered goals. *Nature* 497(7447):74-79.

71.   Dupret D, O'neill J, Pleydell-Bouverie B, & Csicsvari J (2010) The reorganization and reactivation of hippocampal maps predict spatial memory performance. *Nature neuroscience* 13(8):995-1002.

72.	Karlsson MP & Frank LM (2009) Awake replay of remote experiences in the hippocampus. *Nature neuroscience* 12(7):913-918.

73.	Foster DJ & Wilson MA (2006) Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature* 440(7084):680-683.

74.	Diba K & Buzsáki G (2007) Forward and reverse hippocampal place-cell sequences during ripples. *Nature neuroscience* 10(10):1241-1242.

75.	Kaplan R*, et al.* (2016) Hippocampal sharp-wave ripples influence selective activation of the default mode network. *Current Biology* 26(5):686-691.

76.	Zhou J & Seeley WW (2014) Network dysfunction in Alzheimer's disease and frontotemporal dementia: implications for psychiatry. *Biological psychiatry* 75(7):565-573.

77.	Chiong W*, et al.* (2013) The salience network causally influences default mode network activity during moral reasoning. *Brain*.

78.	Menon V & Uddin LQ (2010) Saliency, switching, attention and control: a network model of insula function. *Brain Structure and Function* 214(5-6):655-667.

79.	Crossley NA*, et al.* (2014) The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain* 137(8):2382-2395.

80.	O'Callaghan C, Bertoux M, & Hornberger M (2014) Beyond and below the cortex: the contribution of striatal dysfunction to cognition and behaviour in neurodegeneration. *Journal of Neurology, Neurosurgery and Psychiatry* 85(4):371-378.

81.	Bell PT & Shine JM (2016) Subcortical contributions to large-scale network communication. *Neuroscience & Biobehavioral Reviews* 71:313-322.

82.	Choi EY, Yeo BT, & Buckner RL (2012) The organization of the human striatum estimated by intrinsic functional connectivity. *Journal of neurophysiology* 108(8):2242-2263.

83.	Jarbo K & Verstynen TD (2015) Converging structural and functional connectivity of orbitofrontal, dorsolateral prefrontal, and posterior parietal cortex in the human striatum. *Journal of Neuroscience* 35(9):3865-3878.

84.	Zabelina DL & Andrews-Hanna JR (2016) Dynamic network interactions supporting internally-oriented cognition. *Current Opinion in Neurobiology* 40:86-93.

85.	Rascovsky K*, et al.* (2011) Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* 134(9):2456-2477.

86.	McKhann GM*, et al.* (2011) The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 7(3):263-269.

87.	Mioshi E, Dawson K, Mitchell J, Arnold R, & Hodges JR (2006) The Addenbrooke's Cognitive Examination Revised (ACE R): a brief cognitive test battery for dementia screening. *International Journal of Geriatric Psychiatry* 21(11):1078-1085.

88.   Wear HJ*, et al.* (2008) The Cambridge Behavioral Inventory revised. *Dementia and Neuropsychologia* 2:102-107.

89.   Smith SM (2002) Fast robust automated brain extraction. *Human brain mapping* 17(3):143-155.

90.   Zhang Y, Brady M, & Smith S (2001) Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* 20(1):45-57.

91.   Andersson JLR, Jenkinson M, & Smith S (2007) Non-linear optimisation. (FMRIB technical report TR07JA1 from http://www.fmrib.ox.ac.uk/analysis/techrep).

92.   Smith SM & Nichols TE (2009) Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* 44(1):83-98.

93.   Nichols TE & Holmes AP (2002) Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Human brain mapping* 15(1):1-25.

94.   Jenkinson M, Bannister P, Brady M, & Smith S (2002) Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 17(2):825-841.

95.   Behzadi Y, Restom K, Liau J, & Liu TT (2007) A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* 37(1):90-101.

96.   Dosenbach NU, Fair DA, Cohen AL, Schlaggar BL, & Petersen SE (2008) A dual-networks architecture of top-down control. *Trends in cognitive sciences* 12(3):99-105.

97.   Damoiseaux JS, Viviano RP, Yuan P, & Raz N (2016) Differential effect of age on posterior and anterior hippocampal functional connectivity. *NeuroImage* 133:468-476.

98.   Xia M, Wang J, & He Y (2013) BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one* 8(7):e68910.