BldC delays entry into development to produce a sustained period of vegetative growth in *Streptomyces venezuelae*

Matthew J. Bush[1],*, Mahmoud Al-Bassam[1],#, Govind Chandra[1], Kim C. Findlay[2] and Mark J. Buttner[1]

[1]*Department of Molecular Microbiology*, and [2]*Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK*.

* To whom correspondence should be addressed. Tel: 44 (0) 1603 450757; Fax: 44 (0) 1603 450778; Email:matt.bush@jic.ac.uk

Current address: Department of Paediatrics, University of California, San Diego, La Jolla, CA 92093, USA

Keywords: Morphological differentiation; sporulation; cell division; transcriptional regulation.

Abstract

Streptomycetes are filamentous bacteria that differentiate by producing spore-bearing reproductive structures called aerial hyphae. The transition from vegetative to reproductive growth is controlled by the *bld* (bald) loci, and mutations in *bld* genes prevent the formation of aerial hyphae, either by blocking entry into development (mutations in activators) or by inducing precocious sporulation in the vegetative mycelium (mutations in repressors). One of the *bld* genes, *bldC*, encodes a 68-residue protein with a winged Helix-Turn-Helix (wHTH) DNA-binding motif. Here we exploit the benefits of the new model species, *Streptomyces venezuelae*, which sporulates in liquid culture, to study the biological role of BldC. Using electron microscopy and time-lapse imaging, we show that *bldC* mutants are bald because they initiate development prematurely, bypassing the formation of aerial hyphae. This correlates with premature expression of BldC target genes, showing that BldC acts as a repressor to sustain vegetative growth and delay entry into development.

Introduction

The complex *Streptomyces* life cycle involves two distinct filamentous cell forms: the growing or vegetative hyphae and the reproductive or aerial hyphae, which differentiate into long chains of spores (1-5). Genetic studies identified the regulatory loci that control entry into development, which are called *bld* (bald) genes because null mutations in these loci prevent the formation of aerial hyphae. However, baldness can arise for two different reasons. The larger class of *bld* mutants, which define positive regulators, fail to initiate development, forming colonies of undifferentiated vegetative mycelium. In contrast, a smaller but growing class of *bld* mutants, which define negative regulators, enter development prematurely, inducing sporulation in the vegetative mycelium and bypassing the formation of aerial hyphae. Thus, macroscopically these two classes of mutants look similar, forming smooth colonies that lack the 'hairy' appearance of the wild type, but microscopically it is apparent that they arise for diametrically opposed reasons (4, 6-8).

bldC is known to encode a short, 68 residue protein related to the DNA-binding domain of MerR-family proteins (9), but there has been less insight into its biological role and impact on *Streptomyces* development. In part, this is because previous studies have focussed on the classical model species, *S. coelicolor*, which sporulates only on solid medium. Here we exploit the benefits of the new model species, *Streptomyces venezuelae*, which sporulates in liquid culture (10), to study the biological role of BldC. We show that *bldC* mutants are bald because they enter development prematurely, bypassing the formation of aerial hyphae. This correlates with premature expression of BldC target genes, showing that BldC acts as a repressor to sustain vegetative growth and delay entry into development.

Results

Deletion of *bldC* causes premature initiation of development

We constructed an *S. venezuelae bldC* mutant by replacing the *bldC* coding region with an

apramycin resistance (*apr*) cassette. The resulting mutant was bald, unable to produce the

reproductive aerial hyphae that give mature wild-type *Streptomyces* colonies their

characteristic fuzzy appearance (Fig 1.). However, scanning electron microscopy (SEM) of

mature colonies of the *bldC* mutant showed that most of the biomass consisted of spores,

rather than undifferentiated vegetative hyphae (Fig. 2). Comparison of the growth of the wild

type and the *bldC* mutant on plates over time showed that after 1 day they looked similar

(vegetative growth only) but after 2 days the wild type had produced aerial hyphae while the

bldC mutant was still restricted to vegetative growth. After 3 days, the aerial hyphae of the

wild-type had differentiated into spores, and most of the biomass of the *bldC* mutant had also

differentiated into spores, bypassing aerial mycelium formation. The *bldC* mutant also

seemed to produce higher levels of extracellular matrix than the wild type (Fig. 2). The *bldC*

mutant phenotype was fully complemented by introducing a single copy of the *bldC* gene

under the control of its native promoter, expressed *in trans* from the ΦBT1 integration site

(Figs. 1 and 2).

Using an established microfluidic system and methodology (10), we conducted fluorescence

time-lapse microscopy to further study the developmental defects associated with deletion of

bldC. As in previous studies (6,10), we introduced an FtsZ-YPet translational fusion into the

wild type, mutant and complemented mutant strains, enabling us to monitor each of the two

distinct modes of cell division that occur in *Streptomyces*. In Fig. 3, the scattered single Z-

rings mark the position of vegetative cross-walls, which do not constrict or give rise to cell-

cell separation, but simply divide the vegetative hyphae into long, box-like compartments

(e.g. Figs. 3A + C, panel 2). In contrast, during reproductive growth, long ladders of regularly

spaced Z-rings are synchronously deposited along sporogenic hyphae. These Z-rings mark

the sites of sporulation septa, which do constrict, ultimately leading to the formation of chains

of spores (e.g. Figs. 3A + C, panels 3 +4). Time-lapse imaging of strains harbouring the FtsZ-

YPet fusion showed that the duration of vegetative growth was shorter in the *bldC* mutant

compared to the wild type and the complemented mutant (Fig.3 and Movies S1 A/B, S2 A/B

and S3 A/B). Noticeably, following germination, hyphal outgrowth in the *bldC* mutant was

associated with an immediate increase in FtsZ-YPet expression, leading to the precocious

formation of ladders of Z-rings (Fig. 3B and Movie S2A/B). However, although ladders of Z-

rings were observed as early as 4 hours in the *bldC* mutant, mature spores were not observed

in the corresponding DIC images until 21 hours, the same time mature spores were also seen

in the wild type (Figs. 3A and B). Wild-type patterns of FtsZ expression and sporulation were

restored in the complemented mutant (Fig. 3C and Movie S3A/B). From these data, we

concluded that the overall role of BldC is to sustain vegetative growth and delay entry into

development.

BldC levels are highest early in development

Using an anti-BldC polyclonal antibody, we monitored BldC levels in *S. venezuelae* during

sporulation in liquid culture. Western blotting showed that BldC is abundant throughout the

life cycle, but that BldC levels are highest early on, during vegetative growth (Figure 4).

BldC represses transcription of its target genes

ChIP-chip studies in *S. coelicolor* showed that BldC binds upstream of ~280 genes (11).

These targets include many genes encoding key transcriptional regulators of the *Streptomyces*

developmental cascade (e.g. *bldM*, *whiB*, *whiD*, *whiH*, *whiI*, *sigF* and *bldC* itself), as well as

others encoding proteins involved in chromosome condensation and segregation during

sporulation (e.g. *hupS*, *smeA-sffA*). Schumacher et al. (11) characterised the interaction of *S.*

coelicolor BldC with the promoters of two of its targets, *whiI* and the *smeA-ssfA* operon. *whiI*

encodes an orphan response-regulator that is essential for the later stages of sporulation, when

it forms a functional heterodimer with a second orphan response-regulator, BldM, enabling

WhiI to bind to DNA and regulate the expression of ~40 late sporulation genes (12). The

smeA-sffA operon encodes a small membrane protein (SmeA) that recruits a DNA translocase

(SffA) to sporulation septa (13). Deletion of *smeA-sffA* results in a defect in spore

chromosome segregation and has pleiotropic effects on spore maturation (13).

BldC binds directly to the *whiI* and *smeA* promoters in *S. coelicolor* (11) and ChIP-chip

analysis confirmed that they are also BldC targets in *S. venezuelae* (Table S1, Fig. 5). To

assess the regulatory influence of BldC on the *whiI* and *smeA* promoters, we performed qRT-

PCR using RNA prepared from both wild-type *S. venezuelae* and the *bldC* mutant, examining

the 8 and 10 hour time points when BldC is most abundant in wild-type cells (Fig. 5). In the

wild type, expression of both BldC targets comes on in the 10-h time point. In contrast, in the

bldC mutant, expression of both *whiI* and *smeA* is on in the 8- and 10-h time points.

Furthermore, at 10-h, expression of *whiI* is 15-fold higher in the *bldC* mutant compared to the

wild type and expression of *smeA* is 40-fold higher. Similarly, we conducted qRT-PCR to

assess the regulatory impact of BldC on the expression of three further key BldC targets - the

sigF, *whiD* and *hupS* genes. As we observed for *whiI* and *smeA*, expression of the *sigF*, *whiD*

and *hupS* genes comes on early in the *bldC* mutant (Fig. 6). We conclude that BldC functions

to repress the transcription of these developmental target genes during vegetative growth,

consistent with the premature initiation of development seen in a *bldC* mutant.

Discussion

Canonical *bld* mutations block entry into development and so the resulting colonies do not form aerial hyphae and spores. Such mutations typically define positive regulators such as the response regulator BldM (12) or the sigma factor BldN (14). In contrast, we have shown that *S. venezuelae bldC* mutants are bald because they enter development prematurely, bypassing the formation of aerial hyphae, and that this correlates with premature expression of BldC target genes like *whiI* and *smeA*. Thus, BldC functions as a repressor to sustain vegetative growth and delay entry into development. As such, BldC joins a growing class of Bld regulators known to function as a developmental "brake".

BldD was the first Bld regulator of this alternative class to be clearly recognized. BldD sits at the top of the developmental cascade and represses a large regulon of ~170 sporulation genes during vegetative growth. BldD activity is controlled by the second messenger c-di-GMP, which mediates dimerization of two BldD protomers to generate a functional repressor. In this way, c-di-GMP signals through BldD to repress expression of the BldD regulon, extending vegetative growth and inhibiting entry into development (4, 8, 15). Because it is a BldD-(c-di-GMP) complex that represses the BldD regulon and not BldD alone, engineering the degradation of c-di-GMP *in vivo* also causes a precocious hypersporulation phenotype like that of a *bldD* null mutant (8).

More recently, *bldO* was identified as a second member of this emerging class of *bld* mutant (6-7). In contrast to BldD and BldC, which both control large regulons, BldO functions to repress a single developmental gene, *whiB*. The precocious hypersporulation phenotype of the *bldO* mutant arises from premature expression of *whiB*, and in line with this, constitutive expression of *whiB* alone is sufficient to induce precocious hypersporulation in wild-type *S. venezuelae* (6). WhiA and WhiB act together to co-control the same set of promoters to initiate developmental cell division in *Streptomyces* (16-17). WhiA is constitutively present

throughout the life cycle, but it only binds to its target promoters at the onset of sporulation

(16). This is because WhiA and WhiB function cooperatively and *in vivo* DNA binding by

WhiA depends on WhiB, and *vice versa* (17). As a consequence, the regulation of *whiB*

expression is key in controlling the switch between hyphal growth and sporulation. This

critical role for WhiB is reflected in the extensive developmental regulation to which *whiB*

transcription is subject, being directly repressed by BldC (11), BldD (18) and BldO (6), and

directly activated by BldM (12).

Materials and Methods

Construction and complementation of an *S. venezuelae bldC* null mutant. Using

'Redirect' PCR targeting (19-20), *bldC* mutants were generated in which the coding region

was replaced with a single apramycin resistance (*apr*) cassette. A cosmid library that covers >

98% of the *S. venezuelae* genome (M.J. Bibb and M.J. Buttner, unpublished) is fully

documented at http://strepdb.streptomyces.org.uk/. Cosmid 4O24 was introduced into *E. coli*

BW25113 containing pIJ790 and the *bldC* gene (*sven3846*) was replaced with the *apr-oriT*

cassette amplified from pIJ773 using the primer pairs bldCdis_F and bldCdis_R. The

resulting disrupted cosmids were confirmed by restriction digestion and by PCR analysis

using the flanking primers bldCcon_F and bldCcon_R, and introduced into *S. venezuelae* by

conjugation (Keiser *et al*., 2000). Null mutant derivatives, generated by double crossing over,

were identified by their apramycin-resistant, kanamycin-sensitive and morphological

phenotypes, and their chromosomal structures were confirmed by PCR analysis using the

flanking primers bldCcon_F and bldCcon_R. A representative *bldC* null mutant was

designated SV25. For complementation, *bldC* was amplified with the primers bldCcomp_F

and bldCcomp_R, generating an 846bp fragment carrying the coding sequence and the *bldC*

promoter, and cloned into HindIII-KpnI/Asp718 cut pIJ10770 to create pIJ10618. The

plasmid was introduced into the *bldC* mutant by conjugation and fully complemented all

aspects of the mutant phenotype.

Time-lapse imaging of *S. venezuelae*. Fluorescent time-lapse imaging was conducted

essentially as described previously (6,11). Before imaging, fresh *S. venezuelae* spores for

each of the strains imaged were first prepared by inoculating 30 ml cultures of MYM with 10

µl of the appropriate spore stock or 20 µl of the appropriate mycelial culture. Cells were

cultured at 30 ºC and 250 rpm until fully differentiated (16-24 h for hypersporulating strains,

otherwise 36-40 h). 1 ml of each culture was spun briefly to pellet mycelium, the supernatant

spores were diluted 1:50 in fresh MYM, and 50 µl was transferred to the cell loading well of

a prepared B04A microfluidic plate (Merck-Millipore). The remaining culture was filter-

sterilised to obtain spent MYM that was free of spores and mycelial fragments. The ONIX

manifold was then sealed to the B04A plate before transferring to the environmental

chamber, pre-incubated at 30 ºC. Spores were loaded onto the B04A plate, at 4 psi for 15

seconds using the ONIX microfluidic perfusion system. Fresh MYM medium was set to flow

at 2 psi during the first 3 hours during germination, before the 2-psi flow of spent MYM

medium for the remainder of the experiment. The system was left to equilibrate for 1 h prior

to imaging.

Imaging was conducted using a Zeiss Axio Observer.Z1 widefield microscope equipped with

a sCMOS camera (Hamamatsu Orca FLASH 4), a metal-halide lamp (HXP 120V), a

hardware autofocus (Definitive Focus), a 96-well stage insert, an environmental chamber, a

100x 1.46 NA Oil DIC objective and the Zeiss 46 HE shift free (excitation500/25 nm,

emission 535/30 nm) filter set. DIC images were captured with a 150 ms exposure time, YFP

images were captured with a 100 ms exposure time. Images were taken every 30 min. In all

experiments, multiple x/y positions were imaged for each strain and in each experiment.

Representative images were transferred to the Fiji software package (http://fiji.sc/Fiji),

manipulated and converted into the movie files presented here, as described previously

(Schlimpert *et al*., 2016).

Chromatin immunoprecipitation-microarray (ChIP-chip) analysis. To carry out the

ChIP-chip experiments, cultures of *S. venezuelae* and the congenic *bldC* null mutant strain

SV25 were grown for 12 h in MYM liquid medium. Formaldehyde was added to cultures at a

final concentration of 1% (v/v) and incubation was continued for 30 min. Glycine was then

added to a final concentration of 125 mM to stop the cross-linking. Cultures were left at room

temperature (RT) for 5 min before the mycelium was harvested and washed twice in PBS

buffer pH 7.4. Each mycelial pellet was resuspended in 0.5 ml lysis buffer (10 mM Tris HCl

pH 8.0, 50 mM NaCl) containing 15 mg/ml lysozyme and protease inhibitor (Roche Applied

Science) and incubated at 25 ℃ for 1 h. An equal volume of IP buffer (100 mM Tris HCl pH

8, 250 mM NaCl, 0.5% Triton-X-100, 0.1% SDS) containing protease inhibitor was added

and samples were chilled on ice. Samples were sonicated for 7 cycles of 15 s each at 10

microns to shear the chromosomal DNA into fragments ranging from 300-1000 bp in size.

Samples were centrifuged twice at 13,000 rpm at 4 ℃ for 15 min to clear the cell extract,

after which 10 µl of cell extract was set aside for total DNA extraction. The remainder was

incubated with 10% (v/v) protein A-sepharose (Sigma) for 1 h on a rotating wheel to remove

non-specifically binding proteins. Samples were then centrifuged for 15 min at 4℃ and

13,000 rpm to remove the beads. Supernatants were incubated with 10% (v/v) anti-BldC

antibody (9) overnight at 4 ℃ with rotation. Subsequently, 10% (v/v) protein A-sepharose

was added to precipitate BldC and incubation was continued for 4 h. Samples were

centrifuged at 3500 rpm for 5 min and the pellets were washed four times with 0.5x IP buffer.

Each pellet was incubated overnight at 65 ºC in 150 µl IP elution buffer (50 mM Tris HCl pH

7.6, 10 mM EDTA, 1% SDS) to reverse cross-links, and 10 µl of the total cell extract control

was treated in the same way. Samples were centrifuged at 13,000 rpm for 5 min to remove

the beads. Each pellet was re-extracted with 50 µl TE buffer (10 mM Tris HCl pH 7.4, 1 mM

EDTA) and the supernatant incubated with 0.2 mg/ml Proteinase K (Roche) for 2 h at 55ºC.

The resulting samples were extracted with phenol-chloroform and further purified using

QiaQuick columns, eluting in 50 µl EB buffer (Qiagen). DNA labelling, hybridization to

DNA microarrays and data processing were carried out as described previously (10). All data

is deposited at ArrayExpress (Accession: E-MTAB-6129).

qRT-PCR. Mycelial pellets from MYM cultures were washed in PBS and resuspended in

900 µl lysis solution (400 µl phenol [pH4.3], 100 µl chlorophorm:isoamyl alcohol (24 : 1)

and 400 µl RLT buffer [Qiagen]) with lysing matrix B (MP Biomedicals) and homogenised

using a FastPrep FP120 Cell Disruptor (Thermo Savant). Two pulses of 30 s of intensity 6.0

were applied with cooling down for 1 min on ice between pulses. Supernatants were

centrifuged for 15 min, full-speed on a bench-top centrifuge at 4°C and then treated according

to the instructions given in the RNEasy Kit (Qiagen). The RNA samples were treated with

on-column DNase I (Qiagen), followed by an additional DNase I treatment (Turbo DNA-free,

Ambion) until they were free of DNA contamination (determined by PCR amplification of

hrdB). RNA was quantified and equal amounts (350 ng) of total RNA from each sample was

converted to cDNA using SuperScript II reverse transcriptase and random primers

(Invitrogen). cDNA was then used as template in qRT-PCR performed using the SensiFAST

SYBR No-ROX kit (Bioline). Three technical replicates were used for each gene. Specific

qPCR primers (Table S1, final concentration of 250 nM) were used to amplify the target

genes *whiI, smeA, whiD, sigF* and *hups,* as well as the *hrdB* reference gene. To normalize for

differing primer efficiency, a standard curve was constructed using chromosomal DNA.

Melting curve analysis was used to confirm the production of a specific single product from

each primer pair. qRT-PCR was performed using a CFX96 Touch instrument using hardshell

white PCR plates (BioRad), sealed with thermostable film covers (Thermo). PCR products

were detected with SYBR green fluorescent dye and amplified according to the following

protocol: 95°C, 3 min, then 45 cycles at 95°C 5 sec, 62°C 10 sec and 72°C 7 sec. Melting

curves were generated at 65 to 95°C with 0.5°C increments. The BioRad CFX manager

software was used to calculate starting quantity (SQ) values for *smeA* and *whiI* at each time

point. These values were divided by the mean SQ value derived from the *hrdB* reference at

the corresponding time points, generating a value for relative expression. The resulting values

were normalised against the mean relative expression of the wild type at 8 hours, which was

set to 1.

Western Blotting. Samples of frozen mycelium, originating from 2 ml liquid MYM samples,

were resuspended in 0.4 ml ice-cold sonication buffer [20 mM Tris pH 8.0, 5 mM EDTA, 1 x

EDTA-free protease inhibitors (Roche)] and sonicated (5x 15 sec on/15 sec off) at 4.5 micron

amplitude. Lysates were then centrifuged at 16,000 xg for 15 min at 4°C to remove cell

debris. Total protein concentration was determined using the Bradford assay (Biorad). 1 µg of

total protein from each time point was loaded in triplicate into a microplate (proteinsimple

#043-165) and anti-bldC antibody (9) diluted 1:200. BldC levels, originating from the wild

type strain and the *bldC* mutant negative control were then assayed using the automated

Western blotting machine Wes (ProteinSimple, San Jose, CA), according to the

manufacturer's guidelines.

Scanning electron microscopy. Colonies were mounted on the surface of an aluminum stub

with optimal cutting temperature compound (Agar Scientific Ltd, Essex, UK), plunged into

liquid nitrogen slush at approximately -210°C to cryopreserve the material, and transferred to

the cryostage of an Alto 2500 cryotransfer system (Gatan, Oxford, England) attached to a FEI

NanoSEM 450 field emission gun scanning electron microscope (FEI Ltd, Eindhoven, The

Netherlands). The surface frost was sublimated at -95°C for 3½ mins before the sample was

sputter coated with platinum for 2 min at 10 mA at below -110°C. Finally, the sample was

moved onto the cryostage in the main chamber of the microscope, held at approximately -

130°C, and viewed at 3 kV.

Acknowledgements

We are grateful to Oxford Gene Technology for expert handling of the ChIP samples. This

work was funded by BBSRC grant BB/H006125/1 (to M.J.B.) and by BBSRC Institute

Strategic Programme Grant BB/J004561/1 to the John Innes Centre.

References

1. Flärdh K, Buttner MJ. 2009. *Streptomyces* morphogenetics: dissecting differentiation in a

filamentous bacterium. Nat Rev Microbiol 7: 36-49.

2. McCormick JR, Flärdh K. 2012. Signals and regulators that govern *Streptomyces*

development. FEMS Microbiol Rev 36: 206-231.

3. Jakimowicz D, and van Wezel GP. 2012. Cell division and DNA segregation in

Streptomyces: how to build a septum in the middle of nowhere? Mol Microbiol 85: 393-404.

4. Bush MJ, Tschowri N, Schlimpert S, Flärdh K, Buttner MJ. 2015. c-di-GMP signaling and

the regulation of developmental transitions in *Streptomyces*. Nat Rev Microbiol 13: 749-760.

5. Chater KF. 2016. Recent advances in understanding *Streptomyces*. F1000Res 5: 2795.

6. Bush MJ, Chandra G, Findlay KC, Buttner MJ. 2017. Multi-layered inhibition of

Streptomyces development: BldO is a dedicated repressor of whiB. Mol. Microbiol. 104: 700-

711.

7. Flärdh K, McCormick JR. 2017. The *Streptomyces* O-B one connection: a force within

layered repression of a key developmental decision. Mol. Microbiol 104: 695-699.

8. Tschowri N, Schumacher MA, Schlimpert S, Chinnam NB, Findlay KC, Brennan RG,

Buttner MJ. 2014. Tetrameric c-di-GMP mediates effective transcription factor dimerization

to control *Streptomyces* development. Cell 158: 1136-1147.

9. Hunt AC, Servin-Gonzalez L, Kelemen GH, Buttner MJ. 2005. The bldC developmental

locus of *Streptomyces coelicolor* encodes a member of a family of small DNA-binding

proteins related to the DNA-binding domains of the MerR family. J Bacteriol 187: 716-728.

10. Schlimpert S, Flärdh K, Buttner, MJ. 2016. Fluorescence time-lapse imaging of the

complete *Streptomyces* life cycle using a microfluidic device. J Vis Exp 108: e53863.

11. Schumacher MA, den Hengst CD, Bush MJ, Le TB, Tran NT, Chandra G, Zeng W,

Travis B, Brennan RG, Buttner MJ. 2017. Manuscript in preparation.

12. Al-Bassam MM, Bibb MJ, Bush MJ, Chandra G, Buttner MJ. 2014. Response regulator

heterodimer formation controls a key stage in *Streptomyces* development. PLoS Genet 10:

e1004554.

13. Ausmees N, Wahlstedt H, Bagchi S, Elliot MA, Buttner MJ, Flärdh K. 2007. SmeA, a

small membrane protein with multiple functions in *Streptomyces* sporulation including

targeting of a SpoIIIE/FtsK-like protein to cell division septa. Mol Microbiol 65:1458-1473.

14. Bibb MJ, Domonkos A, Chandra G, Buttner MJ. 2012. Expression of the chaplin and

rodlin hydrophobic sheath proteins in *Streptomyces venezuelae* is controlled by σ^{BldN} and a

cognate anti-sigma factor, RsbN. Mol. Microbiol. 84:1033–1049.

15. Schumacher MA, Zeng W, Findlay KC, Buttner MJ, Brennan RG, Tschowri N. 2017. The

Streptomyces master regulator BldD binds c-di-GMP sequentially to create a functional

$BldD_2$-(c-di-GMP)$_4$ complex. Nucleic Acids Res 45: 6923-6933.

16. Bush MJ, Bibb MJ, Chandra G, Findlay KC, Buttner MJ. 2013. Genes required for aerial

growth, cell division, and chromosome segregation are targets of WhiA before sporulation in

Streptomyces venezuelae. mBio 4: e00684-13.

17. Bush MJ, Chandra G, Bibb MJ, Findlay KC, Buttner MJ. 2016. Genome-wide chromatin

immunoprecipitation sequencing analysis shows that WhiB is a transcription factor that co-

controls its regulon with WhiA to initiate developmental cell division in *Streptomyces*. mBio

7: e00523-16.

18. den Hengst CD, Tran NT, Bibb MJ, Chandra C, Leskiw BK, Buttner MJ. 2010. Genes

essential for morphological development and antibiotic production in *Streptomyces coelicolor*

are targets of BldD during vegetative growth. Mol Microbiol 78: 361-379.

19. Gust B, Challis GL, Fowler K, Kieser T, Chater KF. 2003 PCR-targeted *Streptomyces*

gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene

soil odor geosmin. Proc Natl Acad Sci USA 100**:** 1541-1546.

20. Gust B, Chandra G, Jakimowicz D, Yuqing T, Bruton C, Chater KF. 2004. Lambda red-

mediated genetic manipulation of antibiotic-producing *Streptomyces*. Adv Appl Microbiol

54: 107-128.

Figure Legends

FIG 1. BldC is required for the formation of aerial mycelium.

The phenotypes of wild-type *S.venezuelae*, the *bldC* mutant, the *bldC* mutant carrying the empty vector, and the complemented *bldC* mutant, photographed after four days of growth on MYM solid medium.

FIG 2. Deletion of *bldC* causes premature initiation of development on solid medium.

Scanning electron micrographs showing the phenotype of the *bldC* mutant, compared to wild-type after 1 days, 2 days and 3 days of growth on MYM solid medium. The phenotype of the complemented *bldC* mutant is also shown after 3 days of growth on MYM solid medium.

FIG 3. Deletion of *bldC* causes premature initiation of development in liquid medium.

Time-lapse images (4, 7, 12 and 21 h post-inoculation) of (A) wild-type *S. venezuelae*, (B) the *bldC* mutant and (C) the complemented *bldC* mutant, grown in liquid MYM medium in the microfluidic system. All three strains carry the same FtsZ-YPet translational fusion expressed from the native ftsZ promoter, and both the DIC (upper) and fluorescence (lower) images are shown. Scale Bar = 10µm. For the corresponding movies, please see Supporting Information Movies S1A/B, S2A/B and S3 A/B.

FIG 4. Automated Western blot analysis of BldC.

Equal amounts (1 µg) of total protein were loaded for each sample and BldC was detected with polyclonal antibody (9) using the quantitative 'Wes' capillary electrophoresis and blotting system (ProteinSimple – San Jose, CA). The *S. venezuelae bldC* mutant was used as a negative control. Both the wild-type and *bldC* mutant were grown in MYM liquid medium. (A) quantitation of BldO levels (area under each peak; arbitrary units). (B) virtual Western

blot. All experimental samples were analysed in triplicate and the mean value and its

Standard Error are shown for each sample. Each time-point is indicated in hours, along with

its relation to the developmental stage (V = vegetative growth; F = fragmentation; S =

sporulation), as determined by microscopy. Cultures used to analyse BldC levels were

identical to those used to prepare RNA prior to qRT-PCR analysis (Fig. 6).

FIG 5. BldC regulates the expression of many genes in *S. venezuelae*.

(A) Genome-wide distribution of BldC binding sites identified by ChIP-chip analysis using

anti-BldC polyclonal antibody, conducted during vegetative growth (12 hr) in the wild type.

DNA obtained from immunoprecipitation of BldC was labelled with Cy3 and hybridized to

DNA microarrays together with a total DNA control labelled with Cy5. Data are plotted as

Cy3/Cy5 ratios (y-axis), as a function of chromosome location (x-axis).

(B) ChIP-chip and qRT-PCR data for *whiI* and *smeA*. *Left Panels* - ChIP-chip in wild-type *S.*

venezuelae and the *S. venezuelae ΔbldC* mutant (blue and red dots, respectively). Plots span

approximately 8 kb of DNA sequence. *Right Panels* - mRNA abundance determined by qRT-

PCR in the wild type (white bars) and the *bldC* mutant (black bars). Strains were grown in

MYM liquid medium. Expression values were calculated relative to the accumulation of the

constitutively expressed *hrdB* reference mRNA and normalised to the wild-type value at 8 h.

FIG 6. ChIP-chip and qRT-PCR data for *sigF*, *whiD* and *hupS*. *Left Panels* - ChIP-chip in

wild-type *S. venezuelae* and the *S. venezuelae ΔbldC* mutant (blue and red dots, respectively).

Plots span approximately 8 kb of DNA sequence. *Right Panels* - mRNA abundance

determined by qRT-PCR in the wild type (white bars) and the *bldC* mutant (black bars).

Strains were grown in MYM liquid medium. Expression values were calculated relative to

the accumulation of the constitutively expressed *hrdB* reference mRNA and normalised to the

wild-type value at 8 h.

ΔbldC::apr
attB$_{ΦBT1}$::bldC   WT

ΔbldC::apr
attB$_{ΦBT1}$::pIJ10770   *ΔbldC::apr*

FIG 1

Scale Bar = 10 μm

FIG 2.

FIG 3.

FIG 4.

FIG 5.

sigF

FIG 6.

Supplemental Material

Table S1. ChIP-chip data set for *S.venezuelae* BldC. Each row represents an enriched probe (probeID) with the mid-position (midpos) of each probe on the *S. coilicolor* genome recorded. Enrichment ratios are expressed as the log-fold change (logFC). Probes are listed in order of significance (adjusted p value - adj.P.Val). For each probe, the nearest gene to the left and right is identified (left/rightLocusTag), its distance to the midpos of the probe (left/rightDistance), whether the gene is on the forward (1) or reverse (-1) strand (inStrand) and the predicted function (left/rightProduct) based on annotation in strepdb (http://strepdb.streptomyces.org.uk). If the midpos of a probe falls within a gene, it's gene identifier (inLocusTag), distance to the probe (inDistance), whether the gene is on the forward (1) or reverse (-1) strand (inStrand) and predicted function (inProduct) is also listed.

Table S2. Strains, Plasmids and Oligonucleotide primers used in this study

Movie S1. Time-lapse microscopy of the wild type strain carrying the FtsZ-YPet fusion. DIC (A) and YFP-channel (B) movies are at 5 frames per second. The time following the first image is indicated at the bottom left. Images were taken every 30 minutes (DIC 150 ms; YFP 100 ms). Movies were assembled in the Fiji software package (http://fiji.sc/Fiji).

Movie S2. Time-lapse microscopy of the *bldC* mutant carrying the FtsZ-YPet fusion. DIC (A) and YFP-channel (B) movies are at 5 frames per second. The time following the first image is indicated at the bottom left. Images were taken every 30 minutes (DIC 150 ms; YFP 100 ms). Movies were assembled in the Fiji software package (http://fiji.sc/Fiji).

410 **Movie S3.** Time-lapse microscopy of the complemented strain carrying the FtsZ-YPet fusion.

411 DIC (A) and YFP-channel (B) movies are at 5 frames per second. The time following the first

412 image is indicated at the bottom left. Images were taken every 30 minutes (DIC 150 ms; YFP

413 100 ms). Movies were assembled in the Fiji software package (http://fiji.sc/Fiji).