Genome-wide interaction study of a proxy for stress-sensitivity and its prediction of major depressive disorder

Running title: **Stress-sensitivity proxy and depression**

Aleix Arnau-Soler MSc[1], Mark J. Adams PhD[2], Generation Scotland[3], Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium[4], Caroline Hayward PhD[5], Pippa A. Thomson PhD[1,6]

[1] Medical Genetics Section, University of Edinburgh, Centre for Genomic and Experimental Medicine and MRC Institute of Genetics and Molecular Medicine, Edinburgh, UK

[2] Division of Psychiatry, Deanery of Clinical Sciences, University of Edinburgh, Royal Edinburgh Hospital, Morningside Park, Edinburgh EH10 5HF, UK

[3] A collaboration between the University Medical School and NHS in Aberdeen, Dundee, Edinburgh and Glasgow, Scotland, UK

[4] For a full list of MDD working group of the PGC investigators, see the Supplemental Information

[5] Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

[6] Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK

Correspondence should be addressed to: P. Thomson (Pippa.Thomson@ed.ac.uk)

Centre for Genomic and Experimental Medicine
Institute of Genetics and Molecular Medicine
University of Edinburgh
Western General Hospital
Crewe Road South
Edinburgh, EH4 2XU, UK
Phone: +44(0)1316518756

ABSTRACT

Individual response to stress is correlated with neuroticism and is an important predictor of both neuroticism and the onset of major depressive disorder (MDD). Identification of the genetics underpinning in individual differences in response to negative events (stress-sensitivity) may improve our understanding of the molecular pathways involved, and its association with stress-related illnesses. We sought to generate a proxy for stress-sensitivity through modelling the interaction between SNP allele and MDD status on neuroticism score in order to identify genetic variants that contribute to the higher neuroticism seen in individuals with a lifetime diagnosis of depression compared to unaffected individuals. Meta-analysis of genome-wide interaction studies (GWIS) in UK Biobank (N = 23 092; UKB) and Generation Scotland: Scottish Family Health Study (N = 7 155; GS:SFHS) identified no genome-wide significance SNP interaction. However, gene-based tests identified a genome-wide significant gene, *ZNF366*, a negative regulator of glucocorticoid receptor function implicated in alcohol dependence (p = 1.48x10^{-7}; Bonferroni-corrected significance threshold p < 2.79x10^{-6}). Using summary statistics from the stress-sensitivity term of the GWIS, SNP heritability for stress-sensitivity was estimated at 5.0%. In models fitting polygenic risk scores of both MDD and neuroticism derived from independent GWAS, we show that polygenic risk scores derived from the stress-sensitivity GWIS significantly improved the prediction of MDD. This study may improve interpretation of larger genome-wide association studies of MDD and other stress-related illnesses, and the understanding of the etiological mechanisms underpinning stress-sensitivity.

INTRODUCTION

Stressful life events are known to increase liability to mental illness and disease-related traits[1] including neuroticism[2-4], major depressive disorder (MDD)[5-7], autoimmune diseases[8] and some cancers[9,10]. A greater understanding of the causal mechanism by which negative events affect disease risk or outcome may be beneficial in identifying individuals for targeted support. However, it has been proposed that sensitivity to stress may be an important predictor of response to stress[11,12]. In particular, the effect on an individual may result more from the perceived stress than the event itself, and be dependent on individual differences in stress-sensitivity[13-18]. Studies of 5-HTT and twin studies suggest that stress-sensitivity may, at least in part, be heritable[19-21]. Despite a complex interaction between MDD, neuroticism and stress, multivariate structural equation models have confirmed a genetic effect on perceived stress, overlapping that on MDD or neuroticism, but with a specific genetic component[20]. The inter-relatedness of these traits may offer an approach to identify the genetic variation that affects an individual's stress-sensitivity, and improve genetic prediction of an individual's liability to negative outcomes. By modelling the interaction between SNP allele and MDD status on neuroticism score through genome-wide interaction studies (GWIS), we sought to investigate the genetics of stress-sensitivity.

The personality trait neuroticism is highly heritable (30–50% estimates from twin studies)[22-25], is higher in individuals with depression compared to controls[26,27] and is known to have shared genetic aetiology with depression[28-31]. Neuroticism is strongly correlated with measures of sensitivity to punishment but not reward[32], positively correlated with perceived personal relevance of a stressor[33,34] and has been used previously as a proxy measure of stress-sensitivity[35]. Neuroticism is thought to mediate or interact with the effects of adverse life events on risk of depression[5,36]. It has a

substantial stable component[37], however, there is evidence for change, as well as

stability, across the life span[2-4,38]. Whereas the stable component of neuroticism is

strongly determined by genetics, change in neuroticism score is attributed to the effects

of unshared environment[38]. Persistent change in neuroticism score has been shown in

response to life events[2-4].

Major depressive disorder (MDD) is a complex disorder influenced by both genetic

contributions and environmental risk factors, with heritability estimates from twin and

family studies of between 31-42%[39,40]. Confirmed environmental risk factors for MDD

include maternal infections, childhood maltreatment and negative life events[5-7,41,42].

However, few genetic studies have such information and even fewer prospective studies

exist. Incorporation of stressful life events has been shown to improve the ability to

predict MDD[43,44] and, although stress is an environmental risk factor, it may have an

independent genetic contribution to risk of depression[44-48].

These studies suggest that a genetic variable derived from the difference in neuroticism

levels seen in individuals with MDD compared to controls may allow us to identify

genetic loci important for stress-sensitivity. Further, polygenic risk scores (PRS) derived

from this stress-sensitivity variable may improve prediction of MDD over that based on

MDD or neuroticism PRS alone.

Using unrelated individuals from two large population-based samples, UK Biobank (UKB;

N = 23 092) and Generation Scotland: Scottish Family Health Study (GS:SFHS; N = 7 155),

we sought to identify genes involved in stress-sensitivity by performing GWIS for the

interaction between MDD status and SNP allele on neuroticism score. We identified a

gene significantly associated with stress-sensitivity and show that a PRS derived from

the interaction term of the GWIS, significantly predicts liability to depression

independently of the PRS for MDD and/or neuroticism.

MATERIALS AND METHODS

Cohorts' description

Study ethics are detailed in Supplemental Information: Ethics statements.

UK Biobank (UKB). UKB is a major national health resource that aims to improve the prevention, diagnosis and treatment of a wide range of illnesses. It recruited more than 500 000 participants aged from middle to older age who visited 22 assessment centres across the UK between 2006 and 2010. Data were collected on background and lifestyle, cognitive and physical assessments, sociodemographic factors and medical history. The scientific rationale, study design, ethical approval, survey methods, and limitations are reported elsewhere[49,50]. The present study was conducted on genome-wide genotyping data available from the first release of UKB data. Details of sample processing specific to UKB project are available at http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=155583 and the Axiom array at http://media.affymetrix.com/support/downloads/ manuals/axiom_2_assay_auto_workflow_user_guide.pdf. UKB genotyping and the stringent QC protocol applied to UKB data before it was released can be found at http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=155580. Imputed UKB genotype data[51] (imputed by UKB using a merged panel of the UK10K haplotype reference panel and the 1000 Genomes Phase 3 reference panel) with quality > 0.9 was hard-called using PLINK v1.9[52] and SNPs genotyped on GS:SFHS extracted. Individuals were removed based on UKB genomic analysis exclusion (UKB Data Dictionary item #22010), non-white British ancestry (#22006: genetic ethnic grouping; from those individuals who self-identified as British, PCs analysis was used to remove outliers), high genotype missingness (#22005), genetic relatedness (#22012; no pair of individuals have a KING-estimated kinship coefficient > 0.0442), QC failure in UK BiLEVE study (#22050 and #22051: UK BiLEVE Affymetrix and UK BiLEVE genotype quality controls for samples) and gender mismatch

(#22001: genetic sex). Further, individuals in UKB who were also participants of

GS:SFHS, and their relatives, were excluded to remove any overlap of individuals

between discovery and target samples. A dataset of 109 283 individuals with 557 813

imputed SNPs remained for further analysis, aged 40-79 (57 328 female, 51 954 male;

mean age = 57.1 years, s.d. = 7.99), of which 109 282 had data available for neuroticism

score and 23 092 had data available on MDD status (n_{cases} = 7 834, $n_{controls}$ = 15 258, n_{female}

= 11 510, n_{male} = 11 582; mean age = 57.7 years, s.d. = 8.04). Thus, the final dataset

comprised 23 092 unrelated individuals.

Generation Scotland Scottish Family Health Study (GS:SFHS). GS:SFHS is a family-

based genetic epidemiology study which includes 23 960 participants from ~ 7 000

Scottish family groups collected by a cross-disciplinary collaboration of Scottish medical

schools and the National Health Service (NHS) from Feb 2006 to Mar 2011. Participants

were interviewed and clinically assessed for a wide range of health-related traits

(including high-fidelity phenotyping for Major Depressive Disorder and related

endophenotypes), environmental covariates and linked to routine health records[53,54].

The protocol for recruitment is described in detail in previous publications[55,56]. GS:SFHS

genotyping and quality control is detailed elsewhere[57]. Individuals with more than 2%

missing genotypes and gender discrepancies were removed, as well as population

outliers. SNPs with genotype missingness > 2%, minor allele frequency < 1% and a

Hardy-Weinberg Equilibrium test $p < 1x10^{-6}$ were exclude. Finally, individuals were

removed based on relatedness (pi-hat < 0.05), maximizing retention of case individuals,

using PLINK v1.9[52]. Genome-wide SNP data comprised 7 233 unrelated individuals

genotyped for 560,698 SNPs (n_{female} = 3 476, n_{male} = 3 757; PLINK v1.9[52]), aged 18-92

(mean age = 50.4 years, s.d. = 12.06) of which: 7 190 had clinical data on MDD; 7 196

individuals had data on neuroticism; and 7 155 had data on both neuroticism and MDD,

remained for further analysis.

Phenotype assessment

Neuroticism score (EPQN). Participants in both UKB and GS:SFHS cohorts were assessed

for neuroticism using 12 questions from the Eysenck Personality Questionnaire-Revised

Short Form's Neuroticism Scale (EPQN)[58-61]. Neuroticism can be scored by adding up the

number of "Yes" responses on EPQN. This short scale has a reliability of more than 0.8[62].

EPQN distributions were found to be sufficiently "normal" after assessment for skewness

and kurtosis to be analysed using linear regression (both coefficients were between -1

and 1).

MDD status. In UKB depressive symptoms were assessed by items relating to the

lifetime experience of minor and major depression[58], items from the Patient Health

Questionnaire[63] and items on help-seeking for mental health[61]. Controls were defined as

those who a) answered "No" to the question "Ever depressed for a whole week" or b)

who reported symptoms but duration was too short (<2 weeks) or who did not report

having seen a GP or psychiatrist for nerves, anxiety, tension or depression. In our

unrelated sample: 7 834 participants were diagnosed with MDD (with single, moderate

or recurrent episodes) and 15 258 were controls (N = 23 092).

In GS:SFHS, participants who answered yes to either of two screening questions were

assessed for psychiatric, mood state/psychological distress, personality and cognitive

assessment by trained researchers using the Structured Clinical Interview for DSM-IV

Non-Patient Version (SCID)[64]. In our unrelated sample, of those individuals with data

available for both MDD and EPQN (N = 7 155), used for the GWIS, 2 010 had a lifetime

diagnosis of MDD and 5 145 were controls.

GWIS and derivation of a genetic stress-sensitivity effect

The effect size of an stress-sensitivity effect (β_{SS}) was derived by performing a GWIS for

the effect of the MDD status and SNP allele on EPQN (dependent variable) in both UKB

and GS:SFHS cohorts using PLINK 1.90 (PLINK-command --gxe; fitting MDD diagnosis as

a binary "group" effect)[52]. PLINK-command --gxe estimates the difference in allelic

association with a quantitative trait (EPQN) between two groups (MDD cases vs.

controls) producing effect estimates on each group and a test of significance for the

interaction between SNP allele and MDD status. The interaction p value reflects the

difference between the regression coefficient of the allelic effect in a linear model for

EPQN in MDD cases (β_A) and the same regression coefficient in a linear model for EPQN

in controls (β_B). The stress-sensitivity interaction effect was defined as the difference in

allele effect between MDD cases and control groups.

Considering one SNP, the effect it confers to EPQN can be modelled by MDD status

(control = 0, MDD case = 1) as follows:

$$\begin{cases} MDD = 0; \ EPQN = \beta_0 + \beta_B SNP + \beta_{0c} COV + \varepsilon \\ MDD = 1; \ EPQN = \beta_1 + \beta_A SNP + \beta_{1c} COV + \varepsilon \end{cases}$$

193 This is equivalent to modelling the effect on MDD cases as follows:

$$\begin{cases} MDD = 0; \ EPQN = \beta_0 + \beta_B SNP + \beta_{0c} COV + \varepsilon \\ MDD = 1; \ EPQN = \beta_1 + \beta_B SNP + (\beta_A - \beta_B) SNP + \beta_{1c} COV + \varepsilon \end{cases}$$

194 Or, it can be modelled as a whole as:

$$EPQN = \beta_0 + \beta_2 MDD + \beta_B SNP + (\beta_A - \beta_B) SNP * MDD + \beta_{0c} COV + \beta_{2c} COV * MDD + \varepsilon$$

195 Where COV stands for covariates, β_2 stands for $\beta_1 - \beta_0$, and β_{2c} stands for $\beta_{1c} - \beta_{0c}$.

196   Thus, the interaction effect (β_{SS}) can be estimated as the difference in allelic effect on

197   EPQN between MDD cases (β_A) and controls (β_B) as follows,

$$\hat{\beta}_{SS} = \hat{\beta}_A - \hat{\beta}_B$$

$\hat{\beta}_{SS}$ is therefore defined as the effect size reflecting the genetic stress-sensitivity effect on

MDD cases compared to controls (see Supplemental Figure S1).

Stress-sensitivity GWIS, main additive effect GWASs, meta-analysis and gene-set

analysis.

For GWIS and subsequent analyses, sample specific covariates were applied as follows:

UKB. All phenotypes were adjusted for centre, array and batch as random effects prior to

analyses. Analyses were adjusted for age, sex and 15 informative principal components

(PCs; UKB Data Dictionary items #22009.01 to #22009.15) as fixed effects to take

account of possible population stratification. *GS:SFHS.* All the analyses were adjusted for

age, sex and 20 PCs.

GWAS for MDD and neuroticism, using logistic and linear models of additive allelic

effects respectively, were conducted on the same sample sets for comparison and

generation of matched PRS using PRSice[65].

Results from the GWIS of UKB and GS:SFHS were combined in a sample size weighted

meta-analysis performed using METAL[66]. While the use of standard error weighting is

more common, the different diagnostic schema and MDD prevalence between the two

cohorts (GS:SFHS; 12.2%, UKB: 25.8%)[55,61] may indicate systematic differences in the

measurement of MDD. Generalized gene-based analysis of the meta-analysis was

performed using MAGMA[67] implemented through FUMA[68] (http://fuma.ctglab.nl).

Briefly, SNP summary statistics were mapped to 17 931 protein-coding genes. Individual

SNP *p* values from a gene were combined into a gene test-statistic using a SNP-wise

model and a known approximation of the sampling distribution used to obtain a gene-

based p value. Genome-wide significance was defined at $p = 0.05/17\,931 = 2.79 \times 10^{-6}$.

LD Score regression

The summary statistics from the meta-analysis were used to examine the genetic overlap

between the polygenic architecture of stress-sensitivity, MDD and neuroticism. LD score

regression was used to derive the genetic correlations (r_G) between these traits[69,70] using

meta-analysed GWAS and GWIS summary statistics. SNP-based heritability was also

estimated using LD score regression, which only requires summary statistics from

single-SNP analyses.

Pathway, functional and gene expression analyses

Lead SNPs, independently associated with the phenotype, were identified using PLINK

1.90 by clumping (p threshold $< 2 \times 10^{-5}$; LD r2 > 0.1; physical kb threshold = 500kb; 1000

Genomes Project Phase 1 CEU, GBR, TSI genotype data), and analysed using DEPICT[71].

Further detail is given in the Supplemental Information: DEPICT analyses.

Genes associated with lead SNPs were investigated for evidence of: phenotypic

association in the NCBI dbGaP database of genotypes and phenotypes[72]

(https://www.ncbi.nlm.nih.gov/gap/phegeni), regulatory DNA elements in normal cell

lines and association with expression quantitative trait loci (eQTLs) using the

RegulomeDB database[73] (http://www.regulomedb.org) and the Genotype-Tissue

Expression (GTEx) Portal[74] (http://www.gtexportal.org).

Polygenic profiling

PRS were produced using PRSice[65] and standardized to a mean of 0 and a standard

deviation of 1. PRS for stress-sensitivity (PRS_{SS}) were created by weighting the sum of

the reference alleles in an individual by the stress-sensitivity effect, β_{SS} using GWIS

summary statistics. Additional PRS were generated weighting by MDD main additive

effects (PRS_D) and neuroticism main additive effects (PRS_N) using GWAS summary

statistics from GS:SFHS or UKB. In addition, PRS_D and PRS_N were also generated using

summary statistics from the most recent Psychiatric Genetic Consortium (PGC) MDD

meta-analysis[40] (excluding GS:SFHS, and UKB individuals when required; N = 155 866 &

138 884) and the Genetics of Personality Consortium (GPC) neuroticism meta-

analysis[23,75] (N = 63 661). Generalized linear models were implemented in R 3.1.3[76]. The

direct effect of PRS_{SS} (model 1), PRS_D (model 2) and PRS_N (model 3) on MDD risk were

assessed in independent logistic regression models on GS:SFHS (target cohort) using

GWAS and GWIS statistics from UKB (the largest cohort) as the discovery sample to

weight PRS. Multiple regression models fitting both PRS_D and PRS_N (model 4) and fitting

each of them separately with PRS_{SS} (models 5 and 6) were also calculated. Finally, full

additive multiple regression models fitting PRS weighted by all three effects (full model)

was assessed using both PRS_{SS}, PRS_D and PRS_N at their best-fit in independent models.

Further, results were also assessed using PRS_D and PRS_N weighted by PGC2 MDD[40] and

GPC neuroticism[75] summary statistics. Further detail is given in the Supplemental

Information: Polygenic Profiling. All models were adjusted by sex, age and 20 PCs. A null

model was estimated from the direct effects of all covariates on MDD. Likelihood ratio

tests were used to assess significance of each PRS.

Cross-validation was performed using UKB as target sample and GS:SFHS as discovery

sample. Additional analyses using PRS_D and PRS_N weighted by PGC2 MDD[40] and GPC

neuroticism[75] summary statistics were also tested. MDD status on UKB was adjusted by

centre, array and genotyping batch as random effects and scaled (between 0 and 1) prior

to analysis, giving a quasi-binomial distribution of MDD status on UKB. Models

implemented on UKB (quasi-binomial regression) were adjusted by sex, age and 15 PCs.

Nagelkerke's R^2 coefficients were estimated to quantify the proportion of MDD liability

explained at the observed scale by each model and converted into R^2 coefficients at the

liability scale (prevalence: 12.2% in GS:SFHS[55] and 25.8% in UKB[61]) using Hong Lee's

transformation[77] available from GEAR: GEnetic Analysis Repository[78].

Using stress-sensitivity to stratify depression

GS:SFHS MDD cases (n_{cases} = 2 016; n_{female} = 1 345, n_{male} = 671) have data available on

MDD course (single or recurrent), age of onset (n = 1 964) and episode count (n = 2 016),

as well as on neuroticism (n = 2 010). In addition, a subset were evaluated by Mood

Disorder Questionnaire[79] (MDQ; n = 1 022) and Schizotypal Personality Questionnaire[80]

(SPQ; n = 1 093). The reduced sample number of MDQ and SPQ reflects the later addition

of these questionnaires to the study and does not reflect a particular subgroup of

GS:SFHS.

Difference in PRS_{SS} and PRS_D between MDD cases and controls on GS:SFHS were tested

using a Student's two sample t-test (two tailed). Cases of MDD on GS:SFHS with data

available on each trait analyzed were stratified by quintiles based on PRS_{SS} and PRS_D

(5x5 groups). Post hoc, the effects on each trait of quintiles based on PRS_{SS} and its

interaction effect with quintiles based on PRS_D were assessed using linear regression

models adjusting by sex and age in an attempt to identify a characteristic subtype of MDD

patients with differential stress-sensitivity levels. The same analysis was reproduced

using PRSs as continuous variables.

RESULTS

We confirmed the elevated neuroticism score in MDD cases in our samples. Individuals

with a diagnosis of MDD had significantly higher EPQN scores compared to healthy

controls (all $p < 1.9 \times 10^{-279}$) in both GS:SFHS ($\text{mean}_{\text{controls}} = 3.16$; $\text{mean}_{\text{cases}} = 6.42$) and

UKB ($\text{mean}_{\text{controls}} = 2.79$; $\text{mean}_{\text{cases}} = 5.64$). This difference remained significant after

matching cases with controls for PGC PRS_{D}, sex and age. (GS:SFHS: $\text{mean}_{\text{controls}} = 3.51$;

UKB: $\text{mean}_{\text{controls}} = 2.97$; all $p < 2.7 \times 10^{-158}$; see Supplemental Table S1).

Meta-analysis of stress-sensitivity in UKB and GS:SFHS

No SNPs were associated with stress-sensitivity at the genome-wide significant

threshold ($p < 5 \times 10^{-8}$, Figure 1). However, 14 SNPs from 8 loci achieved suggestive p

value ($p < 1 \times 10^{-5}$) ranging between $p = 8.9 \times 10^{-6}$-5.1×10^{-7} (Meta-analysis: Table 1; UKB

and GS:SFHS: Supplemental Tables S2 & S3; Meta-analysis QQ-plot with λ: Supplemental

Figure S2; UKB and GS:SFHS QQ-plots: Supplemental Figure S3). Traits with prior

evidence of association with the nearest genes to the 8 lead SNPs were identified using

dbGap and are shown in Supplemental Table S4. Comparison between the SNP

association profile along the genome between stress-sensitivity GWIS and MDD GWAS

meta-analyses is shown in Miami plots filtering for the most significant stress-sensitivity

or MDD SNPs ($p < 0.001$; Meta-analysis: Figure 2; UKB and GS:SFHS: Supplemental Figure

S4). No SNP with a p-value < 0.01 had a corresponding p-value in the alternate trait,

suggesting that different variants contribute to depression and stress-sensitivity. Gene-

based test identified *ZNF366* as the only gene achieving genome-wide significance ($p = $

1.48×10^{-7}; Bonferroni-corrected significance threshold $p < 2.79 \times 10^{-6}$; Supplemental Table

S5 & Supplemental Figure S5). Using summary statistics from meta-analysis GWIS

results, stress-sensitivity SNP-based heritability was estimated at 5.0% ($h^2 = 0.0499$, s.e.

= 0.017, p = 1.67x10^{-3}). Conversely, matched SNP-based heritabilities for MDD and

neuroticism were estimated at 9.6% (h^2 = 0.0962, s.e. = 0.0179, p = 3.87x10^{-8}) and 10.1%

(h^2 = 0.1006, s.e. = 0.0076, p = 3.47x10^{-40}) respectively, using summary statistics from

the meta-analysed GWAS of UKB and GS:SFHS.

LD score regression was performed to obtain genetic correlations between stress-

sensitivity, MDD and neuroticism. As previously shown, there was a significant genetic

correlation between MDD and neuroticism (r$_G$ = 0.637, s.e. = 0.0704, p = 1.39x10^{-19}).

However, we found no evidence for a genetic correlation between stress-sensitivity and

MDD (rG = -0.099, s.e. = 0.182, p = 0.585) or between stress-sensitivity and neuroticism

(r$_G$ = 0.114, s.e. = 0.107, p = 0.285).

Pathway enrichment, functional annotation and gene expression analyses

Lead SNPs from the GWIS meta-analysis were investigated using DEPICT. No gene

showed statistically significant links to stress-sensitivity at a DEPICT false discovery rate

(FDR) < 0.05. No significant result was found for either gene set analysis or tissue

enrichment analysis at FDR < 0.05. Evidence of regulatory elements on normal cell

lines/tissues was identified for 5 of the 12 lead SNPs (i.e. rs3762096, rs10987199,

rs2221540, rs246565, rs319924). Two lead SNPs were associated with eQTLs: rs319924

(an intronic SNP in *EYS*) and rs9509508 (an intronic SNP in *LATS2*) and potentially

regulate *LGSN/RP3-407E4.3* (p = 6.31x10^{-12}/ p = 1.15x10^{-5}) and *LATS2* (p = 3.74x10^{-8}),

respectively.

Polygenic risk scores for stress-sensitivity predict MDD liability.

PRS were used to investigate whether common variants affecting stress-sensitivity

predict MDD risk. We generated PRS (PRS$_{SS}$) based on the summary statistics from the

GWIS for stress-sensitivity. PRS_{SS} significantly predicted MDD risk in GS:SFHS using

weights from the larger UKB summary data ($p = 1.8x10^{-3}$; β = 0.087, s.e. = 0.028; best-fit

p threshold = 0.004). This association was cross-validated (nominal significance $p < 0.05$)

in UKB using summary data from the smaller GS:SFHS GWIS ($p = 0.04$; β = 0.027, s.e. =

0.013; best-fit p threshold = $1.15x10^{-4}$; Table 2 and Supplemental Figure S6 & S7). On the

liability scale, the MDD variance explained by PRS_{SS} was modest, 0.243% in GS:SFHS and

0.037% in UKB. This was less than predicted by PRS weighted by the genetic main effects

of MDD or neuroticism (PRS_D: 0.386% in GS:SFHS and 0.211% in UKB; PRS_N: 0.34% in

GS:SFHS and 0.178% in UKB; Table 2 and Supplemental Figure S6).

Due to the known genetic correlations between MDD, neuroticism and stressful life

events[20], models jointly fitting the effects of multiple PRS were analysed. Multiple

regression analyses showed that, compared to PRS_D effects alone, stress-sensitivity effect

significantly explains an additional 0.248% in GS:SFHS (a predictive improvement of

64.3%; $p = 1.6x10^{-3}$; PRS_D: β = 0.113, s.e. = 0.028; PRS_{SS} : β = 0.088, s.e. = 0.028) and an

additional 0.036% in UKB (a predictive improvement of 17.1%; $p = 0.044$; PRS_D: β =

0.064, s.e. = 0.013; PRS_{SS} : β = 0.027, s.e. = 0.013; Figure 3). The inclusion of PRS_{SS} in the

full model, where PRS_{SS} was fitted along with both PRS_D and PRS_N weighted by GWAS

summary statistics derived from the same datasets, remained significant; explaining an

additional 0.245% (a predictive improvement of 35.8%: full model $p = 1.7x10^{-3}$; PRS_D: β =

0.107, s.e. = 0.029; PRS_N: β = 0.123, s.e. = 0.036; PRS_{SS}: β = 0.088, s.e. = 0.028) and an

additional 0.036% (a predictive improvement of 11.2%; full model $p = 0.044$; PRS_D: β =

0.054, s.e. = 0.014; PRS_N: β = 0.048, s.e. = 0.014; PRS_{SS} : β = 0.027 and s.e. = 0.013) in

GS:SFHS and UKB, respectively.

In models fitting PRS_D and PRS_N, the variances explained were non-additive,

demonstrating the partial overlap between MDD risk prediction from PRS_D and PRS_N

main additive effects. This overlap was not seen between variance explained by stress-

sensitivity effect and the variance explained by either PRS_D and/or PRS_N.

Multiple regression analyses fitting PRS_D and PRS_N derived from worldwide consortiums

(Figure 3) showed that the increased sample size from GWAS used to derive PRS_D

resulted in an increment of MDD variance explained by PRS_D (GS:SFHS: from 0.386% to

1.233%; UKB: from 0.21% to 1.281%). However, there was no change in the proportion

of the variance explained by the PRS_{SS} in the full model (PRS_{SS} $p = 1.29\text{x}10^{-3}$ and $p = 0.042$

in GS:SFHS and UKB, respectively). These results suggest that PRS_{SS} explains a

proportion of MDD risk not accounted for by PRS_D or PRS_N at current sample sizes.

Using stress-sensitivity to stratify MDD in GS:SFHS

MDD cases show significantly higher PRS_{SS} ($p = 2\text{x}10^{-3}$) and PRS_D ($p = 1.8\text{x}10^{-4}$) than

controls. Association between MDD-related traits and stress-sensitivity risk quintiles

was assessed on MDD cases in order to identify a subgroup of MDD patients, perhaps

defining a characteristic aetiological subtype of MDD. However, stratification analysis

failed, and no quintile based on PRS_{SS} nor its interaction with quintiles based on PRS_D

showed statistically significant effects on any trait analyzed. Individuals with high PRS_{SS}

were not significantly different from other cases for sex, MDD course, age of onset or

episode count, nor neuroticism, mood disorder or schizotypal personality scores ($p >$

0.05; see Supplemental Table S6). Results remained non significant when PRSs were

used as continuous variables ($p > 0.05$).

DISCUSSION

The existence of genetic variants affecting an individual's risk of depression in response

to stress has been predicted previously[44,47,48] and is consistent with the departure from a

simple additive genetic model seen in twin-studies of recurrent depressive disorder[81].

Through international research efforts such as the PGC and UK Biobank, there are ever-

increasing sample sizes available for understanding the genetics of MDD. These

resources are beginning, and will continue to, identify genome-wide significant loci[82,83]

[40]. However, the lack of environmental data and/or their reliability, makes the study of

genetic individual's response to their negative effects, and their contribution to the onset

of MDD and other stress-related disorders, difficult. As a way to address this limitation,

we generated a proxy for stress-sensitivity through modelling the interaction between

SNP allele and MDD status on neuroticism score in a GWIS approach. Thus, we sought to

identify the genetic underpinnings of individual's sensitivity to stress response (stress-

sensitivity) through those variants that contribute to higher neuroticism levels only in

individuals with a lifetime diagnosis of MDD but not in healthy controls.

We performed a GWIS to identify loci showing differential effects on neuroticism scores

in individuals with and without MDD (so called stress-sensitivity proxy). No SNPs

reached genome-wide significance, but 14 SNPs from 8 loci reached suggestive

significance levels. The closest genes to these SNPs for stress-sensitivity have prior

evidence of association in dbGAP with a wide range of neuropsychiatric traits such as

schizophrenia, bipolar disorder, attention deficit disorder with hyperactivity, mental

competency, intuition, sleep or alcohol drinking (Supplemental Table S4: Traits with

significant evidence of association with suggestive stress-sensitivity genes). However,

enrichment analysis showed no evidence for enrichment of specific pathways or tissues.

The top two loci, *PTP4A1-PHF3-EYS* and *ZNF366* have been previously associated with

alcohol dependence[84-88], alcohol intake (dbGaP: phs000342) and glucocorticoid receptor

function[89-91]. The most significant SNP in this study, rs319924, is an intronic variant in

EYS that is a potential eQTL for *LGSN*[74], a gene previously associated with male-specific

depression[92]. This is of particular interest given previous studies linking alcohol

consumption, stress and the risk of depression[93-98]. Evidence of an eQTL effect was also

predicted for a lead SNP in *LATS2*, a positive regulator of histone methyltransferase

activity[99] a process important in anxiety-related behaviours[100]. The prior association of

the top two loci in this study with alcohol related-phenotypes suggests that genes

involved in the sensitivity to stress may mediate the effects of stress on alcohol

consumption. Some *PHF3* paralogs have been shown to be linked with depression and

modulate stress response [101,102].

Gene-based analysis identified a genome-wide significant association between *ZNF366*

and stress-sensitivity. *ZNF366* (also known as *DC-SCRIPT*) is a corepressor of

transcription found in nuclear receptor complexes including the glucocorticoid receptor.

ZNF366 represses glucocorticoid receptor-mediated transcription in monocyte-derived

dendritic cells[89]; and may act through histone deacetylases to modulate immune

response[90]. There is evidence from a large-scale mRNA display study that *PHF3*, in the

region underlying the most significant peak in the single SNP analysis, may also interact,

directly or indirectly, with the glucocorticoid receptor (IntAct database[91]) but this has

not been confirmed. These results reinforce the hypothesis that our proxy for stress-

sensitivity truly reflects the genetic architecture of sensitivity to respond to stress.

We estimated a significant lower bound on heritability for stress-sensitivity of 5%.

Whilst the known genetic overlap between MDD and neuroticism was detectable, the

lack of genetic correlation with stress-sensitivity indicated that there is no significant

overlap between stress-sensitivity and MDD or neuroticism. This analysis may be limited

by our sample size, however using the largest available meta-analyses of MDD and

neuroticism[40,75] did not decrease the proportion of liability explained by the PRS$_{SS}$. We

note, however, that as such meta-analyses increase in size it is likely, as with the effects

of smoking in schizophrenia[103,104], that the indirect genetic effects of the environment on

the risk of depression will be detected by GWAS. However, it is only through studies such

as ours that the mechanism for the effect of the risk alleles may be clarified.

Further, we show that such genetic information in stress-sensitivity could significantly

improve the proportion of liability to MDD predicted by PRS based only on additive

genetic effects on MDD identified by large GWAS. The summary results from the GWIS

were used to derive a PRS reflecting the genetic difference in stress-sensitivity. Although

not robust to multiple testing correction, this variable significantly predicted liability to

MDD in GS:SFHS (p = 1.8x10^{-3}) and this was replicated in UKB (p = 0.04). This is

consistent with the expectation that the larger the discovery sample (i.e. UKB), the

greater the accuracy of the weighting and the more predictive the PRS[105]. Multiple

regression models suggest that inclusion of PRS weighted by stress-sensitivity

significantly improves MDD prediction over use of either MDD and/or neuroticism

weighted PRS alone (GS:SFHS full model p = 1.7x10^{-3}; UKB full model p = 0.04). However,

we were unable to identify a subgroup of MDD cases with higher PRS$_{SS}$. The polygenic

interaction approach used in our study may, therefore, improve the interpretation of

both positive and negative findings from GWAS studies (i.e. pathways and mechanisms

involved, lack of replication, or negative findings in variants mediating environmental

effects). Added to paralleling recent developments in GWAS analyses, it may maximize

our power to detect gene-by-environment effects in this heterogeneous disorder

Future studies will be required to further investigate the effects of adverse life events in

individuals with high or low polygenic risk scores for stress-sensitivity. However, the

methodology presented allows addressing the genetic response to negative outcomes via

proxy in the absence of prospective environmental data.

Here we identify an independent set of risk variants for an individual's response to

negative outcomes and show that incorporating information across many loci provides

clear and replicable evidence for a genetic effect of stress-sensitivity on MDD risk;

identifying a potential genetic link with alcohol intake. These results require further

study, but may inform treatment of comorbid alcohol dependency and depression.

ACKNOWLEDGMENTS

The 1st author AAS is funded by University of Edinburgh (www.ed.ac.uk) and Medical Research Council for his PhD study at the University of Edinburgh Institute of Genetics and Molecular Medicine (www.ed.ac.uk/igmm). MA is supported by STRADL through a Wellcome Trust Strategic Award (reference 104036/Z/14/Z). Generation Scotland received core support from the Chief Scientist Office of the Scottish Government Health Directorates [CZD/16/6] and the Scottish Funding Council [HR03006]. Genotyping of the GS:SFHS samples was carried out by the Genetics Core Laboratory at the Wellcome Trust Clinical Research Facility, Edinburgh, Scotland and was funded by the Medical Research Council UK and the Wellcome Trust (Wellcome Trust Strategic Award "STratifying Resilience and Depression Longitudinally" (STRADL), Reference 104036/Z/14/Z). The Psychiatric Genomics Consortium MDD Working Group depends on the contributions of many parties.

FINANCIAL DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

1.   Monroe SM. Modern approaches to conceptualizing and measuring human life
stress. *Annu Rev Clin Psychol* 2008; **4**: 33-52.

2.   Jeronimus BF, Ormel J, Aleman A, Penninx BW, Riese H. Negative and positive life
events are associated with small but lasting change in neuroticism. *Psychol Med* 2013;
43(11): 2403-2415.

3.   Jeronimus BF, Riese H, Sanderman R, Ormel J. Mutual reinforcement between
neuroticism and life experiences: a five-wave, 16-year study to test reciprocal causation.
J Pers Soc Psychol 2014; **107**(4): 751-764.

4.   Riese H, Snieder H, Jeronimus BF, Korhonen T, Rose RJ, Kaprio J, *et al*. Timing of
Stressful Life Events Affects Stability and Change of Neuroticism. *European Journal of*
Personality 2014; **28**(2): 193-200.

5.   Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex, and
stressful life events in the prediction of episodes of major depression. *Am J Psychiatry*
2004; **161**(4): 631-636.

6.   Kessler RC. The effects of stressful life events on depression. *Annu Rev Psychol*
1997; **48**: 191-214.

7.   Tennant C. Life events, stress and depression: a review of recent findings. *Aust N Z*
J Psychiatry 2002; **36**(2): 173-182.

8.   Porcelli B, Pozza A, Bizzaro N, Fagiolini A, Costantini MC, Terzuoli L, *et al*.
Association between stressful life events and autoimmune diseases: A systematic review
and meta-analysis of retrospective case-control studies. *Autoimmun Rev* 2016; **15**(4):
325-334.

9.   Lin Y, Wang C, Zhong Y, Huang X, Peng L, Shan G, *et al*. Striking life events
associated with primary breast cancer susceptibility in women: a meta-analysis study. *J*
Exp Clin Cancer Res 2013; **32**(1): 53.

10.   Jafri SHR, Ali F, Mollaeian A, Hasan SM, Hussain R, Akkanti BH, *et al*. Major
stressful life events and risk of developing lung cancer. *Journal of Clinical Oncology* 2017;
35(15_suppl): 1575-1575.

11.   Hasler G, Drevets WC, Manji HK, Charney DS. Discovering endophenotypes for
major depression. *Neuropsychopharmacology* 2004; **29**(10): 1765-1781.

12.   Bogdan R, Nikolova YS, Pizzagalli DA. Neurogenetics of depression: a focus on
reward processing and stress sensitivity. *Neurobiol Dis* 2013; **52**: 12-23.

13.   Lazarus RS, Folkman S. Transactional theory and research on emotions and
coping. *European Journal of Personality* 1987; **1**(3): 141-169.

14.   Smith MA, Riccalton VC, Kelly-Hughes DH, Craw OA, Allen SF, O'Connor DB, *et al*.
The relationship between Type D personality and physical health complaints is
mediated by perceived stress and anxiety but not diurnal cortisol secretion. *Stress* 2018:
1-8.

15.   Herr RM, Barrech A, Riedel N, Gundel H, Angerer P, Li J. Long-Term Effectiveness
of Stress Management at Work: Effects of the Changes in Perceived Stress Reactivity on
Mental Health and Sleep Problems Seven Years Later. *Int J Environ Res Public Health*
2018; **15**(2).

16.   Moore RC, Eyler LT, Mausbach BT, Zlatar ZZ, Thompson WK, Peavy G, *et al*.
Complex interplay between health and successful aging: role of perceived stress,
resilience, and social support. *Am J Geriatr Psychiatry* 2015; **23**(6): 622-632.

17.   Hayman LW, Jr., Lucas T, Porcerelli JH. Cognitive appraisal vs. exposure-based
stress measures: links to perceived mental and physical health in low-income black
women. *J Nerv Ment Dis* 2014; **202**(11): 807-812.

18.   Harris ML, Loxton D, Sibbritt DW, Byles JE. The influence of perceived stress on
the onset of arthritis in women: findings from the Australian Longitudinal Study on
women's health. *Ann Behav Med* 2013; **46**(1): 9-18.

19.   Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE. Genetic sensitivity to the
environment: the case of the serotonin transporter gene and its implications for
studying complex diseases and traits. *Am J Psychiatry* 2010; **167**(5): 509-527.

20.   Rietschel L, Zhu G, Kirschbaum C, Strohmaier J, Wust S, Rietschel M, *et al*.
Perceived stress has genetic influences distinct from neuroticism and depression. *Behav
Genet* 2014; **44**(6): 639-645.

21.   Rietschel L, Streit F, Zhu G, McAloney K, Frank J, Couvy-Duchesne B, *et al*. Hair
Cortisol in Twins: Heritability and Genetic Overlap with Psychological Variables and
Stress-System Genes. *Sci Rep* 2017; **7**(1): 15351.

22.   Glahn DC, Curran JE, Winkler AM, Carless MA, Kent JW, Jr., Charlesworth JC, *et al*.
High dimensional endophenotype ranking in the search for major depression risk genes.
Biol Psychiatry 2012; **71**(1): 6-14.

23.   van den Berg SM, de Moor MH, McGue M, Pettersson E, Terracciano A, Verweij KJ,
et al. Harmonization of Neuroticism and Extraversion phenotypes across inventories
and cohorts in the Genetics of Personality Consortium: an application of Item Response
Theory. *Behav Genet* 2014; **44**(4): 295-313.

24.   Wray NR, Birley AJ, Sullivan PF, Visscher PM, Martin NG. Genetic and phenotypic
stability of measures of neuroticism over 22 years. *Twin Res Hum Genet* 2007; **10**(5):
695-702.

582
583   25.   Vukasovic T, Bratko D. Heritability of personality: A meta-analysis of behavior
584   genetic studies. *Psychol Bull* 2015; **141**(4): 769-785.

585
586   26.   Hirschfeld RM, Klerman GL. Personality attributes and affective disorders. *Am J*
587   *Psychiatry* 1979; **136**(1): 67-70.

588
589   27.   Hirschfeld RM, Klerman GL, Clayton PJ, Keller MB, McDonald-Scott P, Larkin BH.
590   Assessing personality: effects of the depressive state on trait measurement. *Am J*
591   *Psychiatry* 1983; **140**(6): 695-699.

592
593   28.   Levinson DF. The genetics of depression: a review. *Biol Psychiatry* 2006; **60**(2):
594   84-92.

595
596   29.   Middeldorp CM, Cath DC, Van Dyck R, Boomsma DI. The co-morbidity of anxiety
597   and depression in the perspective of genetic epidemiology. A review of twin and family
598   studies. *Psychol Med* 2005; **35**(5): 611-624.

599
600   30.   Smith DJ, Escott-Price V, Davies G, Bailey ME, Colodro-Conde L, Ward J, *et al*.
601   Genome-wide analysis of over 106 000 individuals identifies 9 neuroticism-associated
602   loci. *Mol Psychiatry* 2016; **21**(6): 749-757.

603
604   31.   Luciano M, Hagenaars SP, Davies G, Hill WD, Clarke TK, Shirali M, *et al*. Association
605   analysis in over 329,000 individuals identifies 116 independent variants influencing
606   neuroticism. *Nat Genet* 2018; **50**(1): 6-11.

607
608   32.   Smits DJM, Boeck PD. From BIS/BAS to the big five. *European Journal of*
609   *Personality* 2006; **20**(4): 255-270.

610
611   33.   Schneider TR, Rench TA, Lyons JB, Riffle RR. The influence of neuroticism,
612   extraversion and openness on stress responses. *Stress Health* 2012; **28**(2): 102-110.

613
614   34.   Kim SE, Kim HN, Cho J, Kwon MJ, Chang Y, Ryu S, *et al*. Direct and Indirect Effects
615   of Five Factor Personality and Gender on Depressive Symptoms Mediated by Perceived
616   Stress. *PLoS One* 2016; **11**(4): e0154140.

617
618   35.   Tak LM, Kingma EM, van Ockenburg SL, Ormel J, Rosmalen JG. Age- and sex-
619   specific associations between adverse life events and functional bodily symptoms in the
620   general population. *J Psychosom Res* 2015; **79**(2): 112-116.

621
36.   Hovens JG, Giltay EJ, van Hemert AM, Penninx BW. Childhood Maltreatment and
the Course of Depressive and Anxiety Disorders: The Contribution of Personality
Characteristics. *Depress Anxiety* 2016; **33**(1): 27-34.

37.   Bazana PG, Stelmack RM. Chapter 8 - Stability of Personality Across the Life Span:
A Meta-Analysis. *On the Psychobiology of Personality*. Elsevier: Oxford, 2004, pp 113-144.

38.   McGue M, Bacon S, Lykken DT. Personality Stability and Change in Early
Adulthood. *Developmental Psychology* 1993; **29**(1): 96-109.

39.   Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression:
review and meta-analysis. *Am J Psychiatry* 2000; **157**(10): 1552-1562.

40.   Wray NR, Sullivan PF. Genome-wide association analyses identify 44 risk variants
and refine the genetic architecture of major depression. *bioRxiv* 2017.

41.   Loret de Mola C, de Franca GV, Quevedo Lde A, Horta BL. Low birth weight,
preterm birth and small for gestational age association with adult depression:
systematic review and meta-analysis. *Br J Psychiatry* 2014; **205**(5): 340-347.

42.   Peyrot WJ, Milaneschi Y, Abdellaoui A, Sullivan PF, Hottenga JJ, Boomsma DI, *et al*.
Effect of polygenic risk scores on depression in childhood trauma. *Br J Psychiatry* 2014;
205(2): 113-119.

43.   Mullins N, Power RA, Fisher HL, Hanscombe KB, Euesden J, Iniesta R, *et al*.
Polygenic interactions with environmental adversity in the aetiology of major
depressive disorder. *Psychol Med* 2016; **46**(4): 759-770.

44.   Colodro-Conde L, Couvy-Duchesne B, Zhu G, Coventry WL, Byrne EM, Gordon S, *et
al*. A direct test of the diathesis-stress model for depression. *Mol Psychiatry* 2017.

45.   Kendler KS, Baker JH. Genetic influences on measures of the environment: a
systematic review. *Psychol Med* 2007; **37**(5): 615-626.

46.   Luciano M, Batty GD, McGilchrist M, Linksted P, Fitzpatrick B, Jackson C, *et al*.
Shared genetic aetiology between cognitive ability and cardiovascular disease risk
factors: Generation Scotland's Scottish family health study. *Intelligence* 2010; **38**(3):
304-313.

47.   Silberg J, Rutter M, Neale M, Eaves L. Genetic moderation of environmental risk
for depression and anxiety in adolescent girls. *Br J Psychiatry* 2001; **179**: 116-121.

48.   Kendler KS, Kessler RC, Walters EE, MacLean C, Neale MC, Heath AC, *et al*.
Stressful life events, genetic liability, and onset of an episode of major depression in
women. *Am J Psychiatry* 1995; **152**(6): 833-842.

49.   Biobank U (2010). UK Biobank: protocol for a large-scale prospective
epidemiological resource.

670
671   50.   Biobank U (2007). UK Biobank ethics and governance framework, version 3.0

672
673   51.   Biobank U. Genotype imputation and genetic association studies of UK Biobank.
674   2015.

675
676   52.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, *et al.* PLINK:
677   a tool set for whole-genome association and population-based linkage analyses. *Am J*
678   *Hum Genet* 2007; **81**(3): 559-575.

679
680   53.   Amador C, Huffman J, Trochet H, Campbell A, Porteous D, Generation S, *et al.*
681   Recent genomic heritage in Scotland. *BMC Genomics* 2015; **16**: 437.

682
683   54.   Smith BH, Campbell A, Linksted P, Fitzpatrick B, Jackson C, Kerr SM, *et al.* Cohort
684   Profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its
685   participants and their potential for genetic research on health and illness. *Int J Epidemiol*
686   2013; **42**(3): 689-700.

687
688   55.   Fernandez-Pujals AM, Adams MJ, Thomson P, McKechanie AG, Blackwood DH,
689   Smith BH, *et al.* Epidemiology and Heritability of Major Depressive Disorder, Stratified
690   by Age of Onset, Sex, and Illness Course in Generation Scotland: Scottish Family Health
691   Study (GS:SFHS). *PLoS One* 2015; **10**(11): e0142197.

692
693   56.   Smith BH, Campbell H, Blackwood D, Connell J, Connor M, Deary IJ, *et al.*
694   Generation Scotland: the Scottish Family Health Study; a new resource for researching
695   genes and heritability. *BMC medical genetics* 2006; **7**: 74.

696
697   57.   Kerr SM, Campbell A, Murphy L, Hayward C, Jackson C, Wain LV, *et al.* Pedigree
698   and genotyping quality analyses of over 10,000 DNA samples from the Generation
699   Scotland: Scottish Family Health Study. *BMC Med Genet* 2013; **14**: 38.

700
701   58.   Biobank U (2012). Touchscreen questionnaire.

702
703   59.   Eysenck HJ ES (1975). Manual of the Eysenck Personality Questionnaire. Hodder
704   and Stoughton: London.

705
706   60.   Kendler KS, Myers J. The genetic and environmental relationship between major
707   depression and the five-factor model of personality. *Psychol Med* 2010; **40**(5): 801-806.

708
709   61.   Smith DJ, Nicholl BI, Cullen B, Martin D, Ul-Haq Z, Evans J, *et al.* Prevalence and
710   characteristics of probable major depression and bipolar disorder within UK biobank:
711   cross-sectional study of 172,751 participants. *PLoS One* 2013; **8**(11): e75362.

712
62.   Eysenck SBG, Eysenck, H.J. & Barrett, P. A revised version of the psychoticism
scale. *Personality and Individual Differences* 1985; **6**: 21-29.

63.   Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version
of PRIME-MD: the PHQ primary care study. Primary Care Evaluation of Mental
Disorders. Patient Health Questionnaire. *JAMA* 1999; **282**(18): 1737-1744.

64.   First MB SR, Gibbon M, Williams JB (2002). Structured Clinical Interview for DSM-
IV-TR Axis I Disorders. New York State Psychiatric Institute: New York.

65.   Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software.
Bioinformatics 2015; **31**(9): 1466-1468.

66.   Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of
genomewide association scans. *Bioinformatics* 2010; **26**(17): 2190-2191.

67.   de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set
analysis of GWAS data. *PLoS Comput Biol* 2015; **11**(4): e1004219.

68.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. FUMA: Functional
mapping and annotation of genetic associations. *bioRxiv* 2017.

69.   Bulik-Sullivan BK, Loh PR, Finucane HK, Ripke S, Yang J, Schizophrenia Working
Group of the Psychiatric Genomics C, *et al.* LD Score regression distinguishes
confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015;
47(3): 291-295.

70.   Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh PR, *et al.* An atlas of
genetic correlations across human diseases and traits. *Nat Genet* 2015; **47**(11): 1236-
1241.

71.   Pers TH, Karjalainen JM, Chan Y, Westra HJ, Wood AR, Yang J, *et al.* Biological
interpretation of genome-wide association studies using predicted gene functions. *Nat*
Commun 2015; **6**: 5890.

72.   Mailman MD, Feolo M, Jin Y, Kimura M, Tryka K, Bagoutdinov R, *et al.* The NCBI
dbGaP database of genotypes and phenotypes. *Nat Genet* 2007; **39**(10): 1181-1186.

73.   Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, *et al.*
Annotation of functional variation in personal genomes using RegulomeDB. *Genome*
research 2012; **22**(9): 1790-1797.

74.   Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot
analysis: multitissue gene regulation in humans. *Science* 2015; **348**(6235): 648-660.

75.   Genetics of Personality C, de Moor MH, van den Berg SM, Verweij KJ, Krueger RF,
Luciano M, *et al.* Meta-analysis of Genome-wide Association Studies for Neuroticism, and

the Polygenic Association With Major Depressive Disorder. *JAMA Psychiatry* 2015;
72(7): 642-650.

762
763   76.   Team RC. *R: A language and environment for statistical computing.* R Foundation
764   for Statistical Computing: Vienna, Austria, 2015.

765
766   77.   Lee SH, Wray NR, Goddard ME, Visscher PM. Estimating missing heritability for
767   disease from genome-wide association studies. *Am J Hum Genet* 2011; **88**(3): 294-305.

768
769   78.   Chen GB. Estimating heritability of complex traits from genome-wide association
770   studies using IBS-based Haseman-Elston regression. *Front Genet* 2014; **5**: 107.

771
772   79.   Hirschfeld RM, Williams JB, Spitzer RL, Calabrese JR, Flynn L, Keck PE, Jr., *et al.*
773   Development and validation of a screening instrument for bipolar spectrum disorder:
774   the Mood Disorder Questionnaire. *Am J Psychiatry* 2000; **157**(11): 1873-1875.

775
776   80.   Raine A. The SPQ: a scale for the assessment of schizotypal personality based on
777   DSM-III-R criteria. *Schizophr Bull* 1991; **17**(4): 555-564.

778
779   81.   Polderman TJ, Benyamin B, de Leeuw CA, Sullivan PF, van Bochoven A, Visscher
780   PM, *et al.* Meta-analysis of the heritability of human traits based on fifty years of twin
781   studies. *Nat Genet* 2015; **47**(7): 702-709.

782
783   82.   consortium C. Sparse whole-genome sequencing identifies two loci for major
784   depressive disorder. *Nature* 2015; **523**(7562): 588-591.

785
786   83.   Hyde CL, Nagle MW, Tian C, Chen X, Paciga SA, Wendland JR, *et al.* Identification of
787   15 genetic loci associated with risk of major depression in individuals of European
788   descent. *Nat Genet* 2016; **48**(9): 1031-1036.

789
790   84.   Zuo L, Tan Y, Zhang X, Wang X, Krystal J, Tabakoff B, *et al.* A New Genomewide
791   Association Meta-Analysis of Alcohol Dependence. *Alcohol Clin Exp Res* 2015; **39**(8):
792   1388-1395.

793
794   85.   Cook MN, Baker JA, Heldt SA, Williams RW, Hamre KM, Lu L. Identification of
795   candidate genes that underlie the QTL on chromosome 1 that mediates genetic
796   differences in stress-ethanol interactions. *Physiol Genomics* 2015; **47**(8): 308-317.

797
798   86.   Zuo L, Wang K, Wang G, Pan X, Zhang X, Zhang H, *et al.* Common PTP4A1-PHF3-
799   EYS variants are specific for alcohol dependence. *Am J Addict* 2014; **23**(4): 411-414.

800
801   87.   Zuo L, Zhang X, Deng HW, Luo X. Association of rare PTP4A1-PHF3-EYS variants
802   with alcohol dependence. *J Hum Genet* 2013; **58**(3): 178-179.

804   88.   Zuo L, Zhang CK, Wang F, Li CS, Zhao H, Lu L, *et al.* A novel, functional and
805   replicable risk gene region for alcohol dependence identified by genome-wide
806   association study. *PLoS One* 2011; **6**(11): e26726.

807
808   89.   Hontelez S, Karthaus N, Looman MW, Ansems M, Adema GJ. DC-SCRIPT regulates
809   glucocorticoid receptor function and expression of its target GILZ in dendritic cells. *J*
810   *Immunol* 2013; **190**(7): 3172-3179.

811
812   90.   Lopez-Garcia J, Periyasamy M, Thomas RS, Christian M, Leao M, Jat P, *et al.*
813   ZNF366 is an estrogen receptor corepressor that acts through CtBP and histone
814   deacetylases. *Nucleic Acids Res* 2006; **34**(21): 6126-6136.

815
816   91.   Miyamoto-Sato E, Fujimori S, Ishizaka M, Hirai N, Masuoka K, Saito R, *et al.* A
817   comprehensive resource of interacting protein regions for refining human transcription
818   factor networks. *PLoS One* 2010; **5**(2): e9289.

819
820   92.   Aragam N, Wang KS, Pan Y. Genome-wide association analysis of gender
821   differences in major depressive disorder in the Netherlands NESDA and NTR
822   population-based samples. *J Affect Disord* 2011; **133**(3): 516-521.

823
824   93.   Fidalgo TM, da Silveira ED, da Silveira DX. Psychiatric comorbidity related to
825   alcohol use among adolescents. *Am J Drug Alcohol Abuse* 2008; **34**(1): 83-89.

826
827   94.   Lopez B, Turner RJ, Saavedra LM. Anxiety and risk for substance dependence
828   among late adolescents/young adults. *J Anxiety Disord* 2005; **19**(3): 275-294.

829
830   95.   Low NC, Lee SS, Johnson JG, Williams JB, Harris ES. The association between
831   anxiety and alcohol versus cannabis abuse disorders among adolescents in primary care
832   settings. *Fam Pract* 2008; **25**(5): 321-327.

833
834   96.   Rutledge PC, Sher KJ. Heavy drinking from the freshman year into early young
835   adulthood: the roles of stress, tension-reduction drinking motives, gender and
836   personality. *J Stud Alcohol* 2001; **62**(4): 457-466.

837
838   97.   Schmidt NB, Buckner JD, Keough ME. Anxiety sensitivity as a prospective
839   predictor of alcohol use disorders. *Behav Modif* 2007; **31**(2): 202-219.

840
841   98.   Varlinskaya EI, Kim EU, Spear LP. Chronic intermittent ethanol exposure during
842   adolescence: Effects on stress-induced social alterations and social drinking in
843   adulthood. *Brain Res* 2017; **1654**(Pt B): 145-156.

844
845   99.   Torigata K, Daisuke O, Mukai S, Hatanaka A, Ohka F, Motooka D, *et al.* LATS2
846   Positively Regulates Polycomb Repressive Complex 2. *PLoS One* 2016; **11**(7): e0158562.

847
848   100.   Shen EY, Jiang Y, Javidfar B, Kassim B, Loh YE, Ma Q, *et al.* Neuronal Deletion of
849   Kmt2a/Mll1 Histone Methyltransferase in Ventral Striatum is Associated with Defective

850   Spike-Timing-Dependent Striatal Synaptic Plasticity, Altered Response to Dopaminergic
851   Drugs, and Increased Anxiety. *Neuropsychopharmacology* 2016; **41**(13): 3103-3113.
852

101.   Wong ML, Arcos-Burgos M, Liu S, Velez JI, Yu C, Baune BT, *et al.* The PHF21B gene
is associated with major depression and modulates the stress response. *Mol Psychiatry*
2017; **22**(7): 1015-1025.

102.   Walsh RM, Shen EY, Bagot RC, Anselmo A, Jiang Y, Javidfar B, *et al.* Phf8 loss
confers resistance to depression-like and anxiety-like behaviors in mice. *Nat Commun*
2017; **8**: 15142.

103.   Gage SH, Jones HJ, Taylor AE, Burgess S, Zammit S, Munafo MR. Investigating
causality in associations between smoking initiation and schizophrenia using Mendelian
randomization. *Sci Rep* 2017; **7**: 40653.

104.   Hartz SM, Horton AC, Hancock DB, Baker TB, Caporaso NE, Chen LS, *et al.* Genetic
correlation between smoking behaviors and schizophrenia. *Schizophr Res* 2017.

105.   Dudbridge F. Power and predictive accuracy of polygenic risk scores. *PLoS Genet*
2013; **9**(3): e1003348.

FIGURE LEGENDS

Figure 1. **Manhattan plots showing stress-sensitivity associations.** Manhattan plots

of the GWIS from **A)** UKB, **B)** GS:SFHS and **C)** sample size weighted meta-analysis of UKB

and GS:SFHS. The x-axis is chromosomal position and y-axis is the p value (-log10 p

value) of association with stress-sensitivity effect. Suggestive genome-wide significance

threshold ($p = 1x10^{-5}$) is shown by solid line at y = 5. Genes or closest gene up- and

down-stream from SNP position (/) are annotated. "-": No gene within 100kb of the SNP.

Figure 2. Miami plots showing comparison between association profile between

stress-sensitivity GWIS and MDD GWAS. Miami plots from meta-analysis filter at $p =$

$1x10^{-3}$: **A)** filtering for stress-sensitivity p values (•), **B)** filtering for MDD p values (×).

The x-axis is chromosomal position and y-axis is the p value (-log10 p value) of

association with stress-sensitivity (up; red dots) and MDD p value (down; blue crosses).

Dot line: genome-wide suggestive threshold ($p = 1x10^{-5}$) at the filtered effect; dashed

lines: $p = 0.01$ and 0.05 at unfiltered effect.

Figure 3. MDD is best predicted using multiple PRS. MDD risk explained (R^2

coefficient (%); top bar values) on the liability scale by each PRS; weighted by GWAS

main additive and GWIS stress-sensitivity effects independently and combined. **A)**

GS:SFHS: there is an increment on MDD risk prediction from adding PRS_{SS} to PRS_D model

of 64.3% and 35.8% when adding PRS_{SS} to both MDD and neuroticism PRS together. **B)**

UKB: there is an increment on MDD risk prediction from adding PRS_{SS} to PRS_D model of

17.1% and 11.2% when combining PRS_{SS} with both MDD and neuroticism PRS. **C & D)**

Replication of A & B fitting PRS_D and PRS_N using summary statistics from worldwide

consortiums (i.e. PGC &GPC). Significance codes: p values *** < 0.001 < ** < 0.01 < * <

0.05; derived from likelihood ratio tests. SS stands for stress-sensitivity.

Table 1. Top 25 SNPs from meta-analysis of GWISs.

Rank	CHR	SNP	BP	A1	Z-score	Effect	p	p (EPQN)	p (MDD)	GENE	POSITION
1	6	rs319924	64487247	A	5.024	++	5.05×10^{-7}	0.376	0.637	EYS	Intronic
2	5	rs246565	71809247	A	-4.684	--	2.82×10^{-6}	0.248	0.589	ZNF366	5998bp 5'
3	10	rs2265265	24854876	A	4.604	++	4.15×10^{-6}	0.035	0.084	KIAA1217 / ARHGAP21	18104bp 3' / 17662bp 3'
4	6	rs1057530	64427095	A	-4.556	--	5.21×10^{-6}	0.636	0.840	PHF3/EYS	1677bp 3' / 2781bp 3'
5	16	rs7199110	78790765	A	-4.553	--	5.29×10^{-6}	0.661	0.741	WWOX	Intronic
6	6	rs10485358	64386060	A	-4.546	--	5.46×10^{-6}	0.390	0.902	PHF3	Intronic
7	12	rs10778077	101193988	A	4.54	++	5.62×10^{-6}	0.614	0.430	ANO4	Intronic
8	5	rs13358894	71803446	A	4.527	++	5.99×10^{-6}	0.257	0.651	ZNF366	197bp 5'
9	10	rs2256220	24856314	A	-4.524	--	6.06×10^{-6}	0.134	0.129	KIAA1217 / ARHGAP21	19542bp 3' / 16224bp 3'
10	10	rs3762096	98136250	A	-4.521	--	6.15×10^{-6}	0.437	0.149	TLL2	Intronic
11	11	rs2221540	132716369	A	-4.492	--	7.05×10^{-6}	0.468	0.364	OPCML	Intronic
12	5	rs10043659	71781839	A	4.483	++	7.37×10^{-6}	0.339	0.808	ZNF366	Intronic
13	12	rs10778078	101195088	A	-4.45	--	8.58×10^{-6}	0.599	0.456	ANO4	Intronic
14	9	rs10987199	128968987	A	-4.442	--	8.91×10^{-6}	0.199	0.026	LOC101929116	63416bp 3'
15	5	rs10042132	71789021	A	-4.416	--	1.01×10^{-5}	0.418	0.538	ZNF366	Intronic
16	11	rs10894606	132671611	A	-4.404	--	1.06×10^{-5}	0.438	0.587	OPCML	Intronic
17	12	rs7295089	2440464	A	4.372	++	1.23×10^{-5}	0.266	0.212	CACNA1C	Intronic
18	5	rs9293292	71696942	A	-4.351	--	1.36×10^{-5}	0.126	0.731	PTCD2/ZNF366	41762bp 3' / 42292bp 3'
19	15	rs3097437	27872136	A	4.346	++	1.38×10^{-5}	0.970	0.226	GABRG3	93762bp 3'
20	9	rs1999377	11919732	A	4.344	++	1.40×10^{-5}	0.436	0.064	-	Intragenic
21	5	rs6862221	71754962	A	4.342	++	1.41×10^{-5}	0.543	0.823	ZNF366	Intronic
22	5	rs9293289	71683885	A	-4.323	--	1.54×10^{-5}	0.395	0.510	PTCD2/ZNF366	28705bp 3' / 55349bp 3'
23	11	rs4575282	132719646	A	-4.313	--	1.61×10^{-5}	0.598	0.514	OPCML	Intronic
24	9	rs2417008	128970219	A	-4.3	--	1.71×10^{-5}	0.208	0.026	LOC101929116	62184bp 3'
25	9	rs7021461	128972210	A	4.299	++	1.72×10^{-5}	0.202	0.025	LOC101929116	60193bp 3'

Effect shows effect direction in GS:SFHS and UK Biobank, *p* reflects significance of GWIS stress-sensitivity effect, *p (EPQN)* SNP main effect on neuroticism derived from GWAS meta-analysis of EPQN between UK Biobank and Generation Scotland. *p (MDD)* SNP main effect on MDD derived from GWAS meta-analysis of MDD between UK Biobank and Generation Scotland. *Position* position of the SNP respect to closest gene transcripts within 100kb (including UTRs) from 5 prime (5') or 3prime (3').

Table 2. MDD risk prediction at best-fits.

UKB predicting on GS:SFHS

Weighted effect	Best-fit threshold	# SNPs	R^2 (%)[a]	R^2 (%)[b]	*p*
stress-sensitivity	0.004	1 313	0.175	0.243	1.8×10^{-3}
MDD	0.125	25 833	0.278	0.386	8.3×10^{-5}
EPQN	1×10^{-3}	863	0.245	0.340	2.2×10^{-4}
MDD + EPQN	-	-	0.492	0.684	1.1×10^{-6}
joint models	-	-	0.669	0.928	3.9×10^{-8}

PGC2 & GPC predicting on GS:SFHS

PGC2 MDD	0.227	41 917	0.888	1.233	1.9×10^{-12}
GPC EPQN	0.013	4 234	0.101	0.140	0.018
PGC2 MDD + GPC EPQN	-	-	0.175	1.324	2.7×10^{-12}
joint models	-	-	1.138	1.580	9.7×10^{-14}

GS:SFHS predicting on UKB

Weighted effect	Best-fit threshold	# SNPs	R^2 (%)[a]	R^2 (%)[b]	*p*
stress-sensitivity	1.15×10^{-4}	42	0.024	0.037	0.04
MDD	0.033	8 135	0.135	0.211	1.1×10^{-6}
EPQN	0.07	15 613	0.114	0.178	7.3×10^{-6}
MDD + EPQN	-	-	0.206	0.323	1.3×10^{-8}
joint models	-	-	0.229	0.359	8.7×10^{-9}

PGC2 & GPC predicting on UKB

PGC2 MDD	0.095	22 005	0.818	1.281	2.4×10^{-33}
GPC EPQN	6×10^{-3}	2 182	0.067	0.105	5.6×10^{-4}
PGC2 MDD + GPC EPQN	-	-	0.863	1.352	6.6×10^{-34}
joint models	-	-	0.886	1.388	1.8×10^{-32}

stress-sensitivity stress-sensitivity, *MDD* major depressive disorder, *EPQN* neuroticism score, *joint models* combined effect fitting all 3 PRS weighted by all the effects (i.e. stress-sensitivity, MDD and EPQN), [a]Nagelkerke's R^2 at observed scale, [b]R^2 on the liability scale

900   **Figure 1**
901

902 **Figure 2**

904   **Figure 3**
905