Fennec - Functional Exploration of Natural Networks and Ecological Communities

Markus J. Ankenbrand[1,2], Sonja Hohlfeld[2],

Lorenz Weber[3], Frank Förster[1,2,4], Alexander Keller[1,2,5]

April 24, 2018

[1]Department of Bioinformatics, University of Würzburg, Würzburg, Germany

[2]Center for Computational and Theoretical Biology, University of Würzburg, Würzburg, Germany

[3]Mayflower GmbH, Würzburg, Germany

[4]Fraunhofer Institute for Molecular Biology and Applied Ecology IME, Applied Ecology and Bioresources, Gießen, Germany

[5]Department of Animal Ecology and Tropical Biology, University of Würzburg, Würzburg, Germany

Abstract

Species composition assessment of ecological communities and networks is an important aspect of biodiversity research. Yet often ecological traits of organisms in a community are more informative than scientific names only. Furthermore, other properties like threat status, invasiveness, or human usage are relevant for many studies, but cannot be evaluated from taxonomy alone. Despite public

databases collecting such information, it is still a tedious manual task to enrich community analyses with such, especially for large-scaled data.

Here we present the FENNEC, a web-based workbench that eases the process of mapping publicly available trait data to the user's communities in an automated process. Usage is either by a local self-hosted or a public instance (https://fennec.molecular.eco) covering exemplary traits. The FENNEC aims to motivate public trait data submission and its reuse in meta-analyses. Further, it is an open-source development project with the code freely available to use and community contribution (https://github.com/molbiodiv/fennec).

Keywords

Traits, Community, Ecology, Functional, Database, Software

Introduction

An important task in biodiversity research is the analysis of species composition of ecological communities and networks. This can be done using traditional methods and more recently also with analytical methods designed for large scale sample processing, like DNA metabarcoding (Keller, Danner, et al., 2015) or automated image analysis (Oteros et al., 2015) producing data volumes hard to cope with manually. Therefore, tools for automated taxonomic identification have been developed, e.g. QIIME (Caporaso et al., 2010) and mothur (Schloss et al., 2009).

Ecological or socio-economical hypotheses can however often not be evaluated by looking at taxonomy alone (Junker et al., 2015; Xu et al., 2014), but meta-data

for each community-member (e.g. life-history, size or threat-status) becomes important. In microbial ecology, development of tools has already been initiated that aim to automatically map taxonomy information to functional traits, mostly through mapping on known genomes (Aßhauer et al., 2015; Edgar, 2017; Keller, Horn, et al., 2014; Langille et al., 2013). To our knowledge, it remains to date a manual effort to enrich eukaryotic communities similarly with trait meta-data, although such information is already publicly available.

Databases have been developed that provide trait information for eukaryotes and prokaryotes, e.g. LEDA Traitbase (Kleyer et al., 2008), TRY (Kattge et al., 2011), and BacDive or ProTraits for microbial traits (Brbić et al., 2016; Söhngen et al., 2016). On a higher level, TraitBank (Parr, Wilson, Leary, et al., 2014; Parr, Wilson, Schulz, et al., 2014) aggregates this information from different sources. These sources are of course far from complete, yet the existing data is already highly informative.

Currently these trait data have to be manually searched and mapped to the community, which can comprise hundreds or even thousands of taxa. This is impractical, trait data should be accessible with automatic batch annotation procedures, not only single manual requests. Furthermore, tools for visualization and interactive analysis of community data like Phinch and phyloseq are currently limited with respect to traits as they only accept taxonomy as OTU metadata (Bik et al., 2014; McMurdie et al., 2013). Here we present the Fennec, a web-based workbench that helps researchers enrich their taxonomy-based community and network interaction tables with relevant traits for their research questions. The workbench includes also basic tools for interactive visualization and analysis of trait data.

A public instance is hosted at https://fennec.molecular.eco, which currently holds roughly 1,700,000 organisms and 224,000 trait entries (referring to 73,500

distinct taxa) gathered from various sources. User-provided community and network data can be readily mapped to these traits.

An alternative option to use the FENNEC is to download and host a local instance for user and private traits alongside the ones in the public database. It is freely available at https://github.com/molbiodiv/fennec as an open-source project and we encourage community development and re-use of the code.

Online documentation is available on how to use the public and set up own instances (https://fennec.readthedocs.io). This documentation also includes a case study in pollination ecology to demonstrate how the FENNEC can be used to invastigate specialization of solitary bees on plants with specific growth habit, distribution of invasive taxa as well as occurence of endangered organisms.

Material and Methods

The FENNEC is a web application developed in PHP (https://php.net/) using Symfony (https://symfony.com/) and a JavaScript (ES6) front-end. Server side functionality is bundled in modular web services called via AJAX requests. Layout and interactivity are provided by multiple well-established libraries including bootstrap (https://getbootstrap.com/), jQuery-ui (https://jqueryui.com/), react (https://facebook.github.io/react/), lodash (https://lodash.com/), datatables (https://datatables.net/), and plotly.js (https://plot.ly/javascript/). Code quality is ensured by tests and strategies for continuous integration. Data is stored in a PostgreSQL (https://www.postgresql.org/) database and accessed via the doctrine object-relational mapper (http://www.doctrine-project.org/). User provided data is uploaded and stored in BIOM format (version 1.0) (McDonald et al., 2012) using the biojs-io-biom library (Ankenbrand et al., 2017).

Results

Accessibility

There are three ways to use the FENNEC workbench:

1. Public Instance: We have set up a public instance of the FENNEC available at https://fennec.molecular.eco. Its database currently hosts trait data related to pollination and microbiomes from various sources. The database is subject to constant further extension with more traits, yet our main goal is to maintain high quality of the data available here. User-provided network and community data is private per default, requiring the user to authenticate using a FENNEC or GitHub account.

2. Local Instances: All program code is open-source (MIT License) and freely available at the public repository: https://github.com/molbiodiv/fennec. Alongside the code, ready-to-use (pre-configured) docker containers are available to be run in a virtualization environment (https://www.docker. com). Local instances can be restricted in accessibility for dedicated workgroups or users. Databases can be filled directly with arbitrary trait data not limited to those included in the public instance.

3. Application Programming Interface (API): We also provide an open API that allows third-party programs to make calls to the public instance, or if available also local instances.

Extensive documentation on the code, but also tutorials for users and guides for administrators to host local instances and software developers to use the API are available at the GitHub repository, the public instance and https: //fennec.readthedocs.io.

Figure 1: General structure of FENNEC. Organisms and traits from different sources are stored by the administrator. The user imports a community project e.g. in biom format. Organisms in the community are mapped against those in the database and enriched with traits. The trait composition can be interactively explored or exported as enriched projects for downstream software.

Mapping community and network data to traits

Minimum requirement for using the FENNEC is to provide a community or network table, including taxa as rows and samples (communities)/ taxa (networks) as columns. Cells are considered as abundances but also presence/absence data can be used. Beyond this, users may provide own taxonomy data as an alternative to the default NCBI taxonomy (Federhen, 2012)) and meta-data for the samples. These tables may be uploaded separately as tab-separated text- or combined BIOM-files (Ankenbrand et al., 2017; McDonald et al., 2012) (figure 1). Files can be managed using the project page of FENNEC (figure 2B). Depending on the user input, taxa are mapped using scientific names or common database identifiers (e.g. NCBI-taxonomy-ID, EOL-ID).

Traits to be analyzed can be explored and selected via the web interface, and added as meta-data to the project (figure 2A). All data available for the selected

traits and taxa of interest are automatically linked into the dataset. If multiple values are available for a single trait and organism combination, they are automatically aggregated, i.e. categorical traits are unified and numerical traits are averaged. To make trait usage as transparent and flexible as possible and to facilitate proper attribution along with aggregated trait values, trait citations for individual values are provided alongside actual traits. Citations can be exported as a separate table and are included in any downloaded BIOM file.

After the mapping, communities are enriched with selected meta-data and can be further processed with standard analytical and statistical software. Download options are individual tables or as a single BIOM file including all information (e.g. for fast integration in R with phyloseq (McMurdie et al., 2013) or QIIME (Caporaso et al., 2010)).

To provide basic analytical plots directly in the workbench, we integrated and modified the open-source project Phinch (Bik et al., 2014). This allows quick interactive exploration of species and trait distributions in each sample, groups, or aggregated by trait types (figure 2D).

Data for the public instance

A public instance of FENNEC is hosted at https://fennec.molecular.eco and freely available for direct usage. Taxonomy data in this instance consists of a full representation of the NCBI Taxonomy database (Federhen, 2012 accessed 21/03/2018, >1.7 million taxa). Currently only a small fraction (about 73,500) of the taxa in the database have associated traits. We aim to extend this set of traits but also allow users to contribute traits to the general public database. A mapping of EOL-IDs (according to https://opendata.eol.org/dataset/hierarchy_entries, accessed on 04/04/2017) has also been imported, so that full-text information

about taxa is available where EOL offers such (Parr, Wilson, Leary, et al., 2014). Currently and as a starting seed, trait data from TraitBank (Parr, Wilson, Schulz, et al., 2014), EPPO (EPPO, 2017), the World Crops Database (Bijlmakers, 2017), the cavety-nesting bees and wasps database (Budrys et al., 2014, part of the SCALES project (Henle et al., 2014)), IUCN (IUCN, 2017), and ProTraits (Brbić et al., 2016) have been imported which is subject to continuous extension. We aim to maintain high-quality of these publicly available traits, so that the integration of more traits is a steadily ongoing process. While the bulk of trait data is gathered from databases, in the next release users can also participate in the uploading of trait data, so that this process can be actively supported by the community.

Importing organism and trait data

Traits are imported using a simple table containing mandatory the organism and trait value, a citation and optionally ontology URL as columns, and each entry as a new row. Two trait formats are currently supported: categorical and numerical. Categorical traits may also include an ontology URL for their value, supporting the hierarchical classification characteristics. Numerical trait types may be uploaded with an associated unit.

Currently users can upload own traits as project-specific meta-data or in local instances for everybody accessible if allowed by the administrator. For the public instance we plan a moderation system for a future release. All data will undergo a limited manual verification, which for example ensures that units are correctly standardized, but does not verify the correctness of the underlying data. Therefore, the user-names of uploaders will be permanently linked to the data to be able to address future changes and updates, alongside corresponding

citation information.

For local instances, arbitrary organism and trait data can be imported into the FENNEC database by the local administrator using the command line interface (CLI), which are not subject to a central verification process, but available instantly. This allows creation of instances tailored to specific organism groups and associated research questions, with responsibility for local administrators to ensure quality. Either the global NCBI-taxonomy data can be used or custom taxonomy data provided. Each imported organism receives a unique FENNEC-ID which can be linked on-the-fly to other identifiers like NCBI-Taxonomy-IDs or EOL-IDs. The linked EOL-ID is used to provide dynamic content for each organism using the EOL-API (figure 2C).

Discussion

The FENNEC is a useful tool for automated mapping from taxonomic data to functional meta-data of whole communities. This can be done with user-supplied traits or such data-mined from trait databases. A growing public instance is available for analyses in pollination studies. It can be accessed via a graphical web interface or by calls via an API. Local instances can be tailored to specific organism groups, workgroups and research questions. The workbench provides basic visualization options for mapped data, as well as export options in various formats for use with downstream software.

Outlook and limitations

The main factor restricting FENNECs utility is the currently limited amount of trait data available in a usable format. It became apparent while building the

Figure 2: User interface of Fennec. A: Explore traits stored in the database via the search. B: Upload community data, manage and analyze them. C: Get dynamic content for organisms. D: Visualize data using the "Donut Partition Chart".

FENNEC that a lot of trait data is available online, but the majority does not adhere to the FAIR principles (Findable, Accessible, Interoperable, Reusable) (Mons et al., 2017; Wilkinson et al., 2016). Also licensing of the data is a common problem, data can only be efficiently re-used if it is open and citable in addition to being FAIR (Katz, 2017). It is essential to guarantee trait data collectors that data re-users are able to properly cite all data sources, e.g. by adhering to the FORCE11 data citation principles (Martone, 2014). FENNEC supports this by preserving all relevant information. We therefore encourage trait data collectors to make their data available via existing platforms like TraitBank (Parr, Wilson, Schulz, et al., 2014), and with that also usable for downstream analysis tools like FENNEC and ultimately available to the whole research community.

Conclusion

FENNEC as a tool provides valuable assistance to analyze ecological data with more than just taxonomy. Species traits and meta-data like threat status and economic importance help to answer questions beyond the mere presence or absence of organisms. The public instance can be used as a reference, to try features of FENNEC and analyze datasets with data from other public databases. Self-tailored local instances with private data increases the range of applications.

Current main limitation is trait data availability, but we expect an increase with time, ongoing open-science initiatives and scientific journals requiring public data deposition before publication. Beside developing a public automatic mapping procedure, we also aim to demonstrate the importance of making trait data publicly available and its usefulness in follow up studies. We thus advocate the submission of trait data to public databases.

Despite limitations in data availability, the FENNEC is technically already able

to enrich ecological community analyses with trait information. Its usefulness
is expected to increase due to continued development guided by user feedback,
integration of more analysis tools, better taxonomic resolution, and increasing
availability of suitable trait data.

Acknowledgements

This study was supported by the German Excellence Initiative to the Graduate
School of Life Sciences, University of Würzburg (405/1517) to MJA and the DFG
(KE1743/7-1) to AK. We are grateful to the EOL for quick, open and helpful
responses to our emails. Many thanks to A. Voulgari-Kokota, N. Terhoeven,
J. Schultz and A. Claßen for beta testing, suggestions and valuable feedback.

Author Contributions

MJA and AK conceived the project, as well as designed methodology and
software; MJA and SH wrote the code with support by FF and LW; MJA
analyzed the case study and drafted the manuscript. All authors contributed
critically to the drafts and gave final approval for publication.

References

Ankenbrand, M. J. et al. (2017). "biojs-io-biom, a BioJS component for handling
data in Biological Observation Matrix (BIOM) format". In: *F1000Research* 5,
p. 2348. DOI: 10.12688/f1000research.9618.2.

Aßhauer, K. P. et al. (2015). "Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data". In: *Bioinformatics* 31.17, pp. 2882–2884. DOI: 10.1093/bioinformatics/btv287.

Bijlmakers, H. (2017). *World Crops Database - Fruits, vegetables, cereals and other agricultural crops.* World Crops Database. URL: http://world-crops.com/ (visited on 07/25/2017).

Bik, H. M. and P. Interactive (2014). "Phinch: An interactive, exploratory data visualization framework for –Omic datasets". In: *bioRxiv*, p. 009944. DOI: 10.1101/009944.

Brbić, M. et al. (2016). "The landscape of microbial phenotypic traits and associated genes". In: *Nucleic Acids Research* 44.21, pp. 10074–10090. DOI: 10.1093/nar/gkw964. eprint: /oup/backfile/content_public/journal/nar/44/21/10.1093_nar_gkw964/3/gkw964.pdf.

Budrys, E., A. Budriene, and S. Orlovskyte (2014). *Cavity-nesting wasps and bees database.* URL: http://scales.ckff.si/scaletool/?menu=6&submenu=3 (visited on 09/22/2017).

Caporaso, J. G. et al. (2010). "QIIME allows analysis of high-throughput community sequencing data". In: *Nature Methods* 7.5, pp. 335–336. DOI: 10.1038/nmeth.f.303.

Edgar, R. C. (2017). "SINAPS: Prediction of microbial traits from marker gene sequences". In: *bioRxiv*. DOI: 10.1101/124156. eprint: https://www.biorxiv.org/content/early/2017/04/04/124156.full.pdf.

EPPO (2017). *EPPO Global Database (available online).* URL: https://gd.eppo.int/ (visited on 09/22/2017).

Federhen, S. (2012). "The NCBI Taxonomy database". In: *Nucleic Acids Research* 40 (D1), pp. D136–D143. DOI: 10.1093/nar/gkr1178.

Henle, K. et al. (2014). "Scaling in Ecology and Biodiversity Conservation". In: *Advanced Books* 1, e1169. DOI: 10.3897/ab.e1169.

IUCN (2017). *IUCN Red List of Threatened Species. Version 2017-1*. URL: http://www.iucnredlist.org/ (visited on 09/22/2017).

Junker, R. R., N. Blüthgen, and A. Keller (2015). "Functional and phylogenetic diversity of plant communities differently affect the structure of flower-visitor interactions and reveal convergences in floral traits". In: *Evolutionary Ecology* 29.3, pp. 437–450. DOI: 10.1007/s10682-014-9747-2.

Kattge, J. et al. (2011). "TRY – a global database of plant traits". In: *Global Change Biology* 17.9, pp. 2905–2935. DOI: 10.1111/j.1365-2486.2011.02451.x.

Katz, D. S. (2017). *FAIR is not fair enough*. Daniel S. Katz's blog. URL: https://danielskatzblog.wordpress.com/2017/06/22/fair-is-not-fair-enough/ (visited on 09/13/2017).

Keller, A., N. Danner, et al. (2015). "Evaluating multiplexed next-generation sequencing as a method in palynology for mixed pollen samples". In: *Plant Biology* 17.2, pp. 558–566. DOI: 10.1111/plb.12251.

Keller, A., H. Horn, et al. (2014). "Computational integration of genomic traits into 16S rDNA microbiota sequencing studies". In: *Gene* 549.1, pp. 186–191. DOI: 10.1016/j.gene.2014.07.066.

Kleyer, M. et al. (2008). "The LEDA Traitbase: a database of life-history traits of the Northwest European flora". In: *Journal of Ecology* 96.6, pp. 1266–1274. DOI: 10.1111/j.1365-2745.2008.01430.x.

Langille, M. G. I. et al. (2013). "Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences". In: *Nature Biotechnology* 31.9, pp. 814–821. DOI: 10.1038/nbt.2676.

Martone, M. ((2014). *Joint Declaration of Data Citation Principles*. FORCE11. San Diego CA. URL: https://force11.org/datacitation (visited on 09/13/2017).

McDonald, D. et al. (2012). "The Biological Observation Matrix (BIOM) format or: how I learned to stop worrying and love the ome-ome". In: *GigaScience* 1, p. 7. DOI: 10.1186/2047-217X-1-7.

McMurdie, P. J. and S. Holmes (2013). "phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data." In: *PloS one* 8.4, e61217. DOI: 10.1371/journal.pone.0061217.

Mons, B. et al. (2017). "Cloudy, increasingly FAIR; revisiting the FAIR Data guiding principles for the European Open Science Cloud". In: *Information Services & Use* 37.1, pp. 49–56. DOI: 10.3233/ISU-170824.

Oteros, J. et al. (2015). "Automatic and Online Pollen Monitoring". In: *International Archives of Allergy and Immunology* 167.3, pp. 158–166. DOI: 10.1159/000436968.

Parr, C. S., N. Wilson, P. Leary, et al. (2014). "The Encyclopedia of Life v2: Providing Global Access to Knowledge About Life on Earth". In: *Biodiversity Data Journal* 2, e1079. DOI: 10.3897/BDJ.2.e1079.

Parr, C. S., N. Wilson, K. Schulz, et al. (2014). "TraitBank: Practical semantics for organism attribute data". In: *Semant Web–Interoperability, Usability, Appl an IOS Press J*, pp. 650–1860.

Schloss, P. D. et al. (2009). "Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities". In: *Applied and Environmental Microbiology* 75.23, pp. 7537–7541. DOI: 10.1128/AEM.01541-09.

Söhngen, C. et al. (2016). "BacDive – The Bacterial Diversity Metadatabase in 2016". In: *Nucleic Acids Research* 44 (D1), pp. D581–D585. DOI: 10.1093/nar/gkv983.

Wilkinson, M. D. et al. (2016). "The FAIR Guiding Principles for scientific

data management and stewardship". In: *Scientific Data* 3, sdata201618. DOI:

10.1038/sdata.2016.18.

Xu, Z. et al. (2014). "Which is more important for classifying microbial commu-

nities: who's there or what they can do?" In: *The ISME Journal*, pp. 1–3. DOI:

10.1038/ismej.2014.157.