1   **Genome-wide association analysis of 350,000 Caucasians from the UK Biobank**

2   **identifies novel loci for asthma, hay fever and eczema.**

4   Åsa Johansson[1], Mathias Rask-Andersen[1], Torgny Karlsson[1], Weronica E. Ek[1*]

6   [1] Department of Immunology, Genetics and Pathology, Science for Life

7   Laboratory, Uppsala University, Uppsala. Sweden

*Corresponding author: Weronica E. Ek

Address: box 815, 75108, Uppsala, Sweden

Telephone: +46703519004.

Fax: +46184714931

Email: Weronica.ek@igp.uu.se

Abstract

Even though heritability estimates suggest that the risk of asthma, hay fever and

eczema is largely due to genetic factors, previous studies have not explained a large

part of the genetics behind these diseases. In this GWA study, we include 346,545

Caucasians from the UK Biobank to identify novel loci for asthma, hay fever and

eczema. We further investigate if associated lead SNPs have a significantly larger

effect for one disease compared to the other diseases, to highlight possible disease

specific effects.

We identified 141 loci, of which 41 are novel, to be associated ($P \leq 3 \times 10^{-8}$) with

asthma, hay fever or eczema, analysed separately or as disease phenotypes that

includes the presence of different combinations of these diseases. The largest number

of loci were associated with the combined phenotype (asthma/hay fever/eczema).

However, as many as 20 loci had a significantly larger effect on hay fever/eczema-

only compared to their effects on asthma, while 26 loci exhibited larger effects on

asthma compared with their effects on hay fever/eczema. At four of the novel loci,

TNFRSF8, MYRF, TSPAN8, and *BHMG1,* the lead SNPs were in LD (> 0.8) with

potentially casual missense variants.

Our study shows that a large amount of the genetic contribution is shared between the

diseases. Nonetheless, a number of SNPs have a significantly larger effect on one of

the phenotypes suggesting that part of the genetic contribution is more phenotype

specific. Identified loci and probable causal genes may in the future be used as targets

for treatments of asthma, hay fever and eczema.

Introduction

Asthma, hay fever and eczema are common complex immunological diseases affecting many people worldwide (1). The prevalence for these diseases vary among populations and have an underlying architecture that include both environmental and genetic risk factors (1). Comorbidity between asthma, hay fever and eczema is common, and previous genome-wide association (GWA) studies have, apart from identifying a large number of genetic variants associated with risk of disease (2–8) also found evidence of a genetic overlap between the diseases (6, 9).

Family and twin studies have estimated that the contribution of genetic factors, i.e. the heritability for asthma (1, 10, 11) to be 35-95%, for hay fever (1, 11) to be 33-91%, and for eczema (12) to be as high as 90%. A recent large study estimated the SNP-based heritability, the heritability that can be attributed the genetic variation captured by SNPs in a GWA study, to be 15% for asthma, 22% for hay fever, and 9% eczema (6). The same study performed a GWA study that included the first release of UK Biobank (N=138,354) analysing asthma, hay fever and eczema as a combined phenotype and identified 99 significantly associated loci (6). Many of the identified target genes were predicted to influence the function of immune cells, and only six loci were identified to have disease specific effects (6). Many previous GWA studies for asthma, hay fever and eczema have been conducted in different cohorts that were subsequently meta-analysed with the purpose of increasing statistical power (2, 4, 6–8, 13).

The aim with this study was to explain a larger part of the genetic background of self-reported asthma, hay fever, and eczema as well as identify possible novel disease specific effects. We investigated the genetic background of self-reported asthma, hay

fever, and eczema combined to a single phenotype, similar as to Ferreira *et al* (6)*,* but

in a more homogenous population, as we included both the first and the second

release of UK Biobank (N=346,545), compared to Ferreira (6) that only included the

first UK Biobank release (N=138,354) as part of a large meta-analysis. We also

analysed each disease phenotype independently in larger groups than previous studies

conducted in UK Biobank. We further had a larger power to investigated if associated

lead SNPs had a significantly larger effect for one disease phenotype compared to the

other phenotypes, to highlight possible disease specific effects. Associated SNPs were

functionally annotated to assess likely causal mechanisms.

Although the phenotypes in the UK Biobank are self-reported, the questions are well

defined and identical for all participants.

Results

<u>Association analysis</u>

The UK Biobank includes 502,682 participants, of which 443,068 are Caucasians.

The disease prevalence in the Caucasian participants were 11.7% for asthma and

23.2% for hay fever and/or eczema (combined). As many as 45.8% of the asthmatic

participants had reported having hay fever and/or eczema, and 23.0% of the hay fever

and/or eczema participants had reported being diagnosed with asthma (Figure 2). We

conducted the GWA study using 346,545 unrelated Caucasians (Table 1), who passed

the quality control (QC) for the second UK Biobank genetic data release, and had no

ambiguities with regards to disease status. After QC, 15,688,218 genetic variants

where included in the analyses (see methods). QC and the final number of included

participants in respective analyses are summarized in the methods and in Table 1. We

did not identify any statistically significant associations located on the X-

chromosome.

<u>GWA study for self-reported asthma</u>

After QC, 41,926 self-reported asthma cases (independent on hay fever/eczema

status) and 239,773 controls were included in the GWA analysis. We identified 75

risk loci located > 1 Mb apart and containing at least one significantly associated

genetic variant ($P \leq 3 \times 10^{-8}$ after adjusting for LD-score intercept of 1.065), that were

associated with self-reported asthma, of which 15 loci were found to be novel asthma

loci not previously identified in a GWA study (Table 2; Manhattan plot, Figure 1; S1

Table and S2 Table; quantile-quantile (QQ) plot, S1 Fig). Using approximate

conditional analysis (14), we identified 116 independent significant associations

within these 75 loci (S1 Table). The strongest associations for asthma were found

within the HLA locus on chromosome 6 ($P = 2.06 \times 10^{-100}$), including 14 independent

genetic variants. Several genes within this region have previously been reported to be

associated with asthma (i.e., *HLA-DQB1, HLA-G* and *HLA-DRB1*) (1, 3, 13). Among

the novel asthma loci, some have previously been associated with other similar

phenotypes (S1 Table). For example, *SDK1,* previously annotated to the nearby

CARD11 gene, have been reported to be associated with atopic dermatitis (15), but

this is the first time that the *SDK1* locus has been identified in a GWA study for

asthma. Five of the 15 novel lead SNPs were further replicated in an independent

cohort (P<0.05; Table 2). However, most of the SNPs were not possible to investigate

in the replication cohort, due to a low number of overlapping SNPs between the

cohorts.

Annotation of asthma associated SNPs

Associated SNPs were further functionally annotated to assess likely causal

mechanisms (see Methods). Overlap with GTEx eQTLs were found for 15 of the 75

asthma loci. Of these, four eQTLs (*EEFSEC, ADAM19, HHEX* and *TMEM258*)

overlapped with the novel loci, where increased expression of *TMEM258* in cell

transformed fibroblasts appears to lower the risk for developing asthma (Table 2; S1

Table S1 and S3 Table). In contrast, increased expression of *EEFSEC* in lung tissue

seems to increase the risk for asthma (S3 Table). Similarly, increased expression of

ADAM19 in whole blood and *HHEX* in cell transformed fibroblasts appears to

increase the risk of asthma (S3 Table). 19 probable causal missense variants could be

observed within the 75 significant GWA loci of which four missense variants for the

15 novel loci (S4 Table). The latter are located within *TNFRSF8, MYRF, TSPAN8,*

and *BHMG1.* The association at *TNFRSF8* was represented by only one genetic

variant, rs2230624 (S2 Fig). This SNP is a missense variant in two transcripts for

TNFRSF8 and causes a cysteine to a tyrosine substitution which was predicted as

'probably damaging' by PolyPhen (16) (PolyPhen-score 0.751-0.921) and had a

'deleterious' SIFT-score (17) of 0. The lead SNP at the *MYRF* locus, rs174535, is a

missense variant in five transcripts for *MYRF*. Rs174535 causes a serine to arginine

substitution and was predicted to be 'probably damaging' by PolyPhen(16)

(PolyPhen-score 0.961-1) and had a 'deleterious' SIFT-score (17) of 0.04-0.07.

However, rs174535 is also in LD with the most significant eQTL for *TMEM258* in

cell transformed fibroblasts. The lead SNP in the *BHMG1* locus, rs11671106, is a

missense variant for *BHMG1* and was predicted as 'probably damaging' by PolyPhen

(16) (PolyPhen-score 0.94) and had a 'deleterious' SIFT-score (17) of 0.01. The lead

SNP at the *TSPAN8* locus, rs11178649, was in complete LD with rs3763978 (R^2=1), a

missense variant in three transcripts for *TSPAN8,* which causes a glycine to alanine

substitution which was predicted as 'probably damaging' by PolyPhen (16)

(PolyPhen-score 0.989) and had a 'deleterious' SIFT-score(17) of 0.03

<u>GWA study for self-reported hay fever/eczema</u>

After QC, 84,034 self-reported hay fever and/or eczema cases that were combined as

a single phenotype were included in the analysis. We identified 109 loci to be

associated ($P \leq 3 \times 10^{-8}$, LD-score intercept =1.079) with self-reported hay

fever/eczema, and 22 of these were novel (Table 3; Manhattan plot, Figure 1; S5

Table and S6 Table; QQ-plot, S3 Fig). The strongest association was observed for the

lead SNP rs5743604 ($P= 7.5 \times 10^{-72}$) located within *TLR1*. This SNP has previously

been associated with allergic disease (6, 21). Using conditional analysis, we identified

154 independent significant associations within these 109 loci (Table 3; S5 Table).

Moreover, two of our lead SNPs (rs4845604 and rs9986945, mapped to *RORC* and

SDK1), observed within previously known loci were in low LD ($R^2 \leq 0.05$) with the

previously reported genetic variants, indicating that they represent novel variants

within or close to known loci (S5 Table). The *UBAC2* locus has previously been

reported to be associated with asthma (9), but this is the time that the *UBAC2* locus is

reported to be associated with hay fever and/or eczema. We replicated six of the novel

lead SNPs in an independent eczema GWA study, the EAGLE consortium ($P \leq 0.05$),

(Table 3). However, all SNPs that did not replicate in EAGLE were neither significant

in UK Biobank when analysing eczema separately (Table 3).

Annotation of hay fever/eczema SNPs

For eleven of the 109 hay fever/eczema associated loci, the lead SNP was in LD with

the lead SNP for GTEx eQTLs (Table 3; S3 Table) and 14 overlapped with possible

causal missense variants in genes, including *IL6R, IL7R, IL13* and *SMAD4* (S7

Table).

GWA studies for hay fever and eczema analysed separately

Hay fever and eczema could not be separated for most of the participants, since they

had primarily answered yes or no on whether they had either hay fever or eczema.

However, to investigate hay fever and eczema individually, we also analysed hay

fever (N hay fever cases = 18,915) and eczema (N eczema cases = 7,884) separately

in a smaller subset of UK Biobank participants (Manhattan plot, S4 Fig; QQ-plots, S5

and S6 Fig). A total of 27 and 18 loci were identified for hay fever and eczema,

respectively. One novel hay fever and one novel eczema locus, which has not been

reported in previous GWA studies and that were not significantly associated in the

combined hay fever/eczema analysis, was detected when analysing hay fever and

eczema separately (S8-S11 Tables). The lead SNP, rs12920150 ($P=1.02 \times 10^{-9}$), at the

hay fever locus is located close to *CBLN1* and the lead SNP, rs2485363 ($P=1,20 \times 10^{-8}$

^{8}), at the eczema locus is located downstream of *TAGAP*. This novel eczema locus

was nominally replicated using the summary statistics from the GWA study on

eczema in the EAGLE consortium (P=0.018, OR=1.05 [95% CI 1.02-1.92]). Another

locus that was not detected when analysing hay fever/eczema combined was detected

when analysing eczema separately. The lead SNP for this locus, rs676387

($P=2.26 \times 10^{-10}$), is located within *HSD17B1* (S10 and S11 Tables). This region has

previously been reported to be associated with allergic disease (6) and overlap with an

eQTL for *TUBG2* in skin, where a decreased expression of *TUBG2* seems to lower

the risk for eczema (S3 Table).

<u>GWA study for asthma/hay fever/eczema (combined as a single phenotype)</u>

For the combined analysis of asthma and/or hay fever and/or eczema (N

cases=106,752), we identified 110 significant loci (LD-score intercept=1.081), and 16

of these were novel GWA loci that have not been significantly associated with either

asthma, hay fever or eczema in previous GWA studies (Table 4; Manhattan plot,

Figure 1; S12 and S13 Tables; QQ-plot, S7 Fig). However, 12 of these 16 novel loci

were detected when analysing asthma and hay fever/eczema separately, while the

remaining four novel loci were only found when analysing asthma, hay fever, and/or

eczema together as a single phenotype. Using conditional analysis, we identified 164

independent associations within these 110 loci (Table 4; S12 Table). The most

significant SNP, rs72823641 (P=1.14×10^{-78}), was located within *IL1RL1* and was also

significantly associated with asthma and hay fever/eczema when these phenotypes

were analysed separately (P = 4.09×10^{-61} and P = 9.64×10^{-64}) (S12 and S13 Tables).

This region has previously been associated with allergic diseases (6) (S14 Table). We

also identified five lead SNPs for the combined phenotype asthma and/or hay fever

and/or eczema within previously known loci, which were found to be in low LD

($R^2 \leq 0.05$) with previously reported genetic variants, indicating that they represent

novel variants within known loci. These five lead SNPs mapped to *LPP, IL31,*

LINC00393, CCR7 and *NFATC* (S12 and S14 Tables).

Annotation of Asthma/hay fever/eczema (combined as a single phenotype) SNPs

For 16 of the 110 asthma and/or hay fever and/or eczema associated loci, the lead

SNPs overlapped with a lead SNP for an eQTL (Table 4; S3 and S12 Tables). Among

the novel loci, one overlapped with an eQTL for *HIST1H2BD* in whole blood

(P=1.11×10^{-16}). A decreased level of *HIST1H2BD* seems to increase the risk of this

combined phenotype (S3 Table). Probable causal missense variants could be observed

at 17 out of 110 significant loci and one of these was observed at one of the novel loci

located within *TNFRSF8* and was also identified in the asthma analysis above (S15

Table).

<u>SNP-based heritability</u>

To quantify the SNP-based heritability for asthma and hay fever/eczema, we used LD

score regression analysis (LDSC) (19). These analyses included the same cases and

controls as for the association analysis (see Methods). The SNP-based heritability was

estimated to be 21% for asthma and 16% for hay fever/eczema (Table 5). Our

significant loci, which were located $\geq$ 1 Mb apart and contained at least one

significantly associated genetic variant at $P \leq 3 \times 10^{-8}$, explained 4.2% of the heritability

for asthma and 3.6% of the heritability for hay fever/eczema (Table 5).

<u>Identification of phenotype specific loci (SNP)</u>

In our GWA studies, we included all individuals reporting either asthma (for the

asthma GWA study) or hay fever/eczema (for the hay fever/eczema GWA study) as

cases, independent on if they reported having the other disease phenotype (i.e.,

asthma cases could have reported having asthma and hay fever/eczema or only

asthma). To investigate possible phenotype-specific SNPs, we performed polytomous

(multinomial) logistic regression to identify whether the effect of a locus (lead SNP)

was significantly (FDR$\leq$0.05) larger for one disease phenotype as compared to

another. These effects can therefore be considered as being disease/phenotype

specific. To conduct these analyses, we used four non-overlapping groups: 1) asthma

cases without hay fever/eczema (N=22,858), 2) hay fever/eczema cases without

asthma (N=65,063), 3) asthma cases with hay fever/eczema (only including N=19,299

participants that had reported asthma in combination with hay fever or eczema), and

4) controls without asthma, hay fever and eczema (N=240,817) (Figure 2). Hay fever

and eczema were not separated in this analysis due to the small sample size (Table 1).

Groups were compared in a pairwise fashion (S16 Table).

A total of 154 lead SNPs (see method section for a description of the selection of lead

SNPs), representing the 138 different loci, identified in the GWA study for asthma,

hay fever/eczema, or for asthma/hay fever/eczema, were included in polytomous

logistic regression analyses. To illustrate the specificity in the Venn diagrams (Figure

3), each SNP was assigned to an area that represents either a phenotype-specific effect

(significantly larger in one group of cases) or a shared effect (no significant difference

between the two groups of cases).

In the comparison of asthma-only, i.e. without hay fever and eczema, to hay

fever/eczema-only, i.e. without asthma, 26 loci/SNPs were specific for asthma-only,

i.e. had a significantly higher OR for asthma compared to for hay fever/eczema, while

20 loci/SNPs were specific to hay fever/eczema-only. A major part of the loci/SNPs,

103, showed no significant difference in effect between these disease phenotypes.

(Figure 3; S16 Table).

When comparing subjects with asthma and hay fever/eczema to subjects with asthma,

53 loci/SNPs were specific for asthma with hay fever/eczema. No SNP was specific

for the asthma only group (Figure 3; S16 Table). For the remaining 96 loci/SNPs,

there was no significant difference in effect between subjects with asthma only and

subjects with asthma as well as hay fever/eczema.

Finally, when comparing cases of hay fever/eczema only with cases of hay

fever/eczema combined with asthma (Figure 3; S16 Table), 64 loci/SNPs had

significantly larger effect in the group with hay fever/eczema combined with asthma.

No locus had a larger effect in the hay fever/eczema without asthma group. As many

as 83 loci/SNPs had no detectable difference in effect between these two disease

phenotypes.

For some loci, multiple, possibly independent ($R^2 \leq 0.8$) SNPs were included in the

analyses. For most of the analyses, such independent SNPs within the same locus

showed the same phenotype specificity, or lack of specificity. That is, all independent

SNPs within one locus belong to the same area in the Venn diagram (Figure 3).

However, for a number of loci, the effect for the different independent SNPs showed

different phenotype specificity. This resulted in 149 independent loci/SNPs when

comparing the asthma-only group to the hay fever/eczema-only group and when

comparing subjects with asthma and hay fever/eczema to subjects with asthma only

(Figure 3). For the last group, when comparing cases of hay fever/eczema only with

cases of hay fever/eczema combined with asthma, 147 independent loci/SNPs were

identified and included in the Venn diagram (Figure 3). For example, two

uncorrelated SNPs ($R^2 < 0.05$) were found to be located within the same intron of

IL2RA: rs61839660, which was associated with hay fever/eczema, and rs12722547,

which was associated with asthma in the GWA study. The rs61839660 SNP has a

significantly larger effect in both hay fever/eczema-only and hay fever/eczema with

asthma, compared to asthma-only but no difference in effect between hay

fever/eczema with or without asthma. The effect of rs12722547 was instead

significantly larger in the hay fever/eczema with asthma group, compared to the hay

fever/eczema without asthma group. Rs12722547 also exhibited a trend (nominal P =

0.05) towards having a larger effect in asthma-only compared to hay fever/eczema-

only (S16 Table).

Discussion

In this large GWA study, including 346,545 unrelated Caucasian participants from

UK Biobank, we identified 141 unique loci that are associated with self-reported

asthma, hay fever, and/or eczema when these traits are analysed separately or together

as combined phenotypes. In comparison with previous studies based on UK Biobank

and similar disease phenotypes, our study has several strengths and presents

additional results. Out of all identified loci, as many as 41 are novel to our study and

have not been reported to be associated with the same disease phenotype previously.

Compared to Ferreira *et al* (6) and Zhu *et al* (9), that only included the first release of

UK Biobank, we included the full UK Biobank cohort. We also present five different

GWA studies for five different phenotypes and further had the strength to identify a

number of possible phenotype specific effects that had not been discussed previously.

The largest number of loci were associated with combined phenotype (asthma and/or

hay fever and/or eczema), most likely due to the larger sample size of this group.

However this is in agreement with a shared genetic contribution between diseases, as

has been shown in Ferreira *et al* (6) and Zhu *et al* (9). With this combined phenotype

(asthma and/or hay fever and/or eczema), we identified four novel loci that were not

found for asthma or hay fever/eczema when analysed separately. Three of these loci

appear to be highly relevant to the pathogenies of all three diseases: *SMAD7, KLF2*

and *RIN3*. The variant at the *KLF2* locus is located in the 5' UTR of *KLF2*. This gene

plays a role in processes during development including epithelial integrity,

inflammation, and T-cell viability. Previous studies have found associations between

this locus and lymphocyte percentage of white cells, neutrophil percentage of white

cells, white blood cell counts, monocyte percentage of white cells, and eosinophil

percentage of granulocytes (20). The variant at the *SMAD7* locus is located within an

intron of *SMAD7*. This gene has previously been associated with inflammatory bowel

disease (IBD) (21), colorectal cancer (22), and haemoglobin concentration (20). The

variant at the *RIN3* locus, is located within an intron of *RIN3*, and is also associated

with *RIN3* expression. This gene has previously been associated with myeloid white

cell count, eosinophil basophil counts (20), and chronic obstructive pulmonary

disease (23). It is worth noting that some of our novel loci have previously been

associated with a related phenotype (S1, S5, S8, S10 and S12 Tables). For example,

some of the novel asthma loci has previously been associated with IgE levels,

eosinophil counts or dermatitis and some of the novel hay fever/eczema loci with IgE

levels or eosinophil counts.

For four of the novel loci: near *TNFRSF8, MYRF, TSPAN8,* and *BHMG1;* the lead

SNP was in LD with potentially deleterious missense variants. The lead genetic

variant at the *TNFRSF8* locus, rs2230624, which is associated with asthma as well as

the combined asthma/hay fever/eczema phenotype, is a potentially causal missense

variant that causes a cysteine to a tyrosine substitution in the TNFRSF8 protein. This

protein, which is also referred to as *CD30*, is a receptor that is expressed on activated

T and B cells and has been shown in clinical studies to have a role in the development

of allergic asthma (24). To the best of our knowledge, this is the first time that this

locus has been associated with asthma and allergy in a GWA study. The lead SNP at

the *MYRF* asthma locus, rs174535, is a missense variant within the myelin regulatory

factor protein (*MYRF*) that causes a serine to arginine substitution near the end of the

protein. This gene lies within the fatty acid desaturase (*FADS*) cluster on a fatty acid

synthesis-associated haplotype (25). Variants on this haplotype are also strongly

associated with expression of *FADS1* and *FADS2,* two genes that are involved in the

desaturation of polyunsaturated fatty acids in the biosynthesis of long chain

polyunsaturated fatty acids (LC-PUFAs). One of these variants has previously been

shown to modulate the effect of breast-feeding on asthma (26); another has been

associated with increased risk of inflammation (27). Reduced capacity to desaturase

omega-6 LC-PUFAs due to *FADS* polymorphisms has been shown to be nominally

associated with reduced risk for development of eczema, potentially due to a

pathogenic role of omega-6 LC-PUFAs in development of allergy (28).

For seven of the 41 novel GWA loci, the lead SNP was in LD (> 0.8) with an eQTL.

We could see a positive correlation between expression of *TUBG2* (in skin), *HHEX*

(in cell transformed fibroblasts), *EEFSEC* (in lung and cell transformed

fibroblasts), and *ADAM19* (in whole blood) and risk of disease, as well as a negative

correlation for *TMEM258* and *HIST1H2BD*. Decreased expression of *TMEM258* in

cell transformed fibroblasts was associated with increased risk of asthma.

In transgenic experiments in mice, it has been shown that a lower expression

of *TMEM258* leads to severe intestinal inflammation (29), which agrees with our

results. A possible limitation of this analysis is that it relied solely on the GTEx

database. Additional sources of information on eQTLs may increase the total number

of eQTLs that are associated with asthma, hay fever and eczema.

For 16 loci that were associated with asthma, 20 loci associated with hay

fever/eczema and for 21 loci associated with asthma/hay fever/eczema, we identified

multiple independently associated variants. This indicates that several of the asthma-,

hay fever-, and eczema-associated loci represent multiple independent disease-

associated variants. As an example, the *FLG* locus contains three independent asthma,

and asthma/hay fever/eczema-associated variants. This gene has previously been

shown to contain loss-of-function mutations that are causal for skin barrier deficiency

and strongly predispose to both eczema and asthma (30). The four most prevalent

European *FLG* mutations are c.2282del4, p.R501X, p.R2447X, and p.S3247X (30).

An additional example is the *HLA* region whose association with immune diseases is

particularly complex and which has previously been suggested to include several

independent regulatory factors (31). In our analyses, we identify as many as 21

independent associations within this locus.

As highlighted by this study, as well as previous studies (2–7, 32), many disease-

associated loci overlap between asthma, hay fever and eczema. However, several loci

were only significantly associated with only one of the investigated phenotypes. By

testing for association with hay fever and eczema separately in a smaller set of

participants, we were able to resolve some of these signals. Interestingly, one of the

strongest associations for hay fever/eczema ($P=7.96 \times 10^{-25}$), found within the *FLG*

locus, was more significantly associated with eczema when this phenotype was

analysed separately ($P=1.15 \times 10^{-69}$). In contrast, this variant was not associated with

hay fever when hay fever was analysed separately. It was, however, associated with

asthma ($P=2.37 \times 10^{-27}$). This is in agreement with the previous GWA study by Ferreira

et al, where a SNP at the *FLG* locus was shown to be specifically associated with

eczema (6). However, a different study has shown that mutations within the *FLG*

locus are associated with eczema starting in the first year of life, and that these

mutations are associated with a later development of both asthma and hay fever (33).

This is an example of the typical progression of allergic diseases that often begin

early in life, which is commonly referred to as the atopic march (33–35). When

analysing hay fever separately, we identified one novel locus near *CBLN1*. Studies on

transgenic mice have shown that knock-out of *CBLN1* mimics loss-of-function

mutations that occur in the orphan glutamate receptor, *GRID2* (36). Autoantibodies

against glutamate receptors are involved in the development of autoimmune disease

(37). One novel locus was also identified and replicated when analysing eczema

separately, downstream of *TAGAP*. This locus has previously been associated with

celiac disease (38) and multiple sclerosis (39).

We further investigated our novel asthma, hay fever/eczema and eczema loci in two

independent cohorts. We were only able to replicate six SNPs out of the 15 identified

to be novel for asthma with the summary statistics from the GABRIEL asthma

consortium. The GABRIEL GWA study only included 582, 802 SNPs genotyped with

the Illumina Human610 quad array, and therefore, a large number of SNPs did not

overlap between our studies. However, for most of the loci where we identified the

same SNP or a proxy in LD (R2≥0.8), we did find a nominal replication (P≤0.05)

(Table 2). The GABRIEL study was based on childhood asthma while the UK

Biobank asthma is based on adult asthma and these two disease phenotypes may

therefore have some different underlying genetic effects. It is also important to

remember that a lack in replication may also be due to a lower power to detect

associated SNPs in GABRIEL due to a smaller sample size. Five lead SNPs identified

for the combined analysis hay fever/eczema replicated in the EAGLE study (Table 3).

The lack of replication is most probably due to differences between phenotypes.

While the EAGLE study only included eczema cases, our study also included hay

fever. All SNPs that did not replicate in EAGLE, was neither statistically significant

when analysing eczema independently in UK Biobank (Table 3). We also tried to

replicate the four novel loci that was only identified in the combined analysis,

asthma/hay fever/eczema, using the GABRIEL and EAGLE cohorts. However, none

were nominally replicated (P‹0.05) which is probably due to the smaller sample sizes

in GABRIEL and EAGLE, and most importantly due to the differences in disease

phenotypes between the cohort.

Out of all asthma and/or allergic disease-associated loci that have been reported to the

GWAS catalog as of December 2, 2018, the majority (N=108) were nominally

replicated in our study (P≤0.05; S14 Table). Twelve associations were not possible to

test due to lack of data, i.e. neither the reported SNP nor any SNP in LD with the

reported SNP were presented in our data (S14 Table). Asthma, hay fever, and eczema

are known to be heterogeneous diseases in which environmental factors play an

important role (1). Genetic variants associated with asthma, hay fever and eczema are

likely to be population specific (40). It is therefore possible that population-specific

variants are not detected in our study. Many of the previous associations that were not

replicated in our study have been identified in studies that have used a somewhat

different phenotype (41, 42), populations of different ancestry (15, 43) or small

sample sizes (< 10,000) (43, 44). Research findings from studies on smaller cohorts

are more likely to be false positives, especially when no replication of primary

findings has been performed, and are thereby less likely to represent true causative

mechanisms (45) (for more information, see S14 Table). A recent GWA study by Zhu

et al (9), which was also conducted on the UK Biobank cohort, however using a

different combination of allergies as a phenotype, reported seven novel allergy-

associated loci, five of which were replicated in our study. These loci where not

available in the GWAS catalog at the time of writing this article and are therefore not

included in S14 Table. The two loci that did not replicate in our study where mapped

to *ALG9* on chromosome 11(rs659529) and to *EVI5* on chromosome 1 (rs12743520).

In previous GWA studies for asthma, the disease phenotype commonly contained

other disease phenotypes as well, e.g. participants with asthma commonly also report

hay fever/eczema. In contrast, our polytomous logistic regression approach allowed

for identification of genetic variants with differing effects between the different sub-

phenotypes. These effects can therefore be considered as being disease/phenotype

specific. This was achieved by subdividing the participants in four non-overlapping

groups depending on asthma and hay fever/eczema status. The SNPs that were

included in these analyses were selected from our main GWA analyses, but not

including the two SNPs identified for the hay fever and eczema phenotypes analysed

separately since we did not have power (large enough sample size) enough to include

hay fever and eczema separately in these analyse. This means that a locus that was

defined as specific for asthma-only has already been associated with any of the

combined phenotypes and/or with asthma, independent of hay fever /eczema status.

The association for such variants may have been due to comorbidity between asthma

and the other diseases, e.g. a larger fraction of asthma cases in the hay fever/eczema

group compared to the controls, or that the effect of the asthma-only specific variants

was only partly diluted by being combined with other disease phenotypes. A large

number of loci exhibited differential effects between hay fever/eczema-only and

asthma-only. As many as 20 loci had a significantly larger effect on hay

fever/eczema-only compared to their effects on asthma while 26 loci exhibited larger

effects on asthma compared with their effects on hay fever/eczema (Figure 3). Among

the loci that were specific for asthma-only, we find *ADAM19* and *ADAMTSL3* which

are proteins with multiple biological roles within the cell and believed to be important

in a number of diseases, including asthma (46). Among the loci that were specific for

hay fever/eczema we find the toll like receptor loci, *TLR1/TLR10*, which also showed

a larger effect on hay fever compared to asthma-only in the Ferreira *et al* study (6).

Most associated variants at this locus are located within the promoter region of *TLR1*,

which encodes the toll-like receptor 1. This protein constitutes a component of the

innate immune response to microbial pathogens (47). Several loci that overlap

between asthma-only and hay fever/eczema-only were annotated to genes related to

tumor necrosis factor (TNF) function, such as *TNFAIP3, TNFAIP8, TNFRSF11A,*

TNFRSF14, TNFRSF6B, TNFRSF8, TNFSF4. These proteins are mainly expressed in

immune cells and regulate immune response and inflammation as well as

proliferation, apoptosis and embryogenesis (48).

The largest number of phenotype-specific loci was observed for the group of cases

with asthma and hay fever/eczema (Figure 3; S16 Table), a group of cases that has not

been included in similar analyses in previous studies (6). This is a group of

participants with an allergic disease in combination with asthma, which could to some

degree represent participants with allergic asthma. The number of phenotype-specific

loci is considerable larger in our study compared to previous studies that have

performed similar analyses, such as the study by Ferreira *et al.*(6),which only

identified six disease specific loci. This is not surprising since our analyses included

larger sample sizes: N= 65,063, N=22,858, and N=19,299 compared to N= 33,305,

N=12,268, and N=6,276 in the study by Ferreira *et al* (6) for the three sub-groups

included in the analyses of disease-specific effects. In addition, since only genome-

wide significant SNPs were taken forward to the polytomous logistic regression

analyses, we used the False Discovery Rate by Benjamini-Hochberg to adjust for

multiple testing. This increases the power to pinpoint as many positive findings as

possible, still with a small false-discovery rate (5% in our case), compared to the

more conservative Bonferroni method used in the previous study by Ferreira *et al* (6) .

The previous study, also separated hay fever and eczema, and compared the three

groups hay fever-only, eczema-only and asthma-only. Since different subgroups of

cases were analysed in our study our results do not disagree with that of Ferreira *et al*

(6) that found six disease-specific SNPs: near *FLG, RPTN-HRNR* (close to *FLG*),

IL2RA, IL1RL2- IL8R1, WDR36-CAMK4 and *GSDMB;* where five of them were

significantly different between hay fever and eczema.

The SNP-based heritability was estimated to be 21% for asthma and 16% for hay

fever/eczema. These percentages represent the portion of heritability that can be

captured by the common genetic variants that were included in the GWA study. The

SNP-based heritability for asthma has previously been estimated at 15%, which is

slightly lower than the estimate from our study and probably due to a smaller sample

size, and/or a difference in disease definition in the previous study (6). In comparison

to the high estimates for the heritability (33-95%) from family and twin studies (1, 10,

12), this suggests that a major contribution to the genetic risk for asthma, hay fever

and eczema might not be identified in studies using common genetic variants, or need

cohorts with even larger sample sizes. However, heritability estimates from family

and twin studies have been suggested to be overestimated (49–51) due to the fact that

these estimates often are based on simplistic models that ignore shared environmental

factors. Our estimate might also be lower due to the presence of disease-associated

rare variants that are not captured by the SNP-based heritability estimate.

A possible limitation of the present study is the self-reported phenotypes, which might

lead to a recall bias and misclassification. Another limitation is that the UK Biobank

cohort traits are not independent since there are shared cases between asthma, hay

fever and eczema and completely shared controls. However, findings presented in this

article apply to a single large population of individuals of similar age. Population

stratification was also controlled for by filtering for Caucasian participants, including

ancestry derived principal components and adjusting for the LD-score intercept in our

analyses. Participants of the UK Biobank are also more likely to be exposed to more

similar environmental factors, compared to the participants of previous meta-analyses

that utilise a large number of smaller cohorts from different countries and age-groups.

Analysing hay fever and eczema as a combined phenotype is another limitation in our

study, which prohibits identification of hay fever- and eczema-specific SNPs. We

therefore refer to SNPs as phenotype-, rather than disease-specific in the polytomous

logistic regression analyses. However, both hay fever and eczema are IgE mediated

hypersensitivities and therefore probably share similar physiology (52).

Conclusion

In summary, we describe 15 novel loci for asthma, 22 novel loci for hay fever and/or

eczema and an additional four novel loci were found when analysing asthma, hay

fever and eczema together. Two novel loci were also identified when analysing hay

fever and eczema separately. Pinpointing candidate genes for common diseases are

important for tailor-made studies that want to prioritize candidate genes for

developing novel therapeutic strategies. This study further highlights a large amount

of shared genetic contribution to these diseases, indicating that the comorbidity

between asthma, hay fever and eczema is partly due to shared genetic factors.

However, we also show that a number of SNPs have a significantly larger effect on

one of the phenotypes, suggesting that part of the genetic contribution is phenotype

specific.

Methods

Study population

The UK Biobank includes 502,682 participants recruited from all across the UK. Participants were between 37 and 73 years old at time of recruitment between 2006 and 2010. Most participants visited the centre once, but some individuals visited the centre at up to three times. Participants answered questions about self-reported medical conditions, diet, and lifestyle factors. A total of 820,967 genotyped SNPs and up to 90 million imputed variants is available for most participants. The UK Biobank study was approved by the National Research Ethics Committee (REC reference 11/NW/0382). An application for using data from UK Biobank has been approved (application nr: 15479). We included 346,545 unrelated Caucasians (see selection of participants and sample QC below) with genotypes from the second UK Biobank genotype release (Table 1).

Disease phenotypes: asthma and hay fever/eczema

Self-reported asthma as well as self-reported hay fever and/or eczema (combined) were assessed using the UK Biobank touch screen question number (Data field 6152), which asked the participants the following question: *has a doctor ever told you that you have had any of the following conditions? (You can select more than one answer):* 1) asthma and 2) hay fever, allergic rhinitis or eczema, 3) none of the above or 4), prefer not to answer. Because hay fever and eczema diagnosis could not be separated we called this variable hay fever/eczema (i.e., participants reported hay fever and/or eczema). All participants were also invited to participate in an interview. At first, nurses (trained UK Biobank staff-member) confirmed with each participant that the information they provided on the screen or questionnaire was correct if they had answered that a doctor had told them they had one or more of the following

diseases: heart attack, angina, stroke, high blood pressure, blood clot in leg, blood clot

in lung, emphysema/chronic bronchitis, asthma, or diabetes. Due to the confirmation

of asthma cases, the overlap in asthma variables between the touch-screen

questionnaire and verbal interview was very high. For asthma, only 622 individuals

were removed due to conflicting answers between the touch-screen and verbal

interview. Using a drop-down menu, the nurses could also add other diagnoses. These

diagnoses (UK Biobank data field 20002) were used to define hay fever and eczema

cases separately. However, the disease prevalence in this variable appears to be

largely underreported as many individuals reported hay fever or eczema in the touch-

screen questionnaire but did not report hay fever or eczema during the interview. For

this reason, the touch-screen data variables hay fever/eczema, with a much larger

sample size (Table 1) compared to hay fever and eczema separately, was used as one

of the primary phenotypes analysed in this study. For hay fever/eczema, 4,881

individuals were removed due to conflicting answers between the touch-screen

questionnaire and the interview, for individuals reported they had hay fever during the

interview but not on the touch-screen questionnaire (N=2,143), or reported they had

eczema in the interview but not in the touch-screen (N=2,738). We further removed

22 individuals who had asked to be removed from the UK Biobank.

Controls

Controls (N=239,773) were selected as individuals answering "none of the above" in

question 6152, and who did not report asthma, hay fever or eczema in variable

number 20002. The same controls were used for all phenotypes.

Genotyping

The UK Biobank Axiom array had been used to genotype 438,417 of the 502,682 UK

Biobank participants. The other 49,994 samples (all from the interim release) had

been genotyped on the closely related UK BiLEVE array. The UK BiLEVE cohort

and the rest of the UK Biobank differ only in small details of the DNA processing

stage. The two arrays have 95% common marker content. We included a variable for

array type (UK BiLEVE or UK Biobank Axiom) as covariate. SNPs in UK Biobank

were imputed using UK10K (53) and 1000 genomes phase 3 (54) as reference panels.

Imputation in the second release resulted in 92,693,895 SNPS (released in June 2017).

However, because the UK Biobank reported problems with imputation quality for a

subset of the SNPs (caused by mismatch in coordinates between the UK10 and the

1000 genomes reference panels), we followed the recommendation to only include

genetic variants included on the HRC panel (55) (N=39,727,058).

<u>Quality control</u>

Quality control of genotype data and imputation of genotypes had already been

carried out centrally by UK Biobank. From the imputed dataset, we only included

SNPs in the HRC panel with a MAF ≥ 0.01. We removed SNPs deviating from

Hardy-Weinberg (P-value $< 1 \times 10^{-20}$) and markers with more than 5% missing

genotype data. We only included SNPs with an imputation quality >0.3. After QC, a

total of 15,688,218 autosomal SNPs and SNPs on the X-chromosome were included

in our analyses. We only included Caucasian participants who were clustering

according to the genetic principal components (56,180 non-Caucasians were

removed: individuals listed in UK Biobank data file 22006). We further removed first

and second-degree relatives (N=32,751), using kinship data (estimated genetic

relationship > 0.044), and participants with sex discordance, high

heterozygosity/missingness (individuals listed in UK Biobank data field 22010 and

22027), and participants with more than 5% missing genotypes. After QC and

exclusion, 346,545 unrelated Caucasian participants remained.

<u>Genome-wide association study</u>

A GWA study were performed for each phenotype using logistic regression and an

additive genetic model implemented in PLINK version 1.90 (56). We performed a

GWA studies for five sets of phenotypes: 1) asthma (independent on hay fever and

eczema status), 2) hay fever/eczema (hay fever and/or eczema independent on asthma

status), 3) hay fever and/or eczema and/or asthma, as well as 4) hay fever

(independent on asthma and eczema status) and 5) eczema (independent on asthma

and hay fever status). The same controls, that have reported that they did not have any

of the disease phenotypes, were used for all analyses (N=239,773). The following

covariates were included in our analysis: Townsend deprivation index (TDI) (as a

proxy for socioeconomic status), sex, age, smoking, and the first ten ancestry derived

principal components. In addition, to adjust for the different genotyping chips, we

included a binary indicator variable for UK Biobank Axiom versus UK BiLEVE

genotyping array. We calculated the LD-score intercept, using the LD score

regression software (LDSC) (19), for each phenotype and adjusted the summary

statistics accordingly (19). The genome-wide significance threshold was set to 3×10^{-8},

as suggested for GWA studies that include variants with a minor allele frequency

(MAF) ≥ 0.01(57), which was the threshold used in our study. Individual loci were

defined as regions with at least one significantly associated SNP (P $\leq 3 \times 10^{-8}$). Start

and stop positions for each locus were where no additional significantly associated

SNPs could be found (upstream for start position, or downstream for stop position)

within 1 Mb.

<u>Identification of additional independent variants within associated loci</u>

To identify independently associated variants within each defined locus (significant

SNPs (P ≤ 3x10^{-8}), within 1 Mb), we used an approximate conditional analyses

implemented in GCTA (14). LD calculations were based on 5,000 randomly selected

Caucasian participants from UK Biobank (after sample QC). For each locus, the most

significant top SNP was identified and the summary statistics of all SNPs within the

same locus was adjusted by the effect of the lead SNP. After adjusting for the lead

SNP, we identified the most significantly associated SNP within the locus that

remained significant (P ≤ 3x10^{-8}). In the next step, we once again adjusted the

summary statistics of all SNPs within the same locus, by including the effect of both

the original lead SNP and the conditional lead SNP form the first iteration. This

process was thereafter repeated until no other SNPs within the locus were found

significant after adjusting for all previously detected independent lead SNPs.

<u>Determining the novelty status of significant loci</u>

To determine whether significant loci were novel to any of the diseases, we compiled

a list of all asthma, hay fever, eczema and allergy risk SNPs with genome-wide

significant association (≤ 5x10^{-8}) reported in the NHGRI-EBI GWAS catalog

(downloaded December 2, 2018). We also searched for GWA study results using

PubMed and bioRxiv. We classified a locus to be 'novel, if the locus was > 1 Mb

from any of the previously reported loci/variants for the disease. We also estimated

LD between each lead SNP and all genome-wide significant associations found in the

NHGRI-EBI GWAS catalog, to determine whether the lead SNP was a novel variant

in a known locus (if the locus was < 1 Mb from any of the previously reported

loci/variants for the disease). We considered our associated SNP to be a novel variant

if R^2 were smaller than 0.05 between our top associated variant and previously

reported variants within the same locus. A locus was also reported as novel for a

specific disease (i.e., asthma) if previous GWA studies only reported association to a

different allergic disease (for example hay fever). If the locus was previously reported

for a combined phenotype, i.e. in studies combining different allergic diseases,

including the one tested, it was not reported as a novel locus.

<u>Annotation of target genes and identification of causal genetic effects</u>

To identify likely target genes for associated variants, we first reported the closest

gene(s) to the lead SNP for each locus and reported if the SNP was intronic or exonic

using the Human Genome Browser (GRCh37). We also performed additional

analyses to potentially better define plausible target genes. To examine the

relationship between the lead SNP for each locus and gene expression we used the

Genotype-Tissue Expression (GTEx) database (58) to find evidence of overlap with

expression quantitative loci (eQTLs). We downloaded significant eQTLs from the

Genotype-Tissue Expression (GTEx) database. First, we selected GTEx SNPs that

overlapped with the UK Biobank SNPs and used a conservative significance threshold

$P \leq 2.3 \times 10^{-9}$ for cis effects (<1 Mb) form the GTEx data, in agreement with

previous studies (6). Second, we identified the most significant eQTL SNP for each

tissue and gene in the GTEx dataset. Third, we estimated the LD between the lead

eQTL SNPs and our lead GWA study SNPs. A lead GWA SNP in LD ($R^2 > 0.8$) with

a lead GTEx eQTL SNP was considered to overlap with the eQTL. Only cells or

tissues that were relevant for our disease phenotypes were considered when searching

for eQLTs, including EBV-transformed lymphocyte, transformed fibroblasts, whole

blood, lung and skin (sun exposed and not sun exposed).

713 We also used the Bioconductor biomaRt (59) package in R for functional annotation

714 of associated SNPs. In BiomaRt, lead SNPs, and all SNPs in LD ($R^2>0.8$) with a lead

715 SNP, were cross-referenced against: Ensembl Genes, Ensembl Variation, and

716 Ensembl Regulation version 91 (Accessed 9 December 2017 using the human

717 assembly GRCh37). Here we checked whether the lead SNPs were in LD ($R^2>0.8$)

718 with a potentially functional genetic variant by investigated regulatory features for the

719 SNPs (i.e., promoters, enhancers etc.), binding motifs (i.e., if any of the SNPs were

720 found within a motif for a transcription factor), and if the SNPs were possibly

721 damaging variants (i.e., missense, stop gained, stop lost, or splice acceptor/donor

722 variants) and if the variants were predicted to be deleterious by SIFT or PolyPhen.

724 Replication

725 We replicated our novel asthma, hay fever/eczema and eczema loci in two

726 independent cohorts the EAGLE eczema consortium and the GABRIEL asthma

727 consortium ($P\leq0.05$). Our novel loci identified for asthma was replicated using the

728 summary statistics from the GABRIEL consortium which consisted of 10,365

729 physician-diagnosed asthmatic cases and 16,100 healthy controls (60). All individuals

730 in GABRIEL were genotyped for 582,892 SNPs using the Illumina Human610 quad

731 array. More information on this cohort has been published elsewhere (60). The

732 EAGLE consortium GWA summary statistics consists of 21,000 atopic dermatitis

733 (eczema) cases and 96,000 controls (61) and were used to replicate novel loci for hay

734 fever/eczema and eczema analysed separately. Further information about this cohort

735 has been published previously (61). If the lead SNP from our study was not found in

736 GABRIEL or EAGLE, we search for a proxy in LD (≥0.8) with the lead SNP.

738 SNP-based heritability

To quantify the SNP-based heritability for asthma and for hay fever/eczema

(combined as a single phenotype) we used LD score regression software (LDSC) (19)

including the same cases and controls as for the association analysis for each

phenotype (19). To calculate the heritability on the liability scale, we needed to adjust

for disease prevalence. Since this was a population-based study, we set the Caucasian

population and sample prevalence to the one calculated for each disease in UK

Biobank. We included 1,108,908 HapMap SNPs to calculate the heritability for

asthma and hay fever/eczema. We also removed all significant loci from each

individual GWA study result to estimate how much of the heritability was explained

by the significant loci reported in this study.

<u>Identification of phenotype-specific loci (SNP)</u>

To identify possible phenotype-specific SNPs, we performed polytomous

(multinomial) logistic regression to identify whether the effect of a locus (lead SNP)

was significantly (FDR≤0.05) larger for one disease phenotype as compared to

another. These effects can therefore be considered as being disease/phenotype

specific. To conduct these analyses, we used four non-overlapping groups: 1) asthma

cases without hay fever/eczema (N=22,858), 2) hay fever/eczema cases without

asthma (N=65,063), 3) asthma cases with hay fever/eczema (only including N=19,299

participants that had reported asthma in combination with hay fever or eczema), and

4) controls without asthma, hay fever and eczema (N=240,817) (Figure 2). Hay fever

and eczema were not separated in this analysis due to the small sample size (Table 1).

We performed polytomous logistic regression for all possibly independent ($R^2<=0.8$)

associated lead SNPs identified in the asthma, hay fever/eczema or asthma/hay

fever/eczema GWA studies. For some regions, different SNPs, that represent the same

signal (R^2>0.8 between the SNPs), were identified in the different GWA studies. For

these regions, only the SNP with the lowest P-value from the original GWA study

was included in these analyses. For regions where, different lead SNPs were

identified in the different GWA studies, and where these lead SNPs were not in strong

LD (R^2<=0.8), all lead SNPs were included in the analyses.

The polytomous (multinomial) logistic regression was performed with the response

variable, *Y,* being categorically distributed with K=4 non-overlapping

groups/outcomes (the four non-overlapping groups are explained above). Out of

K·(K-1)/2=6 comparisons in total, there are K-1=3 independent comparisons. The

logit function is defined as the logarithm of the quotient between the probability of a

given outcome (e.g., P(Y=1)) and the probability of a reference or pivot outcome (i.e.,

P(Y=4) in our case). This function is assumed to be linear in all explanatory variables,

including covariates and the specific SNP under consideration. Note that the beta

estimates (i.e., the log-odds ratios) are unique for each comparison. The polytomous

(multinomial) regression was performed using multinom in the R library nnet for the

three independent odds: P(*Y*=1)/P(*Y*=4), P(*Y*=2)/P(*Y*=4), and P(*Y*=3)/P(*Y*=4). Beta

estimates, standard errors, and p-values (two-sided, normal approximation) for the

remaining comparisons between phenotypic outcomes (i.e., P(*Y*=1)/P(*Y*=2),

P(*Y*=1)/P(*Y*=3), and P(*Y*=2)/P(*Y*=3)) were calculated from the model output such that,

e.g., $beta_{12} = beta_{14} - beta_{24}$ and $se_{12}^2 = se_{14}^2 + se_{24}^2$, where the first subscript denotes

the outcome of interest while the second subscript denotes the reference outcome.

To determine whether the lead SNPs were specific to one disease phenotype or shared

among phenotypes, we identified for which disease phenotype the OR was the highest

(we used the value of the OR rather than the most significant P-value in order not to

be influenced by the different power in the phenotype groups due to different sample-

sizes), and whether the OR was significantly (FDR $\leq$ 0.05) higher compared to the

other disease phenotypes. As a threshold for significance, we used an FDR

(Benjamini-Hochberg) value of 0.05, corresponding to a nominal P-value of < 0.017

in the three sets of cases vs cases analyses. In our analyses, an FDR adjustment is to

prefer (in favour of Bonferroni) due to its power to pinpoint as many positive findings

as possible, while retaining a low false-discovery rate (5% in our case).

Results were plotted as Venn diagrams to show the pair-wise overlap between disease

phenotypes. If two SNPs from the same locus that were not in LD with each other (R^2

<= 0.8) were assigned to the same area, the locus only occurs once in the Venn

diagram. However, for a few loci, multiple unlinked (R^2 <= 0.8) SNPs from the same

locus were assigned to different areas. Such loci were included at multiple locations

in the Venn diagram together with the name of the SNP (i.e., gene_SNP).

Ethics

UK Biobank was given ethical approval by the North West Multicentre Research

Ethics Committee, the National Information Governance Board for Health and Social

Care and the Community Health Index Advisory Group. UK Biobank holds a generic

Research Tissue Bank approval granted by the National Research Ethics Service

(http://www.hra.nhs.uk/) that lets applicants conduct research on UK Biobank data

without obtaining ethical approvals for each separate project. Access to UK Biobank

genetic and phenotypic data was given through the UK Biobank Resource under

Application Number 15479. All participants provided signed consent to participate in

UK Biobank.

Data availability

The genotypes and phenotypes included in the current study are available from the

UK Biobank data, which can be accessed by researchers upon application

(https://www.ukBiobank.ac.uk/). Summary statistics and codes used for this project

can be accessed by contacting the corresponding author.

URLs

UK Biobank, http://www.ukBiobank.ac.uk; PLINK, https://www.cog-

genomics.org/plink2 ; NHGRI-EBI GWAS Catalog, https://www.ebi.ac.uk/gwas/;

Software tool for LD Score estimation and estimation of variance components from

summary statistics, https://github.com/bulik/ldsc/; GCTA,

http://cnsgenomics.com/software/gcta/; BiomaRt, http://www.bioconductor.org;

GTEx, https://www.gtexportal.org/home/; DGIdb, http://www.dgidb.org/.

Acknowledgement

We acknowledge all the participants and the administrative staff at the UK Biobank.

The computations were performed on resources provided by SNIC through Uppsala

Multidisciplinary Centre for Advanced Computational Science (UPPMAX) under

projects b2016021, b2017059, sens2017538, and sens2017541. The work was

supported by grants from the Swedish Society for Medical Research (SSMF), the

Kjell and Märta Beijers Foundation, Göran Gustafssons Foundation, the Swedish

Medical Research Council (Project Number 2015-03327), the Marcus Borgström

Foundation, the Åke Wiberg Foundation, the Borgström Hedström Foundation and

the Swedish Heart-Lung Foundation.

Author contributions

Planned the study (WEE and ÅJ), analysed the data (WEE, ÅJ, TK), literature search

(WEE), Figures (WEE, MRA, ÅJ), data interpretation (WEE, ÅJ, TK, MRA), writing

of manuscript (WEE, ÅJ, TK, MRA).

References

1. Ober,C. and Yao,T.C. (2011) The genetics of asthma and allergic disease: A 21st

century perspective. *Immunol. Rev.*, **242**, 10–30.

2. Marenholz,I., Esparza-Gordillo,J., Ruschendorf,F., Bauerfeind,A., Strachan,D.P.,

Spycher,B.D., Baurecht,H., Margaritte-Jeannin,P., Saaf,A., Kerkhof,M., *et al.*

(2015) Meta-analysis identifies seven susceptibility loci involved in the atopic

march. *Nat. Commun.*, **6**, 8804.

3. Ferreira,M.A.R., Matheson,M.C., Tang,C.S., Granell,R., Ang,W., Hui,J.,

Kiefer,A.K., Duffy,D.L., Baltic,S., Danoy,P., *et al.* (2014) Genome-wide

association analysis identifies 11 risk variants associated with the asthma with

hay fever phenotype. *J. Allergy Clin. Immunol.*, **133**.

4. Hinds,D.A., McMahon,G., Kiefer,A.K., Do,C.B., Eriksson,N., Evans,D.M., St

Pourcain,B., Ring,S.M., Mountain,J.L., Francke,U., *et al.* (2013) A genome-wide

association meta-analysis of self-reported allergy identifies shared and allergy-

specific susceptibility loci. *Nat. Genet.*, **45**, 907–11.

5. Portelli,M. a, Hodge,E. and Sayers,I. (2015) Genetic risk factors for the

development of allergic disease identified by genome-wide association. *Clin.*

Exp. Allergy, **45**, 21–31.

6. Ferreira,M.A., Vonk,J.M., Baurecht,H., Marenholz,I., Tian,C., Hoffman,J.D.,

Helmer,Q., Tillander,A., Ullemar,V., Van Dongen,J., *et al.* (2017) Shared

genetic origin of asthma, hay fever and eczema elucidates allergic disease

biology. *Nat. Genet.*, **49**, 1752–1757.

7. Demenais,F., Margaritte-Jeannin,P., Barnes,K.C., Cookson,W.O.C., Altmüller,J.,

Ang,W., Barr,R.G., Beaty,T.H., Becker,A.B., Beilby,J., *et al.* (2018)

Multiancestry association study identifies new asthma risk loci that colocalize

with immune-cell enhancer marks. *Nat. Genet.*, 10.1038/s41588-017-0014-7.

8. Waage,J., Standl,M., Curtin,J.A., Jessen,L.E., Thorsen,J., Tian,C., Schoettler,N.,

Flores,C., Abdellaoui,A., Ahluwalia,T.S., *et al.* (2018) Genome-wide association

and HLA fine-mapping studies identify risk loci and genetic pathways

underlying allergic rhinitis. *Nat. Genet.*, 10.1038/s41588-018-0157-1.

9. Zhu,Z., Lee,P.H., Chaffin,M.D., Chung,W., Loh,P.R., Lu,Q., Christiani,D.C. and

Liang,L. (2018) A genome-wide cross-trait analysis from UK Biobank highlights

the shared genetic architecture of asthma and allergic diseases. *Nat. Genet.*,

10.1038/s41588-018-0121-0.

10. Ullemar,V., Magnusson,P.K.E., Lundholm,C., Zettergren,A., Melén,E.,

Lichtenstein,P. and Almqvist,C. (2016) Heritability and confirmation of genetic

association studies for childhood asthma in twins. *Allergy Eur. J. Allergy Clin.*

Immunol., **71**, 230–238.

11. Fagnani,C., Annesi-Maesano,I., Brescianini,S., D'Ippolito,C., Medda,E.,

Nisticò,L., Patriarca,V., Rotondi,D., Toccaceli,V. and Stazi,M.A. (2008)

Heritability and shared genetic effects of asthma and hay fever: an Italian study

of young twins. *Twin Res. Hum. Genet.*, **11**, 121–131.

12. Bataille,V., Lens,M. and Spector,T.D. (2012) The use of the twin model to

investigate the genetics and epigenetics of skin diseases with genomic,

transcriptomic and methylation data. *J. Eur. Acad. Dermatol. Venereol.*, **26**,

1067–1073.

13. Moffatt,M.F., Gut,I.G., Demenais,F., Strachan,D.P., Bouzigon,E., Heath,S., von

Mutius,E., Farrall,M., Lathrop,M. and Cookson,W.O.C.M. (2010) A large-scale,

consortium-based genomewide association study of asthma. *N. Engl. J. Med.*,

363, 1211–21.

14. Yang,J., Ferreira,T., Morris,A.P., Medland,S.E., Madden,P.A.F., Heath,A.C.,

Martin,N.G., Montgomery,G.W., Weedon,M.N., Loos,R.J., *et al.* (2012)

Conditional and joint multiple-SNP analysis of GWAS summary statistics

identifies additional variants influencing complex traits. *Nat. Genet.*, **44**, 369–

375.

15. Hirota,T., Takahashi,A., Kubo,M., Tsunoda,T., Tomita,K., Sakashita,M.,

Yamada,T., Fujieda,S., Tanaka,S., Doi,S., *et al.* (2012) Genome-wide association

study identifies eight new susceptibility loci for atopic dermatitis in the Japanese

population. *Nat. Genet.*, **44**, 1222–1226.

16. Adzhubei,I.A., Schmidt,S., Peshkin,L., Ramensky,V.E., Gerasimova,A., Bork,P.,

Kondrashov,A.S. and Sunyaev,S.R. (2010) A method and server for predicting

damaging missense mutations. *Nat. Methods*, **7**, 248–249.

17. Ng, C.,P. and Henikoff,S. (2001) Predicting deleterious amino acid substitutions.

Genome Res., **11**, 863–874.

18. Pickrell,J.K., Berisa,T., Liu,J.Z., Ségurel,L., Tung,J.Y. and Hinds,D.A. (2016)

Detection and interpretation of shared genetic influences on 42 human traits. *Nat.*

Genet., **48**, 709–717.

19. Bulik-Sullivan,B.K., Loh,P.-R., Finucane,H.K., Ripke,S., Yang,J., Patterson,N.,

Daly,M.J., Price,A.L. and Neale,B.M. (2015) LD Score regression distinguishes

confounding from polygenicity in genome-wide association studies. *Nat. Genet.*,

47, 291–295.

20. Astle,W.J., Elding,H., Jiang,T., Allen,D., Ruklisa,D., Mann,A.L., Mead,D.,

Bouman,H., Riveros-Mckay,F., Kostadima,M.A., *et al.* (2016) The Allelic

Landscape of Human Blood Cell Trait Variation and Links to Common Complex

Disease. *Cell*, **167**, 1415–1429.e19.

21. De Lange,K.M., Moutsianas,L., Lee,J.C., Lamb,C.A., Luo,Y., Kennedy,N.A.,

Jostins,L., Rice,D.L., Gutierrez-Achury,J., Ji,S.G., *et al.* (2017) Genome-wide

association study implicates immune activation of multiple integrin genes in

inflammatory bowel disease. *Nat. Genet.*, **49**, 256–261.

22. Tenesa,A., Farrington,S.M., Prendergast,J.G.D., Porteous,M.E., Walker,M.,

Haq,N., Barnetson,R.A., Theodoratou,E., Cetnarskyj,R., Cartwright,N., *et al.*

(2008) Genome-wide association scan identifies a colorectal cancer

susceptibility locus on 11q23 and replicates risk loci at 8q24 and 18q21. *Nat.*

Genet., **40**, 631–637.

23. Cho,M.H., Castaldi,P.J., Hersh,C.P., Hobbs,B.D., Barr,R.G., Tal-Singer,R.,

Bakke,P., Gulsvik,A., San José Estépar,R., Van Beek,E.J.R., *et al.* (2015) A

genome-wide association study of emphysema and airway quantitative imaging

phenotypes. *Am. J. Respir. Crit. Care Med.*, **192**, 559–569.

24. Polte,T., Behrendt,A.-K. and Hansen,G. (2006) Direct evidence for a critical role

of CD30 in the development of allergic asthma. *J. Allergy Clin. Immunol.*,

10.1016/j.jaci.2006.07.014.

25. Ameur,A., Enroth,S., Johansson,A., Zaboli,G., Igl,W., Johansson,A.C.,

Rivas,M.A., Daly,M.J., Schmitz,G., Hicks,A.A., *et al.* (2012) Genetic adaptation

of fatty-acid metabolism: a human-specific haplotype increasing the biosynthesis

of long-chain omega-3 and omega-6 fatty acids. *Am J Hum Genet*, **90**, 809–820.

26. Standl,M., Sausenthaler,S., Lattka,E., Koletzko,S., Bauer,C.P., Wichmann,H.E.,

Von Berg,A., Berdel,D., Krämer,U., Schaaf,B., *et al.* (2012) FADS gene cluster

modulates the effect of breastfeeding on asthma. Results from the GINIplus and

LISAplus studies. *Allergy Eur. J. Allergy Clin. Immunol.*, **67**, 83–90.

27. Martinelli,N., Girelli,D., Malerba,G., Guarini,P., Illig,T., Trabetti,E., Sandri,M.,

Friso,S., Pizzolo,F., Schaeffer,L., *et al.* (2008) FADS genotypes and desaturase

activity estimated by the ratio of arachidonic acid to linoleic acid are associated

with inflammation and coronary artery disease. *Am. J. Clin. Nutr.*, **88**, 941–949.

28. Barman,M., Johansson,S., Hesselmar,B., Wold,A.E., Sandberg,A.S. and

950 Sandin,A. (2013) High Levels of Both n-3 and n-6 Long-Chain Polyunsaturated

951 Fatty Acids in Cord Serum Phospholipids Predict Allergy Development. *PLoS*

952 *One*, **8**.

953 29. Graham,D.B., Lefkovith,A., Deelen,P., de Klein,N., Varma,M., Boroughs,A.,

954 Desch,A.N., Ng,A.C.Y., Guzman,G., Schenone,M., *et al.* (2016) TMEM258 Is a

955 Component of the Oligosaccharyltransferase Complex Controlling ER Stress and

956 Intestinal Inflammation. *Cell Rep.*, **17**, 2955–2965.

957 30. Esparza-Gordillo,J., Matanovic,A., Marenholz,I., Bauerfeind,A., Rohde,K.,

958 Nemat,K., Lee-Kirsch,M.A., Nordenskjöld,M., Winge,M.C.G., Keil,T., *et al.*

959 (2015) Maternal Filaggrin Mutations Increase the Risk of Atopic Dermatitis in

960 Children: An Effect Independent of Mutation Inheritance. *PLoS Genet.*, **11**.

961 31. Simmonds,M. and Gough,S. (2007) The HLA Region and Autoimmune Disease:

962 Associations and Mechanisms of Action. *Curr. Genomics*, **8**, 453–465.

963 32. Ferreira,M.A.R., Vonk,J.M., Baurecht,H., Marenholz,I., Tian,C., Hoffman,J.D.,

964 Helmer,Q., Tillander,A., Ullemar,V., Lu,Y., *et al.* (2018) Eleven loci with new

965 reproducible genetic associations with allergic disease risk. *J. Allergy Clin.*

966 *Immunol.*, 10.1016/j.jaci.2018.03.012.

967 33. Schuttelaar,M.L.A., Kerkhof,M., Jonkman,M.F., Koppelman,G.H., Brunekreef,B.,

968 De Jongste,J.C., Wijga,A., McLean,W.H.I. and Postma,D.S. (2009) Filaggrin

969 mutations in the onset of eczema, sensitization, asthma, hay fever and the

970 interaction with cat exposure. *Allergy Eur. J. Allergy Clin. Immunol.*, **64**, 1758–

971 1765.

972 34. Spergel,J.M. (2010) From atopic dermatitis to asthma: The atopic march. *Ann.*

973 *Allergy, Asthma Immunol.*, 10.1016/j.anai.2009.10.002.

974 35. Illi,S., Von Mutius,E., Lau,S., Nickel,R., Grüber,C., Niggemann,B., Wahn,U.,

975 Wahn,V., Groeger,M., Zepp,F., *et al.* (2004) The natural course of atopic

dermatitis from birth to age 7 years and the association with asthma. *J. Allergy*

Clin. Immunol., 10.1016/j.jaci.2004.01.778.

36. Hirai,H., Pang,Z., Bao,D., Miyazaki,T., Li,L., Miura,E., Parris,J., Rong,Y.,

Watanabe,M., Yuzaki,M., *et al.* (2005) Cbln1 is essential for synaptic integrity

and plasticity in the cerebellum. *Nat. Neurosci.*, 10.1038/nn1576.

37. Gleichman,A.J. (2012) Glutamate receptor autoimmune diseases: Understanding

the mechanisms, determining the epitopes, and developing better tests. *ProQuest*

Diss. Theses.

38. Hunt,K.A., Zhernakova,A., Turner,G., Heap,G.A.R., Franke,L., Bruinenberg,M.,

Romanos,J., Dinesen,L.C., Ryan,A.W., Panesar,D., *et al.* (2008) Newly

identified genetic risk variants for celiac disease related to the immune response.

Nat. Genet., 10.1038/ng.102.

39. Sawcer,S., Hellenthal,G., Pirinen,M., Spencer,C.C.A., Patsopoulos,N.A.,

Moutsianas,L., Dilthey,A., Su,Z., Freeman,C., Hunt,S.E., *et al.* (2011) Genetic

risk and a primary role for cell-mediated immune mechanisms in multiple

sclerosis. *Nature*, 10.1038/nature10251.

40. Gupta,J., Johansson,E., Bernstein,J.A., Chakraborty,R., Khurana Hershey,G.K.,

Rothenberg,M.E. and Mersha,T.B. (2016) Resolving the etiology of atopic

disorders by using genetic analysis of racial ancestry. *J. Allergy Clin. Immunol.*,

10.1016/j.jaci.2016.02.045.

41. Sleiman,P.M.A., Flory,J., Imielinski,M., Bradfield,J.P., Annaiah,K., Willis-

Owen,S.A.G., Wang,K., Rafaels,N.M., Michel,S., Bonnelykke,K., *et al.* (2010)

Variants of DENND1B associated with asthma in children. *N. Engl. J. Med.*,

10.1056/NEJMoa0901867.

42. Israel,E., Lasky-Su,J., Markezich,A., Damask,A., Szefler,S.J., Schuemann,B.,

Klanderman,B., Sylvia,J., Kazani,S., Wu,R., *et al.* (2015) Genome-wide

association study of short-acting β2-agonists a novel genome-wide significant

locus on chromosome 2 near ASB3. *Am. J. Respir. Crit. Care Med.*,

10.1164/rccm.201408-1426OC.

43. Himes,B.E., Hunninghake,G.M., Baurley,J.W., Rafaels,N.M., Sleiman,P.,

Strachan,D.P., Wilk,J.B., Willis-Owen,S.A.G., Klanderman,B., Lasky-Su,J., *et*

al. (2009) Genome-wide Association Analysis Identifies PDE4D as an Asthma-

Susceptibility Gene. *Am. J. Hum. Genet.*, 10.1016/j.ajhg.2009.04.006.

44. Yucesoy,B., Kaufman,K.M., Lummus,Z.L., Weirauch,M.T., Zhang,G., Cartier,A.,

Boulet,L.-P., Sastre,J., Quirce,S., Tarlo,S.M., *et al.* (2015) Genome-Wide

Association Study Identifies Novel Loci Associated With Diisocyanate-Induced

Occupational Asthma. *Toxicol. Sci.*, **146**, 192–201.

45. Ioannidis,J.P.A. (2005) Why most published research findings are false. *PLoS*

Med., 10.1371/journal.pmed.0020124.

46. Brocker,C.N., Vasiliou,V. and Nebert,D.W. (2009) Evolutionary divergence and

functions of the ADAM and ADAMTS gene families. *Hum. Genomics*,

10.1186/1479-7364-4-1-43.

47. Rock,F.L., Hardiman,G., Timans,J.C., Kastelein,R.A. and Bazan,J.F. (1998) A

family of human receptors structurally related to Drosophila Toll. *Proc. Natl.*

Acad. Sci., 10.1073/pnas.95.2.588.

48. Clark,J., Vagenas,P., Panesar,M. and Cope,A.P. (2005) What does tumour

necrosis factor excess do to the immune system long term? In *Annals of the*

Rheumatic Diseases.

49. Zuk,O., Hechter,E., Sunyaev,S.R. and Lander,E.S. (2012) The mystery of missing

heritability: Genetic interactions create phantom heritability. *Proc. Natl. Acad.*

Sci., **109**, 1193–1198.

50. Visscher,P.M., McEvoy,B. and Yang,J. (2010) From Galton to GWAS:

quantitative genetics of human height. *Genet. Res. (Camb).*, **92**, 371–379.

51. Golan,D., Lander,E.S. and Rosset,S. (2014) Measuring missing heritability:

Inferring the contribution of common variants. *Proc. Natl. Acad. Sci.*, **111**,

E5272–E5281.

52. Pawankar,R., Mori,S., Ozu,C. and Kimura,S. (2011) Overview on the

pathomechanisms of allergic rhinitis. *Asia Pac. Allergy*,

10.5415/apallergy.2011.1.3.157.

53. UK10K Consortium (2015) The UK10K project identifies rare variants in health

and disease. *Nature*, **526**, 82–90.

54. Huang,J., Ellinghaus,D., Franke,A., Howie,B. and Li,Y. (2012) 1000 Genomes-

based imputation identifies novel and refined associations for the Wellcome

Trust Case Control Consortium phase 1 Data. *Eur. J. Hum. Genet.*, **20**, 801–805.

55. McCarthy,S., Das,S., Kretzschmar,W., Delaneau,O., Wood,A.R., Teumer,A.,

Kang,H.M., Fuchsberger,C., Danecek,P., Sharp,K., *et al.* (2016) A reference

panel of 64,976 haplotypes for genotype imputation. *Nat. Genet.*, **48**, 1279–1283.

56. Purcell,S., Neale,B., Todd-Brown,K., Thomas,L., Ferreira,M.A.R., Bender,D.,

Maller,J., Sklar,P., de Bakker,P.I.W., Daly,M.J., *et al.* (2007) PLINK: A tool set

for whole-genome association and population-based linkage analyses. *Am. J.*

Hum. Genet., **81**, 559–575.

57. Fadista,J., Manning,A.K., Florez,J.C. and Groop,L. (2016) The (in)famous

GWAS P-value threshold revisited and updated for low-frequency variants. *Eur.*

J. Hum. Genet., **24**, 1202–1205.

58. Carithers,L.J. and Moore,H.M. (2015) The Genotype-Tissue Expression (GTEx)

Project. *Biopreserv. Biobank.*, **13**, 307–308.

59. Durinck,S., Moreau,Y., Kasprzyk,A., Davis,S., De Moor,B., Brazma,A. and

Huber,W. (2005) BioMart and Bioconductor: A powerful link between

biological databases and microarray data analysis. *Bioinformatics*, **21**, 3439–

3440.

60. Moffatt,M.F., Gut,I.G., Demenais,F., Strachan,D.P., Bouzigon,E., Heath,S., von

Mutius,E., Farrall,M., Lathrop,M. and Cookson,W.O.C.M. (2010) A large-scale,

consortium-based genomewide association study of asthma. *N. Engl. J. Med.*,

363, 1211–1221.

61. Paternoster,L., Standl,M., Waage,J., Baurecht,H., Hotze,M., Strachan,D.P.,

Curtin,J.A., Bønnelykke,K., Tian,C., Takahashi,A., *et al.* (2015) Multi-ancestry

genome-wide association study of 21,000 cases and 95,000 controls identifies

new risk loci for atopic dermatitis. *Nat. Genet.*, **47**, 1449–1456.

1064

Figure Legends

Figure 1. Manhattan plots for asthma, for hay fever and/or eczema, and for asthma and/or hay fever and/or eczema (combined) for autosomal chromosomes. The black horizontal line indicates the genome wide threshold (3×10^{-8}). The black regions represent novel loci found in this study.

Figure 2. Comorbidity between asthma and hay fever/eczema.

Figure 3. Venn diagram showing the phenotype-specificity of the GWA loci, based on the results from the polytomous logistic regression analyses. The Venn diagram show loci (SNPs) that are specific (significantly larger effect) to or shared between (no

significant difference in effects) between two non-overlapping groups of cases. The name of each locus is denoted by the most likely gene(s). At some of the loci (e.g. *IL2RA, LPP,* and *IL4R*), more than one independent (R2<0.8) lead SNP has been analysed in the polytomous logistic regression. If those showed different specificity pattern, they have been included twice in the figure with the name of respective lead SNP(s) also included in the locus name. P-values and estimates for the genes can be found in Table S16 where the area number 1 (green in the figure) indicates specificity for the asthma only, 2 (blue in the figure) specificity for hay fever/eczema only and area number 3 (white in the figure) specificity for asthma with hay fever/eczema (significantly larger estimate in the asthma with hay fever/eczema group of cases).

S1 Figure QQ-plot for Asthma in UK Biobank. The red line denotes the expected null-line of no association.

S2 Figure Regional plot for the missense variant rs2230624 in asthma.

S3 Figure QQ-plot for Hay fever and/or Eczema (combined) in UK Biobank. The red line denotes the expected null-line of no association.

S4 Figure Manhattan plots for Hay Fever and/or Eczema (combined), Hay Fever (only) and Eczema (only) analysed in UK Biobank for autosomal chromosomes. The black horizontal line indicates the genome wide threshold ($3x10^{-8}$). The black regions in the hay fever and/or eczema (combined) plot represent novel loci found in this study and the black regions in the hay fever (only) and eczema (only) plot represent two novel loci, not found in previous GWA studies or in our combined hay fever / eczema analysis.

S5 Figure QQ-plot for Hay fever (only) in UK Biobank. The red line denotes the expected null-line of no association.

S6 Figure QQ-plot for Eczema (only) in UK Biobank. The red line denotes the expected null-line of no association.

S7 Figure QQ-plot for Asthma and/or Hay fever and/or Eczema (combined) in UK Biobank. The red line denotes the expected null-line of no association.

Tables

Table 1. Baseline characteristics of Caucasian participants in UK Biobank.

	Asthma	Hay fever /eczema	Asthma/hay fever/eczema combined[b]	Hay fever	Eczema	Controls[c]
N Caucasians[a] (prior to QC)	51,645	102,862	130,865	22,919	9,578	294,477
N total included after QC[d]	41,934	84,050	106,772	18,915	7,884	239,773
N (%) males after QC	21,730 (51.8%)	42,639 (50.7%)	55,124 (51.6%)	8,692 (46.0%)	3,365 (42,7%)	138,666 (57.8%)
Age year span (mean)	38-70 (56.1)	39-72 (55.4)	38-72 (55.7)	40-77 (55.0)	40-70 (55.0)	39-73 (57.2)
Townsend deprivation index range (mean)	-6.3-10.6 (-1.3)	-6.3-10.6 (-1.6)	-6.3-10.6 (-1.5)	-6.3-10.4 (-1.7)	-6.3-9.6 (-1.9)	-6.3-10.9 (-1.5)
% Ever smoked (N yes / N no)	60.4% (31,040 / 20,390)	58.5% (59,983 / 42,616)	59.3% (77,335/53,121)	56.8% (12,986/9,879)	60.9% (5,815/3,737)	60.4% (177,212/116,275)

[a]The total number of Caucasians is N = 443,068.

[b]Asthma or hay fever and/or eczema combined as one phenotype.

[c]The same controls were used in all analyses.

[d] We removed first and second-degree relatives, using kinship data (estimated genetic relationship > 0.044), and participants with sex discordance, high heterozygosity, and participant with more than 5% missing SNP genotypes, resulting in 346,545 individuals after QC.

Table 2. Summary results for the 15 novel loci significantly associated with self-reported asthma in UK Biobank ($P \leq 3 \times 10^{-8}$) with replication in the GABRIEL cohort.

lead SNP	locus[a] chr:start-end (kbp)	N snps (total[b]/ independent[c])	MAF[d]	Minor/ major allele	OR (95% CI) for minor allele	P	Likely target gene (s)	GABRIEL P OR (95% CI) for effective allele (minor/effective allele)
rs2230624	1:12,175-12,175	1/1	0.02	A/G	0.80 (0.75-0.86)	1.01×10^{-10}	*TNFRSF8*[e]	No proxy
rs2296618	1:198,656-198,670	5/1	0.13	G/A	0.93 (0.91-0.96)	8.03×10^{-9}	*PTPRC*[f]	No proxy
rs10934853	3:127,886-128,075	3/1	0.27	A/C	0.95 (0.94-0.97)	2.20×10^{-8}	*EEFSEC*[g]	P=0.006 OR=1.06 (1.02-1.11) A/C
rs6778937	3:176,708-176,868	28/1	0.28	C/T	0.95 (0.93-0.97)	2.54×10^{-9}	*TBL1XR1*[f]	No proxy
rs11466773	5:156,930-156,988	7/1	0.06	T/C	1.09 (1.06-1.13)	6.32×10^{-9}	*ADAM19*[g]	No proxy
rs2614266	6:135,691-135,818	6/1	0.44	A/T	1.05 (1.03-1.06)	8.90×10^{-9}	*AHI1*[f]	No proxy
rs10215232	7:3,062-3,153	12/1	0.12	G/C	0.93 (0.91-0.95)	1.53×10^{-9}	*SDK1*[f]	rs9986945[h], R^2=1.0[i] P=0.03 OR=1.07 (1.01-1.14) T/G
rs41283642	9:101,915-101,989	3/1	0.03	T/C	0.86 (0.82-0.90)	1.27×10^{-11}	*TGFBR1*[g]	No proxy
rs2497318	10:94,34-94,44	24/1	0.45	T/C	0.95 (0.94-0.97)	3.21×10^{-10}	*HHEX*[g]	Rs10882091[h], R^2=0.81[i] P=0.84 OR=1.00 (0.97-1.05) C/T
rs174535	11:61,543-61,623	49/1	0.35	C/T	0.95 (0.93-0.96)	1.02×10^{-11}	*MYRF*[e], *TMEM258*[g]	rs102275[h], R^2=1.0[i] P=0.045

								OR=1.04 (1.00-1.09) C/T
rs11178649	12:71,409-71,585	103/1	0.41	T/G	0.95 (0.93-0.96)	2.68×10^{-11}	*TSPAN8*[e]	rs1051334[h] $R^2=1.0$[i] P=0.04 OR=0.95 (0.92-0.99) (G/T)
rs4761592	12:94,556-94,604	17/1	0.15	T/C	0.93 (0.92-0.95)	1.27×10^{-9}	*PLXNCl*[f]	rs3912394[h], $R^2=0.85$[i] P=0.021 OR=1.07 (1.01-1.13) T/C
rs9316059	13:44,475-44,490	5/1	0.20	T/A	1.06 (1.04-1.08)	1.35×10^{-8}	*LINC00284*[f]	rs3764147[h], $R^2=0.93$[i] P=0.58 OR=1.01 (0.97-1.06) A/G
rs4842921	15:84,556-84,556	1/1	0.39	A/G	0.96 (0.94-0.97)	2.63×10^{-8}	*ADAMTSL3*[f]	No proxy
rs11671106	19:46,219-46,370	28/1	0.35	T/C	0.95 (0.94-0.97)	8.29×10^{-10}	*BHMG1*[e]	rs7250497[h], $R^2=0.97$[i] P=0.05 OR=1.04 (1.00-1.09) G/A

More details can be found in S1-S3 and S4 Tables.

[a] Defined as SNPs located < 1 Mb apart containing at least one significantly associated genetic variant at $P \leq 3 \times 10^{-8}$.

[b] Total number of SNPs with $P \leq 3 \times 10^{-8}$ within loci.

[c] Total number of independent associations within the locus, based on conditional analysis (14).

[d] Minor allele frequency.

[e] Lead SNP is in LD ($R^2 > 0.8$) with a missense variant.

[f] Gene(s) closest to the lead SNP.

[g] Lead SNP is in LD ($R^2 > 0.8$) with the lead eQTL SNP. Information on tissue type can be found in S3 Table.

[h] proxy SNP in LD (>0.8) with lead SNP.

[i] R^2 between lead SNP in UK Biobank and proxy SNP in GABRIEL.

Table 3. Summary results for the 22 novel loci significantly associated with self-reported hay fever and/or eczema in UK Biobank (P ≤ 3x10^{-8}) with replication in the EAGLE cohort

Lead SNP	Locus[a] chr:start-stop (kbp)	N snps (total[b]/ independent[c])	MAF[d]	Minor/ major allele	P (OR)[e] [95%CI] hay fever/eczema	P (OR)[e] [95%CI] hay fever	P (OR)[e] [95%CI] eczema	Likely target gene	EAGLE P OR (95% CI) estimated for the minor allele (minor/major allele)
rs1201113	1:12,100-12,147	3/1	0.12	A/G	1.05x10^{-8} (0.95) [0.93-0.97]	4.67x10^{-3} (0.95) [0.92-0.99]	1.69x10^{-2} (0.94) [0.90-0.99]	*TNFRSF*[f]	P=0.18 OR=1.04 (0.98-1.10) A/G
rs906363	1:212,858-212,877	6/1	0.15	C/T	4.44x10^{-9} (1.05) [1.03-1.07]	8.56x10^{-4} (0.97) [1.02-1.08]	2.54x10^{-4} (1.09) [1.04-1.13]	*BATF3*[f]	P=1.52x10^{-6} OR=1.19 (1.07-1.17) C/T
rs13405815	2:28,623-28,644	9/1	0.46	T/C	3.01x10^{-9} (0.97) [0.95-0.98]	4.04x10^{-5} (0.96) [0.94-0.98]	3.70x10^{-3} (0.95) [0.92-0.98]	*RP11-373D23.3*[g]	Proxy rs6547850[h], R^2=1.0[i] P=0.0014 OR=0.95 (0.92-0.98) T/G
rs10185028	2:61,112-61,161	6/1	0.23	G/A	2.74x10^{-8} (1.04) [1.03-1.05]	9.47x10^{-4} (1.04) [1.02-1.07]	1.17x10^{-2} (1.05) [1.01-1.09]	*REL*[f]	P=0.0007 OR=1.08 (1.03-1.12) G/A
rs11717778	3:112,526-112,693	144/1	0.34	A/G	6.04x20^{-11} (0.96) [0.95-0.97]	2.67x10^{-4} (0.96) [0.94-0.98]	5.38x10^{-6} (0.92) [0.89-0.96]	*CD200R1L*[f]	P=0.0007 OR=0.94 (0.91-0.98) A/G
rs62379371	5:133,439-133,639	4/1	0.05	A/G	6.08x10^{-14} (0.90) [0.87-0.92]	7.83x10^{-6} (0.89) [0.84-0.94]	4.18x10^{-5} (0.85) [0.78-0.92]	*TCF7*[f]	P=0.62 OR=0.95 (0.77-1.17) A/G
rs13185930	5:137,461-137,605	10/1	0.25	A/G	1.12x10^{-8} (1.04) [1.03-1.05]	3.19x10^{-2} (1.03) [1.00-1.05]	4.64x10^{-1} (1.01) [0.98-1.05]	*GFRA3*[f]	P=0.42 OR=1.02 (0.98-1.06) A/G

rs2229768	6:25,823-26,239	24/1	0.24	C/T	8.20×10^{-11} (1.05) [1.03-1.06]	8.99×10^{-6} (1.06) [1.03-1.08]	2.74×10^{-1} (1.02) [0.98-1.06]	*U91328.19* [g]	P=0.80 OR=1.00 (0.96-1.04) C/T
rs1998266	6:36,349-36,380	5/1	0.14	T/C	1.70×10^{-8} (0.95) [0.94-0.97]	1.23×10^{-3} (0.95) [0.92-0.98]	1.68×10^{-3} (0.93) [0.88-0.97]	*ETV7* [f]	P=0.08 OR=0.95 (0.91-1.00) T/C
rs2746438	6:135,624-135,950	36/1	0.44	T/A	5.70×10^{-11} (1.04) [1.03-1.05]	9.24×10^{-7} (1.05) [1.03-1.08]	2.73×10^{-5} (1.07) [1.04-1.11]	*AHI1* [f]	P=0.21 OR=1.02 (0.99-1.06) T/A
rs3918226	7:150,690-150,690	1/1	0.08	T/C	5.65×10^{-10} (0.93) [0.91-0.95]	1.54×10^{-7} (0.90) [0.86-0.93]	6.09×10^{-1} (0.98) [0.93-1.05]	*NOS3* [f]	P=0.33 OR=1.04 (0.97-1.11) T/C
rs6986151	8:101,514-101,519	2/1	0.20	C/T	6.19×10^{-9} (1.04) [1.03-1.06]	5.39×10^{-6} (1.06) [1.04-1.09]	2.44×10^{-3} (1.06) [1.02-1.11]	*ANKRD46* [f]	P=0.31 OR=1.00 (0.95-1.05) C/T
rs1330303	9:16,715-16,756	2/1	0.35	T/C	5.21×10^{-10} (0.96) [0.95-0.97]	2.95×10^{-8} (0.94) [0.92-0.96]	3.90×10^{-1} (0.99) [0.95-1.02]	*BNC2* [f]	P=0.11 OR=1.01 (0.99-1.06) T/C
rs4743311	9:101,790-101,820	3/1	0.25	G/A	1.69×10^{-8} (1.04) [1.03-1.05]	7.93×10^{-6} (1.06) [1.03-1.08]	1.27×10^{-1} (1.03) [0.99-1.07]	*COL15A1* [f]	P=0.88 OR=1.00 (0.96-1.04) G/A
rs12343737	9:117,804-117,834	2/1	0.10	T/C	2.09×10^{-8} (0.95) [0.93-0.95]	1.28×10^{-6} (0.92) [0.89-0.95]	9.59×10^{-2} (0.96) [0.91-1.01]	*TNC* [f]	P=0.25 OR=0.97 (0.92-1.03) T/C
rs10986320	9:127,022-127,095	13/1	0.37	C/G	8.81×10^{-9} (1.04) [1.02-1.05]	4.13×10^{-5} (1.05) [1.02-1.07]	7.32×10^{-2} (1.03) [1.00-1.07]	*NEK6* [f]	P=0.20 OR=0.98 (0.95-1.01) C/G

rs4076542	11:2,237-2,296	4/1	0.37	A/G	1.74x10⁻⁸ (1.04) [1.02-1.05]	4.40x10⁻³ (1.04) [1.02-1.06]	7.78x10⁻¹ (1.02) [0.98-1.05]	*ASCL2*[f]	P=0.70 OR=0.99 (0.96-1.03) A/G
rs4939490	11:60,793-60,793	4/1	0.39	G/C	2.15x10⁻⁸ (0.97) [0.96-0.98]	2.15x10⁻³ (0.97) [0.95-0.99]	9.76x10⁻² (0.97) [0.94-1.01]	*CD6*[f]	P=0.25 OR=1.02 (0.99-1.06) G/C
rs3116590	13:50,808-50,811	2/1	0.21	G/A	1.00x10⁻⁸ (1.04) [1.03-1.06]	1.11x10⁻² (1.03) [1.01-1.06]	7.24x10⁻³ (1.05) [1.01-1.10]	*DLEU1*[f]	P=0.38 OR=1.02 (0.98-1.06) G/A
rs4771332	13:99,839-100,070	7/1	0.31	T/C	6.21x10⁻⁹ (0.96) [0.95-0.98]	6.09x10⁻³ (0.97) [0.95-0.99]	1.29x10⁻¹ (0.97) [0.94-1.01]	*UBAC2*	P=0.40 OR=0.98 (0.95-1.02) C/T
rs4381563	15:75,399-75,448	7/1	0.34	A/T	3.37x10⁻⁹ (0.96) [0.95-0.98]	2.98x10⁻³ (0.97) [0.95-0.99]	3.37x10⁻² (0.96) [0.93-1.00]	*PPCDC*[f]	P=0.89 OR=1.00 (0.97-1.04) A/T
rs6066184	**20:45,232-45,716**	**33/1**	**0.26**	**G/C**	**6.55x10⁻¹⁶ (0.95) [0.93-0.96]**	**5.25x10⁻¹¹ (0.92) [0.90-0.94]**	**2.03x10⁻² (0.96) [0.92-0.99]**	***EYA2*[f]**	**P=0.02 OR=1.05 (1.01-1.09) G/C**

More details can be found in S3 and S5-S7 Table.

[a] Defined as SNPs located < 1 Mb apart containing at least one significantly associated genetic variant at $P \leq 3 \times 10^{-8}$.

[b] Total number of SNPs with $P \leq 3 \times 10^{-8}$ within loci.

[c] Total number of independent associations within the locus, based on conditional analysis (14).

[d] Minor allele frequency.

[e] OR for minor allele.

[f] Gene(s) closest to the lead SNP.

[g] Lead SNP is in LD ($R^2 > 0.8$) with the lead eQTL SNP. Information on tissue type can be found in S3 Table.

[h] proxy SNP in LD (>0.8) with lead SNP.

[i] R^2 between lead SNP in UK Biobank and proxy SNP in EAGLE.

Table 4. Summary results for the 16 novel loci significantly associated with self-reported asthma and/or hay fever and/or eczema (combined) in UK Biobank (P $\leq 3 \times 10^{-8}$).

Lead SNP	Locus[a] chr:start-stop (kbp)	N snps (total[b]/ independent[c])	MAF[d]	Minor/ major allele	P (OR[e]) [95%CI] combined	P (OR[e]) [95%CI] asthma	P (OR[e]) [95%CI] hay fever/ eczema	Likely target gene
rs2230624	1:12,080-12,175	2/2	0.02	A/G	2.64×10^{-9} (0.87) [0.84-0.91]	1.01×10^{-10} (0.80) [0.75-0.86]	1.99×10^{-7} (0.88) [0.84-0.92]	*TNFRSF8*[f]
rs7410883	1:198,640-198,670	5/1	0.11	C/T	7.00×10^{-9} (0.95) [0.93-0.97]	2.26×10^{-8} (0.93) [0.91-0.95]	1.78×10^{-6} (0.96) [0.94-0.97]	*PTPRC*[g]
rs9816107	3:112,526-112,693	160/1	0.34	A/C	6.80×10^{-11} (0.96)	1.51×10^{-6} (0.96) [0.95-0.97]	8.39×10^{-11} (0.96) [0.95-0.97]	*CD200R1L*[g]
rs62379371	5:133,439-133,639	4/1	0.05	A/G	1.13×10^{-13} (0.91) [0.89-0.93]	1.23×10^{-7} (0.91) [0.88-0.94]	6.08×10^{-14} (0.90) [0.87-0.92]	*VDAC1*[g]
rs9379828	6:26,038-26,184	13/1	0.37	G/C	2.71×10^{-9} (1.03) [1.02-1.05]	1.07×10^{-10} (1.05) [1.04-1.07]	1.42×10^{-6} (1.03) [1.02-1.04]	*HIST1H2BD*[h]
rs1330303	9:16,715-16,715	1/1	0.35	T/C	2.84×10^{-8} (0.97) [0.96-0.98]	0.016 (0.98) [0.97-1.00]	5.21×10^{-10} (0.96) [0.95-0.97]	*BNC2*[g]
rs41283642	9:101,915-101,915	1/1	0.03	T/C	7.03×10^{-9} (0.92) [0.89-0.94]	1.27×10^{-11} (0.86) [0.82-0.90]	6.42×10^{-6} (0.93) [0.90-0.96]	*TGFBR1*[g]
rs3758212	9:127,002-127,178	14/1	0.37	T/C	2.99×10^{-9} (1.03) [1.02-1.05]	1.80×10^{-5} (1.04) [1.02-1.05]	2.18×10^{-8} (1.04) [1.02-1.05]	*NEK6*[g]
rs2505504	**10:43,728-43,763**	**17/1**	**0.29**	**A/G**	**5.45×10^{-9} (1.04)** [1.02-1.05]	**1.96×10^{-6} (1.04)** [1.02-1.06]	**4.61×10^{-7} (1.03)** [1.02-1.05]	***RASGEF1A***[g]

rs7114923	11:2,237-2,305	21/1	0.37	T/C	3.74x10⁻⁹ (1.03) [1.02-1.05]	1.20x10⁻⁵ (1.04) [1.02-1.05]	2.65x10⁻⁸ (1.04) [1.02-1.05]	*ASCL2*[g]	
rs3116590	13:50,808-50,808	1/1	0.21	G/A	1.52x10⁻⁸ (1.04) [1.03-1.05]	0.0064 (1.03) [1.01-1.05]	1.00x10⁻⁸ (1.04) [1.03-1.06]	*DLEU1*[g]	
rs61975764	**14:93,014-93,014**	**1/1**	**0.47**	**A/G**	**2.65x10⁻⁸ (1.03) [1.02-1.04]**	**1.46x10⁻⁷ (1.04) [1.03-1.06]**	**7.84x10⁻⁷ (1.03) [1.02-1.04]**	***RIN3*[g]**	
rs4381563	15:75,275-75,448	19/1	0.33	A/T	7.81x10⁻⁹ (0.97) [0.96-0.98]	0.00027 (0.97) [0.96-0.99]	3.37x10⁻⁹ (0.96) [0.95-0.97]	*PPCDC*[g]	
rs12956924	**18:46,451-46,451**	**1/1**	**0.31**	**A/G**	**1.52x10⁻⁸ (1.03) [1.02-1.05]**	**2.64x10⁻⁵ (1.04) [1.02-1.05]**	**2.09x10⁻⁷ (1.03) [1.02-1.05]**	***SMAD7*[g]**	
rs10419921	**19:16,412-16,412**	**1/1**	**0.30**	**T/C**	**1.67x10⁻⁸ (1.03) [1.02-1.05]**	**5.62x10⁻⁶ (1.04) [1.02-1.06]**	**1.09x10⁻⁶ (1.03) [1.02-1.04]**	***KLF2*[g]**	
rs6066184	20:45,228-45,716	26/2	0.26	G/C	1.19x10⁻¹³ (0.95) [0.94-0.97]	4.27x10⁻⁶ (0.96) [0.94-0.98]	6.55x10⁻¹⁶ (0.95) [0.93-0.96]	*EYA2*[d]	

Loci found only when analysing all three diseases as one phenotype are marked as bold. More details can be found in S3, S12-S13 and S15 Tables.

[a] Defined as SNPs located<1 Mb apart containing at least one significantly associated genetic variant at $P \leq 3\text{x}10^8$.

[b] Total number of SNPs with $P \leq 3\text{x}10^{-8}$ within loci.

[c] Total number of independent associations within the locus, based on conditional analysis (14).

[d] Minor allele frequency.

[e] OR for minor allele.

[f] Lead SNP is in LD ($R^2 > 0.8$) with a missense variant.

[g] Gene(s) closest to the lead SNP.

[h] Lead SNP is in LD ($R^2 > 0.8$) with the lead eQTL SNP. Information on tissue type can be found in S3 Table.

Table 5. SNP-based heritability in UK Biobank for asthma and hay fever/eczema (combined as one phenotype) estimated with LDSC (19).

| Phenotype | N cases | N controls | Prevalence used in LD-score[a] | | (A) All SNPs | | (B) Without significant loci | | h2 explained by significant loci |
			Population	Sample	h2	SE	h2	SE	Absolute terms: (A) - (B)
Asthma	41,926	239,751	0.117	0.117	0.210	0.017	0.168	0.010	0.042
Hay fever/ Eczema	84,034	239,751	0.232	0.232	0.160	0.010	0.124	0.007	0.036

[a]LDSC requires values for population and sample prevalence when estimating SNP heritability. In these analyses we used the prevalence in the UK Biobank cohort (Table 1) as the sample prevalence. However, due to the cross-sectional design of the UK Biobank we used the same values as population prevalence.

Abbreviations

SNP Single nucleotide polymorphism

GWA study Genome Wide Association Study