Probing the evolutionary robustness of anti-virulence treatments targeting iron

2 **uptake in *Pseudomonas aeruginosa***

4 Chiara Rezzoagli[1, *], David Wilson[1], Michael Weigert[1], Stefan Wyder[1], Rolf Kümmerli[1, *]

6 [1] Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland

8 * Corresponding authors:

Email: chiara.rezzoagli@uzh.ch (CR), rolf.kuemmerli@uzh.ch (RK)

Abstract

Antivirulence treatments are therapeutic strategies that inhibit the expression or functioning of bacterial virulence factors. They hold great promise to be both effective and exert weaker selection for resistance than conventional antibiotics. However, the evolutionary robustness argument, based on the idea that anti-virulence treatments disarm rather than kill pathogens, is controversial, and there is a lack of studies testing the potential for resistance evolution under conditions relevant for infections. Here, we approach this issue by probing the evolutionary robustness of two anti-virulence treatments, gallium and flucytosine, targeting the iron scavenging pyoverdine of the opportunistic human pathogen *Pseudomonas aeruginosa*. We first let bacteria evolve under different drug concentrations in replicated populations over 20 days, using human serum as an ex-vivo model. We then applied a combination of phenotypic and genotypic screens to show that resistance against flucytosine quickly arose and spread in all populations. Genetic analysis revealed that mutations in *upp*, a gene encoding an enzyme required for flucytosine activation, are responsible for resistance evolution. Conversely, resistance against gallium arose only sporadically and not in all populations. Resistance mechanisms were based on mutations in transcriptional regulators, which resulted in the upregulation of pyocyanin, a redox-active molecule promoting siderophore-independent iron acquisition. Our work highlights that mutants resistant against anti-virulence treatments can easily arise, however, selective spreading varies considerably between treatments. This demonstrates that there is no 'yes' or 'no' answer to the question of whether anti-virulence treatments are evolutionarily robust. Instead, evolutionary robustness is a relative measure on a sliding scale, with specific treatments occupying different positions on this scale.

Author Summary

There is an urgent need for alternative therapeutic options against multi-drug resistant bacteria. Antivirulence approaches, which target bacterial virulence factors thereby

reducing the ability to establish and maintain infections, are proposed as an evolutionary robust way to manage infections. Conversely to conventional antibiotics, there is great hope that "disarming" rather than killing pathogens will exert weaker selection for resistance. Here we compared the evolutionary robustness of two proposed antivirulence drugs, gallium and flucytosine, both targeting the pyoverdine mediated iron-uptake in the opportunistic human pathogen *Pseudomonas aeruginosa*. Pyoverdine is a strong iron chelator and an important virulence factor during acute infections, and has thus been identified as a promising target for antivirulence drugs. After exposing bacteria to treatments for 20 days in human blood serum, as an *ex-vivo* infection model, we found that resistant mutants against the two anti-virulence drugs readily arose. However, extensive selective spreading of resistant clones was only observed for flucytosine but not for gallium treatment. Our study demonstrates that anti-virulence treatments are not evolutionarily robust *per se*, but vary substantially in their propensity to select for resistance. We thus need standardized protocols to identify those treatments that are both efficient in controlling infections and withstand resistance evolution.

Introduction

There is currently much interest in therapeutic approaches that inhibit the expression or functioning of bacterial virulence factors [1–8]. Virulence factors are structures and molecules that allow bacteria to establish and maintain infections, and to cause host damage [9,10]. Examples of virulence factors include flagella and pili to adhere to the host tissue, secreted enzymes, tissue-damaging toxins and siderophores to scavenge iron from the host [11]. Approaches that target these traits are called anti-virulence treatments. There is great hope that disarming rather than directly killing pathogens is an efficient and evolutionarily robust way to manage infections [2,12–15]. In particular, it is thought that anti-virulence treatments exert weaker selection for resistance than conventional antibiotics because pathogens are not killed directly. However, empirical evidence for the evolutionary robustness of anti-virulence treatments is controversial with positive and negative reports

64 currently balancing each other out [16–21]. Part of the controversy is that it remains unclear what a resistance phenotype is with regard to anti-virulence therapies, and that there is a

66 shortage of studies examining resistance evolution under realistic conditions in parallel replicates both at the phenotypic and genetic level.

Here, we tackle this issue by probing the evolutionary robustness of two proposed

anti-virulence treatments, gallium and flucytosine, using them as experimental test cases to demonstrate how resistance evolution can be examined and quantified. The two anti-

virulence drugs target both the iron-scavenging pyoverdine of the opportunistic human pathogen *Pseudomonas aeruginosa* [19,22,23], which is an important virulence factor for this

species during acute infections [19,24–30]. In the context of infections, the host actively withholds iron from pathogens by binding it to strong iron-chelating proteins, such as

transferrin or lactoferrin [31]. In order to scavenge iron from these sources, opportunistic pathogenic bacteria secrete siderophores with high affinity for this essential nutrient. Given

the importance of iron for bacterial growth, it has been proposed that drugs interfering with iron uptake could be effective therapeutics to control infection [32]. Gallium and flucytosine

both fulfill this role, albeit through different modes of action. Gallium is an iron-mimic and binds irreversibly to pyoverdine outside the cell, thereby rendering the secreted molecules

useless for iron uptake [19,22,30]. While gallium is an anti-virulence drug at low dosages, it can function as an antibiotic at high concentrations. Then, gallium penetrates into bacterial

cells, where it is thought to replace iron in enzymes such as catalases and cytochromes, leading to cell death [33,34]. Flucytosine, meanwhile, is an anti-fungal drug that was

repurposed for the use against bacterial pathogens [23]. This drug enters the pathogen, where it is enzymatically activated to a fluorinated ribonucleotide. This active form is then

incorporated in the RNA and inhibits, via a yet unknown mechanism, the *pvdS* iron starvation sigma factor controlling pyoverdine synthesis [23,35].

Different lines of reasoning have been advocated for why these two drugs should be

92   evolutionarily robust. Flucytosine, while stalling pyoverdine production, does not seem to

compromise the growth of *P. aeruginosa* in in-vitro cultures [23]. If growth of the pathogen is

94   not affected, evolutionary theory predicts that resistant mutants, resuming virulence factor

production, should not experience a selective advantage and thus should not spread [15].

96   This reasoning for evolutionary robustness should not apply to gallium, which has been

shown to compromise bacterial growth due to the induced iron starvation [19,22,36,37].

98   Conversely, gallium and also flucytosine are supposed to be evolutionarily robust because

they suppress either the functioning (gallium) or the synthesis (flucytosine) of a secreted

100   virulence factor, which can be shared as a public good between pathogen individuals (iron-

loaded pyoverdine can be taken up by all bacteria with a matching receptor; [38,39]).

102   Resistant mutants should not spread in this case because they bear the cost of resumed

virulence factor production, whereas the benefit of increased virulence factor availability is

104   shared with everyone else in the population, including the drug-susceptible individuals

[12,14,16,20]. In addition, gallium is supposed to be evolutionarily robust because it targets a

106   secreted virulence factor outside the cell. The reasoning here is that common resistance

mechanisms, such as the prevention of drug influx, increased drug efflux, and intra-cellular

108   drug degradation cannot operate outside the cell. Thus, an extra-cellular mode of drug action

is supposed to reduce the number of mutational targets that can confer resistance [19].

To probe whether these principles apply to gallium and flucytosine, we exposed

112   replicated populations of *P. aeruginosa* to two different concentrations of these two drugs in

human blood serum. Together with a drug-free control treatment, we let the treated

114   populations evolve for 20 consecutive days in eight-fold replication, by transferring a fraction

of the evolving cultures to fresh human serum on a daily basis. Human serum has recently

116   been established as an *ex-vivo* model to mimic realistic host conditions, as for instance

experienced during bloodstream infections [40]. Following experimental evolution, we

118   screened evolved populations and randomly isolated clones for the evolution of resistance

phenotypes. We focused on three potential resistance phenotypes, which include the (partial

120   or full) restoration of growth, the (partial or full) restoration of virulence factor production, or

the evolution of a bypassing mechanism for iron uptake [15,19]. Finally, we sequenced the

122   whole genome of evolved clones to uncover the genetic basis of potential resistance

mechanisms.

Results

Gallium and flucytosine affect both growth and pyoverdine production in

human serum.

128   Since human serum is a novel *ex-vivo* model in the context of anti-virulence

experiments, we first assessed whether (a) pyoverdine production is beneficial in this media,

130   and (b) how gallium and flucytosine treatment affects the growth and pyoverdine production

of *P. aeruginosa*. To test (a), we compared the growth of our wildtype strain PAO1 to its

132   siderophore-negative mutants PAO1 Δ*pvdD* (deficient for pyoverdine production) and PAO1

Δ*pvdD* Δ*pchEF* (deficient for the production of pyoverdine and the secondary siderophore

134   pyochelin) in human serum across a range of iron limitations (S1 Fig). Across all conditions,

we observed that growth was significantly reduced in the siderophore-deficient mutants

136   compared to the wildtype (ANOVA for PAO1 Δ*pvdD*: t_{49}= -8.13, $p < 0.0001$, for PAO1 Δ*pvdD*

Δ*pchEF*: t_{49}= -7.92, $p < 0.0001$), confirming that human serum is an iron-limited media, and

138   that pyoverdine is important for growth.

140   To test (b), we subjected PAO1 cultures to a range of drug concentrations under

standardized conditions. The resulting dose-response curves reveal that both drugs

142   significantly reduce bacterial growth (Fig 1A and 1B). For gallium, growth reduction

manifested only at relatively high concentrations, whereas already the lowest concentration

144 of flucytosine caused a substantial growth reduction, which then remained fairly constant

across the concentration gradient. This experiment further confirmed that both drugs affect

146 the production of pyoverdine (Fig 1C and 1D), consistent with previous findings in standard

laboratory media. For flucytosine, the lowest concentration tested completely stalled

148 pyoverdine production. Gallium treatment affected pyoverdine production in a more complex

way, but consistent with previous findings [19]: at intermediate gallium concentrations,

150 pyoverdine is up-regulated to compensate for the gallium-induced pyoverdine inhibition, and

down-regulated at higher concentrations, when pyoverdine-mediated signaling becomes

152 impaired [22]. Our results thus reveal that both drugs affect pyoverdine production in the

predicted way in human serum, and thus act as anti-virulence drugs.

Do bacteria evolve population-level resistance to antivirulence treatments?

156 We carried out an experimental evolution with eight independent lines per treatment

for a period of 20 days, with daily transfers of a sub-fraction of the evolved cultures to fresh

158 human serum. Our experimental design comprised five treatment regimes: a no-drug control

treatment, a low and a high concentration treatment for both gallium and flucytosine. The

160 concentrations used (50 µM and 280 µM for gallium and 10 µg/ml and 140 µg/ml for

flucytosine) were inferred from the dose-response curves (Fig 1). Following experimental

162 evolution, we measured whether evolved populations showed a resistance phenotype either

through recovery of growth and/or a restoration of the pyoverdine production profile under

164 the treatment regimes they evolved in. For growth (Fig 2A), we found that the evolved lines

all grew significantly better than the treated ancestral wildtype (Welch's two sample t-tests,

166 for gallium low: $t_{11.9}$ = -4.96, p = 0.0003; for gallium high: $t_{13.3}$ = -6.48, p < 0.0001; for

flucytosine low: $t_{12.2}$ = -5.09, p = 0.0002; for flucytosine high $t_{7.5}$ = -11.79, p < 0.0001).

Regarding pyoverdine profiles, we observed no significant restoration of pyoverdine

170 production for the lines evolved under low gallium concentration (comparison with the treated

ancestor, Welch's two sample t-test: $t_{8.8} = 0.94$, $p = 0.3719$) (Fig 2B). Conversely, lines evolved under the other three drug regimes all showed a significant increase in pyoverdine production relative to the treated ancestor (Fig 2B) (Welch's two sample t-test for gallium high: $t_{13.1} = -3.69$, $p = 0.0026$; for flucytosine low: $t_{7.2} = -7.64$, $p = 0.0001$; for flucytosine high: $t_{9.6} = -54.65$, $p < 0.0001$). While the increase was moderate for the gallium high treatment, there was full restoration of pyoverdine production in both flucytosine treatments (no significant difference relative to the ancestral untreated wildtype, ANOVA for flucytosine low: $t_{88} = -1.31$, $p = 0.1944$; for flucytosine high: $t_{88} = 0.42$, $p = 0.6766$).

Next, we asked whether the observed restoration of growth and pyoverdine production indeed indicates resistance evolution, or whether these patterns could also be explained by media adaptation. For that purpose, we analyzed the growth and pyoverdine production of lines evolved in the absence of drugs. We found that these control lines also showed a significant increase in pyoverdine production relative to the untreated ancestor (Welch's two sample t-test, $t_{9.1} = -4.03$, $p = 0.0047$, Fig 2B). Moreover, some of the untreated evolved lines also showed significantly improved growth, although the overall growth increase across lines was not significant ($t_{9.1} = -1.61$, $p = 0.1424$, Fig 2A). These analyses demonstrate the difficulty of identifying resistance profiles of non-killing anti-virulence treatments, as it is hard to disentangle resistance evolution from media adaptation at the level of the entire population.

Screening for resistance profiles in evolved single clones

A more detailed way of screening for resistance profiles is to pick individual clones and measure whether their drug dose-response changed compared to the ancestor. In a first step, we thus isolated five random clones from each evolved line (i.e. 40 per treatment), and individually analyzed their growth and pyoverdine production. These analyses revealed high variation among clones in both growth and pyoverdine production in human serum (S2 Fig),

198   suggesting that most evolved populations were heterogeneous, consisting of multiple

different genotypes. For in-depth analysis, we then randomly picked 16 single clones (four

200   per drug treatment) and tested whether these evolved clones differ in their drug dose

response curve compared to the ancestral wildtype.

We observed that three out of eight clones subjected to gallium (Fig 3A-3D) and all

204   eight clones subjected to flucytosine showed a significantly altered dose response (Fig 3E-

3H). Clones GL_2 and GL_3, evolved under low gallium, showed a significant increase in

206   pyoverdine production under intermediate gallium concentrations (between 8 and 128 µM),

which goes along with an improved growth performance for GL_2, but not GL_3. In contrast,

208   clone GH_1, evolved under high gallium concentration, did not show an altered pyoverdine

production response, but grew significantly better when exposed to gallium (Fig 3A-3D). For

210   the eight clones evolved under the flucytosine regime, changes in the dose-response curves

were both striking and uniform: growth and pyoverdine production were both no longer

212   affected by the drug (Fig 3E-3H). Since these dose-response curve analyses directly include

a control for media adaptation (i.e. the no-drug treatment), our results indicate that all eight

214   clones evolved complete resistance to flucytosine. For gallium, on the other hand, our data

suggest that three out of the eight clones exhibited a phenotype that is compatible with at

216   least partial resistance.

218   **Linking phenotypes to genotypes**

To test whether there is a genetic basis for the putative resistance profiles described

220   in Fig 3, we sequenced the whole genome of all 16 clones. Following read mapping to the

PAO1 genome, we called sequence variants, both SNPs and INDELs, which have emerged

222   during evolution under anti-virulence drug exposure (Table 1). For the clones evolved with

flucytosine, we observed that all of them had acquired mutations in the coding sequence of

224   *upp.* There were four different types of mutations, including two different non-synonymous

SNPs, a 15-bp deletion and a 1-bp insertion (S1 Table). The *upp* gene encodes for a uracil

226 phosphoribosyl-transferase, which is one of the enzyme required for the activation of

flucytosine in the cell [41,42].

For the clones evolved under gallium treatment, the mutational pattern was more

230 heterogeneous (Table 1). While no mutations were detected for several clones (GH_2,

GH_3, and GH_4), the three clones with significantly altered dose responses had indeed

232 mutations that could be relevant in this context: clone GH_1 featured a 3-nt deletion in the

coding sequence of *mvaU*, whereas the clones GL_2 and GL_3 were mutated in the *vfr*

234 gene. Both *mvaU* and *vfr* encode transcriptional regulators involved in the regulation of

virulence factor expression, including proteases, pyocyanin and pyoverdine.

In addition, several clones from the gallium and flucytosine treatments had mutations

238 in either *dipA* (dispersion-induced phosphodiesterase A; GL_1, GL_4, FH_4) or *morA*

(motility regulator; GL_3, FH_2). The repeated yet unspecific appearance of these mutations

240 could suggest that they represent general adaptations to human serum. Taken together, our

sequencing analysis identify three potential targets of selection that could be associated with

242 resistance evolution: the gene *upp* for flucytosine, and the genes encoding for the global

transcriptional virulence regulators *vfr* and *mvaU* for gallium. In the next section, we will

244 investigate the possible underlying mechanisms of how mutations in these genes could

confer resistance in more detail.

246 **Table 1. List of mutations in evolved single clones**.

Treatment	Clone	Gene[1]	Description	Mutation	Type	Position[2]
Gallium Low	GL_1	*dipA*	dispersion-induced phosphodiesterase A	CA→C	INDEL	5642855-5642856
	GL_2	*vfr*	transcriptional regulator	C→T	SNP	706108
	GL_3	*vfr*	transcriptional regulator	C→T	SNP	706108
		PA3801	conserved hypothetical protein	A→G	SNP	4260811
		morA	motility regulator	G→T	SNP	5158144
	GL_4	*dipA*	dispersion-induced phosphodiesterase A	CA→C	INDEL	5642855-5642856
Gallium High	GH_1	*mvaU*	transcriptional regulator	GAGC→G	INDEL	3016276-3016279
	GH_2	none				
	GH_3	none				
	GH_4	none				
Flucytosine Low	FL_1	*upp*	uracil phosphoribosyl-transferase	T→C	SNP	5213244
		yfiR	tripartite signaling complex	C→T	SNP	1214975
		PA1369	hypothetical protein	C→T	SNP	1483680
		PA2770-PA2771	intergenic region	G→A	SNP	3129202
	FL_2	*upp*	uracil phosphoribosyl-transferase	GAGAAGATCTCCGGGA→G	INDEL	5213011-5213037
	FL_3	*upp*	uracil phosphoribosyl-transferase	A→C	SNP	5212855
		upp	uracil phosphoribosyl-transferase	A→G	SNP	5213146
	FL_4	*upp*	uracil phosphoribosyl-transferase	T→C	SNP	5213244
		groEL	protein chaperone	G→A	SNP	4916838
		PA2770-PA2771	intergenic region	G→A	SNP	3129202
Flucytosine High	FH_1	*upp*	uracil phosphoribosyl-transferase	G→GC	INDEL	5212852
		upp	uracil phosphoribosyl-transferase	A→C	SNP	5212855
		fliF	flagella M-ring outer membrane protein precursor	CA→C	INDEL	1194060-1194061
	FH_2	*upp*	uracil phosphoribosyl-transferase	A→C	SNP	5212855
		morA	motility regulator	G→A	SNP	5159713
	FH_3	*upp*	uracil phosphoribosyl-transferase	A→C	SNP	5212855
		fliF	flagella M-ring outer membrane protein precursor	TCGTCC→T	INDEL	1193365-1193370
	FH_4	*upp*	uracil phosphoribosyl-transferase	A→C	SNP	5212855
		dipA	dispersion-induced phosphodiesterase A	GA→G	INDEL	5643059-5643060

248   [1]Only mutations not found in the ancestor wildtype PAO1 are reported. Common mutations among all samples and the ancestor are listed in S3

Table.

250   [2] Position on PAO1 reference genome

Evolution of bypassing mechanisms for iron acquisition under gallium

252 **treatment**

Since gallium induces iron starvation via the quenching of pyoverdine molecules

254 outside the cell, it was proposed that bacteria could evolve bypassing mechanisms to

guarantee iron uptake in a siderophore-independent manner [19]. In human serum, where

256 iron is bound to transferrin or heme, we can envisage two such bypassing mechanisms.

First, bacteria could upregulate the production of pyocyanin, which can reduce ferric to

258 ferrous iron outside the cell, thereby facilitating direct pyoverdine-independent iron uptake

[18]. Second, bacteria could upregulate protease production to acquire iron from transferrin

260 or heme through protease-induced hydrolysis [28,43]. To test whether such bypassing

mechanisms evolved in our experiment, we quantified the production of pyocyanin and

262 proteases in the eight clones evolved under gallium treatment.

264 We indeed found that three clones displayed significantly increased pyocyanin

production per cell compared to the ancestral wildtype, and these were exactly the clones

266 with mutations in either *mvaU* or *vfr* (Fig 4A; ANOVA for GH_1: $t_{79} = 9.64$, $p < 0.0001$; for

GL_2: $t_{99} = 6.13$, $p < 0.0001$; for clone GL_3: $t_{99} = 14.8$, $p < 0.0001$). Since these clones are

268 mutated in global transcriptional regulator *mvaU* or *vfr*, which control directly (*mvaU*) or

indirectly (*vfr*) the expression of pyocyanin [44,45], a causal link between mutations in these

270 genes and the observed increased pyocyanin production is likely. Hence, the upregulation of

pyocyanin could represent a plausible by-passing mechanism, being under positive selection

272 in the presence of gallium.

274 In contrast, we found little evidence that increased protease production could be an

efficient by-passing mechanism in human serum. This is because six of the evolved clones

276 actually showed reduced and not increased production of extracellular proteases (Fig 4B).

Moreover, the two clones (GL_1 and GL_4) with significantly increased protease production

278   (ANOVA, for GL_1: $t_{10} = 13.22$, $p < 0.0001$; for GL_4: $t_{10} = 11.60$, $p < 0.0001$, Fig 4B) did not

show an altered drug dose-response curve, suggesting that increased protease activity in

280   these clones is not an evolutionary response to drug exposure.

282   **Inactivation of Upp is responsible for resistance to flucytosine**

As pointed out above, the enzyme Upp (uracil phosphoribosyl-transferase) is

284   essential for the activation of flucytosine within the cell. The natural function of Upp is to

convert uracil to UMP in the salvage pathway of pyrimidine (Fig 5A), with UMP representing

286   a nucleotide precursor. Interestingly, *P. aeruginosa* can also produce UMP through the

conversion of L-glutamine and L-aspartate [46] (Fig 5A). This multi-pronged metabolic

288   network led us to hypothesize that *upp* might not be essential for the pyrimidine metabolism,

and that the inactivation of this gene through mutations could confer resistance to

290   flucytosine.

292   To test this hypothesis, we compared the flucytosine dose-response curve of the

wildtype strain to an isogenic (transposon) mutant (MPAO1 Δ*upp*) with a non-functional *upp*

294   gene. Consistent with the patterns of the evolved clones (Fig 3G), we found that MPAO1

Δ*upp* was completely insensitive to flucytosine: neither growth (Fig 5B) nor pyoverdine

296   production (Fig 5C) were affected by the drug under the range of flucytosine concentration

tested (Welch's two sample *t*-test for difference between wildtype and mutant, for growth: $t_{4.3}$

298   = -5.56, $p = 0.0041$; for pyoverdine production: $t_{5.0} = -48.80$, $p < 0.0001$). These results

strongly suggest that the inactivation of *upp* is a simple and efficient way to evolve resistance

300   to flucytosine.

302   **Discussion**

New treatment approaches against the multi-drug resistant ESKAPE pathogens, to

304   which *P. aeruginosa* belongs, are desperately needed [8,47,48]. In this context, anti-

virulence treatments that disarm rather than directly kill the bacteria have attracted particular

306 interest, because such approaches have been proposed to be both effective in managing

infections and sustainable in the sense that resistance should not easily evolve [2,5,12–15].

308 Promising approaches include the inhibition of quorum sensing molecules regulating

virulence factor production [3], the suppression of toxins [49,50], or siderophore molecules

310 required for iron-scavenging within the host [19,22,23,37,51]. In our study, we probed the

evolutionary robustness argument by focusing on two anti-virulence drugs (gallium and

312 flucytosine) targeting siderophore production of *P. aeruginosa*, as experimental test cases.

Using a combination of replicated experimental evolution and phenotypic and genotypic

314 analysis, our results do not support the unanimous argument for the evolutionary robustness

of anti-virulence drugs currently prevailing in the literature. Instead, we found repeated and

316 consistent evolution of complete resistance against flucytosine, based on a mechanism that

prevents drug activation inside the cell. The situation is more promising for gallium, where

318 different variants of partially resistant mutants, based on iron uptake bypassing mechanisms,

only sporadically arose. Our work highlights that the question "are anti-virulence treatments

320 evolutionarily robust?" cannot be answered with "yes" or "no". Instead, we show that

evolutionary robustness is a relative measure with specific treatments lying on different

322 positions on the continuum between these two extremes. Our task is to assess the relative

position of each novel treatment on this continuum.

Our work on flucytosine identifies two issues of anti-virulence approaches, which have been

326 largely neglected so far. The first issue is that anti-virulence treatments can have

considerable negative off-target effects, which turned out to be detrimental for the

328 evolutionary robustness of flucytosine treatment (Fig 5A). Flucytosine undergoes several

enzymatic modifications within the cell, finally resulting in fluorinated ribonucleotides, which

330 compromise RNA and DNA functionality [52]. Thus, flucytosine not only inhibits pyoverdine

synthesis, but likely compromises cell viability more generally, setting the stage for selection

to favor mutants with alleviated fitness costs under drug exposure. Our results support that exactly this scenario occurred during experimental evolution in human serum through

mutations in *upp.* Fig 5A shows that the essential pyrimidine nucleotide precursor UMP can be synthesized through the salvage pathways reutilizing exogenous free bases and

nucleosides, or via a *de novo* biosynthesis pathway using L-glutamine or L-aspartate. This latter pathway requires higher amounts of energy (phosphate and reducing power), and this

is why organisms preferentially use the salvage pathway. However, the salvage pathway generates the harmful fluoro-UMP under flucytosine treatment, which provides a selective

advantage to mutants that inhibit the salvage pathway (through mutations in *upp*) and switch to the *de novo* biosynthesis pathway. The second issue is that whether or not anti-virulence

drugs affect pathogen fitness can depend on the ecological context. While Imperi et al. [23] showed that flucytosine does not significantly reduce bacterial growth in trypticase soy broth

dialysate (TSBD), we demonstrate significant fitness effects of this drug in human serum. Although these variable outcomes might not be surprising from an ecological perspective,

they have significant consequences for the evolutionary robustness argument of anti-virulence drugs: it may vary from one infectious context to another.

     While flucytosine failed our hopes of evolutionary robustness, gallium is more promising

350   because we observed no pervasive pattern of resistance evolution. Nonetheless, this anti-virulence treatment is not beyond any doubts, as we characterized several clones that have

352   evolved putative bypassing mechanism through which gallium-mediated pyoverdine quenching can be circumvented. We found one evolved clone with a mutation in *mvaU*,

354   encoding a positive regulator of pyocyanin production. This mutant indeed overproduced the redox-active molecule pyocyanin, which is able to reduce iron extracellularly, possibly

356   promoting iron uptake in a pyoverdine-independent way [18]. We further isolated two independent clones with mutations in *vfr*, encoding a global virulence factor regulator [10].

358   These mutants showed generally increased pyocyanin expression (Fig 4A) and increased

pyoverdine production (Fig 3C) under drug exposure. The molecular interpretation of these

360 patterns is that mutations in Vfr are known to stimulate PQS (Pseudomonas Quinolon Signal)

synthesis, which is inhibited by a functional Vfr. PQS in turn is known to positively regulate

362 pyocyanin synthesis. At the same time, it was shown that secreted PQS serves as an iron

chelator, increasing iron limitation thereby promoting pyoverdine synthesis [45,53].

Important to note, however, is that the occurrence of resistant mutants alone does not

366 necessarily mean that they spread and fix in the population. So, we might ask why mutants

with *mvaU* or *vfr* were not more prevalent given their gallium-resistant phenotypes. One

368 plausible evolutionary answer is that pyocyanin itself is secreted and might serve as a public

good. Thus, the benefit of its action (reducing iron extracellularly) could be shared between

370 resistant and susceptible wildtype cells. This scenario would corroborate the evolutionary

argument that resistant mechanisms, which are based on public good traits, should not fix in

372 a population [19,54–56]. This is a promising argument, supporting the idea that gallium is a

relatively robust anti-virulence drug. Nonetheless, the down-side of our finding is that the

374 putative by-passing mechanism observed involves the up-regulation of a potent toxin, which

is itself considered a virulence factor [57]. While several studies have suggested that the

376 quenching of particular virulence factors can lead to a phenotypic up-regulation of other

regulatorily-linked virulence factors [30,58], we here show that the same can happen as a by-

378 product of resistance evolution (as hypothesized by [59]). Whether and to what extent this

pattern affects virulence in a host should receive particular attention in future studies.

By using a replicated experimental evolution design, combined with phenotypic and genetic

382 analyses, our study helps to resolve the current debate in the literature on whether or not

anti-virulence approaches are evolutionarily robust. The proponents of the evolutionary

384 robustness argument relied so far on evolutionary theory and proof-of-principle studies to

support their idea that putatively resistant mutants cannot invade population of susceptible

386   bacteria [14,16,19,20]. Their main argument is that resistant mutants, restoring virulence

factor production, cannot spread because they alone carry the cost of virulence factor

388   synthesis, whilst sharing the benefit with all cells in the population, including the susceptible

ones. This prediction indeed held in competition experiments with engineered mimics of

390   resistant mutants under well-mixed conditions, where virulence factors are easily shared

[16,20]. Our experiments with gallium partially support their argument as mutants with up-

392   regulated pyoverdine or pyocyanin production did neither seem to have fixed in populations

(S2 Fig), nor did they appear in all replicates (Table 1). Nonetheless, *vfr* mutants for instance

394   appeared independently in two replicates, which suggests that these mutants experienced a

certain selective advantage. This advantage could be explained by the fact that human

396   serum is a viscous medium limiting the diffusion of secreted compounds [60], such that *vfr*-

mutants could experience a local advantage by increasing pyoverdine and pyocyanin

398   production. The local advantage might allow them to increase to a certain frequency,

potentially leading to a stable co-existence with the drug-susceptible wildtype strain. This

400   phenomenon, known as negative-frequency dependent selection, has been described for

several microbial public goods [54,61,62].

Conversely, the proponents of the argument that anti-virulence therapies are not

404   evolutionarily robust rely on a number of experiments where mutants with resistant

phenotypes were isolated [17,18,63,64]. Our experiments with flucytosine also support their

406   line of argumentation, as we found pervasive resistance evolution against this treatment.

This matches the results by Maeda et al. [17], who showed that resistance to the quorum-

408   quenching compound C-30 (brominated furanone) evolves repeatedly via upregulation of a

drug efflux pump. The spread of these mutants can be explained by the fact that their

410   treatment did not solely interfere with quorum quenching but also compromised the ability of

cells to grow in the adenosine medium, which requires a functional quorum sensing system

412   [65]. Their study together with our work show that as soon as there are negative off-target

effects, affecting the fitness of individual cells, resistance mechanisms will arise and spread.

414 Taken together, our study contributes to the solution of the debate by demonstrating that the

details of the mode of action together with the ecological context determine whether and to

416 what extent an anti-virulence drug can be evolutionarily robust.

418 In conclusion, our work advances research on anti-virulence drugs on multiple fronts. First, it

shows that resistant phenotypes are difficult to define, as they can involve the restoration of

420 growth, the resumption of virulence factor production, and/or the activation of a bypassing

mechanism. Detailed phenotypic and genotypic analyses, as those proposed in our study,

422 are required to disentangle background adaptation from resistance evolution. Second, we

show that anti-virulence approaches are neither completely evolution-proof nor does the

424 notion "all roads lead to resistance" apply [66]. A detailed evolutionary analysis for each

individual drug is required to find out where it is situated on the continuum between the two

426 extremes. Third, we advocate the application of more rigorous evolutionary approaches to

test for resistance evolution. While there are rigorous standards when it comes to describing

428 the precise molecular mode of action of a novel antibacterial [21,67], there is much room for

improvement for standards regarding the quantification and characterization of resistance

430 evolution [68,69].

Material and Methods

Strains and culturing conditions

434   We used the commonly used and genetically well-characterized *P. aeruginosa* PAO1

436   wildtype strain for all experiments. For some assays, we further used control strains, which

are listed in S2 Table. Overnight cultures were grown in 8 ml Luria–Bertani (LB) medium in

438   50-ml Falcon tubes and incubated at 37°C, 200 rpm for 18 hours. For all experiments, we

washed overnight cultures with 0.8% NaCl saline solution and adjusted them to OD_{600} = 2.5.

440   Bacteria were further diluted to a final starting of OD_{600} = 2.5 x 10^{-3}.

Growth assays in human serum

442   To understand to what extent iron is a limiting factor for bacterial growth in human

444   serum (from Sigma-Aldrich, Switzerland), we grew PAO1 and the corresponding

siderophore-deficient mutants in human serum supplemented with either: (a) 50 mM HEPES

446   to buffer the medium at physiological pH; (b) 50 mM HEPES together with the strong iron

chelator human apo-transferrin (100 µg/ml) and its co-factor $NaHCO_3$ (20 mM) to further

448   increase iron limitation; or (c) 50 mM HEPES together with $FeCl_3$ 20µM to decrease iron

limitation (S1 Fig). For the main experiments, we decided to use medium variant (b),

450   representing the most strongly iron limited medium. All chemicals where purchased from

Sigma-Aldrich, Switzerland.

Growth and virulence factor inhibition curves

454   We used gallium ($GaNO_3$) and flucytosine (5-Fluorocytosine) as anti-virulence drugs

targeting pyoverdine production (purchased from Sigma-Aldrich). To assess the extent to

456   which these two compounds inhibit PAO1 growth and pyoverdine production, we subjected

bacterial cultures to a range of antibacterial concentrations. For each compound we tested 8

458   concentrations, which included 0, 8, 16, 32, 64, 128, 256, 512 µM and 0, 2.5, 5, 10, 20, 40,

80, 140 µg/ml for gallium and flucytosine, respectively. Overnight cultures of bacteria were

460   grown and diluted as described above and used to inoculate 200 µl of iron-limited human

serum on 96-well plates. We then incubated the plates at 37 °C in a Tecan Infinite M-200

462   plate reader (Tecan Group Ltd., Switzerland), and tracked growth by measuring OD at 600

nm and pyoverdine-associated fluorescence (excitation at 400 nm, emission at 460 nm)

464   every 15 minutes for 24 hours. Plates were shaken for 15 seconds (3 mm orbital

displacement) prior to each reading event.

Experimental evolution

468   We exposed wildtype cultures of PAO1 to experimental evolution for a period of 20

days under five different selective regimes in eight-fold replication each. We had a no-drug

470   control treatment in addition to a low and a high concentration drug treatment for both gallium

and flucytosine. The concentrations used were inferred from the dose-response curves (Fig

472   1), and were 50 µM and 280 µM for gallium and 10 µg/ml and 140 µg/ml for flucytosine.

Experimental evolution took place in 190 µl iron-limited human serum distributed on the inner

474   wells of a 96-well plate. The plate also contained wells with media but no bacteria as

negative media controls. To initiate experimental evolution, an overnight culture of PAO1 was

476   grown as described above, and individual wells were inoculated with 10 µl of diluted culture

to a final density of 10^6 cells per well (10^{-3} final dilution). The plate was incubated statically at

478   37°C in the plate reader for 23.5 hours, and OD_{600} was measured every 15 minutes prior to a

shaking event (15 seconds at amplitude 3). After this growth period, cultures were

480   appropriately diluted in 0.8% NaCl and transferred to a new plate containing fresh media.

During the first 10 days of experimental evolution, the culture dilution factor was 10^{-3}.

482   Because there was a general increase in culture growth over this time period, we increased

culture dilution factor to $2*10^{-3}$ from day 11 to 20. Finally, to account for the fact that the

484   positive controls (bacteria in human serum without treatments) grew better than the anti-

bacterial treated cultures, we used dilution factors of $2*10^{-3}$ (from day 1 to 10) and $4*10^{-3}$

486   (from 11-20) for this treatment. Following each transfer, we preserved cultures by adding 100

        µl of a 50% glycerol-LB solution, and storing the mixture at -80°C.

488   **Population level resistance profiles**

                To test whether evolved populations had adapted to the presence of the

490   antibacterials by restoring growth and/or pyoverdine production, we exposed all evolved

        lineages to the same drug concentration they experienced during experimental evolution,

492   under standardized conditions in 5-fold replication each. To compare the performance of the

        evolved lineages relative to the ancestor wildtype, we simultaneously grew the wildtype

494   under drug and no-drug treatments. Plates were incubated at 37° C, shaken at 160 rpm for

        24 hours. Subsequently, we measured OD_{600} and pyoverdine-associated fluorescence using

496   the plate reader, and scaled all values relative to the untreated wildtype.

498   **Single clone isolation and inhibition curves**

                To isolate random clones from evolved lineages, we streaked out an aliquot of the

500   final day freezer stocks of the experimental evolution onto LB plates. After overnight

        incubation at 37°C, we randomly picked five colonies from each evolved lineage (i.e. 200

502   clones in total). To assess the variation in growth and pyoverdine production of these

        evolved clones, we measured their performance under the treatment conditions they

504   experienced during experimental evolution in 3-fold replication, in the same way we did for

        the population-level analysis. Moreover, we performed an in-depth analysis for 16 randomly

506   picked single clones (four per drug treatment). In particular, we repeated the drug-inhibition

        experiments using the same concentrations and the identical protocol as described above for

508   the ancestral wildtype.

510   **Sequencing analysis**

                We isolated the genomic DNA of the 16 evolved clones and performed whole

512   genome re-sequencing. We used the GenElute Bacterial Genomic DNA kit (Sigma Aldrich)

for DNA isolation. DNA concentrations were assayed using the Quantifluor dsDNA sample kit

514   (Promega) and samples were sent to the Functional Genomics Center Zurich for library

preparation (Nextera XT) and sequencing. Sequencing was performed on the Illumina HiSeq

516   4000 platform with single-end 125 base pair reads. Adapter sequences were clipped using

Trimmomatic v0.33 [70] and reads trimmed using Flexbar v2.5 [71]. We then aligned the

518   reads to the PAO1 reference genome using BWA v0.7.12 [72] and applied GATK v3.5

[73] indel   realignment,   duplicate   removal   and HaplotypeCaller   SNP/INDEL

520   discovery according to the GATK Best Practices recommendations, generating a variant call

format (VCF) file, which was then filtered as following: (i) variants with coverage < 20 reads

522   were discarded; (ii) variants with a Fisher Strand (FS) score > 30.0 were discarded to ensure

there is no strand bias in the data; (iii) variants with a QD value < 2.0 were discarded

524   (confidence value that there is a true variation at a given site); and (iv) clustered variants (≥ 3

variants in 35nt window) were discarded as they likely present sequencing or alignment

526   artifacts. This filtering process yielded a list of potential SNPs and small INDELs, which we

annotated using snpEff 4.1g [74] and then screened manually, compared to the sequenced

528   genome of our ancestor wildtype for relevant mutations in gene coding and intergenic

regions (see S3 Table).

Pyocyanin and protease assays

532   To establish links between the detected mutations and phenotypes, we screened the

eight sequenced clones evolved under gallium for changes in their proteases and pyocyanin

534   production profiles. To quantify pyocyanin production, we inoculated cells from overnight

cultures (diluted to a starting of $OD_{600} = 10^{-6}$) into 1 ml of LB. Following incubation for 24

536   hours at 37°C under shaken conditions (160 rpm), we measured pyocyanin in the cell-free

supernatant through absorbance at OD_{691} nm using the plate reader (protocol adapted from

538   [19]). We further quantified protease production of the same eight clones in human serum

(inoculated from overnight culture to a starting $OD_{600} = 2.5 \times 10^{-3}$). Following growth for 24

540   hours at 37°C under shaken conditions (160 rpm), we isolated the supernatants containing

proteases through centrifugation of the cultures at 3700 rpm for 15 minutes. We then set up

542   the proteolytic reaction mix (adapted from [75]) containing 0.1 ml of azocasein solution (30

mg/ml), 0.3 ml of 50 mM phosphate buffer (pH 7.5), and 0.1 ml culture supernatant. The

544   samples were incubated at 37°C for 2 hours, during which hydrolysis of the azocasein

releases the azo-dye, generating an absorbance signal at 366 nm, which is proportional to

546   the proteolytic activity. The proteolytic reaction was stopped adding 0.5 ml 20% (w/v)

trichloroacetic acid (TCA). We then centrifuged the mixture at 12000 rpm for 10 min, and

548   quantified the proteolytic activity via absorbance reading in the plate reader.

550   **Statistical analysis**

    Statistical analysis was carried out in RStudio (version 0.99.896, with R version

552   3.3.0). We analysed kinetic growth curves and pyoverdine production profiles using the *grofit*

package [76]. Because growth patterns often followed non-logistic trajectories, we fitted non-

554   parametric model (Splines) curves to estimate curve parameters, such as yield and integral

(area under the curve). Integral is particularly useful in the context of non-parametric growth

556   trajectories because it combines information on the lag-phase, growth rate and yield into one

growth parameter. To analyse dose-response curves we used the *drc* package [77] version

558   3.0-1. Model selection was conducted by comparison of different non-linear models and the

best fit was chosen using the following criteria: the log likelihood value, Akaike's information

560   criterion (AIC), the estimated residual standard error and the *p*-value from a lack-of-fit test. It

turned out that Weibull and Gompertz functions provided the best fit for gallium and flucytosin

562   dose-response curves, respectively. For all analyses, we scaled growth and pyoverdine

production relative to the untreated ancestral wildtype (=1). We used general linear mixed

564   effect models to compare whether parameters of growth or pyoverdine profiles differ in

evolved cultures treated with or without antibacterials. We further used Welch's two-sample *t*-

566   test to compare whether growth and pyoverdine levels of evolved lines differ from the

ancestor wildtype. Statistical analysis of the dose-response curve for the evolved single

568 clones was carried out by first fitting spline curves to the dose-response data, followed by the

estimation of integrals for these spline fits, and the comparison of the scaled fits relative to

570 the ancestor wildtype using a one-way ANOVA (Analysis of variance). Protease and

pyocyanin production of evolved single clones and the ancestor wildtype were corrected for

572 cell number (OD_{600}) and analysed using ANOVA.

574 **Data availability**

Whole-genome sequencing data have been deposited in the ArrayExpress database

576 at EMBL-EBI (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-6110.

578 **Acknowledgments**

We thank Adin Ross-Gillespie for advice, and the Functional Genomics Center Zurich for

580 technical support with the strain sequencing.

582 **References**

1. Escaich S. Antivirulence as a new antibacterial approach for chemotherapy. Curr Opin

584 Chem Biol. 2008;12: 400–408. doi:10.1016/j.cbpa.2008.06.022

2. Rasko DA, Sperandio V. Anti-virulence strategies to combat bacteria-mediated

586 disease. Nat Rev Drug Discov. 2010;9: 117–128. doi:10.1038/nrd3013

3. LaSarre B, Federle MJ. Exploiting Quorum Sensing To Confuse Bacterial Pathogens.

588 Microbiol Mol Biol Rev. 2013;77: 73–111. doi:10.1128/MMBR.00046-12

4. Maura D, Ballok AE, Rahme LG. Considerations and caveats in anti-virulence drug

590 development. Curr Opin Microbiol. 2016;33: 41–46. doi:10.1016/j.mib.2016.06.001

5. Vale PF, McNally L, Doeschl-Wilson A, King KC, Popat R, Domingo-Sananes MR, et

592 al. Beyond killing. Evol Med Public Heal. 2016; 148–157. doi:10.1093/emph/eow012

6. Johnson BK, Abramovitch RB. Small Molecules That Sabotage Bacterial Virulence.

594   Trends Pharmacol Sci. 2017;38: 339–362. doi:10.1016/j.tips.2017.01.004

7.   Rampioni G, Visca P, Leoni L, Imperi F. Drug repurposing for antivirulence therapy

596   against opportunistic bacterial pathogens. Emerg Top Life Sci. 2017;

doi:10.1042/ETLS20160018

598 8.   Dickey SW, Cheung GYC, Otto M. Different drugs for bad bugs: antivirulence

strategies in the age of antibiotic resistance. Nat Rev Drug Discov. 2017;16: 1–15.

600   doi:10.1038/nrd.2017.23

9.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. Common

602   virulence factors for bacterial pathogenicity in plants and animals. Science (80-).

1995;268: 1899–902. doi:10.1126/science.7604262

604 10.   Balasubramanian D, Schneper L, Kumari H, Mathee K. A dynamic and intricate

regulatory network determines Pseudomonas aeruginosa virulence. Nucleic Acids

606   Res. 2013;41: 1–20. doi:10.1093/nar/gks1039

11.   Wu HJ, Wang AHJ, Jennings MP. Discovery of virulence factors of pathogenic

608   bacteria. Curr Opin Chem Biol. 2008;12: 93–101. doi:10.1016/j.cbpa.2008.01.023

12.   André JB, Godelle B. Multicellular organization in bacteria as a target for drug therapy.

610   Ecol Lett. 2005;8: 800–810. doi:10.1111/j.1461-0248.2005.00783.x

13.   Baron C. Antivirulence drugs to target bacterial secretion systems. Curr Opin

612   Microbiol. 2010;13: 100–105. doi:10.1016/j.mib.2009.12.003

14.   Pepper JW. Drugs that target pathogen public goods are robust against evolved drug

614   resistance. Evol Appl. 2012;5: 757–761. doi:10.1111/j.1752-4571.2012.00254.x

15.   Allen RC, Popat R, Diggle SP, Brown SP. Targeting virulence: can we make evolution-

616   proof drugs? Nat Rev Microbiol. 2014;12: 300–308. doi:10.1038/nrmicro3232

16.   Mellbye B, Schuster M. The sociomicrobiology of antivirulence drug resistance: a

618   proof of concept. MBio. 2011;2: e00131-11. doi:10.1128/mBio.00131-11

17.   Maeda T, García-Contreras R, Pu M, Sheng L, Garcia LR, Tomás M, et al. Quorum

620   quenching quandary: resistance to antivirulence compounds. ISME J. 2012;6: 493–

501. doi:10.1038/ismej.2011.122

622   18.   García-Contreras R, Lira-Silva E, Jasso-Chávez R, Hernández-González IL, Maeda T,

Hashimoto T, et al. Isolation and characterization of gallium resistant Pseudomonas

624   aeruginosa mutants. Int J Med Microbiol. 2013;303: 574–582.

doi:10.1016/j.ijmm.2013.07.009

626   19.   Ross-Gillespie A, Weigert M, Brown SP, Kümmerli R. Gallium-mediated siderophore

quenching as an evolutionarily robust antibacterial treatment. Evol Med Public Heal.

628   2014;2014: 18–29. doi:10.1093/emph/eou003

20.   Gerdt JP, Blackwell HE. Competition Studies Confirm Two Major Barriers That Can

630   Preclude the Spread of Resistance to Quorum-Sensing Inhibitors in Bacteria. ACS

Chem Biol. 2014;9: 2291–2299. doi:10.1021/cb5004288

632   21.   Sully EK, Malachowa N, Elmore BO, Alexander SM, Femling JK, Gray BM, et al.

Selective Chemical Inhibition of agr Quorum Sensing in Staphylococcus aureus

634   Promotes Host Defense with Minimal Impact on Resistance. PLoS Pathog. 2014;10.

doi:10.1371/journal.ppat.1004174

636   22.   Kaneko Y, Thoendel M, Olakanmi O, Britigan BE, Singh PK. The transition metal

gallium disrupts Pseudomonas aeruginosa iron metabolism and has antimicrobial and

638   antibiofilm activity. J Clin Invest. 2007;117: 877–888. doi:10.1172/JCI30783

23.   Imperi F, Massai F, Facchini M, Frangipani E, Visaggio D, Leoni L, et al. Repurposing

640   the antimycotic drug flucytosine for suppression of Pseudomonas aeruginosa

pathogenicity. Proc Natl Acad Sci U S A. 2013;110: 7458–63.

642   doi:10.1073/pnas.1222706110

24.   Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. Pyoverdin is essential for

644   virulence of Pseudomonas aeruginosa. Infect. Immun. 1996. pp. 518–523.

25.   Takase H, Nitanai H, Hoshino K, Otani T. Impact of siderophore production on

646   Pseudomonas aeruginosa infections in immunosuppressed mice. Infect Immun.

2000;68: 1834–1839.

648 26. Harrison F, Browning LE, Vos M, Buckling A. Cooperation and virulence in acute

Pseudomonas aeruginosainfections. BMC Biol. 2006;4: 21. doi:10.1186/1741-7007-4-

650 21

27. Cornelis P, Dingemans J. Pseudomonas aeruginosa adapts its iron uptake strategies

652 in function of the type of infections. Front Cell Infect Microbiol. 2013;3: 75.

doi:10.3389/fcimb.2013.00075

654 28. Bonchi C, Imperi F, Minandri F, Visca P, Frangipani E. Repurposing of gallium-based

drugs for antibacterial therapy. BioFactors. 2014;40: 303–312. doi:10.1002/biof.1159

656 29. Granato ET, Harrison F, Kümmerli R, Ross-Gillespie A. Do Bacterial "Virulence

Factors" Always Increase Virulence? A Meta-Analysis of Pyoverdine Production in

658 Pseudomonas aeruginosa As a Test Case. Frontiers in Microbiology. 2016. p. 1952.

doi:https://doi.org/10.3389/fmicb.2016.01952

660 30. Weigert M, Ross-Gillespie A, Leinweber A, Pessi G, Brown SP, Kümmerli R.

Manipulating virulence factor availability can have complex consequences for

662 infections. Evol Appl. 2017;10: 91–101. doi:10.1111/eva.12431

31. Valenti P, Berlutti F, Conte MP, Longhi C, Seganti L. Lactoferrin Functions: Current

664 Status and Perspectives. J Clin Gastroenterol. 2004;38: S127-9.

32. Smith DJ, Lamont IL, Anderson GJ, Reid DW. Targeting iron uptake to control

666 Pseudomonas aeruginosa infections in cystic fibrosis. Eur Respir J. 2013;42: 1723–

36. doi:10.1183/09031936.00124012

668 33. Chitambar CR. Gallium-containing anticancer compounds. Future Med Chem. 2012;4:

1257–1272. doi:10.4155/fmc.12.69

670 34. Hijazi S, Visca P, Frangipani E. Gallium-Protoporphyrin IX Inhibits Pseudomonas

aeruginosa Growth by Targeting Cytochromes. Front Cell Infect Microbiol. 2017;7: 1–

672 15. doi:10.3389/fcimb.2017.00012

35. Visca P, Imperi F, Lamont IL. Pyoverdine siderophores: from biogenesis to

674 biosignificance. Trends Microbiol. 2007;15: 22–30. doi:10.1016/j.tim.2006.11.004

36. Banin E, Lozinski A, Brady KM, Berenshtein E, Butterfield PW, Moshe M, et al. The potential of desferrioxamine-gallium as an anti-*Pseudomonas* therapeutic agent. Proc Natl Acad Sci U S A. 2008;105: 16761–16766. doi:10.1073/pnas.0808608105

37. DeLeon K, Balldin F, Watters C, Hamood A, Griswold J, Sreedharan S, et al. Gallium Maltolate Treatment Eradicates Pseudomonas aeruginosa Infection in Thermally Injured Mice. Antimicrob Agents Chemother. 2009;53: 1331–1337. doi:10.1128/AAC.01330-08

38. Inglis RF, Scanlan P, Buckling A. Iron availability shapes the evolution of bacteriocin resistance in Pseudomonas aeruginosa. ISME J. 2016;10: 2060–2066. doi:10.1038/ismej.2016.15

39. Griffin AS, West SA, Buckling A. Cooperation and competition in pathogenic bacteria. Nature. 2004;430: 1024–1027. doi:10.1038/nature02744

40. Bonchi C, Frangipani E, Imperi F, Visca P. Pyoverdine and proteases affect the response of pseudomonas aeruginosa to gallium in human serum. Antimicrob Agents Chemother. 2015;59: 5641–5646. doi:10.1128/AAC.01097-15

41. Edlind TD, Katiyar SK. Mutational Analysis of Flucytosine Resistance in Candida glabrata. Antimicrob Agents Chemother. 2010;54: 4733–4738. doi:10.1128/AAC.00605-10

42. Beck DA, O'Donovan GA. Pathways of pyrimidine salvage in Pseudomonas and former Pseudomonas: Detection of recycling enzymes using high-performance liquid chromatography. Curr Microbiol. 2008;56: 162–167. doi:10.1007/s00284-007-9050-3

43. Doring G, Pfestorf M, Botzenhart K, Abdallah MA. Impact of Proteases on Iron Uptake of Pseudomonas aeruginosa Pyoverdin from Transferrin and Lactoferrin. 1988;56: 291–293.

44. Li C, Wally H, Miller SJ, Lu C. The multifaceted proteins MvaT and MvaU, members of the H-NS family, control arginine metabolism, pyocyanin synthesis, and prophage activation in Pseudomonas aeruginosa PAO1. J Bacteriol. 2009;191: 6211–6218.

702   doi:10.1128/JB.00888-09

45.   Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR, Lamont IL, et al. The

704   Pseudomonas aeruginosa 4-Quinolone Signal Molecules HHQ and PQS Play

Multifunctional Roles in Quorum Sensing and Iron Entrapment. Chem Biol. 2007;14:

706   87–96. doi:10.1016/j.chembiol.2006.11.014

46.   Isaac JH, Holloway BW. Control of pyrimidine biosynthesis in Pseudomonas

708   aeruginosa. J Bacteriol. 1968;96: 1732–41.

47.   Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens.

710   Expert Rev Anti Infect Ther. 2013;11: 297–308. doi:10.1586/eri.13.12

48.   Brown D. Antibiotic resistance breakers: can repurposed drugs fill the antibiotic

712   discovery void? Nat Rev Drug Discov. 2015;14: 821–832. doi:10.1038/nrd4675

49.   Lu C, Maurer CK, Kirsch B, Steinbach A, Hartmann RW. Overcoming the Unexpected

714   Functional Inversion of a PqsR Antagonist in Pseudomonas aeruginosa: An In Vivo

Potent Antivirulence Agent Targeting pqs Quorum Sensing. Angew Chemie Int Ed.

716   2014;53: 1109–1112. doi:10.1002/anie.201307547

50.   Henry BD, Neill DR, Becker KA, Gore S, Bricio-Moreno L, Ziobro R, et al. Engineered

718   liposomes sequester bacterial exotoxins and protect from severe invasive infections in

mice. Nat Biotechnol. 2015;33: 81–88. doi:10.1038/nbt.3037

720   51.   Kelson AB, Carnevali M, Truong-Le V. Gallium-based anti-infectives: targeting

microbial iron-uptake mechanisms. Curr Opin Pharmacol. 2013;13: 707–716.

722   doi:10.1016/j.coph.2013.07.001

52.   Harbers E, Chaudhuri NK, Heidelberger C. Studies on fluorinated pyrimidines. VIII.

724   Further biochemical and metabolic investigations. J Biol Chem. 1959;234: 1255–62.

53.   Lin J, Zhang W, Cheng J, Yang X, Zhu K, Wang Y, et al. A Pseudomonas T6SS

726   effector recruits PQS-containing outer membrane vesicles for iron acquisition. Nat

Commun. 2017;8: 14888. doi:10.1038/ncomms14888

728   54.   Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. Bacterial cheating drives the

population dynamics of cooperative antibiotic resistance plasmids. Mol Syst Biol.

730   2013;9: 683. doi:10.1038/msb.2013.39

55.   Medaney F, Dimitriu T, Ellis RJ, Raymond B. Live to cheat another day⬚: bacterial

732   dormancy facilitates the social exploitation of β -lactamases. ISME J. 2015;10: 778–

787. doi:10.1038/ismej.2015.154

734   56.   Kelsic ED, Zhao J, Vetsigian K, Kishony R. Counteraction of antibiotic production and

degradation stabilizes microbial communities. Nature. 2015;521: 516–519.

736   doi:10.1038/nature14485

57.   Gellatly SL, Hancock REW. Pseudomonas aeruginosa: New insights into

738   pathogenesis and host defenses. Pathog Dis. 2013;67: 159–173. doi:10.1111/2049-

632X.12033

740   58.   Beceiro A, Tomás M, Bou G. Antimicrobial resistance and virulence: A successful or

deleterious association in the bacterial world? Clin Microbiol Rev. 2013;26: 185–230.

742   doi:10.1128/CMR.00059-12

59.   Lau GW, Hassett DJ, Ran H, Kong F. The role of pyocyanin in Pseudomonas

744   aeruginosa infection. Trends Mol Med. 2004;10: 599–606.

doi:10.1016/j.molmed.2004.10.002

746   60.   Kümmerli R, Griffin AS, West S a, Buckling A, Harrison F. Viscous medium promotes

cooperation in the pathogenic bacterium Pseudomonas aeruginosa. Proc Biol Sci.

748   2009;276: 3531–3538. doi:10.1098/rspb.2009.0861

61.   Ross-Gillespie A, Gardner A, West SA, Griffin AS. Frequency dependence and

750   cooperation: theory and a test with bacteria. Am Nat. 2007;170: 331–342.

doi:10.1086/519860

752   62.   Raymond B, West SA, Griffin AS, Bonsall MB. The dynamics of cooperative bacterial

virulence in the field. Science (80-). 2012;337: 85–88. doi:10.1126/science.1218196

754   63.   Garcia-Contreras R, Maeda T, Wood TK. Resistance to Quorum-Quenching

Compounds. Appl Environ Microbiol. 2013;79: 6840–6846. doi:10.1128/AEM.02378-

756   13

64.   Kalia VC, Wood TK, Kumar P. Evolution of resistance to quorum sensing inhibitors.
758   Microb Ecol. 2014;68: 13–23. doi:10.1007/s00248-013-0316-y

65.   Dandekar AA, Chugani S, Greenberg EP. Bacterial Quorum Sensing and Metabolic
760   Incentives to Cooperate. Science. 2012;338: 264–266. doi:10.1126/science.1227289

66.   Breidenstein EBM, de la Fuente-Núñez C, Hancock REW. Pseudomonas aeruginosa:
762   all roads lead to resistance. Trends Microbiol. 2011;19: 419–426.
         doi:10.1016/j.tim.2011.04.005

764   67.   Ling LL, Schneider T, Peoples AJ, Spoering AL, Engels I, Conlon BP, et al. A new
         antibiotic kills pathogens without detectable resistance. Nature. 2015;517: 455–459.
766   doi:10.1038/nature14098

68.   Perron GG, Zasloff M, Bell G. Experimental evolution of resistance to an antimicrobial
768   peptide. Proc R Soc B Biol Sci. 2006;273: 251 LP-256. doi:10.1098/rspb.2005.3301

69.   Hochberg ME, Jansen G. Bacteria: Assessing resistance to new antibiotics. Nature.
770   2015;519: 158. doi:10.1038/519158e

70.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
772   data. Bioinformatics. 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170

71.   Dodt M, Roehr J, Ahmed R, Dieterich C. FLEXBAR—Flexible Barcode and Adapter
774   Processing for Next-Generation Sequencing Platforms. Biology (Basel). 2012;1: 895–
         905. doi:10.3390/biology1030895

776   72.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler
         transform. Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

778   73.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
         Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation
780   DNA sequencing data. Genome Res. 2010;20: 1297–1303.
         doi:10.1101/gr.107524.110

782   74.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly

784   (Austin). 2012;6: 80–92. doi:10.4161/fly.19695

75.   Chessa JP, Petrescu I, Bentahir M, Van Beeumen J, Gerday C. Purification, physico-

786   chemical characterization and sequence of a heat labile alkaline metalloprotease

isolated from a psychrophilic Pseudomonas species. Biochim Biophys Acta - Protein

788   Struct Mol Enzymol. 2000;1479: 265–274. doi:10.1016/S0167-4838(00)00018-2

76.   Kahm M, Hasenbrink G, Ludwig J. grofit□: Fitting Biological Growth Curves with R. J

790   Stat Softw. 2010;33. doi:10.18637/jss.v033.i07

77.   Ritz C, Baty F, Streibig JC, Gerhard D. Dose-Response Analysis Using R. 2015; 1–13.

792   doi:10.1371/journal.pone.0146021

 Figure captions

**Fig 1. Gallium and flucytosine affect both growth and pyoverdine production of *P.*
aeruginosa in human serum (HS)**. Both anti-virulence drugs reduce growth of bacterial
cultures in a dose-dependent manner (A, B), albeit following different patterns: gallium curbs
bacterial growth only at relatively high concentrations (A), whereas flucytosine already
reduces growth at low concentrations (B). Both drugs further affect pyoverdine production (C,
D). When increasing gallium exposure, bacteria first upregulate pyoverdine production at
intermediate drug concentrations, but then down-scale investment levels at high drug
concentrations (C). In contrast, flucytosine administration leads to an almost complete
abolishment of pyoverdine production even at the lowest drug concentration. All data are
scaled relative to the drug-free treatment. Error bars denote standard errors of the mean
across 6 (for flucytosine) and 18 (for gallium) replicates. Dose-response curves were fitted
using either Weibull (for gallium) or Gompertz (for flucytosine) functions (see material and
methods for details). Vertical lines indicate the drug concentrations used in the experimental
evolution.

**Fig 2. Population level growth and pyoverdine production after evolution in human
serum**. PAO1 cultures were exposed to either no treatment, low (50 µM) or high (280 µM)
gallium, low (10 µg/ml) or high (140 µg/ml) flucytosine concentrations during a 20-day
experimental evolution experiment in eight-fold replication. Following evolution, we assessed
growth and pyoverdine production of evolved populations (displayed on the x-axis) and
compared their performance relative to the untreated (gray solid line, set to 1) and treated
ancestral wildtype (black dashed line). (A) Compared to the treated ancestral wildtype,
growth of evolved populations significantly increased under all treatment regimes. Growth
also increased in some but not all of the non-treatment lines. (B) Pyoverdine production of
evolved populations significantly increased relative to the untreated ancestral wildtype under
all conditions, also in the no treatment lines. This indicates that increased pyoverdine

production might be a general response to growth in human serum, which makes it difficult to

822   disentangle resistance evolution from media adaptation. Error bars show the standard error

of the mean across 5 independent replicates.

Fig 3. Changes in dose-response curves for evolved single clones indicate resistance

826   **evolution**. 16 randomly picked clones, four per treatment, were exposed to a range of drug

concentrations to test whether their dose-response altered during evolution compared to the

828   ancestral wildtype (black circles and lines). (A and B) Growth dose-response curves under

gallium treatment show that two evolved clones (GL2 and GH1) are significantly less

830   inhibited than the ancestral wildtype. (C and D) Pyoverdine dose-response curves under

gallium treatment show that two evolved clones (GL2 and GL3) make significantly more

832   pyoverdine than the ancestral wildtype. (E and F) Growth dose-response curves under

flucytosine treatment show that all evolved clones grow significantly better than the ancestral

834   wildtype, and are in fact no longer affected by the drug. (G and H) Pyoverdine dose-response

curves under flucytosine treatment show that all evolved clones produce significantly more

836   pyoverdine than the ancestral wildtype, and are in fact no longer affected by the drug.

Growth and pyoverdine production were measured after 24 h. For each clone, values are

838   scaled relative to its performance in human serum without drugs (absolute values of

pyoverdine and growth in the absence of treatment are reported in S3 Fig). We used spline

840   functions to fit dose-response curves, and used the integral (area under the curve) to

quantify the overall dose response of each clone across the concentration gradient. Error

842   bars denote standard errors of the mean across 6 replicates. Asterisks represent significance

levels: $* = p < 0.05$; $*** = p < 0.0001$, based on linear model with $df = 45$).

Fig 4. Upregulation of pyocyanin or protease production as potential bypassing

846   **mechanisms for iron acquisition under gallium treatment**. The eight sequenced clones

evolved under gallium treatments (low: 50 µM, high: 280 µM) were screened for their change

848   in the secretion of pyocyanin (A) and proteases (B) relative to the ancestral wildtype. Standard protocols were used for the phenotypic screens in drug free media (see Material

850   and Methods for details). All values are corrected for cell number, and scaled relative to the ancestor wildtype (black line). We included the strains PAO1 Δ*rhlR* (deficient for pyocyanin

852   production) and PAO1 Δ*lasR* (deficient for protease production) as negative controls in the respective assays (dashed lines). Error bars denote standard errors of the mean across 3

854   (for proteases) and 8 (for pyocyanin) replicates. Asterisks represent significance levels: * = p < 0.05; *** = p < 0.0001, based on ANOVA.

Fig 5. Upp is a non-essential enzyme, and mutations in its gene result in flucytosine

resistance. (A) Flucytosine interferes with the pyrimidine metabolism in *P. aeruginosa.* The drug enters the cell through the transporter CodB (not shown), where it is first converted to

fluorouracil by the cytosine deaminase CodA, and then to fluoro-UMP by the uracil phosphoribosyl-transferase Upp. Fluoro-UMP is a modified nucleotide precursor, the action

of which results in RNA molecules with compromised functionality. Through an as yet unknown mechanism, fluoro-UMP also arrests pyoverdine synthesis in *P. aeruginosa.*

Importantly, the nucleotide-precursor UMP can also be produced through an alternative *de-novo* pathway from the amino acids L-glutamine and L-aspartate, making Upp a non-

essential enzyme in this bacterium. Experiments with the transposon mutant MPAO1 *Δupp* (deficient for Upp production) indeed demonstrate that the lack of Upp no longer affects

strain growth (B) and pyoverdine production (C). This demonstrates that the inactivation of *upp* is a simple and efficient way to evolve resistance to flucytosine. Experiments were

carried out in human serum across a range of flucytosine concentrations. Growth and pyoverdine production of MPAO1 *Δupp* (gray-dashed lines) and its corresponding wildtype

MPAO1 (black-solid lines) were measured after 24 hours for each treatment separately in 6-fold replication. All values are scaled relative to the drug-free treatment. For statistical

analysis, we compared the integrals of the dose-response curves between the mutant and

the wildtype strain.

Supporting information captions

878   **S1 Fig. Human Serum is an iron limited media.** To show that pyoverdine production is

beneficial in human serum, we compared growth of *P. aeruginosa* PAO1 wildtype and of

880   siderophore-deficient mutants PAO1 *ΔpvdD* and PAO1 *ΔpvdD ΔpchEF* in human serum with

different supplements, which results in three different iron regimes. When adding only 50 mM

882   HEPES to buffer the medium at physiological pH (middle panel) we observed that growth of

siderophores mutants was significantly reduced compared to the wildtype (ANOVA, for PAO1

884   *ΔpvdD*: t_{15}= -7.47, p < 0.0001; for PAO1 *ΔpvdD ΔpchEF*: t_{15}= -6.87, p < 0.0001), confirming

that human serum is an iron-limited media. To further increase iron limitation (left panel), we

886   added the strong iron chelator human apo-transferrin (100 µg/ml) and its co-factor $NaHCO_3$

(20 mM); conversely to increase iron-availability in human serum we added 20 µM of iron

888   chloride (right panel). We found that growth decreased when iron became more limiting

(ANOVA, for human serum + transferrin: t_{49}= -25.26, p < 0.0001), while growth increased

890   when extra iron was added (for human serum + iron chloride t_{49}= 14.60, p < 0.0001).

However, the wildtype PAO1 grew significantly better compared to the siderophores mutants,

892   in all three iron regimes (ANOVA, for PAO1 *ΔpvdD*: t_{49}= -8.12, p < 0.0001; for PAO1 *ΔpvdD*

ΔpchEF: t_{49} = -7.91, p < 0.0001), confirming that siderophores are important for iron

894   scavenging in human serum. All growth data are scaled relative to the wildtype growth in

human serum supplemented with HEPES 50 mM (middle panel). Error bars denote standard

896   errors of the mean across 6 replicates.

898   **S2 Fig. Growth and pyoverdine production profiles of evolved single clones under**

treatment regimes. We were interested in examining the variation in growth and pyoverdine

900   production profiles among evolved clones. For that purpose, we streaked out all evolved

cultures on LB agar, and picked 5 random clones per evolved population. Overall, we

902   isolated 160 clones, 40 clones per treatment. Single clones were tested under the treatment

regime experienced during experimental evolution, and growth and pyoverdine production

904   were measured after 24 hours. Panels show growth and pyoverdine production of evolved

clones relative to the untreated wildtype (grey line) under gallium (A, C) and flucytosine

906   treatment (B, D). We found heterogeneous growth and pyoverdine patterns under all

treatment regimes, suggesting diversification in evolving populations. Data points show

908   means across three independent replicates. Dashed lines depict the mean growth or the

mean pyoverdine production of the ancestral strain under treatment. Labeled single clones

910   marked in red refer to the clones used for in-depth analysis and sequencing.

912   **S3 Fig. Growth and pyoverdine production of evolved single clones in human serum**

with no treatment. We quantified the growth (A) and pyoverdine production (B) of the 16

914   evolved single clones we used for in-depth analysis and whole-genome sequencing.

Specifically, we grew the clones for 24 hours in human serum and compared their

916   performance to the ancestor wildtype in absence of the treatments. This allowed us to test for

media adaptation. (A) We observed that some of the clones from both treatments showed

918   significantly improved growth compared to the ancestral wildtype. This was the case for

clones GL1 (t_{43} = 2.04, p = 0.0475), GL_4 (t_{43} = 2.04, p = 0.0475), GH_1 (t_{43} = 3.13, p =

920   0.0125), FL_1 (t_{43} = 5.88, p < 0.0001) and all clones evolved under high flucytosine (FH_1: t_{43}

= 3.09, p = 0.0054; FH_2:, t_{43} = 3.03, p = 0.0054; FH_3: t_{43} = 4.38, p = 0.0002; FH_4, t_{43} =

922   2.82, p = 0.0071). Interestingly, two of the clones evolved under gallium low treatment

showed reduced growth compared to PAO1 (GL_2: t_{43} = -3.35, p = 0.0033; GL_3: t_{43} = -3.36,

924   p = 0.0033). (B) Regarding per capita pyoverdine production (pyoverdine fluorescence

divided by growth), we found that two clones showed increased levels of pyoverdine

926   production under gallium treatment (GL_1: t_{43} = 2.79, p = 0.0200; GL_4: t_{43} = 2.69, p =

0.0200), and three clones of the flucytosine high treatment did so too (FH_1: t_{43} = 2.43, p =

928   0.0356; FH_2: t_{43} = 2.29, p = 0.0356; for FH_4: t_{43} = 2.35, p = 0.0356). In contrast, there were

also a number clones with significantly reduced pyoverdine production compared to PAO1

930   (GH_1: t_{43}= -3.14, p = 0.0045; GH_3: t_{43}= -3.10, p = 0.0045; GH_4: t_{43}= -3.46, p = 0.0045;

FL_4: t_{43}= -2.02, p=0.0488). All data are scaled relative to the ancestor wildtype. Error bars denote standard error of the mean across 6 replicates. Asterisks represent significance codes (* = p<0.05; ** = p<0.001, *** = p<0.0001 based on ANOVA, corrected for multiple pairwise comparison with the false discovery rate method).

S4 Fig. Alteration of the fluorescent properties of pyoverdine when bound to gallium. The fluorescent signal of pyoverdine becomes inflated when gallium binds to it [19]. To take this signal bias into account, we quantified the bias in fluorescence signal as a function of gallium concentration. Specifically, we supplemented iron-limited human serum with 200 μM of purified pyoverdine across a range of gallium concentrations (8 to 512 μM, as used for the main experiments). We found that the signal bias can be explained by a 3 parameters-logistic function. We used this function to correct for signal bias in all our analyses.

S1 Table. Effects of mutation on the Upp protein sequence of evolved single clones.

S2 Table. List of control strains used in this study.

S3 Table. Differences to the PAO1 reference genome shared among all sequenced evolved clones and the ancestor wildtype.

Inhibition curves in human serum

A

cytidine

cytosine

CMP

CodA — uridine

CDP → CTP

uracil → UMP

Upp

PyrG

UDP → UTP

PyrF

Orotidine 5-P ← Orotate ← - - - - - L-glutamine/ L-aspartate

PyrE

DNA
RNA

→ Salvage pathway (blue)

→ De novo pathway (green)

→ Enzymes not characterized (black)

B

p=0.004

Relative growth (y-axis: 0.0, 0.5, 1.0, 1.5, 2.0)

Flucytosine (μg/ml) (x-axis: 10, 100)

C

p<0.0001

Relative pyoverdine (y-axis: 0.0, 0.5, 1.0, 1.5, 2.0)

Flucytosine (μg/ml) (x-axis: 10, 100)