Learning naturalistic temporal structure in the posterior medial network

Mariam Aly[1,2]*, Janice Chen[1,3], Nicholas B. Turk-Browne[1,4,5], Uri Hasson[1,4]

[1] Princeton Neuroscience Institute; Princeton University; Princeton, NJ, 08544; USA

[2] Department of Psychology; Columbia University; New York, NY, 10027; USA

[3] Department of Psychological & Brain Sciences; Johns Hopkins University; Baltimore, MD, 21218; USA

[4] Department of Psychology; Princeton University; Princeton, NJ, 08544; USA

[5] Department of Psychology; Yale University; New Haven, CT, 06520; USA

*Please address correspondence to:
Mariam Aly
Department of Psychology
406 Schermerhorn Hall
1190 Amsterdam Ave
Columbia University
New York, NY 10027
email: ma3631@columbia.edu
phone: 212-854-5318

Impact statement: Aly et al., show that the posterior medial network is engaged by temporally structured events, whether the temporal structure comes from world knowledge or is learned anew in recent experience.

Abstract

The posterior medial network is at the apex of a temporal integration hierarchy in the brain, integrating information over many seconds of viewing intact, but not scrambled, movies. This has been interpreted as an effect of temporal structure. Such structure in movies depends on pre-existing event schemas, but temporal structure can also arise de novo from learning. Here we examined the relative role of schema-consistent temporal structure and arbitrary but learned temporal structure on the human posterior medial network. We tested whether, with repeated viewing, the network becomes engaged by scrambled movies with temporal structure. Replicating prior studies, posterior medial regions were immediately locked to stimulus structure upon exposure to intact but not scrambled movies. However, for temporally structured scrambled movies, functional coupling within the network increased across stimulus repetitions, rising to the level of intact movies. Thus, temporal structure is a key determinant of network dynamics and function.

Keywords: temporal regularities, statistical learning, network dynamics, precuneus, hippocampus

Introduction

The brain's activity is influenced by events across multiple timescales, from seconds to minutes ago (Chaudhuri et al., 2015; Hasson et al., 2015; Murray et al., 2014). Prior studies that have investigated the integration of information over time have often relied on naturalistic stimuli, such as movies or stories (e.g., Hasson et al., 2004, 2008; Honey et al., 2012a,b; Lerner et al., 2011). These stimuli are presented either intact or temporally scrambled (divided into randomly re-ordered segments). Scrambling can be done at different temporal granularities (e.g., at the word, sentence, or paragraph levels), and which level of scrambling disrupts activity in a given brain region indicates its timescale of processing. For example, sensory areas track instantaneous physical parameters (e.g., speech volume) and are unaffected by even the finest scrambling, indicating that they have short processing timescales. Conversely, higher-order areas in the posterior medial network (Ranganath and Ritchey, 2012), which overlap with the default mode network (Raichle, 2015), are sensitive to temporal context extending tens of seconds in the past (e.g., a previous sentence can affect the response to an incoming word). They can be disrupted by even coarse scrambling, indicating that they have long processing timescales (Hasson et al., 2015).

Why do higher-order areas respond in a reliable manner to intact, but not scrambled, movies and stories? Both intact and scrambled movies and stories contain events that unfold over time, but only intact stimuli are predictable over long timescales: they share similarities with learned representations of real-life events we repeatedly encounter (i.e., schemas), which have temporal structure on the order of seconds or minutes. The structure conferred by schemas can influence how new information is interpreted, as well as the ability to remember that information and predict upcoming events (Gilboa & Marlatte, 2017; van Kesteren et al., 2012). For example, if food is served to someone at a restaurant, we know that she is having a meal, that she ordered after viewing the menu, and that she will ask for the bill upon finishing. Scrambling the order of events will interfere with our ability to integrate information over time and predict what will happen next, because existing schema

no longer apply (e.g., we might see someone order food *after* watching her eat it). Nevertheless, we are capable of learning new event sequences that may initially violate expectations (e.g., paying *before* we eat at a canteen) by extracting temporal regularities across repetitions, a process known as statistical learning (Aslin & Newport, 2012; Schapiro et al., 2017). Here, we explore whether brain regions that accumulate information over long timescales for schema-consistent movies can become engaged by novel event sequences over repeated viewings.

We scanned individuals with fMRI while they watched three 90-s movie clips, six times each, from *The Grand Budapest Hotel* (Figure 1). One clip ("Intact") was viewed in its original format: a temporally structured clip consistent with pre-existing schemas of how real-life events unfold. Another clip ("Scrambled-Fixed") comprised short segments that were randomly re-ordered, but their order was identical across all repetitions, creating incoherent but stable temporal structure. A final clip ("Scrambled-Random") also comprised scrambled segments, but the segments were randomly re-ordered for each repetition, creating incoherent and unstable temporal structure. In contrast to studies in which scrambled movies were presented in the same scrambled order twice (e.g., Hasson et al., 2008; Honey et al., 2012a), the six repetitions used here allowed us to search for statistical learning of temporal structure over repeated presentations.

To examine brain activity dynamics for these intact and scrambled movie clips, we conducted several analyses that provide complementary insights. We first examined the consistency of temporal dynamics in the brain across subjects using *inter-subject correlation*. Inter-subject correlation reveals neural responses that are locked to the structure of the external stimulus and are anatomically shared across individuals. This analysis consists of calculating the correlation between one brain region's activity timecourse in an individual and that same brain region's activity timecourse in other individuals who viewed the same external stimulus. A recent extension of inter-subject correlation, *inter-subject functional correlation*, enables identification of stimulus-locked responses that are functionally shared across individuals at the level of brain networks (Simony et al., 2016). This analysis

consists of calculating the correlation between one brain region's activity timecourse in an individual and a *different* brain region's activity timecourse in other individuals.

To search for neural responses that are locked to stimulus structure, but are idiosyncratic (i.e., not shared across individuals), we also calculated *intra-subject correlation* across repeated viewings of a stimulus (e.g., Hasson et al., 2008; Honey et al., 2012a). This analysis consists of calculating the correlation between a single brain region's activity timecourse for different repetitions of a given stimulus, within an individual. Finally, we examined *intra-subject functional correlation* — the correlations between the activity timecourses of different brain regions within an individual (often called functional connectivity). Intra-subject functional correlation analyses can identify neural responses which are shared across different nodes of a network irrespective of whether they are shared across individuals or stable across repeated viewings for a given individual.

Inter- and intra-subject analyses, in tandem, can reveal important information about how a stimulus is processed in the brain. High intra-subject correlations, alongside low inter-subject correlations, would indicate that each individual is processing the stimulus in a unique way. Low intra-subject correlations, alongside low inter-subject correlations, may indicate that each individual is processing the stimulus in a unique way that is unstable (i.e., changes) over stimulus repetitions.

Likewise, within-region (inter- and intra-subject correlation) and between-region (inter- and intra-subject functional correlation) analyses offer complementary information, because regions can change how they communicate with one another, whether or not a given region on its own shows changes in activity (Al-Aidroos et al., 2012; Córdova et al., 2016). These analyses therefore offer a complementary window into learning in the brain.

In sum, by comprehensively examining temporal dynamics in the posterior medial network over movie clip repetitions, we tested how the network's responses change with exposure to novel and schematic temporal structure. The critical question was in what ways the posterior medial network could become engaged during viewing of novel temporal structure (in scrambled movies), or

if, alternatively, it is only engaged by temporal structure that is consistent with pre-existing schema (in intact movies).

Materials and Methods

Subjects

Thirty individuals (12 male) from the Princeton University community participated for monetary compensation (age: $M = 23.0$ years, $SD = 4.2$; education: $M = 15.3$ years, $SD = 3.2$; all right-handed). The sample size was determined *a priori* based on prior fMRI studies using naturalistic stimuli and employing analysis techniques identical to those in the current study (e.g., Andric et al., 2016; Chen et al., 2016, 2017; Hasson et al., 2004, 2008; Honey et al., 2012b; Lerner et al., 2011; Simony et al., 2016). Although the sample size varies across these studies, the average is about 15 subjects. Because we had two counterbalancing conditions (see below), we opted to run 15 subjects in each, for a total sample size of 30.

We only tested individuals who had not previously seen the movie used in this study. Informed consent was obtained from all subjects, and the study was approved by the Institutional Review Board at Princeton University.

Design and Procedure

Three 90-s clips (referred to as A, B, and C below) were selected from *The Grand Budapest Hotel*, and viewed six times each. Each clip included multiple characters, dialogue, music, and movement through several rooms and hallways. Each clip was assigned to one of three conditions (Figure 1). All subjects viewed all three clips. The accompanying audio (dialogue and music) was delivered with MRI-compatible Sensimetrics in-ear headphones. To make the stimuli for each condition, we divided each 90-s clip into short segments, keeping the divisions at natural breaks as much as possible (e.g., an editor's cut, the end of a spoken sentence). Clip A was divided into 27 segments (2.5–4.4 s). Clip B was divided into 24 segments (2.4–5.5 s). Clip C was divided into 26 segments (2.7–4.6 s).

The *Intact* clip was viewed in its original format, as a temporally continuous clip with unfolding events consistent with schemas learned over the course of our lifetime (e.g., what happens when an elevator button is pushed, or what happens when walking into a hotel). This clip was reconstructed in its original order from the shorter segments, so that any audiovisual artifacts introduced by editing would be present for the Intact condition as well as the Scrambled conditions described below. For all subjects, the Intact clip was clip A. For the *Scrambled-Fixed* clip, the segments were randomly re-ordered at first (disrupting consistency with event schemas in semantic memory), but then viewed in the same scrambled order for all six repetitions of the clip (thus, segment transitions were perfectly predictable after the first presentation, providing the basis for learning temporal structure over repetitions). For the *Scrambled-Random* clip, the segments were randomly re-ordered separately for each of the six repetitions of the clip (thus, segment transitions were unpredictable, and segment-level temporal structure was eliminated). There were two counterbalancing conditions: For half of the subjects, the Scrambled-Fixed clip was clip B and the Scrambled-Random clip was clip C, and vice versa for the other half. All subjects in a given counterbalancing condition viewed identical stimuli, i.e., the Scrambled-Fixed clip was scrambled in the same way for all subjects (and for all repetitions), and the Scrambled-Random clip was scrambled in the same way for all subjects for any given repetition (but scrambled differently for different repetitions).

To summarize the design: The Intact and Scrambled-Fixed clips were both repeated identically for all six repetitions. The former was in its original, continuous format; thus, temporal expectations existed on the very first exposure because of consistency with real-world event schemas. The latter became temporally predictable over repetitions because of deterministic transitions between re-ordered movie segments. The Scrambled-Fixed and Scrambled-Random clips both contained random events inconsistent with real-world predictions, but the latter was viewed in a different scrambled order every time. Although the Scrambled-Random clip contained temporal structure on short

timescales (e.g., within the ~3s duration of each segment), longer timescale structure across segments

was present only in the Intact and Scrambled-Fixed clips.

Figure 1. Stimuli and design. Subjects viewed 90-s clips from *The Grand Budapest Hotel*. Each of three clips was assigned to one of three conditions. Each clip was viewed six times. Numbers below each screenshot refer to a segment within each clip, with segments being continuous chunks 2.4–5.5 s long. The Intact clip was viewed in its original format. The segments for the Scrambled-Fixed clip were randomly re-ordered, but viewed in the same scrambled order for all repetitions. The segments for the Scrambled-Random clip were also randomly re-ordered, but the segments were viewed in a different order for each repetition.

The experiment was divided into three runs, each ~10 minutes long. Each run began with 4.5 s

of blank time and then an additional 6-s countdown screen to alert subjects to the upcoming start of

the movie clips. The rest of each run comprised six movie clips in total (clips A, B, and C viewed twice each), with 3 s of fixation and then 6 s of countdown between each clip. Across runs, A, B, and C were each viewed in each of the six serial positions in a run (e.g., A was viewed in positions one and five in run one, positions two and four in run two, and positions three and six in run three). The same clip was never viewed back-to-back within a run. The within-run lag between repetitions of a clip was matched across the clips: in one run, there was a lag of one between repetitions, in another there was a lag of two, and in another there was a lag of three between repetitions of the same clip. All subjects viewed A, B, and C in the same order, but the assignment of clip to condition differed across subjects.

Subjects were instructed to view the clips, but there was no other task during the scan. It was emphasized that attention should be paid at all times. To verify this, we surprised subjects with a post-scan free recall test, which required them to type everything they could remember about each of the clips. Subjects were generally able to recall the main events and characters in each clip and (varying amounts of) perceptual details.

fMRI Methods

Acquisition. Data were collected on a 3T Siemens Prisma scanner with a 64-channel head/neck coil. Functional images were obtained with a multiband EPI sequence (TR = 1.5 s, TE = 39 ms, flip angle = 50°, acceleration factor = 4, shift = 3, voxel size = 2.0-mm iso), with 60 oblique axial slices acquired in an interleaved order. Whole-brain high-resolution (1.0-mm iso) T1-weighted structural mages were acquired with an MPRAGE sequence. Field maps were collected for registration, consisting of 40 oblique axial slices (3-mm iso).

Software. Preprocessing, registration, and permutation tests for whole-brain analyses were conducted using FEAT, FLIRT, and command-line functions in FSL (http://fsl.fmrib.ox.ac.uk/fsl/). All other analyses were performed with custom Matlab scripts.

Preprocessing. The first three volumes of each run were discarded to allow for T1 equilibration. Preprocessing steps included brain extraction, motion correction, high-pass filtering (max period =

140 s), spatial smoothing (3-mm FWHM Gaussian kernel), and registration to standard MNI space (via an intermediate registration to the subject's anatomical image with boundary-based registration).

The fields maps were preprocessed using a custom script, in accordance with the FUGUE user guide from FSL. First, the two field map magnitude images were averaged together and skull stripped. The field map phase image was converted to rad/s and smoothed with a 2-mm Gaussian kernel. The resulting phase and magnitude images were included in the preprocessing step of FEAT analyses to unwarp the functional images and aid registration to anatomical space. Following registration, the distortion-corrected functional images were compared to the originals to ensure that unwarping was effective. In all cases, using the field maps reduced distortion in anterior temporal and frontal regions.

After preprocessing, the filtered 4D functional images for each run were divided into volumes that corresponded to each of the six clips presented within that run. Four volumes were then cropped from the beginning of each clip to adjust for hemodynamic lag. Finally, the timecourse for each clip was z-scored within each subject and region of interest (ROI).

ROI definition. We focused our analyses primarily on the precuneus, as a key node in the posterior medial network (Hasson et al., 2015; Ranganath and Ritchey, 2012), and a region that is consistently and robustly recruited by long-timescale regularities (Baldassano et al., 2017; Chen et al., 2016, 2017; Hasson et al., 2010, 2015; Honey et al., 2012b; Lerner et al., 2011; Utevsky et al., 2015). In addition to examining activity within the precuneus, we investigated its interactions with three other ROIs in the posterior medial network: the hippocampus, angular gyrus, and posterior cingulate cortex (Ranganath & Ritchey, 2012); we could not include medial prefrontal cortex because of excessive signal dropout. For completeness, we also examined all pairwise interactions between the four posterior medial ROIs. These ROIs were defined bilaterally with the Harvard-Oxford Structural Atlas. The precuneus and posterior cingulate ROIs were edited to not contain overlapping voxels, with the precuneus ROI comprising dorsomedial parietal cortex and the posterior cingulate ROI comprising ventromedial parietal cortex.

We also examined a control ROI in early visual cortex, chosen for its relatively short temporal integration window in prior studies (e.g., Chen et al., 2016; Hasson et al., 2008). The ROI was defined functionally from a separate group of 25 subjects watching two minutes of an audiovisual movie (Chen et al., 2016), including voxels bilaterally along the collateral sulcus with the highest inter-subject correlation (rs > 0.45).

The timecourse of activity was averaged across all voxels in an ROI, and these mean ROI timecourses were submitted to further analysis. We conducted within-region and between-region analyses, within-individual (intra-subject) and between individuals (inter-subject), for complementary insights into how the brain changes with learning. These analyses, along with their logic, are described in more detail below.

Within-region correlations. We examined the consistency of BOLD responses to the movie clips in two ways, both of which have been used in prior studies of temporal integration with naturalistic stimuli (e.g., Hasson et al., 2005, 2008; Honey et al., 2012a,b; Lerner et al., 2011): (1) intra-subject correlation over clip repetitions, and (2) inter-subject correlation for the same clip repetition. As noted above, for these analyses we focused on the precuneus; subsequent analyses incorporated additional regions.

To assess the consistency of precuneus activity to a given clip within an individual (i.e., intra-subject correlation), the timecourse of activity for repetition 1 of each clip was compared to that of repetition 6 of the same clip using Pearson correlation. For the Intact and Scrambled-Fixed clips, the order of segments within each clip was the same across repetitions, thus brain activity was being compared for objectively identical stimuli in repetitions 1 and 6. For the Scrambled-Random clip, segment order was different across all repetitions, thus brain activity was being compared for *different* stimuli in repetitions 1 and 6. Thus, this condition serves as a baseline because correlations should approach zero even in early sensory areas.

Planned comparisons between conditions were conducted at the group level with random-effects paired *t*-tests, after Fisher transformation of the correlation coefficients to ensure normality. Two-tailed *p*-values and 95% confidence intervals are reported for this and all other analyses, as well as the correlation values as a measure of effect size.

To assess the consistency of precuneus activity to a given clip across individuals (i.e., inter-subject correlation), we correlated each subject's timecourse of activity for repetition 1 of each clip with the mean of all *other* subjects' timecourses for that same repetition and clip. Because subjects viewed the same stimulus but may have had different thoughts and emotions triggered by it, correlating timecourses across individuals highlights the stimulus-locked brain activity that is shared across people irrespective of these individual differences. In contrast, the intra-subject correlation analysis described above can identify idiosyncratic responses that are particular to a person and/or asynchronous across people (Hasson et al., 2004, 2009).

For this analysis, we focused on the first repetition of each clip because this allowed a measure of the consistency of precuneus dynamics between subjects upon initial exposure, before learning could influence processing, and additionally because this allowed us to compare our results to published studies with only one repetition (e.g., Chen et al., 2016; Hasson et al., 2004; Lerner et al., 2011). Nevertheless, for completeness, we also examined the inter-subject correlation for the last repetition of each clip (Supplementary Figure S1).

There were two clip-to-condition counterbalancing subgroups (i.e., clip B was the Scrambled-Fixed clip for half of subjects and Scrambled-Random for the other half), so inter-subject correlation was calculated within each subgroup and then pooled. In this way, we obtained the correlation between brain activity of different people viewing identical stimuli (this includes identical segment ordering for the scrambled clips). Specifically, the brain activity of each of the subjects numbered 1-15 was correlated against the average of the other 14 subjects numbered 1-15 (these subjects watched identical clips). Likewise, the brain activity of each of the subjects numbered 16-30 was correlated

against the average of the other 14 subjects numbered 16-30 (these subjects watched identical clips). Thus, each subject had a correlation value for his/her brain activity vs. the mean of all other subjects in his/her counterbalancing group. All 30 correlations were then used for group analyses. We conducted inter-subject analyses for the Intact clip within the two subgroups of 15 subjects in the same manner (even though all subjects saw the same clip). Thus, this analysis was consistent and equivalently powered across conditions.

The resulting inter-subject correlation data are not independent, however: For n subjects in the same counterbalancing group, any pair of subjects will share $n - 2$ timecourses in the calculation of the mean "others" timecourse. For example, the "others" timecourse for subject 1 contains data from subjects 2-15, and the "others" timecourse for subject 2 contains data from subjects 1, 3-15. Thus, data for subjects 3-15 were used to calculate the inter-subject correlation for both subject 1 and subject 2. The inter-subject correlation values for different subjects are therefore not independent. Because of this, we conducted non-parametric permutation tests to compare the mean inter-subject correlation for different conditions. For each subject, we took the difference between the two conditions of interest (after Fisher-transforming the correlations) and then flipped the sign of the difference (effectively swapping condition assignments) for some random subset of subjects before obtaining the group mean. This was repeated 1000 times, and the distribution of mean differences for the randomly flipped data was compared to the observed mean difference between conditions to obtain p-values. Two-tailed p-values were obtained by finding the smaller tail of the null distribution vs. the observed mean and multiplying that value by two.

In addition to this approach, we also calculated confidence intervals (CIs) on the mean difference between conditions using a bootstrap approach. For each subject, we took the difference between the two conditions of interest (after Fisher-transforming the correlations), randomly sampled subjects with replacement, and calculated the mean from this sample. This was repeated 1000 times, and the points below and above which 2.5% of the samples lay (i.e., the bounds that contained 95% of

the data) were the bootstrap confidence intervals. Both of these approaches (i.e., the permutation test and the bootstrapped CIs) yielded comparable results to paired-samples *t*-tests.

Between-region correlations. Intra-subject functional correlation between ROIs was examined across all repetitions of each clip to see how neural coupling changes with learning. For each repetition and clip, the timecourse of BOLD activity in the precuneus was correlated with that of the other ROIs, for the same repetition and clip. After Fisher transformation, four types of analyses were conducted to assess changes in intra-subject functional correlation: (1) a 2 x 3 repeated-measures analysis of variance (ANOVA) with repetition (1 or 6) and condition (Intact, Scrambled-Fixed, Scrambled-Random) as factors; (2) planned comparisons of repetitions 1 vs. 6 within condition using paired *t*-tests; (3) a 6 x 3 repeated-measures ANOVA with repetition (1-6) and condition (Intact, Scrambled-Fixed, Scrambled-Random) as factors; and (4) a test of whether the slope across repetitions 1-6 within condition was different from zero, using one-sample *t*-tests.

Together, these four analyses offered a complete picture of learning-related changes in functional correlation: Analyses 1 and 2 allowed us to examine intra-subject functional correlation at the beginning vs. end of learning (repetitions 1 vs. 6), whereas analyses 3 and 4 allowed us to examine the nature of change over the course of all six repetitions. The ANOVAs (analyses 1 & 3) allowed us to examine whether differences between conditions and repetitions existed, whereas the more-targeted within-condition analyses (analyses 2 & 4) allowed us to examine learning-related changes in each condition separately.

For completeness, we also examined intra-subject functional correlation between all pairs of posterior medial ROIs. To visualize changes in functional coupling, we made network graphs for the first vs. last repetitions using a Matlab toolbox *circularGraph*, where each posterior medial ROI was a node, and the edges represented the strength of coupling between all pairs of ROIs.

To assess the selectivity of changes in intra-subject functional correlation within the posterior medial ROIs, we conducted two additional analyses. First, we examined intra-subject functional

correlation between the precuneus and a control ROI in early visual cortex. Second, we conducted whole-brain, voxelwise intra-subject functional correlation analyses using the precuneus as the seed ROI. For each subject, clip, and repetition, the timecourse of BOLD activity in the precuneus was correlated with the timecourse of every other voxel in the brain. A contrast map of repetition 6 minus repetition 1 was generated for each subject and condition. Within each condition, these contrast maps were analyzed at the group level using random-effects non-parametric tests (randomise in FSL), and corrected for family-wise error (FWE) across all voxels. The resulting maps were thresholded at $p < 0.05$ corrected.

We also conducted functional correlation analyses between subjects. Inter-subject functional correlation (Simony et al., 2016) is analogous to the inter-subject correlation analyses described above, but the timecourse of activity for a given ROI and subject is correlated to the mean of all other subjects' timecourses for a different ROI (within repetition and condition). For example, to calculate inter-subject functional correlation between the hippocampus and precuneus, we first obtained the correlation between the hippocampal timecourse for a particular individual and the mean precuneus timecourse across all other individuals. Then, we calculated the correlation between the precuneus timecourse for the same individual and the mean hippocampal timecourse across all other individuals. The average of these calculations is taken as the measure of hippocampus-precuneus inter-subject functional correlation for that subject. The analysis is repeated for all subjects, and the resulting data are submitted to group-level analysis. As with the inter-subject correlation analysis, we calculated inter-subject functional correlation within each subgroup of 15 subjects that viewed identical stimuli, and then pooled across subjects. The resulting data were analyzed using permutation tests and bootstrap methods, as above.

<center>**Results**</center>

Within-Region Correlations

Prior studies have found that temporal dynamics within posterior medial regions, such as the precuneus, are more consistent for intact vs. scrambled movies and stories. Our first aim was to replicate these findings — specifically, to show that temporal fluctuations in the precuneus are more highly correlated for intact vs. scrambled stimuli when measured: (1) within an individual across repetitions of these clips, i.e., intra-subject correlation (e.g., Hasson et al., 2004, 2008; although only two repetitions were used in that study), and (2) between individuals upon initial exposure to these clips, i.e., inter-subject correlation (e.g., Honey et al., 2012b; Lerner et al., 2011).

For the intra-subject correlation analysis, it is important to note that the Scrambled-Random clip was an objectively different stimulus on the first and last repetitions (i.e., the segments within the clip were presented in a different order), but the Intact and Scrambled-Fixed stimuli were objectively identical on the first and last repetition. Thus, the intra-subject correlation analysis for the Scrambled-Random condition is expected to yield low correlations due to changes in the stimulus across runs. The intra-subject correlation for the Scrambled-Random clip is therefore presented here only for completeness, as it is not expected to be significantly different from zero.

As shown in Figure 2A, and consistent with prior studies, there was higher intra-subject correlation between precuneus activity for the first and last repetitions of the Intact clip compared to the correlation between the first and last repetitions of the Scrambled-Fixed clip (r_{intact} = 0.21, $r_{S\text{-Fix}}$ = 0.05; t_{29} = 2.78, p = .009, 95% CI: 0.05–0.32). The correlation for the Intact clip was also higher than that for the Scrambled-Random clip (r_{intact} = 0.21, $r_{S\text{-Rnd}}$ = -0.05; t_{29} = 4.26, p = .0002, 95% CI: 0.15–0.44), although, as noted above, the correlation for the Scrambled-Random clip was expected to be near zero because the stimulus is different on the first vs. last repetition. Response reliability for the Intact clip was also significantly greater than zero (t_{29} = 4.13, p = .0003, 95% CI: -0.12–0.36).

<center>16</center>

Figure 2. Temporal dynamics in the precuneus are more consistent for intact vs. scrambled movies. (A) The timecourses of BOLD activity in the precuneus for the first and last repetitions of each clip were extracted and correlated within subject. The mean intra-subject correlation was higher for the Intact clip compared to the Scrambled-Fixed and Scrambled-Random clips, which did not themselves differ. **(B)** Reliability was also assessed by correlating, for the first repetition, each subject's timecourse of BOLD activity with the mean of all other subjects. The mean inter-subject correlation was higher for the Intact clip compared to the Scrambled-Fixed and Scrambled-Random clips, which did not themselves differ. Dots indicate individual subjects. Error bars are ± 1 SEM. ** $p < .01$, *** $p < .001$.

Next, we compared the reliability of intra-subject precuneus dynamics for the Scrambled-Fixed and Scrambled-Random clips. Neither of these clips were consistent with pre-existing event schemas, but the Scrambled-Fixed clip contains predictable segment sequences over repetitions (and thus, temporal structure on the order of tens of seconds). However, learning of temporal structure in the Scrambled-Fixed clip would not be expected to increase the across-repetition reliability of a region: in fact, the region should be *less* similar to itself over repetitions if learning changes how it processes information. Intra-subject correlations over repetitions should also be low for the

Scrambled-Random clip, but for a different reason: because the clip is viewed in a different order every time, there is a different stimulus for repetitions 1 and 6. Indeed, the intra-subject correlation between precuneus activity in repetitions 1 and 6 was not different from zero in either condition (Scrambled-Fixed: $r = 0.05$; $t_{29} = 1.10$, $p = .28$, 95% CI: -0.05–0.16; Scrambled-Random: $r = -0.05$ $t_{29} = 1.43$, $p = .16$, 95% CI: -0.13–0.02) and the conditions were not different from one another ($t_{29} = 1.64$, $p = .11$, 95% CI: -0.03–0.25).

A similar pattern of results was obtained when examining the reliability of precuneus activity between subjects on the first repetition of each clip, before learning-related changes could occur (Figure 2B; for the same analysis for the last repetition of each clip, see Supplementary Figure S1). In this case, we examined inter-subject correlation between individuals watching identical stimuli, i.e., the scrambled movies were watched in the same scrambled order for these different subjects. The inter-subject correlation for the Intact clip was higher than that for the Scrambled-Fixed clip ($r_{intact} = 0.58$, $r_{S-Fix} = 0.24$; 95% bootstrapped CI: 0.28–0.57; $p = 0$ from permutation test) and the Scrambled-Random clip ($r_{intact} = 0.58$, $r_{S-Rnd} = 0.36$; 95% bootstrapped CI: 0.19–0.40; $p = 0$ from permutation test). The inter-subject correlation for the Scrambled-Fixed clip was marginally lower than that for the Scrambled-Random clip (95% bootstrapped CI: -0.27–0.01; $p = .09$ from permutation test). However, because the stimuli in these conditions did not differ in the first repetition (i.e., before the scrambled order had been repeated), this difference is not meaningful. Finally, inter-subject correlation in the precuneus was reliably above zero for all three conditions, as expected because all subjects being compared viewed identical stimuli in all conditions (all ps = 0 from permutation tests).

Together, these within-region analyses replicate and extend prior work showing greater response reliability in posterior medial regions for intact vs. scrambled movies and stories (e.g., Hasson et al., 2008; Honey et al., 2012a,b; Lerner et al., 2011).

Between-Region Correlations

Low inter-subject correlation (Figure 2B), as observed for the Scrambled-Fixed and Scrambled-Random clips, indicates that temporal dynamics in the precuneus are not locked to the external stimulus that is viewed by all subjects. However, intra-subject dynamics were also unstable (Figure 2A). Changing temporal dynamics over clip repetitions could indicate learning, but they can also occur for other reasons (e.g., fatigue). Thus, to further search for learning effects, we examined network functional correlations during each clip repetition within each individual (c.f. Andric et al., 2016). Analyses of intra-subject functional correlation can be useful for understanding how brain regions respond to naturalistic stimuli. For example, functional coupling between the hippocampus and medial prefrontal cortex is modulated by prior exposure to schema-relevant information while watching an intact movie (van Kesteren et al., 2010). We hypothesized that learning of temporal structure in the Scrambled-Fixed clip would manifest as enhanced intra-subject functional coupling, over clip repetitions, between the precuneus and other regions in the posterior medial network: hippocampus, angular gyrus, and posterior cingulate cortex (Figure 3).

We expected that changes in network intra-subject functional correlations over repetitions would differ between the three conditions. To verify this, we conducted 2 (repetition: 1 or 6) by 3 (condition: Intact, Scrambled-Fixed, Scrambled-Random) repeated-measures ANOVAs, looking specifically for an interaction. Such an interaction was found for coupling between hippocampus and precuneus ($F_{2, 58} = 6.12$, $p = .004$). Follow-up planned comparisons revealed a reliable increase in intra-subject functional correlation from repetition 1 to 6 for the Scrambled-Fixed clip ($r_{Rep1} = 0.17$, $r_{Rep6} = 0.42$, $t_{29} = 4.38$, $p = .0001$, 95% CI: 0.16–0.43) but not the Intact clip ($r_{Rep1} = 0.42$, $r_{Rep6} = 0.43$, $t_{29} = 0.33$, $p = .74$, 95% CI: -0.11–0.15) or the Scrambled-Random clip ($r_{Rep1} = 0.23$, $r_{Rep6} = 0.32$, $t_{29} = 1.98$, $p = .06$, 95% CI: -0.005–0.28). Indeed, while intra-subject functional correlation was initially higher for the Intact clip than the Scrambled-Fixed clip in repetition 1 ($r_{Intact} = 0.42$, $r_{S-Fix} = 0.17$, $t_{29} = 5.08$, $p = .00002$, 95% CI:

0.16–0.38), this difference disappeared by repetition 6 (r_{Intact} = 0.43, r_{S-Fix} = 0.42, t_{29} = 0.0001, p = .9999, 95% CI: -0.13–0.13).

Figure 3. Temporal structure enhances intra-subject functional correlation within the posterior medial network. (A) The timecourse of BOLD activity in the precuneus was correlated within-subject with that of the hippocampus for the first and last repetition of each clip. Functional correlations did not change over repetitions of the Intact or Scrambled-Random clips, but more than doubled over repetitions of the Scrambled-Fixed clip, rising to the level of the Intact clip. A similar pattern of results was obtained for **(B)** the precuneus and angular gyrus and **(C)** the precuneus and posterior cingulate. Dots indicate individual subjects. Error bars are ± 1 SEM of the within-subject difference between repetitions. * p < .05, ** p < .01, *** p < .001.

To obtain additional evidence that increased intra-subject functional correlation from the first to last repetition of the Scrambled-Fixed clip reflected learning, we examined these functional correlations across all six repetitions. If these changes are in fact indicative of learning, enhancements in intra-subject functional correlation should be monotonic, increasing gradually with exposure to the temporal structure (Supplementary Figure S2). We conducted a 6 (repetition: 1-6) by 3 (condition: Intact, Scrambled-Fixed, Scrambled-Random) repeated-measures ANOVA, which again revealed a significant interaction for precuneus-hippocampus intra-subject functional correlation ($F_{10, 290}$ = 2.82, p = .002). This interaction was driven by a significant monotonic increase in intra-subject functional correlation over repetitions for the Scrambled-Fixed clip, manifested as a slope greater than zero (β = 0.05, t_{29} = 4.06, p = .0003, 95% CI: 0.02–0.07). The slope was not different from zero for the Intact clip (β = 0.004, t_{29} = 0.43, p = .67, 95% CI: -0.02–0.02) and was numerically smaller for the Scrambled-Random clip (β = 0.03, t_{29} = 2.17, p = .04, 95% CI: 0.002–0.06).

Neither ANOVA revealed a significant repetition x condition interaction for precuneus-angular gyrus intra-subject functional correlation (repetitions 1 vs. 6: $F_{2, 58}$ = 1.87, p = .16; repetitions 1 through 6: $F_{10, 290}$ = 0.96, p = .47) or precuneus-posterior cingulate cortex intra-subject functional correlation (repetitions 1 vs. 6: $F_{2, 58}$ = 2.08, p = .13; repetitions 1 through 6: $F_{10, 290}$ = 1.78, p = .06). Nevertheless, within the Scrambled-Fixed condition, intra-subject functional correlation increased from repetitions 1 vs. 6 (precuneus-angular gyrus: r_{Rep1} = 0.39, r_{Rep6} = 0.65, t_{29} = 5.20, p = .00001, 95% CI: 0.23–0.54; precuneus-posterior cingulate cortex: r_{Rep1} = 0.40, r_{Rep6} = 0.63, t_{29} = 5.24, p = .00001, 95% CI: 0.20–0.46) and the slope across all six repetitions was greater than zero (precuneus-angular gyrus: β = 0.07, t_{29} = 4.68, p = .00006, 95% CI: 0.04–0.09; precuneus-posterior cingulate cortex: β = 0.06, t_{29} = 4.76, p = .00005, 95% CI: 0.03–0.08). The interactions failed to reach significance because of (numerically smaller) increases in precuneus-angular gyrus intra-subject functional correlation for both the Intact clip (repetitions 1 vs. 6: r_{Rep1} = 0.53, r_{Rep6} = 0.64, t_{29} = 3.23, p = .003, 95% CI: 0.08–0.37; slope across

repetitions: $\beta = 0.04$, $t_{29} = 3.43$, $p = .002$, 95% CI: 0.02–0.06) and the Scrambled-Random clip (repetitions 1 vs. 6: $r_{Rep1} = 0.44$, $r_{Rep6} = 0.56$, $t_{29} = 2.42$, $p = .02$, 95% CI: 0.03–0.40; slope across repetitions: $\beta = 0.04$, $t_{29} = 2.42$, $p = .02$, 95% CI: 0.007–0.08), and in precuneus-posterior cingulate cortex intra-subject functional correlation for the Scrambled-Random clip (repetitions 1 vs. 6: $r_{Rep1} = 0.44$, $r_{Rep6} = 0.55$, $t_{29} = 2.62$, $p = .01$, 95% CI: 0.05–0.41; slope across repetitions: $\beta = 0.04$, $t_{29} = 2.48$, $p = .02$, 95% CI: 0.007–0.08).

As with precuneus-hippocampus intra-subject functional correlation, both precuneus-angular gyrus intra-subject functional correlation and precuneus-posterior cingulate intra-subject functional correlation were higher for the Intact clip compared to the Scrambled-Fixed clip on repetition 1 (precuneus-angular gyrus: $r_{Intact} = 0.53$, $r_{S-Fix} = 0.39$, $t_{29} = 2.20$, $p = .04$, 95% CI: 0.01–0.34; precuneus-posterior cingulate cortex: $r_{Intact} = 0.62$, $r_{S-Fix} = 0.40$, $t_{29} = 3.83$, $p = .0006$, 95% CI: 0.14–0.48), but this difference disappeared by repetition 6 (precuneus-angular gyrus: $r_{Intact} = 0.64$, $r_{S-Fix} = 0.65$, $t_{29} = 0.27$, $p = .79$, 95% CI: -0.14–0.18; precuneus-posterior cingulate cortex: $r_{Intact} = 0.67$, $r_{S-Fix} = 0.63$, $t_{29} = 1.49$, $p = .15$, 95% CI: -0.04–0.26).

We visualized these effects (and others not reported above) by plotting graphs of all possible pairwise relationships between posterior medial ROIs (Figure 4). There was overall increased functional coupling in the network from repetitions 1 to 6 of the Scrambled-Fixed clip, but less consistent enhancement for the Intact and Scrambled-Random clips.

Figure 4. Changes in posterior medial network functional correlations from first to last repetition. Graph visualization of intra-subject functional correlation between all pairs of posterior medial ROIs, including the results from Figure 3, as well as pairs that did not include precuneus. The edges depict the *t* value comparing intra-subject functional correlation in repetitions 1 vs. 6, with thicker lines indicating greater enhancement over time. Solid lines are statistically significant changes, whereas dashed lines are not statistically significant. The most consistent enhancement in intra-subject functional correlation was for the Scrambled-Fixed clip (all six edges were statistically significant), compared to the Intact clip (two statistically significant edges) and the Scrambled-Random clip (three statistically significant edges). PCC = posterior cingulate cortex, HPC = hippocampus, PCN = precuneus, ANG = angular gyrus.

To assess the selectivity of enhanced functional coupling within the posterior medial network for the Scrambled-Fixed clip, we conducted two additional analyses. First, we examined intra-subject functional correlation between the precuneus and a control region in early visual cortex (Figure 5). Unlike the posterior medial network, there were no changes in intra-subject functional correlation between the precuneus and early visual cortex over repetitions in any condition (all *ps* > .46). Moreover, there were no main effects of repetition or condition, nor repetition by condition interactions in either ANOVA (i.e., repetitions 1 vs. 6 and repetitions 1 through 6, all *ps* > .15).

Figure 5. Functional correlation between the precuneus and early visual cortex. Intra-subject functional coupling between precuneus and visual cortex did not increase over repetitions in any condition. Dots are individual subjects. Error bars are ± 1 SEM of the within-subject difference between repetitions.

Next, we conducted whole-brain analyses of intra-subject functional correlation with the precuneus as a seed ROI, for the first and last repetitions of the Intact, Scrambled-Fixed, and Scrambled-Random clips. There were no reliable increases in intra-subject functional correlation for repetition 1 vs. 6 of the Intact or Scrambled-Random clips, correcting for multiple comparisons. In contrast, for the Scrambled-Fixed clip, significant increases in intra-subject functional correlation was observed in the hippocampus, angular gyrus, and medial parietal cortex (Figure 6), consistent with the ROI analyses of the posterior medial network. Although the medial parietal cluster overlaps with the seed ROI, this result indicates *changes* in intra-subject functional correlation over repetitions and thus is not guaranteed.

Figure 6. Changes in whole-brain precuneus (intra-subject) functional correlations from first to last repetition. Contrast depicts regions that increase in functional coupling with the precuneus (purple) from the first to last repetition of each clip ($p < .05$ corrected). No regions showed enhanced intra-subject functional correlation for the Intact (top) and Scrambled-Random (bottom) clips. For the Scrambled-Fixed clip (middle), greater precuneus intra-subject functional correlation (in green) was observed in the hippocampus and angular gyrus (and a medial parietal region that overlapped with the seed). The black mask (right) indicates to which voxels this analysis was applied (where > 90% of subjects had coverage).

What is being learned in the Scrambled-Fixed condition that increases functional coupling in the posterior medial network? One possibility is that stronger intra-subject functional correlation reflects the regions' activity becoming more locked to the stimulus. That is, upon initial exposure, the Scrambled-Fixed clip may seem incomprehensible and unpredictable, and thus is not consistently processed by different regions within the network. The extraction of temporal regularities over repetitions may allow posterior medial regions to focus on the same, regular aspects of the stimulus. This explanation can be tested by examining whether components of functional correlations that are locked to the stimulus increase over learning. Such stimulus-locked components can be measured

with inter-subject functional correlation (Simony et al., 2016). In inter-subject functional correlation analyses, correlations are calculated between the timecourse of activity in a given region in one subject and the timecourses of activity in other regions averaged over all other subjects. As with inter-subject correlations (the same region between subjects), inter-subject functional correlation analysis isolates dynamics of brain activity attributable to the external stimulus viewed by all subjects, as opposed to components that are idiosyncratic to each individual or asynchronous across individuals. We calculated inter-subject functional correlation between all pairs of posterior medial network ROIs and found no increase across repetitions in any condition (Supplementary Figure S3). Indeed, the only robust effects were *decreases* from repetition 1 to repetition 6 for HPC-PCN (Intact: r_{Rep1} = 0.31, r_{Rep6} = 0.07, p =0 from permutation test, 95% bootstrapped CI: 0.19–0.30; Scrambled-Fixed: r_{Rep1} = 0.11, r_{Rep6} = 0.03, p = .04 from permutation test, 95% bootstrapped CI: 0.007–0.16), PCC-PCN (Intact: r_{Rep1} = 0.37, r_{Rep6} = 0.12, p = 0 from permutation test, 95% bootstrapped CI: 0.16–0.36), and HPC-PCC (Intact: r_{Rep1} = 0.21, r_{Rep6} = 0.03, p = 0 from permutation test, 95% bootstrapped CI: 0.12–0.28; Scrambled-Random: r_{Rep1} = 0.05, r_{Rep6} = -0.09, p = .002 from permutation test, 95% bootstrapped CI: 0.08–0.21). This pattern of results — increased network coupling for the Scrambled-Fixed condition within but not between subjects — indicates that learning in the posterior medial network did not increase how much its responses were locked to the external stimulus. Moreover, this lack of an increase for inter-subject between-region correlations mirrors the lack of an increase in inter-subject within-region correlations (Supplementary Figure S1).

An alternative explanation, consistent with the lack of inter-subject enhancements in correlated temporal dynamics, is that what is being learned may be idiosyncratic to individuals. For example, over repetitions, each individual may converge on a unique meaning of the Scrambled-Fixed clip or learn particular segment transitions. This account predicts that the temporal dynamics within a region will be more similar for clip repetitions toward the end vs. beginning of learning. That is, many learning-related changes may occur early during exposure, with a particular mode of processing or

interpretation of the movie arrived at by the end of learning. Thus, for the Scrambled-Fixed clip, the correlation between within-region temporal dynamics for repetitions 1 vs. 2 may be lower than that for repetitions 5 vs. 6. However, this was not found: the within-region intra-subject correlation was not different for repetitions 1 and 2 compared to repetitions 5 and 6 of the Scrambled-Fixed clip (ANG: r_{Rep1-2} = -0.02, r_{Rep5-6} = -0.04, t_{29} = 0.25, p = .80, 95% CI: -0.14–0.18; HPC: r_{Rep1-2} = 0.0003, r_{Rep5-6} = 0.02, t_{29} = 0.32, p = .75, 95% CI: -0.11–0.08; PCC: r_{Rep1-2} = 0.06, r_{Rep5-6} = -0.01, t_{29} = 1.05, p = .30, 95% CI: -0.08–0.24; PCN: r_{Rep1-2} = 0.15, r_{Rep5-6} = 0.08, t_{29} = 0.91, p = .37, 95% CI: -0.10–0.25).

There were also no differences between these repetition pairs for the Scrambled-Random clip (all ps > .62); no differences are expected here because the Scrambled-Random clip was viewed in a different order for each repetition, thus brain activity is being compared for objectively different stimuli.

For the Intact clip, we found greater similarity *early* in learning in precuneus and hippocampus (precuneus: r_{Rep1-2} = 0.44, r_{Rep5-6} = 0.13, t_{29} = 5.33, p = .0001, 95% CI: 0.22–0.48; hippocampus: r_{Rep1-2} = 0.18, r_{Rep5-6} = 0.08, t_{29} = 2.34, p = .03, 95% CI: 0.13–0.20). This effect for the Intact clip was not reliable in angular gyrus (r_{Rep1-2} = 0.19, r_{Rep5-6} = 0.08, t_{29} = 1.78, p = .09, 95% CI: -0.02–0.25) or posterior cingulate (r_{Rep1-2} = 0.17, r_{Rep5-6} = 0.08, t_{29} = 1.54, p = .13, 95% CI: -0.03–0.24). These results are inconsistent with the idea of gradually arriving at a particular manner of processing the movie clip.

Discussion

The brain contains a hierarchy of regions that respond to information over varying timescales (Chaudhuri et al., 2015; Hasson et al., 2008; Murray et al., 2014). Unlike sensory regions, in which activity at any given moment is predominantly driven by the immediate environment, higher-order regions in the posterior medial network integrate events over many seconds or minutes (Baldassano et al., 2017; Chen et al., 2016; Hasson et al., 2015; Simony et al., 2016). Prior studies have often explored temporal integration by analyzing within-region dynamics for naturalistic stimuli. The reliability of a

region's temporal dynamics is calculated either by examining how highly that region's activity is correlated between different subjects watching the same stimulus, or within a subject over repetitions of the same stimulus (e.g., Hasson et al., 2004, 2008; Honey et al., 2012a,b; Lerner et al., 2011). These studies find that the reliability of temporal fluctuations in posterior medial regions, whether measured between subjects or within a subject, is highest for intact movies and stories and is reduced when these stimuli are temporally scrambled.

Higher within-region reliability for intact vs. scrambled stimuli may occur because intact movies and stories contain rich temporal structure that can exploit pre-existing event schemas: Over a lifetime, we have learned, via statistical learning, which events tend to follow each other when, for example, we walk into a hotel or push a button for an elevator. However, statistical learning can also take place much more rapidly, allowing temporal structure to be learned within a relatively short period of time (Aslin & Newport, 2012; Schapiro et al., 2017). Events with novel structure may initially violate expectations because they do not align with pre-existing schemas, but can become predictable after only a handful of repetitions (Turk-Browne et al., 2010). The goal of this study was to explore whether recently learned temporal structure is sufficient to engage the posterior medial network.

Learning of temporal structure by regions in the posterior medial network may not manifest as increases in the consistency of within-region dynamics across stimulus repetitions, however, because a region may become *less* similar to itself over stimulus repetitions if learning occurs. Learning may instead be revealed as changes in between-region functional coupling: For example, the brain's network dynamics change from the first to the second viewing of a movie (Andric et al., 2016). One possibility is that regions within a network may communicate more, or differently, with one another when a stimulus is recognized as potentially significant because it contains temporal structure. We therefore examined both within- and between-region dynamics to look for evidence of learning.

We found that temporal dynamics within posterior medial regions were more reliable for movie clips that were consistent vs. inconsistent with pre-existing schemas (also see Keidel et al., 2017). This was true both within individuals over clip repetitions and between individuals within repetition, replicating and extending prior work with intact vs. scrambled stimuli (e.g., Hasson et al., 2008; Honey et al., 2012a; Lerner et al., 2011). However, over repetitions of a scrambled movie with fixed temporal structure, intra-subject functional coupling (i.e., functional connectivity) between posterior medial regions gradually increased, rising to the level of the intact movie. Such increased coupling may be a signature of learning temporal structure, and might have been missed by prior studies because scrambled stimuli were presented at most twice in the same order, and functional correlations in the posterior medial network were not examined (e.g., Hasson et al., 2008; Honey et al., 2012a). Thus, examining within-region dynamics may not be sufficient to reveal learning: Between-region functional coupling can show increases in network engagement when within-region dynamics suggest no such effect.

Although enhancements in posterior medial network intra-subject functional correlation was strongest for the Scrambled-Fixed clip, a subset of region pairs showed enhanced coupling over repetitions for the Intact and/or Scrambled-Random clips. Such increases may occur because there is some temporal structure to be learned even here. In the Intact clip, information about the specific sequence of events can be learned when the movie is watched several times. That is, while individuals can draw upon event schemas to make sense of what is happening in a general sense for the intact movie, repeated viewings allow learning of movie-specific information, e.g., names of specific characters, and how they move and interact over space and time. In the Scrambled-Random clip, there was also the possibility of extracting some structure over repetitions, as regularities existed on a short timescale *within* each segment that comprised the clip. Nevertheless, the Scrambled-Fixed clip offered the most opportunity for learning: In addition to what could be learned in the Scrambled-Random clip, there were deterministic transitions between clip segments, which were unknown on the first

repetition. Long time-scale temporal structure may therefore be key for engaging the posterior medial network, even if it is inconsistent with pre-existing event schemas.

What is the content of functionally coupled representations in the posterior medial network? That is, as intra-subject functional correlation increases over the course of learning temporally structured but scrambled clips, what do posterior medial regions come to jointly represent? We examined two possibilities — increased stimulus locking and converging idiosyncratic interpretations — but found no evidence for either. That is, posterior medial network dynamics were largely unique to each individual and repetition rather than shared across individuals or repetitions. For example, inter-subject correlation and inter-subject functional correlation, used to isolate stimulus-locked temporal fluctuations, either decreased or were unaffected (Supplementary Figure S1 and S3). Moreover, within-region, intra-subject temporal dynamics were no more similar for clip repetitions at the end vs. beginning of learning, suggesting that individuals did not gradually arrive at a particular mode of processing the movie clips. Thus, network activity was not strictly tied to the stimulus that was commonly viewed by subjects, but was related in part to idiosyncratic cognitive processes for each individual and even for each repetition for a given individual.

What is enhanced functional coupling reflecting then? One possibility is that the network becomes more engaged, but in different ways over repetitions, when a stimulus has been recognized as meaningful, perhaps as a result of detecting temporal structure from prior experience. For example, what is attended may change on each repetition of the clip as a result of prior statistical learning, helping to enable the extraction of new information (Zhao et al., 2013). These switches in attention may manifest as enhanced functional coupling between regions that is dissociable from stimulus-locked correlations. Indeed, such "background" correlations (i.e., task-based functional coupling not attributable to stimuli) have been observed in the visual system when attention switches between objects (Al-Aidroos et al., 2012; Córdova et al., 2016). A similar mechanism may operate in the posterior medial network, allowing us to learn novel, episode-specific details in a familiar event

schema. This could explain how, for example, we can remember unique moments from the last time we went to a restaurant, as opposed to only the moments common to all restaurant visits.

A study from van Kesteren and colleagues (2010) offers further insight about what enhanced functional correlations for temporally structured events could mean. During viewing of an intact movie's conclusion, coupling between the hippocampus and ventromedial prefrontal cortex was higher in individuals who had viewed a scrambled version of the preceding part of the movie, relative to individuals who had viewed the intact version of the preceding part (which provided schematic structure). The authors suggest that functional coupling between hippocampus and ventromedial prefrontal cortex might be enhanced when there is greater difficulty integrating novel information because pre-existing schema are not available. Intriguingly, this was observed alongside higher correlations in inter-subject temporal dynamics in the ventromedial prefrontal cortex for the schema consistent vs. inconsistent group.

We were not able to examine the ventromedial prefrontal cortex in the current study (because of signal dropout, see Figure 6), but this perspective would predict reductions in functional coupling between the hippocampus and ventromedial prefrontal cortex over repetitions of the Scrambled-Fixed clip, as temporal structure is learned and the ease of integrating information increases (i.e., as a new schema potentially begins to form). Such a decrease in functional correlation would serve as a contrast to the increasing posterior medial network coupling found in the current study, and would provide a dissociation among different components of the default mode network (e.g., Andrews-Hanna et al., 2010).

An alternative perspective is that the decrease in inter-subject correlation over repetitions, both within-region (Supplementary Figure S1) and between regions (Supplementary Figure S3) indicates that subjects are gradually becoming disengaged from the movies being viewed. If so, increases in within-subject functional correlations might just indicate greater disengagement of the "default mode" (Raichle, 2015) from the external world, as subjects become fatigued and turn to

internal thoughts rather than processing external stimulus. However, if this were the case, such increased functional correlation should occur for all three clips being viewed and for an equivalent amount, rather than being largest for the Scrambled-Fixed clip. Moreover, if high functional correlation between posterior medial regions indicates disengagement from external stimuli because of fatigue, then it is not clear why such high correlations were observed on the very first exposure to the intact movie clip (Figure 3).

The increased coupling between the hippocampus and other regions in the posterior medial network over repetitions of temporally structured, scrambled clips contributes to the literature on time and sequence coding in hippocampus (e.g., DuBrow and Davachi, 2014, 2016; Hsieh et al., 2014; MacDonald et al., 2011; Mankin et al., 2012; Manns et al., 2007; Pastalkova et al., 2008; see Davachi and DuBrow, 2015; Eichenbaum, 2013; Ranganath and Hsieh, 2015) and studies showing a role for the hippocampus in memory for intact movies (e.g., Chen et al., 2017; Lehn et al., 2009; Gelbard-Sagiv et al., 2008). This is also consistent with studies of statistical learning that have found learning-related changes in hippocampal activity (Bornstein & Daw, 2013; Hindy et al., 2016; Schapiro et al., 2012; Turk-Browne et al., 2009, 2010). The stimuli in these studies were groupings of shapes, objects, faces, and/or scenes, showing that the hippocampus can learn temporal regularities even among arbitrary stimuli that cannot exploit pre-existing schemas. Our results complement this work by showing changes in hippocampal coupling with other brain regions during the learning of naturalistic temporal regularities.

The statistical learning literature emphasizes that a key consequence of extracting temporal structure is the ability to generate accurate predictions (e.g., Schapiro et al., 2012; Turk-Browne et al., 2010). Indeed, the hippocampus exhibits anticipatory signals that can impact goal-directed behavior (e.g., Bornstein and Daw, 2013; Brown et al., 2016; Hindy et al., 2016; Johnson and Redish, 2007; Pfeiffer and Foster, 2013; for review, see Buckner, 2010; Redish, 2016). Given the increased coupling of posterior medial regions with the hippocampus, they may be contributing to hippocampal prediction.

For example, integration of long-timescale information in posterior medial regions may allow the hippocampus to generate predictions about events in the more distant future. Indeed, providing such a temporal history to the hippocampus helps it extract regularities and generate predictions (Schapiro et al., 2017). Another possibility is that predictive signals in the hippocampus are communicated to posterior medial regions to help them with temporal integration.

In conclusion, the posterior medial network is critical for remembering event sequences and integrating information over long timescales (Hasson et al., 2015; Ranganath and Ritchey, 2012). We found that repeated viewing of novel temporal sequences leads to increased functional coupling between the precuneus, hippocampus, angular gyrus, and posterior cingulate cortex. This occurred in the absence of evidence for increased stimulus-locked responses, suggesting idiosyncratic and labile temporal dynamics. These findings highlight how repeated exposure to temporal structure can induce changes in network organization over time, which may be a necessary precursor for learning.

Acknowledgments

This work was funded by NIH grants R01-EY021755 to N.B.T.B. and R01-MH094480 to U.H. This project was also made possible through the support of a grant from the John Templeton Foundation. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of these agencies. The authors have no competing interests to declare.

Conflict of Interest

None

Author Contributions

Mariam Aly: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing – original draft preparation, Writing – review & editing, Visualization, Project administration.

Janice Chen: Conceptualization, Methodology, Investigation, Writing – review & editing, Project administration.

Nicholas Turk-Browne: Conceptualization, Methodology, Resources, Writing – review & editing, Supervision, Funding acquisition

Uri Hasson: Conceptualization, Methodology, Resources, Writing – review & editing, Supervision, Funding acquisition

References

Al-Aidroos N, Said CP, Turk-Browne NB. (2012). Top-down attention switches coupling between low-level and high-level areas of human visual cortex. Proc Natl Acad Sci., 109: 14675-14680.

Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, Buckner RL. (2010). Functional-anatomic fractionation of the brain's default network. Neuron 65: 550-562.

Andric M, Goldin-Meadow S, Small SL, Hasson U. (2016). Repeated movie viewings produce similar local activity patterns but different network configurations. NeuroImage, 142: 613-627.

Aslin RN, Newport EL. (2012). Statistical learning: From acquiring specific items to forming general rules. Curr Dir Psychol Sci 21: 170-176.

Baldassano C, Chen J, Zadbood A, Pillow JW, Hasson U, Norman KA. (2017). Discovering event structure in continuous narrative perception and memory. Neuron 95: 709-721.

Bornstein AM, Daw ND. (2013). Cortical and hippocampal correlates of deliberation during model-based decisions for rewards in humans. PLoS Comp Biol 9, e1003387: 1-19.

Brown TI, Carr VA, LaRocque KF, Favila SE, Gordon AM, Bowles B, Bailenson JN, Wagner AD. (2016). Prospective representation of navigational goals in the human hippocampus. Science 352: 1323-1326.

Buckner RL. (2010). The role of the hippocampus in prediction and imagination. Annu Rev Psychol 61: 27-48.

Chaudhuri R, Knoblauch K, Gariel M-A, Kennedy H, Wang X-J. (2015). A large-scale circuit mechanism for hierarchical dynamical processing in the primate cortex. Neuron 88: 419-431.

Chen J, Honey CJ, Simony E, Arcaro MJ, Norman KA, Hasson U. (2016). Accessing real-life episodic information form minutes versus hours earlier modulates hippocampal and high-order cortical dynamics. Cerebral Cortex 26: 3428-3441.

Chen J, Leong YC, Honey CJ, Yong CH, Norman KA, Hasson U. (2017). Shared memories reveal shared structure in neural activity across individuals. Nat Neurosci 20: 115-125.

Córdova NI, Tompary A. Turk-Browne NB. (2016). Attentional modulation of background connectivity between ventral visual cortex and the medial temporal lobe. Neurobio Learn Mem 134:115-122.

Davachi L, DuBrow S. (2015). How the hippocampus preserves order: The role of prediction and context. Trends Cogn Sci 19: 92-99.

DuBrow S, Davachi L. (2014). Temporal memory is shaped by encoding stability and intervening item reactivation. J Neurosci 34: 13998-14005.

DuBrow S, Davachi L. (2016). Temporal binding within and across events. Neurobiol Learn Mem. http://dx.doi.org/10.1016/j.nlm.2016.07.011

Eichenbaum H. (2013). Memory on time. Trends Cogn Sci 17: 81-88.

Gelbard-Sagiv H, Mukamel R, Harel M, Malach R, Fried I. (2008). Internally generated reactivation of single neurons in human hippocampus during free recall. Science 322: 96-101.

Gilboa A, Marlatte H. (2017). Neurobiology of schemas and schema-mediated memory. Trends Cogn Sci 21: 618-631.

Hasson U, Avidan G, Gelbard H, Vallines I, Harel M, Minshew N, Behrmann M. (2009). Shared and idiosyncratic cortical activation patterns in autism revealed under continuous real-life viewing conditions. Autism Research 2: 220-231.

Hasson U, Chen J, Honey CJ. (2015). Hierarchical process memory: Memory as an integral component of information processing. Trends Cogn Sci 19: 304-313.

Hasson U, Malach R, Heeger DJ. (2010). Reliability of cortical activity during natural stimulation. Trends Cogn Sci 14: 40-48.

Hasson U, Nir Y, Levy I, Fuhrmann G, Malach R. (2004). Intersubject synchronization of cortical activity during natural vision. Science 303:1634-1640.

Hasson U, Yang E, Vallines I, Heeger DJ, Rubin N. (2008). A hierarchy of temporal receptive windows in human cortex. J Neurosci 28: 2539-2550.

Hindy NC, Ng FY, Turk-Browne NB. (2016). Linking pattern completion in the hippocampus to predictive coding in visual cortex. Nat Neurosci. doi: 10.1038/nn.4284

Honey CJ, Thesen T, Donner TH, Silbert LJ, Carlson CE, Devinsky O, Doyle WK, Rubin N, Heeger DJ, Hasson U. (2012a). Slow cortical dynamics and the accumulation of information over long timescales. Neuron 76: 423-434.

Honey CJ, Thompson CR, Lerner Y, Hasson U. (2012b). Not lost in translation: Neural responses shared across languages. J Neurosci 32:15277-15283.

Hsieh L-T, Gruber MJ, Jenkins LJ, Ranganath C. (2014). Hippocampal activity patterns carry information about objects in temporal context. Neuron 81: 1165-1178.

Johnson A, Redish AD. (2007). Neural ensembles in CA3 transiently encode paths forward of the animal at a decision point. J Neurosci 27: 12176-12189.

Keidel JL, Oedekoven CSH, Tut AC, Bird CM. (2017). Multiscale integration of contextual information during a naturalistic task. Cerebral Cortex. https://doi.org/10.1093/cercor/bhx218

Lehn H, Steffenach H-A, van Strien NM, Veltman DJ, Witter MP, Håberg AK. (2009). A specific role of the human hippocampus in recall of temporal sequences. J Neurosci 29: 3475-3484.

Lerner Y, Honey CJ, Silbert LJ, Hasson U. (2011). Topographic mapping of a hierarchy of temporal receptive windows using a narrated story. J Neurosci 31: 2906-2915.

MacDonald CJ, Lepage KQ, Eden UT, Eichenbaum H. (2011). Hippocampal "time cells" bridge the gap in memory for discontiguous events. Neuron 71: 737-749.

Mankin EA, Sparks FT, Slayyeh B, Sutherland RJ, Leutgeb S, Leutgeb JK. (2012). Neuronal code for extended time in the hippocampus. Proc Natl Acad Sci 109: 19462-19467.

Manns JR, Howard MW, Eichenbaum H. (2007). Gradual changes in hippocampal activity support remembering the order of events. Neuron 56: 530-540.

Murray JD, Bernacchia A, Freedman DJ, Romo R, Wallis JD et al. (2014). A hierarchy of intrinsic timescales across primate cortex. Nature Neuroscience 17: 1661-1663.

Pastalkova E, Itskov V, Amarasingham A, Buzsáki G. (2008). Internally generated cell assembly sequences in the rat hippocampus. Science 321: 1322-1327.

Pfeiffer BE, Foster DJ. (2013). Hippocampal place-cell sequences depict future paths to remembered goals. Nature 497: 74-79.

Raichle ME. (2015). The brain's default mode network. Annu Rev Neurosci 38: 433-447.

Ranganath C, Ritchey M. (2012). Two cortical systems for memory-guided behavior. Nat Rev Neurosci, 13: 713-726.

Redish AD. (2016). Vicarious trial and error. Nat Rev Neurosci 17: 147-159.

Schapiro AC, Kustner LV, Turk-Browne NB. (2012). Shaping of object representations in the human medial temporal lobe based on temporal regularities. Curr Biol 22: 1622-1627.

Schapiro AC, Turk-Browne NB, Botvinick MM, Norman KA. (2017). Complementary learning systems within the hippocampus: A neural network modeling approach to reconciling episodic memory with statistical learning. Phil Trans R Soc B. 372: 20160049.

Simony E, Honey CJ, Chen J, Lositsky O, Yeshurun Y, Wiesel A, Hasson U. (2016). Dynamic reconfiguration of the default mode network during narrative comprehension. Nat Communications 7:12141, p.1-13.

Turk-Browne NB, Scholl BJ, Chun MM, Johnson MK. (2009). Neural evidence of statistical learning: Efficient detection of visual regularities without awareness. J Cogn Neurosci 21: 1934-1945.

Turk-Browne NB, Scholl BJ, Johnson MK, Chun MM. (2010). Implicit perceptual anticipation triggered by statistical learning. J Neurosci 30: 11177-11187.

Utevsky AV, Smith DV, Huettel SA. (2014). Precuneus is a functional core of the default-mode network. J Neurosci 34: 932-940.

van Kesteren MTR, Fernández G, Norris DG, Hermans EJ. (2010). Persistent schema-dependent hippocampal-neocortical connectivity during memory encoding and postencoding rest in humans. Proc Natl Acad Sci 107: 7550-7555.

van Kesteren MTR, Ruiter DJ, Fernández G, Henson RN. (2012). How schema and novelty augment memory formation. Trends Neurosci 35: 211-219.

Zhao J, Al-Aidroos N, Turk-Browne NB. (2014). Attention is spontaneously biased toward regularities. Psych Sci 24: 667-677.

Figure S1. Inter-subject temporal dynamics across clip repetitions. (A) Response reliability was assessed by correlating, for the first and last repetition separately, each subject's timecourse of BOLD activity in the precuneus with the mean precuneus timecourse across all other subjects (i.e., inter-subject correlation). **(B)** Statistical analyses for repetition 1 are reported in the main text; the data are shown here again for comparison to repetition 6. Inter-subject correlation was higher for repetition 1 vs. repetition 6 of the Intact clip ($r_{Rep1} = 0.58$, $r_{Rep6} = 0.26$; 95% bootstrapped CI: 0.30–0.51; $p = 0$ from permutation test). Inter-subject correlation was also higher for repetition 1 vs. repetition 6 of the Scrambled-Random clip ($r_{Rep1} = 0.36$, $r_{Rep6} = 0.09$; 95% bootstrapped CI: 0.13–0.45; $p = .002$ from permutation test). In contrast, repetition 1 and repetition 6 were not different from one another for the Scrambled-Fixed clip ($r_{Rep1} = 0.24$, $r_{Rep6} = 0.19$; 95% bootstrapped CI: -0.07–0.21; $p = .36$ from permutation test). For repetition 6, inter-subject correlation was reliably above zero for all three conditions (Intact: $r_{Rep6} = 0.26$; 95% bootstrapped CI: 0.18–0.38; $p = 0$ from permutation test; Scrambled-Fixed: $r_{Rep6} = 0.19$; 95% bootstrapped CI: 0.12–0.28; $p = 0$ from permutation test; Scrambled-Random: $r_{Rep6} = 0.09$; 95% bootstrapped CI: 0.006–0.18; $p = .04$ from permutation test). Correlations were higher for the Intact clip vs. the Scrambled-Random clip on repetition 6 (95% bootstrapped CI: 0.04–0.32; $p = .02$ from permutation test). The Intact and Scrambled-Fixed clips were not different from one another on repetition 6 (95% bootstrapped CI: -0.04–0.20; $p = .22$ from permutation test), and neither were the Scrambled-Fixed and Scrambled-Random clips (95% bootstrapped CI: -0.008–0.21; $p = .12$ from permutation test). Dots indicate individual subjects. Error bars are ± 1 SEM. * $p < .05$, ** $p < .01$, *** $p < .001$.

Figure S2. Intra-subject functional correlation across all six repetitions of each clip. (A) Intra-subject functional correlation between precuneus and hippocampus increased monotonically across all six repetitions of the Scrambled-Fixed clip. ($\beta = 0.05$ $t_{29} = 4.06$, $p = .0003$, 95% CI: 0.02–0.07). The slope was not different from zero for the Intact clip ($\beta = 0.004$, $t_{29} = 0.43$, $p = .67$, 95% CI: -0.02–0.02) and was numerically smaller for the Scrambled-Random clip ($\beta = 0.03$, $t_{29} = 2.17$, $p = .04$, 95% CI: 0.002–0.06). **(B)** Intra-subject functional correlation between precuneus and angular gyrus increased monotonically across all six repetitions of the Scrambled-Fixed clip ($\beta = 0.07$, $t_{29} = 4.68$, $p = .00006$, 95% CI: 0.04–0.09). There were also numerically smaller increases for the Intact ($\beta = 0.04$, $t_{29} = 3.43$, $p = .002$, 95% CI: 0.02–0.06) and Scrambled-Random clips ($\beta = 0.04$, $t_{29} = 2.42$, $p = .02$, 95% CI: 0.007–0.08). **(C)** Intra-subject functional correlation between precuneus and posterior cingulate cortex increased monotonically for the Scrambled-Fixed clip ($\beta = 0.06$, $t_{29} = 4.76$, $p = .00005$, 95% CI: 0.03–0.08). There was a numerically smaller increase for the Scrambled-Random clip ($\beta = 0.04$, $t_{29} = 2.48$, $p = .02$, 95% CI: 0.007–0.08). The slope for the Intact clip was not different from zero ($\beta = 0.02$, $t_{29} = 1.58$, $p = .12$, 95% CI: -0.006–0.05). Dots indicate individual subjects. Error bars are ± 1 SEM. * $p < .05$, ** $p < .01$, *** $p < .001$.

Figure S3. Inter-subject functional correlation. (A) The timecourse of BOLD activity in the precuneus for a given subject was correlated with the mean timecourse of BOLD activity in the hippocampus for all other subjects. The timecourse of BOLD activity in the hippocampus for that same subject was also correlated with the mean timecourse of BOLD activity in the precuneus for all subjects. The average of these correlations was the inter-subject functional correlation value for the hippocampus and the precuneus for that subject. Inter-subject functional correlation decreased from repetition 1 to repetition 6 of the Intact clip (r_{Rep1} = 0.31, r_{Rep6} = 0.07, p =0 from permutation test, 95% bootstrapped CI: 0.19–0.30) and the Scrambled-Fixed clip (r_{Rep1} = 0.11, r_{Rep6} = 0.03, p = .04 from permutation test, 95% bootstrapped CI: 0.007–0.16), and did not change for the Scrambled-Random clip (r_{Rep1} = 0.03, r_{Rep6} = 0.001, p = .49 from permutation test, 95% bootstrapped CI: -0.06–0.11). **(B)** Inter-subject functional correlation between the precuneus and angular gyrus did not change over repetitions in any condition (Intact: r_{Rep1} = 0.23, r_{Rep6} = 0.14, p = .09 from permutation test, 95% bootstrapped CI: -0.006–0.20; Scrambled-Fixed: r_{Rep1} = 0.03, r_{Rep6} = 0.09, p = .16 from permutation test, 95% bootstrapped CI: -0.15–0.02; Scrambled-Random: r_{Rep1} = 0.04, r_{Rep6} = -0.04, p = .15 from permutation test, 95% bootstrapped CI: -0.008–0.17). **(C)** Inter-subject functional correlation between the precuneus and posterior cingulate cortex decreased over repetitions for the Intact clip (r_{Rep1} = 0.37, r_{Rep6} = 0.12, p = 0 from permutation test, 95% bootstrapped CI: 0.16–0.36), but did not change over repetitions of the Scrambled-Fixed clip (r_{Rep1} = 0.06, r_{Rep6} = 0.07, p = .77 from permutation test, 95% bootstrapped CI: -0.09–0.07) or the Scrambled-Random clip (r_{Rep1} = -0.0005, r_{Rep6} = -0.07, p = .10 from permutation test, 95% bootstrapped CI: -0.01–0.14).