Title: Carbon pool trends and dynamics within a subtropical peatland during long-term restoration.

Paul Julian II[1], Stefan Gerber[2], Alan L. Wright[1], Binhe Gu[3] and Todd Z. Osborne[2,4]

[1] University of Florida, Soil and Water Sciences Department, Ft. Pierce, FL 34945
*Corresponding Author: pjulian@ufl.edu; ORCID: 0000-0002-7617-1354
[2] University of Florida, Soil and Water Sciences Department, Gainesville, FL, 32611

[3] South Florida Water Management District, Water Quality Bureau, West Palm Beach, FL, 33406

[4] University of Florida, Whitney Laboratory for Marine Bioscience, St Augustine, FL 32080

Abstract

Background: The Florida Everglades has undergone significant ecological change spanning the continuum of disturbance to restoration. While the restoration effort is not complete and the ecosystem continues to experience short duration perturbations, a better understanding of long-term C dynamics of the Everglades is needed to facilitate new restoration efforts. This study evaluated temporal trends of different aquatic carbon (C) pools of the northern Everglades Protection Area over a 20-year period to gauge historic C cycling patterns. Dissolved inorganic C (DIC), dissolved organic C (DOC), particulate organic C (POC), and surface water carbon dioxide ($pCO_{2(aq)}$) were investigated between May 1, 1994 and April 30, 2015.

Results: Annual mean concentrations of DIC, DOC, POC, and $pCO_{2(aq)}$ significantly decreased through time or remained constant across the Water Conservation Areas (WCAs). Overall, the magnitude of the different C pools in the water column significantly differed between regions. Outgassing of CO_2 was dynamic across the Everglades ranging from 420 to 2001 kg CO_2 yr-1. Overall the historic trend in CO_2 flux from the marsh declined across our study area while $pCO_{2(aq)}$ largely remained somewhat constant with the exception of Water Conservation Area 2 which experienced significant declines in $pCO_{2(aq)}$. Particulate OC concentrations were consistent between WCAs, but a significantly decreasing trend in annual POC concentrations were observed.

Conclusions: Hydrologic condition and nutrient inputs significantly influenced the balance, speciation, and flux of C pools across WCAs suggesting a subsidy-stress response in C dynamics relative to landscape scale responses in nutrient availability. The interplay between nutrient inputs and hydrologic condition exert a driving force on the balance between DIC and DOC production via the metabolism of organic matter which forms the base of the aquatic foodweb. Along the restoration trajectory as water quality and hydrology continues to improve it is expected that C pools will respond accordingly.

Keywords: Inorganic carbon, organic carbon, carbon dioxide, particulate carbon, hydrology, nutrients

Introduction

Wetlands, including peatlands, are net C sinks that store a large amount of the global C created by an unbalanced accumulation of C via plant productivity and export from decomposition of organic matter via C dioxide (CO_2) or methane (CH_4) to the atmosphere. Carbon forms such as dissolved inorganic C (DIC), particulate organic C (POC) and DOC can be transported laterally through run-off (Updegraff et al. 1995; Freeman et al. 2004; Billett and Moore 2008), and contribute to wetlands' carbon budgets. This type of export C, more specifically organic species from wetlands, represents a significant regional redistribution of terrestrial C and exerts important controls on aquatic productivity in downstream waterbodies. Dissolved organic and inorganic C may thereby enter a feedback loop with aquatic productivity within a wetland (Carpenter and Pace 1997; Hanson et al. 2003).

Net aquatic production (NAP), or total metabolic balance of an aquatic ecosystem, is the difference between gross primary production and ecosystem respiration in the water column. As a result of maintaining a metabolic balance, different "pools" of aquatic carbon are influenced by ecosystem processes related to ecosystem metabolism and function. Aquatic ecosystems with low nutrients such as total phosphorus (TP) and high DOC from either allochthonous (external) or autochthonous (internal) sources tend to be net heterotrophic, while those exhibiting low DOC and high TP tend towards net autotrophy (Hanson et al. 2003; Waiser and Robarts 2004). Therefore, the supply of various C pools and other water quality indicators are important in understanding ecosystem function.

Most wetlands and lake ecosystems are heterotrophic with respect to aquatic metabolism with

surface waters acting as net sources of CO_2 to the atmosphere by decomposition of organic

matter (Kling et al. 1991; Cole et al. 1994; Waiser and Robarts 2004). In contrast, some coastal

wetlands and a small number of lakes are considered autotrophic (Kling et al. 1992; Cole et al.

1994; Duarte and Cebrian 1996; Gu et al. 2011; Hopkinson et al. 2012). The net C source in

wetland and littoral zones could be sustained through C fixation by emergent vegetation and its

subsequent decomposition. Thus, while the surface water may be heterotrophic, the wetland

ecosystem as a whole (water + soil), can still function as a net C sinks resulting in overall C

accretion.

The major sources of total dissolved CO_2 contributing to aquatic systems are (1) diffusion of

atmospheric CO_2 from the atmosphere, (2) hydrologic transport from upstream, (3) groundwater

discharge, (4) CO_2 derived from organic matter decomposition within the water column and the

soil and (5) the dissolution of C minerals (Telmer and Veizer 1999; Finlay 2003). In some river

and small stream systems, metabolism of C inputs appears to be greater than *in-situ* primary

production resulting in higher aqueous dissolved CO_2 than CO_2 in the atmosphere (Cole et al.

1994). A total metabolic balance is driven by factors that govern both primary production and

respiration. Productivity can be influenced by nutrients, light availability, thermal cycle,

hydrodynamics (i.e. current, velocity, mixing, etc.), internal loading of organic matter and food

web structure. Meanwhile ecosystem respiration may be strongly subsidized by external loading

of organic matter and nutrients (Odum 1956; Kling et al. 1991; Wetzel 1992; Cole et al. 2000;

Hessen et al. 2004; Elser et al. 2007; Gu et al. 2011). Therefore, the combination and interaction

of these factors influence the partitioning and conversion of C into distinct pools.

The objectives of this study were to evaluate different aquatic C pools within the Everglades

ecosystem. The first objective of this study was to evaluate long-term regional trends in aquatic

C (DIC, DOC, POC and pCO$_2$) within the Everglades as large scale restoration efforts progress

to restore quality, quantity, and timing of water to the Everglades ecosystem. The first hypothesis

is as the system recovers from degraded water quality impacts, C pools will decline or level out.

The decrease may be directly caused by reduced nutrient availability to primary producers and

possibly modified by altered (increasing or decreasing) organic matter turnover in the water

column and soils. The second objective is to estimate CO$_2$ flux from the Everglades Protection

Area (EPA) marsh to the atmosphere and investigate long-term trends, with the hypotheses that

aquatic CO$_2$ fluxes will also decline concurrently with the decline in other C pools as a response

to restoration.

Methods

Study Area

The Everglades ecosystem is a complex system of marshes, canals and levees with water control

structures covering approximately 10,000 km^2 of former contiguous Everglades marsh and

currently divided into large, distinct shallow impoundments (Bancroft et al. 1992; Light and

Dineen 1994). Surface water is delivered primarily from the north and west to the EPA through

water control structures connecting the Everglades Agricultural Area (EAA), stormwater

treatment areas (STAs) and the C-139 basin and through urban areas along the eastern edge of

the EPA. Surface water from these land areas are delivered to the EPA through water control

structures, but the timing and distribution of the surface water inflows from the upstream

watersheds to the EPA are based on complex series of operational decisions accounting for

natural and environmental system requirements, water supply for urbanized and natural areas,

aquifer recharge, and flood control (Julian et al. 2016).

This study was centered on the northern third of the EPA, known as the water conservation areas

(WCAs; Fig 1) that receive the majority of surface water inputs through the highly managed

surface water system via a combination of canals and water control structures. Water

Conservation Area 1 is unique in the distinct point source inputs of surface water that moves

around the marsh edge via the perimeter canal. As a result of this limited interaction of the

WCA-1 marsh interior with mineral-rich canal drainage water, WCA-1 is the sole remaining

soft-water ecosystem in the Everglades and predominately rainfall driven hydrology in the marsh

interior (Newman and Hagerthey 2011). Both WCA-2 and WCA-3 surface hydrology is

controlled by a system of levees and water control structures along the perimeter. Additionally

canals, levees and water control structures bisect WCA-3 effectively dividing the area into four

hydrologically distinct areas (DeBusk et al. 2001; Bruland et al. 2006). Soil within the WCAs are

Histosols, and encompass both Loxahatchee and Everglades peat formations with depths ranging

from ~1 to 2 m with a mosaic of aquatic sloughs, expanses of wet prairie, strands of sawgrass

(*Cladium jamaicense* Crantz), and patches of brush and tree islands (Gleason et al. 1974; Brandt

et al. 2000; DeBusk et al. 2001; Bruland et al. 2006)

Source of Data

Water quality data was retrieved from the South Florida Water Management District (SFWMD)

online database (DBHYDRO; www.sfwmd.gov/dbhydro) for sites within the WCAs identified in

Fig. 1 between water year 1995 to 2015 (May 1, 1994 – April 30, 2015). Only stations with

greater than four samples per year and three years of monitoring data were considered for this

analysis resulting in 44 monitoring stations across the EPA. Surface water grab samples were

typically collected between 06:00 to 17:00 local time, with most samples collected at

approximately 09:00. Water quality parameters used include alkalinity, pH, temperature, specific

conductivity, DOC, total phosphorus (TP) and total nitrogen (TN). In order to quantify air-water

net CO_2 exchange between surface waters and the atmosphere additional data on wind speed and

atmospheric CO_2 partial pressure was obtained. Wind speed data measured 10 meters above

ground elevation was retrieved from metrological stations within the EPA identified by Fig. 1.

Monthly atmospheric partial pressure carbon dioxide ($pCO_{2(atm)}$) data was retrieved from the

National Oceanic & Atmospheric Administration (NOAA) online global monitoring division

webpage (www.esrl.noaa.gov/gmd) using the Key Biscayne monitoring location (25.6654° N,

80.1580° W; NAD1983). Stage elevation data was retrieved from the SFWMD DBHYDRO for

stage monitoring site identified in Fig. 1 between water year (WY) 1979 to 2015 (May 1, 1978 –

April 30, 2015).

Water quality data were screened based on laboratory qualifier codes, consistent with FDEP's

quality assurance rule (Florida Administrative Code 2008). Any datum associated with a fatal

qualifier indicating a potential data quality problem were removed from the analysis. For

purposes of data analysis and summary statistics, data reported as less than method detection

limit (MDL) were assigned a value of one-half the MDL, unless otherwise noted.

Data Analysis

Dissolved inorganic C concentrations were calculated from the relationship between water

column alkalinity, pH and temperature as outlined by Wetzel and Likens (2000). This

methodology of quantifying DIC concentrations is consistent with previous studies where they

determined that the calculated DIC concentration based on alkalinity and relevant parameters can

be used to estimate DIC concentration in the absence of direct measure DIC concentration (Gu et

al. 2008, 2011). Particulate organic C concentration were calculated from the difference between

total organic C (TOC) and DOC. The concentration of dissolved free CO_2 within the water

column were calculated using the pH and CO_2 fraction relationship presented in (Wetzel and

Likens 2000)(Table 1, equation 1). Surface water pCO_2 was calculated using Henry's Law

(Table 1, equation 2) where K_H is the dissolution constant of CO_2 corrected for water

temperature (Table 1, equation 3). Atmospheric concentration of CO_2 [$CO_{2(atm)}$] above the

stagnant layer were estimated from Henry's Law using monthly atmospheric CO_2 partial

pressure from the Key Biscayne NOAA monitoring location (Table 1, equation 4). Due to the

limited availability of wind data CO_2 flux calculations were limited to a nine-year period

(WY1999 - 2008). During the nine-year period the WCAs experienced changes in climate (i.e.

drought and flood), water quality and system operations due to the construction and operation of

the Everglades Stormwater Treatment Areas. Therefore, to extend the period of record of flux

calculations comparisons of calculations with differences in wind speed data were performed

(Appendix 1). As a result of this comparison, a period of record mean wind speed value of 2.87

m s^{-1} was substituted for wind speed in the flux calculations to estimate the flux of CO_2 between

the atmosphere and surface water, equation 5 (Table 1) was used which incorporated CO_2

diffusion coefficient (Table 1, equation 6) and surface boundary layer thickness (Table 1,

equation 7). For sites with dense macrophyte coverage, it was assumed that the emergent

macrophytes would reduce wind speed at the air-water interface to effectively zero similar to

(Hagerthey et al. 2010). Water column DIC concentrations and surface water pCO_2 ($pCO_{2(aq)}$)

were calculated for all station with the necessary data between WY1995 to WY2015.

Regional comparison was conducted using annual mean DIC, DOC, POC and $pCO_{2(aq)}$

concentrations for each WCA based on the Florida water year (May-April). Station specific trend

analysis was using Kendall's τ correlation analysis and Thiel-Sen's slope estimate (zyp R-

package; (Bronaugh and Werner 2013). Annual mean DIC, DOC, POC, $pCO_{2(aq)}$ and CO_2 flux

were compared between WCAs using the Kruskal-Wallis rank sum test and Dunn's test of

multiple comparisons (Dunn's test R-package; (Dinno 2015) using stations with greater than six

samples collected in a given WY. Annual mean DOC and DIC ratio were computed using mass

per volume concentrations and compared between WCAs using the Kruskal-Wallis rank sum test

and Dunn's test of multiple comparisons. Unless otherwise noted all statistical operations were

performed using the base stats R-package. All statistical operations were performed with R©

(Ver 3.1.2, R Foundation for Statistical Computing, Vienna Austria) at a critical level of

significance of α = 0.05.

Results

During this study, calculated DIC concentrations ranged from 1.9 mg C L^{-1} to 144.0 mg C L^{-1}

across the WCAs. Statistically significantly declining annual DIC trends were observed at a total

of 9 stations during this study, no trend was apparent for 32 of the stations during this study, the

remaining didn't have sufficient data to conduct a trend analysis (Fig 2). Most notable

significantly decreasing trends were observed at stations along the eutrophication gradient with

WCA-2 (DeBusk et al. 2001). Regional annual mean DIC concentrations between WY1995 and

2015 significantly declined in WCA-2 at a rate of – 0.53 mg L^{-1} yr^{-1}, no temporal trend was

apparent in WCA-1 and WCA-3 (Table 2). Dissolved inorganic C concentration significantly

differed between WCAs ($χ^2$=54.8, df=2, ρ<0.001) with DIC concentration being lowest within

WCA-1 (17.2 ± 0.6 mg C L^{-1}) and greatest in WCA-2 (67.1 ± 0.7 mg C L^{-1}) (Fig. 3). Monthly

mean DIC and stage elevation was significantly negatively correlated for WCA-3; negatively

correlated with mean monthly water temperature for WCA-2 and WCA-3; positively correlated

with mean monthly TP concentration for WCA-1 and WCA-2; TN and specific conductance for

all regions of the study area (Table 4).

Dissolved organic C concentrations ranged from 0.5 mg C L^{-1} to 65.9 mg C L^{-1} across WCAs.

Significantly declining trend were observed at the individual site scale, with a total of 9 sites

with significantly declining trends, meanwhile no trend was apparent for 33 of the stations in this

study (Fig 2). Regional annual mean DOC concentrations significantly declined in WCA-2 at a

rate of $- 0.37$ mg L^{-1} yr^{-1}, no temporal trend was apparent in WCA-1 and WCA-3 (Table 2).

Annual mean DOC concentrations significantly differed between WCAs (Fig. 4, χ^2=41.1 df=2,

ρ<0.01), with WCA-1 and WCA-3 regional concentrations being similar (19.4 ± 0.2 mg C L^{-1}

and 22.1 ± 0.2 mg C L^{-1}, respectively) and WCA-2 being different between WCA-1 and WCA-3

with a higher observed regional average DOC concentration (35.8 ± 0.3 mg C L^{-1}) (Fig. 3).

Monthly mean DOC concentrations were negatively correlated with mean monthly stage

elevations forWCA-2 and WCA-3; no correlated with mean monthly surface water temperature;

positively correlated with TP for WCA-2; and positively correlated for TN and specific

conductivity for all regions (Table 4). Annual mean DOC:DIC values significantly differed

between WCAs (Fig. 5, χ^2=52.8, df=2, ρ<0.01) with WCA-1 having the highest observed ratio

of 1.95 ± 0.07 followed by WCA-2 (0.55 ± 0.004) and WCA-3 (0.46 ± 0.003).

Particulate organic C concentrations ranged from 0 to 40.3 mg C L^{-1} across the WCAs.

Significantly declining trends in annual mean POC concentrations were observed at 24 stations,

meanwhile no trend was apparent for 9 of the stations in this study (Fig 3). Regionally annual

mean POC concentrations across all three WCA's significantly declined at a rate of -0.26 to -

0.008 mg C L^{-1} yr^{-1} (Table 2). Annual mean POC concentrations did not significantly differ

across the WCAs (Fig 4, χ^2=4.36, df=2, ρ=0.11), however the multiple comparison test

determined a significant difference of POC concentration between WCA-1 and WCA-2 (z-score

= -2.06, ρ<0.05). Regional average concentrations ranging from 0.7 ± 0.04 mg C L^{-1} (WCA-1) to

1.6 ± 0.20 mg C L^{-1} (WCA-2) with all regions experiencing significantly declining trends (Table

2). Mean monthly POC concentrations were only correlated with monthly mean TP and specific

conductivity for WCA-2 and WCA-3 (Table 4).

Calculated pCO$_{2(aq)}$ concentrations ranged from 135.0 µatm to 2.1 atm (2.1 x 10^6 µatm) across

the WCAs. Significantly declining trends in annual mean pCO$_{2(aq)}$ were observed at 7 stations,

while no trend was apparent at 34 stations during the course of this study (Fig 2). At several

stations, concurrent declines in pCO$_{2(aq)}$, DIC, DOC and POC were observed potentially

indicating a change in C dynamics at these specific locations within the Everglades system.

Regionally annual mean pCO$_{2(aq)}$ significantly declined across WCA-2 at a rate of -792 µatm

yr^{-1} (Table 2) and no significant temporal trend was apparent for WCA-1 and WCA-3 (Table 2).

Annual mean pCO$_{2(aq)}$ significantly differed between WCAs (Fig 4, χ^2=13.4, df=2, ρ<0.01) with

WCA-2 and WCA-3 being statistically similar (z-score=0.67, ρ=0.25; Fig 4). Annual mean

pCO$_{2(aq)}$ concentrations followed a decreasing north-to-south trend with WCA-1 having the

greatest concentration (52,379 ± 6,445 µatm) and WCA-3 having the lowest concentration

(24,963 ± 983 µatm). Mean monthly pCO$_{2(aq)}$ were positively correlated with surface water

temperature within WCA-2 and positively correlated with TP for WCA-1 and WCA-2 (Table 4).

The calculated water-air CO_2 flux ranged from -6.2 to 36,361 mg m^{-2} d^{-1} across the WCAs.

Annual mean regional flux range from 285 to 876 mg m^{-2} d^{-1}. Annual mean CO_2 flux

significantly declined during the course of the study for all regions (Table 2). Similar to $pCO_{2(aq)}$

regional trends, annual mean flux was significantly different between WCAs (χ^2=26.2, df=2,

ρ<0.01) with annual mean flux rates being significantly different between WCA-1 and WCA-2

and WCA-3 and similar between WCA-2 and WCA-3 (z-score=0.008, ρ=0.50). Flux rates were

greatest for WCA-1, followed by WCA-2 and WCA-3 (Table 3). Using the area of each region

(WCA-1: 567 km^2; WCA-2: 537 km^2 and' WCA-3: 2,368 km^2) and the annual mean daily CO_2

flux rate extrapolated to an annual estimate for each region an estimated CO_2 flux to the

atmosphere is 1,261 $\pm$ 459 kg CO_2 yr^{-1} during this study, with the expectation that mass transport

from the marsh will decreases as indicated by the annual CO_2 flux trends.

Monthly $pCO_{2(atm)}$ concentration ranged from 354 µatm to 404 µatm at the Key Biscayne

monitoring location between WY1995 to WY2015 with a mean $\pm$ standard error of 380 $\pm$ 0.8

µatm during this period. During this period, WY mean $pCO_{2(atm)}$ concentrations significantly

increased (τ=1.00, ρ<0.001) during the study at a rate of 2.0 µatm year^{-1}.

Discussion

Dissolved Inorganic Carbon and pCO$_{2(aq)}$

Dissolved inorganic C concentrations were consistent with those reported by (Gu et al. 2008)

who investigated the role of fire on C balance within WCA-2. Across the entire period of record

DIC concentrations gradually decreased both at the regional (Fig 3 and Table 2) and individual

station scale (Fig 2) with concurrent changes in water quality systems wide (Julian et al. 2016).

Dissolved inorganic C is an essential source of C to benthic macrophytes and other autotrophic

species (Raven et al. 1982; Sand-Jensen and Frost-Christensen 1998). Changes in water column

nutrient concentrations, DIC concentrations, and water quantity influence wetland productivity

which in turn influences biomass turnover, C demand in the aquatic ecosystems, and ecosystem

function (Findlay et al. 2002; Bossio et al. 2006; Corstanje et al. 2007).

Each WCA has unique biogeochemical properties and hydrologic dynamics therefore DIC

concentrations and C dynamics observed during this study were expected to vary between WCAs

(Fig. 4). The interior portions of the WCA-1 marsh are hydrologically dominated by rainfall with

very little surface water flows penetrating beyond the outer edge of the marsh (Harvey and

McCormick 2009). This hydrologic setting results in low water column pH, low alkalinity and

oligotrophic conditions with respect to TP (Julian et al. 2016). As such, these conditions allow

for greater flux of CO_2 from the marsh within WCA-1 relative to the other portions of the EPA

presumably due to a combination of organic matter decomposition and dissolution of calcium

carbonate driven by low pH conditions. Meanwhile, a different set of drivers are present for

WCA-2 which historically received large quantities of storm water run-off from the EAA

resulting in large areas considered to be eutrophic (DeBusk et al. 2001). However, hydrologic

restorations efforts and the construction and operation of the Everglades STAs have reduced

storm water inputs and TP into WCA-2 (Julian 2015). Unlike the other two WCAs, WCA-3 is

hydrologically and physically compartmentalized resulting in a mosaic of areas with drastically

different hydroperiods (north versus south), nutrient inputs and cycling patterns (Reddy et al.

1998; Bruland et al. 2006). The strength of the groundwater connection within WCAs is variable

as is the thickness of peat/soil and depth to the lime rock bedrock across the landscape (Scheidt

and Kalla 2007). As in stream ecosystems, wetland DIC is derived from several sources

including the dissolution of carbonate, respiration by aquatic plants and heterotrophs by the

consumption of organic matter, shallow groundwater inputs from elevated levels of soil CO_2 and

atmospheric draw-down (Wetzel 1992; Palmer et al. 2001). In some stream ecosystems it has

been observed that DIC is predominately supplied and controlled by drainage of CO_2-rich

shallow groundwater into the water column (Palmer et al. 2001; Finlay 2003). In wetlands

respiration by aquatic plants and heterotrophs is the dominate source and potential control of

DIC concentrations (Richey et al. 2002; Hagerthey et al. 2010). This DIC source and control

pathway has also been reported for estuarine environments (Raymond et al. 2000). Therefore, in

the Everglades, respiration during the metabolism of organic matter is the primary source of DIC

in surface waters, with organic matter, nutrient gradients and DOC concentrations controlling C

turnover as indicated by the vary large pCO$_{2\text{ (aq)}}$ concentrations observed (Fig. 3).

(Odum et al. 1979) hypothesized that in wetlands, plant productivity will be greatest when

periodic short duration flooding provides subsidies of nutrients and water. Alternatively,

prolonged flooding will cause physiological stress to the plants and limit nutrient subsidies thus

limiting productivity. Using this subsidy-stress conceptual model, WCA-2 had a larger subsidy

of water as indicated by longer hydroperiods and deeper water depths in combination with the

highest concentration of available nutrients (DeBusk et al. 2001; Julian et al. 2016). Increased

nutrient concentrations accelerate accretion of organic matter within the marshes by stimulation

of net primary production and organic matter decomposition via microbial metabolism which in

turn contributed to elevated DIC production, utilization and turnover (Reddy et al. 1993; Qualls

and Richardson 2000; Fisher and Reddy 2010). Furthermore, the positive correlation between

TN, specific conductance and DIC concentration (Table 4) could provide further evidence of

nutrient subsidy stimulated DIC production. Specific conductivity has been used as a tracer of

surface water with higher conductivity water representing higher available nutrient canal water

penetrating portions of the marsh while lower specific conductivity water represents interior

marsh surface water with relatively low nutrient concentrations (Harwell et al. 2008; Surratt et al.

2008). Alternatively, the correlation between TN and DIC could also be linked to the

productivity of periphyton and blue-green algae which fix atmospheric nitrogen during

metabolism. Regardless, there is an apparent hydrologic factor involved with DIC production

and uses as suggested by the wetland subsidy-stress model of (Odum et al. 1979) in that WCA-1

is hydrologically isolated and WCA-3 is hydrologically fractured (i.e. compartmentalized),

which in part regulates ecosystem level productivity, DIC production and carbon turnover (Table

4).

Dissolved Organic Carbon and Particulate Organic Carbon

Dissolved organic C concentrations during this study were consistent with concentrations

observed in other studies within the Everglades system (Qualls and Richardson 2003; Aiken et

al. 2011; Lu et al. 2014). Wetlands are major sinks of C that sequester $0.003 - 2.2$ kg C m^{-2} yr^{-1},

with DOC typically representing <1% of the TOC in soil (i.e. bulk soil + porewater), but ~90%

of the TOC in surface water (Kadlec and Wallace 2009; Kayranli et al. 2010). Particulate organic

C concentrations observed during this study corroborate the consensus that DOC is the dominate

form of organic C in the water column, typically an order of magnitude greater than POC

concentrations (Fig 4, Table 3). Furthermore, suspended particulate concentrations in the interior

portions of the EPA are relatively low (Julian et al. 2016), providing more evidence which

suggests that most of the organic C is in the dissolved fraction (Lu et al. 2003).

The size of the DOC pool is orders of magnitude larger than the POC pool within the northern

portion of the EPA. Nutrients can be a controlling factor in the production of organic C via

aquatic plant and algae production, and decomposition. Surface water nutrient concentrations

were positively correlated with both DOC and POC (Table 4), suggesting higher primary

productivity in the water column and rates of organic C leaching and litterfall decomposition in

areas of higher nutrients occurs within the Everglades system which correspond to result

presented by previous studies (DeBusk and Reddy 1998; D'Angelo and Reddy 1999; Stern et al.

2007). Additionally, marsh TP concentrations in the WCAs have significantly decreased since

WY1979 with WCA-2 having the largest period of record trend decrease (WY1979-WY2015; -

0.48 μg L^{-1} yr^{-1}) in concentrations (Julian et al. 2016). The decrease in nutrient concentrations

also corresponded with significant decreases in DOC, POC, DIC and pCO$_2$ (this study; Table 2)

with WCA-2 observing the largest concurrent decreases across the EPA. Availability of nutrients

and electron acceptors plays an important role in wetland C cycling with available nutrients

limiting the growth of microbial fauna (Drake et al. 1996; D'Angelo and Reddy 1999; Wright

and Reddy 2001).

While nutrients play an important role in DIC production and C turnover, hydrologic conditions

also play a significant role in the cycling and production of organic C within wetlands. Dissolved

organic C was negatively correlated with stage elevation, suggesting a possible dilution of DOC

within marshes (Table 4). (Lu et al. 2003) observed a similar correlation with DOC in the

southern portion of ENP, suggesting that rainfall diluted DOC concentrations as water levels

increases along flow transects within the Everglades system. This process has also been

demonstrated in other forested and wetland ecosystems (Fraser et al. 2001; Kawasaki et al. 2005;

Ågren et al. 2007). Some studies have attributed this decreases in DOC being caused by flushing

and export of DOC from the system as seen in forested ecosystems (Inamdar et al. 2004).

Additionally, (Lu et al. 2003) observed that canal inflow water is a major source of dissolved

organic matter in wetlands, the majority of which is DOC. As the canal surface water flows

penetrated through marsh DOC is leached from periphyton, vegetation, senescent plant material,

detritus and soil which increased the marsh DOC concentrations. The strong correlation between

DOC and specific conductance provides evidence towards this DOC dynamic (Table 4). Using

radio-isotopes, (Stern et al. 2007) determined that DOC turnover rates are typically longer than

the wetland water residence time, suggesting that most of the DOC will persist in the water

column as the water flows. Therefore, the temporal and spatial variation in marsh DOC

concentrations and to some extent POC concentrations are driven in part by the source of water

and factors contributing to the flux of DOC from ecosystem components to the overlying water

column.

Particulate OC has been used to estimate or gauge carbon burial in oceanic, estuarine, lake and

stream ecosystems (Schindler et al. 1997; Barth et al. 1998; Robertson et al. 1999; Allison et al.

2007). Rivers and streams are net transporters of POC to lake and estuarine ecosystems. In deep

lakes and marine environments POC is an important source of C and quickly mineralized.

Accumulation rates can vary several orders of magnitude across systems ranging from 0.9 g C m^{-}

^{2} yr^{-1} in riverine tributaries (Kao and Liu 1996) to 378 g C m^{-2} yr^{-1} in oceanic systems (Suess

1980). As discussed above, DOC is the dominate fraction in wetland ecosystems (Wetzel 1984)

however C accumulation rates in the Everglades is on par with most riverine ecosystems at an

average rate of 205 g C m^{-2} yr^{-1} (range: 86 – 424 g C m^{-2} yr^{-1} (Reddy et al. 1993). Although in

river ecosystems, POC concentrations and C accumulation rates can be relatively high due to

allochthonous inputs of C while POC concentrations within the Everglades are relatively low and

C accumulation rates are relatively moderate suggesting dominate authocthonous production of

C.

Dissolved Inorganic Carbon: Dissolved Organic Carbon Ratio

Each WCA experiences different hydrologic conditions. Some studies have attributed differences

in DOC:DIC ratios as indicators of hydrologic influence on dissolved C flux dynamics in stream

and river ecosystems (Elder et al. 2000; Palmer et al. 2001; Kawasaki et al. 2005). The data

suggest that the hydrologic driver is also present in WCAs. Despite similar ecological features

(i.e. ridge and slough), C loads and cycling differ between WCAs (Fig 4). Furthermore,

DOC:DIC ratios significantly differ between areas with the DOC:DIC ratio for WCA-1 deviating

from WCA-2 and WCA-3 indicating greater DOC concentrations relative to DIC within WCA-1.

Meanwhile, DOC:DIC ratios within WCA-2 and WCA-3 indicate a more balanced dissolved C

flux (Fig 5) but differ slightly presumable due to differences in water quality conditions (Table

3). The difference of WCA-1 to the other WCAs is primarily attributed to differences in

hydrologic condition with interior portions of WCA-1 receiving very little surface water flow

from canals but rather rainfall, which has low alkalinity concentrations and relatively low pH

(Julian et al. 2016). Meanwhile, WCA-2 and WCA-3 are driven largely by surface water and

groundwater flow (Harvey et al. 2004; Harvey and McCormick 2009).

In lake ecosystems decomposition, mineralization and sedimentation processes would reduce

DOC and increase DIC, which would overall decreases the DOC:DIC ratios. Meanwhile, surface

water and groundwater inputs into lakes can potentially increase DOC:DIC ratios depending on

the source and magnitude of flow (Elder et al. 2000). Lake waters are commonly supersaturated

with respect to CO_2 as a result of CO_2 generated from the decomposition of organic matter

imported from upstream, resulting in a net flux of C to the atmosphere (Cole et al. 1994, 2001).

A similar process occurs in wetland, wetlands especially the Everglades receive copious

quantities of allochthonous C and produce large quantities of autochthonous C from biomass

turnover therefore allowing the water column to become supersaturated with CO_2 resulting in

large CO_2 fluxes (Table 3).

$CO_{2(aq)}$ versus $CO_{2(atm)}$

Over the past century, the mean global $CO_{2(atm)}$ has risen from approximately 280 µatm to over

368 µatm (Keeling and Whorf 1994; Baldocchi et al. 2001). Similarly, atmospheric CO_2

concentrations increases across the study period at the Key Biscayne monitoring location. While

this location is ~60 kilometers from the study location (center of WCA-3 to Key Biscayne),

atmospheric CO_2 concentrations observed at Key Biscayne are comparable to data collected from

locations within ENP (J.G. Barr, *Unpublished data* and G. Starr, *Unpublished data*) at three

separate monitoring locations with a more limited period of record. In contrast, annual mean

p$CO_{2(aq)}$ concentrations significantly declined within the WCAs (Table 2). Even though *p*$CO_{2(aq)}$

is orders of magnitude greater than that of the *p*$CO_{2(atm)}$ this diverging relationship is unexpected

and more work is needed to explore to explain this phenomenon.

Wetlands release large quantities of C as CO_2 (Table 3) and methane due to anaerobic conditions

and the decomposition of organic matter. In wetlands, C sequestration typically outpaces C

release making wetland soils the world's largest C sinks (Kayranli et al. 2010). Here, the

difference between *$CO_{2(aq)}$ versus $CO_{2(atm)}$* indicates increasing flux to the atmosphere. This does

not mean that these ecosystems are net sources as this can be compensated by a particulate flux

from emergent vegetation or by allochthonous carbon. However, the temporal pattern suggests

long-term changes.

Wetland C reserves are at risk of becoming atmospheric C sources due to changes in climate and

hydrologic conditions (Gorham 1991). Climate change in the form of increased temperatures and

altered precipitation patterns (Kundzewicz et al. 2008; Whitehead et al. 2009) can potentially

decrease wetlands ability to store C. Increased temperatures could stimulate organic matter

decomposition and reduced rainfall and surface water flow into wetlands due to a drier climate

could lead to compaction of organic soils allowing for rapid subsidence (Reddy et al. 2006;

Kayranli et al. 2010). If the projected future climate is expected to be drier than past climates,

WCA-1 could be the most affected by climate change due to its isolated hydrology, low DIC and

high CO_2 flux rates. Currently, it is not certain how climate change will influence C storage and

other ecosystem processes of the Everglades and more direct research is needed to explore the

effects of climate change and sea-level rise on C storage and processes in the Everglades

ecosystem.

Conclusion

Hydrologic condition, nutrient inputs, and nutrient cycling significantly influence the balance,

speciation, and flux of C from wetland ecosystems consistent with the hypothesized subsidy-

stress model proposed by (Odum et al. 1979) for wetland vegetation. Within the Everglades

ecosystem the interplay between nutrient inputs and hydrologic condition exert a driving force

on the balance between DIC and DOC production via the metabolism of organic matter. As

Everglades restoration efforts progress and water quality continues to improve within the

Everglades ecosystem the C cycle and associated C pools will also respond in kind. More

specifically, impacted portions of the WCAs that have achieved a long-term TP concentration of

10 µg P L^{-1} recovering from impacted to unimpacted (Julian 2015) also exhibited significant

declines in DIC concentrations. However, in light of expected climate change (i.e. altered

precipitation, warmer air temperatures, etc.), the mechanisms of C pool control and production

could change as a result of drier climates could significantly impact rainfall driven portions of

the Everglades such as WCA-1. Perhaps future research can identify direct linkages of DOC,

DIC and climate change trends as forecasts for the Everglades Ecosystem.

Acknowledgements

We would like to thank the anonymous peer-reviewer(s) and editor(s) for their efforts and

constructive review of this manuscript. Additionally, we would like to acknowledge all of the

current and past South Florida Water Management District staff involved in the collection and

laboratory analysis of the data used in this manuscript.

Conflict of Interest Statement

The authors declare that they have no conflict of interest.

Funding

Support to write this manuscript was provided by the University of Florida Soil and Water

Sciences Wetland Biogeochemistry Laboratory Fellowship.

Authors' Contributions

PJ designed the study, performed data analysis including necessary calculations and statistics

analysis, and wrote the manuscript. SG assisted in data analysis and provided editorial assistance.

BG, ALW and TZO provided substantial editors assistance and review. All authors read and

approved the final manuscript.

References

Ågren A, Buffam I, Jansson M, Laudon H (2007) Importance of seasonality and small streams for the landscape regulation of dissolved organic carbon export. J Geophys Res Biogeosciences 112:G03003. doi: 10.1029/2006JG000381

Aiken GR, Hsu-Kim H, Ryan JN (2011) Influence of Dissolved Organic Matter on the Environmental Fate of Metals, Nanoparticles, and Colloids. Environ Sci Technol 45:3196–3201. doi: 10.1021/es103992s

Allison MA, Bianchi TS, McKee BA, Sampere TP (2007) Carbon burial on river-dominated continental shelves: Impact of historical changes in sediment loading adjacent to the Mississippi River. Geophys Res Lett 34:L01606. doi: 10.1029/2006GL028362

Baldocchi D, Falge E, Gu L, et al (2001) FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem–Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities. Bull Am Meteorol Soc 82:2415–2434. doi: 10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2

Bancroft GT, Hoffman W, Sawicki RJ, Ogden JC (1992) The Importance of the Water Conservation Areas in the Everglades to the Endangered Wood Stork (Mycteria americana). Conserv Biol 6:392–398. doi: 10.1046/j.1523-1739.1992.06030392.x

Barth JAC, Veizer J, Mayer B (1998) Origin of particulate organic carbon in the upper St. Lawrence: isotopic constraints. Earth Planet Sci Lett 162:111–121. doi: 10.1016/S0012-821X(98)00160-5

Billett MF, Moore TR (2008) Supersaturation and evasion of CO2 and CH4 in surface waters at Mer Bleue peatland, Canada. Hydrol Process 22:2044–2054. doi: 10.1002/hyp.6805

Bossio DA, Fleck JA, Scow KM, Fujii R (2006) Alteration of soil microbial communities and water quality in restored wetlands. Soil Biol Biochem 38:1223–1233. doi: 10.1016/j.soilbio.2005.09.027

Brandt LA, Portier KM, Kitchens WM (2000) Patterns of change in tree islands in Arthur R. Marshall Loxahatchee National Wildlife Refuge from 1950 to 1991. Wetlands 20:1–14. doi: 10.1672/0277-5212(2000)020[0001:POCITI]2.0.CO;2

Bronaugh DB, Werner A (2013) Zhang + Yue-Pilon trends package. CRAN R-Project

Bruland GL, Grunwald S, Osborne TZ, et al (2006) Spatial Distribution of Soil Properties in Water
Conservation Area 3 of the Everglades. Soil Sci Soc Am J 70:1662. doi: 10.2136/sssaj2005.0134

Carpenter SR, Pace ML (1997) Dystrophy and Eutrophy in Lake Ecosystems: Implications of Fluctuating
Inputs. Oikos 78:3. doi: 10.2307/3545794

Cole JJ, Caraco NF, Kling GW, Kratz TK (1994) Carbon Dioxide Supersaturation in the Surface Waters of
Lakes. Science 265:1568–1570.

Cole JJ, Cole JJ, Caraco NF, Caraco NF (2001) Carbon in catchments: connecting terrestrial carbon losses
with aquatic metabolism. Mar Freshw Res 52:101–110.

Cole JJ, Pace ML, Carpenter SR, Kitchell JF (2000) Persistence of net heterotrophy in lakes during nutrient
addition and food web manipulations. Limnol Oceanogr 45:1718–1730. doi:
10.4319/lo.2000.45.8.1718

Corstanje R, Reddy KR, Prenger JP, et al (2007) Soil microbial eco-physiological response to nutrient
enrichment in a sub-tropical wetland. Ecol Indic 7:277–289. doi: 10.1016/j.ecolind.2006.02.002

D'Angelo EM, Reddy KR (1999) Regulators of heterotrophic microbial potentials in wetland soils. Soil Biol
Biochem 31:815–830. doi: 10.1016/S0038-0717(98)00181-3

DeBusk WF, Newman S, Reddy KR (2001) Spatio–temporal patterns of soil phosphorus enrichment in
Everglades Water Conservation Area 2A. J Environ Qual 30:1438–1446.

DeBusk WF, Reddy KR (1998) Turnover of detrital organic carbon in a nutrient-impacted Everglades
marsh. Soil Sci Soc Am J 62:1460–1468.

Dinno A (2015) Dunn's test of multiple comparisons using rank sums. CRAN R-Project

Drake HL, Aumen NG, Kuhner C, et al (1996) Anaerobic Microflora of Everglades Sediments: Effects of
Nutrients on Population Profiles and Activities. Appl Environ Microbiol 62:486–493.

Duarte CM, Cebrian J (1996) The fate of marine autotrophic production. Limnol Oceanogr 41:1758–
1766.

Elder JF, Rybicki NB, Carter V, Weintraub V (2000) Sources and yields of dissolved carbon in northern
Wisconsin stream catchments with differing amounts of peatland. Wetlands 20:113–125. doi:
10.1672/0277-5212(2000)020[0113:SAYODC]2.0.CO;2

Elser JJ, Bracken MES, Cleland EE, et al (2007) Global analysis of nitrogen and phosphorus limitation of
primary producers in freshwater, marine and terrestrial ecosystems. Ecol Lett 10:1135–1142.
doi: 10.1111/j.1461-0248.2007.01113.x

Findlay SEG, Dye S, Kuehn KA (2002) Microbial growth and nitrogen retention in litter of Phragmites
australis compared to Typha angustifolia. Wetlands 22:616–625. doi: 10.1672/0277-
5212(2002)022[0616:MGANRI]2.0.CO;2

Finlay JC (2003) Controls of streamwater dissolved inorganic carbon dynamics in a forested watershed.
Biogeochemistry 62:231–252. doi: 10.1023/A:1021183023963

Fisher MM, Reddy KR (2010) Estimating the Stability of Organic Phosphorus in Wetland Soils. Soil Sci Soc
Am J 74:1398. doi: 10.2136/sssaj2009.0268

Florida Administrative Code (2008) Chapter 62-160 Quality Assurance.

Fraser CJD, Roulet NT, Moore TR (2001) Hydrology and dissolved organic carbon biogeochemistry in an
ombrotrophic bog. Hydrol Process 15:3151–3166. doi: 10.1002/hyp.322

Freeman C, Fenner N, Ostle NJ, et al (2004) Export of dissolved organic carbon from peatlands under
elevated carbon dioxide levels. Nature 430:195–198. doi: 10.1038/nature02707

Gleason PJ, Cohen AD, Smith WG, et al (1974) The environmental significance of Holocene sediments
from the Everglades and saline tidal plain. In: Environments of south Florida: Present and past.
Miami Geological Society: Coral Gables, FL, pp 287–341

Gorham E (1991) Northern peatlands: role in the carbon cycle and probable responses to climatic
warming. Ecol Appl 1:182–195.

Gu B, Miao S, Edelstein C, Dreschel T (2008) Effects of a prescribed fire on dissolved inorganic carbon
dynamics in a nutrient-enriched Everglades wetland. Fundam Appl Limnol Arch Für Hydrobiol
171:263–272. doi: 10.1127/1863-9135/2008/0171-0263

Gu B, Schelske CL, Coveney MF (2011) Low carbon dioxide partial pressure in a productive subtropical
lake. Aquat Sci 73:317–330. doi: 10.1007/s00027-010-0179-y

Hagerthey SE, Cole JJ, Kilbane D (2010) Aquatic metabolism in the Everglades: Dominance of water
column heterotrophy. Limnol Oceanogr 55:653–666.

Hanson PC, Bade DL, Carpenter SR, Kratz TK (2003) Lake metabolism: Relationships with dissolved
organic carbon and phosphorus. Limnol Oceanogr 48:1112–1119. doi:
10.4319/lo.2003.48.3.1112

Harvey JW, Krupa SL, Krest JM (2004) Ground Water Recharge and Discharge in the Central Everglades.
Ground Water 42:1090–1102. doi: 10.1111/j.1745-6584.2004.tb02646.x

Harvey JW, McCormick PV (2009) Groundwater's significance to changing hydrology, water chemistry,
and biological communities of a floodplain ecosystem, Everglades, South Florida, USA.
Hydrogeol J 17:185–201. doi: 10.1007/s10040-008-0379-x

Harwell MC, Surratt DD, Barone DM, Aumen NG (2008) Conductivity as a tracer of agricultural and urban
runoff to delineate water quality impacts in the northern Everglades. Environ Monit Assess
147:445–462. doi: 10.1007/s10661-007-0131-3

Hessen DO, Ågren GI, Anderson TR, et al (2004) Carbon sequestration in ecosystems: the role of
stoichiometry. Ecology 85:1179–1192. doi: 10.1890/02-0251

Hopkinson CS, Cai W-J, Hu X (2012) Carbon sequestration in wetland dominated coastal systems — a
561   global sink of rapidly diminishing magnitude. Curr Opin Environ Sustain 4:186–194. doi:
562   10.1016/j.cosust.2012.03.005

563   Inamdar SP, Christopher SF, Mitchell MJ (2004) Export mechanisms for dissolved organic carbon and
564   nitrate during summer storm events in a glaciated forested catchment in New York, USA. Hydrol
565   Process 18:2651–2661. doi: 10.1002/hyp.5572

566   Julian P (2015) Appendix 3A-6: Water Year 2010-2014 annual total phosphorus criteria compliance
567   assessment. In: 2015 South Florida Environmental Report. South Florida Water Management
568   District, West Palm Beach, FL,

569   Julian P, Payne GG, Xue SK (2016) Chapter 3A: Water Quality in the Everglades Protection Areas. In: 2016
570   South Florida Environmental Report. South Florida Water Management District, West Palm
571   Beach, FL,

572   Kadlec RH, Wallace SD (2009) Treatment wetlands. CRC Press, Boca Raton, FL

573   Kao S-J, Liu K-K (1996) Particulate organic carbon export from a subtropical mountainous river (Lanyang
574   Hsi) in Taiwan. Limnol Oceanogr 41:1749–1757. doi: 10.4319/lo.1996.41.8.1749

575   Kawasaki M, Ohte N, Katsuyama M (2005) Biogeochemical and hydrological controls on carbon export
576   from a forested catchment in central Japan. Ecol Res 20:347–358. doi: 10.1007/s11284-005-
577   0050-0

578   Kayranli B, Scholz M, Mustafa A, Hedmark Å (2010) Carbon Storage and Fluxes within Freshwater
579   Wetlands: a Critical Review. Wetlands 30:111–124. doi: 10.1007/s13157-009-0003-4

580   Keeling CD, Whorf TP (1994) Atmospheric CO 2 records from sites in the SIO air sampling network,
581   Trends' 93: A Compendium of Data on Global Change TA Boden, et al. Rep. Oak Ridge National
582   Laboratory, Oak Ridge, Tennessee

583   Kling GW, Kipphut GW, Miller MC (1991) Arctic Lakes and Streams as Gas Conduits to the Atmosphere:
584   Implications for Tundra Carbon Budgets. Science 251:298.

585   Kling GW, Kipphut GW, Miller MC (1992) The flux of CO2 and CH4 from lakes and rivers in arctic Alaska.
586   Hydrobiologia 240:23–36. doi: 10.1007/BF00013449

587   Kundzewicz ZW, Mata LJ, Arnell NW, et al (2008) The implications of projected climate change for
588   freshwater resources and their management. Hydrol Sci J 53:3–10. doi: 10.1623/hysj.53.1.3

589   Light SS, Dineen JW (1994) Water control in the everglades: a historical perspective. In: Davis S, Ogden J
590   (eds) Everglades: The ecosystem and its restoration. St. Lucie Press, Delray Beach, FL, pp 47–84

591   Lu H, Yang L, Shabbir S, Wu Y (2014) The adsorption process during inorganic phosphorus removal by
592   cultured periphyton. Environ Sci Pollut Res 21:8782–8791. doi: 10.1007/s11356-014-2813-z

593   Lu XQ, Maie N, Hanna JV, et al (2003) Molecular characterization of dissolved organic matter in
594   freshwater wetlands of the Florida Everglades. Water Res 37:2599–2606. doi: 10.1016/S0043-
595   1354(03)00081-2

596   Newman S, Hagerthey SE (2011) Water Conservation Area 1: A Case Study of Hydrology, Nutrient, and
597   Mineral Influences on Biogeochemical Processes. Crit Rev Environ Sci Technol 41:702–722. doi:
598   10.1080/10643389.2010.530910

599   Odum EP, Finn JT, Franz EH (1979) Perturbation Theory and the Subsidy-Stress Gradient. BioScience
600   29:349–352. doi: 10.2307/1307690

601   Odum HT (1956) Primary production in flowing waters. Limnol Ocean 1:102–117.

602   Palmer SM, Hope D, Billett MF, et al (2001) Sources of organic and inorganic carbon in a headwater
603   stream: Evidence from carbon isotope studies. Biogeochemistry 52:321–338. doi:
604   10.1023/A:1006447706565

605   Qualls RG, Richardson CJ (2000) Phosphorus enrichment affects litter decomposition, immobilization,
606   and soil microbial phosphorus in wetland mesocosms. Soil Sci Soc Am J 64:799–808.

607   Qualls RG, Richardson CJ (2003) Factors controlling concentration, export, and decomposition of
608   dissolved organic nutrients in the Everglades of Florida. Biogeochemistry 62:197–229. doi:
609   10.1023/A:1021150503664

610   Raven J, Beardall J, Griffiths H (1982) Inorganic C-sources for Lemanea, Cladophora and Ranunculus in a
611   fast-flowing stream: Measurements of gas exchange and of carbon isotope ratio and their
612   ecological implications. Oecologia 53:68–78. doi: 10.1007/BF00377138

613   Raymond PA, Bauer JE, Cole JJ (2000) Atmospheric CO2 evasion, dissolved inorganic carbon production,
614   and net heterotrophy in the York River estuary. Limnol Oceanogr 45:1707–1717. doi:
615   10.4319/lo.2000.45.8.1707

616   Reddy KR, DeLaune RD, DeBusk WF, Koch MS (1993) Long-term nutrient accumulation rates in the
617   Everglades. Soil Sci Soc Am J 57:1147–1155.

618   Reddy KR, Osborne TZ, Inglett KS, Corstanje R (2006) Influence of water levels on subsidence of organic
619   soils in the upper St. Johns River Basin. Saint Johns River Water Management District, Palatka, FL
620   USA

621   Reddy KR, Wang Y, DeBusk WF, et al (1998) Forms of soil phosphorus in selected hydrologic units of the
622   Florida Everglades. Soil Sci Soc Am J 62:1134–1147.

623   Richey JE, Melack JM, Aufdenkampe AK, et al (2002) Outgassing from Amazonian rivers and wetlands as
624   a large tropical source of atmospheric CO2. Nature 416:617–620. doi: 10.1038/416617a

625   Robertson AI, Bunn SE, Boon PI, Walker KF (1999) Sources, sinks and transformations of organic carbon
626   in Australian floodplain rivers. Mar Freshw Res 50:813. doi: 10.1071/MF99112

627   Sand-Jensen K, Frost-Christensen H (1998) Photosynthesis of amphibious and obligately submerged
628   plants in CO2-rich lowland streams. Oecologia 117:31–39. doi: 10.1007/s004420050628

629   Scheidt D, Kalla PI (2007) Everglades ecosystem assessment: water management and quality,
630   eutrophication, mercury contamination, soil and habitat: monitoring for adapyive management:
631   a R-EMAP status report. United States Environmental Protection Agency, Athens, GA

632   Schindler DW, Curtis PJ, Bayley SE, et al (1997) Climate-Induced Changes in the Dissolved Organic Carbon
633   Budgets of Boreal Lakes. Biogeochemistry 36:9–28.

634   Stern J, Wang Y, Gu B, Newman J (2007) Distribution and turnover of carbon in natural and constructed
635   wetlands in the Florida Everglades. Appl Geochem 22:1936–1948. doi:
636   10.1016/j.apgeochem.2007.04.007

637   Suess E (1980) Particulate organic carbon flux in the oceans—surface. Nature 288:261.

638   Surratt DD, Waldon MG, Harwell MC, Aumen NG (2008) Time-series and spatial tracking of polluted
639   canal water intrusion into wetlands of a National Wildlife Refuge in Florida, USA. Wetlands
640   28:176–183. doi: 10.1672/07-74.1

641   Telmer K, Veizer J (1999) Carbon fluxes, pCO2 and substrate weathering in a large northern river basin,
642   Canada: carbon isotope perspectives. Chem Geol 159:61–86. doi: 10.1016/S0009-
643   2541(99)00034-0

644   Updegraff K, Pastor J, Bridgham SD, Johnston CA (1995) Environmental and Substrate Controls over
645   Carbon and Nitrogen Mineralization in Northern Wetlands. Ecol Appl 5:151–163. doi:
646   10.2307/1942060

647   Waiser M, Robarts R (2004) Net heterotrophy in productive prairie wetlands with high DOC
648   concentrations. Aquat Microb Ecol 34:279–290. doi: 10.3354/ame034279

649   Wetzel RG (1992) Gradient-dominated ecosystems: sources and regulatory functions of dissolved
650   organic matter in freshwater ecosystems. In: Salonen K, Kairesalo T, Jones RI (eds) Dissolved
651   Organic Matter in Lacustrine Ecosystems. Springer Netherlands, pp 181–198

652   Wetzel RG (1984) Detrital Dissolved and Particulate Organic Carbon Functions in Aquatic Ecosystems.
653   Bull Mar Sci 35:503–509.

654   Wetzel RG, Likens GE (2000) The Inorganic Carbon Complex: Alkalinity, Acidity, CO2, pH, Total Inorganic
655   Carbon, Hardness, Aluminum. In: Limnological Analyses. Springer New York, pp 113–135

656   Whitehead PG, Wilby RL, Battarbee RW, et al (2009) A review of the potential impacts of climate change
657   on surface water quality. Hydrol Sci J 54:101–123. doi: 10.1623/hysj.54.1.101

658   Wright AL, Reddy KR (2001) Heterotrophic Microbial Activity in Northern Everglades Wetland Soils. Soil
659   Sci Soc Am J 65:1856. doi: 10.2136/sssaj2001.1856

Figures and Tables

Fig. 1. Map of surface water quality, stage and weather monitoring location within the Everglades Protection Area relative to surrounding features including the Everglades Stormwater Treatment Area, Everglades Agricultural Area and Water Conservation Areas.

Fig. 2. Annual Kendall trend analysis results for each monitoring location with enough data (greater than six samples per year and three water years) for dissolved inorganic carbon, dissolved organic carbon, particulate organic carbon and surface water pCO_2 ($pCO_{2(aq)}$). Significantly increasing or decreasing trends were determined based on a ρ-value <0.05 and positive or negative Kendall τ values to denote direction. No trend results were identified as ρ-values>0.05.

Fig 3 Annual mean dissolved inorganic carbon concentrations, dissolved organic carbon, particulate organic carbon and surface water $pCO_{2(aq)}$ for interior portions of the Everglades Protection Area between water years 1995 and 2015 (May 1, 1994 – April 30, 2015).

Fig. 4. Boxplot of annual mean dissolved inorganic concentration, dissolved organic carbon, particulate organic carbon and surface water $pCO_{2(aq)}$ concentration for each region of the Everglades Protection Area. Letters above box-and-whicker plots indicate statistical differences based on Dunn test results.

Fig. 5. Boxplot of Dissolved Organic Carbon and Dissolved Inorganic Carbon annual mean concentration ratio for each region of the Everglades Protection Area. Graphical representation of this data can be found in Supplemental Information.

Gulf of Mexico

Atlantic Ocean

△ Water Quality Station

● Stage Station

■ Meterologic Station

○ Water Control Structures

— Canals

Wildlife Management Areas

Everglades Stormwater Treatment Areas

Restoration Strategies Features (Circa 2017)

Everglades Agrictural Area

N

0   10   20   30
Kilometers

WCA-1

WCA 2A

WCA 2B

WCA-3A

WCA-3B

Table 1. Equations used to calculate surface water pCO_2 concentrations and CO_2 flux rates.

Parameter	Units	Equation	Reference	Equation
Dissolved free CO_2 ($CO_{2(aq)}$)	mol m^{-3}	$\left[CO_{2(aq)}\right] = \dfrac{DIC}{12.01} * \left[7.0x10^6 \times \exp(-2.414 \times pH)\right]$	(Wetzel and Likens 2000)	1
Surface Water pCO_2	atm	$pCO_2 = \dfrac{\left[CO_{2(aq)}\right]}{K_H}$	(Gu et al. 2008)	2
CO_2 Dissolution Constant (K_H)	mol m^{-3} atm^{-1}	$K_H = 0.0279T^2 - 2.3895T + 75.819$	(Gu et al 2008)	3
CO_2 ($CO_{2(atm)}$) at the top of the stagent layer.	mol m^{-3}	$\left[CO_{2(atm)}\right] = K_H * pCO_{2(atm)}$	(Gu et al. 2011)	4
CO_2 Flux (F_{CO2})	mol m^{-2} s^{-1}	$F_{CO_2} = \left(\left[CO_{2(atm)}\right] - \left[CO_{2(aq)}\right]\right) \times \dfrac{D_{CO_2}}{z}$	(Gu et al. 2011)	5
CO_2 Diffusion Coefficient (D_{CO2})	m^{-2} s^{-1}	$D_{CO_2} = \dfrac{(-12.2048 + 0.04752T)}{10^9}$	(Akgerman and Gainer 1972)	6
Surface boundary layer thickness (z)	m	$z = \dfrac{10^{(2.56 - 0.133\ W)}}{10^6}$	(Kling et al. 1992)	7

DIC= Dissolved Inorganic Carbon; K_H = Henry's Law CO_2 dissolution constant; T=Temperature (°C)

Table 2. Trend analysis results expressed as Kendall's τ, ρ-value (Thiel-Sen Slope) for annual mean dissolved inorganic carbon (DIC), dissolved organic carbon (DOC), particulate organic carbon (POC), surface water $p\mathrm{CO_2}$ ($p\mathrm{CO_{2(aq)}}$) and $\mathrm{CO_2}$ flux rate for each region of the Everglades Protection Area between water years 1995 and 2015 (May 1, 1994 – April 30, 2015).

	Thiel-Sen Slope Units	WCA1	WCA2	WCA3
DIC	mg L^{-1} Yr^{-1}	-0.30, 0.06 (-0.32)	-0.34, <0.05 (-0.53)	-0.14, 0.39 (-0.13)
DOC	mg L^{-1} Yr^{-1}	-24, 0.15 (-0.12)	-0.39, <0.05 (-0.37)	-0.11, 0.50 (-0.08)
POC	mg L^{-1} Yr^{-1}	-0.54, <0.01 (-0.05)	-0.58, <0.01 (-0.09)	-0.59, <0.01 (-0.08)
$p\mathrm{CO_{2(aq)}}$	µatm Yr^{-1}	-0.30, 0.06 (-1454)	-0.50, <0.01 (-792)	-0.29, 0.07 (-670)
$\mathrm{CO_2}$ Flux	mg m^{-2} d^{-1} Yr^{-1}	-0.31, <0.05 (-25.0)	-0.40, <0.05 (-7.4)	-0.34, <0.05 (-8.3)

Table 3. Mean ± standard error surface water parameters observed during this study within each portion of the Everglades Protection Area between water years 1995-2015 (May 1, 1994 – April 30, 2015).

Parameter	Units	WCA-1	WCA-2	WCA-3
Alkalinity	mg CaCO$_3$ L^{-1}	44.1 ± 0.9	248.4 ± 1.4	168.4 ± 0.9
Water Temperature	°C	23.4 ± 0.1	24.1 ± 0.1	24.1 ± 0.1
pH	SU	6.6 ± 0.01	7.3 ± 0.01	7.2 ± 0.01
Specific Conductance	μS cm^{-1}	213.6 ± 3.8	936.5 ± 5.9	517.5 ± 3.6
Dissolved Inorganic Carbon	mg C L^{-1}	17.2 ± 0.6	67.2 ± 0.7	47.0 ± 0.5
Dissolved Organic Carbon	mg C L^{-1}	19.4 ± 0.2	35.8 ± 0.3	21.1 ± 0.2
Particulate Organic Carbon	mg C L^{-1}	0.65 ± 0.05	1.64 ± 0.20	1.04 ± 0.09
Total Phosphorus	μg P L^{-1}	9.1 ± 0.3	25.5 ± 1.2	8.9 ± 0.2
Total Nitrogen	mg N L^{-1}	1.19 ± 0.01	2.03 ± 0.02	1.33 ± 0.01
pCO$_{2,aq}$	μatm	54,379 ± 6,445	26,714 ± 1,020	24,963 ± 984
CO$_2$ Flux	mg m^{-2} d^{-1}	876.2 ± 96.4	285.1 ± 10.1	304.8 ± 13.4

Table 4. Spearman correlation analysis results of monthly mean surface water *carbon variables* and various water quality parameters. Values are represented as spearman's rho (ρ-value). Correlations that are not statistically significant are identified in bold text.

	Parameter	WCA-1	WCA-2	WCA-3
Dissolved Inorganic Carbon	Stage Elevation	**0.19 (0.14)**	**-0.25 (0.06)**	-0.49 (<0.01)
	Water Temperature	**0.06 (0.63)**	-0.31 (<0.01)	-0.57 (<0.01)
	Total Phosphorus	0.39 (<0.01)	0.34 (<0.05)	**-0.23 (0.07)**
	Total Nitrogen	0.62 (<0.01)	0.65 (<0.01)	0.59 (<0.01)
	Specific Conductance	0.74 (<0.01)	0.67 (<0.01)	0.65 (<0.01)
Dissolved Organic Carbon	Stage Elevation	**0.07 (0.61)**	-0.33 (<0.05)	-0.28 (<0.05)
	Water Temperature	**-0.08 (0.52)**	**-0.14 (0.30)**	**-0.03 (0.79)**
	Total Phosphorus	**0.25 (0.05)**	0.38 (<0.01)	**0.009 (0.95)**
	Total Nitrogen	0.78 (<0.01)	0.67 (<0.01)	0.72 (<0.01)
	Specific Conductance	0.74 (<0.01)	0.52 (<0.01)	0.60 (<0.01)
Particulate Organic Carbon	Stage Elevation	**-0.23 (0.10)**	**-0.007 (0.96)**	**0.14 (0.28)**
	Water Temperature	**-0.15 (0.26)**	**-0.12 (0.37)**	**-0.18 (0.18)**
	Total Phosphorus	**0.08 (0.55)**	0.40 (<0.01)	0.37 (<0.01)
	Total Nitrogen	**0.19 (0.16)**	**0.17 (0.23)**	**-0.11 (0.41)**
	Specific Conductance	**0.30 (<0.05)**	0.04 (0.79)	-0.32 (<0.05)
$pCO_{2,aq}$	Stage Elevation	**-0.05 (0.20)**	**0.25 (0.06)**	**0.19 (0.14)**
	Water Temperature	**0.25 (0.05)**	0.35 (<0.01)	**0.21 (0.10)**
	Total Phosphorus	0.30 (<0.05)	0.48 (<0.01)	**0.22 (0.09)**
	Total Nitrogen	**0.22 (0.10)**	**0.08 (0.51)**	**0.09 (0.48)**
	Specific Conductance	**0.12 (0.38)**	**0.02 (0.91)**	**-0.13 (0.31)**