Title:

Biased gene retention in the face of massive nuclear introgression obscures species relationships

Authors:

Evan S. Forsythe[1], Andrew D. L. Nelson[1], Mark A. Beilstein[1*]

Affiliations:

[1]School of Plant Sciences, University of Arizona, Tucson, AZ 85721, USA.

Corresponding Author:

Mark A. Beilstein; 1140 E. South Campus Dr.; Forbes 303; Tucson, AZ 85721; (520) 626-1562;

mbeilstein@email.arizona.edu

Keywords:

Introgression | Arabidopsis | Phylogenomics | Cytonuclear interactions

Abstract:

Phylogenomic analyses are recovering previously hidden histories of hybridization, revealing the genomic consequences of these events on the architecture of extant genomes. We exploit a suite of genomic resources to show that introgressive hybridization occurred between close relatives of Arabidopsis, impacting our understanding of species relationships in the group. The composition of introgressed and retained genes indicates that selection against incompatible cytonuclear and nuclear-nuclear interactions likely acted during introgression, while neutral processes also contributed to genome composition through the retention of ancient haplotype blocks. We also developed a divergence-based test to distinguish donor from recipient lineages without the requirement of additional taxon-sampling. Finally, to our great surprise, we find that cytonuclear discordance appears to have arisen via extensive nuclear, rather than cytoplasmic, introgression, meaning that most of the genome was displaced during introgression, while only a small proportion of native alleles were retained.

Significance:

Hybridization can lead to the transfer of genes across species boundaries, impacting the evolution of the recipient species through a process known as introgression (IG). IG can facilitate sharing of adaptive alleles but can also result in deleterious combinations of incompatible foreign alleles (i.e. epistatic incompatibility). How hybrids overcome these epistatic hurdles remains an open question. Here, we characterize IG in Arabidopsis and its closest relatives. Interestingly, our analyses favor an evolutionary scenario in which the vast majority of nuclear genes were displaced by foreign alleles during the evolution of *Capsella* and *Camelina*, obscuring species relationships. Simultaneously, a subset of nuclear genes resisted displacement, thereby

minimizing epistatic incompatibilities between the organellar and nuclear genomes, suggesting

one potentially fundamental mechanism for overcoming barriers to hybridization.

Background:

Hybridization is a driving force in plant evolution[1], occurring naturally in ~10% of all

plants, including 22 of the world's 25 most important crops[2]. Botanists have long realized that

through backcrossing to parents, hybrids can serve as bridges for the transfer of genes between

species, a process known as introgression (IG). As more genome sequences become available,

comparative analyses have revealed the watermarks of historical IG events in plant and animal

genomes[3–5]. Cytonuclear discordance is a hallmark of many IG events, occurring, in part,

because nuclear and cytoplasmic DNA differ in their mode of inheritance. In plants, this discord

is often referred to as "chloroplast capture," which has been observed in cases where IG of the

chloroplast genome occurs in the near absence of nuclear IG or via nuclear IG to a maternal

recipient[6]. Moreover, unlinked nuclear and cytoplasmic IG creates an interaction interface for

independently evolving nuclear and cytoplasmic alleles, either of which may have accumulated

mutations that result in incompatibilities with deleterious effects when they are united in hybrids.

Such incompatibilities could exert a selective pressure that influences which hybrid genotypes

are permissible thereby favoring the co-introgression of alleles for interacting genes[7].

Disentangling IG from speciation is particularly important because IG may facilitate the

transfer of adaptive traits. Robust statistical techniques[5,8–15] have been developed to detect the

signatures of historical introgression (IG) in extant and extinct genomes. While existing

techniques are able to identify the taxa that exchanged genes during IG using a four-taxon

system, most methods do not explicitly distinguish which taxon served as donor and which as

recipient during IG (i.e. polarization of IG directionality), an important distinction considering

that IG impacts the evolution of the recipient lineage[4,6]. The existing methods that do polarize IG

are only able to do so when there is a fifth taxon available, which diverged from its sister taxon

involved in IG[11], prior to the proposed IG event.

The wealth of genomic and functional data in Arabidopsis[16], combined with publicly

available genome sequence for 26 species make the plant family Brassicaceae an ideal group for

comparative genomics. Phylogeny of the group has been the focus of numerous studies[17–23],

providing a robust estimate of its evolutionary history. While the genus *Arabidopsis* is well

circumscribed[20,24], the identity of its closest relatives remains an open question. Phylogenetic

studies to date recover three monophyletic groups: clade A, including the sequenced genomes of

A. thaliana[16] and *A. lyrata*[25]; clade B, including the *B. stricta* genome[26]; and clade C, including

the genomes of *Capsella rubella*, *C. grandiflora*[27], and *Camelina sativa*[28] (Supplementary

Information). Analyses using nuclear markers strongly support A(BC), which is most often cited

as the species tree[17,19,21–23]. Organellar markers strongly support B(AC)[18,19,29,30] (Fig. 1a-b and

Table S1). The genome sequences listed above can be used to explore the processes underlying

this incongruence.

Here, we exploit a suite of genomic resources to explore a putative chloroplast capture

event involving Arabidopsis and its closest relatives by inferring gene trees for markers in all

three cellular genomes from six available whole genome sequences. We document cytonuclear

discordance and ask if it arose through IG of organelles or extensive IG of nuclear genes.

Further, using a new divergence-based approach, we ask: Which lineage was the recipient of

introgressed alleles? Finally, we explore the extent to which neutral processes, such as physical

linkage as well as non-neutral processes, such as selection against incompatible alleles at

interacting loci, shaped the recipient genome.

Results:

Gene tree incongruence within and between organelle and nuclear genomes. We searched

for incongruent histories present within and among nuclear and organellar genomes in

representative species from each clade. We included *Cardamine hirsuta*[31] and *Eutrema*

salsugineum[32] as outgroups. We considered three processes capable of producing incongruent

histories: duplication and loss, incomplete lineage sorting (ILS), and IG. In addition, we assessed

the possible contribution of phylogenetic error or 'noise'.

Given the well-known history of whole genome duplication in Brassicaceae, we took

extensive measures to minimize the possibility that duplication and loss biased our inferences.

We identified single-copy nuclear genes as well as genes that were retained in all species post-

duplication (see Discussion). In the chloroplast, we found 32 single-copy genes, while in

mitochondria we identified eight. Maximum likelihood (ML) analyses of these yielded well-

supported B(AC) trees (Fig. 1a and Fig. S2d-g). We identified 10,193 single-copy nuclear genes

using *Orthofinder*[35] (denoted as 'full single-copy dataset') (Fig. S1a-c). These genes were

indicated as single-copy by *Orthofinder* because they form clusters that include exactly one locus

from each species (with the exception of *C. sativa*, see **Methods**). These single-copy genes span

the eight chromosomes of *C. rubella* (Fig. S1d), whose karyotype serves as an estimate of the

ancestral karyotype for these species[36]. ML analyses yielded 8,490 (87.6%) A(BC), 774 (8.0%)

B(AC), and 429 (4.4%) C(AB) trees (Fig. 1c-f and Table S2).

The most parsimonious explanation for our single-copy genes is that they were either not

duplicated in our focal species or, if duplicated, were returned to single-copy before a speciation

occurred, thus behaving as unduplicated in a phylogenetic context, meaning that any observed

incongruent topologies resulted from a process other that duplication. However, while not

parsimonious, it is important to consider the possibility that ancestral duplication, paralog

retention through two speciation events, and lineage specific loss events led to hidden out-

paralogs in our dataset. To further reduce the probability that this series of events contributed to

incongruent gene trees, we further filtered our dataset to include only genes that were previously

indicated as reliable single-copy markers in angiosperms[33,34]. This filter reduced our single-copy

dataset to 2,098 genes (Fig. S1e-f). We combined this dataset with genes that were duplicated

during whole genome duplication[37] but did not undergo loss in focal species to yield a dataset of

2,747 genes, which we denote as 'conservatively single-copy', so named because they are the

genes that are least likely to contain hidden out-paralogs. ML analyses of these genes yielded

2,236 (86.5%) A(BC), 236 (9.1%) B(AC), and 114 (4.4%) C(AB) trees (Fig. 1b-f), consistent

with our results from the full single-copy dataset.

To ask whether phylogenetic noise contributed to incongruent nuclear gene tree

topologies, we also filtered our single-copy nuclear gene tree results to contain only trees in

which the observed topology was supported by at least 70% bootstrap support (BS) and found

that B(AC) and C(AB) trees were still present (Fig. 1f). Together, these analyses confirm the

incongruent histories present in the organellar and nuclear genomes and indicate that

incongruence cannot be fully explained by gene duplication and loss or by phylogenetic noise.

Contribution of introgression to incongruent gene trees. A number of approaches have been

developed to determine the relative contributions of ILS and IG to gene tree incongruence.

Comparative genomic approaches are based on the *D*-statistic[5,9], which is typically applied to

whole genome alignments and is calculated by determining the frequency of site patterns. It was

not feasible to construct accurate whole genome alignments among our taxa, and thus we used

multiple sequence alignments from single-copy genes to calculate D- and F-statistics. Analyses

of both full and conservatively single-copy gene alignments indicated that introgression occurred

(Table S3; positive D and F). Since phylogenomic analyses often focus on comparisons of gene

trees rather than site-patterns, we also applied the rational of the D-statistic to gene trees, using

gene tree topologies as proxies for site patterns to calculate a related statistic, referred to here as

D_{GT} (see **Methods**). Consistent with D and F, D_{GT} indicated that ILS is sufficient to explain the

frequency of C(AB) but not the observed frequencies of A(BC) and B(AC) in the nuclear

genome (Table S4; positive D_{GT}).

Coalescent based approaches[14,15] use gene trees to distinguish between organismal

histories that are tree-like (incongruencies among trees arise from ILS) and network-like

(incongruencies result from ILS + IG). We analyzed our gene tree data in *PhyloNet*[15] and found

that reticulate networks were favored over tree-like evolution (Fig. S2j-q; ΔAIC $\geq$ 87.80 and

ΔBIC $\geq$ 73.50). Similarly, *Tree Incongruence Checking in R* (*TICR*)[14] indicated that a simple

tree-like history fit the data poorly because the concordance factors for a significant proportion

of quartets departed from expectation (Fig. S2r-u; $p = 0.00058$; χ^2 test). In sum, both

comparative genomic and coalescent based approaches support an evolutionary history that

includes IG.

Recovery of the species branching order and introgression events. To uncover which

lineages were affected by IG, we determined the relative timing of the B(AC) and A(BC)

branching events by calculating node depths (Fig. 2)[38]. IG nodes are expected to be younger than

speciation nodes[12,38,39] because IG produces incongruent trees when it occurs between non-sister

species subsequent to speciation[4,5,9] (illustrated by Fig. 2a). Therefore, we calculated the depth of

the node uniting clade A with clade C in nuclear B(AC) trees and compared it with the depth of

the node uniting the B and C clades in nuclear A(BC) trees (Fig. 2a-c, N.D.). We calculated node

depths using four separate measures to account for potential biases (Fig. 2d-g). To account for

selection on amino acids, we used synonymous divergence (dS) (Fig. 2d). To account for

potential differing rates of evolution across the genome, we normalized dS using the divergence

between the clade of interest and an outgroup (i.e. 'relative node depth')[12] (Fig. 2e). To account

for potential differences in rates of evolution between lineages, we also calculated node depths

from ultrametric trees in which the rates of evolution had been smoothed across the tree using a

penalized likelihood approach[40] (Fig. 2f, Fig. S3, and Table. S5). To account for potential

intragene discordance due to recombination within a gene, we divided each gene alignment into

200nt windows, inferred a neighbor joining tree for each window, and only calculated node

depth from windows that were concordant with the ML tree for the gene, thus minimizing the

probability of recombination within the loci from which node depth is calculated (Fig. 2g, Fig

S4). For all four node depth measures, the node depth for A(BC) was significantly shallower than

for B(AC) (Fig. 2d-g, Fig. S3, Fig S4 and Table S6; $p<2.2e-16$, Wilcoxon), indicating that IG

rather than speciation produced the observed A(BC) nuclear gene trees. This result is insensitive

to the removal of the deepest nodes in both A(BC) and B(AC) bins (Fig. S3o-t). Hence, node

depth data suggest that A and C diverged from each other prior to the exchange of genes between

clade B and C via IG. This surprising result stands in opposition to previously published trees

inferred from single or concatenated nuclear genes, which strongly favor A(BC)[20,22–24]. However,

it bolsters the argument that B(AC) best represents the species branching order despite the low

frequency of these genes in the nucleus (similar to [38]), and further suggests that the vast majority

of nuclear genes in either B or C arrived there via IG. We discuss the implications of this finding

on the concept of the species branching order (see **Discussion**). It should be noted that our

downstream analyses of selection and neutral processes (Fig. 4, Fig. S6, and Table S6) are

framed in the context of nuclear introgression but would remain equally valid if cytonuclear

discordance arose via organellar introgression.

Identification of unidirectional introgression donor and recipient linages. We next asked

whether transfer of genetic material during IG was unidirectional and, if so, which of the two

clade ancestors was the donor and which was the recipient of introgressed alleles. Existing

methods for polarizing the direction of IG require additional taxa with specific phylogenetic

positioning relative to the introgression event[5,11]. No such taxa are available for our inferred

introgression event; therefore, existing polarization methods are not applicable to our data.

Instead, we present a divergence-based approach to infer directionality of IG, calculated from

pairwise sequence divergence between taxa involved in IG and a sister taxon by comparing

divergence values obtained from introgressed loci *vs*. non-introgressed loci (see **Methods**).

We calculated the rate of pairwise dS for all pairs of species and used these to determine

the average dS between pairs of clades (B vs. C = $dS(B,C)$; A vs. C = $dS(A,C)$; A vs. B =

$dS(A,B)$) (Fig. S5). We denoted dS values with $_{SP}$ when obtained from B(AC) trees (our inferred

species branching order) and $_{IG}$ when obtained from A(BC) trees (IG branching order) (Fig. 3a

and b). We compared $dS(B,C)_{IG}$, $dS(A,C)_{IG}$, and $dS(A,B)_{IG}$ to $dS(B,C)_{SP}$, $dS(A,C)_{SP}$, and

$dS(A,B)_{SP}$, respectively, to ask if divergence is consistent with unidirectional IG from B to C

(Fig. 3a) or from C to B (Fig. 3b), or with bidirectional IG. We found that $dS(B,C)_{SP} > dS(B,C)_{IG}$

(p<2.2e-16, Wilcoxon), $dS(A,C)_{SP} < dS(A,C)_{IG}$ (p=2.365e-12), and $dS(A,B)_{SP} = dS(A,B)_{IG}$

(p=0.1056), indicating unidirectional IG from clade B to clade C (Fig. 3c and Fig. S5). This

result is consistent with the *Phylonet* network shown in Fig. S2m and one shown in Fig. S2n,

which respectively indicate that 96.6% and 90.5% of sampled nuclear alleles were introgressed

from clade B to C.

The role of cytonuclear interactions during introgression. The IG that occurred during the

evolution of clade C resulted in a genome in which the majority of nuclear alleles were displaced

by alleles from clade B, while native organellar genomes were maintained. We asked whether we

could detect patterns within the set of nuclear genes that were also maintained alongside

organelles during IG. We hypothesized that during the period of exchange, selection would favor

the retention of alleles that maintain cytonuclear interactions, especially when replacement with

the paternal allele is deleterious[7]. Using Arabidopsis Gene Ontology (GO) data[41], we asked if

B(AC) nuclear genes were significantly enriched for chloroplast and mitochondrial-localized GO

terms, indicating that these genes are more likely to be retained than are other nuclear genes. We

calculated enrichment (E) for each GO category by comparing the percentage of B(AC) nuclear

genes with a given GO term to the percentage of A(BC) genes with that term (see **Methods**).

Positive E indicates enrichment among B(AC) genes; negative E indicates enrichment among

A(BC) genes. B(AC) nuclear genes are significantly enriched for chloroplast (E=0.10,

p=0.00443, 1-tail Fisher's) and mitochondrial localized (E=0.13, p=00250) GO terms (Fig. 4a

and Table S6). Enrichment was also detected at the level of organelle-localized processes such as

photosynthesis (E=0.29, p=0.01184), including the light (E=0.44, p=0.00533) and dark (E=0.65,

p=0.04469) reactions. The opposite enrichment pattern exists for nuclear localized genes (E=-

0.06, p=0.00936) (Fig. 4a). In sum, these results suggest a role for selection in shaping which

genes were displaced during IG.

The role of nuclear-nuclear interactions during introgression. We also asked if interactions

between/among nuclear genes influenced the likelihood of replacement by foreign alleles. Using

Arabidopsis protein-protein interaction data[42], we constructed an interaction network of the full

set of single-copy nuclear genes (Fig 4b). To assess whether genes with shared history are

clustered in the network, we calculated its assortativity coefficient (A) (**Methods**). We assessed

significance by generating a null distribution for A using 10,000 networks of the same size and

shape with randomized topology assignments. In our empirical network, A was significantly

positive (A=0.0885, p=0.00189, Z-test), and hence topologies are clustered (Fig. 4c), indicating

that selection acted against genotypes containing interactions between maternal and paternal

alleles.

The role of physical linkage during introgression. While it appears gene function exerted

influence on nuclear IG, we also wondered whether blocks of genes with similar histories were

physically clustered on chromosomes. We looked for evidence of haplotype blocks using the *C.*

rubella genome map (Fig. 4d). Previous studies in this group estimate linkage disequilibrium to

decay within 10kb[43,44], creating blocks of paternal or maternal genes around that size. We

assessed the physical clustering of genes with shared history by two measures: 1) number of

instances in which genes with the same topology are located within 10kb of each other (Fig.

S6a), and 2) number of instances in which neighboring genes share topology, regardless of

distance (Fig. S6b). The second measure provides a simple measure of clustering without

requiring an estimate of ancestral linkage. We compared both measures to a null distribution

generated from 10,000 replicated chromosome maps in which the topology assignments were

randomized across the marker genes. By both measures, we found significant clustering of

A(BC) (measure 1: p=3.022e-8; measure 2: p=1.41364e-10, Z-test) and B(AC) (measure 1:

p=0.003645; measure 2: p=1.7169e-11) genes (Fig. S6c-h). The observed clustering indicates

that haplotype blocks of co-transferred and un-transferred genes are detectable in extant

genomes, pointing to physical linkage as a factor influencing whether genes are transferred or

retained.

Discussion:

Phylogenomic studies in plants face unique challenges. The prevalence of gene and

genome duplication complicates the detection of orthologs, and thus choosing markers that

minimize duplication is extremely important when applying tests of IG originally developed for

animals[5]. Since duplication history cannot be definitively known, we can never be sure that

cryptic duplication has not introduced phylogenetic incongruence into our dataset; this is a risk in

any phylogenetic study, especially in plants. We acknowledge that all nuclear genes have

undergone duplication at some point in Brassicaceae[37] and address this challenge by specifically

targeting genes least likely to have undergone duplication during the speciation and introgression

events we detected. If duplication was biasing the results we obtained from our full single-copy

dataset, we expected that the proportion of B(AC) trees would have decreased in our

conservatively single-copy dataset. However, the proportions we observed were not substantially

impacted by our conservative single-copy filter. In fact, the proportion of B(AC) genes was

slightly higher in the conservatively single-copy genes, the opposite of what we would expect if

duplication was creating incongruent trees. Moreover, results of the D-, F-, and D_{GT}-statistics

from both datasets significantly indicated IG (Table S3, and Table S4), another indication that

biases associated with cryptic duplication and loss are not driving our conclusions of IG.

We applied several methods to distinguish between IG and ILS. Like all applications of

D and related statistics, it's important to acknowledge that ancestral population structure may

produce signatures that mimic IG[45]. However, when this possibility was thoroughly explored in

the case of Neanderthal, IG remained the favored hypothesis[46]. Here, regardless of the measure

or approach employed, our results (Fig. S2, Table S3, and Table S4), were always consistent

with an explanation of IG rather than ILS or duplication and loss. While we appreciate the

limitations of each approach, here we argue that the consistent finding of IG favors this

hypothesis over all others.

Our initial interpretation of the observed phylogenetic incongruence was that A(BC)

resulted from simple speciation events and B(AC) resulted from IG between clades A and C, a

pattern we referred to as cytoplasmic IG. However, in light of recent findings from

mosquitos[38,47], we thought it important to consider alternative hypotheses. Using the same

approach that revealed IG in mosquitos, we calculated the mean node depth for each of the

alternative topologies we recovered for nuclear genes. In addition, we employed several

strategies to account for the effects of selection (Fig. 2d), effective population size variation

across the genome (Fig. 2e), linage-specific effects (Fig. 2f). and intragenic recombination (Fig.

2g) on our node depth calculations. In all cases, our node depth comparisons rejected the

hypothesis that the node uniting clades A and C on B(AC) trees resulted from an introgression

event, and instead indicated that the node uniting clades B and C on A(BC) trees resulted from

an introgression between clades B and C. Based on these results, we suggest that the 'true'

species branching order is B(AC).

There is growing debate about the efficacy of bifurcating phylogenies in describing

organismal evolution, prompting the development of powerful network frameworks that

highlight reticulation in species relationships. While our analysis reinforces the importance of

considering reticulation, we also show that bifurcating trees should not be entirely abandoned in

the face of reticulation. The presence of reticulation does not preclude the occurrence of simple

bifurcating speciation events, it simply means some bifurcations result from speciation while

others result from IG. Therefore, some gene trees will have nodes representing speciation events

while other genes trees will have a node or nodes that represent IG. We define the 'true' species

branching order as the topology of the gene tree in which all nodes represent speciation events,

even if this history does not represent the majority of the genome. Our finding of massive

nuclear IG leads to a dilemma regarding which branching order should be used in future

comparative studies in this group. For many (if not most) practical purposes, it is reasonable to

continue to use A(BC) because it represents the history of most of the genome. However, studies

using this topology should bear in mind that this history is more complicated than simple

speciation and consider the potential implications. Integrating all available information into a

useful model for studying trait evolution represents a future goal in systematics.

We demonstrate the use of several complementary techniques to identify the taxa that

exchanged genes during IG, many of which operate in a four-taxon (or four-clade) context.

However, most methods do not explicitly distinguish which taxon served as donor and which as

recipient during IG. The existing methods that do polarize IG are only able to do so when there is

a fifth taxon (or clade)[11]. The divergence-based approach presented here can be applied to infer

the directionality of IG in a four-taxon case when additional taxa are not available. It should be

noted that our goal in the present study was to present the conceptual framework of divergence-

based polarization of IG and to lay the groundwork for further development of these types of

methods. It was not our goal, here, to mathematically derive the test or to explore parameter

robustness. For example, factors such as population size and structure, divergence time, size of

loci, rate of evolution[48], and extent of linkage disequilibrium[49] have been demonstrated to affect

existing statistics for inferring IG[9,45] but have not been explored here. We have also not explored

the power of our test to polarize IG when it is asymmetrical but not strictly unidirectional, all of

the above representing important next steps toward understanding the conditions under which

divergence-based phylogenetic methods can accurately recover the direction of IG.

Applied to genomic data, our test infers IG of nuclear genes from clade B to clade C.

Since cytoplasmic inheritance is matrilineal in Brassicaceae, we conclude that clade C was the

maternal recipient of paternal clade B nuclear alleles. While we can only postulate about the

specific crosses and backcrosses that occurred during IG, it is likely that F1 hybrids arose from a

clade C maternal parent and clade B paternal parent. We find evidence that selection acted

during the backcrosses that followed, resulting in resistance of organelle interacting nuclear

genes to replacement by paternal alleles. Maternal nuclear alleles that function in chloroplasts or

mitochondria in fundamental processes were not replaced at the same rate as maternal alleles

localized to other areas of the cell or for other functions. These genes may constitute a core set

whose replacement by paternal alleles is deleterious. We also find evidence that selection acted

to maintain nuclear-nuclear interactions. In general, our results suggest that epistatic interactions

between genes exerted selective pressure that influenced which genes were displaced and which

were retained. Whether this type of selection drove the displacement or retention of entire

haplotype blocks via hitchhiking remains a future question.

In summary, our comparative genomic analyses are consistent with an evolutionary

history in which massive unidirectional nuclear IG, driven by selection and influenced by

linkage, underlie the original observation of "chloroplast capture." The species branching order

in this group is more accurately reflected by B(AC), and thus similar to the findings of [38], nuclear

IG obscured speciation such that the latter was only recoverable from extensive genomic data.

What makes IG here particularly interesting is that its impact on the genome is evident despite

the fact that it must have occurred prior to the radiation of clade A 13 – 9 million years ago[20,22].

Hence, it's likely that, as additional high-quality genomes become available, comparative

analyses will reveal histories that include nuclear IG, even when the genomes considered are

more distantly related. In short, our findings explore the genomic battle underlying chloroplast

capture to reveal an onslaught of alleles via unidirectional IG. A core set of nuclear genes

resisted displacement by exogenous alleles; purifying selection removed genotypes with

chimeric epistatic combinations that were deleterious, just as Bateson-Dobzhansky-Muller first

described[7,50]. Will other IG events reveal similar selective constraints as those we detail? If so, it

could point us toward key interactions between cytoplasmic and nuclear genomes that lead to

successful IG, thereby refining our understanding of the factors governing the movement of

genes among species.

Methods:

Phylogenomic pipeline

Clustering of putative orthologs. Coding sequences (CDS) for *Arabidopsis thaliana*, *A. lyrata*,

Capsella rubella, *C. grandiflora*, *Boechera stricta*, and *Eutrema salsugineum* were obtained

from *Phytozome*[16,25–27,32,51]; *Camelina sativa* and *Cardamine hirsuta* were obtained from *NCBI*

[28,31]. Datasets were processed to contain only the longest gene model when multiple isoforms

were annotated per locus. CDS were translated into amino acid (AA) sequences using the

standard codon table. The resulting whole proteome AA sequences for the eight species were

used as input to cluster orthologs via *Orthofinder* (version 1.1.4)[35] under default parameters (Fig.

S1a). Two different filtering strategies with varying stringency were applied to the resulting

clusters to yield two dataset partitions referred to as 'full single-copy dataset' and

'conservatively single-copy dataset'. Both filtering strategies are described below.

Full single-copy dataset filtering. The full single-copy dataset was identified by sorting

Orthofinder results to include only clusters that contained exactly one sequence per species,

except in the case of *C. sativa*. Clusters containing one to three sequences from *C. sativa* were

also retained as single-copy (Fig. S1b) because it is a hexaploid of relatively recent origin. Thus,

clusters with up to three *C. sativa* paralogs (*i.e.* homeologs) were retained, and we expected these

homeologs to form a clade under phylogenetic analysis (see **Multiple sequence alignment and**

gene tree inference of nuclear genes). Gene clusters that yielded trees deviating from this

expectation were omitted from further analysis. The full single-copy dataset also contains groups

classified as retained duplicates (Fig. S1c). Retained duplicate clusters contain exactly two

sequences per species (three to six in *C. sativa*). The *A. thaliana* sequences in each cluster

represent known homeologs from the *α* whole genome duplication that occurred at the base of

Brassicaceae[37], and thus is shared by all sampled species in this study. We retained only those

gene clusters that produced trees in which the paralogs formed reciprocally monophyletic clades

(Fig. S1c).

Conservative single-copy dataset filtering. We also used a more stringent set of criteria to

develop a conservatively single-copy dataset. For this dataset, we compared the results obtained

from *Orthofinder* with results from previously published assessments of plant single-copy or low

copy gene families[33,34]. The criteria and taxon sampling of our *Orthofinder* filtering and the

filtering strategies of the two previous analyses differed, meaning each analysis provides its own

level of stringency. Moreover, both previous analyses included *A. thaliana*, allowing for direct

comparison with our results. We filtered our clusters to include only those genes recovered by

both *Orthofinder* and in at least one published analysis. We refer to these as conservatively

single-copy. Conservatively single-copy genes plus the retained duplicates described above

constitute the conservatively single-copy dataset. CP and MT gene datasets were filtered using

the same criteria used to filter the full single-copy dataset.

Multiple sequence alignment and gene tree inference of nuclear genes. For single-copy

genes, we generated AA-guided multiple sequence alignment of CDS using the *MAFFT*

algorithm (version 6.850)[52], implemented using *ParaAT* (version 1.0)[53], under the default

settings for both. Multiple sequence alignments of CDS for each gene cluster were used to infer

maximum likelihood gene trees using *RAxML* (version 8)[54] under the general time reversible

model with gamma distributed rate heterogeneity. Support values for nodes were calculated from

100 bootstrap replicates using rapid bootstrapping.

Assembly and annotation of mitochondria and chloroplast genomes. Whole genome

sequence reads for *A. lyrata*, *B. stricta*, *C. rubella*, *C. grandiflora,* and *C. sativa* were acquired

from *NCBI's Sequence Read Archive* (SRA). The run IDs of SRA files used to assemble

organelle genomes for each species were: *A. lyrata* (DRR013373, DRR013372); *B. stricta*

(SRR3926938, SRR3926939); *C. rubella* (SRR065739, SRR065740); *C. grandiflora*

(ERR1769954, ERR1769955); *C. sativa* (SRR1171872, SRR1171873). Both SRAs for each

species were independently aligned to the *Arabidopsis thaliana* mitochondrial (MT) genome

(Ensembl 19) using *HiSat2*[55] with default settings for paired-end reads within *CyVerse's*

Discovery Environment[56]. 15-30X coverage was recovered for each alignment. Mapped read

alignment files were converted from BAM to SAM using *SAMtools*[57]. MT consensus sequences

were generated (base pair call agreement with 75% of all reads) from each alignment within

Geneious (version 7.0; Biomatters)[58]. Each MT consensus sequence was annotated based on the

A. thaliana MT genome annotation (Ensembl 19). CDSs were then extracted using gffread from

the *Cufflinks* package[59]. The same method was used to assemble the *B. stricta* CP genome. All

other chloroplast genome sequences were publicly available.

Multiple sequence alignment and tree inference from chloroplast and mitochondria

markers. Single-copy CP and MT genes were identified, aligned, and used to infer phylogeny as

described previously for nuclear genes. Individual gene tree results are presented in Fig. S2d-e.

We also generated concatenated alignments for both the CP and MT genes using

SequenceMatrix[60]. We inferred trees (Fig. 1a-b) from both concatenated alignments using

RAxML with the same parameters described above.

Downstream analyses

Gene tree topology analysis. Tree sorting was performed in batch using the *R* packages, *Ape*[61],

Phangorn[62], and *Phytools*[63]. Gene trees from the retained duplicates were midpoint rooted and

split at the root into two subtrees, each of which contained a sequence from all eight analyzed

species. Subtrees were analyzed as individual trees alongside all other single-copy gene families

as described below. First, each gene tree was rooted at *E. salsugineum*. Next trees were sorted by

considering the topological arrangement of the A, B, and C lineages. For example, a tree was

categorized A(BC) if *B. stricta*, *C. rubella*, *C. grandiflora*, and *C. sativa* formed a monophyletic

clade. Thus, the branch in the tree leading to the monophyletic clade (the branch uniting *B.*

stricta, *C. rubella*, *C. grandiflora*, and *C. sativa* in the above example) was considered the

topology-defining branch. Statistical support for any given tree was summarized as the bootstrap

value along the topology-defining branch.

Since the focus of our analysis was on topological incongruence of A, B, and C clades,

our topology assessment was not designed to detect topological arrangements within A, B, and C

clades or in other parts of the trees. If a gene cluster failed to form either a monophyletic A or C

clade following phylogenetic analysis, it was marked as 'other topology' and removed from

further downstream analysis. Exact topologies of all trees, including those recorded as 'other

topology', are summarized in Table S2.

Applying D, F, and D_{GT} statistics to assess the effects of incomplete lineage sorting and

introgression. To determine whether the observed gene tree incongruences could have been

caused primarily by incomplete lineage sorting (ILS), we calculated Patterson's D-statistic (D)

(also known as the ABBA-BABA or 4-taxon test)[5,9]. D is typically applied to whole genome

alignments of three in-group taxa and one out-group taxon. It is calculated by scanning the

alignment to identify site patterns consistent with two possible resolutions of ILS (ABBA and

BABA). Due to the relatively deep divergence and numerous chromosomal rearrangements

between genomes used here, it was not feasible to construct accurate whole genome alignments.

Instead, we identified ABBA and BABA site patterns within single-gene multiple sequence

alignments used to infer gene trees. We calculated D and F using the total number ABBA and

BABA sites from all nuclear gene alignments (or subsets of nuclear genes corresponding to

individual chromosomes or conservatively single-copy genes). We excluded *C. sativa* sequences

from this analysis due to the presence of multiple *C. sativa* paralogs in some trees. We

considered only biallelic sites in which the two outgroups, *E. salsugineum* and *C. hirsuta,* have

the same allele. We also required individual species within each clade to have the same allele.

For example, an ABBA site would be one in which *E. salsugineum, C. hirsuta, A. thaliana, A.*

lyrata, C. rubella, C. grandiflora, and *B. stricta* display T, T, G, G, G, G, and T, respectively.

Note that all members of clade A and C share the derived allele. An example of a BABA site

would be T, T, G, G, T, T, and G, respectively. In this case, members of clades A and B share the

derived allele. We also tallied AABB sites, (e.g. T, T, T, T, G, G, and G, respectively), in which

clades B and C share the derived allele, although AABB sites are not a component of D or F.

We calculated D and F according to the equations from[64]. All site counts and statistics are shown

in Table S3.

We also applied the rationale of D to gene tree topology counts by calculated a related

statistic, D_{GT}. We used gene tree topologies as proxies for site patterns. Since B(AC) and C(AB)

trees were closest in frequency in the nuclear genome, we asked whether their frequencies were

statistically significantly different using D_{GT}. B(AC) trees and C(AB) trees were treated as

ABBA and BABA sites, respectively, while A(BC) was treated as AABB. D_{GT} was then

calculated as follows:

`DGT = (∑(B(AC) trees) - ∑(C(AB) trees)) / (∑(B(AC) trees) + ∑(C(AB)`
473   `trees))`
474

We calculated D_{GT} for the set of all nuclear genes as well as for subsets of genes present

on each of *C. rubella's* nuclear chromosomes[36]. Results from all D_{GT} calculations are given in

Table S4.

Phylogenetic network reconstruction and introgression analysis. To evaluate the likelihood

that the observed incongruence was caused by IG, we also reconstructed maximum likelihood

phylogenetic networks using InferNetwork_ML in *PhyloNet* (version 3.6.1)[15]. We input all

nuclear gene trees (Fig. S1d, *Full single-copy genes* dataset) and implemented InferNetwork_ML

using the command '`InferNetwork_ML (all) h -n 100 -di -o -pl 8;`', where h is

the number of reticulations allowed in a given network. The method ignores gene tree branch

lengths, utilizing gene tree topologies alone to infer reticulation events. We performed separate

analyses using $h = 0$ (a tree), $h = 1$, and $h = 2$, outputting the 100 most likely trees/networks

(designated with –n) from each analysis. We followed the analysis strategies of[65], manually

inspecting networks to identify those with edges consistent with both the major nuclear topology

[A(B,C)] as well as the major CP and MT topology [B(A,C)] (Fig. S2l-o). Additionally, we

reported the most likely tree/network from each analysis (Fig. S2k, p-q). As an additional means

of asking whether ILS alone adequately explains incongruence, we performed Tree Incongruence

Checking in R (TICR)[14]. We used a population tree inferred from *PhyloNet* (h = 0) (Fig. S2j)

with a table of concordance factors for all quartets. We performed the *TICR* test as implemented

in the *R* package, *phylolm*[66], according to the methods outlined in:

https://github.com/crsl4/PhyloNetworks.jl/wiki/TICR-test:-tree-versus-network%3F.

Identification of introgressed topology and species branching order. In order to identify the

topology most likely to represent IG, we measured node depths on trees displaying the A(BC)

B(AC). As above, *C. sativa* sequences were not considered in order to avoid complications

associated with paralogous sequences. For each nuclear gene tree, we calculated pairwise

synonymous divergence (*dS*) between taxa on the tree using *PAML* (version 4.8)[67]. To infer the

pairwise distance between two clades on the tree, we took the average *dS* score between each

combination of taxa present in the two clades. For example, the depth of the node uniting clades

A and C on B(AC) trees would be the average of *dS*(*A. thaliana*, *C. rubella*), *dS*(*A. lyrata*, *C.*

rubella), *dS*(*A. thaliana*, *C. grandiflora*), and *dS*(*A. lyrata*, *C. grandiflora*). To calculate

normalized *dS,* each *dS* node depth (as described above) was divided by the average pairwise *dS*

of each ingroup species versus the outgroup, *C. hirsuta.*

We also calculated node depths from ultrametric gene trees. Before measuring node

depths, gene trees were smoothed to ultrametric trees using semiparametric penalized likelihood

rate smoothing[40]. We implemented the rate smoothing algorithm designated by the *chronopl*

function in the *Ape* package. We tested six values of the smoothing parameter (λ), which

controls the tradeoff between parametric and non-parametric formulation of rate smoothing, to

assess the sensitivity of node depths to different values of λ. We calculated node-depth on

ultrametric trees for nodes representing T_1 and T_2 on each given topology (Fig. S3a). We plotted

the frequency distributions of node depths (Fig, S3b) as well as descriptive statistics (Fig. S3c-t).

In order to account for intragenic recombination, we split each gene alignment into 200nt

alignments, the goal being to reduce the probability of recombination occurring in the middle of

our alignment. For each window, we calculated a distance matrix and inferred a neighbor joining

"window tree" using *Ape* in R[61]. We calculated the depth of the T_1 node for each window

displaying either A(BC) or B(AC) from the distance matrix by averaging the pairwise distance

values similar to our treatment of *dS* node depths above. We documented the number of

discordant windows in alignments for A(BC) (Fig. S4a) and B(AC) (Fig. S4b) trees and used

boxplots to compare distributions of A(BC) and B(AC) node depths (Fig. 2g and Fig. S4c).

Divergence-based polarization of introgression. For each nuclear gene tree from our

Brassicaceae dataset, we calculated pairwise synonymous divergence (*dS*) between taxa on the

tree using *PAML* (version 4.8)[67]. To infer the pairwise distance between two clades on the tree,

we took the average *dS* score between each combination of taxa present in the two clades. We

excluded *C. sativa* sequences from this analysis due to the presence of multiple *C. sativa*

paralogs in some trees. We define *dS* between clades B and C, clades A and C, and clades A and

B as *dS(B,C)*, *dS(A,C),* and *dS(A,B)*, respectively (Fig. 3 and Fig. S5). For example, to calculate

the distance between clade A and clade C (*dS(A,C)*) for a given tree, we used the following

equation:

dS(A,C) = (dS(A.thaliana,C. rubella) + dS(A.thaliana,C. grandiflora) +
536   dS(A.lyrata,C. rubella) + dS(A.lyrata,C.grandiflora)) / 4
537

We calculated *dS(X,Y)* for both the species branching order, B(AC), and the introgression

tree, A(BC) (*dS(X,Y)$_{SP}$* vs. *dS(X,Y)$_{IG}$,* respectively). Frequency distributions of each value were

determined.

GO category enrichment analysis. *Gene Ontology* (GO)[41] data for Arabidopsis were obtained

from *The Arabidopsis Information Resource* (www.arabidopsis.org)[16]. We determined the GO

terms associated with the Arabidopsis genes present in our full single-copy data set. For each GO

term, the percentage of B(AC) trees containing the GO term was compared to the percentage of

A(BC) trees containing it. Comparisons were quantified with an enrichment score (*E*). For

example, we used the following equation to ask if B(AC) or A(BC) topology genes are enriched

for CP localization:

550   *E* = ((% B(AC) trees that are CP localized) –
551   (% A(BC) trees that are CP localized)) /
552   (% B(AC) + A(BC) topology genes that are CP localized)
553

Positive *E* indicates enrichment for a given GO category among B(AC) trees, while negative *E*

indicates enrichment among A(BC) trees (Table S6).

Network analysis of protein-protein interactions. Experimentally curated protein-protein

interaction data for Arabidopsis were downloaded from *Arabidopsis thaliana Protein Interaction*

Network (AtPIN) (version 2.6.70)[42]. Interaction data were filtered to contain only genes included

in the full single-copy data set. An undirected interaction network was visualized and analyzed

using the *igraph* package (http://igraph.org) in *R*. Each node in the graph represents a single-

copy nuclear gene family while each edge in the graph indicates a physical interaction in

Arabidopsis. Nodes were colored by gene tree topology and diameter of nodes are proportional

to bootstrap support values for the gene tree (see Fig. S2a-c).

We asked if genes displaying the same topology are clustered with each other in the

network by calculating nominal assortativity[68]. Assortative mixing/clustering of gene tree

topology results across the network was quantified by the assortativity coefficient (*A)* of the

network. Positive *A* indicates clustering of genes with the same topology, while negative *A*

indicates over-dispersal. We calculated the observed *A* for our network as well as a null

distribution of *A* generated by randomly assigning a topology to nodes in 10,000 replicates of our

network.

Mapping of gene coordinates to *A. thaliana* and *C. rubella* nuclear genomes. Topology

results were mapped to the nuclear genome of *C. rubella* using the gene coordinates from the

GFF file associated with the genome assembly. Genome maps were visualized using the *R*

package, *Sushi*[69], made available through *Bioconductor*[70]. Colored horizontal lines indicate genes

displaying each topology. The length of each line represents the bootstrap support value found at

the topology-defining branch in the gene tree (see Fig. S2a-c).

Detection of linkage disequilibrium. Topology results mapped to the *C. rubella* genome were

used to ask if genes displaying the same topology are clustered together linearly along

chromosomes. We assessed the physical clustering of A(BC), B(AC), and C(AB) genes with two

measures: 1) number of instances in which genes with the same topology are located within 10kb

of each other (Fig. S6a), and 2) number of instances in which neighboring genes share topology,

regardless of distance (Fig. S6b). We established a null distribution for both measurements by

generating 10,000 maps of the *C. rubella* genome in which observed location of single-copy

genes and the overall gene tree frequencies were maintained, but the assignment of topologies to

genes was randomized across chromosomes. Measure 1 and measure 2 were calculated for each

of the 10,000 replicates to obtain null distributions.

Statistical Analyses

All statistical tests were performed in *R* (version 3.4). Below, we describe methods used

to assess the significance of our results. Our general strategy was to provide sufficient

information to enable readers to make their own interpretations of the data; toward that goal, we

have included Bonferroni corrected and uncorrected (raw) *p*-values for each experiment where

corrections could be applied (Tables S5 and Table S5 or within supplemental text). The

conclusions we draw are statistically robust, and thus are not affected by whether significance is

assessed by raw or Bonferroni corrected *p*-values. The fact that the majority of the *p*-values in

support of our conclusions are significant shows that we are not 'cherry picking'. Thus, our

results are unlikely to have been affected by type-one error that can be associated with multiple

tests. Therefore, in order to avoid inflation of type-two error, we report raw *p*-values in the main

body of the manuscript.

***D, F,* and *D_{GT}*-statistics.** We calculated *D, F,* and *D_{GT}* for both the full single-copy and

conservatively single-copy data sets. Confidence intervals were obtained by resampling either

dataset to generate 10,000 bootstrap replicates, recalculating *$D/F/D_{GT}$* for each replicate. The

resulting distributions were compared using the Z-test. To account for potential autocorrelation

bias caused by non-independence of linked genes, $D/F/D_{GT}$ were also calculated using block

bootstrapping. For D and F, block bootstrapping was achieved by simply bootstrap resampling

from the available gene alignments and recalculating D/F with each replicate. For D_{GT} block

bootstrapping was accomplished by splitting the dataset into 100 equal size blocks of

neighboring genes based on position along *C. rubella* chromosomes. Blocks were then bootstrap

resampled 10,000 times and D_{GT} was recalculated with each replicate to obtain a distribution. *P*-

values from analyses of the whole genome were Bonferroni adjusted for four comparisons for

D_{GT}.

Phylogenetic network reconstruction and introgression analysis. *PhyloNet* models were

statistically compared by calculating AIC and BIC scores for each tree/network with the following

expressions:

620
621   `AIC = 2k – 2(log L)`
622
623   `BIC =  (log(n) *  k) – 2(log L)`
624

where k is the number of free parameters in the model, n is the number of input gene trees, and L

is the maximum likelihood value of the model. We compared hypotheses by calculated

difference in AIC and BIC scores for each given tree/network relative to the most likely network

(ΔAIC and ΔBIC).

Node depth based test of species branching order. Frequency distributions of node depths

were plotted. Two-tailed *T*-tests and Wilcoxon rank sum tests were performed to assess

differences in distribution means and medians, respectively. *P*-values were Bonferroni corrected

for six comparisons.

Divergence based test of IG directionality. Frequency distributions of node depths were

plotted. Two-tailed Wilcoxon rank sum tests were performed to assess differences in distribution

medians. *P*-values were Bonferroni corrected for three comparisons.

GO category enrichment. Enrichment of GO categories was assessed by comparing GO

categories of A(BC) genes versus B(AC) genes. For each GO category, two-by-two contingency

tables were constructed and used to perform Fisher's exact tests. Results from two-tailed and

one-tailed tests are reported. *P*-values from primary comparisons were Bonferroni corrected for

three comparisons.

Protein-protein interaction network. Clustering in the interaction network was quantified with

an assortativity coefficient (A)[68]. To assess significance of the observed A, we randomly assigned

one of the three topologies (keeping the frequency of each topology the same as in the original

data set) to genes in 10,000 copies of the network. We computed A for each of the 10,000

networks to obtain a null distribution of A and used the null distribution to perform a two-tailed

Z-test.

Haplotype block linear clustering. We quantified linear clustering of topologies by counting

the number of occurrences of proximal and neighboring genes in the observed data. We assessed

the significance of the observed values by generating null distributions from 10,000 datasets in

which the topologies were randomized. We used the null distributions to perform two-tailed *Z*-

tests. *P*-values were Bonferroni corrected for six comparisons.

Data Availability:

Gene tree data are linked to the online version of the paper. Scripts and input files used to

perform analyses are available at:

https://github.com/EvanForsythe/Brassicaceae_phylogenomics.

References:

1. Stebbins, G. L. The Significance of Hybridization for Plant Taxonomy and Evolution. *Taxon* **18,** 26–35 (1968).

2. Yakimowski, S. B. & Rieseberg, L. H. The role of homoploid hybridization in evolution: A century of studies synthesizing genetics and ecology. *Am. J. Bot.* **101,** 1247–1258 (2014).

3. Rieseberg, L. H., Whitton, J. & Linder, C. R. Molecular marker incongruence in plant hybrid zones and phylogenetic trees. *Acta Bot. Neerl.* **45,** 243–262 (1996).

4. Dasmahapatra, K. K. *et al.* Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. *Nature* **487,** 94–98 (2012).

5. Green, R. E. *et al.* A Draft Sequence of the Neandertal Genome. *Science (80-.).* **328,** 710–722 (2010).

6. Rieseberg, L. H. & Soltis, D. E. Phylogenetic consequences of cytoplasmic gene flow in plants. *Evol. trends Plants* **5,** 65–84 (1991).

7. Sloan, D. B., Havird, J. C. & Sharbrough, J. The on-again, off-again relationship between mitochondrial genomes and species boundaries. *Mol. Ecol.* **26,** 2212–2236 (2017).

8. Joly, S., McLenachan, P. A. & Lockhart, P. J. A Statistical Approach for Distinguishing Hybridization and Incomplete Lineage Sorting. *Am. Nat.* **174,** E54–E70 (2009).

9. Durand, E. Y., Patterson, N., Reich, D. & Slatkin, M. Testing for Ancient Admixture between Closely Related Populations. *Mol. Biol. Evol.* **28,** 2239–2252 (2011).

10. Stolzer, M. *et al.* Inferring duplications, losses, transfers and incomplete lineage sorting with nonbinary species trees. *Bioinformatics* **28,** i409–i415 (2012).

11. Pease, J. B. & Hahn, M. W. Detection and Polarization of Introgression in a Five-Taxon Phylogeny. *Syst. Biol.* **64,** 651–662 (2015).

12. Rosenzweig, B. K., Pease, J. B., Besansky, N. J. & Hahn, M. W. Powerful methods for detecting introgressed regions from population genomic data. *Mol. Ecol.* 2387–2397 (2016). doi:10.1111/mec.13610

13. Than, C., Ruths, D. & Nakhleh, L. PhyloNet: a software package for analyzing and reconstructing reticulate evolutionary relationships. *BMC Bioinformatics* **9,** 322 (2008).

14. Stenz, N. W. M., Larget, B., Baum, D. A. & Ané, C. Exploring tree-like and non-tree-like patterns using genome sequences: An example using the inbreeding plant species Arabidopsis thaliana (L.) heynh. *Syst. Biol.* **64,** 809–823 (2015).

15. Than, C., Ruths, D. & Nakhleh, L. PhyloNet: a software package for analyzing and reconstructing reticulate evolutionary relationships. *BMC Bioinformatics* **9,** 322 (2008).

16. Lamesch, P. *et al.* The Arabidopsis Information Resource (TAIR): Improved gene annotation and new tools. *Nucleic Acids Res.* **40,** 1202–1210 (2012).

17. Bailey, C. D. *et al.* Toward a global phylogeny of the Brassicaceae. *Mol. Biol. Evol.* **23,** 2142–2160 (2006).

18. Beilstein, M. A., Al-Shehbaz, I. A. & Kellogg, E. A. Brassicaceae phylogeny and trichome evolution. *Am. J. Bot.* **93,** 607–619 (2006).

19. Beilstein, M. A., Al-Shehbaz, I. A., Mathews, S. & Kellogg, E. A. Brassicaceae phylogeny inferred from phytochrome A and ndhF sequence data: tribes and trichomes revisited. *Am. J. Bot.* **95,** 1307–27 (2008).

20. Beilstein, M. A., Nagalingum, N. S., Clements, M. D., Manchester, S. R. & Mathews, S. Dated molecular phylogenies indicate a Miocene origin for Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **107,** 18724–18728 (2010).

709   21.   Couvreur, T. L. P. *et al.* Molecular phylogenetics, temporal diversification, and principles
710   of evolution in the mustard family (Brassicaceae). *Mol. Biol. Evol.* **27,** 55–71 (2010).
711   22.   Huang, C.-H. *et al.* Resolution of Brassicaceae Phylogeny Using Nuclear Genes Uncovers
712   Nested Radiations and Supports Convergent Morphological Evolution. *Mol. Biol. Evol.*
713   **33,** msv226- (2015).
714   23.   Oyama, R. K. *et al.* The shrunken genome of Arabidopsis thaliana. *Plant Syst. Evol.* **273,**
715   257–271 (2008).
716   24.   Al-Shehbaz, I. a. & O'Kane, S. L. Taxonomy and Phylogeny of Arabidopsis
717   (Brassicaceae). *Arab. B.* **6,** 1–22 (2002).
718   25.   Hu, T. T. *et al.* The Arabidopsis lyrata genome sequence and the basis of rapid genome
719   size change. **43,** 476–481 (2011).
720   26.   Lee, C.-R. *et al.* Young inversion with multiple linked QTLs under selection in a hybrid
721   zone. *Nat. Ecol. Evol.* **1,** 0119 (2017).
722   27.   Slotte, T. *et al.* The Capsella rubella genome and the genomic consequences of rapid
723   mating system evolution. *Nat. Genet.* **45,** 831–5 (2013).
724   28.   Kagale, S. *et al.* The emerging biofuel crop Camelina sativa retains a highly
725   undifferentiated hexaploid genome structure. *Nat. Commun.* **5,** 3706 (2014).
726   29.   Franzke, A., German, D., Al-Shehbaz, I. A. & Mummenhoff, K. Arabidopsis family ties:
727   molecular phylogeny and age estimates in Brassicaceae. *Taxon* **58,** 425–427 (2009).
728   30.   Koch, M., Haubold, B. & Mitchell-Olds, T. Molecular systematics of the brassicaceae:
729   Evidence from coding plastidic matK and nuclear Chs sequences. *Am. J. Bot.* **88,** 534–544
730   (2001).
731   31.   Gan, X. *et al.* The Cardamine hirsuta genome offers insight into the evolution of
732   morphological diversity. *Nat. Plants* **2,** 16167 (2016).
733   32.   Yang, R. *et al.* The Reference Genome of the Halophytic Plant Eutrema salsugineum.
734   *Front. Plant Sci.* **4,** 1–14 (2013).
735   33.   De Smet, R. *et al.* Convergent gene loss following gene and genome duplications creates
736   single-copy families in flowering plants. *Proc. Natl. Acad. Sci. U. S. A.* **110,** 2898–903
737   (2013).
738   34.   Duarte, J. M. *et al.* Identification of shared single copy nuclear genes in Arabidopsis,
739   Populus, Vitis and Oryza and their phylogenetic utility across various taxonomic levels.
740   *BMC Evol. Biol.* **10,** 61 (2010).
741   35.   Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
742   comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16,** 157
743   (2015).
744   36.   Schranz, M. E., Windsor, A. J., Song, B.-H., Lawton-Rauh, A. & Mitchell-Olds, T.
745   Comparative genetic mapping in Boechera stricta, a close relative of Arabidopsis. *Plant*
746   *Physiol.* **144,** 286–98 (2007).
747   37.   Bowers, J. L., Chapman, B. A., Rong, J. & Paterson, A. H. Unraveling angiosperms
748   genome evolution by phylogenitc analysis of chromosomal duplications events. *Nature*
749   **422,** 433–438 (2003).
750   38.   Fontaine, M. C. *et al.* Extensive introgression in a malaria vector species complex
751   revealed by phylogenomics. *Science (80-.).* **347,** 1258522–1258522 (2015).
752   39.   Lee-Yaw, J. A., Grassa, C. J., Joly, S., Andrew, R. L. & Rieseberg, L. H. An evaluation of
753   alternative explanations for widespread cytonuclear discordance in annual sunflowers
754   (Helianthus). *New Phytol.* (2018). doi:10.1111/nph.15386

40.   Sanderson, M. J. Estimating Absolute Rates of Molecular Evolution and Divergence
Times: A Penalized Likelihood Approach. *Mol. Biol. Evol.* **19,** 101–109 (2002).

41.   Blake, J. A. *et al.* Gene ontology consortium: Going forward. *Nucleic Acids Res.* **43,**
D1049–D1056 (2015).

42.   Brandão, M. M., Dantas, L. L. & Silva-Filho, M. C. AtPIN: Arabidopsis thaliana protein
interaction network. *BMC Bioinformatics* **10,** 454 (2009).

43.   Song, B. H. *et al.* Multilocus patterns of nucleotide diversity, population structure and
linkage disequilibrium in Boechera stricta, a wild relative of Arabidopsis. *Genetics* **181,**
1021–1033 (2009).

44.   Kim, S. *et al.* Recombination and linkage disequilibrium in Arabidopsis thaliana. *Nat.*
Genet. **39,** 1151–1155 (2007).

45.   Eriksson, A. & Manica, A. Effect of ancient population structure on the degree of
polymorphism shared between modern human populations and ancient hominins. *Proc.*
Natl. Acad. Sci. **109,** 13956–13960 (2012).

46.   Lohse, K. & Frantz, L. A. F. Neandertal admixture in eurasia confirmed by maximum-
likelihood analysis of three genomes. *Genetics* **196,** 1241–1251 (2014).

47.   Wen, D., Yu, Y., Hahn, M. W. & Nakhleh, L. Reticulate evolutionary history and
extensive introgression in mosquito species revealed by phylogenetic network analysis.
Mol. Ecol. **25,** 2361–2372 (2016).

48.   Martin, S. H., Davey, J. W. & Jiggins, C. D. Evaluating the use of ABBA-BABA statistics
to locate introgressed loci. *Mol. Biol. Evol.* **32,** 244–257 (2015).

49.   Plagnol, V. & Wall, J. D. Possible ancestral structure in human populations. *PLoS Genet.*
2, (2006).

50.   Orr, H. A. Dobzhansky, Bateson, and the genetics of speciation. *Genetics* **144,** 1331–1335
(1996).

51.   Goodstein, D. M. *et al.* Phytozome: A comparative platform for green plant genomics.
Nucleic Acids Res. **40,** 1178–1186 (2012).

52.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol. Biol. Evol.* **30,** 772–780 (2013).

53.   Zhang, Z. *et al.* ParaAT: A parallel tool for constructing multiple protein-coding DNA
alignments. *Biochem. Biophys. Res. Commun.* **419,** 779–781 (2012).

54.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30,** 1312–3 (2014).

55.   Kim, D. *et al.* Transcript-level expression analysis of RNA-seq experiments with HISAT,
StringTie and Ballgown. *Nat. Protoc.* **11,** 1650–1667 (2016).

56.   Merchant, N. *et al.* The iPlant Collaborative: Cyberinfrastructure for Enabling Data to
Discovery for the Life Sciences. *PLoS Biol.* **14,** 1–9 (2016).

57.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,**
2078–2079 (2009).

58.   Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop software platform
for the organization and analysis of sequence data. *Bioinformatics* **28,** 1647–1649 (2012).

59.   Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.*
28, 511–515 (2010).

60.   Vaidya, G., Lohman, D. J. & Meier, R. SequenceMatrix: Concatenation software for the
fast assembly of multi-gene datasets with character set and codon information. *Cladistics*

801   **27,** 171–180 (2011).

802 61. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of phylogenetics and evolution in R
803   language. *Bioinformatics* **20,** 289–290 (2004).

804 62. Schliep, K. P. phangorn: Phylogenetic analysis in R. *Bioinformatics* **27,** 592–593 (2011).

805 63. Revell, L. J. phytools: An R package for phylogenetic comparative biology (and other
806   things). *Methods Ecol. Evol.* **3,** 217–223 (2012).

807 64. Zheng, Y. & Janke, A. Gene flow analysis method, the D-statistic, is robust in a wide
808   parameter space. *BMC Bioinformatics* **19,** 1–19 (2018).

809 65. Wen, D., Yu, Y., Hahn, M. W. & Nakhleh, L. SOM: Reticulate evolutionary history and
810   extensive introgression in mosquito species revealed by phylogenetic network analysis.
811   *Mol. Ecol.* **25,** 2361–2372 (2016).

812 66. Tung Ho, L. S. & Ané, C. A linear-time algorithm for gaussian and non-gaussian trait
813   evolution models. *Syst. Biol.* **63,** 397–408 (2014).

814 67. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24,**
815   1586–91 (2007).

816 68. Newman, M. E. J. Mixing patterns in networks. *Phys. Rev. E* **67,** 026126 (2003).

817 69. Phanstiel, D. H., Boyle, A. P., Araya, C. L. & Snyder, M. P. Sushi.R: Flexible,
818   quantitative and integrative genomic visualizations for publication-quality multi-panel
819   figures. *Bioinformatics* **30,** 2808–2810 (2014).

820 70. Gentleman, R. *et al.* Bioconductor: open software development for computational biology
821   and bioinformatics. *Genome Biol.* **5,** R80 (2004).

822 71. Alexander, P. J., Windham, M. D., Govindarajulu, R., Al-Shehbaz, I. a. & Bailey, C. D.
823   Molecular Phylogenetics and Taxonomy of the Genus Boechera and Related Genera
824   (Brassicaceae: Boechereae). *Syst. Bot.* **35,** 559–577 (2010).

825 72. Bailey, C. D., Al-shehbaz, I. A. & Rajanikanth, G. Generic Limits in Tribe Halimolobeae
826   and Description of the New Genus Exhalimolobos (Brassicaceae). **32,** 140–156 (2017).

827 73. Slotte, T., Ceplitis, A., Neuffer, B., Hurka, H. & Lascoux, M. Intrageneric phylogeny of
828   Capsella (Brassicaceae) and the origin of the tetraploid C. bursa-pastoris based on
829   chloroplast and nuclear DNA sequences. *Am. J. Bot.* **93,** 1714–1724 (2006).

830 74. Galasso, I., Manca, A., Braglia, L., Ponzoni, E. & Breviario, D. Genomic fingerprinting of
831   *Camelina* species using cTBP as molecular marker. *Am. J. Plant Sci.* **6,** 1184–1200
832   (2015).

833 75. Copetti, D. *et al.* Extensive gene tree discordance and hemiplasy shaped the genomes of
834   North American columnar cacti. *Proc. Natl. Acad. Sci.* **114,** 201706367 (2017).

836 **Supplementary Information** is linked to the online version of the paper.

837 **Acknowledgments:** The data reported in this paper are provided in the Supplementary

838 Information. Scripts used to perform analyses are available at:

839 https://github.com/EvanForsythe/Brassicaceae_phylogenomics. This work was funded by NSF

840 grants 1409251, 1444490, and 1546825 to MAB. We thank M. J. Sanderson, M. M. McMahon,

841 E. Lyons, D.B. Sloan, M.P Simmons, R. N. Gutenkunst, A. E. Baniaga, and S. M. Lambert for

Author Contributions: E.S.F and M.A.B conceived the study. A.D.L.N performed organellar

genome assembly. E.S.F performed all other analyses. E.S.F and M.A.B wrote the manuscript

with input from A.D.L.N. All authors approved of manuscript before submission.

Author Information: The authors declare no competing financial interests. Correspondence and

requests for materials should be addressed to M.A.B. at mbeilstein@email.arizona.edu.

Figure 1 | Incongruent gene tree topologies are observed within and between nuclear and

organellar genomes. a. Chloroplast and **b.** mitochondria ML trees with branch support from 100

bootstrap replicates. Scale bars represent mean substitutions/site. **c-f.** ML gene tree topologies

inferred from nuclear single-copy genes rooted by *E. salsugineum.* **c.** A(BC), **d.** B(AC) and **e.**

C(AB) topologies. **f.** Numbers and frequencies of gene trees displaying A(BC) (orange), B(AC)

(green), and C(AB) (purple). Single-copy genes are shown categorized by dataset and by level of

bootstrap support.

859
Figure 2 | Node depths indicate extensive introgression led to transfer of nuclear genes. a.
Model depicting expected node depths (N.D.) for genes undergoing IG (left) or speciation
(right). Speciation history is represented by thick grey bars. Individual gene histories are
represented by black branches. Blunt ended branches represent a native allele that was replaced
by an IG allele. Vertical arrow indicates expected difference in node depth. **b-c.** The informative
node depths on A(BC) (**b**) and B(AC) (**c**) trees. **d-f.** Boxplots depicting observed median and
quartile node depths measured from dS (**d**), normalized dS (**e**), ultrametric gene trees (**f**), and
concordant windows within gene alignments (**g**).
868
869
870
871

Figure 3 | Unidirectional introgression led to transfer of nuclear genes from clade B to

clade C. a-b. Model depicting pairwise *dS* divergence between clades A, B, and C. Arrows point

to nodes on the species tree (B(AC)) and the IG tree (A(BC)) indicated with *SP* and *IG*

subscripts, respectively. Expected node depths under IG from clade B to clade C **(a)** or from

clade C to B **(b)**. Vertical arrows depict expected differences between gene trees representing

speciation and IG. **c.** Observed *dS* distances on speciation gene trees (green boxes; $dS(B,C)_{SP}$,

$dS(A,C)_{SP}$, and $dS(A,B)_{SP}$) and IG gene trees (orange boxes; $dS(B,C)_{IG}$, $dS(A,C)_{IG}$, and $dS(A,B)_{IG}$).

Arrows indicate observed differences between *SP* and *IG* comparing *dS(B,C), dS(A,C),* and

dS(A,B). Horizontal bars above boxes in **c** represent distribution comparisons. **p<0.01, *NS*

p>0.05.

884
Figure 4 | The genomic consequences of epistasis and genetic linkage during IG.
a. Enrichment (*E*) for GO terms = (% B(AC) genes – % A(BC) genes) / (% B(AC) + A(BC)
genes). **b.** Protein-protein interaction network for Arabidopsis protein complexes. Node fill, gene
tree topology; node diameters proportional to bootstrap support (Fig. S2a-c). **c.** Assortativity
coefficient (*A*) of the network. Null distribution of *A* (grey curve); dotted line, observed *A*. **d.**
Nuclear genes mapped to *C. rubella*. Vertical lines, genes (colored by topology). Line heights
proportional to bootstrap support (Fig. S2a-c). *******p*<0.01, *p<0.05.

892
893
894