Title: Improved management facilitates return of an iconic fish species

Authors: Brian R. MacKenzie[1*], Kim Aarestrup[2], Kim Birnie-Gauvin[2], Massimiliano Cardinale[3], Mads Christoffersen[1], Henrik S. Lund[4], Iñigo Onandia[5], Gemma Quilez-Badia[6], Mark R. Payne[1], Andreas Sundelöf[3], Claus Sørensen[7], Michele Casini[3]

Affiliations:

[1]National Institute for Aquatic Resources (DTU Aqua), Technical University of Denmark, Kemitorvet, Building 150, DK 2800 Kongens Lyngby, Denmark;

[2]National Institute for Aquatic Resources (DTU Aqua), Technical University of Denmark, Silkeborg, Denmark;

[3]Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of Marine Research, Turistgatan 5, Lysekil 45330, Sweden

[4]Danish Fishermen Producer Organisation, Nordensvej 3, Taulov, DK 7000 Fredericia, Denmark

[5]AZTI Tecnalia, Marine Research Division, Herrera Kaia Portualdeaz/g, 20110 Pasaia, BasqueCountry, Spain

[6]Catalan Association for Responsible Fishing, Barcelona, Catalonia, Spain

[7]Marinexperten, H.C.Ørstedsvej 9, DK 9900 Frederikshavn, Denmark

*Correspondence to: brm@aqua.dtu.dk.

Abstract:

Species declines and losses of biota are often associated with shifting baselines in perceived historical abundances, and/or neglect or abandonment of recovery actions aimed at ecological restoration. Such declines are frequently accompanied by contractions in the geographical distribution of the species, with associated negative ecological impacts and diminishing socio-economic benefits. Here we show using citizen science and other data that after 50-60 years of near total absence, the iconic top predator and highly migratory species bluefin tuna, *Thunnus thynnus*, returned by the hundreds if not thousands in waters near Denmark, Norway and Sweden

during August-October 2015-2017. The re-utilisation of this former habitat is part of a

geographically more widespread expansion of the summer foraging area to the northern part of

the northeast Atlantic Ocean, encompassing waters from east Greenland to west Sweden. The

remarkable return to the Skagerrak, Kattegat and North Sea has been facilitated by improved

fishery management for bluefin tuna and its prey. Bluefin tuna biomass in the northeast Atlantic

and Mediterranean has been increasing since a recovery plan was implemented in the late 2000s,

and biomasses of two key prey species (herring, *Clupea harengus*; mackerel, *Scomber scombrus*)

recovered during the late 1980s-1990s. The reappearance of bluefin tuna in the Skagerrak-

Kattegat and other waters of northern Europe, despite a recent history of mismanagement and

illegal fishing in the northeast Atlantic and Mediterranean which led to a critical population

decline, offers hope that other marine ecological recoveries are possible under improved

management of fisheries and ecosystems.

Short title: Return of an iconic fish species

One Sentence Summary: Improved management promotes the return of an ocean icon to

northern Europe.

Keywords: bluefin tuna, range, recovery, prey, management

Significance Statement:

Commercial fisheries are often perceived being in a state of decline and collapse, putting food

and economic security at risk. Such declines are frequently accompanied by contractions in stock

distribution, negative ecological impacts and diminishing socio-economic benefits. Here we

present an example based on one of the world's most valuable and iconic fish species, bluefin

tuna, which demonstrates that effective management of both bluefin tuna and its prey has been a

key factor leading to a remarkable reoccupation of formerly lost habitat. This reappearance,

following decades of absence, occurred despite the bluefin tuna stock having had a recent, long

history of unsustainable and illegal exploitation. Marine ecological recovery actions can be

successful, even in situations which may initially appear intractable.

Author Contributions: BRM designed research, compiled data and wrote the manuscript; all

authors contributed to the design of the study and/or edited drafts of the manuscript. KA, KBG,

MCH, and GQB assisted with data collection in Denmark and MCA, IO and AS assisted with

data collection in Sweden. CS and HSL assisted with data collection from recreational and

commercial fishers. MRP produced satellite imagery temperature products.

Introduction:

Recovering the biomass and spatial range of depleted fish stocks is challenging for fisheries

managers and conservation ecologists (1–5). Once lost, former biomasses and ranges often

disappear from human memory, thereby reducing motivation for and impairing recovery efforts

(6–8). Biomass and range recovery often take longer than anticipated, even when fishing is

reduced below sustainable levels, due to several factors such as bycatch fishing mortality (i. e.

captures as bycatch in other fisheries), and changes in stock productivity (1–4, 9, 10). Species

that are highly migratory and whose migrations take them into the high seas and multiple fishing

jurisdictions, such as tunas and billfishes, are potentially even more vulnerable to depletion and

prolonged recovery times than stocks under single or few fishery jurisdictions (2, 11).

Here we report a recovery of the former range of distribution by a highly migratory, highly

valuable, iconic top predator fish species (bluefin tuna, *Thunnus thynnus*). This species was the

target of unsustainable exploitation for many years in the 1990s-2000s, and its biomass declined

to record low levels in the late 2000s (12). As one of the world's most valuable fish species, it

has also been the subject of much scientific, public and conservation scrutiny (13).

Bluefin tuna spawn in sub-tropical regions (e.g., Mediterranean Sea) and then migrate north long

distances as adults to summer and autumn foraging areas (14, 15). In the northeast Atlantic,

historical adult foraging areas are located in the Bay of Biscay, on the northwest European

continental shelf off Ireland and the UK, and in the Norwegian Sea-North Sea-Skagerrak-

Kattegat-Øresund (the latter are hereafter referred to as northern European waters;

Supplementary Figure S1 showing ICES areas and sea names (14, 16)). Bluefin tuna occupy

these waters in summer-early autumn before migrating southwards to overwintering areas. This

long-distance migratory behaviour is a part of the species' life-history, having evolved through

generations (15).

However, the migrations to northern European waters stopped almost completely in the early-

mid-1960s and bluefin tuna have been extremely rare in the area since the 1970s (16, 17); the

species has not supported targeted commercial or recreational fisheries in the area since then (12)

(Supplementary Figure S2). The reasons for both the disappearance and the long period before

reappearance are unclear, but likely due to a combination of overexploitation of juveniles and

adults of both the tuna and their prey, and changing oceanographic conditions (16, 18–20).

Similar large changes in range of fish stocks have been reported in the past for some other fish

species as populations decline and expand (3, 4, 21, 22). Theoretical and conceptual models

attribute such changes to density – dependent effects on habitat utilisation, biological rates and

properties such as growth rates, habitat availability and cannibalism, as well as social

interactions and collective memory within populations (19, 21, 23).

In this report, we describe the re-appearance of bluefin tuna in northern European waters using

citizen science data (i. e., observations from non-scientists pursuing activities on or near the sea)

and our own observations, and we discuss possible reasons why the species has returned to this

region. Given the high level of illegal and unsustainable exploitation of this species in other parts

of its range in the recent past (mid-late 1990s until ca. 2008-2010; (12)), the recovery of the

habitat and range of this species is extraordinary and could become a classic example of how

recoveries can occur under a suitable combination of fishery management regulations and

ecosystem conditions.

Results:

Bluefin tuna observations in 2015-2017:

We found and received many reports of bluefin tuna in the region. The observations are

summarized in Figure 1 and Supplementary Table S2. The reports included observations of

single individual tunas and of schools of various sizes from a few specimens (2-10) to hundreds.

The species is relatively easy to identify and distinguish in this area, mainly because of their

surface jumping behaviour, body shape, color and size. The sizes of tuna observed were usually

large (ca. 2 – 3 m, corresponding approximately to ages 8+, based on length-age relationships for

this population) and the jumping or surface-breaking behaviour is characteristic of this species.

Most observations we present are based on individuals, which are partially or entirely out of the

water due to jumping and surface swimming, which facilitated reliable identification; the

observations are supported in many cases by photographs or videos available on public social

media and angling or news media websites in Scandinavia. Some of these photographs are

shown in Figure 2 and as Supplementary Figure S3, and some links to online videos of bluefin

tuna jumping are listed in Supplementary Table S3. Other observations occurred from a bluefin

tuna tagging study conducted in the area in September 2017 (24).

Our observations were made in 2015-2017. Most observations occurred in 2016 and 2017,

although we have records of sightings of 100s-1000s in 2015. The locations of most of the

sightings we obtained were in the central part of the eastern Skagerrak, along the Swedish west

coast in the Skagerrak, and in the Kattegat all the way down to the Øresund (Figure 1). For

example, recreational fishing tour boats, individual recreational fishermen and the Swedish Coast

Guard observed individual bluefin tuna swimming at the surface and jumping clear of the water

near a Swedish national marine park (Kosterhavet) in 2016. One bluefin tuna was caught by a

Danish recreational angler north of Skagen, Denmark in the Skagerrak on Sept. 19, 2016 and

released alive after capture. This fish was measured in the water to be 3.03 m and estimated to

weigh app. 400-450 kg (25).

At nearly the same time (Sept. 17, 2016) and ca. 7 deg. latitude farther north, bluefin tuna were

captured as part of a targeted commercial fishing operation in the Norwegian Sea, off Ona

(62.8603°N 6.5543°E), Møre, Norway, approximately halfway between Bergen and Trondheim.

Each of the tuna caught (N = 190) weighed ca. 170-300 kg (26–28). Several Danish commercial

fishermen, including one of the co-authors of this investigation (HSL), reported that they

repeatedly saw schools in a localized area north of Skagen on several days during ca. two weeks

in mid-late September, 2016; the total number of tunas observed on some days was in the

hundreds. During 2017, we had many more observations than in 2015 or 2016. Several of the

observations in 2017 were made as part of a bluefin tuna capture-tag-release project conducted in

the area where many bluefin tuna were seen in 2016. During the 8 fishing days, bluefin tuna

swimming at the surface or jumping were seen by anglers and/or scientists every day; several

anglers, many of whom are professional fishing guides or commercial fishermen for other

species, also reported seeing them at depth on their echosounders. Several bluefin tuna were also

captured and reported as bycatch by commercial fishing vessels in the Skagerrak-Kattegat in

August-September, 2017 (Figure 1).

These 2017 sightings and bycatches in the Skagerrak-Kattegat are supported by other presence

data from neighboring areas. These data include commercial catches by Norwegian vessels

along the Norwegian coast during August 2017 (29), a bycatch by a Danish fishing vessel 50-60

miles east of the Shetland Islands (North Sea) on Aug. 5-6, 2017 (30), and a stranded bluefin

tuna (2.2 m long) found on the Dutch coast on July 31, 2017 (31).

The aggregate seasonality of our presence data in the Skagerrak-Kattegat was from August 23

(2017) – Dec. 4. (2016). The cumulative amount of reports indicates that the species was present

in large numbers for at least 2 months during late summer-autumn.

Ecosystem conditions:

The longest available time series of potential prey biomasses are from ICES stock assessments

for herring and mackerel stocks in the North Sea, Norwegian Sea and northeast Atlantic. These

data show that the biomasses have been high since the late 1980s-early 1990s for these three

stocks in the northeast Atlantic Ocean. The sum of the three biomasses has been at record-high

levels since the early 2000s (Figure 3).

At the smaller spatial scale of the Skagerrak-Kattegat, demersal research surveys in late

summer-autumn show that catch rates (considered to be a relative indicator of biomass) for four

potential prey species (herring, sprat, mackerel, anchovy) were relatively low in 2016 and 2017.

The most abundant of these species is usually herring; however its abundance peaked in 2011

and has declined to low levels since then, including in 2015-2017. Other demersal and the

pelagic surveys also show that prey abundance in 2015-2017 was approximately average or even

below-average (Supplementary Figure S4).

August-October surface temperatures have been well above long-term average since 1994 when

a significant regime shift occurred (Figure 3; STARS test; $P < 0.0001$ (32)); this shift is evident

in the larger northeast Atlantic region and the Skagerrak-Kattegat sub-region where our tuna

sightings have been made. However temperatures during 2015-2017 were not unusually warm

compared to other years during the most recent regime.

Discussion:

Bluefin tuna appear to have re-entered former foraging habitat in northern European waters,

which they vacated about 50-60 years ago. The observations are identical to those reported in

historical fishery reports, newspapers and scientific literature from the 1920s-1960s (Figure 2),

when bluefin tuna were common in these waters (e. g., (33, 34)); the jumping and surface-

breaking swimming behaviour is typical for bluefin tuna foraging on prey species (33, 35). Our

observations indicate that bluefin tuna were abundant throughout the combined Skagerrak-

Kattegat and coastal Norwegian Sea region during especially 2016-2017. Since neither Denmark

nor Sweden has a fishing quota, and there are no surveys potentially monitoring the distribution

and abundance of bluefin tuna, the public observations are essential for providing evidence of

their return to these waters.

Our observations in the Skagerrak-Kattegat during these three years are supported by and

consistent with many other reports of bluefin tuna occurring in northern parts of the northeast

Atlantic since at least 2012. In that year, bluefin tuna were observed for the first time known to

science in waters off east central Greenland (Denmark Strait) as bycatch during mackerel fishing

(36). Bluefin tuna were also caught as bycatch in that area in subsequent years (37, 38). Bluefin

tuna have also occurred more frequently in waters near Iceland and Norway since the early-mid

2010s. The increased abundance in these waters has led to re-establishment of commercial

fishing quotas (2014 onwards) and catches for Iceland and Norway (39). For example, one

Norwegian fishing vessel captured 21 bluefin tunas as bycatch in September 2015 in the

Norwegian Sea (ca. 80 nautical miles NW of Ålesund) (40). We noted that during the same year

and month, large numbers of bluefin tuna were also seen in the Skagerrak-Kattegat by citizen

scientists, including one individual bluefin tuna found stranded on the Swedish west coast

(Figure 3). In 2013, one dead bluefin tuna (2.2 m long) was found on a Dutch beach (41); prior

to this stranding and the one in 2017 (31), no bluefin tuna had been reported stranded on Dutch

beaches since 1920 (41) when tunas were more common in Dutch coastal waters (42). The

collective volume of reports of bluefin tuna in the wider northern northeast Atlantic (i. e., from

Sweden westwards to east Greenland) indicates that the species has been present since at least

2012 onwards, with a documented re-appearance in the Skagerrak-Kattegat starting in 2015. The

return to this region, and more generally the northern part of the northeast Atlantic, is therefore

not likely a local, short-term anomaly, but more consistent with, and part of a widespread

increase in range since at least 2012. Below we discuss potential reasons for this change in

spatial distribution.

Cause of reappearance:

Given the long absence from the Skagerrak-Kattegat and neighboring waters, one must ask why

bluefin tuna has finally returned now. We believe that the factor which has likely contributed

most to the return is improved bluefin tuna fishery management since ca. the mid-late 2000s.

Several changes were made during this period, including reductions in quotas, increases in

minimum landing sizes from 6 to 10 kg and then to 30 kg so that a much larger share of juveniles

can now survive long enough to reach maturity (assumed to be at an age of 4 years, or ca. 25 kg

(12)), improved catch reporting requirements and documentation, a shorter fishing season during

the main spawning period in the Mediterranean Sea and strengthened fishery surveillance and

enforcement (12, 13).

Prior to implementation of these changes, the stock was overexploited both legally (because

ICCAT's contracting parties were adopting total allowable catches higher than those

recommended by ICCAT's scientific body [Scientific Committee on Research and Statistics,

SCRS] as being scientifically sustainable in the long term), and illegally (e.g., landings often

exceeded, sometimes even by 2-fold, the quotas adopted by the contracting parties during many

years in the late 1990s and early 2000s). Before the new regulations were implemented,

historical exploitation of juveniles in southern parts of the stock range was high since the 1950s

and has been considered to be a major factor leading to the disappearance and subsequent

continued absence of bluefin tuna from northern parts of the range (20). In addition, the parent

stock biomass in the early-mid-2000s was perceived to be declining (12) and at a rate that if

continued could have met criteria for listing this stock as "critically endangered" according to

IUCN (International Union for the Conservation of Nature) criteria (43).

Implementation of the new fishery management regulations, and a real and effective monitoring

control and surveillance system, appears to have had positive effects on the Mediterranean stock

of bluefin tuna. Shortly after, several stock indicators of abundance started increasing, including

the biomass of adults and the production rate of new young bluefin tuna "recruits" (12, 13).

Moreover standardized catch rates of adults and juveniles in several regions of the northeast

Atlantic and Mediterranean Sea by commercial fishing vessels have increased since the

implementation of these new regulations (39).

As the stock has increased, bluefin tuna appears to have expanded its migratory range, a pattern

common among recovering fish stocks (3, 22), to explore new feeding habitats and to reduce

density-dependent competition for prey, including into some northern areas beyond formerly

documented distribution ranges such as Denmark Strait (east of Greenland (36)). This

exploratory foraging behaviour may have led them to return to the northern European shelf

waters, including the Norwegian Sea, North Sea and Skagerrak-Kattegat where they apparently

have found sufficient prey for foraging.

A second reason for the return to these waters may be the relatively high biomasses of potential

energy-rich prey species such as mackerel, herring, and sprat. Herring and mackerel dominate

pelagic fish biomass in the northern part of the northeast Atlantic Ocean and their biomasses

have recovered to or beyond historical estimates (Figure 3)(44). Some of these species (herring

and mackerel) were also overexploited in the 1950s-1970s leading to local collapses and fishery

closures for these species. These declines may have been a factor inhibiting earlier return of

bluefin tuna to this region. However following implementation of more sustainable fishing

practices, the biomasses of these species recovered to long-term average levels by the early-mid

1990s and have remained high or increased further since then (Figure 3). However, the tunas did

not reappear in large numbers until the early-mid 2010s (i. e., ca. 10-15 years after the prey

biomass indicators recovered). This delay coincides with the period when bluefin tuna were

exploited at levels well beyond those considered sustainable and when overall biomass was

declining, leading to development and implementation of a recovery plan (12, 45, 46). The

delayed return, despite high prey biomass in the North and Norwegian Seas, suggests that the

main reason for the reappearance of bluefin tuna in the region was the increase in tuna biomass

itself, and the time required for individuals to re-learn former migration pathways and foraging

habitats (19, 47). It is possible therefore that if bluefin tuna had been more abundant in the 1990s

and 2000s, they would have already appeared in high abundance in the Norwegian-North Sea-

Skagerrak-Kattegat area several years earlier than now.

Ocean temperature conditions are also known to affect tuna distributions and migrations (18, 48,

49), and have been generally warmer since 1994. However temperatures in 2015-2017 were not

exceptionally warm and were in fact colder than in some earlier years in the recent warm regime

(post-1994). Notably, bluefin tuna have been present in the Skagerrak-Kattegat during many

earlier years (e. g., 1920s-60s) when temperatures were lower than during the post-1994 regime

(compare Figure 3 and Supplementary Figure S2), and have occupied colder areas farther north

in the past (e. g., in the Norwegian Sea and along the west Norwegian coast; see Ref.(34) for

temperature data). We conclude that temperatures in the Skagerrak-Kattegat or northeast Atlantic

region appear to have had little direct role on the re-appearance of bluefin tuna, although they

may have had indirect effects via changes in abundance, migratory behaviour or distribution of

local prey species. However the exact mechanisms by which temperature may have acted are

unclear and remain speculative.

In summary, both food and temperature conditions have been higher than average for many years

before the tuna returned to the Skagerrak-Kattegat and probably also the wider North Sea-

Norwegian Sea-Skagerrak-Kattegat region. We consider it unlikely that either of these factors in

2015-2017 were the main direct drivers for the recent appearance of bluefin tuna in the

Skagerrak-Kattegat region, although adequate prey and temperature conditions likely induced

exploratory foraging bluefin tuna to remain in and potentially return to this region, once it was

re-discovered. Consequently local stocks of prey species such as the western Baltic spring-

spawning herring, which migrate through the Skagerrak-Kattegat when bluefin tuna are

seasonally present, probably stimulate bluefin tuna to remain and forage in the region after they

have arrived and explored in the Norwegian-North Sea region.

The recent increase in overall bluefin tuna biomass in the northeast Atlantic-Mediterranean Sea

has probably led to an increase in exploratory and more distant foraging behaviour by some

individuals. Some of these individuals and schools of bluefin tuna have apparently encountered

the larger biomass of potential prey species, including mackerel and herring in the region, and

have in turn transferred or shared migratory and habitat knowledge with other individuals. These

mechanisms are known to be key elements of the discovery of new habitats and connectivity

among geographic regions in migratory populations (5, 19, 21, 23) and may have led to the

reappearance of bluefin tuna in the Skagerrak-Kattegat and more generally in the northern

northeast Atlantic, including towards east Greenland and Iceland (36), where mackerel

biomasses have been increasing (50, 51). New tagging studies in the region will help address

these mechanisms in future.

Although the mechanism by which bluefin tuna have learned to migrate to the Skagerrak-

Kattegat-North Sea-Norwegian Sea region is unclear,these newly-acquired feeding migrations

may have direct benefits for the population spawner biomass, egg production and overall

sustainability of the stock. Historical feeding in the North and Norwegian Seas allowed bluefin

tuna to rapidly increase their overall condition factors due mainly to consumption of energy-rich

prey species (17, 52). Fish in better condition typically have higher and better quality

reproductive output, potentially leading to higher recruitment per spawner and are less likely to

skip spawning (53). Development of new annual feeding migrations to this area may therefore

have benefits for population fecundity and recruitment.

Future perspectives:

The return of bluefin tuna to the Skagerrak-Kattegat in 2015-2017 seems to be part of a longer-

lasting seasonal re-occupation of these and other northern European waters. If this situation

continues in the future, it will open many new possibilities for both scientific understanding of

species biology and dynamics, and for socio-economy opportunities as for example sustainable

fisheries (commercial and recreational), catch and release angling, and eco-tourism. Regarding

science, a priority should be to investigate the migration behaviour and population origin of the

bluefin tunas which have appeared in these waters using a combination of modern tagging,

genetics, otolith and modelling methods. Bluefin tuna in the Atlantic are managed as two stocks

(western and eastern stocks, the latter including the Mediterranean) with separate quotas and

regulations (12). Historically, bluefin tuna caught in the northern European region had migrated

both from the northeast and northwest Atlantic (14), although the historical proportions are not

known with certainty. Any future commercial or recreational catches should be assigned to the

correct stock. Such assignment would need for example genetic or otolith-based evidence (54,

55). New sustainable commercial and recreational fisheries would support and diversify local

fishery economies in primarily rural areas of the region. Furthermore, given the recent interest in

the general public for the return of this species to northern European waters (56, 57) (e. g., 7

video clips made by citizen scientists have been viewed on social media > 510,000 times since

September 2015: Supplementary Table S3), and the possibility that jumping bluefin tuna can be

seen from relatively small boats operating within minutes to a few hours of shore, there is

potential that the species could create and contribute to the eco-tourism industry.

A pre-requisite for realizing these scientific and socio-economic opportunities is that the recent

fishery management regulations for both bluefin tuna and their prey, and their compliance,

continues in future. In general, it presently appears that efforts in the past decade towards

sustainable fishery management both for the bluefin tuna and its prey species are having positive

benefits for these stocks. Should this be true, bluefin tuna may become a regular summer

component of local fish communities and food webs (as it historically has been; (52)), and

contribute to small but lucrative commercial and recreational fisheries and eco-tourism

economies. If our working hypothesis that the reappearance is due mainly to recovered bluefin

tuna and prey stocks, then management actions which ensure a modest-high biomass of large

individuals in the tuna population will be needed.

We note that commercial fishing quotas for the NE Atlantic-Mediterranean stock will increase

gradually until 2020 by 13,295 t or 58 % relative to the 2017 quota (58). The population

abundance estimate used as basis for this increase has many uncertainties of unknown magnitude

(39, 59, 60). As a result, should the stock not be, or grow, as large as expected, then the higher

quotas could lead to a population decline. If this happens, then it is possible that the abundance

of the larger individuals containing the migratory memory and knowledge will decline; this

could lead to a retraction from the northern area again. We further note that the size range of

bluefin tunas seen in the area in 2015-2017 is relatively narrow (generally 2-3 m), suggesting

that the migrants which have arrived during 2015-2017 are from a relatively small number of

year-classes born in the early-mid 2000s (assuming a stock-specific length-at-age relationship

(61)). If the number of year classes observed in northern European waters is indeed small, and if

post mid-2000 year classes are weak-average, then the combination of ongoing natural mortality,

higher landings and fewer new recruits could lead to reduced immigration to northern European

waters in the future. Continued monitoring of stock size, distribution and habitat characteristics

(e. g, prey abundance and size), including tagging, is needed to improve our understanding of

factors affecting bluefin tuna ecology in the region.

The re-establishment of summer migration to the Skagerrak-Kattegat, together with the recent

increase in overall stock biomass, indicates that, as with some other recovering large iconic fish

stocks (3), improved management, enforcement and compliance can yield positive benefits when

ecosystem conditions for stock production are suitable (3, 9). These observations apply even for

species such as bluefin tuna which has historically suffered from much illegal and over-fishing

and whose highly migratory behaviour takes them into multiple fishing jurisdictions including

international waters. The reappearance of bluefin tuna in the Skagerrak-Kattegat and neighboring

seas (e. g., North Sea, Norwegian Sea) serves as an example of the benefits of implementing

effective recovery actions, despite a decades-long absence from the region and a highly

unsustainable fisheries exploitation situation. In this case, implementation has not been too late

to promote recovery. Similar efforts with other populations and species could also yield positive

outcomes. These findings, as well as some other recent and ongoing major marine recoveries (3,

62, 63), offer some optimism for the long-term recovery and sustainability of commercially-

exploited fish stocks, the ecosystems in which they live, and the economic sectors which they

(could) support.

Materials and Methods:

General:

As there are presently extremely limited, if any, commercial or recreational fisheries (i. e.,

Denmark and Sweden have no quota, but Norway does) for bluefin tuna in the Skagerrak-

Kattegat and no scientific fisheries (surveys) for this species in the region. Therefore the main

source of information available for documenting presence was reports from the public, including

commercial and recreational fisher observations. We consider all these records to be "citizen

science" and compiled observations from reports in newspapers, social media and via direct

contact of the public with us. Details for the data compilation are available in Supplementary

Material. We supplemented the citizen science observations with our own observations in 2017

as part of a dedicated tagging project in the area (24) (observational details in Supplementary

Material). All observations were then organized chronologically and visualized to display their

spatial-temporal distribution. To place our work in the historical context of past fisheries for

bluefin tuna in the region, we compiled and plotted officially reported landings data from ICES

and other historical sources (summarized in (34)) for the years 1903-2014, which was the last

year for which landings data are available from ICES. We use officially reported data from ICES

instead of from ICCAT because the former are available at higher spatial resolution for our

region of interest.

Ecosystem conditions (e. g., local food concentrations, temperature) are known to affect

distributions of bluefin tuna (18, 36). We derived estimates of the inter-annual variability in

abundance of major prey species and sea surface temperatures (i. e. those most likely

experienced by bluefin tuna during summer foraging on the continental shelf) for the northeast

Atlantic to evaluate whether food and/or temperature conditions were unusually favorable in

2015-2017 compared to earlier years. Further details of the data sources and compilation are

presented in Supplementary Material.

Acknowledgments: We thank Jeppe Olesen for GIS and mapping expertise and members of the

public without whom this investigation could not have been possible. The work was partly

supported by the Nordic Council of Ministers (project no. 141-2016-Tuna; NordTun), ICCAT

(Tagging programme on bluefin tuna GBYP Phase 7) and has been conducted using MyOcean

Products.

References:

1.   Worm B, et al. (2009) Rebuilding global fisheries. *Science (80-)* 325:578–585.

2.   Murawski SA (2010) Rebuilding depleted fish stocks: the good, the bad, and, mostly, the

ugly. *ICES J Mar Sci J du Cons* 67(9):1830–1840.

3.   Rose GA, Rowe S (2015) Northern cod comeback. *Can J Fish Aquat Sci* 72(12):1789–

1798.

4.   Cardinale M, et al. (2011) Historical spatial baselines in conservation and management of

marine resources. *Fish Fish* 12(3):289–298.

5.   Reuchlin-Hugenholtz E, Shackell NL, Hutchings JA (2015) The potential for spatial

distribution indices to signal thresholds in marine fish biomass. *PLoS One*

10(3):e0120500.

6.   Pauly D (1995) Anecdotes and shifting baseline syndrome of fisheries. *Trends Ecol Evol*

10:430.

7.   Jackson JBC, et al. (2001) Historical overfishing and the recent collapse of coastal

ecosystems. *Science (80-)* 293:629–638.

8.   Lotze HK (2014) Lessons from historical ecology and management. *The Sea; Vol. 16:*

Marine Ecosystem-Based Management (568 Pp.), eds Fogarty M, McCarthy JJ (Harvard

University Press, Cambridge, MA), pp 16–55.

9.   Britten GL, Dowd M, Kanary L, Worm B (2017) Extended fisheries recovery timelines in

a changing environment. *Nat Commun* 8:15325.

10.   Szuwalski CS, Vert-Pre KA, Punt AE, Branch TA, Hilborn R (2015) Examining common

assumptions about recruitment: a meta-analysis of recruitment dynamics for worldwide

marine fisheries. *Fish Fish* 16(4):633–648.

11.   Cullis-Suzuki S, Pauly D (2010) Failing the high seas: A global evaluation of regional

fisheries management organizations. *Mar Policy* 34(5):1036–1042.

12.   ICCAT (2014) *Report of the 2014 Atlantic bluefin tuna stock assessment session, Madrid,*

Spain, September 22-27, 2014 (www.iccat.org) (ICCAT, Madrid, Spain).

13.   Fromentin J-M, Bonhommeau S, Arrizabalaga H, Kell LT (2014) The spectre of

uncertainty in management of exploited fish stocks: The illustrative case of Atlantic

bluefin tuna. *Mar Policy* 47:8–14.

14.   Mather FJ, Mason JM, Jones AC (1995) Historical document: life history and fisheries of

Atlantic bluefin tuna. *NOAA Tech Memo NMFS-SEFSC* 370:1–165.

15.   Cury P, Anneville O, Bard FX, Fonteneau A, Roy C (1998) Obstinate north Atlantic

bluefin tuna (Thunnus thynnus thynnus): an evolutionary perspective to consider

spawning migration. *ICCAT CollVolSciPapers (Proc.of ICCAT Tuna Symp 1998, Part 1)*

50:239–247.

16.   Fromentin JM (2009) Lessons from the past: investigating historical data from bluefin

tuna fisheries. *Fish Fish* 10(2):197–216.

17.   Tiews K (1978) On the disappearance of bluefin tuna in the North Sea and its ecological

implications for herring and mackerel. *Rapp P-v Reun Cons.i nt Explor Mer* 172:301–309.

18.   Fromentin J-M, Reygondeau G, Bonhommeau S, Beaugrand G (2013) Oceanographic

changes and exploitation drive the spatio-temporal dynamics of Atlantic bluefin tuna

(Thunnus thynnus). *Fish Oceanogr* 23:147–156.

19.   De Luca G, Mariani P, MacKenzie BR, Marsili M (2014) Fishing out collective memory

of migratory schools. *J R Soc Interface* 11(95). doi:10.1098/rsif.2014.0043.

20.   Cort JL, Abaunza P (2015) The Fall of the Tuna Traps and the Collapse of the Atlantic

Bluefin Tuna, Thunnus thynnus (L.), Fisheries of Northern Europe from the 1960s. *Rev*

Fish Sci Aquac 23(4):346–373.

21.   MacCall A (1990) *Dynamic biogeography of fish populations* (Washington Sea Grant,

Seattle).

22.   Worm B, Tittensor DP (2011) Range contraction in large pelagic predators. *Proc Natl*

Acad Sci U S A 108(29):11942–11947.

23. Petitgas P, Secor DH, McQuinn I, Huse G, Lo N (2010) Stock collapses and their

recovery: mechanisms that establish and maintain life-cycle closure in space and time.

ICES J Mar Sci 67(9):1841–1848.

24. MacKenzie BR, et al. (2017) *Final project report: Electronic tagging of adult bluefin*

tunas by sport fishery or hand lines in the North Sea, off the coast of Sweden and/or other

countries. (ICCAT GBYP 04/2017 Tagging Programme 2017, Atlantic-wide Research

Programme on Bluefin Tuna (Kongens Lyngby) Available at:

http://www.iccat.int/GBYP/Docs/Tagging_Phase_7_Consortium_DTU.pdf.

25. Bursell J (2016) 400 kilo+ tun genudsat i dansk farvand. *Fisk og Fri*. Available at:

http://www.fiskogfri.dk/ [Accessed September 23, 2016].

26. Aarøy TA (2016) Fikk 190 makrellstørjer på en gang. *Sysla (online newspaper)*. Available

at: http://sysla.no/2016/09/17/havbruk/fikk-190-makrellstorjer-pa-en-gang_157057/.

27. Tangen M (2016) Hillersøy med storkast for makrelstørje (English: The fishing vessel

Hillersøy with a big catch of bluefin tuna). *Kystbloggen.com*. Available at:

http://www.webinntekt.com/blogg/?p=1534 [Accessed January 6, 2017].

28. Tangen Ø (2016) *Historisk storkast etter makrellstørje i 2016 (video)* (Norway) Available

at: https://www.youtube.com/watch?v=eXGZ4nUhk6w.

29. Bruland W (2017) – Då eg såg den i nota tok eg eit hopp

(https://www.nrk.no/sognogfjordane/_-da-eg-sag-den-i-nota-tok-eg-eit-hopp-1.13656987).

NRK (Norwegian State Radio Telev. Available at: https://www.nrk.no/sognogfjordane/_-

da-eg-sag-den-i-nota-tok-eg-eit-hopp-1.13656987 [Accessed May 15, 2018].

30. Aaberg M (2017) Kæmpetun sætter dobbelt-rekord: Gik til 125.000 kroner på auktion

(https://www.dr.dk/nyheder/regionale/midtvest/kaempetun-saetter-dobbelt-rekord-gik-til-

125000-kroner-paa-auktion). *Danmarks Radio Nyheder (Danish Natl Radio Telev News)*.

Available at: https://www.dr.dk/nyheder/regionale/midtvest/kaempetun-saetter-dobbelt-

rekord-gik-til-125000-kroner-paa-auktion [Accessed May 15, 2018].

31. Zeezoogdieren (2017) Dode tonijn bij Brouwersdam (Dead tuna at Brouwersdam,

Netherlands; https://www.zeezoogdieren.org/wordpress/2017/07/31/dode-tonijn-bij-

brouwersdam/). *Zeezoogdieren.org*. Available at:

https://www.zeezoogdieren.org/wordpress/2017/07/31/dode-tonijn-bij-brouwersdam/

[Accessed May 15, 2018].

32. Rodionov S, Overland J (2005) Application of a sequential regime shift detection method

to the Bering Sea ecosystem. *ICES J Mar Sci* 62(3):328–332.

33. Svendsen L (1949) *Tun: fiskeri af tunfisk* (J. Fr. Clausen, Copenhagen).

34. MacKenzie BR, Myers RA (2007) The development of the northern European fishery for

north Atlantic bluefin tuna (Thunnus thynnus) during 1900-1950. *Fish Res* 87:229–239

(doi:10.1016/j.fishres.2007.01.013).

35. Bauer R, Bonhommeau S, Brisset B, Fromentin J (2015) Aerial surveys to monitor bluefin

tuna abundance and track efficiency of management measures. *Mar Ecol Prog Ser*

534:221–234.

36. MacKenzie BR, Payne MR, Boje J, Hoyer JL, Siegstad H (2014) A cascade of warming

impacts brings bluefin tuna to Greenland waters. *Glob Chang Biol* 20(8):2484–2491.

37. Olsen JI (2014) Den største makrelfangst ved Grønland

(http://naalakkersuisut.gl/da/Naalakkersuisut/Nyheder/2014/08/290814-Den-stoerste-

makrelfangst-ved-Groenland). *Naalakkersuisut (Government of Greenland)*. Available at:

http://naalakkersuisut.gl/da/Naalakkersuisut/Nyheder/2014/08/290814-Den-stoerste-

makrelfangst-ved-Groenland [Accessed May 15, 2018].

38.   Sommer K (2015) Årets første tunfisk fanget (https://knr.gl/da/nyheder/%C3%A5rets-

f%C3%B8rste-tunfisk-fanget). *KNR (Kalaallit Nunaata Radioa;*

GreenlandicBroadcasting Corp. Available at: https://knr.gl/da/nyheder/årets-første-

tunfisk-fanget [Accessed May 15, 2018].

39.   ICCAT (2017) *Report of the 2017 ICCAT bluefin stock assessment meeting (Madrid,*

Spain 20-28 July, 2017) (Madrid, Spain).

40.   Aam S (2015) Fekk storfangst med makrellstørje (Sept. 17, 2015). *Sunnmørsposten*.

Available at: http://www.smp.no/nyheter/article11575190.ece.

41.   Zeezoogdieren (2013) Monstervis op strand Ritthem (Monster fish on Ritthem beach;

https://www.zeezoogdieren.org/wordpress/2013/10/22/monstervis-op-strand-ritthem/).

Zeezoogdieren.org. Available at:

https://www.zeezoogdieren.org/wordpress/2013/10/22/monstervis-op-strand-ritthem/

[Accessed May 15, 2018].

42.   Bennema FP (2018) Long-term occurrence of Atlantic bluefin tuna Thunnus thynnus in

the North Sea: contributions of non-fishery data to population studies. *Fish Res* 199:177–

185.

43.   MacKenzie BR, Mosegaard H, Rosenberg AA (2009) Impending collapse of bluefin tuna

in the northeast Atlantic and Mediterranean. *Conserv Lett* 2(1):26–35.

44.   ICES (2017) *Report of the ICES Advisory Committee. ICES Advice - Books 1-10.*
520   (Copenhagen, Denmark) Available at: www.ices.dk.

521   45.   ICCAT (2006) *Recommendation by ICCAT to establish a multi-annual recovery plan for*
522   *bluefin tuna in the eastern Atlantic and Mediterranean.*

523   46.   ICCAT (2010) *Report of the 2010 Atlantic bluefin tuna stock assessment session (Madrid,*
524   *Spain - September 6 to 12, 2010)* (ICCAT, Madrid).

525   47.   Mariani P, Křivan V, MacKenzie BR, Mullon C (2015) The migration game in habitat
526   network: the case of tuna. *Theor Ecol* 9(2):219–232.

527   48.   Dufour F, Arrizabalaga H, Irigoien X, Santiago J (2007) Climate impacts on albacore and
528   bluefin tunas migrations phenology and spatial distribution. *Prog Oceanogr* 86(1–2):283–
529   290.

530   49.   Druon J-N, et al. (2016) Habitat suitability of the Atlantic bluefin tuna by size class: an
531   ecological niche approach. *Prog Oceanogr* 142:30–46.

532   50.   Jansen T, et al. (2016) Ocean warming expands habitat of a rich natural resource and
533   benefits a national economy. *Ecol Appl* 26(7). doi:10.1002/eap.1384.

534   51.   Astthorsson OS, Valdimarsson H, Gudmundsdottir A, Oskarsson GJ (2012) Climate-
535   related variations in the occurrence and distribution of mackerel (Scomber scombrus) in
536   Icelandic waters. *ICES J Mar Sci* 69:1289–1297.

537   52.   Mariani P, Andersen KH, Lindegren M, Mackenzie BR (2017) Trophic impact of Atlantic
538   bluefin tuna migrations in the North Sea. *ICES J Mar Sci* 74(6):1552–1560.

539   53.   Marshall CT (2009) Implementing information on stock reproductive potential in fisheries
540   management: the motivation, challenges and opportunities. *Fish Reproductive Biology:*

Implications for Assessment and Management, eds T J, Fogarty MJ, Megrey BA,

Moksness E (Blackwell Publishing Ltd), pp 395–420.

54.   Kumar G, Kocour M (2015) Population Genetic Structure of Tunas Inferred from

Molecular Markers: A Review. *Rev Fish Sci Aquac* 23(1):72–89.

55.   Rooker J, et al. (2014) Crossing the line: migratory and homing behaviors of Atlantic

bluefin tuna. *Mar Ecol Prog Ser* 504:265–276.

56.   Christiansen M, Gyldenkræne P (2016) Nordjysk lystfisker jubler: der er blåfinnede tun

overalt. *Danmarks Radio Nyheder*. Available at:

http://www.dr.dk/nyheder/indland/nordjysk-lystfisker-jubler-der-er-blaafinnede-tun-

overalt [Accessed September 26, 2016].

57.   Fjeldseth Ø (2016) Vil ha sportsfiske etter størje. *NJFF Nyheter (Norges Jeger- og Fisk ;*

Nor Hunt Sportfish Assoc. Available at: https://www.njff.no/nyheter/Sider/NJFF-vil-ha-

sportsfiske-etter-storje.aspx.

58.   ICCAT (2018) *Report for the biennial period, 2016-2017, Part II (2017), Vol. 1,*

Proceedings of the 25th regular meeting of the Commission (Madrid, Spain).

59.   ICCAT (2017) *Addendum to the detailed report of the 2017 bluefin tuna stock assessment*

session: Updates to the bluefin tuna stock assessment models adopted during the 2017

bluefin tuna species group meeting (Madrid, Spain).

60.   Collette BB (2017) Bluefin tuna science remains vague. *Science (80-)* 358(6365):879.2-

880.

61.   Cort JL (1991) Age and growth of the bluefin tuna (Thunnus thynnus thynnus) of the

Northeast Atlantic. *ColVolSciPapICCAT* 35:213–230.

62. Lefcheck JS, et al. (2018) Long-term nutrient reductions lead to the unprecedented

recovery of a temperate coastal region. *Proc Natl Acad Sci* 115(14):3658–3662.

63. Reusch TBH, et al. (2018) The Baltic Sea as a time machine for the future coastal ocean.

Sci Adv 4(5):eaar8195.

64. Donlon CJ, et al. (2012) The Operational Sea Surface Temperature and Sea Ice Analysis

(OSTIA) system. *Remote Sens Environ* 116:140–158.

65. Rayner NA, et al. (2003) Global analyses of sea surface temperature, sea ice and night

marine air temperature since the late nineteenth century. *JGeophysRes* 108:4407–doi:

10.1029/2002JD002670 (updates available.

Figure legends (figures are located after legends):

Figure 1. Locations where individual or schools of bluefin tuna were observed in Skagerrak-

Kattegat-Øresund in 2015-2017. Colour contours: averaged sea surface temperature as derived

from satellite imagery (64) during Sept. 7-21 in each of the three years. Black contour lines:

bottom topography. During 2017, bluefin tuna were also captured for tagging purposes and in

commercial fisheries as bycatches. These catches are indicated with different symbols.

Figure 2. Photographic documentation of presence and jumping behaviour of bluefin tuna in the

Skagerrak and Kattegat in a historical period (1947 (33)) and in 2016 (four lower photographs).

The historical photographs illustrate the similarity of historical bluefin tuna size, shape and

behaviour with that observed in 2015-2017. Additional photographs and links to web-based

video clips on social media are available in Supplementary Figure S3 and Table S3. All images

are reproduced with permission from photographers.

Figure 3. Indicators of potential prey biomass and temperature conditions in the Skagerrak-

Kattegat and neighboring regions. A: interannual variability in spawning stock biomass of key

prey species (and their sum) for bluefin tuna in northern European continental shelf regions. The

stocks are autumn-spawning herring in the North Sea, spring-spawning herring in the

Norwegian-Barents Sea, and mackerel in the northeast Atlantic (44). B: mean ln CPUE (ln

kg/hour + 0.001; $\pm$ 2 x standard error) for herring, sprat, mackerel and anchovy in the Skagerrak-

Kattegat during August-September research vessel surveys (International Bottom Trawl Survey,

IBTS). C: inter-annual variability in late summer-autumn (mean of August, September and

October) sea surface temperature in the Skagerrak-Kattegat and a larger area of the northeast

Atlantic (20^{o} W $- 13^{o}$ E; $50^{o} - 66^{o}$ N) for 1870-2017 (black solid line with dots and black dashed

line with squares, respectively) and regime-specific mean temperatures (red: Skagerrak-Kattegat;

blue: northeast Atlantic) for different statistically significant regimes (32). Data source: Hadley

Climate Research Unit, UK (65)).

605   Figure 1.

608   Figure 2.

Figure 3.

Title: Improved management facilitates return of an iconic fish species

Authors: Brian R. MacKenzie[1*], Kim Aarestrup[2], Kim Birnie-Gauvin[2], Massimiliano
Cardinale[3], Mads Christoffersen[1], Henrik S. Lund[4], Iñigo Onandia[5], Gemma Quilez-Badia[6],
Mark R. Payne[1], Andreas Sundelöf[3], Claus Sørensen[7], Michele Casini[3]

Affiliations:

[1]National Institute for Aquatic Resources (DTU Aqua), Technical University of Denmark,
Kemitorvet, Building 150, DK 2800 Kongens Lyngby, Denmark;

[2]National Institute for Aquatic Resources (DTU Aqua), Technical University of Denmark,
Silkeborg, Denmark;

[3]Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of
Marine Research, Turistgatan 5, Lysekil 45330, Sweden

[4]Danish Fishermen Producer Organisation, Nordensvej 3, Taulov, DK 7000 Fredericia, Denmark

[5]AZTI Tecnalia, Marine Research Division, Herrera Kaia Portualdeaz/g, 20110 Pasaia,
BasqueCountry, Spain

[6]Catalan Association for Responsible Fishing, Barcelona, Catalonia, Spain

[7]Marinexperten, H.C.Ørstedsvej 9, DK 9900 Frederikshavn, Denmark

*Correspondence to: brm@aqua.dtu.dk.

20 Supplementary Materials:

22 Supplementary Text:
23

Methods:

This section provides details how bluefin tuna observations were compiled, how they should be
interpreted, and the sources and processing of ecosystem data (prey abundances, sea surface
temperatures)

Bluefin tuna observations:

We compiled observations of bluefin tuna from primarily public sources such as social media (i.
e., Facebook, YouTube) and websites representing both commercial fishermen and anglers in
Denmark and Sweden. We also obtained and used information sent to us by the public following
announcements on the DTU Aqua website in 2016 and 2017 and sent to Danish commercial and

sportsfishermens' organizations and contact by SLU Aqua with Swedish sportsfishermen that we
were interested in receiving sighting observations. We supplemented these citizen science
observations with reports of commercial catches and bycatches in the Norwegian Sea, North Sea,
Skagerrak, Kattegat and Øresund. The time period covered was August-October in 2016 and
2017. However during the course of this data collection, we received or found reports of
observations in 2015 which we used to support our overall results and conclusions. In many
cases, the sightings were supported by photographs or video recordings of bluefin tuna. The
information provided by members of the public included date and location of the observation,
how many bluefin tuna were observed (e. g., single individual, school, approximate number of
schools and number of fish per school), behaviour (jumping over water surface, breaking water
surface), and prey escape behaviours observed at the surface. The observations were entered into
a database and visualized geographically to illustrate their spatial distribution in relation to
distance from land, bottom topography and sea surface temperature. We are aware that the
citizen science data may not necessarily represent the true spatial distribution of the bluefin
tunas; details of the data collection and their interpretation are presented in Supplementary
Material.

We also made our own observations in 2017. These observations were gathered during a bluefin
tuna tagging study conducted in the eastern Skagerrak-Kattegat. Fishing operations were
conducted on 8 days in September, 2017 by ca. 30-40 angling boats. As part of the tagging
study, we requested anglers to record sightings of bluefin tuna jumping and swimming at the
surface. These sightings, along with the hooked, tagged and released, and hooked-escaped
bluefin tunas were used in our compilation. Full details of the tagging operation are described
elsewhere (1).

Contribution of citizen science to bluefin tuna ecology:

For reasons explained in the text, our investigation relies heavily on input from the public to
document the species presence in this region. As with all citizen science reports, there is a
possibility for some false, biased and otherwise incorrect reporting. We believe that such
records are not likely present in our compilation because of the nearly simultaneous nature of the
records over a wide area (e. g., the many sightings reported in the eastern Skagerrak-Kattegat on
nearly the same day in 2016 as a large commercial catch in central Norwegian coastal waters (2);
the sightings, capture, tagging and release of bluefin tunas in the eastern Skagerrak-Kattegat
during September 8-24, 2017 (1) and a commercial catch of 78 tunas in Norway in late August,
2017 (3)), the similarity of the reported behaviour to historical sightings of tuna in the region (e.
g .,(4); see Figure 2 of main manuscript and Supplementary Figure S3) and our own observations
during the 2017 tagging operation, and the distinguishable features of bluefin tuna behaviour and
size that reduce the likelihood of misidentification with other species.

Moreover, some of the citizen science reports were made by highly reputable observers,
including on-duty officers of national coast guard services or off-duty members of our research
vessels while participating in recreational sea-based activities. Their observations and reports
were identical to those made by other members of the public, by commercial and recreational
fishermen targeting other species and by ourselves during our 2017 tagging study. For
example, some fishermen observed tuna while trawling for pelagic fish (e. g., herring and
mackerel) that are prey for bluefin tuna in this region. Lastly, several of the reports and our
interviews via email or telephone include statements by the observers that they had never seen

such behaviour before despite years and even decades of activity on the sea and that they had
knowledge of the species' former presence in the area from older generations. We are confident
therefore that our observations represent a valid and reliable source of documentary evidence of
the presence of the species in these waters.

We are aware however that the reports based on citizen reporting reflect the spatial distribution
of where the reporting observers were located, and also our own efforts to obtain such
information, which started in 2016. As a result, such data do not necessarily represent the full
spatial distribution of where the tuna were located because (1) some tuna may have been
observed in other areas, but not reported to us, (2) some tuna may have been present in other
areas (and of course other depths), but not seen by any human observers; and (3) observers were
surely present over a much wider area than indicated by our few reports, but it is not possible to
know which of those observers saw or did not see bluefin tuna. The spatial and temporal
distribution of tuna based on our observations must therefore be interpreted cautiously, and we
cannot exclude the possibility that bluefin tuna were present over a much wider area or at other
times than is indicated by our data. We have tried to minimize such observer bias by making
broad contact to the public and especially commercial fishermen (e. g., via their associations).
Nevertheless, to obtain a more representative distribution in the area, alternative methods would
need to be employed such as acoustic surveys, aerial surveying via airplanes (5, 6) or with drones
(7) or tagging with electronic tags (8, 9) and subsequent modelling (10, 11). In addition,
increased public awareness of the species in the area and the need for its documentation could
also increase public reporting of bluefin tuna observations and the reliability of distributional
maps. Such combined survey- citizen science methods could also potentially be used to derive
estimates of relative abundance in the region, which is not possible with our dataset.

Estimates of abundance of potential prey for bluefin tuna:

We estimated abundances of potential prey for bluefin tuna in the region from regional stock
assessments for main prey species and from fishery research vessel surveys. We used the North
Sea herring, Norwegian spring-spawning herring and Northeast Atlantic mackerel total stock
biomass estimates from the ICES assessments as indicators of potential prey for bluefin tuna in
the region of our study. These stocks occupy large areas (see ICES stock management area map,
Supplementary Figure S1), which overlap with the historical distribution of bluefin tuna in the
region (12–15); moreover tuna which enter the Skagerrak and Kattegat historically passed
through the northern North Sea and Norwegian Sea on their way to this region (12, 14) and
would potentially encounter these prey during the migration and while foraging for prey. These
biomass estimates are based on stock assessments of the various stocks (16).

We also used scientific research vessel surveys to estimate prey abundances more locally in the
Skagerrak-Kattegat and where bluefin tuna were observed in 2015-2017. Hydro-acoustic and
bottom trawl surveys are conducted annually in the region as part of population status monitoring
in the region for fisheries management purposes (17–20). The surveys are conducted in
February-March (demersal survey in Kattegat-Belt Sea), late June-early July (hydro-acoustic
pelagic survey in Skagerrak-Kattegat), and August-September (demersal survey in Skagerrak-
Kattegat). The three surveys when considered in aggregate provide information about the
relative abundance of potential prey (herring, sprat, mackerel) among years. The demersal
survey in August-September is conducted when bluefin tuna were present in our area and we

present its results in the main article; survey estimates at other times of year are presented as
Supplementary Information.

The main characteristics of the surveys (depth sampled, geographic location, year and seasonal
coverage) are summarized in Supplementary Table S1. The notable feature for all the surveys is
that sampling methods and gear within each survey are the same throughout the time periods
shown here (17–20). As seen in the Table, only the acoustic survey is directly designed to
estimate abundance and biomass of pelagic fish species (e. g., herring, sprat); this survey is used
as input to ICES stock assessments for herring in the western Baltic Sea and in the North Sea
(17). The other two surveys are designed to capture demersal fish species as part of the
International Bottom Trawl Survey (IBTS) (21) and Baltic International Trawl Survey (BITS)
(18). However these surveys regularly capture pelagic fish species, including herring and sprat,
and these data can indicate relative trends and fluctuations in biomass, even though they may not
necessarily represent true abundances due to lower catchability for pelagic fishes which are
distributed higher in the water column than the demersal sampling gear. For the acoustic survey,
we use the total abundances of all size groups estimated on the survey in specific strata of the
survey (i. e., Kattegat, and waters along the Swedish west coast in the northeastern part of the
Skagerrak (19)). For the demersal surveys, we calculated annual geometric means of catch-per-
unit-effort (CPUE) using biomass/trawl hour as a relative biomass metric. Full details of the
survey methods and sampling gear are available in literature (17–20, 22).

Estimation of sea surface temperature:

Bluefin tuna occupy mainly surface waters (i. e., above the seasonal thermocline) when feeding
on continental shelves in summer as in the region of our study. This habitat is also the depth
layer predominantly occupied by their main prey (e. g., herring, sprat, and mackerel) in the
region. We assumed that sea surface temperature (SST) as estimated by satellite imagery is an
approximate indicator of the temperatures available for and experienced by bluefin tuna while
foraging in the region.

We calculated the average SST for the region for the months of August, September and October
for the time period 1870-2017 using a large international database of in situ and satellite-derived
observations(23) available online (http://wps-web1.ceda.ac.uk/ui/home) at 1 degree monthly
resolution. This time series allowed us to evaluate whether 2015-2017 were exceptionally warm
summers relative to historical variability and trends. We calculated mean temperatures for both
the Skagerrak-Kattegat (8^o – 13^o E; 55^o – 59^o N) and a larger area of the northeast Atlantic Ocean
(20^o W – 13^o E; 50^o – 66^o N) through which bluefin tuna migrate when entering the Norwegian-
North Sea-Skagerrak-Kattegat. We evaluated whether regime shifts in temperature occurred
using the STARS algorithm (24) for a significance level P = 0.05; settings used for testing were
Huber parameter = 1 and series length = 10. Higher resolution (0.05 degree daily) satellite-based
estimates of SST (25) in the Skagerrak and Kattegat region were averaged temporally over the
main period where tuna were observed (7th – 21st September during 2015-2017) and used to
characterize the thermal environment in which the fish were observed.

Supplementary references:

1. MacKenzie BR, et al. (2017) *Final project report: Electronic tagging of adult bluefin tunas by sport fishery or hand lines in the North Sea, off the coast of Sweden and/or other countries. (ICCAT GBYP 04/2017 Tagging Programme 2017, Atlantic wide Research Programme on Bluefin Tuna* (Kongens Lyngby) Available at: http://www.iccat.int/GBYP/Docs/Tagging_Phase_7_Consortium_DTU.pdf.

2. Martinussen TM (2016) "Hillersøy" fikk 191 størjer på ett kast. *Fiskeribladet.no*. Available at: https://fiskeribladet.no/nyheter/?artikkel=49030.

3. Bjerkeland Ø (2017) Tok 78 makrellstørjer i ett kast vest for Ona. *RBNETT.no*. Available at: http://www.rbnett.no/nyheter/2017/09/15/Tok-78-makrellstørjer-i-ett-kast-vest-for-Ona-15312201.ece.

4. Svendsen L (1949) *Tun: fiskeri af tunfisk* (J. Fr. Clausen, Copenhagen).

5. Lutcavage M, Kraus S, Hoggard W (1997) Aerial survey of giant bluefin tuna, Thunnus thynnus, in the great Bahama Bank, Straits of Florida, 1995. *Fish Bull* 95(2):300–310.

6. Bauer R, Bonhommeau S, Brisset B, Fromentin J (2015) Aerial surveys to monitor bluefin tuna abundance and track efficiency of management measures. *Mar Ecol Prog Ser* 534:221–234.

7. Kiszka J, Mourier J, Gastrich K, Heithaus M (2016) Using unmanned aerial vehicles (UAVs) to investigate shark and ray densities in a shallow coral lagoon. *Mar Ecol Prog Ser* 560:237–242.

8. Wilson SG, et al. (2015) Tracking the fidelity of Atlantic bluefin tuna released in Canadian waters to the Gulf of Mexico spawning grounds. *Can J Fish Aquat Sci* 72(11):1700–1717.

9. Richardson DE, et al. (2016) Discovery of a spawning ground reveals diverse migration strategies in Atlantic bluefin tuna (Thunnus thynnus). *Proc Natl Acad Sci U S A* 113(12):3299–304.

10. Chapman EW, Jorgensen C, Lutcavage ME (2011) Atlantic bluefin tuna (Thunnus thynnus): a state-dependent energy allocation model for growth, maturation, and reproductive investment. *Can J Fish Aquat Sci* 68(11):1934–1951.

11. Mariani P, Křivan V, MacKenzie BR, Mullon C (2015) The migration game in habitat network: the case of tuna. *Theor Ecol* 9(2):219–232.

12. Mather FJ, Mason JM, Jones AC (1995) Historical document: life history and fisheries of Atlantic bluefin tuna. *NOAA Tech Memo NMFS-SEFSC* 370:1–165.

13. Fromentin JM (2009) Lessons from the past: investigating historical data from bluefin tuna fisheries. *Fish Fish* 10(2):197–216.

14. Tiews K (1978) On the disappearance of bluefin tuna in the North Sea and its ecological

197   implications for herring and mackerel. *Rapp P-v Reun Cons.i nt Explor Mer* 172:301–309.

198   15.   MacKenzie BR, Myers RA (2007) The development of the northern European fishery for
199   north Atlantic bluefin tuna (Thunnus thynnus) during 1900-1950. *Fish Res* 87:229–239
200   (doi:10.1016/j.fishres.2007.01.013).

201   16.   ICES (2016) *Report of the ICES Advisory Committee. ICES Advice - Books 1-10.*
202   (Copenhagen, Denmark).

203   17.   ICES (2016) *First interim report of the ICES Working Group on International Pelagic*
204   *Surveys (WGIPS)* (Copenhagen) Available at: www.ices.dk.

205   18.   ICES (2016) Second interim report of the Baltic International Fish Survey Working Group
206   (WGBIFS), 30 March - 3 April, 2016, Rostock, Germany. *ICES C 2016/SSGIEOM 07*:1–
207   597.

208   19.   Stæhr K-J (2016) *Acoustic herring survey report for RV "Dana" 2nd June - 5th July,*
209   *2016* (Charlottenlund, Denmark).

210   20.   Eigaard OR, Støttrup JG, Hofmann E, Hovgård H (2003) *DFU's standardtrawl:*
211   *konstruction og sammenlignende fiskeri (DFU-rapport No. 126-03) (in Danish)*
212   (Charlottenlund, Denmark).

213   21.   ICES (2012) *Manual for the International Bottom Trawl Surveys. Series of ICES Survey*
214   *Protocols. SISP 1-IBTS VIII* (Copenhagen, Denmark) Available at: www.ices.dk.

215   22.   ICES (2013) *Manual for the Baltic International Trawl Surveys (BITS) - Addendum 1:*
216   *WGBIFS BITS Manual 2013* (Copenhagen, Denmark) Available at: www.ices.dk.

217   23.   Rayner NA, et al. (2003) Global analyses of sea surface temperature, sea ice and night
218   marine air temperature since the late nineteenth century. *JGeophysRes* 108:4407–doi:
219   10.1029/2002JD002670 (updates available.

220   24.   Rodionov S, Overland J (2005) Application of a sequential regime shift detection method
221   to the Bering Sea ecosystem. *ICES J Mar Sci* 62(3):328–332.

222   25.   Donlon CJ, et al. (2012) The Operational Sea Surface Temperature and Sea Ice Analysis
223   (OSTIA) system. *Remote Sens Environ* 116:140–158.

226   **Supplementary Figures:**

227   Figure S1. ICES fishery stock management areas.

Figure S2. Reported catches of bluefin tuna in ICES areas II, III and IV corresponding
approximately to respectively the Norwegian Sea, North Sea, and Skagerrak-Kattegat-Øresund
(see Figure S1 for map of ICES stock management areas). Catch data officially reported to ICES
from 1903-2014 are from ICES databases available online (www.ices.dk). Additional catch data
from before 1927 were compiled from historical fishery reports, catch records, museum records
and other documents as summarized in (15).

Figure S3. Photographs showing bluefin tuna in the Skagerrak-Kattegat during September 2016.
Photos provided with permission by members of the public. Collage 1: R. Waje, collage 2: H.
Karlsson, collage 3: J. Wolner.

Collage 1 – R. Waje.

Collage 2: H. Karlsson

Collage 3: J. Wolner

Figure S4. A: mean ln CPUE (ln kg/hour + 0.001; ± 2 x standard error) for herring, sprat and
anchovy in autumn (late October-early November) research trawl surveys in Kattegat-Øresund-
Belt Sea (north of 55 N.) during 1994-2017. B: estimates of prey biomass as derived from
research vessel hydro-acoustic surveys in the Kattegatduring 2008-2017 (see Table S1 for survey
details).

262
263

264
265 **Supplementary Tables:**

266 Table S1. Overview of research vessel surveys which provided estimates of biomass of potential
267 prey for bluefin tuna in the Skagerrak-Kattegat-Øresund. The surveys are part of international
268 bottom-trawl and acoustic surveys in the region designed and coordinated internationally by
269 ICES.

Characcteristic	Survey vessel		
	RV Havfisken (Denmark)	RV Dana (Denmark)	RV Argos (Sweden)
Sampling gear	Demersal trawl	Hydro-acoustics with pelagic and demersal calibration trawls	Demersal trawl
Year coverage	1994-2017 (October-November)	2008-2017	1992-2017 (August-September)
Location	Kattegat, Øresund, Belt Sea north of 55 N.	Skagerrak, Kattegat	Skagerrak, Kattegat
Target species	Demersal fish community	Herring and sprat	Demersal fish community

Table S2. Observational data for sightings, commercial bycatches, and research tag-releases of
bluefin tuna individuals and schools in the Skagerrak, Kattegat and Øresund during 2015-2017.
The data were recorded by members of the public including anglers ("citizen scientists"),
commercial fishermen and research scientists. .Observers' identities are known to the authors.
<mark>Table will be submitted to journal as an excel file supplement.</mark>

Table S3. Examples of videos on social media of bluefin tuna *Thunnus thynnus* swimming and
jumping at surface in the Skagerrak-Kattegat and off Norway during 2015-2017. Also indicated
is the number of views of each videoclip. The total number of views was 510,240 as of April
23, 2018.

Link	Location	Date recorded or uploaded	Videomaker	Comments and notes	Views as of July 12, 2017
https://www.youtube.com/watch?v=IKOF843Ew74	Skagerrak	Sept. 10, 2015 (recorded) Sept. 11, 2015 (uploaded)	Thomas Kolmorgen; Fiskeavisen.dk		26,166
https://www.youtube.com/watch?v=Rd9mRhHTpt0	Swedish west coast, near Måskeår	Sept. 15 (recorded), 2016	Micael Karlsson		46,027
https://www.youtube.com/watch?v=eXGZ4nUhk6w	Ona, Møre, Norway	Sept. 16-17, 2016 (recorded) Oct. 11, 2016 (uploaded)	Magnus Tangen	Short documentary (7:22) of commercial fishing on Norwegian vessel Hillersøy, which captured 190 tunas in one haul.	230,506
https://youtu.be/ESL0vS_NZRk	Skagerrak	Sept. 18, 2016 (recorded) Sept. 21, 2016 (uploaded)	Uffe Nielsen	Exact location not specified. Can see tuna in distance.	118,127
https://youtu.be/GSmDHQyDud8	West coast of Sweden	Uploaded Sept. 17, 2016	Bo Svensson	Very clear and sharp video; calm	6205

				water, easy to see tuna	
https://www.youtube.com/watch?v=WoadCif97qk	Hordaland, Norway	Sept. 10, 2017	Martin Wiech	Several large tuna visible jumping	3209
https://www.facebook.com/Oresundsakvariet/videos/1538892282843170/?q=%C3%B8resundsakvariet%20tun	Northern Øresund, Denmark	Sept. 22, 2017	Berit Lyrup	Very clear video of several large tuna jumping few metres from boat	80,000+

283 **Author Contributions**: BRM designed research, compiled data and wrote the manuscript; all
284 authors contributed to the design of the study and/or edited drafts of the manuscript. KA, KBG,
285 MCh, and GQB assisted with data collection in Denmark and MCa, IO and AS assisted with
286 data collection in Sweden. CS and HSL assisted with data collection from recreational and
287 commercial fishers. MRP produced satellite imagery temperature products.