1 **Title:** Networks of genetic similarity reveal non-neutral processes shape strain structure in
2 *Plasmodium falciparum*
3
4

5 Qixin He, Shai Pilosof, Kathryn E. Tiedje, Shazia Ruybal-Pesántez, Yael Artzy-Randrup,
6 Edward B. Baskerville, Karen P. Day & Mercedes Pascual
7
8 **Author information:**
9
10 **Affiliations**
11 Department of Ecology and Evolution, University of Chicago,
12 1101 E 57th Street, Chicago, IL, USA 60637
13 Qixin He, Shai Pilosof, Edward B. Baskerville, Mercedes Pascual
14

15 School of BioSciences, Bio21 Institute/University of Melbourne, Melbourne, Australia
16 30 Flemington Rd, Parkville, VIC3010, Australia
17 Kathryn E. Tiedje, Shazia Ruybal-Pesántez, Karen P. Day
18
19 IBED, University of Amsterdam
20 Postbus 94240, 1090 GE Amsterdam
21 Yael Artzy-Randrup
22
23 **Corresponding author**
24 Correspondence to Mercedes Pascual: pascualmm@uchicago.edu.
25

26 **Document order:**
27 Abstract, Introduction, Results, Discussion, methods, References, figure legends (5), Data
28 Availability, Acknowledgements

29   **Abstract**
30
Pathogens compete for hosts through patterns of cross-protection conferred by immune
responses to antigens. In *Plasmodium falciparum* malaria, the *var* multigene family encoding
for the major blood-stage antigen *Pf*EMP1 has evolved enormous genetic diversity through
ectopic recombination and mutation. With 50-60 *var* genes per genome, it is unclear whether
immune selection can act as a dominant force in structuring *var* repertoires of local
populations. The combinatorial complexity of the *var* system remains beyond the reach of
existing strain theory, and previous evidence for non-random structure cannot demonstrate
immune selection without comparison to neutral models. We develop two neutral models that
encompass malaria epidemiology but exclude competitive interactions between parasites.
These models, combined with networks of genetic similarity, reveal non-neutral strain
structure in both simulated systems and an extensively sampled population in Ghana. The
unique population structure we identify underlies the large transmission reservoir
characteristic of highly endemic regions in Africa.

Introduction

A central question in ecology and evolution regards the extent to which non-neutral processes structure diversity[1–4]. It remains a challenge to identify signature patterns that reveal an important role of ecological interactions in facilitating and stabilizing species coexistence in ecosystems with high diversity, such as tropical rain forests[5,6]. These ecological interactions specifically depend on trait differences between species, so that the assembly of diversity would not simply reflect stochastic colonization and extinction events of equivalent species as under neutrality. Here we address whether competitive interactions act as a non-neutral stabilizing force that promotes coexistence in another highly diverse system: *Plasmodium falciparum* populations as an ensemble of diverse strains in regions of high malaria transmission.

Recurrent malaria infections in endemic regions do not generate sterilizing immunity towards subsequent infection[7]; this suggests the existence of a large number of strains of the pathogen. A vast reservoir of *P. falciparum* exists in local human populations in Africa in the form of asymptomatic infections, hosts that carry the parasite without manifestation of the disease[8]. An understanding of the antigenic diversity of the parasite in such reservoirs, including whether and how this diversity is structured into strains, is fundamental to understanding immunity patterns and developing intervention strategies in the transmission dynamics of *P. falciparum* malaria.

The high transmission rates of endemic regions suggest frequency-dependent competition among parasites for hosts, through the cross-protection conferred by the adaptive immune system[9,10]. Since the success of an infection depends on the immunological memory of a host, new and rare antigenic types have a fitness advantage in the transmission system relative to common ones. In ecological theory, traits that confer such frequency-dependent advantage are known to promote coexistence via the formation of distinct niches, and to oppose in so doing the opposite, destabilizing, effect of absolute fitness differences in average growth rates[11,12]. The antigenic variation in pathogens maps conceptually to such stabilizing trait differences in ecological competition. Interestingly, "immune selection", a form of balancing selection, is already recognized as an important evolutionary force promoting the diversification and persistence of the *var* gene family, whose ancient origin predates the speciation of *P. falciparum*[13,14]. The role of immune selection is much less recognized and understood however for the faster time scales of ecology/epidemiology, and for the higher organizational levels of either the repertoires of *var* genes that constitute a parasite or the population structure of coexisting strains[15].

In high transmission regions, the extensive diversity of the *var* gene family[16] exhibits low amino acid similarity encoded by different *var* genes and a very low percentage of genes shared between parasites, both locally and regionally (e.g., less than 0.3% in Africa[15,17]). Previous work, known as strain theory, has posited that the non-random association of gene variants within genomes results from selection against recombinants through cross-immunity[18], akin to emergent niches of limiting similarity[19] or selection towards divergent local adaptations[20].

We are therefore interested in addressing whether signatures of immune selection can be detected at the repertoire level in such a diverse system. Existing models for strain theory[18,21,22] do not provide sufficient guidance on expected empirical patterns because they incorporate so far limited *var* gene diversity compared to observed numbers for *P.*

falciparum. This is the case for both earlier formulations with a few distinct loci[21,22] and a
recent extension for multi-copy gene families[23] that have predicted the emergence of
dominant strains with minimum genetic overlap. It is therefore unclear whether population
structure can emerge at realistic, high genetic diversity, especially under extensive
recombination rates. Existing strain theory also lacks a neutral counterpart, a neutral
hypothesis to disentangle, and statistically test, patterns generated by the acquisition of
specific immunity from those resulting simply from the basic demography of the
transmission system. In this study, we address these limitations by extending an individual-
based stochastic model to incorporate realistic mutation and recombination processes and
generate levels of diversity comparable to those of the *var* gene family in hyper-endemic
malaria regions. In addition, two process-based neutral models are formulated, which include
the same epidemiological and evolutionary processes of the full model, but replace specific
immunity by either: (i) no immune memory or (ii) generalized protection acquired via the
number of previous infections. Last, we propose the application of network properties to
identify selection signatures based on comparisons of *var* repertoire structures under immune
selection versus neutrality. We demonstrate that the structure of genetic similarity networks
contains clear signatures of neutral versus non-neutral processes, and that immune selection
plays an important role in shaping the empirical strain structure of *var* gene repertoires in a
local *P. falciparum* population from Bongo District (BD), Ghana. This structure differs from
what is expected on the basis of either previous strain theory[23] or recent ecological theory[24,25],
in which niche differentiation takes the form of clusters of strains or species with limited
overlap among them. An ensemble of network properties rather than a given clustering metric
is therefore needed. We discuss our results in the context of other attempts to identify niche
differentiation on the basis of genomic data in microbial ecology[26] and trait differences in
community ecology[24,25].

Results

Extended individual-based stochastic model of *var*

We extended the individual-based stochastic *var* model of Artzy-Randrup *et al.*[23] to incorporate more realistic mutation and recombination processes, which allows us to reach levels of diversity comparable to those of the *var* gene family (Fig. 1; Methods). In the model, each parasite *var* genome consists of a repertoire of 60 copies of *var* genes (Fig. 1a). Each *var* gene is considered a linear combination of two epitopes based on the empirical description of two hypervariable regions in the *var* tag region of the DBLα domain[27]. The transmission system is composed of a pool of these gene variants and a local human population open to immigrant parasites, in which we track transmission and infections. Simulations start with a static pool of *var* genes consisting of random combinations of the two epitopes (Fig. 1b). During transmission events, mitotic recombination and mutation generate new *var* genes, making the overall pool of epitopes effectively open to innovation, whereas meiotic recombination shuffles the composition of *var* genes of two or more repertoires in co-infections during the vector stage of transmission (Fig. 1c). The epitopes in the genes represent components of the *Pf*EMP1 molecule that are recognized and remembered by the immune system of the host; they are the 'traits' that effectively mediate competition for hosts at the population level.

In the immune selection model, individuals gain protection against specific epitope variants, through expression of the corresponding genes in an infection. Therefore, the rate at which hosts clear the infections increases with a higher number of specific epitopes seen from past infections. We then developed two neutral models and compared the repertoire structures they generate with those of the model with specific immunity. The first neutral model assumes generalized immunity in which protection is acquired as a function of the number of previous exposures irrespective of their specific antigenic identity; the second one is a completely neutral model in which infections propagate and recover, but hosts are blind to any history of exposure so that repertoires do not compete for hosts (Fig. 1d). These two neutral models include all the epidemiological processes, except for specific immunity towards the *var* genes that a host has been exposed to, and the resulting cross-protection. The epidemiological phenotype under immune selection is the duration of naïve infection (D). The parameter, D, directly influences the basic reproduction number (or fitness) of the parasite, R_0, and thus we match the infection period in the complete-neutrality model to the average duration that emerges in the corresponding immune selection model for each specific set of parameters (Fig. 1d; Methods). Similarly, we match the distribution of infection duration in the generalized immunity model to the emergent distribution of the full model. Therefore, repertoires under these neutral models do not exhibit fitness differences related to their specific genetic composition, while repertoires under the specific immunity model do differ in fitness as a result of the aggregated history of infection of the host population (Supplementary Fig. 1). The resulting Entomological Inoculation Rate (EIR), a measure of the force of infection experienced by individual human hosts measured as the number of infectious bites per host per year, is comparable among the three models. However, since the complete neutrality model does not include any mechanism to generate the typical empirical age infection distribution with higher prevalence of asymptomatic infection in children[8], prevalence in this model tends to be higher than that in the corresponding immune selection and generalized immunity models (Supplementary Fig. 2).

A network representation of var repertoire diversity

The reticulate evolutionary pattern of *var* genes, generated by frequent mitotic and meiotic
recombination within and between parasite genomes[28,29], respectively, precludes the
application of traditional population genetics tests for balancing selection (e.g., Tajima's *D*).
Hence, we develop an application of network theory to study the evolution of *var* repertoire
structures, and show that the structure of genetic similarity networks contains clear signatures
of neutral versus non-neutral processes. We analyzed the genetic structure of the parasite
population using networks in which nodes are *var* repertoires, weighted edges encode the
degree of overlap between the epitopes of these repertoires, and the direction of an edge
indicates whether one repertoire can outcompete the other (Methods). Comparisons of
structure across the similarity networks generated under the three models reveal distinctive
features of immune selection, although the specific features that distinguish immune selection
vary under different epidemiological settings (described below).
181
Because *var* genes exhibit different diversity levels across different endemic regions[17], we
investigated the influence of *var* gene pool size (i.e., the number of *var* genes in the global
pool; Fig. 1b) on the immune selection signatures. Since the two most relevant
epidemiological parameters, transmission intensity and duration of a naïve infection,
determine the intensity of competition among *var* repertoires, we vary them systematically to
address their influence on signatures of immune selection. Higher transmission and longer
duration of a naïve infection intensify competition among repertoires. They also increase the
rate of meiotic recombination among repertoires in mosquitoes. It follows that signature
patterns of immune selection should be most evident with increasing values of duration, gene
pool size and transmission, for conditions representative of high endemicity.
192
To explore selection signatures in networks generated under different intensity of competition
between strains, we use a suite of network metrics (see complete list in Supplementary Table
1, and Supplementary Fig. 3 for the low competition scenarios). If a process akin to limiting
similarity underlies population structure, networks are expected to be partitioned into
disconnected clusters of highly similar repertoires that occupy separate niches in
antigenic/genetic space. One way to quantify the partitioning of a network is by calculating
maximum modularity (Q)[30]. Under high competition, when the local *var* gene pool is of a
medium size characteristic of endemic regions of Asia/Pacific (~1200-2400 different *var*
genes)[31], the selection case differs notably from those of the two neutral models: repertoires
are typically grouped into well-defined modules (expressed as high Q and module F_{st} values,
Fig. 2a, b), whereas in networks resulting from the two neutral models, nodes are typically
connected to form star-shaped or tree-like structures. This qualitative difference in structure
resembles the prediction of non-overlapping strains in classic strain theory, where the
disconnected clusters are analogous to niches in immune space consisting of highly similar
repertoires.
208
In addition, because competition at the repertoire level in the selection case promotes
matching competitiveness of two given connected repertoires, it results in reciprocally-
connected directed edges of similar weights. In contrast, in the two neutral models,
repertoires with lower number of unique genes are not removed by selection, and when one
repertoire outcompetes another, there is only one directed edge between the pair. We use 3-
node motifs to capture this variation in competitiveness. For example, a binary in-tree motif
(A->B<-C) reflects that repertoire B is outcompeted by A and C, whereas a complete graph
motif in which three repertoires are all reciprocally connected (A<->B<->C<->A) indicates a
balanced, reciprocal competition. We find that networks of the selection model have a high
proportion of such reciprocal motifs compared to those of the two neutral models. Binary in-

tree or out-tree motifs are instead the most common in the neutral models, reflecting parent-
offspring evolutionary relationship, resulting from recombination where the recombinants are
not purged by immune selection (Fig. 2c, f).
222
Under a regime with a larger initial gene pool that matches the diversity levels of endemic
regions in Africa (~12,000-24,000 different *var* genes)[15], repertoires have a lower genetic
overlap compared to medium size gene pool (see Supplementary Fig. 4). This pattern follows
naturally from increased gene pool diversity because repertoires can be formed from a larger
number of gene combinations. Although such low overlap can indicate a non-random
structure[15], it cannot per se distinguish selection from the two neutral models. Accordingly,
module F_{sr} is low in all three cases and is not a good indicator of selection (Fig. 2e), despite
the selection model possessing more separate components than neutral models (Fig. 2d).
Nonetheless, networks generated with immune selection can still be differentiated from those
generated under neutral models using motif composition (Fig. 2f), as well as other network
metrics (see Fig. 3 and Supplementary Fig. 3). In particular, the weaker similarity between
repertoires entails a less clear network partitioning than that of lower gene pool diversity
systems. We therefore use betweenness centrality as a property reflecting their limiting
similarity: this metric measures the importance of a repertoire in a network by calculating the
proportion of shortest paths connecting any pair of nodes that go through it[32]. For the
networks generated with the neutral models, betweenness centrality varies little among
repertoires, with no highly central ones (Fig. 2d). This is because the persistence of each
repertoire is independent of the antigenic composition of other repertoires given the lack of
specific competition. By contrast, in the selection case some repertoires are clearly more
central than others (Fig. 2d), reflecting the non-random persistence of antigenic niches,
connected through these hubs via a series of recombination events.
244
Network classification at the repertoire and gene levels
To apply the findings from network theory to empirical data, we first asked whether networks
produced with the agent-based model can be classified into the processes that generate them
— immune selection vs. generalized immunity or complete neutrality— using an ensemble of
networks properties (Supplementary Table 1). Because competition is mainly among
repertoires that are highly similar, we inspected whether the quantile of edges ranked by
similarity values that are left in the networks influences the accuracy of classification. We
found that the power of classification (measured by the proportion of correct classifications
[true positives and true negatives]; Methods) remains largely unaltered as long as the bottom
20% in similarities of edges are removed (see supplementary Table 2). With a medium gene
pool size, there is a positive correlation between the transmission intensity and our ability to
classify networks correctly, reflecting an increasing divergence between the different kinds of
networks (Fig. 3a). With a high gene pool size, the classification always differentiates the
selection scenario from the two neutral ones correctly (Fig. 3c). The proportion of different
kinds of motif structures is the most powerful metric for the classification in high and
medium diversities. While metrics related to node degree contribute more in classification
under medium gene pool size, metrics based on reciprocity and distance are more important
for high gene pool size. Overall, a higher number of network features contributes evenly to
the classification under the high diversity scenario than under the medium diversity one (Fig.
3b, d).
265
Most endemic regions have seasonal transmission of malaria. Therefore, we repeated the
classification analysis using networks generated from the agent-based model that included a
seasonal transmission pattern with a high gene pool size (Methods). Here again, we found

that classification power remains largely unaltered as long as the bottom 20% of links are
removed. Unlike in non-seasonal networks, the classification can also differentiate
generalized immunity from complete neutrality scenarios (Supplementary Table 2), due to
the differences in the distribution of duration of infections between these two scenarios (see
Supplementary Fig. 1). Specifically, in the scenario of generalized immunity, early infections
in naïve hosts persist long enough to overcome the bottleneck in transmission during the dry
season, and are therefore carried on into the wet season. In contrast, infections under
complete neutrality have relatively short durations that limit persistence through the dry
season. Therefore, strain composition has a faster turnover rate under complete neutrality
than under generalized immunity. These different dynamics leave different signatures in the
networks, including motif compositions (Supplementary Fig. 5).
280
In addition to the effects of immune selection at the repertoire level, frequency-dependent
competition also works at the level of *var* genes in the population. Specifically, frequency-
dependent competition will limit the abundance of genes that are similar to many others in
their epitope composition — and are thus readily recognized by the immune system — while
favoring the abundance of genes with a unique composition of epitope alleles. We can test
this prediction using a network in which nodes are genes and edges encode similarity in
allelic composition (Fig. 4a). In the immune selection case, we find a characteristic negative
correlation between node degree (number of genes similar to a focal gene) and the frequency
of genes in the host population, because only genes that occupy a different niche can reach
high frequencies. This effect is absent in the neutral models (Fig. 4b-d).
291
Comparison with empirical data
Deep genetic sampling of local populations in BD, Ghana allows application of these
theoretical findings to examine the role of immune selection in nature. Gene similarity
networks were built from *var* DBLα domain tags sequenced from 1,248 *P. falciparum*
isolates in Ghana (Methods). An isolate refers to a complete sample of parasites from a host,
which may contain multiple infections (i.e., multiplicity of infection, or MOI>1). We restrict
our analyses to the upsB/upsC group of the DBLα domain because this subset is known to
exhibit frequent ectopic recombination within itself relative to the more conserved upsA
group[29]. This group is therefore less prone to generate the above negative correlation
spuriously out of differences in recombination rates, and provides a more appropriate
counterpart to our theory, which does not consider functional differences between *var* gene
variants. The resulting gene similarity network exhibits a strong negative correlation between
var DBLα type frequency and number of similar neighbors, providing evidence for
frequency-dependent competition (Fig. 5a).
306
We then examine the similarity network at the level of repertoires, by calculating shared
DBLα types between different repertoires in a subsample of 161 isolates whose MOI is equal
to one (i.e., whose *var* genes most likely compose a single repertoire; Methods;
Supplementary Fig. 5). We applied our network classification method to ask whether immune
selection played an important role in shaping empirical population structure. We generated a
library of simulated networks under parameter ranges representative of the sampling area. In
order to represent seasonality in the empirical data correctly, we ran the simulations using a
seasonal biting rate that is characteristic to the local area based on Appawu *et al.*[33] (Methods).
The resulting values of EIR [~25-170] encompass empirical observations for the region[33]. We
sampled 94 and 67 parasites every October and the next June respectively to construct the
similarity networks that represents the empirical sampling (Methods). Classification of the
empirical network obtained with the top 10% of edges in similarities, and therefore, the

potentially strongly interacting repertoires, indicates its resemblance to networks generated
with the immune selection scenario based on discriminant functions (Fig. 5b). This
classification result is robust up to a similarity strength cutoff of the top 25% of the edges.
Beyond this point, when weaker edges are considered, results assign the classification to
generalized immunity (Supplementary Fig. 6). The lower 75% edge weights in the empirical
networks represent, however, the sharing of at most one gene between pairs of repertoires,
which provides little information on strongly interacting repertoires. This is discussed below.
326
Discussion
The findings provide clear evidence for the role of frequency-dependent competition as a
result of immune selection in structuring antigen composition in a natural population of *P.*
falciparum. Network patterns of repertoire similarity differed significantly from those
obtained by the demography of transmission alone. The importance of immune selection at
the *var* gene level has been recognized before based on the ancient origin[13,14] and sequence
diversity of the gene family itself[16]. An agglutination experiment by Bull *et al.*[34] also showed
that children who have broader protection against *Pf*EMP1 variants tend to be infected by
only rare variants. Our results provide an additional piece of evidence of selection at the gene
level, in that common *var* gene variants can only persist if they are genetically distinct. That
immune selection operates at the level of the genes does not necessarily imply an effect at the
higher level of organization of repertoires or strains, especially given large gene pools and
high recombination rates. We have shown that under extreme genetic diversity and intense
recombination, immune selection profoundly structures repertoire diversity, and that this
structure can be detected and characterized using network metrics.
342
The network signatures we have identified present conceptual similarity to traditional tests of
balancing selection developed in population genetics or community ecology (as summarized
in Supplementary Table 3), thus filling the gap of available tests for highly recombinant gene
families. Specifically, a higher number of components/communities with even sizes is
observed under the immune selection scenario, in analogy to evenness measures in ecology
or Tajima's D in population genetics. Another key pattern is limiting similarity. It is akin to
high linkage disequilibrium (i.e., non-random association) among genes under balancing
selection, such as HLA genes[35]. In ecology, early competition models such as the standard
Lotka-Volterra equations showed that stable coexistence requires a given degree of niche
divergence[19]. The limit of coexistence is now understood to depend on both stabilizing
(frequency-dependent) niche differences and destabilizing (fixed) fitness differences in
average growth rates[11]. Moreover, more recent theory on niche differentiation has described
the emergence of clusters of similar species in trait space, where the limited overlap is now
between these groups and not individual species[24]. This pattern is a long-lived transient state
which can be stabilized when immigration[25] or evolution[24] are introduced explicitly. It is the
basis for the gap statistics method to distinguish simulated communities under niche
differentiation from null models based on two clustering measures, k-means dispersion[36] and
Ripley's K function[37]. Clustering on the basis of genomic data was also applied to this end in
microbiome communities[26] although limitations of the null model confound the results[38]. The
coexistence pattern of highly similar and divergent species is analogous to the population
structure we find in our model under a low to medium diversity of the gene pool, consistent
with previous studies[23], as well as the genetic patterns resulting from balancing selection in
evolutionary genetics (Supplementary Table 3). In that regime, here too, competition-driven
clustering is identified by our network metrics. In particular, modularity detects communities
that are more highly linked within than between; module F_{ST}, introduced here, quantifies
linkage based on genetic distinctiveness of a focal group weighted by its genetic diversity.

Importantly, clustering is no longer a characteristic property in our system at high diversity of
the gene pool, comparable to that observed in endemic regions. Thus, cluster measures alone
are unable to differentiate immune selection from neutral processes, and we suspect the same
will be the case in ecological systems if trait spaces underlying interactions are high-
dimensional[39]. Existing models in community ecology have considered so far only a low
number of dimensions.

For the *var* system, we have therefore relied on an ensemble of network properties. Motif
compositions provide a detailed profile of similarity patterns among triplets, which were
found to explain the largest proportion of the variance between the selection and neutral
models (Fig. 3). In addition, reciprocity in the network captures a unique signature of
balancing selection in multi-copy gene families, which does not have a counterpart in
ecology or population genetics measures. This is because the fitness of repertoires depends
on the number of unique genes under immune selection, given functional equivalence
between individual genes. Competition among genomes not only selects for those with
diverse genes, but also for the ones with more genes. Accordingly, genomes in West Africa,
where competition is intense maintain more than 50 unique *var* genes, whereas those in South
America, where transmission, and therefore competition, is low, are 10 *var* genes shorter[40].
More generally, these network properties can be applied to other multicopy gene repertoires
for antigenic variation, such as *vsg* genes in *Trypanosoma brucei* or *msg* genes in
Pneumocystis carinii[41].

Detecting signatures of selection requires a comparison to dynamic neutral models. The
observation of high modularity or low pairwise similarity among repertoires does not
guarantee the importance of immune selection, as low gene pool size or low transmission can
also produce similar patterns under neutral models (see network structures in Supplementary
Fig. 3). It is likely that generalized immunity, neutrality and immune selection are all at work,
either together or at different stages or genes. We have classified networks to these three
scenarios separately to investigate whether a distinct population structure exists for *var*
repertoires that reflects an important role of immune selection. In other pathogen systems, the
role of neutral vs. non-neutral forces were also investigated, which revealed unforeseen
patterns. For example, Nicoli *et al.*[42] found that strain replacement is more likely under
generalized immunity than strain-specific immunity when vaccination is applied in Pertussis.
Cobey & Lipsitch[43] demonstrated that weak specific immunity together with generalized
immunity permit the coexistence of strains with weak competence in *Streptococcus*
pneumoniae.

Another interesting feature of the *var* repertoire dynamics is the positive feedback between
increasing competition and the generation of inferior competitors. In endemic regions with
higher rates of transmission, hosts exhibit higher average MOI[44]. The rate of generation of
recombinant genomes is positively correlated with MOI because mosquitoes are more likely
to pick up multiple *var* repertoires from the host. Since unfit repertoires are generated at a
higher rate under higher transmission, immune selection leaves stronger signatures in high
transmission regimes, compared to lower transmission (Fig. 2,3). Conversely, ectopic
recombination can create new antigenic variants, conferring a fitness advantage to strains
with rare variants. Therefore, in areas of high transmission, there is a high generation of
allelic diversity, which is maintained by selection. These two forms of recombination, caused
by different molecular mechanisms, result in different directions of selection.

We have explored parameter space broadly in terms of transmission intensity and gene pool
size, and have also considered seasonal and non-seasonal transmission. Additional variations
of model structure will be addressed in future work. In particular, the rules that represent
within-host dynamics and regulate the expression of multiple infections and their interactions
with host immune responses, can be examined in more detail. The genetic multiplicity of *P.
falciparum* was found to potentially prolong asexual infections[45,46], probably due to cross-
reactive immune responses[47,48]. Although for our main results, we did not consider a cost
associated with higher MOI, the introduction of a more prolonged infection within multiply
infected hosts and across a range of parameters, indicated the robustness of the described
network structures.

Another extension of the model should relax the functional equivalence of all *var* genes[9,49–51].
In the empirical network analysis, when edges are included that represent the lower part of
the similarity distribution (the 75% weakest edge weights, corresponding to links between
repertoires pairs that only share one gene), the empirical pattern resembles that obtained
under generalized immunity (Supplementary Fig. 6). Since the majority of the one-gene
sharing edges involves the most common *var* types, the pattern could arise from these genes
having more critical functions than other *var* types, violating model assumptions of
functional equivalence. The pattern could also arise from differential selection pressure on
the genes. While our model assumes that parasites express all *var* genes in the repertoire
during the infection, ensuring equal selection pressure on all the genes, empirical experiments
have shown evidence of a structured switching pattern, in which slow switching *var* genes
express more prominently than fast switching ones, preventing exhaustion of the repertoire in
one infection[52–55]. Thus the activation hierarchy could bias *var* genes towards differential
strength of immune selection. Other open areas for further investigation of *var* evolution
include phenotypic mapping of sequence diversity to immunity, population structure over
time, and its influence on responses of the malaria system to interventions.

Early motivation for strain theory was the recognition that organization of *Pf*EMP1 variants
(and their underlying genes) into persistent largely non-overlapping sets can deeply alter our
understanding of epidemiology and control, for example by viewing *P. falciparum*'s apparent
large reproductive number (R_0) as resulting from a large ensemble of strains with much lower
reproductive numbers[21]. With the sheer number of existing and ever-changing variants and
repertoires, previous definitions of strains as long-lived entities do not apply at high
endemicity. The resulting population structure nevertheless exhibits limited similarity, in the
form of sparse small clusters and/or isolated individual repertoires interspersed into voids in
antigenic/genetic space, instead of well-defined niches. This emergent structure provides an
image of competition at the 'limit' of limiting similarity because of immense diversity. The
resulting coexistence and diversity at the different hierarchical levels of genes and repertoires
would enable the large reservoir of asymptomatic infections that makes malaria so resilient to
elimination in high transmission regions. As such, monitoring *var* gene diversity and
structure in responses to control efforts becomes central to understanding malaria
epidemiology and to creating a theory of control grounded in the reality of the complexity of
the system.

Methods
464
The extended *var* evolution model

Model parameters and symbols are summarized in Supplementary Table 4. The diversity of
var genes is represented at three organizational levels corresponding respectively to alleles
(epitopes), genes, and repertoires. Specifically, each parasite genome consists of *g var* genes
(in the main text, $g = 60$). The specific combination of the *var* genes is referred to as a *var*
repertoire throughout the paper. Each *var* gene is in turn a linear combination of *l* loci (in the
main text, $l = 2$) encoding epitopes that are connected linearly and each epitope can be
viewed as a multi-allele locus with *n* possible alleles. Immune selection in the model is a
result of specific immunity to epitope variants (alleles), which represent components of the
*Pf*EMP1 molecule that are recognized and remembered by the immune system of the host[56];
In effect, these epitopes serve as traits that mediate competition for hosts at the population
level, since individuals gain protection against specific alleles expressed by the parasite
during an infection (Fig. 1a).

Initiation of the simulation

To initiate the *var* gene pool *G*, a random allele for each epitope is chosen from the *n* alleles
to form each gene (Supplementary Table 4). In the simplest case, if there are two epitopes in
a *var*, then a particular *var* $g_i = \{L_{i1}, L_{i2}\}$, where L_{i1}, L_{i2} are random numbers from $U(1,n)$.
With n_i possible alleles at each epitope, the total number of possible genes is $\prod n_i$. However,
we chose *G* at least five times less than $\prod n_i$ so that not all combinations of alleles
constituting a gene would be available. This choice is based on the fact that not all
combinations of alleles form viable proteins. In the beginning of the simulation run, twenty
hosts were selected and infected with distinctive parasite genomes that consist of sets of *g var*
genes randomly selected from the pool *G*. The size of the host population, *H,* is kept at a
constant size (i.e., when a host dies, a new host is born). For the age structure in the
simulations, we fitted an exponential distribution to the reported Bongo population
demography and estimated an average life span of 30 years. Therefore, each host has a death
rate of $d = 1/30$ per year.

Repertoire transmission

Vectors (mosquitoes) are not explicitly modeled. Instead, we set a biting rate *b* so that the
average waiting time to the next biting event is equal to $1/(b*H)$. The force of infection is
kept the same across host age in the model. When a biting event occurs, two hosts are
randomly selected, one donor and one recipient. If the donor has infectious parasite
repertoires, the receiver is infected with a probability of *p* (i.e., transmission probability). If
the donor is infected with multiple strains in the blood stage, then the transmission
probability of each strain is *p* divided by the number of infectious repertoires. A delay is
applied between a transmission event and a repertoire becoming infectious to account for
stages of the life cycle in the vector and in the human host that are not modeled explicitly
(see below).

Meiotic recombination

Meiotic recombination occurs between strains in the sexual stage of the parasite's life cycle.
When multiple strains are transmitted to the donor, these strains have a $(1- P_r)$ probability to
remain as the original strain, and a P_r probability to become a recombinant strain, with P_r
calculated as follows,

$P_r = 1 - 1/N_s$ (1),

where N_s is the number of strains transmitted to the donor. Although the association of
physical locations and major groups of *var* genes is established, orthologous gene pairs
between two strains are often unknown. Therefore, we implement recombination between
strains as a process in which g genes are randomly selected out of all the original genes from
the two strains pooled together. Because physical locations of *var* genes can be mobile, this
assumption is a reasonable simplification of the meiotic recombination process.

Ectopic recombination within the strain in the asexual blood stage

Var genes often change their physical locations through ectopic recombination and gene
conversions. These processes occur at both sexual and asexual stages. However, ectopic
recombination is observed more often in the asexual stage where the parasites spend most of
their life cycle[6]. Therefore, we only model ectopic recombination among genes within the
same genome during the asexual stage. Two genes are first selected from the repertoire. Then
the location of the recombination breakpoint is randomly chosen, so that loci to the right of
the breakpoint are either swapped (recombination) or copied (gene conversion). In the current
implementation, we assume all breakpoints result into recombination rather than gene
conversion. Finally, newly recombined genes have a probability P_f to be functional (i.e.,
viable) defined by the similarity of the parental genes:

$P_f(x) = \tau^{\frac{x(\delta-x)}{\delta-1}}$ (2)

(Eq.3 in Drummond et al.[57]), where x is the number of mutations between the recombined
gene and one of the parental genes, δ is the genetic difference between the two parental genes
and τ is the recombinational tolerance. If the recombined gene is selected to be non-
functional, then the parental gene will be kept. Otherwise, the recombined gene will
substitute the parental gene so that a repertoire with a new gene is formed.

Mutation

Mutations occur at the level of epitopes. While infecting a host, each epitope has a rate of
mutation, μ, to mutate to a new allele so that n increases by one. New mutations can die from
lack of new transmissions, proliferate through new transmissions of the repertoire within
which they mutated, incorporate into other genes through ectopic recombination, or be
transferred to a different repertoire through meiotic recombination.
538
Within-host dynamics

Each strain is individually tracked through its entire life cycle, encompassing the liver stage,
asexual blood stage, and the transmission and sexual stages. Because we do not explicitly
model mosquitoes, we delay the expression of each strain in the receiver host by 14 days to
account for the time required for the sexual stage in the mosquito and the liver stage.
Specifically, the infection of the host is delayed 7 days to account for the time required for
gametocytes to develop into sporozoites within mosquitoes. When a host is infected, the
parasite remains in the liver stage for additional 7 days[58] before being released as merozoites
into the bloodstream, invading red blood cells and starting the expression of the *var*
repertoire. The expression of genes in the repertoire is sequential and the infection ends when
the whole repertoire is depleted. During the expression of the repertoire, the host is
considered infectious with the active strain. The expression order of the repertoire is
randomized for each infection, while the deactivation rates of the genes are controlled by the
host immunity. When one gene is actively expressed, host immunity 'checks' whether it has

seen any epitopes in the infection history. In the immune selection model, the deactivation
rate changes so that the duration of the active period of a gene is proportional to the number
of unseen epitopes. Duration of infection is not varied a priori as a function of age but is
instead determined by whether a given host has seen the particular strain in the past.
Therefore, the duration of infection of a particular repertoire in a particular host is,

$$Total\ duration = \frac{D}{g} \times \sum_{j=1}^{g} (No.\ new\ epitopes)/l \qquad (3)$$

After the gene is deactivated, the host adds the deactivated gene alleles to their immunity
memory. A new gene from the repertoire then becomes immediately active, and the strain is
cleared from the host when the whole repertoire of *var* genes is depleted. The immunity
towards a certain epitope wanes at a rate $w = 1/100$ per day[59]. In the current implementation,
we assume no cost associated with multiplicity of infection (MOI); that is, the duration of
infection is not correlated with the number of genomes a host is infected with (blue line). We
also explored another version of within-host dynamics, in which the duration of infection
increases as a function of MOI. The two scenarios of within-host dynamics produce
qualitatively similar network structures (see Supplementary Fig. 7).

Implementation of the simulation

The simulation is an individual-based, discrete-event, continuous-time stochastic model in
which all known possible future events are stored on a single event queue along with their
putative times, which may be at fixed times or drawn from a probability distribution. When
an event is triggered, it may trigger the addition or removal of future events on the queue, or
the modification of their rates, thus causing a recalculation of their putative time. This
approach is adapted from the next-reaction method[60], which optimizes the Gillespie first-
reaction method[61] by storing all events on an indexed binary heap. This data structure is
simple to implement and sufficiently fast and compact to store all events in the system, down
to individual state transitions for each infection course within each host. Specifically,
modifying the putative time for an event on the queue is O(log N), and heap storage is O(N),
where N is the number of events.

Statistical analyses

Selection versus neutral models

In order to disentangle signatures of immune selection from those of transmission per se in
parasite population structures, we designed neutral models in which hosts do not build
specific immunity towards alleles or genes, in addition to the selection model described
above. In the complete neutrality model (Fig. 1d), when hosts are infected, the duration of
infection is determined by the deactivation rate of each gene, which follows a Poisson
distribution of a constant rate; thus, hosts do not build immunity after an infection. The rate
of deactivation is calculated to match the average duration of infection of the corresponding
selection model, while maintaining the rest of the parameters (e.g., *G*, *b*) (Supplementary Fig.
1). In the generalized immunity model, the duration of infection decreases as the number of
past infections increases, similarly to the selection model. However, the identity of the alleles
does not play a role. We therefore match the average curve of duration of infection vs.
number of past infections to that of the corresponding selection scenario (See Fig. 1).

Diversity metrics, as well as epidemiological parameters, are calculated after each run to
compare between scenarios (see Supplementary Note 1 and Supplementary Fig. 8). Diversity
is quantified using common measures from ecology, including Shannon diversity[62] ($H =$

$-\sum_{i=1}^{S} p \ln(p_i)$), Simpson's diversity and evenness[63] ($D = \frac{1}{\sum_{i=1}^{S} p_i^2}$, $E = S/\sum_{i=1}^{S} p_i^2$), beta

diversity (i.e., turnover in composition of *var* genes or repertoires among parasite samples in

time; $\sum_i^{S} |x_{ij} - x_{ik}| / \sum_i^{S} |x_{ij} + x_{ik}|$), as well as within-repertoire diversity at the allelic and

genetic levels. Within-repertoire diversity is calculated as the number of unique alleles or

genes divided by the potential maximum number of unique alleles or genes (e.g., 60 genes

and 120 alleles if the genome size is 60 and the number of epitopes is 2). Entomological

inoculation rate (EIR), prevalence, and multiplicity of infection (MOI) are also compared

among model runs under different parameter settings and scenarios.

Building of similarity networks

In addition to diversity, similarity networks based on allelic composition at the gene or

repertoire levels are built to investigate parasite population structure. For this purpose, 150

parasites are sampled at 120-day intervals in the hosts, to subsample the simulations in a way

that is meaningful for later empirical application of network analyses. Directional similarity

networks for *var* genes or parasite genomes (i.e., *var* repertoires) are built with edges

encoding the proportion of shared unique alleles. Specifically,

$S_{ij} = \frac{a}{U_i} \,; S_{ji} = \frac{a}{U_j}$,

where *a* is the shared number of unique alleles (or genes) between repertoires *i* and *j*, and U_i

and U_j are the total number of unique alleles (or genes) in repertoires *i* and *j* respectively.

This directional index of genetic similarity is designed to represent the relative asymmetric

competition between two repertoires, as explained in Supplementary Fig. 9.

Calculation of Network properties

For the inspection of network structures from repertoires that have strong similarities, we

retained the top 1% of edges. 34 network properties are calculated to detect selection

signatures, as well as to distinguish these from patterns generated by pure transmission

dynamics or generalized immunity. These properties include metrics of transitivity, degree

distributions, component sizes, diameters, reciprocity, and proportion of 3-node graph motifs

(see Supplementary Table 1 for a complete list of properties and definitions). For inspection

of modular structures, an additional metric is introduced and named "module F_{ST}". This

metric quantifies to what extent the strain modules inferred from repertoire similarity

networks are genetically different from each other, by comparing the genetic diversity within

and between modules[64,65].

Simulations and machine learning algorithms for classification

For each combination of parameters (i.e., initial gene pool size *G*, biting rate *b*, and duration

of naïve infection *D*), 100 simulations were run to calculate the distribution of the network

properties under the scenarios of immune selection, complete neutrality and generalized

immunity. We investigated the accuracy of network classification under different quantiles of

retained edges ranked by similarities. We explored the range of retained links from 80% to

10%. The properties are then transformed into non-correlated principle components.

Discriminant analyses[66] are performed on the retained principle components that explain

more than 90% of the variance, to design functions that maximize the differences among

networks generated under different scenarios while minimizing the within-scenario variance.

The accuracy of the discriminant functions is assessed by the proportion of correct

classifications (i.e., power). Here, we use the lower two posterior probabilities of

classification assignment among the three scenarios as the false positive rates. A similar
approach is followed to build a classifier for empirical networks. Details are given below.

Comparisons with empirical data

Data sampling

The empirical data analyzed here was collected from a study performed across two catchment
areas in Bongo District (BD), Ghana located in the Upper East Region near the Burkina Faso
border. Malaria in BD is hyperendemic and is characterized by marked seasonal transmission
of *P. falciparum* during the wet season between June and October. This age-stratified serial
cross-sectional study was conducted over two sequential seasons. The first survey was
completed at the end of the wet season in October 2012, followed by a second survey at the
end of the dry season between mid-May and June 2013. Details on the study population, data
collection procedures and epidemiology have been published elsewhere[8]. Briefly, after
obtaining informed consent, finger prick blood samples were collected for parasitological
assessment for *P. falciparum* by blood smears, and dried blood spots for molecular
genotyping[8]. Sequencing was conducted for all isolates that are microscopically positive but
asymptomatic, with no strong bias in sequencing parasite genomes across host ages. The
study was reviewed and approved by the ethics committees at the Navrongo Health Research
Center, Ghana; the Noguchi Memorial Institute for Medical Research, Ghana; New York
University, United States; the University of Melbourne, Australia; and the University of
Chicago, United States.
660

PCR amplification and var DBLα sequence analysis

The DBLα domain of *P. falciparum var* genes were amplified from genomic DNA using
universal degenerate primers, as previously described[67]. Amplicons were pooled and
barcoded libraries were sequenced on an Illumina MiSeq sequencer using the 2x300 paired-
end cycle protocol, MiSeq Reagent kit v3 chemistry (NYUGTC, New York USA; AGRF,
Melbourne Australia). A custom pipeline was developed to de-multiplex and remove PCR
and sequencing artefacts from the DBLα sequence tags. Reads were demultiplexed into
individual fastq files for each isolate using flexbar v2.5[68] and paired based on valid
combinations of MID tags in the forward and reverse reads. A minimum read length of 100nt
and a maximum uncalled bases threshold of 15 were used. The resulting paired fastq files
were then merged using PEAR v.0.9.10[69] to ensure the resulting merged fastq files had
appropriate base quality scores allowing for filtering of low quality reads. The minimum
assembly length was set to 100nt and the minimum overlap required between a read pair was
set to 20nt. Low quality reads were filtered if they had more than one expected error using
the fastq_filter option of Usearch v8.1.1832[70]. Next, chimeras were filtered using Uchime
denovo and then the filtered reads were clustered using the cluster_fast function of Usearch
after the removal of singletons to reduce the impact of errors. A threshold of 96% identity[15]
was used to cluster the reads. To increase the overall quality of the sequences, the resulting
clusters were removed if they contained less than 15 reads to remove low support reads. The
representative read from each cluster was kept for the remaining stages of the pipeline. Next,
any non-DBLα sequences were filtered out with a domain score threshold of 80. Finally, as a
quality check the remaining reads were aligned to the reference *var* DBLα sequences of the
3D7, Dd2 and HB3 laboratory clones from experimental sequence data. To subsequently
determine DBLα types shared between isolates, the cleaned DBLα reads were clustered using
a pipeline based on the USEARCH software suite version 8.1.1831[70]. Initially duplicate reads
were removed and the remaining reads were sorted by how many duplicates were present
using the derep_prefix command. The remaining reads were then clustered at 96% pairwise

identity using the usearch cluster_fast command. Finally, the original unfiltered reads were
aligned back to the centroids of the clusters and an OTU table was generated using the
usearch_global command before a binary version of the table was generated.

Building of empirical networks and model prediction

Empirical networks are built from *var* DBLα types sequenced and processed from 1284 *P.*
falciparum isolates from individuals residing in BD, Ghana. Following a previously
published analysis framework, the DBLα types are translated into all six reading frames and
classified into either upsA or upsB/upsC (i.e., non-upsA) groups[67]. Gene networks are built
based on pairwise similarities of unique upsB/upsC DBLα types that are above 0.6. The
choice of the threshold is based on the average within-class sequence similarity of the 24
DBLα subclasses (see %ID in Fig. 2 of Rask et al.[16]). Since infections by multiple parasite
genomes (multiplicity of infection [MOI] larger than 1) are very common in malaria endemic
regions, we selected isolates with a total number of upsB/upsC DBLα types ranging from 40-
55 copies to maximize the probability of selecting hosts with single-clone infections, which
reduced the number of isolates to 161 (see main text for rationale of focusing on upsB/upsC
DBLα types; Supplementary Data 1). Because of the need to consider as many repertoires as
possible in our network analyses, we considered all isolates regardless of their age. The
repertoire similarity network is built among these isolates (Supplementary Fig. 5).

In order to build a classifier for the empirical network, a library of simulated networks was
generated for parameter ranges representative of Ghana: global *var* gene pools from 10,000
to 20,000 (according to the sampling counts), duration of naïve infection equal to 1 year, and
mean biting rates ranging from 0.1 to 0.5 person per day. This selection of parameters results
in EIR values similar to those reported for Upper East Region of Ghana in Appawu et al.[33]
(see Supplementary Fig. 2). We generated a vector of monthly seasonal biting amplitudes
relative to the mean biting rate, which resembles the local mosquito population of lowland in
Appawu *et al.*[33]. The simulated networks are then constructed by sampling 161 random
isolates from two periods seven months apart from each other (the end of October and the
beginning of June), resembling the sampling regime of the empirical data. The trained
discriminant functions from simulated networks[66] are then applied to empirical networks to
predict whether the network is generated from immune selection dominant forces or pure
neutral forces. The Bayesian posterior probability of classification is calculated by assuming
Gaussian densities of prior distributions of each class.

721
722

References

1. Rosindell, J., Hubbell, S. P. & Etienne, R. S. The Unified Neutral Theory of Biodiversity and Biogeography at Age Ten. *Trends Ecol. Evol.* **26,** 340–348 (2011).

2. Levine, J. M. & HilleRisLambers, J. The importance of niches for the maintenance of species diversity. *Nature* **461,** 254–257 (2009).

3. Fijarczyk, A. & Babik, W. Detecting balancing selection in genomes: limits and prospects. *Mol. Ecol.* **24,** 3529–3545 (2015).

4. Chisholm, R. A., Fung, T., Chimalakonda, D. & O'Dwyer, J. P. Maintenance of biodiversity on islands. *Proc. R. Soc. B Biol. Sci.* **283,** 20160102 (2016).

5. Cavender-Bares, J., Kozak, K. H., Fine, P. V. A. & Kembel, S. W. The merging of community ecology and phylogenetic biology. *Ecol. Lett.* **12,** 693–715 (2009).

6. Levine, J. M., Bascompte, J., Adler, P. B. & Allesina, S. Beyond pairwise mechanisms of species coexistence in complex communities. *Nature* **546,** 56–64 (2017).

7. Day, K. P. & Marsh, K. Naturally acquired immunity to Plasmodium falciparum. *Immunol. Today* **12,** A68–A71 (1991).

8. Tiedje, K. E. *et al.* Seasonal Variation in the Epidemiology of Asymptomatic Plasmodium falciparum Infections across Two Catchment Areas in Bongo District, Ghana. *Am. J. Trop. Med. Hyg.* **97,** 199–212 (2017).

9. Barry, A. E. *et al.* The Stability and Complexity of Antibody Responses to the Major Surface Antigen of Plasmodium falciparum Are Associated with Age in a Malaria Endemic Area. *Mol. Cell. Proteomics MCP* **10,** (2011).

10. Bull, P. C. *et al.* Parasite antigens on the infected red cell surface are targets for naturally acquired immunity to malaria. *Nat. Med.* **4,** 358–360 (1998).

11. Chesson, P. General Theory of Competitive Coexistence in Spatially-Varying Environments. *Theor. Popul. Biol.* **58,** 211–237 (2000).

12. Adler, P. B., HilleRisLambers, J. & Levine, J. M. A niche for neutrality. *Ecol. Lett.* **10,**

95–104 (2007).

13. Zilversmit, M. M. *et al.* Hypervariable antigen genes in malaria have ancient roots. *BMC*

Evol. Biol. **13,** 110 (2013).

14. Larremore, D. B. *et al.* Ape parasite origins of human malaria virulence genes. *Nat.*

Commun. **6,** 8368 (2015).

15. Day, K. P. *et al.* Evidence of strain structure in Plasmodium falciparum var gene

repertoires in children from Gabon, West Africa. *Proc. Natl. Acad. Sci.* **114,** E4103–

E4111 (2017).

16. Rask, T. S., Hansen, D. A., Theander, T. G., Pedersen, A. G. & Lavstsen, T. Plasmodium

falciparum Erythrocyte Membrane Protein 1 Diversity in Seven Genomes – Divide and

Conquer. *PLOS Comput. Biol.* **6,** e1000933 (2010).

17. Chen, D. S. *et al.* A Molecular Epidemiological Study of var Gene Diversity to

Characterize the Reservoir of Plasmodium falciparum in Humans in Africa. *PLOS ONE*

6, e16629 (2011).

18. Gupta, S., Ferguson, N. & Anderson, R. Chaos, Persistence, and Evolution of Strain

Structure in Antigenically Diverse Infectious Agents. *Science* **280,** 912–915 (1998).

19. Macarthur, R. & Levins, R. The Limiting Similarity, Convergence, and Divergence of

Coexisting Species. *Am. Nat.* **101,** 377–385 (1967).

20. Nosil, P., Funk, D. J. & Ortiz-Barrientos, D. Divergent selection and heterogeneous

genomic divergence. *Mol. Ecol.* **18,** 375–402 (2009).

21. Gupta, S. *et al.* The maintenance of strain structure in populations of recombining

infectious agents. *Nat. Med.* **2,** 437–442 (1996).

22. Buckee, C. O., Recker, M., Watkins, E. R. & Gupta, S. Role of stochastic processes in

maintaining discrete strain structure in antigenically diverse pathogen populations. *Proc.*

Natl. Acad. Sci. **108,** 15504–15509 (2011).

23. Artzy-Randrup, Y. *et al.* Population structuring of multi-copy, antigen-encoding genes in

Plasmodium falciparum. *Elife* **1,** e00093 (2012).

24. Scheffer, M. & Nes, E. H. van. Self-organized similarity, the evolutionary emergence of

groups of similar species. *Proc. Natl. Acad. Sci.* **103,** 6230–6235 (2006).

25. Rocha, R. D., Riolo, M. & Ostling, A. M. Competition and immigration lead to clusters

of similar species, not trait separation. *bioRxiv* 264606 (2018). doi:10.1101/264606

26. Jeraldo, P. *et al.* Quantification of the relative roles of niche and neutral processes in

structuring gastrointestinal microbiomes. *Proc. Natl. Acad. Sci.* **109,** 9692–9698 (2012).

27. Bull, P. C. *et al. Plasmodium falciparum* antigenic variation. Mapping mosaic *var* gene

sequences onto a network of shared, highly polymorphic sequence blocks. *Mol.*

Microbiol. **68,** 1519–1534 (2008).

28. Freitas-Junior, L. H. *et al.* Frequent ectopic recombination of virulence factor genes in

telomeric chromosome clusters of P. falciparum. *Nature* **407,** 1018–1022 (2000).

29. Claessens, A. *et al.* Generation of Antigenic Diversity in Plasmodium falciparum by

Structured Rearrangement of Var Genes During Mitosis. *PLOS Genet.* **10,** e1004812

(2014).

30. Newman, M. E. J. & Girvan, M. Finding and evaluating community structure in

networks. *Phys. Rev. E* **69,** 026113 (2004).

31. Barry, A. E. *et al.* Population Genomics of the Immune Evasion (var) Genes of

Plasmodium falciparum. *PLOS Pathog.* **3,** e34 (2007).

32. Freeman, L. C. A Set of Measures of Centrality Based on Betweenness. *Sociometry* **40,**

35–41 (1977).

33. Appawu, M. *et al.* Malaria transmission dynamics at a site in northern Ghana proposed

for testing malaria vaccines. *Trop. Med. Int. Health* **9,** 164–170 (2004).

34. Bull, P. C., Lowe, B. S., Kortok, M. & Marsh, K. Antibody Recognition of Plasmodium

falciparum Erythrocyte Surface Antigens in Kenya: Evidence for Rare and Prevalent

Variants. *Infect. Immun.* **67,** 733–739 (1999).

35. Bakker, P. I. W. de *et al.* A high-resolution HLA and SNP haplotype map for disease

association studies in the extended human MHC. *Nat. Genet.* **38,** 1166 (2006).

36. MacQueen, J. B. Some Methods for classification and Analysis of Multivariate

Observations. *5th Berkeley Symp. Math. Stat. Probab.* **1,** 281–297 (1967).

37. Ripley, B. D. *Spatial Statistics*. (Wiley, New York, 1981).

38. D'Andrea, R. & Ostling, A. Can Clustering in Genotype Space Reveal "Niches"? *Am.*

Nat. **187,** 130–135 (2015).

39. Clark, J. S. Individuals and the Variation Needed for High Species Diversity in Forest

Trees. *Science* **327,** 1129–1132 (2010).

40. Albrecht, L. *et al.* The South American Plasmodium falciparum var gene repertoire is

limited, highly shared and possibly lacks several antigenic types. *Gene* **453,** 37–44

(2010).

41. Deitsch, K. W., Lukehart, S. A. & Stringer, J. R. Common strategies for antigenic

variation by bacterial, fungal and protozoan pathogens. *Nat. Rev. Microbiol.* **7,** 493–503

(2009).

42. Nicoli, E. J., Ayabina, D., Trotter, C. L., Turner, K. M. E. & Colijn, C. Competition,

coinfection and strain replacement in models of Bordetella pertussis. *Theor. Popul. Biol.*

103, 84–92 (2015).

43. Cobey, S. & Lipsitch, M. Niche and Neutral Effects of Acquired Immunity Permit

Coexistence of Pneumococcal Serotypes. *Science* **335,** 1376–1380 (2012).

44. Trevino, S. G. *et al.* High-resolution single-cell sequencing of malaria parasites. *bioRxiv*

130864 (2017). doi:10.1101/130864

45. Nassir, E. *et al.* Impact of genetic complexity on longevity and gametocytogenesis of

Plasmodium falciparum during the dry and transmission-free season of eastern Sudan.

Int. J. Parasitol. **35,** 49–55 (2005).

46. Bruce, M. C. *et al.* Genetic diversity and dynamics of Plasmodium falciparum and P.

vivax populations in multiply infected children with asymptomatic malaria infections in

Papua New Guinea. *Parasitology* **121,** 257–272 (2000).

47. Klein, E. Y., Graham, A. L., Llinás, M. & Levin, S. Cross-Reactive Immune Responses

as Primary Drivers of Malaria Chronicity. *Infect. Immun.* **82,** 140–151 (2014).

48. Bruce, M. C. *et al.* Cross-Species Interactions Between Malaria Parasites in Humans.

Science **287,** 845–848 (2000).

49. Kraemer, S. M. *et al.* Patterns of gene recombination shape var gene repertoires in

Plasmodium falciparum: comparisons of geographically diverse isolates. *BMC Genomics*

8, 45 (2007).

50. Kyriacou, H. M. *et al.* Differential var gene transcription in Plasmodium falciparum

isolates from patients with cerebral malaria compared to hyperparasitaemia. *Mol.*

Biochem. Parasitol. **150,** 211–218 (2006).

51. Janes, J. H. *et al.* Investigating the Host Binding Signature on the Plasmodium falciparum

PfEMP1 Protein Family. *PLOS Pathog.* **7,** e1002032 (2011).

52. Horrocks, P., Kyes, S., Pinches, R., Christodoulou, Z. & Newbold, C. Transcription of a

subtelomerically located var gene variant in Plasmodium falciparum appears to require

the truncation of an adjacent var gene. *Mol. Biochem. Parasitol.* **134,** 193–199 (2004).

53. Frank, M. *et al.* Frequent recombination events generate diversity within the multi-copy

variant antigen gene families of Plasmodium falciparum. *Int. J. Parasitol.* **38,** 1099–1109

(2008).

54. Recker, M. *et al.* Antigenic Variation in Plasmodium falciparum Malaria Involves a

Highly Structured Switching Pattern. *PLOS Pathog.* **7,** e1001306 (2011).

55. Noble, R. *et al.* The antigenic switching network of Plasmodium falciparum and its

implications for the immuno-epidemiology of malaria. *eLife* **2,** (2013).

56. Blomqvist, K. *et al.* A Sequence in Subdomain 2 of DBL1α of Plasmodium falciparum

Erythrocyte Membrane Protein 1 Induces Strain Transcending Antibodies. *PLoS ONE* **8,**

e52679 (2013).

57. Drummond, D. A., Silberg, J. J., Meyer, M. M., Wilke, C. O. & Arnold, F. H. On the

conservative nature of intragenic recombination. *Proc. Natl. Acad. Sci. U. S. A.* **102,**

5380–5385 (2005).

58. Hermsen, C. C. *et al.* Detection of Plasmodium falciparum malaria parasites in vivo by

real-time quantitative PCR. *Mol. Biochem. Parasitol.* **118,** 247–251 (2001).

59. Collins, W. E., Warren, M., Skinner, J. C. & Fredericks, H. J. Studies on the relationship

between fluorescent antibody response and the ecology of malaria in Malaysia. *Bull.*

World Health Organ. **39,** 451–463 (1968).

60. Gibson, M. A. & Bruck, J. Efficient Exact Stochastic Simulation of Chemical Systems

with Many Species and Many Channels. *J. Phys. Chem. A* **104,** 1876–1889 (2000).

61. Gillespie, D. T. A general method for numerically simulating the stochastic time

evolution of coupled chemical reactions. *J. Comput. Phys.* **22,** 403–434 (1976).

62. Shannon, C. E. A Mathematical Theory of Communication. *SIGMOBILE Mob Comput*

Commun Rev **5,** 3–55 (2001).

63. Pielou, E. C. The measurement of diversity in different types of biological collections. *J.*

Theor. Biol. **13,** 131–144 (1966).

64. Takahata, N. & Satta, Y. Footprints of intragenic recombination at HLA loci.

Immunogenetics **47,** 430–441 (1998).

65. Charlesworth, D. Balancing Selection and Its Effects on Sequences in Nearby Genome

Regions. *PLOS Genet.* **2,** e64 (2006).

66. Jombart, T., Devillard, S. & Balloux, F. Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. *BMC Genet.* **11,** 94

(2010).

67. Ruybal-Pesántez, S. *et al.* Population genomics of virulence genes of Plasmodium

falciparum in clinical isolates from Uganda. *Sci. Rep.* **7,** 11810 (2017).

68. Dodt, M., Roehr, J. T., Ahmed, R. & Dieterich, C. FLEXBAR—Flexible Barcode and

Adapter Processing for Next-Generation Sequencing Platforms. *Biology* **1,** 895–905

(2012).

69. Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate Illumina

Paired-End reAd mergeR. *Bioinformatics* **30,** 614–620 (2014).

70. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics **26,** 2460–2461 (2010).

Acknowledgements

This research was supported by the Fogarty International Center at the National Institutes of Health [Program on the Ecology and Evolution of Infectious Diseases (EEID), Grant number: R01-TW009670]. S.P. was supported by a James S. McDonnell Foundation 21st Century Science Initiative -- Postdoctoral Program in Complexity Science-Complex Systems Fellowship Award and by a Fulbright Fellowship from the U.S. Department of State. We wish to thank the participants, communities and the Ghana Health Service in Bongo District Ghana for their willingness to participate in this study. We would also like to thank the personnel at the Navrongo Health Research Centre for sample collection and parasitological assessment/expertise. We are grateful to Abraham R. Oduro, Anita Ghansah, and Kwadwo Koram for their helpful input related to the field study, to Gerry Tonkin-Hill for the development of the Illumina sequencing cleaning and clustering pipelines, and to Michael F. Duffy and Andrew P. Dobson for their insightful comments on an earlier version of the manuscript. We appreciate the support of the University of Chicago through computational resources at the Midway cluster.

Author contributions

Q.H., S.P. and M.P. conceived and designed the study. Y.A.R. and Q.H. worked on the model. E.B.B. and Q.H. wrote the simulation code. K.E.T. and K.P.D. designed and led the data collection in Ghana. K.E.T. and S.R.P. performed the molecular experiments and genetic sequencing. Q.H. and S.P. analyzed the data. Q.H., S.P. and M.P. wrote the paper. All authors contributed to the final version of the manuscript.

Competing financial interests

The Authors declare no Competing Financial or Non-Financial Interests.

913   **Figure Legends**
914

915
Figure 1. Schematic illustration of the *var* gene model. a, Each parasite genome (ovals)
consists of a repertoire of *g* copies of *var* genes. Each *var* gene (depicted by different colors
within each parasite) is in turn represented as a linear combination of epitopes (depicted by
different shapes), with each epitope having many possible variants (alleles, depicted by
different colors). **b**, The local population receives *var* repertoires from a fixed global *var*
gene pool through migration. **c**, At each transmission event, one donor and one receiver host
are selected at random from the host pool. Each parasite genome in the donor host is
transmitted to the mosquito with probability of 1/(number of genomes). During the sexual
stage of the parasite (within mosquitoes), different parasite genomes can exchange *var*
repertoires through meiotic recombination to generate novel recombinant repertoires. The
receiver host can receive either recombinant genomes or original genomes. During the
asexual reproduction stage of the parasite (within the blood stage of infection), *var* genes
within the same genome exchange epitope alleles through mitotic (ectopic) recombination.

Also, epitopes can mutate. These two processes generate new *var* genes. Each *var* gene is
expressed sequentially and the infection ends when all the *var* genes in the repertoires have
been expressed. A new transmission event may occur throughout the period of expression of
var genes as the result of biting events. **d,** Comparison of the three different models. Only the
immune selection model includes specific immunity, through the dependence of infection
duration on the memory of previous alleles that have been seen by a given host. For
meaningful comparisons, all the models have the same mean duration of infection.
936

Figure 2. **Population structure for the three different models quantified using network**
properties. Shown here are repertoire similarity networks and representative network metrics
across scenarios under different diversity regimes generated from model simulations with
high competition (high duration of naive infection and high transmission rates). Upper panel,
medium diversity (gene pool size of 1,200) and lower panel, high diversity (gene pool size of
24,000). **a, d**, comparisons of repertoire similarity networks of 150 randomly sampled
parasite *var* repertoires from one time point under three scenarios. Edge width is relative to
the strength of genetic similarity between pairs of repertoires. Only the top 1% of edges are
drawn and used in the analysis (see Supplementary Fig. 3 for distribution of edge weights).
Within the largest component of each network, the size of each repertoire is relative to its
normalized betweenness centrality. The value of maximum modularity Q is calculated using
edge-betweenness[30] and is shown at the lower right corner of each network. The modules
obtained in these networks represent groups of highly similar repertoires (strain modules),
which are conceptually similar to geographically isolated populations with limited gene flow.
We therefore calculate the pairwise F_{ST} of strain modules, to quantify how different strain
modules are from each other, providing a measure of limiting similarity that compares
within-module and between-module diversity[64]. **b, e**, pairwise module F_{ST} distributions of
100 repertoire similarity networks per scenario. **c, f**, distributions of the proportion of
occurrences of 3-node graph motifs for the three models across 100 repertoire similarity
networks. Each box represents the second quartile and third quartile of the distribution, with
median value indicated by the line. The whiskers represent the ranges of the first and the
fourth quartiles. Outliers are plotted as circles.

962
Figure 3. Importance of network features for the classification and the power of correct
classification of the selection model. Results are for different levels of duration of naïve
infection, biting rate, and *var* gene pool size (medium diversity [1,200-2,400] in **a-b**, and
high diversity [12,000-24,000] in **c-d**). **a**, **c**, The shade of colored squares indicates the
proportion of correct assignments of the immune selection scenario. Variance contribution of
network features in the first linear discriminant function (LD1) are displayed with color-
codes corresponding to the feature groups (**b**, **d**; see Supplementary Table 1 for feature
groups). Only top features that explain at least 90% of the variance are shown. In simulations
with a gene pool of medium diversity (**a**), the proportion of correct assignments of the
selection model increases with increasing infection duration and biting rate. When the genetic
pool is of high diversity, the selection model is almost always perfectly assigned (**c**), while
neutral and generalized immunity models are harder to differentiate, even under high
transmission and long infection durations (see supplementary table 2). The proportions of
different motifs are the most important features that discriminate scenarios in both high and
medium diversity. In the high diversity scenarios, there are a higher number of important
features that equally contribute to the classification (**d**), as compared to the medium diversity
scenarios (**b**). The classifiers shown in this figure are built with the top 10% edge weights of
the repertoire similarity matrices.
981

982
Figure 4. The relationship between gene frequencies and number of genes similar to a
focal gene is different between selection and null models. a, In a gene similarity network,
each node represents a unique gene circulating in the population and the edges encode the
sharing of at least one allele between genes. The size of the node is proportional to its
frequency in the population and the node degree k depicts the number of genes that share at
least one allele with the focal gene. There is a negative correlation between a gene's
frequency and degree in the immune selection scenario (**b,** r = -0.412, t-test: p-value < 2.2e-
16), but not in the complete neutrality (**c,** r = 0.016, t-test: p-value = 0.17) or generalized
immunity scenarios (**d,** r = -0.002, t-test: p-value = 0.89). G = 24,000, b = 0.5, D = 1 year.
992

Figure 5. **Empirical investigation of the Ghana data shows resemblance to the immune selection scenario.** (**a**) Negative correlation between DBLα type frequencies and their number of similar genes for the upsB/upsC *var* genes in the parasite population (r = -0.040, p<2.2e-16). The number of similar genes is calculated as the degree (k) of a focal gene in the gene similarity network, for amino acid similarities above 0.6. Histograms on the top and right of the plot show the distributions of k and DBLα-type frequencies, respectively. (**b**) Classification of networks generated with the agent-based model using Discriminant Analyses of Principle Components[66] onto a 2-D space formed by the two linear discriminant (LD) functions using the top 10% edge weights. The empirical network is more likely to be generated under an immune selection regime (posterior probability [PP] = 1), as opposed to neutrality (PP = 3.57E-8) or generalized immunity (PP = 2.15E-9). The classification relies on comparisons of 34 network properties (see Supplementary Table 1) trained with 7000 simulated networks and verified on test sets of 800 networks (i.e., 100 combinations of different parameter settings for each of the scenarios were run; infected hosts were sampled at October and the next May per year as empirical data at the stationary stage of the simulations (i.e., the last 26 years in the simulations); similarity networks were then built for randomly sampled parasites from the sampled hosts). Accuracy of network classifications is above 0.99 for each scenario (See Supplementary Table 2 for comparisons of accuracy and the classifications of the empirical network using different percentage of top edges in the network, and Supplementary Fig 5 for motif properties of the empirical network compared with simulated ones).

1016