Transcriptome dynamics at the *Arabidopsis* graft junction reveal an inter-tissue recognition mechanism that activates vascular regeneration

Charles W Melnyk[1,2], Alexander Gabel[3], Thomas J Hardcastle[4], Sarah Robinson[5], Shunsuke Miyashima[6], Ivo Grosse[3], Elliot M Meyerowitz[1,7]

1- Sainsbury Laboratory, University of Cambridge, Bateman Street, Cambridge CB2 1LR, UK

– Department of Plant Biology, Swedish University of Agricultural Sciences, Almas allé 5, 756 51 Uppsala, Sweden

- Institute of Computer Science, Martin Luther University Halle–Wittenberg, 06120 Halle (Saale), Germany

– Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, United Kingdom

- Institute of Plant Science, University of Bern, Altenbergrain 21, 3013 Bern, Switzerland

– Graduate School of Biological Sciences, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

– Howard Hughes Medical Institute and Division of Biology and Biological Engineering, California Institute of Technology, 1201 East California Boulevard, Pasadena, CA 91125, USA

Contact: charles.melnyk@slu.se

ABSTRACT

The ability for cut tissues to join together and form a chimeric organism is a

remarkable property of many plants, however, grafting is poorly characterized at the

molecular level. To better understand this process we monitored genome-wide

temporal and spatial gene expression changes in grafted *Arabidopsis thaliana*

hypocotyls. Tissues above and below the graft rapidly developed an asymmetry such

that many genes were more highly expressed on one side than the other. This

asymmetry correlated with sugar responsive genes and we observed an accumulation

of starch above the graft that decreased along with asymmetry once the sugar-

transporting vascular tissues reconnected. Despite the initial starvation response

below the graft, many genes associated with vascular formation were rapidly

activated in grafted tissues but not in cut and separated tissues indicating that a

recognition mechanism activated that was independent of functional vascular

connections. Auxin which is transported cell-to-cell, had a rapidly elevated response

that was symmetric, suggesting that auxin was perceived by the root within hours of

tissue attachment to activate the vascular regeneration process. A subset of genes

were expressed only in grafted tissues, indicating that wound healing proceeded via

different mechanisms depending on the presence or absence of adjoining tissues. Such

a recognition process could have broader relevance for tissue regeneration, inter-

tissue communication and tissue fusion events.

INTRODUCTION

For millennia people have cut and rejoined plants through grafting. Generating such

chimeric organisms combines desirable characteristics from two plants, such as

disease resistance, dwarfing and high yields, or can propagate plants and avoid the

delays entailed by a juvenile state (1). Agriculturally, grafting is becoming more

relevant as a greater number of plants and species are grafted to increase productivity

and yield (2). However, our mechanistic understanding of grafting and the biological

processes involved, including wound healing, tissue fusion and vascular formation,

remain limited.

Plants have efficient mechanisms to heal wounds and cuts, in part through the

production of wound-induced pluripotent cells termed callus. The callus fills the gap

or seals the wound, and later, differentiates to form epidermal, mesophyll and

vascular tissues (3). In grafted *Arabidopsis* hypocotyls, tissues adhere 1-2 days after

grafting and the phloem, the tissue that transports sugars and nutrients, connects after

three days (4, 5). The xylem, tissue that transports water and minerals, connects after

seven days (4). Plant hormones are important regulators of vascular formation, and at

the graft junction, both auxin and cytokinin responses increase in the vascular tissue

(4-6). Auxin is important for differentiation of vascular tissues whereas cytokinin

promotes vascular stem cells, termed cambium, to divide and proliferate in a process

known as secondary growth (7, 8). Auxin is produced in the upper parts of a plant and

moves towards the roots via cell-to-cell movement. Auxin exporters, including the

PIN proteins, transport auxin into the apoplast, whereas auxin importers, such as the

AUX and LAX proteins, assist with auxin uptake into adjacent cells (8). Disrupting

this transport, such as by mutating *PIN1*, inhibits healing of a wounded stem (9).

Blocking auxin transport with the auxin transport inhibitor TIBA (2,3,5-

triiodobenzoic acid) in the shoot inhibits vascular formation and cell proliferation at

the *Arabidopsis* graft junction (6). In addition to auxin, other compounds, including

sugars, contribute to vascular formation. The localised addition of auxin to callus

induces phloem and xylem but requires the presence of sugar (10, 11). In plants,

sugars are produced in the leaves and transported through the phloem to the roots

(12). The role of sugars in vascular formation and wound healing is not well

established, however, sugars promote cell division and cell expansion (13), processes

important for development including vascular formation.

The molecular and cellular mechanisms for wound healing, tissue reunion and graft

formation remain largely unknown. One emerging theme is that the top and bottom of

the cut do not behave similarly. Such tissue asymmetry occurs in other plant tissues,

most notably leaves. Developing leaf primordia have an inherent asymmetry that is

thought to derive from positional signals from the shoot apical meristem that specifies

differences between the top and the bottom of the leaf. The molecular mechanisms

that establish asymmetry are not well described, though one hypothesis is that auxin

contributes towards or acts on the putative signal (14). Asymmetry also appears in cut

Arabidopsis inflorescence stems where the transcription factor *RAP2.6L* expresses

exclusively below the cut whereas the transcription factor *ANAC071* expresses

exclusively above the cut (9). Both were important for stem healing and *ANAC071*

and a close homologue, *ANAC096,* were important for graft formation (6).

Asymmetry also exists in genetic requirements, since *ALF4* and *AXR1*, two genes

involved in auxin perception, are important below but not above the graft junction for

phloem connection (4). However, *ANAC071* is expressed symmetrically around the

hypocotyl graft junction three days after grafting (6) so the extent of asymmetry and

the mechanistic basis for it during wound healing remains largely uncharacterised.

Previous efforts have characterised wound healing and tissue reunion using

transcriptomic analyses. Mechanical wounding altered approximately 8% of the

Arabidopsis transcriptome and showed a high degree of overlap with transcriptomic

changes elicited by pathogen attack and abiotic stress (15). Stem wounding and

wound-induced callus formation altered the expression of hundreds or thousands of

genes (9, 16, 17), whereas grafting grape vines, lychee trees and hickory trees induced

hundreds or thousands of differentially expressed genes involved in hormone

response, wound response, metabolism, cell wall synthesis and signal transduction (5,

18-21). These grafting studies provide limited information, as tissues from above and

below the graft junction were not isolated to test whether these tissues behaved

differently, and controls were not performed to distinguish how grafting and tissue

fusion might differ from a response associated with cut tissues that remained

separated. Here, we perform an in-depth analysis to describe the spatial and temporal

transcriptional dynamics that occur during healing of cut *Arabidopsis* tissues that are

joined (grafted) or left unjoined (separated). We find that the majority of genes

differentially expressed are initially asymmetrically expressed at the graft junction

and that many of these genes are sugar responsive, which correlates with severing of

the phloem tissue and the accumulation of starch above the junction. However, genes

associated with cell division and vascular formation activate on both sides of the graft

and, similarly, auxin responsiveness activates equally on both sides. We propose that

the continuous transport of substances, including auxin, independent of functional

vascular connections, promoted division and differentiation, while the enhanced auxin

response and blocked of sugar transport provided a unique physiological condition to

activate genes specific to graft formation that promote wound healing.

RESULTS

Genes are asymmetrically expressed around the graft

Previous analyses revealed an asymmetry in several genes important for tissue
reunion or graft formation and identified a number of genes expressed above a cut
that were not expressed below (4, 9). To investigate whether asymmetry was a
common feature of grafting and tissue reunion, we generated RNA deep sequencing
libraries from *Arabidopsis thaliana* hypocotyl tissues immediately above and
immediately below the graft junction 0, 6, 12, 24, 48, 72, 120, 168 and 240 hours after
grafting (HAG)(Figure 1A). Prior to RNA extraction, we separated top and bottom
tissues at the graft junction. We found that the strength required to break apart the
graft junction increased linearly (Figure S1) similarly to previously reported breaking
strength dynamics of grafted *Solanum pennellii* and *Solanum lycopersicum* (22, 23).
When pulling apart grafts to separate top and bottom for sample preparation, grafts
broke cleanly with minimal tissue from one half present in the other half (Figure S1,
Movie S1, S2). We measured the amount of tissue from tops adherent to bottoms and
vice versa (Figure S1) and found less than 4% cross-contamination. In addition to
grafting, we also prepared libraries from ungrafted hypocotyls ("intact" treatment)
and cut plants that had not been reattached ("separated" treatment)(Figure 1A). We
herein refer to tissues harvested above the graft junction or from the shoot side of
separated tissue as "top" and that from below the graft or from the root side of
separated tissue as "bottom" (Figure 1A).
156
RNAs that were differentially expressed as a consequence of grafting (compared to
intact hypocotyls) equally in tops and bottoms of grafts (symmetrically expressed) or
were more highly expressed in one tissue than the other (asymmetrically expressed),
were identified by performing a pairwise analysis of the protein-coding transcriptome
datasets that were differentially expressed relative to the intact group. Several
thousand RNAs were identified that fit either pattern of expression including the
transcript of the cambial markers *HCA2* that was induced symmetrically, and *ANT*
that was induced asymmetrically (Figure 1B-G, Figure S2). 6 to 48 hours after
grafting, the number of graft-differentially expressed genes that were asymmetrically
expressed was roughly 3-fold greater than those symmetrically expressed indicating
that tissues above the cut changed their expression dynamics relative to below the cut.
However, at 72 hours the numbers were nearly equal, and by 120 hours, the number
of symmetrically differentially expressed genes was 3-fold greater than those
asymmetrically expressed (Figure 1B). As a second approach, we performed a

 hierarchical clustering analysis that indicated that the grafted top and grafted bottom

172 became most similar after 72 hours (Figure S3), consistent with the symmetry

173 analysis (Figure 1B). Thus, graft healing and tissue reunion promoted a shift from

174 asymmetry to symmetry (Figure 1B).

Figure 1. Grafting rapidly activates genes in both asymmetric and symmetric patterns. (A) Intact, cut and separated, or cut and grafted *Arabidopsis* tissues were harvested approximately 0.5mm above (top), 0.5mm below (bottom) the cut site or for intact plants 1mm segments spanning the region where cuts were made in grafted and separated plants. (B) Pairwise analysis between the grafted top and grafted bottom identified sets of protein-coding genes symmetrically or asymmetrically expressed at the graft junction, with an FDR < 0.05 and a likelihood of symmetric/asymmetric expression of greater than 50%. Asymmetrically expressed genes were further divided into those whose RNAs were higher in the top (orange dotted line) or higher in the bottom (purple dotted line), again with an FDR < 0.05 and a likelihood of asymmetric expression greater than 50%. (C,D) Expression profiles for transcripts of cambium-associated genes that initially activate symmetrically (*HCA*) or asymmetrically (*ANT*) plotted for intact, separated and grafted samples. (E-G) *HCA2* transcription

upregulates above the graft junction (E) and below the graft junction. (F)
pHCA2::RFP was grafted to Col-0 roots (E) or Col-0 shoots (F) to avoid ambiguity of
signal origin at the junction. *HCA2* was also upregulated in separated shoots but not in
intact samples or in high levels in separated roots (G). HAG, hours after grafting.
HAS, hours after separation. White triangle denotes initial fluorescent signal, dashed
lines denote the graft junction.

Sugar response correlates with asymmetric gene expression

The shift from asymmetry to symmetry could be due to phloem reconnection at 72

hours ((4), Figure 2A) and the resumption of hormone, protein and sugar transport.

We tested a role for sugar by grafting in the presence of exogenous sucrose which has

previously been reported to affect grafting success (24). Low levels of exogenous

sucrose lowered grafting efficiency (Figure 2A), suggesting that differential sugar

responses at the graft junction might be important for vascular reconnection.

Expression of *ApL3*, a gene whose expression is induced by sugar (25), was rapidly

upregulated in separated tops and grafted tops, whereas expression of *DIN6*, *GDH1*

and *STP1*, genes whose expression is repressed by sugar (25-27), was rapidly

upregulated in separated bottoms and grafted bottoms (Figure 2B, Figure S4). These

observations were consistent with sugar accumulation in the grafted top and sugar

depletion in the grafted bottom. The expression of these genes returned to levels

similar to intact samples by 120 hours and, with the exception of *ApL3*, the grafted

samples normalised expression more rapidly than the separated tissues. Genes

associated with photosynthesis increase expression in separated bottoms 24 hours

after cutting, a common response to starvation (13), but likely too late to affect sugar

levels before 24 hours (Figure S4). A transcriptional overlap analysis with RNAs

from known glucose-responsive genes (Table S1) revealed a strong overlap with

genes differentially expressed by grafting. RNAs from known glucose-induced genes

were upregulated in separated tops and grafted tops, whereas transcripts from known

glucose-repressed genes were upregulated in separated bottoms and grafted bottoms

(Figure 2C, Figure S4). This trend was not observed with genes differentially

expressed by mannitol treatment (Figure S4), suggesting the effect was specific to

metabolically active sugars. To further investigate this effect, we stained grafted,

separated and intact plants with Lugol solution to assay for the presence of starch.

Staining above the graft junction increased 48-72 hours after grafting (Figure 2D). By

120 hours, staining was equal on both sides of the graft whereas in separated tops

staining became stronger after 72 hours (Figure 2D, Figure S4). We concluded that

starch accumulated above the graft junction, but after 72 hours, this asymmetry disappeared. To test whether the accumulation of starch and increased sugar responsiveness could explain the observed transition from asymmetry to symmetry, we compared our datasets to previously published genes that are induced by starvation or are induced by sucrose re-addition (Table S1). In early time points, 20-31% of asymmetrically expressed genes are genes known to respond to sugars compared to 2-5% of symmetrically expressed genes (Table 1). However, at 72 hours, the overlap between asymmetrically expressed genes and sugar responsive genes reduced substantially (Table 1).

Figure 2. Asymmetric changes in accumulation of sugar-responsive RNAs and of starch occur at the graft junction; equalizing sucrose between the grafted segments reduces and delays phloem reconnection. (A) *pSUC2::GFP*-expressing *Arabidopsis* shoots were grafted to Col-0 wild type roots and GFP movement to the roots was monitored over 7 days for phloem connection in the presence or absence of various concentrations of sucrose. (B) Expression profiles for transcripts of a sugar-repressed gene (*GDH1*), and a sugar-induced gene (*ApL3*) were plotted for intact, separated and grafted samples. (C) Transcriptional overlap between previously published glucose-induced or glucose repressed genes and our dataset. The number in brackets represents the number of glucose-responsive genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or down-regulated in our dataset relative to intact samples. Asterisks represent a significant difference ($p < 0.05$) between the ratio of up- and down-regulated genes in a previously published transcriptome dataset compared to the ratio

250 of all up- and down- regulated genes in our grafting dataset at a certain time point. (D)
251 Lugol staining of grafted plants at various time points revealed dark brown staining
252 associated with starch accumulation. HAG, hours after grafting.
253

254 **Table 1. Sugar response overlaps with asymmetry.** Symmetrically and
255 asymmetrically differentially expressed genes were compared to previously published
256 sugar-responsive genes (28)[1] and the percent overlap calculated (* p<.05). HAG,
257 hours after grafting.
258

HAG	Sugar Induced[1]	Graft Bottom=Top	Overlap	%	Graft Top>Bottom	Overlap	%	Graft Bottom>Top	Overlap	%
6	2243	4988	263	5	6679	1403	21*	4971	99	2
12	2243	3473	165	5	7111	1444	20*	5657	231	4
24	2243	4135	171	4	6873	1414	21*	4942	195	4
48	2243	3689	197	5	6601	1241	19*	4915	218	4
72	2243	10421	1005	10*	2459	228	9*	2019	138	7
120	2243	15012	1063	7	1510	220	15*	941	115	12*
HAG	Sugar Repressed[1]	Graft Bottom=Top	Overlap	%	Graft Top>Bottom	Overlap	%	Graft Bottom>Top	Overlap	%
6	1998	4988	107	2	6679	53	1	4971	1563	31*
12	1998	3473	68	2	7111	112	2	5657	1526	27*
24	1998	4135	88	2	6873	72	1	4942	1525	31*
48	1998	3689	113	3	6601	93	1	4915	1427	29*
72	1998	10421	530	5	2459	111	5	2019	538	27*
120	1998	15012	979	7	1510	84	6	941	210	22*

259
260

Vascular formation and cell division activates on both sides of the graft

Since the grafted bottom samples exhibited a starvation response up to 48 hours after

grafting, we reasoned that pathways associated cell division and cell differentiation

would be delayed or inhibited. We looked at the expression of markers associated

with vascular formation and cell division in the transcriptome datasets. Cambium,

phloem and provascular markers activated within 6 hours in grafted top samples, but

activation was only delayed 0-24 hours in grafted bottom samples depending on the

gene (Figure 1B-C, Figure 3, Figure S5, Figure S6). Expression of phloem markers

peaked in both grafted tops and grafted bottoms at 72 hours (Figure 3, Figure S5,

Figure S6), the time when phloem reconnections form in grafted *Arabidopsis* (Figure

1, (4, 5)). Notably, the early phloem marker *NAC020* activated before the mid phloem

marker *NAC086* which activated before the late phloem marker *NEN4*, consistent

with the dynamics of phloem transcriptional activation during primary root

development and leaf vascular induction (Figure S5)(29, 30). Certain markers

associated with xylem formation, such as *VND7* and *BFN1*, activated early in the

grafted top. Other xylem markers, such as *IRX3* and *CESA4*, activated late in grafted

samples. By 120 hours after grafting, xylem markers were activated in top and

bottom, consistent with when the first xylem strands differentiate at the graft junction

(4). Genes associated with cell division were activated by 12 hours in the grafted top

and by 24 hours in the grafted bottom (Figure 3, Figure S5). On the other hand,

control genes whose expression does not typically vary between tissues and

treatments (31) were not differentially expressed in grafted tops or bottoms (Figure

S6). The RNAseq expression data appeared to correlate well with transcriptional

fluorescent reporters for both activation dynamics and the localisation of expression

(Figure 1D-F, Figure S2).

The similar activation dynamics of vascular differentiation genes in grafted tops and

grafted bottoms prompted us to test whether this phenomenon occurred with other

known developmental processes. We obtained lists of genes whose expression is

associated with various biological processes from previous publications (Table S1)

and tested how many of the genes differentially expressed in our transcriptomes

overlapped with the previously published lists. Differentially expressed genes in

grafted samples and separated tops partially overlapped with those whose expression

is associated with phloem, xylem and procambium formation (Figure 4, Figure S7).

There was a high overlap between *Arabidopsis* inflorescence stem healing and

grafting, as well as between vascular induction in leaf disk cultures and grafting

(Figure 4). Various genes expressed in a cell type-specific manner also showed a high

transcriptional overlap with grafting, including phloem, endodermis and protoxylem

(Figure 4, Figure S7). In nearly all cases, the separated top, grafted top and grafted

bottom samples showed similar activation dynamics. The separated bottom samples

were exceptional though since gene expression associated with vascular development

and cell-specific processes was downregulated (Figure 4, Figure S7). We also

compared our datasets to RNAs expressed in longitudinal cross sections of the

Arabidopsis root (32). There was little overlap between grafted bottoms and sections

from the root meristemic zone, whereas overlap existed between grafted tops and the

root meristemic zone in early time points, and between grafting and the root

maturation zone (Figure S8). Our analysis also revealed that two genes expressed in

cambium, *WOX4* and *PXY*, were induced by grafting but the primary root marker

WOX5 and lateral root marker *LBD18* were not (Figure 3, Figure S5, Figure S6).

Figure 3. Transcriptional dynamics of genes associated with provasculature, phloem, xylem and cell division. Expression levels were plotted over time for intact, separated and grafted samples.

Figure 4. Transcriptional overlap between previously published vascular datasets and the grafting datasets. Genes whose transcripts are associated with various cell types or biological processes were taken from previously published datasets (see Table S1) and compared to the transcriptomic datasets generated here. The number in brackets represents the number of cell type-specific or process-specific genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0

for differentiation expressed genes (DEG) up- or down-regulated in our dataset
relative to intact samples. Asterisks represent a significant difference (p < 0.05)
between the ratio of up- and down- regulated genes in a previously published
transcriptome dataset compared to the ratio of all up- and down- regulated genes in
our grafting dataset at a certain time point.

Auxin response is symmetric at the graft

The rapid activation of vascular markers in the grafted bottoms despite the starvation

response promoted us to investigated whether other mobile substances such as

phytohormones could play a role in gene activation. We compared lists of genes

known to respond to cytokinin, ethylene or methyl jasmonate (33) and found no

substantial overlap between these lists and genes differentially expressed by grafting

(Figure S9)(Table S1). Abscisic acid-responsive and brassinosteroid-responsive genes

showed overlap with genes differentially expressed in our datasets, but this overlap

was of a similar magnitude in both separated and grafted datasets suggesting the

effect was not specific to grafting (Figure S9). Auxin responsive transcripts were

exceptional though, as they showed a substantial overlap with RNAs differentially

expressed in our data sets that varied depending on the treatment (Figure 5A-B,

Figure S9). Auxin-induced genes were upregulated in separated tops, grafted bottoms

and grafted tops whereas they were repressed in separated bottoms (Figure 8). Auxin

responsive genes such as *IAA1* and *IAA2* (34) were induced to similar levels in grafted

tops and grafted bottoms at 24 hours. To further investigate whether auxin response

was uniform between grafted tops and grafted bottoms, we grafted the auxin-

responsive fluorescent reporter *p35S:DII-Venus* whose fluorescent protein is degraded

in the presence of auxin (35). DII-Venus fluoresced in the separated bottoms but did

not fluoresce in grafted bottoms 14 hours after cutting (Figure 5C) indicating

separated bottoms had a low level of auxin response but grafted tops, grafted bottoms

and separated tops had a high level of auxin response.

Figure 5. Auxin response is symmetric at the graft junction. (A,D) Expression
profiles for various auxin-responsive genes (*IAA1*, *IAA2*) or auxin transporter genes
(*PIN1* and *ABCB1* genes) were plotted for intact, separated and grafted samples. (B)
Overlap between previously published auxin-induced or auxin-repressed RNAs and
our dataset. The number in brackets represents the number of auxin-responsive genes
identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for
differentiation expressed genes (DEG) up- or down-regulated in our dataset relative to
intact samples. Asterisks represent a significant difference ($p < 0.05$) between the
ratio of up- and down- regulated genes in a previously published transcriptome
dataset compared to the ratio of all up- and down- regulated genes in our grafting
dataset at a certain time point. (C) Grafted and separated plants expressing the auxin
responsive *p35S::DII-Venus* transgene that is degraded in the presence of auxin reveal
a reduction of auxin response in cut bottoms, but not in grafted bottoms. HAG, hours
after grafting.

Tissue fusion imparts a unique physiological response that differs from tissue separation

We hypothesized that the symmetric auxin response and asymmetric sugar response at the graft junction could allow a unique transcriptional response since neither separated plants nor intact plants had similar response dynamics to sugars and auxin as measured by genome-wide gene response, starch accumulation and *p35S::DII-Venus* expression (Figure 2, Figure 5). To uncover protein-coding genes differentially expressed specifically by grafting, we performed an empirical Bayesian analysis (36) where at each time point we computed likelihoods for each gene for each possible pattern of differential expression in the intact, grafted top, grafted bottom, separated top and separated bottom tissues (Figure 6A). These likelihoods were used to define a similarity score between pairs of genes, which was used to cluster genes with similar patterns of gene expression across the five tissues (37). Clusters were then identified by the predominant pattern of gene expression observed within a cluster. This analysis produced 113 clusters containing at least 10 RNAs at a single time point (Table S2; Dataset S1). Approximately 6000 genes were differentially expressed in at least one tissue whereas between 1000 and 4000 genes were not differentially expressed (Figure 6B).

To simplify the analysis, we considered clusters where gene expression was grouped into patterns consisting of one comparison. At early time points, the cluster containing genes differentially expressed in grafted tops, grafted bottoms, separated tops and separated bottoms had high numbers that decreased with time and could represent an general wound response (Figure 6C). Similarly, clusters containing separated tops and grafted tops and clusters containing separated bottoms and grafted bottoms initially had high numbers that decreased with time. This observation indicated the grafted top was initially transcriptionally similar to the separated top, whereas the grafted bottom was initially transcriptionally similar to the separated bottom. After the 48-hour time point, clusters containing genes only differentially expressed in separated tops or only differentially expressed in separated bottoms increased in numbers, suggesting these tissues gained a unique pattern of gene expression. The cluster containing genes differentially expressed in grafted tops and grafted bottoms increase in numbers throughout the healing process (Figure 6C). We searched for cluster categories that contained genes only differentially expressed by grafting and found very few genes

downregulated by grafting or upregulated only in the grafted top (Figure 7A). Instead,

there were clusters that contained several hundred differentially expressed genes

present either in the grafted bottom only, or present in both grafted bottom and grafted

top (Figure 7A). Genes whose expression changed only in the grafted bottom sample

were prevalent early during grafting and were most common at 48 hours, whereas

genes activated in both top and bottom became prevalent at 48 hours and were most

common at 120 hours (Figure 7A). We performed a gene ontology (GO) analysis on

the genes whose expression was grafting-specific and found that genes coding for

RNAs differentially expressed only in the grafted bottom sample were enriched in the

immune response and chitin response biological process categories (Table S3).

Previously published chitin-induced RNAs had a high proportion of overlap with

differentially expressed graft bottom-specific genes (Figure 7E, Figure S10).

Grafting-specific RNAs expressed symmetrically, in both the grafted top and grafted

bottom, were enriched in vascular-related biological processes (Table S3). Previously

published phloem-enriched, endodermal-enriched, vascular-induction and stem-

wounding associated RNAs had a high proportion of overlap with differentially

expressed graft-specific genes (Figure 7C-D, Figure S10). Since few genes were

grafting-specific and grafted tissues were initially transcriptionally quite similar to

separated tissues (Figure 6C, Figure 7A), we reasoned that tissues separated for short

periods (<48 hours) could be grafted with similar reconnection dynamics as tissues

that had been grafted immediately. To test this hypothesis, plants were cut at day 0,

and grafted at 1-5 days after separation. Separated tissues did not speed up vascular

reconnection, and instead, it always took three days from the point of tissue

attachment for vascular connections to form (Figure S10). Furthermore, the shoot lost

competence to graft 2-3 days after separation whereas the root remained competent to

graft up to 5 days after separation (Figure S10). Together, it appears the grafted shoot

and root have a unique physiological response that differ from separated plants and

that tissue attachment is required to activate graft formation and tissue fusion.

Figure 6. Clustering the transcriptome at each time point, based on likelihoods of
all possible patterns of differential expression. (A) Cartoon depicting the Bayesian
segmentation. (B) Analysis of differential behaviour produced 113 categories
containing at least 10 genes whose expression was expressed in a specific differential
pattern in at least one time point (see Table S2). One group is comprised of genes
whose transcript levels are not substantially changed in the five tissues (unchanged)
whereas the other group is comprised of the sum of the other 112 groups (genes
whose transcript levels changed after treatment in a least one tissue) over the time
points tested. (C) Major categories in the segmentation revealed RNAs whose levels
changed in all the treatments listed relative to intact samples. Note that a gene can be
represented only in one category for a given time point, that category in which the
transcript level changes best fit the category.

Figure 7. A subset of genes is differentially expressed only during graft formation when compared to 0 hour grafts, or intact or separated tissues. (A) Certain genes were only differentially expressed in grafted tops, grafted bottoms or both in grafted tops and grafted bottoms. (B) Expression profiles for a graft bottom-specific (*ERF6*) or a graft top and bottom differentially regulated gene (*RTM2*) were plotted for intact, separated and grafted samples. (C) Grafting-specific genes are also expressed in other processes of tissue fusion, such as stem healing. Here, 246 previously published RNAs whose differential expression is associated with stem healing were compared to our dataset to assess transcriptional overlap with the grafting-specific genes. Asterisks represent a significant high overlap (p <0.05) of previously published genesets that are also differentially expressed in the grafted samples at a certain time point. (D-E) Genes differentially expressed in grafted tops and grafted bottoms show high overlap to 324 previously published genes whose transcripts are associated with phloem, whereas genes expressed in grafted bottoms show high overlap to 286 previously published genes whose encoded RNAs are associated with chitin treatment. Asterisks represent a significant high overlap (p <0.05) of previously published genes that are also differentially expressed in the grafted samples at a certain time point.

DISCUSSION

To better understand how plants graft, we analysed in depth an RNA deep sequencing dataset that spatially and temporally distinguished genes activated by cutting followed by tissue attachment or continuous tissue separation. Cutting promoted a similar wound response in both grafted and separated tissues, however, by 72 hours after cutting the grafted and separated tissues became transcriptionally dissimilar (Figure 6C) indicating that tissue fusion was mechanistically different from healing an unattached cut surface. During graft formation, tissues had a very high transcriptional overlap with genes differentially expressed by inflorescence stem healing and by vascular induction in leaves (Figure 4, Figure S7) (9, 29) suggesting grafting is closely related to tissue reunion and vascular formation processes. Graft formation

had little transcriptional overlap with lateral root formation (Figure S6)(32) and

appeared to follow a pathway similar to secondary root growth (Figure S6, Figure S8)

since cambium markers *WOX4* and *PXY* that are specific to secondary growth were

expressed during grafting (38)(Figure S5). Grafted tops initially showed a short-

lasting and small transcriptional overlap with genes expressed during primary root

formation which may be related to the accumulation of substances activating

adventitious root formation, a common response in failed grafts or in cut shoots

(Figure S4C). Thus, we conclude that grafting proceeds via a pathway involving

secondary growth with radial meristems activating in the mature cambium to heal the

wound. Vascular formation genes including those specifying cambium and phloem

were activated early, followed by an activation of cell division genes, suggesting that

the start of cellular differentiation preceded activation of cell division. Xylem identity

genes showed an early and a late activation peak (Figure 3, Figure S5). There is no

visible xylem differentiation at the graft junction during the first peak of expression

(4) and these genes might be suppressed by phloem differentiation genes such as *APL*

and *CLE41* that activate early after grafting and are known to suppress protoxylem

formation (39, 40)(Figure S6)(Table S4). Alternatively, the first peak could represent

programmed-cell death that does not lead to xylem differentiation. The second

expression peak of xylem-expressed genes at 120 hours occurred after the

differentiation of functional phloem and coincided with the differentiation of xylem

strands at the graft junction (4). Previous studies highlighted the importance of callus

and pericycle cells during regeneration (16, 41), but we see little evidence that genes

expressed in the pericycle or during callus formation have high transcriptional overlap

with genes differentially expressed by grafting (Figure S4). Expression profiles for all

protein-coding genes can be found in table S4.

We observe only a slight delay in phloem, cambium and cell division activation

below the graft junction compared to above it (Figure 3, Figure S5, Figure S6)

whereas several genes associated with vascular formation, such as *HCA2* (42) and

TMO6 (43)*, activated equally in both grafted top and grafted bottom at 6 hours after

grafting (Figure 1D-G, Figure 3). These data indicate that, at least transcriptionally,

the grafted root rapidly responded to the presence of the grafted shoot and this

response was independent of functional vascular connections. This response was not

present in separated roots, indicating that attachment was key for recognition. Sugars

are known activators of cell division and cell elongation (13) and in our datasets, a

large proportion of asymmetrically expressed genes are genes that are sugar-

responsive. However, sugars are transported in the phloem (12) that is severed upon

grafting whereas the grafted root exhibited a sugar starvation response and showed

similar sugar-response dynamics as the separated root. Instead, we infer that not sugar

but auxin, or some other molecular that is transported in the absence of vascular

connections, could be the signal in the grafted bottom that activates *HCA2*, *TMO6* as

well as cell division, phloem- and cambium-related genes. Alternatively, separation

might cause a build-up of molecules from the root that suppress vascular induction

genes such as *HCA2* and *TMO6*.

Given auxin's role in vascular patterning and formation ((38), it is a strong candidate

for the activating signal. Auxin response was largely symmetric, particularly from 12

hours after grafting (Figure 6) consistent with previous findings that the auxin-

inducible *DR5* and *ANAC071* genes are symmetrically expressed around the graft

junction (4-6, 44). One idea is that auxin transport was not substantially interrupted by

the grafting process, and instead, where opposing tissues adhered, auxin moved

regardless of vascular connections since auxin is transported from cell to cell through

the apoplast (8). The genes encoding auxin efflux proteins PIN1 and ABCB1 were

transcriptionally activated above the graft junction (Figure 5D), similar to putative

Pisum sativum PIN1 protein accumulating above a cut stem prior to vascular

reconnection (45), and could reflect a role for these proteins in exporting auxin across

the cut. Consistent with these observations, adding an auxin transport inhibitor to

grafted *Arabidopsis* shoots prevented the expression of grafting-induced genes below

the graft junction (6). Since auxin is fundamental for patterning vasculature (38), it is

plausible that auxin movement across the graft occurs prior to sugar movement since

the flow of auxin could assist with reconnecting the phloem. Although auxin response

was symmetric, our previous work demonstrated that the auxin response factors *ALF4*

and *AXR1* affected grafting only below the graft junction (4). Mutating *ALF4* below

the graft junction more strongly reduced auxin response than mutating *ALF4* above

the junction (4). Thus, proteins such as ALF4 or AXR1 might enhance or promote

rootstock-specific auxin response and vascular regeneration that could be particularly

important when there is incomplete attachment, cellular damage or inefficient

transport. All higher plants transport auxin from shoot to root, yet not all plant species

can be successfully grafted (3) so the response to auxin rather than the transport itself

may be a determining factor in the ability to graft. A role for sugars is not completely

ruled out though, since the magnitude of differential expression was often lower in the

grafted bottom and exogenous sugar reduced grafting efficiency (Figure 2A).

Pericycle cells require auxin to divide and the addition of sugars enhances the rate of

divisions (46), suggesting that the presence of sugars in the grafted top enhanced cell

division and differentiation. Another hypothesis that warrants testing is whether graft

healing activates via a mechanical signal provided by the physical presence of the

opposing tissues.

Our analyses identified two groups of genes whose expression changes were unique

to graft formation in our experiments (Figure 7). One group activated shortly after

grafting below the graft junction and was enriched in immune-responsive and chitin-

responsive genes (Table S3, Figure 7, Figure S10). The breakdown products of cell

walls are potent elicitors of defence responses (47), so it is possible that the grafted

bottom upregulates pathways specific to wound damage response. This group was not

upregulated in separated bottoms though, so the unique physiological state of the

grafted root, indicated by the presence of auxin response but the absence of sugar

response, could have promoted their upregulation. The second group activated both

above and below the graft junction and became highly expressed later during graft

formation (Figure 7). This group was enriched in RNAs associated with vascular

development (Table S3, Figure 7) and we suggest that the products of these genes are

involved in the vascular reconnection processes between the two tissues. Despite

many transcriptional similarities between separated tops and grafted tissues, tissues

had to be attached for at least three days for phloem connections to form, regardless

of when cutting occurred (Figure S10). Thus, it appears that RNAs expressed in the

separated top or separated bottom are insufficient to drive graft formation. Instead

attachment and the genes activated by this recognition process including grafting-

specific RNA changes (Figure 7A) or RNAs expressed in the grafted bottom, grafted

top and separated top (Figure 6C) are those that contribute to a response that

distinguishes attached from separated plant tissues. Future work should focus on these

RNAs to identify the pathways required for grafting that could be modified to

improve graft formation, wound healing and vascular regeneration. Likewise, the

rapid transcriptional changes below the graft indicate a recognition system that

promotes tissue regeneration. Identifying the exogenous cue that triggers recognition

and understanding how it is perceived should be priorities, as should understanding

whether this phenomenon applies more broadly to inter-tissue communication, tissue

regeneration or tissue fusion events, such as parasitic plant infections (48), epidermal

fusions (49, 50) or petal fusions (51).

METHODS

Plant material and microscopy

Arabidopsis thaliana accession Columbia was used throughout. The *p35S::GFP-ER*

(52), *pSUC2::GFP* (53), *pUBQ10::PM-tdTomato* (54), *pARR5::GFP* (55),

pANT::H2B-YFP (56), *pLOG4::n3GFP* (57), *pCASP1::NLS-GFP* (58), *p35S::DII-*

Venus (35) lines have been previously published. For the construction of

pHCA2::RFP, a 2.9kb 5' upstream region of the *HCA2* gene was cloned into

pDONRp4-p1R donor vector, and recombined with tagRFPer into a destination vector

by the Multisite Gateway system (59). *Arabidopsis thaliana* micrografting and

grafting assays were performed according to previously published protocols (60, 61).

Fluorescent images were taken on a Zeiss LSM-700 or LSM-780 confocal

microscope. Black and white fluorescent images of graft junctions were taken on a

Zeiss V12 dissecting microscope. FIJI software (Fiji.sc) was used to process images.

A micro-extensometer was used for all breaking force graft measurements according

to a previously published protocol (62).

RNAseq sample and library preparation

Grafted wild type *Arabidopsis thaliana* accession Col-0 were harvested at the

respective time points and care taken to separate grafts by gently pulling plants apart.

Approximately 0.5mm of tissue was taken above or below each cut site and kept

separate. Intact plants had 1mm of tissue taken in a similar location on the hypocotyl

as separated or grafted plants. Grafted, separated or intact tissues were pooled into

groups of approximately 24 tissues . Tissues were ground using a microcentrifuge

pestle frozen in liquid nitrogen. RNA was extracted using an RNeasy Kit (Qiagen,

UK) following the manufacturer's instructions. 90-100ng of RNA was used to prepare

RNAseq libraries using the TruSeq® Stranded mRNA LT kit (Illumina, UK)

according to the manufacturer's instructions. The final PCR was for 15 cycles and 11-

12 barcoded samples were randomly mixed to make a total of 7 mixes for 7 flow

lanes, one mix per lane. Samples were sequenced on the HiSeq 4000 platform

(Illumina, UK) with Paired End 75bp transcriptome sequencing (BGI Tech Solutions,

Shenzhen, China).

Iodine staining

Arabidopsis seedlings were placed in a fixation solution (3.7% formaldehyde, 50%

ethanol, 5% acetic acid) for 1 hour at room temperature, then transferred to 70%

ethanol for 10 minutes. Afterwards, plants were transferred to 96% ethanol and stored

at -20°C for up to a week. Samples were rehydrated in 50% ethanol for 1 hour at

room temperature, transferred to distilled water for 30 minutes, then stained for 10

minutes in Lugol solution (Sigma) at room temperature. Plants were rinsed with water

and mounted on microscope slides. Images were taken on a Zeiss Axioimager.M2

microscope.

Bioinformatic analyses

The reads acquired through high-throughput sequencing were quality trimmed with

sickle (63) and aligned using the eXpress tool to protein-coding gene sequences

acquired from TAIR10 using Bowtie2. Library scaling factors were inferred from the

sum of the number of reads assigned to the genes in the lowest seventy-five

percentiles of expressed genes for each library (64). Analyses of the data were carried

out using the R package baySeq (36)and clustering based on the posterior

probabilities acquired from this package. The gene ontology analysis (GO)

enrichment analysis on grafting-specific genes was done with a customized R script

using the package GOstats (65).

ACKNOWLEDGEMENTS

We thank Niko Geldner, Dolf Weijers, Paul Tarr, Yka Helariutta, Ruth Stadler and

The Arabidopsis Information Resource for providing seeds. Funding for this work

was provided by Gatsby Charitable Trust grants GAT3272/C and GAT3273-PR1, by

a Knut and Alice Wallenberg Academy Fellowship KAW2016.0274 (to C.W.M), and

by the Howard Hughes Medical Institute and Gordon and Betty Moore Foundation

grant GBMF3406 (to E.M.M.)

REFERENCES

1. Goldschmidt EE (2014) Plant grafting: new mechanisms, evolutionary implications. *Front Plant Sci* 5:727.
2. Lee J, *et al.* (2010) Current status of vegetable grafting: Diffusion, grafting techniques, automation. *Sci Hortic-Amsterdam* 127:93-105.
3. Melnyk CW (2017) Plant grafting: insights into tissue regeneration. *Regeneration (Oxf)* 4(1):3-14.
4. Melnyk CW, Schuster C, Leyser O, & Meyerowitz EM (2015) A developmental framework for graft formation and vascular reconnection in Arabidopsis thaliana. *Curr Biol* 25(10):1306-1318.
5. Yin H, *et al.* (2012) Graft-union development: a delicate process that involves cell-cell communication between scion and stock for local auxin accumulation. *J Exp Bot* 63(11):4219-4232.
6. Matsuoka K, *et al.* (2016) Differential Cellular Control by Cotyledon-Derived Phytohormones Involved in Graft Reunion of Arabidopsis Hypocotyls. *Plant Cell Physiol* 57(12):2620-2631.
7. Matsumoto-Kitano M, *et al.* (2008) Cytokinins are central regulators of cambial activity. *P Natl Acad Sci USA* 105(50):20027-20031.
8. Leyser O (2011) Auxin, self-organisation, and the colonial nature of plants. *Curr Biol* 21(9):R331-337.
9. Asahina M, *et al.* (2011) Spatially selective hormonal control of RAP2.6L and ANAC071 transcription factors involved in tissue reunion in Arabidopsis. *P Natl Acad Sci USA* 108(38):16128-16132.
10. Wetmore RH & Rier JP (1963) Experimental induction of vascular tissues in callus of angiosperms. *Am J Bot* 50(5):418-430.
11. Aloni R (1980) Role of auxin and sucrose in the differentiation of sieve and tracheary elements in plant tissue cultures. *Planta* 150(3):255-263.
12. Lough TJ & Lucas WJ (2006) Integrative plant biology: Role of phloem long-distance macromolecular trafficking. *Annu Rev Plant Biol* 57:203-232.
13. Wang L & Ruan YL (2013) Regulation of cell division and expansion by sugar and auxin signaling. *Front Plant Sci* 4:163.
14. Chitwood DH, Guo M, Nogueira FT, & Timmermans MC (2007) Establishing leaf polarity: the role of small RNAs and positional signals in the shoot apex. *Development* 134(5):813-823.
15. Cheong YH, *et al.* (2002) Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in Arabidopsis. *Plant Physiol* 129(2):661-677.
16. Iwase A, *et al.* (2011) The AP2/ERF transcription factor WIND1 controls cell dedifferentiation in Arabidopsis. *Curr Biol* 21(6):508-514.
17. Ikeuchi M, *et al.* (2017) Wounding triggers callus formation via dynamic hormonal and transcriptional changes. *Plant Physiol*.
18. Cookson SJ, *et al.* (2013) Graft union formation in grapevine induces transcriptional changes related to cell wall modification, wounding, hormone signalling, and secondary metabolism. *J Exp Bot* 64(10):2997-3008.
19. Cookson SJ, *et al.* (2014) Heterografting with nonself rootstocks induces genes involved in stress responses at the graft interface when compared with autografted controls. *J Exp Bot* 65(9):2473-2481.
20. Zheng BS, *et al.* (2010) cDNA-AFLP analysis of gene expression in hickory (Carya cathayensis) during graft process. *Tree Physiol* 30(2):297-303.

21. Chen Z, *et al.* (2017) Transcriptome changes between compatible and
incompatible graft combination of Litchi chinensis by digital gene expression
profile. *Sci Rep* 7(1):3954.
22. Lindsay DW, Yeoman MM, & Brown R (1974) An analysis of the
development of the graft union in Lycopersicon esculentum. *Ann Bot-London*
38:639-646.
23. Moore R (1984) Graft formation in Solanum pennellii (Solanaceae). *Plant*
Cell Rep 3(5):172-175.
24. Marsch-Martinez N, *et al.* (2013) An efficient flat-surface collar-free grafting
method for Arabidopsis thaliana seedlings. *Plant Methods* 9(1):14.
25. Villadsen D & Smith SM (2004) Identification of more than 200 glucose-
responsive Arabidopsis genes none of which responds to 3-O-methylglucose
or 6-deoxyglucose. *Plant Mol Biol* 55(4):467-477.
26. Thum KE, Shin MJ, Palenchar PM, Kouranov A, & Coruzzi GM (2004)
Genome-wide investigation of light and carbon signaling interactions in
Arabidopsis. *Genome Biol* 5(2):R10.
27. Cordoba E, Aceves-Zamudio DL, Hernandez-Bernal AF, Ramos-Vega M, &
Leon P (2015) Sugar regulation of SUGAR TRANSPORTER PROTEIN 1
(STP1) expression in Arabidopsis thaliana. *J Exp Bot* 66(1):147-159.
28. Osuna D, *et al.* (2007) Temporal responses of transcripts, enzyme activities
and metabolites after adding sucrose to carbon-deprived Arabidopsis
seedlings. *Plant J* 49(3):463-491.
29. Kondo Y, *et al.* (2016) Vascular Cell Induction Culture System Using
Arabidopsis Leaves (VISUAL) Reveals the Sequential Differentiation of
Sieve Element-Like Cells. *Plant Cell* 28(6):1250-1262.
30. Furuta KM, *et al.* (2014) Plant development. Arabidopsis NAC45/86 direct
sieve element morphogenesis culminating in enucleation. *Science*
345(6199):933-937.
31. Czechowski T, Stitt M, Altmann T, Udvardi MK, & Scheible WR (2005)
Genome-wide identification and testing of superior reference genes for
transcript normalization in Arabidopsis. *Plant Physiol* 139(1):5-17.
32. Brady SM, *et al.* (2007) A high-resolution root spatiotemporal map reveals
dominant expression patterns. *Science* 318(5851):801-806.
33. Nemhauser JL, Hong F, & Chory J (2006) Different plant hormones regulate
similar processes through largely nonoverlapping transcriptional responses.
Cell 126(3):467-475.
34. Abel S, Nguyen MD, & Theologis A (1995) The PS-IAA4/5-like family of
early auxin-inducible mRNAs in Arabidopsis thaliana. *J Mol Biol* 251(4):533-
549.
35. Brunoud G, *et al.* (2012) A novel sensor to map auxin response and
distribution at high spatio-temporal resolution. *Nature* 482(7383):103-106.
36. Hardcastle TJ & Kelly KA (2010) baySeq: empirical Bayesian methods for
identifying differential expression in sequence count data. *BMC*
Bioinformatics 11:422.
37. Hardcastle TJ & Papatheodorou I (2017) clusterSeq: methods for identifying
co-expression in high-throughput sequencing data. *bioRxiv* 188581.
38. De Rybel B, Mahonen AP, Helariutta Y, & Weijers D (2016) Plant vascular
development: from early specification to differentiation. *Nat Rev Mol Cell*
Biol 17(1):30-40.

39.   Bonke M, Thitamadee S, Mahonen AP, Hauser MT, & Helariutta Y (2003)
743   APL regulates vascular tissue identity in Arabidopsis. *Nature* 426(6963):181-
744   186.

745   40.   Ito Y*, et al.* (2006) Dodeca-CLE peptides as suppressors of plant stem cell
746   differentiation. *Science* 313(5788):842-845.

747   41.   Sugimoto K, Jiao Y, & Meyerowitz EM (2010) Arabidopsis regeneration from
748   multiple tissues occurs via a root development pathway. *Dev Cell* 18(3):463-
749   471.

750   42.   Guo Y, Qin G, Gu H, & Qu LJ (2009) Dof5.6/HCA2, a Dof transcription
751   factor gene, regulates interfascicular cambium formation and vascular tissue
752   development in Arabidopsis. *Plant Cell* 21(11):3518-3534.

753   43.   Gardiner J, Sherr I, & Scarpella E (2010) Expression of DOF genes identifies
754   early stages of vascular development in Arabidopsis leaves. *Int J Dev Biol*
755   54(8-9):1389-1396.

756   44.   Pitaksaringkarn W, Ishiguro S, Asahina M, & Satoh S (2014) ARF6 and
757   ARF8 contribute to tissue reunion in incised Arabidopsis inflorescence stems.
758   *Plant Biotechnol-Nar* 31(1):49-53.

759   45.   Sauer M*, et al.* (2006) Canalization of auxin flow by Aux/IAA-ARF-
760   dependent feedback regulation of PIN polarity. *Genes Dev* 20(20):2902-2911.

761   46.   Skylar A, Sung F, Hong F, Chory J, & Wu X (2011) Metabolic sugar signal
762   promotes Arabidopsis meristematic proliferation via G2. *Dev Biol* 351(1):82-
763   89.

764   47.   Souza CA*, et al.* (2017) Cellulose-Derived Oligomers Act as Damage-
765   Associated Molecular Patterns and Trigger Defense-Like Responses. *Plant
766   Physiol* 173(4):2383-2398.

767   48.   Musselman LJ (1980) The biology of Striga, Orobanche, and other root-
768   parasitic weeds. *Annual Review of Phytopathology* 18:463-489.

769   49.   Lolle SJ, Hsu W, & Pruitt RE (1998) Genetic analysis of organ fusion in
770   Arabidopsis thaliana. *Genetics* 149(2):607-619.

771   50.   Becraft PW, Stinard PS, & McCarty DR (1996) CRINKLY4: A TNFR-like
772   receptor kinase involved in maize epidermal differentiation. *Science*
773   273(5280):1406-1409.

774   51.   Zhong J & Preston JC (2015) Bridging the gaps: evolution and development
775   of perianth fusion. *New Phytol* 208(2):330-335.

776   52.   Nelson BK, Cai X, & Nebenfuhr A (2007) A multicolored set of in vivo
777   organelle markers for co-localization studies in Arabidopsis and other plants.
778   *The Plant journal* 51(6):1126-1136.

779   53.   Imlau A, Truernit E, & Sauer N (1999) Cell-to-cell and long-distance
780   trafficking of the green fluorescent protein in the phloem and symplastic
781   unloading of the protein into sink tissues. *Plant Cell* 11(3):309-322.

782   54.   Segonzac C*, et al.* (2012) The shoot apical meristem regulatory peptide CLV3
783   does not activate innate immunity. *Plant Cell* 24(8):3186-3192.

784   55.   Yanai O*, et al.* (2005) Arabidopsis KNOXI proteins activate cytokinin
785   biosynthesis. *Curr Biol* 15(17):1566-1571.

786   56.   Randall RS*, et al.* (2015) AINTEGUMENTA and the D-type cyclin CYCD3;1
787   regulate root secondary growth and respond to cytokinins. *Biol Open*.

788   57.   De Rybel B*, et al.* (2014) Plant development. Integration of growth and
789   patterning during vascular tissue formation in Arabidopsis. *Science*
790   345(6197):1255215.

58.   Roppolo D, *et al.* (2011) A novel protein family mediates Casparian strip
formation in the endodermis. *Nature* 473(7347):380-383.

59.   Siligato R, *et al.* (2016) MultiSite Gateway-Compatible Cell Type-Specific
Gene-Inducible System for Plants. *Plant Physiol* 170(2):627-641.

60.   Melnyk CW (2017) Grafting with Arabidopsis thaliana. *Methods Mol Biol*
1497:9-18.

61.   Melnyk CW (2017) Monitoring Vascular Regeneration and Xylem
Connectivity in Arabidopsis thaliana. *Methods Mol Biol* 1544:91-102.

62.   Robinson S, *et al.* (2017) An automated confocal micro-extensometer enables
in vivo quantification of mechanical properties with cellular resolution.
bioRxiv 183533.

63.   Joshi NA & Fass JN (2011) Sickle: A sliding-window, adaptive, quality-based
trimming tool for FastQ files (Version 1.33)).

64.   Hardcastle TJ, Kelly KA, & Baulcombe DC (2012) Identifying small
interfering RNA loci from high-throughput sequencing data. *Bioinformatics*
28(4):457-463.

65.   Falcon S & Gentleman R (2007) Using GOstats to test gene lists for GO term
association. *Bioinformatics* 23(2):257-258.

66.   Drost HG & Paszkowski J (2017) Biomartr: genomic data retrieval with R.
Bioinformatics 33(8):1216-1217.

67.   Carlson M (2017) org.At.tair.db: Genome wide annotation for Arabidopsis), R
package version 3.4.1.

 SUPPORTING INFORMATION

 SI Materials and Methods

Plant material and grafting

Arabidopsis thaliana accession Columbia was used throughout. The *p35S::GFP-ER*

(52), *pSUC2::GFP* (53), *pUBQ10::PM-tdTomato* (54), *pARR5::GFP* (55),

pANT::H2B-YFP (56), *pLOG4::n3GFP* (57), *pCASP1::NLS-GFP* (58), *p35S::DII-*

Venus (35) lines have been previously published. For the construction of

pHCA2::RFP, a 2.9kb 5' upstream region of the *HCA2* gene was cloned into

pDONRp4-p1R donor vector, and recombined with tagRFPer into a destination vector

by the Multisite Gateway system (59). The following primers were used for *pHCA2*

cloning: attB4_HCA2(-)2958

GGGACAACTTTGTATAGAAAGTTGtcgatacgcgggacagatatac

attB1_HCA2ProEnd ggggACTGCTTTTTTGTACAAACTTGttttgtgttctgtatgtttg.

Arabidopsis thaliana micrografting was performed according to a previously

published protocol (60). Briefly, seven day old *Arabidopsis* seedlings were grown

vertically on Murashige and Skoog (MS) medium + 1% bacto agar (pH5.7; no

sucrose) in short day conditions (8 hours of 80-100 µmol m^{-2} s^{-1} light) at 20°C.

Seedlings were placed on one layer of 2.5x4cm sterile Hybond N membrane (GE

Healthcare) on top of two 8.5cm circles of sterile 3 Chr Whatman paper (Scientific

Laboratory Supplies) moisten with sterile distilled water in a 9cm petri dish. In a

laminar flow hood using a dissecting microscope, one cotyledon was removed and a

transvers cut through the hypocotyl was made with a vascular dissecting knife (Ultra

Fine Micro Knife; Fine Science Tools). Grafts were assembled by aligning the two

cut halves and joining them together, after which, the petri dishes was sealed with

parafilm and placed vertically under short day conditions at 20°C. For grafting on a

microscope coverslip to image the graft junction, a 10cm square Petri dish modified

by gluing a microscope coverslip in place of a section of plastic from the back. On top

of the microscope cover slip was placed a 2.5x4 cm rectangle of Hybond N

membrane. At the edges and base of the Petri dish three 3 x 8cm strips of Whatman

paper were placed. Sterile water moistened both Whatman paper and Hybond N.

After which, roots were placed on the Hybond N membrane and hypocotyls on the

coverslip. Grafting then proceeded as above. Graft junctions were imaged through the

coverslip with a Plan-Apochromat 20X/0.8 objective on a Zeiss LSM-700 or LSM-

780 confocal microscope.

Fluorescent assays and microscopy

To test the effect of sugars on grafting, *Arabidopsis thaliana* Col-0 plants were grown

on 1/2MS+ 1% bacto agar (pH5.7; no sucrose) for seven days in short day conditions.

Wild type roots were grafted to scions expressing *pSUC2::GFP* using the protocol

described above but either water or water containing 0.25%, 0.5% 1% or 2% sucrose

was added to the grafted plates. Roots were observed for fluorescence 2-7 days after

grafting with a Zeiss V12 dissecting microscope equipped with a GFP filter. Roots

were scored daily and the same plants were observed during the 7-day assay as

previously described (61).

Fluorescent images were taken on a Zeiss LSM-700 or LSM-780 confocal microscope

with a Zeiss Plan-Apochromat 20X/0.8 dry objective. A 488nm argon laser (Zeiss

780) or 488nm solid-state laser (Zeiss 700) was used for excitation of GFP and YFP.

A 561nm solid-state laser was used for excitation of the tdTomato fluorescent protein.

A T-PMT detector obtained bright-field transmitted light. Black and white fluorescent

images of graft junctions were taken on a Zeiss V12 dissecting microscope fitted with

a Hamamatsu EM-CCD camera and RFP and YFP filters. FIJI software (Fiji.sc) was

used to process images. Image contrast and brightness were adjusted for controls and

samples equally. For longitudinal images of the graft junction, z-stack projections are

shown and made with the average intensity function in FIJI from stacks containing the

hypocotyl vascular tissues, mesophyll and epidermis.

RNAseq sample and library preparation

Wild type *Arabidopsis thaliana* accession Col-0 were grafted as above taking care to

switch shoot and root between different plants. All grafting and cutting was

performed in the morning to minimize circadian effects. For the 0 hour time points,

plants were transferred from 1/2MS plates to the grafting plates and immediately

harvested. Only intact plants were harvested at the 0 hour time since at this point,

there would be insufficient time to reasonably expect the separated or grafted samples

to be transcriptionally different (the time between cutting and freezing is less than two

minutes). For all other time points, after cutting plants were left on grafting plates for

the respective amount of time. Tissues were harvested and care taken to separate

grafts by gently pulling plants apart. Approximately 0.5mm of tissue was taken above

or below each cut site and kept separate. Intact plants had 1mm of tissue taken in a

similar location on the hypocotyl as separated or grafted plants. Grafted, separated or

intact tissues were pooled into groups of approximately 24 tissues (1 plates with 24

plants) which were immediately placed in 96% ethanol on dry ice. After harvesting,

microcentrifuge tubes were briefly centrifuged and the ethanol removed before

storing at -80°C. Plants were grafted over two months to get sufficient material.

Tissues were ground in the microcentrifuge tube using a microcentrifuge pestle frozen

in liquid nitrogen. RNA was extracted using an RNeasy Kit (Qiagen, UK) following

the manufacturer's instructions including on column DNase digestion. RNA was

eluted from the column with 50ul of sterile water. Quality and quantity of RNA was

checked using an Agilent 2200 TapeStation and High Sensitivity (HS) RNA

screentapes (Agilent, UK). After RNA extraction, two to four biological replicates

were combined (50-100 plants) to get sufficient RNA. 90-100ng of RNA was used to

prepare RNAseq libraries using the TruSeq® Stranded mRNA LT kit (Illumina, UK)

according to the manufacturer's instructions. The final PCR was for 15 cycles and

samples were resuspended in 23ul of distilled water. Quantity and quality of DNA

libraries was checked on the Agilent 2200 TapeStation using D1000 screentapes

(Agilent, UK). Each sample had two libraries prepared from grafted tissues or

separated tissues at different times so that independent biological replicates were

made. Samples were diluted to 10nM and 11-12 barcoded samples randomly mixed to

make a total of 7 mixes for 7 flow lanes, one mix per lane. Samples were sequenced

on the HiSeq 4000 platform (Illumina, UK) with Paired End 75bp transcriptome

sequencing (BGI Tech Solutions, Shenzhen, China).

Iodine staining

Arabidopsis seedlings were placed in a fixation solution (3.7% formaldehyde, 50%

ethanol, 5% acetic acid) for 1 hour at room temperature, then transferred to 70%

ethanol for 10 minutes. Afterwards, plants were transferred to 96% ethanol and stored

at -20°C for up to a week. Samples were rehydrated in 50% ethanol for 1 hour at

room temperature, then transferred to distilled water for 30 minutes. Samples were

then transferred to a solution of Lugol solution (Sigma) and stained for 10 minutes at

room temperature. Plants were rinsed with water, then mounted on microscope slides.

Images were taken on a Zeiss Axioimager.M2 microscope with a PlanApochromat

20x objective and SPOT Flex camera (Imsol, UK).

Pairwise and Bayseq analyses

The reads acquired through high-throughput sequencing were quality trimmed with

sickle (63) to increase the read quality before mapping. Reads were aligned to

protein-coding gene sequences acquired from TAIR10 using Bowtie2. Read

assignment was performed using the eXpress tool, which also provided effective gene

lengths for use in normalisation. Library scaling factors were inferred from the sum of

the number of reads assigned to the genes in the lowest seventy-five percentiles of

expressed genes for each library (64).

Analyses of the data were carried out using the R package baySeq (36)and clustering

based on the posterior probabilities acquired from this package. For each timepoint,

all possible patterns of differential expression between the graft types were

considered, where a 'pattern' defines similarity and difference between different

experimental conditions. For example,

'{Col_cut_bottomGenes=Col:Col_bottomGenes=ungraftedGenes},{Col_cut_topGenes=Col:Col_topGenes}'

defines a pattern in which gene expression is equivalent in the separated bottoms, the

grafted bottoms and the intact plant, but different to the equivalently expressed

separated top and grafted top. The total number of possible patterns for five

experimental conditions (as in this analysis) is fifty-two.

For a given timepoint, posterior likelihoods on the likelihood of each pattern of

expression are calculated for every gene with greater than ten reads across all

experimental conditions. The patterns were then modified to include orderings

(denoted by < or >), for example, the pattern described would lead to the ordered

pattern

'{Col_cut_bottomGenes=Col:Col_bottomGenes=ungraftedGenes}>{Col_cut_topGenes=Col:Col_topGenes}'

in which gene expression is equivalent in the separated bottoms, the grafted bottoms

and the intact plant and greater than the equivalently expressed separated top and

grafted top. In total, 541 ordered patterns exist in this data set. Posterior likelihoods

for an ordered pattern were assigned to that of the unordered pattern for genes in

which the (normalised) mean expressions within the equivalently expressed groups

conformed to the ordering, and to zero otherwise.

Based on the posterior likelihoods for the ordered patterns, a similarity score s_{ij} was

established between two genes i and j as the sum over the products of their likelihoods

of each ordered pattern. A single-link agglomerative clustering of genes, in which a

gene will join a cluster if it has a greater than 50% similarity to any gene within that

cluster was then performed based on these similarity scores. We label each cluster

according to the predominant ordered pattern with high likelihood amongst the genes

that comprise it. The change in size of these clusters over time is shown for the major

clusterings in Figure 6.

We can also find likelihoods on comparisons between pairs of experimental

conditions by summing the likelihoods over combinations of patterns. Figure 1B

shows the number of genes identified at each time point in a pairwise analysis

between the grafted top and grafted bottom samples. The likelihood of symmetric

expression (i.e., expression which is equivalent across the graft junction) is calculated

as the sum of the likelihoods of all patterns in which the grafted top and grafted

bottom samples are equivalent. Conversely, asymmetric expression is calculated as

the sum of the likelihoods of all patterns in which the grafted top and grafted bottom

samples are not equivalent. Additional sets can be formed by considering the ordering

of the grafted top and grafted bottoms samples. Sets of genes are identified at each

time point with an FDR of less than 0.05 and a likelihood of symmetric/asymmetric

expression greater than 50%. Genes in this analysis were only included if they were

differentially expressed relative to intact samples.

Gene overlap analyses of up and down regulated genes

To measure if the ratio of up- and down- regulated genes from a previously published

dataset is significantly different to the ratio of up- and down-regulated genes in our

grafting dataset we only took into account genes that are differentially expressed at a

certain time point. A gene was called differentially expressed at a certain time point if

the marginal likelihood, calculated by BaySeq, was greater than 0.9 and if the

absolute $\log_2$-foldchange was greater than 1. Hence, we only consider genes that are

significantly two-fold up- or down- regulated. This definition of differentially

expressed genes was also used to filter the published datasets according to the

expression values in our transcriptome dataset. Hence, some genes were filtered out

from the original published datasets because they did not show a significant up- or

down- regulation during a certain time point in our expression data based on our

criteria. The histograms (Figure 2,4, 5, S4, S7, S8, S9) show the relative number of

up- and down- regulated genes from the published datasets during a certain time point

and a certain condition (separated top, separated bottom, grafted top, grafted bottom)

based on the number of genes in the published dataset after filtering. To calculate the

significance of the difference of the ratios between the published DEGs and all up-

and down- regulated genes, we used a two-sided Fisher's exact test. To correct for

multiple testing we used the Benjamini-Yekutieli (BY) correction method. Hence, the

asterisks in the barplots highlight that the corrected p-value is below 0.05.

1013

Dealing with probe ids from microarray datasets

Due to the fact that some published datasets only used probe ids instead of gene ids to

represent their differentially expressed genes we first had to match these probe ids to

their corresponding gene ids. This step was done with the R package biomartr (66). If

one probe id matched more than one gene id we used all the corresponding gene ids

and tested afterwards if these genes were actually differentially expressed in our

dataset. In some cases, one probe id was represented by more than one gene id.

Hence, some gene sets contained slightly more gene ids than published probe ids. In

contrast, some probe ids did not match to currently existing gene ids. Hence, some

gene sets contained slightly fewer gene ids than published probe ids.

1024

Gene overlap analyses of gene sets involved in graft formation

For this analysis we used the previously calculated gene sets from baySeq of

differentially expressed genes clustered for each time point into grafted bottom,

grafted top, and grafted bottom and top and calculated their overlap to previously

published data sets. The significance of the number of overlapping genes between the

grafted samples and the published datasets was determined by a one-sided Fisher's

exact test, to prove if the overlap is greater than expected. The resulting p-values were

corrected for multiple testing by using the Benjamini-Yekutieli method.

This procedure was also applied to generate Table 1 to study the overlaps of

symmetrically and asymmetrically expressed genes in the grafting dataset with

previously published sugar-responsive genes.

1036

GO enrichment analysis

The gene ontology analysis (GO) enrichment analysis on grafting-specific genes was

done with a customized R script using the package GOstats (65). Gene ontology

annotation was used from the Bioconductor package org.At.tair.db (67). The p-values

calculated by a hypergeometric test were corrected for multiple testing with the

Bonferroni correction. A GO category was called enriched if the corrected p-value

was below 0.05.

1044

1045 A detailed description, the required data and R scripts to reproduce the clustering

1046 (hierarchical clustering and PCA), the statistical analyses regarding the overlap

1047 studies, and the GO enrichment analysis are available via the GitHub repository

1048 (https://github.com/AlexGa/GraftingScripts).

1049

Breaking force measurements

A micro-extensometer was used for all breaking force graft measurements according

to a previously published protocol (62). Briefly, grafted Col-0 plants were attached to

a moving plate using tough tags (0.94X0.50 inches, white, 679 catalog no. TTSW-

1000, DiversifiedBiotech) and a cyanoacrylate glue. The plates were moved apart and

the force was measured using a force sensor (Futek LSB200 10g load cell, Futek

Inc.). Upon breaking the force dropped. The maximum force measured before

breaking was recorded. Images were captured using a Leica SP5 confocal microscope.

During the experiment single z-place images were captured and made into a movie in

ImageJ.

1060

To measure levels of tissue contamination between grafted top and bottom, grafted

green fluorescent protein(*p35S::GFP*) and red fluorescent protein (*pUBQ10::PM-*

tdTomato) expressing plants were pulled apart manually and imaged on a Zeiss LSM

700 confocal microscope. Z-stacks comprising the majority of the hypocotyl were

made and regions approximately 0.5mm above and 0.5mm below the graft junction

were made into projections using the average intensity function on FIJI software, and

then the mean intensity quantified for red and green channels on FIJI. The mean

intensity of one colour in the tissue tested for contamination was divided by the mean

intensity of the same colour in the tissue expressing that transgenes to get a

percentage of contamination. A region away from the cut site was also quantified to

get a percentage of spectral overlap between red and green channels. The percent

spectral overlap was then subtracted from the percentage contamination to get an

overall percentage for how much contamination was present.

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

Movie S1. Breaking weight measurements at the graft junction, related to figure 1. A micro-extensometer pulled apart an Arabidopsis plant 9 days after grafting while force was measured using a force sensor and images captured using a Zeiss Sp5 confocal microscope. Here, the transmitted light field is shown. Shoot is on top and root towards the bottom.

Movie S2. Breaking weight measurements at the graft junction, related to figure 1. A micro-extensometer pulled apart an *Arabidopsis* plant 9 days after grafting while force was measured using a force sensor and images captured using a Zeiss Sp5 confocal microscope. The same experiment is shown in Movie S1, but here, the GFP and RFP channels are shown. Shoot is on top (*p35S::GFP* expressing) and root towards the bottom (*pUBQ10::PM-tdTomato* expressing).

Table S1. Details of previously published datasets used to compare to the grafting datasets.

Table S2. Numbers and categories associated with the Bayseq analysis. Categories are defined as grafted (Col:Col), separated (Col_cut) or intact (ungrafted).

Table S3. GO analysis for biological process (BP). Shown are the top 20 BP GO terms for the grafting-specific genes. Time point selected are those when there are the most genes in grafted bottom samples (48hrs) or grafted top + bottom samples (120 hrs).

Table S4. Normalised reads for all the protein-coding genes in the datasets. By entering the ATG number of interest, a plot is made which shows a differential gene expression profile for the gene of interest.

Dataset S1. Details of the Bayesian segmentation analysis providing files containing the ATGs for every cluster for every time point.

1132

Figure S1. Experimental approach, related to figure 1. (A) The breaking weight
required to pull apart grafted plants measured by a micro-extensometer. (B-C)
Contamination of top in bottom or bottom in top was less than 4% as measured by
grafting green fluorescent protein-expressing plants (*p35S::GFP*) to tomato
fluorescent protein-expressing plants (*pUBQ10::PM-tdTomato*) and measuring
amounts of fluorescence at the different wavelengths of emission in the top segments
relative to the bottom. Images show different plants prior to and after pulling with the
percent contamination indicated.

1141

1142

Figure S2. Comparison between RNAseq transcriptome expression profiles and transcriptional fluorescent reporters, related to figure 1. RNAseq expression profiles for various genes upregulated during graft formation were plotted for intact, separated and grafted samples (left panels). Transcriptional reporter-expressing plants were cut and separated, cut and grafted, or left intact. After cutting, plants were imaged and z-projections made at various time points (right panels). For *pCASP1::GFP*, we did not observe a signal in intact plants grafted on glass slides (see Materials and Methods), but observed a signal with 3/5 plants 7 days after grafting on Whatman-nylon membrane, the same condition used for transcriptome library preparation. HAG, hours after grafting. HAS, hours after separation. Dashed lines denote the graft junction.

1154

1155

1156

Figure S3. Hierarchical cluster analysis and principle component analysis (PCA)
for the transcriptome libraries, related to figure 1. (A) Hierarchical clustering of
samples based on log10 transformed TPM values. Similarity between samples was
measured by 1 – spearman correlation coefficient. (B) PCA of expression data shows
clustering of similar samples. Especially the grafted top and grafted bottom samples
are very similar from 120 hours onwards.

1163

Figure S4. Asymmetry is a feature observed with sugar response and starch accumulation, related to figure 2. (A) Expression profiles for RNAs coded by sugar-repressed genes (*DIN6*, *STP1*) or photosynthetic gene (*RBSC3B*) were plotted for intact, separated and grafted samples. (B) Transcriptional overlap between previously published mannitol-induced, mannitol-repressed, sugar-induced or sugar-repressed genes and our datasets. The numbers in brackets represents the number of glucose or mannitol-responsive genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentially expressed genes (DEG) up or down regulated in our dataset relative to intact samples. Asterisks represent a significant difference (p < 0.05) between the ratio of up- and down- regulated genes in a previously published transcriptome dataset compared to the ratio of all up- and down-regulated genes in our grafting dataset at a certain time point. (C) Lugol staining of grafted plants at various time points reveals dark brown staining associated with starch accumulation. Upper grafted panels are the same as those present in Figure 2 and shown here to compare with controls. HAG, hours after grafting.

1179

1180 **Figure S5. Transcriptional dynamics of genes associated with cambium, phloem,**
1181 **xylem and cell division, related to figure 3.** Expression profiles in grafted tops or
1182 grafted bottoms for various genes of interest were plotted, normalised to intact
1183 samples and plotted on a $\log_2$ scale.

1184

1185

1186

Figure S6. Accumulation of RNA from genes associated with provasculature and
phloem, and of RNAs from WOX transcription factors, related to figure 3. (A)
RNA accumulation profiles for various genes of interest were plotted over time as
measured in grafted tops or grafted bottoms, normalised to intact samples and plotted
on a $\log_2$ scale. The grafting datasets could also be used to investigate the
transcriptional dynamics of related genes, such as the sequential activation of WOX
transcription factors at the graft junction. (B) Expression levels of a primary root
specific transcript (of the *WOX5* gene) or a lateral root specific transcript (from the
LBD18 gene) were plotted over time for intact, separated and grafted samples.
1196

1197

Figure S7. Transcriptional overlap between previously published datasets and the grafting datasets, related to Figure 4. Genes whose transcripts are associated with various cell types or biological processes were taken from previously published datasets (see Table S1) and compared to the datasets generated here. The number in brackets represents the number of cell type-specific or process-specific genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or down-regulated in our dataset relative to intact samples. Asterisks represent a significant difference ($p < 0.05$) between the ratio of up- and down- regulated genes in a previously published transcriptome dataset compared to the ratio of all up- and down- regulated genes in our grafting dataset at a certain time point.

1209

Figure S8. Transcriptional overlap between previously published root
transcriptomic datasets and the grafting RNA-seq dataset, related to Figure 4.
Genes whose transcripts are associated with various root regions were taken from
previously published datasets (see Table S1) and compared to the datasets generated
here. Root layer is indicated in the bottom right of each histogram with reference to
the cartoon. The number in brackets represents the number of root layer-specific
genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0
for differentiation expressed genes (DEG) up- or down-regulated in our dataset
relative to intact samples. Asterisks represent a significant difference ($p < 0.05$)
between the ratio of up- and down- regulated genes in a previously published
transcriptome dataset compared to the ratio of all up- and down- regulated genes in
our grafting dataset at a certain time point.
1222

1223

1224

Figure S9. Transcriptional overlap between previously published hormone
responsive RNA datasets and the grafting RNA datasets, related to Figure 5.
Genes whose differential expression is associated with various hormone responses
were taken from previously published datasets (see Table S1) and compared to the
genes represented in the RNA-seq datasets generated. The number in brackets
represents the number of cell type-specific or process-specific genes identified in the
previous dataset, and overlap is presented as a ratio out of 1.0 for differentiation
expressed genes (DEG) up- or down-regulated in our dataset relative to intact
samples. Asterisks represent a significant difference ($p < 0.05$) between the ratio of
up- and down- regulated genes in a previously published transcriptome dataset
compared to the ratio of all up- and down- regulated genes in our grafting dataset at a
certain time point.

1237

1238

1239

Figure S10. A subset of genes is differentially expressed only during graft formation, related to Figure 7. (A) *pSUC2::GFP* expressing *Arabidopsis* shoots were grafted to Col-0 wild type roots 0-5 days after transferring to grafting plates and kept intact until grafting (intact treatment). GFP movement to the roots for phloem connection assays was monitored over 7 days. (B) Plants were cut at day 0 and grafted at days 0-5 after cutting. (C) Shoots were cut at day 0 and grafted at days 0-5 after cutting. Roots were kept intact until immediately before grafting. (D) Roots were cut at day 0 and grafted at days 0-5 after cutting. Shoots were kept intact until immediately before grafting. (E) Genes differentially expressed in grafted tops and grafted bottoms show overlap with previously published genes whose transcripts are associated with the endodermis, vascular induction and grafting (see Table S1 for treatment information). Asterisks represent significant high overlap (p <0.05) of previously published gene sets that are also differentially expressed in the grafted samples at a certain time point.

1254