1   **Gene expression dynamics and inter-tissue recognition mechanisms during**
2   **tissue fusion at the *Arabidopsis* graft junction**
3

4   Charles W Melnyk[1,2], Alexander Gabel[3], Thomas J Hardcastle[4], Sarah Robinson[5],
5   Shunsuke Miyashima[6], Ivo Grosse[3], Elliot M Meyerowitz[1,7]
6
1- Sainsbury Laboratory, University of Cambridge, Bateman Street, Cambridge
CB2 1LR, UK
2 – Department of Plant Biology, Swedish University of Agricultural Sciences,
Almas allé 5, 756 51 Uppsala, Sweden
3 - Institute of Computer Science, Martin Luther University Halle–Wittenberg,
06120 Halle (Saale), Germany
4 – Department of Plant Sciences, University of Cambridge, Downing Street,
Cambridge CB2 3EA, United Kingdom
5 - Institute of Plant Science, University of Bern, Altenbergrain 21, 3013 Bern,
Switzerland
6 – Graduate School of Biological Sciences, Nara Institute of Science and
Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan
7 – Howard Hughes Medical Institute and Division of Biology and Biological
Engineering, California Institute of Technology, 1201 East California Boulevard,
Pasadena, CA 91125, USA
22
23   Contact: charles.melnyk@slu.se

ABSTRACT

The ability for cut tissues to join together and form a chimeric organism is a

remarkable and agriculturally important property of many plants, however, it is

poorly characterized at the molecular level. To better understand grafting we

monitored temporal and spatial changes in gene expression in grafted tissues.

Tissues above and below the graft rapidly developed asymmetry such that many

genes were more highly expressed on one side than the other. This asymmetry

correlated with sugar responsive genes and we observed an accumulation of

starch above the graft that, after three days, decreased along with asymmetry

once vascular connections were re-established. Despite an initial starvation

response below the graft, many genes associated with cell division and vascular

formation were rapidly activated in grafted tissues but not in cut and separated

tissues indicating a recognition mechanism that was independent of functional

vascular connections. Auxin response in grafted plants differed from that in cut

and separated plants as response was enhanced and symmetric, suggesting that

auxin was perceived by the root within hours of reattachment to activated

vascular regeneration. Such a recognition process could have broader relevance

for wound healing, inter-tissue communication and tissue fusion events.

INTRODUCTION

For millennia people have cut and rejoined plants through grafting. Generating

such chimeric organisms combines desirable characteristics from two plants,

such as disease resistance, dwarfing and high yields, or can propagate plants and

avoid the delays entailed by a juvenile state (Goldschmidt, 2014). Agriculturally,

grafting is becoming more relevant as a greater number of plants and species are

grafted to increase productivity and yield (Lee et al., 2010). However, our

mechanistic understanding of grafting and the biological processes involved,

including wound healing, tissue fusion and vascular formation, remain limited.

Plants have efficient mechanisms to heal wounds and cuts, in part through the

production of wound-induced pluripotent cells termed callus. The callus fills the

gap or seals the wound, and later, differentiates to form epidermal, mesophyll

and vascular tissues (Melnyk, 2017c). In grafted *Arabidopsis* hypocotyls, tissues

adhere 1-2 days after grafting and the phloem, the tissue that transports sugars

and nutrients, connects after three days (Yin et al., 2012; Melnyk et al., 2015).

The xylem, tissue that transports water and minerals, connects after seven days

(Melnyk et al., 2015). Plant hormones are important regulators of vascular

formation, and at the graft junction, both auxin and cytokinin responses increase

in the vascular tissue (Yin et al., 2012; Melnyk et al., 2015; Matsuoka et al., 2016).

Auxin is important for differentiation of vascular tissues whereas cytokinin

promotes vascular stem cells, termed cambium, to divide and proliferate in a

process known as secondary growth (Matsumoto-Kitano et al., 2008; Leyser,

2011). Auxin is produced in the upper parts of a plant and moves towards the

roots via cell-to-cell movement. Auxin exporters, including the PIN proteins,

transport auxin into the apoplast, whereas auxin importers, such as the AUX and

LAX proteins, assist with auxin uptake into adjacent cells (Leyser, 2011).

Disrupting this transport, such as by mutating *PIN1*, inhibits healing of a

wounded stem (Asahina et al., 2011). Blocking auxin transport with the auxin

transport inhibitor TIBA (2,3,5-triiodobenzoic acid) in the shoot inhibits vascular

formation and cell proliferation at the *Arabidopsis* graft junction (Matsuoka et al.,

2016). In addition to auxin, other compounds, including sugars, contribute to

vascular formation. The localised addition of auxin to callus induces phloem and

xylem but requires the presence of sugar (Wetmore & Rier, 1963; Aloni, 1980).

In plants, sugars are produced in the leaves and transported through the phloem

to the roots (Lough & Lucas, 2006). The role of sugars in vascular formation and

wound healing is not well established, however, sugars promote cell division and

cell expansion (Wang & Ruan, 2013), processes important for development

including vascular formation.

The molecular and cellular mechanisms for wound healing, tissue reunion and

graft formation remain largely unknown. One emerging theme is that the top and

bottom of the cut do not behave similarly. Such tissue asymmetry occurs in other

plant tissues, most notably leaves. Developing leaf primordia have an inherent

asymmetry that is thought to derive from positional signals from the shoot apical

meristem that specifies differences between the top and the bottom of the leaf.

The molecular mechanisms that establish asymmetry are not well described,

though one hypothesis is that auxin contributes towards or acts on the putative

signal (Chitwood et al., 2007). Asymmetry also appears in cut *Arabidopsis*

inflorescence stems where the transcription factor *RAP2.6L* expresses

exclusively below the cut whereas the transcription factor *ANAC071* expresses

exclusively above the cut (Asahina et al., 2011). Both were important for stem

healing and *ANAC071* and a close homologue, *ANAC096*, were important for graft

formation (Matsuoka et al., 2016). Asymmetry also exists in genetic

requirements, since *ALF4* and *AXR1*, two genes involved in auxin perception, are

important below but not above the graft junction for phloem connection (Melnyk

et al., 2015). However, *ANAC071* is expressed symmetrically around the

hypocotyl graft junction three days after grafting (Matsuoka et al., 2016) so the

extent of asymmetry and the mechanistic basis for it during wound healing

remains largely uncharacterised.

Previous efforts have characterised wound healing and tissue reunion using

transcriptomic analyses. Mechanical wounding altered approximately 8% of the

Arabidopsis transcriptome and showed a high degree of overlap with

transcriptomic changes elicited by pathogen attack and abiotic stress (Cheong et

al., 2002). Stem wounding and wound-induced callus formation altered the

expression of hundreds or thousands of genes (Asahina et al., 2011; Iwase et al.,

2011; Ikeuchi et al., 2017), whereas grafting grape vines, lychee trees and

hickory trees induced hundreds or thousands of differentially expressed genes

involved in hormone response, wound response, metabolism, cell wall synthesis

and signal transduction (Zheng et al., 2010; Yin et al., 2012; Cookson et al., 2013;

Cookson et al., 2014; Chen et al., 2017). These grafting studies provide limited

information, as tissues from above and below the graft junction were not isolated

to test whether these tissues behaved differently, and controls were not

performed to distinguish how grafting and tissue fusion might differ from a

response associated with cut tissues that remained separated. Here, we perform

an in-depth analysis to describe the spatial and temporal transcriptional

dynamics that occur during healing of cut *Arabidopsis* tissues that are joined

(grafted) or left unjoined (separated). We find that the majority of genes

differentially expressed are initially asymmetrically expressed at the graft

junction and that many of these genes are sugar responsive, which correlates

with severing of the phloem tissue and the accumulation of starch above the

junction. However, genes associated with cell division and vascular formation

activate on both sides of the graft and, similarly, auxin responsiveness activates

equally on both sides. We propose that the continuous transport of substances,

including auxin, independent of functional vascular connections, promoted

division and differentiation, while the enhanced auxin response and blocked of

sugar transport provided a unique physiological condition to activate genes

specific to graft formation that promote wound healing.

RESULTS

Genes are asymmetrically expressed around the graft

Previous analyses revealed an asymmetry in the genetic requirements for tissue

reunion or graft formation and identified several genes expressed above a cut

that were not expressed below (Asahina et al., 2011; Melnyk et al., 2015). To

investigate whether asymmetry was a common feature of grafting and tissue

reunion, we generated RNA deep sequencing libraries from *Arabidopsis thaliana*

hypocotyl tissues immediately above and immediately below the graft junction 0,

6, 12, 24, 48, 72, 120, 168 and 240 hours after grafting (HAG)(Figure 1A). Prior

to RNA extraction, we separated top and bottom tissues at the graft junction. We

found that the strength required to break apart the graft junction increased

linearly (Figure S1) similarly to previously reported breaking strength dynamics

of grafted *Solanum pennellii* and *Solanum lycopersicum* (Lindsay et al., 1974;

Moore, 1984). When pulling apart grafts to separate top and bottom for sample

preparation, grafts broke cleanly with minimal tissue from one half present in

the other half (Figure S1, Movie S1, S2). We measured the amount of tissue from

tops adherent to bottoms and *vice versa* (Figure S1) and found less than 4%

cross-contamination. In addition to grafting, we also prepared libraries from

ungrafted hypocotyls ("intact" treatment) and cut plants that had not been

reattached ("separated" treatment)(Figure 1A). We herein refer to tissues

harvested above the graft junction or from the shoot side of separated tissue as

"top" and that from below the graft or from the root side of separated tissue as

"bottom" (Figure 1A).

RNAs that were differentially expressed as a consequence of grafting (compared

to intact hypocotyls) equally in tops and bottoms of grafts (symmetrically

expressed) or were more highly expressed in one tissue than the other

(asymmetrically expressed), were identified by performing a pairwise analysis of

the protein-coding transcriptome datasets that were differentially expressed

relative to the intact group. Several thousand RNAs were identified that fit either

pattern of expression including the transcript of the cambial markers *HCA2* that

was induced symmetrically, and *ANT* that was induced asymmetrically (Figure

1B-G, Figure S2). 6 to 48 hours after grafting, the number of graft-differentially

expressed genes that were asymmetrically expressed was roughly 3-fold greater

than those symmetrically expressed indicating that tissues above the cut

changed their expression dynamics relative to below the cut. However, at 72

hours the numbers were nearly equal, and by 120 hours, the number of

symmetrically differentially expressed genes was 3-fold greater than those

asymmetrically expressed (Figure 1B). As a second approach, we performed a

hierarchical clustering analysis that indicated that the grafted top and grafted

bottom became most similar after 72 hours (Figure S3), consistent with the

symmetry analysis (Figure 1B). Thus, graft healing and tissue reunion promoted

a shift from asymmetry to symmetry (Figure 1B).

Figure 1. Grafting rapidly activates genes in both asymmetric and symmetric patterns. (A) Intact, cut and separated, or cut and grafted *Arabidopsis* tissues were harvested approximately 0.5mm above (top), 0.5mm below (bottom) the cut site or for intact plants 1mm segments spanning the region where cuts were made in grafted and separated plants. (B) Pairwise analysis between the grafted top and grafted bottom identified sets of protein-coding genes symmetrically or asymmetrically expressed at the graft junction, with an FDR < 0.05 and a likelihood of symmetric/asymmetric expression of greater than 50%. Asymmetrically expressed genes were further divided into those whose RNAs were higher in the top (orange dotted line) or higher in the bottom (purple dotted line), again with an FDR < 0.05 and a likelihood of asymmetric expression greater than 50%. (C,D) Expression profiles for transcripts of cambium-associated genes that initially activate symmetrically (*HCA*) or asymmetrically (*ANT*) plotted for intact, separated and grafted samples.

(E-G) *HCA2* transcription upregulates above the graft junction (E) and below the
graft junction. (F) *pHCA2::RFP* was grafted to Col-0 roots (E) or Col-0 shoots (F)
to avoid ambiguity of signal origin at the junction. *HCA2* was also upregulated in
separated shoots but not in intact samples or in high levels in separated roots
(G). HAG, hours after grafting. HAS, hours after separation. White triangle
denotes initial fluorescent signal, dashed lines denote the graft junction.
209
Sugar response correlates with asymmetric gene expression

The shift from asymmetry to symmetry could be due to phloem reconnection at

72 hours ((Melnyk et al., 2015), Figure 2A) and the resumption of hormone,

protein and sugar transport. We tested a role for sugar by grafting in the

presence of exogenous sucrose which has previously been reported to affect

grafting success (Marsch-Martinez et al., 2013). Low levels of exogenous sucrose

lowered grafting efficiency (Figure 2A), suggesting that differential sugar

responses at the graft junction might be important for vascular reconnection.

Expression of *ApL3*, a gene whose expression is induced by sugar (Villadsen &

Smith, 2004), was rapidly upregulated in separated tops and grafted tops,

whereas expression of *DIN6*, *GDH1* and *STP1*, genes whose expression is

repressed by sugar (Thum et al., 2004; Villadsen & Smith, 2004; Cordoba et al.,

2015), was rapidly upregulated in separated bottoms and grafted bottoms

(Figure 2B, Figure S4). These observations were consistent with sugar

accumulation in the grafted top and sugar depletion in the grafted bottom. The

expression of these genes returned to levels similar to intact samples by 120

hours and, with the exception of *ApL3*, the grafted samples normalised

expression more rapidly than the separated tissues. Genes associated with

photosynthesis increase expression in separated bottoms 24 hours after cutting,

a common response to starvation (Wang & Ruan, 2013), but likely too late to

affect sugar levels before 24 hours (Figure S4). A transcriptional overlap analysis

with RNAs from known glucose-responsive genes (Table S1) revealed a strong

overlap with genes differentially expressed by grafting. RNAs from known

glucose-induced genes were upregulated in separated tops and grafted tops,

whereas transcripts from known glucose-repressed genes were upregulated in

separated bottoms and grafted bottoms (Figure 2C, Figure S4). This trend was

not observed with genes differentially expressed by mannitol treatment (Figure

S4), suggesting the effect was specific to metabolically active sugars. To further

investigate this effect, we stained grafted, separated and intact plants with Lugol

solution to assay for the presence of starch. Staining above the graft junction

increased 48-72 hours after grafting (Figure 2D). By 120 hours, staining was

equal on both sides of the graft whereas in separated tops staining became

stronger after 72 hours (Figure 2D, Figure S4). We concluded that starch

accumulated above the graft junction, but after 72 hours, this asymmetry

disappeared. To test whether the accumulation of starch and increased sugar

responsiveness could explain the observed transition from asymmetry to

symmetry, we compared our datasets to previously published genes that are

induced by starvation or are induced by sucrose re-addition (Table S1). In early

time points, 20-31% of asymmetrically expressed genes are genes known to

respond to sugars compared to 2-5% of symmetrically expressed genes (Table

1). However, at 72 hours, the overlap between asymmetrically expressed genes

and sugar responsive genes reduced substantially (Table 1).

Figure 2. Asymmetric changes in accumulation of sugar-responsive RNAs
and of starch occur at the graft junction; equalizing sucrose between the
grafted segments reduces and delays phloem reconnection. (A) *pSUC2::GFP*-
expressing *Arabidopsis* shoots were grafted to Col-0 wild type roots and GFP

movement to the roots was monitored over 7 days for phloem connection in the
presence or absence of various concentrations of sucrose. (B) Expression profiles
for transcripts of a sugar-repressed gene (*GDH1*), and a sugar-induced gene
(*ApL3*) were plotted for intact, separated and grafted samples. (C)
Transcriptional overlap between previously published glucose-induced or
glucose repressed genes and our dataset. The number in brackets represents the
number of glucose-responsive genes identified in the previous dataset, and
overlap is presented as a ratio out of 1.0 for differentiation expressed genes
(DEG) up- or down-regulated in our dataset relative to intact samples. Asterisks
represent a significant difference (p < 0.05) between the ratio of up- and down-
regulated genes in a previously published transcriptome dataset compared to
the ratio of all up- and down- regulated genes in our grafting dataset at a certain
time point. (D) Lugol staining of grafted plants at various time points revealed
dark brown staining associated with starch accumulation. HAG, hours after
grafting.
272

273 **Table 1. Sugar response overlaps with asymmetry.** Symmetrically and
274 asymmetrically differentially expressed genes were compared to previously
275 published sugar-responsive genes (Osuna et al., 2007)[1] and the percent overlap
276 calculated (* p<.05). HAG, hours after grafting.
277

HAG	Sugar Induced[1]	Graft Bottom=Top	Overlap	%	Graft Top>Bottom	Overlap	%	Graft Bottom>Top	Overlap	%
6	2243	4988	263	5	6679	1403	21*	4971	99	2
12	2243	3473	165	5	7111	1444	20*	5657	231	4
24	2243	4135	171	4	6873	1414	21*	4942	195	4
48	2243	3689	197	5	6601	1241	19*	4915	218	4
72	2243	10421	1005	10*	2459	228	9*	2019	138	7
120	2243	15012	1063	7	1510	220	15*	941	115	12*
HAG	Sugar Repressed[1]	Graft Bottom=Top	Overlap	%	Graft Top>Bottom	Overlap	%	Graft Bottom>Top	Overlap	%
6	1998	4988	107	2	6679	53	1	4971	1563	31*
12	1998	3473	68	2	7111	112	2	5657	1526	27*
24	1998	4135	88	2	6873	72	1	4942	1525	31*
48	1998	3689	113	3	6601	93	1	4915	1427	29*
72	1998	10421	530	5	2459	111	5	2019	538	27*
120	1998	15012	979	7	1510	84	6	941	210	22*

278
279

Vascular formation and cell division activates on both sides of the graft

Since the grafted bottom samples exhibited a starvation response up to 48 hours

after grafting, we reasoned that pathways associated cell division and cell

differentiation would be delayed or inhibited. We looked at the expression of

markers associated with vascular formation and cell division in the

transcriptome datasets. Cambium, phloem and provascular markers activated

within 6 hours in grafted top samples, but activation was only delayed 6-24

hours in grafted bottom samples depending on the gene (Figure 1B-C, Figure 3,

Figure S5, Figure S6). Expression of phloem markers peaked in both grafted tops

and grafted bottoms at 72 hours (Figure 3, Figure S5, Figure S6), the time when

phloem reconnections form in grafted *Arabidopsis* (Figure 1, (Yin et al., 2012;

Melnyk et al., 2015)). Notably, the early phloem marker *NAC020* activated before

the mid phloem marker *NAC086* which activated before the late phloem marker

NEN4, consistent with the dynamics of phloem transcriptional activation during

primary root development and leaf vascular induction (Figure S5)(Furuta et al.,

2014; Kondo et al., 2016). Certain markers associated with xylem formation,

such as *VND7* and *BFN1*, activated early in the grafted top. Other xylem markers,

such as *IRX3* and *CESA4*, activated late in grafted samples. By 120 hours after

grafting, xylem markers were activated in top and bottom, consistent with when

the first xylem strands differentiate at the graft junction (Melnyk et al., 2015).

Genes associated with cell division were activated by 12 hours in the grafted top

and by 24 hours in the grafted bottom (Figure 3, Figure S5). On the other hand,

control genes whose expression does not typically vary between tissues and

treatments (Czechowski et al., 2005) were not differentially expressed in grafted

tops or bottoms (Figure S6). The RNAseq expression data appeared to correlate

well with transcriptional fluorescent reporters for both activation dynamics and

the localisation of expression (Figure 1D-F, Figure S2).

The similar activation dynamics of vascular differentiation genes in grafted tops

and grafted bottoms prompted us to test whether this phenomenon occurred

with other known developmental processes. We obtained lists of genes whose

expression is associated with various biological processes from previous

publications (Table S1) and tested how many of the genes differentially

expressed in our transcriptomes overlapped with the previously published lists.

Differentially expressed genes in grafted samples and separated tops partially

overlapped with those whose expression is associated with phloem, xylem and

procambium formation (Figure 4, Figure S7). There was a high overlap between

Arabidopsis inflorescence stem healing and grafting, as well as between vascular

induction in leaf disk cultures and grafting (Figure 4). Various genes expressed in

a cell type-specific manner also showed a high transcriptional overlap with

grafting, including phloem, endodermis and protoxylem (Figure 4, Figure S7). In

nearly all cases, the separated top, grafted top and grafted bottom samples

showed similar activation dynamics. The separated bottom samples were

exceptional though since gene expression associated with vascular development

and cell-specific processes was downregulated (Figure 4, Figure S7). We also

compared our datasets to RNAs expressed in longitudinal cross sections of the

Arabidopsis root (Brady et al., 2007). There was little overlap between grafted

bottoms and sections from the root meristemic zone, whereas overlap existed

between grafted tops and the root meristemic zone in early time points, and

between grafting and the root maturation zone (Figure S8). Our analysis also

revealed that two genes expressed in cambium, *WOX4* and *PXY*, were induced by

grafting but the primary root marker *WOX5* and lateral root marker *LBD18* were

not (Figure 3, Figure S5, Figure S6).

Figure 3. Transcriptional dynamics of genes associated with
provasculature, phloem, xylem and cell division. Expression levels were
plotted over time for intact, separated and grafted samples.

Figure 4. Transcriptional overlap between previously published vascular
datasets and the grafting datasets. Genes whose transcripts are associated
with various cell types or biological processes were taken from previously
published datasets (see Table S1) and compared to the transcriptomic datasets
generated here. The number in brackets represents the number of cell type-
specific or process-specific genes identified in the previous dataset, and overlap
is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or
down-regulated in our dataset relative to intact samples. Asterisks represent a
significant difference (p < 0.05) between the ratio of up- and down- regulated
genes in a previously published transcriptome dataset compared to the ratio of
all up- and down- regulated genes in our grafting dataset at a certain time point.

Auxin response is symmetric at the graft

The rapid activation of vascular markers in the grafted bottoms despite the

starvation response promoted us to investigated whether other mobile

substances such as phytohormones could play a role in gene activation. We

compared lists of genes known to respond to cytokinin, ethylene or methyl

jasmonate (Nemhauser et al., 2006) and found no substantial overlap between

these lists and genes differentially expressed by grafting (Figure S9)(Table S1).

Abscisic acid-responsive and brassinosteroid-responsive genes showed overlap

with genes differentially expressed in our datasets, but this overlap was of a

similar magnitude in both separated and grafted datasets suggesting the effect

was not specific to grafting (Figure S9). Auxin responsive transcripts were

exceptional though, as they showed a substantial overlap with RNAs

differentially expressed in our data sets that varied depending on the treatment

(Figure 5A-B, Figure S9). Auxin-induced genes were upregulated in separated

tops, grafted bottoms and grafted tops whereas they were repressed in

separated bottoms (Figure 8). Auxin responsive genes such as *IAA1* and *IAA2*

(Abel et al., 1995) were induced to similar levels in grafted tops and grafted

bottoms at 24 hours. To further investigate whether auxin response was uniform

between grafted tops and grafted bottoms, we grafted the auxin-responsive

fluorescent reporter *p35S:DII-Venus* whose fluorescent protein is degraded in the

presence of auxin (Brunoud et al., 2012). DII-Venus fluoresced in the separated

bottoms but did not fluoresce in grafted bottoms 14 hours after cutting (Figure

5C) indicating separated bottoms had a low level of auxin response but grafted

tops, grafted bottoms and separated tops had a high level of auxin response.

Figure 5. Auxin response is symmetric at the graft junction. (A,D) Expression
profiles for various auxin-responsive genes (*IAA1*, *IAA2*) or auxin transporter
genes (*PIN1* and *ABCB1* genes) were plotted for intact, separated and grafted
samples. (B) Overlap between previously published auxin-induced or auxin-

repressed RNAs and our dataset. The number in brackets represents the number
of auxin-responsive genes identified in the previous dataset, and overlap is
presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or
down-regulated in our dataset relative to intact samples. Asterisks represent a
significant difference (p < 0.05) between the ratio of up- and down- regulated
genes in a previously published transcriptome dataset compared to the ratio of
all up- and down- regulated genes in our grafting dataset at a certain time point.
(C) Grafted and separated plants expressing the auxin responsive *p35S::DII-*
Venus transgene that is degraded in the presence of auxin reveal a reduction of
auxin response in cut bottoms, but not in grafted bottoms. HAG, hours after
grafting.

Tissue fusion imparts a unique physiological response that differs from

tissue separation

We hypothesized that the symmetric auxin response and asymmetric sugar

response at the graft junction could allow a unique transcriptional response

since neither separated plants nor intact plants had similar response dynamics

to sugars and auxin as measured by genome-wide gene response, starch

accumulation and *p35S::DII-Venus* expression (Figure 2, Figure 5). To uncover

protein-coding genes differentially expressed specifically by grafting, we

performed an empirical Bayesian analysis (Hardcastle & Kelly, 2010) where at

each time point we computed likelihoods for each gene for each possible pattern

of differential expression in the intact, grafted top, grafted bottom, separated top

and separated bottom tissues (Figure 6A). These likelihoods were used to define

a similarity score between pairs of genes, which was used to cluster genes with

similar patterns of gene expression across the five tissues (Hardcastle &

Papatheodorou, 2017). Clusters were then identified by the predominant pattern

of gene expression observed within a cluster. This analysis produced 113

clusters containing at least 10 RNAs at a single time point (Table S2; Dataset S1).

Approximately 6000 genes were differentially expressed in at least one tissue

whereas between 1000 and 4000 genes were not differentially expressed

(Figure 6B).

To simplify the analysis, we considered clusters where gene expression was

grouped into patterns consisting of one comparison. At early time points, the

cluster containing genes differentially expressed in grafted tops, grafted bottoms,

separated tops and separated bottoms had high numbers that decreased with

time and could represent an initial general wound response (Figure 6C).

Similarly, clusters containing separated tops and grafted tops or separated

bottoms and grafted bottoms initially had high numbers that decreased with

time. This observation indicated the grafted top was initially transcriptionally

similar to the separated top, whereas the grafted bottom was initially

transcriptionally similar to the separated bottom. After the 48-hour time point,

clusters containing genes only differentially expressed in separated tops or only

differentially expressed in separated bottoms increased in numbers, suggesting

these tissues gained a unique pattern of gene expression. The cluster containing

genes differentially expressed in grafted tops and grafted bottoms increase in

numbers throughout the healing process (Figure 6C). We searched for cluster

categories that contained genes only differentially expressed by grafting and

found very few genes downregulated by grafting or upregulated only in the

grafted top (Figure 7A). Instead, there were clusters that contained several

hundred differentially expressed genes present either in the grafted bottom only,

or present in both grafted bottom and grafted top (Figure 7A). Genes whose

expression changed only in the grafted bottom sample were prevalent early

during grafting and were most common at 48 hours, whereas genes activated in

both top and bottom became prevalent at 48 hours and were most common at

120 hours (Figure 7A). We performed a gene ontology (GO) analysis on the genes

whose expression was grafting-specific and found that genes coding for RNAs

differentially expressed only in the grafted bottom sample were enriched in the

immune response and chitin response biological process categories (Table S3).

Previously published chitin-induced RNAs had a high proportion of overlap with

differentially expressed graft bottom-specific genes (Figure 7E, Figure S10).

Grafting-specific RNAs expressed symmetrically, in both the grafted top and

grafted bottom, were enriched in vascular-related biological processes (Table

S3). Previously published phloem-enriched, endodermal-enriched, vascular-

induction and stem-wounding associated RNAs had a high proportion of overlap

with differentially expressed graft-specific genes (Figure 7C-D, Figure S10). Since

few genes were grafting-specific and grafted tissues were initially

transcriptionally quite similar to separated tissues (Figure 6C, Figure 7A), we

reasoned that tissues separated for short periods (<48 hours) could be grafted

with similar reconnection dynamics as tissues that had been grafted

immediately. To test this hypothesis, plants were cut at day 0, and grafted at 1-5

days after separation. Separated tissues did not speed up vascular reconnection,

and instead, it always took three days from the point of tissue attachment for

vascular connections to form (Figure S10). Furthermore, the shoot lost

competence to graft 2-3 days after separation whereas the root remained

competent to graft up to 5 days after separation (Figure S10). Together, it

appears the grafted shoot and root have a unique physiological response that

differ from separated plants and that tissue attachment is required to activate

graft formation and tissue fusion.

Figure 6. Clustering the transcriptome at each time point, based on
likelihoods of all possible patterns of differential expression. (A) Cartoon
depicting the Bayesian segmentation. (B) Analysis of differential behaviour
produced 113 categories containing at least 10 genes whose expression was
expressed in a specific differential pattern in at least one time point (see Table

S2). One group is comprised of genes whose transcript levels are not
substantially changed in the five tissues (unchanged) whereas the other group is
comprised of the sum of the other 112 groups (genes whose transcript levels
changed after treatment in a least one tissue) over the time points tested. (C)
Major categories in the segmentation revealed RNAs whose levels changed in all
the treatments listed relative to intact samples. Note that a gene can be
represented only in one category for a given time point, that category in which
the transcript level changes best fit the category.
474

Figure 7. A subset of genes is differentially expressed only during graft
formation when compared to 0 hour grafts, or intact or separated tissues.
(A) Certain genes were only differentially expressed in grafted tops, grafted
bottoms or both in grafted tops and grafted bottoms. (B) Expression profiles for
a graft bottom-specific (*ERF6*) or a graft top and bottom differentially regulated
gene (*RTM2*) were plotted for intact, separated and grafted samples. (C)
Grafting-specific genes are also expressed in other processes of tissue fusion,
such as stem healing. Here, 246 previously published RNAs whose differential
expression is associated with stem healing were compared to our dataset to
assess transcriptional overlap with the grafting-specific genes. Asterisks
represent a significant high overlap (p <0.05) of previously published genesets
that are also differentially expressed in the grafted samples at a certain time
point. (D-E) Genes differentially expressed in grafted tops and grafted bottoms
show high overlap to 324 previously published genes whose transcripts are
associated with phloem, whereas genes expressed in grafted bottoms show high
overlap to 286 previously published genes whose encoded RNAs are associated
with chitin treatment. Asterisks represent a significant high overlap (p <0.05) of
previously published genes that are also differentially expressed in the grafted
samples at a certain time point.
495

DISCUSSION

To better understand how plants graft, we analysed in depth an RNA deep sequencing dataset that spatially and temporally distinguished genes activated by cutting and tissue attachment versus genes activated by cutting and continuous tissue separation. We observed that during graft formation, tissues had a very high transcriptional overlap with genes differentially expressed by inflorescence stem healing and by vascular induction in leaves (Figure 4, Figure S7) (Asahina et al., 2011; Kondo et al., 2016) suggesting grafting is closely related to these processes. Graft formation had little transcriptional overlap with lateral root formation (Figure S6)(Brady et al., 2007) and appeared to follow a pathway similar to secondary root growth (Figure S6, Figure S8) since cambium-specific markers *WOX4* and *PXY* were expressed during grafting and secondary growth but are not expressed in the root tip (De Rybel et al., 2016)(Figure S5). Grafted tops initially showed some transcriptional overlap with genes expressed during primary root formation which may be related to the accumulation of substances activating adventitious root formation, a common response in grafts that do not properly form. However, this response was short lasting in grafted tops where adjoining tissues could form vascular connections to re-establish substance transport. Thus, we believe that grafting proceeds via a pathway involving secondary growth with radial meristems activating in the mature cambium to heal the wound. Vascular formation genes including those specifying cambium and phloem were activated early, followed by an activation of cell division genes, suggesting that the start of cellular differentiation preceded activation of cell division. Xylem identity genes showed an early and a late activation peak (Figure 3, Figure S5). There is no visible xylem differentiation at the graft junction during the first peak of expression (Melnyk et al., 2015) and these genes might be suppressed by phloem differentiation genes such as *APL* and *CLE41* that activate early after grafting and are known to suppress protoxylem formation (Bonke et al., 2003; Ito et al., 2006)(Figure S6)(Table S4). Alternatively, the first peak could represent programmed-cell death that does not lead to xylem differentiation. The second expression peak of xylem-expressed genes at 120 hours occurred after the differentiation of functional phloem and coincided with the differentiation of xylem strands at the graft junction (Melnyk et al., 2015).

Previous studies highlighted the importance of callus and pericycle cells during

regeneration (Sugimoto et al., 2010; Iwase et al., 2011), but we see little evidence

that genes expressed in the pericycle or during callus formation have high

transcriptional overlap with genes differentially expressed by grafting (Figure

S4). Expression profiles for all protein-coding genes can be found in table S4.

We observe only a slight delay in phloem, cambium and cell division activation

below the graft junction compared to above it (Figure 3, Figure S5, Figure S6)

whereas some genes, such as *HCA2* and *TMO6,* activated equally in both grafted

top and grafted bottom at 6 hours after grafting (Figure 1D-G, Figure 3). These

data indicate that, at least transcriptionally, the grafted root rapidly responded

to the presence of the grafted shoot. This response was not present in separated

roots, indicating that attachment was key for recognition. Sugars are known

activators of cell division and cell elongation (Wang & Ruan, 2013) and in our

datasets, a large proportion of asymmetrically expressed genes are genes that

are sugar-responsive. However, sugars are transported in the phloem (Lough &

Lucas, 2006) that is severed upon grafting whereas the grafted root exhibited a

sugar starvation response and showed similar sugar-response dynamics as the

separated root. Instead, we believe that not sugar but auxin, or some other

molecular that is transported in the absence of vascular connections, could be

the signal in the grafted bottom that activates *HCA2*, *TMO6* as well as cell

division, phloem and cambium related genes.

Given auxin's role in vascular patterning and formation ((De Rybel et al., 2016), it

is a strong candidate for the activating signal. Auxin response was largely

symmetric, particularly from 12 hours after grafting (Figure 6) consistent with

previous findings that the auxin-inducible *DR5* and *ANAC071* genes are

symmetrically expressed around the graft junction (Yin et al., 2012;

Pitaksaringkarn et al., 2014; Melnyk et al., 2015; Matsuoka et al., 2016). One idea

is that auxin transport was not substantially interrupted by the grafting process,

and instead, where opposing tissues adhered, auxin moved regardless of

vascular connections since auxin is transported from cell to cell through the

apoplast (Leyser, 2011). The genes encoding auxin efflux proteins PIN1 and

ABCB1 were transcriptionally activated above the graft junction (Figure 5D),

similar to putative *Pisum sativum* PIN1 protein accumulating above a cut stem

prior to vascular reconnection (Sauer et al., 2006), and could reflect a role for

these proteins in exporting auxin across the cut. Consistent with these

observations, adding an auxin transport inhibitor to grafted *Arabidopsis* shoots

prevented the expression of grafting-induced genes below the graft junction

(Matsuoka et al., 2016). Since auxin is fundamental for patterning vasculature

(De Rybel et al., 2016), it is plausible that auxin movement across the graft

occurs prior to sugar movement since the flow of auxin could assist with

reconnecting the phloem. Although auxin response was symmetric, our previous

work demonstrated that the auxin response factors *ALF4* and *AXR1* affected

grafting only below the graft junction (Melnyk et al., 2015). Mutating *ALF4* below

the graft junction more strongly reduced auxin response than mutating *ALF4*

above the junction (Melnyk et al., 2015). Thus, proteins such as ALF4 or AXR1

might enhance or promote a rootstock-specific auxin response and could be

particularly important when there is incomplete attachment, cellular damage or

inefficient transport. All higher plants transport auxin from shoot to root, yet not

all plant species can be successfully grafted (Melnyk, 2017c) so the response to

auxin rather than the transport itself may be a determining factor in the ability to

graft. A role for sugars is not completely ruled out though, since the magnitude of

differential expression was often lower in the grafted bottom and exogenous

sugar reduced grafting efficiency (Figure 2A). Pericycle cells require auxin to

divide and the addition of sugars enhances the rate of divisions (Skylar et al.,

2011), suggesting that the presence of sugars in the grafted top enhanced cell

division and differentiation. Another hypothesis that warrants testing is whether

graft healing activates via a mechanical signal provided by the physical presence

of the opposing tissues.

Our analyses identified two groups of genes whose expression changes were

unique to graft formation in our experiments (Figure 7). One group activated

shortly after grafting below the graft junction and was enriched in immune-

responsive and chitin-responsive genes (Table S3, Figure 7, Figure S10). The

breakdown products of cell walls are potent elicitors of defence responses

(Souza et al., 2017), so it is possible that the grafted bottom upregulates
pathways specific to wound damage response. This group was not upregulated
in separated bottoms though, so the unique physiological state of the grafted
root, indicated by the presence of auxin response but the absence of sugar
response, could have promoted their upregulation. The second group activated
both above and below the graft junction and became highly expressed later
during graft formation (Figure 7). This group was enriched in RNAs associated
with vascular development (Table S3, Figure 7) and we suggest that the products
of these genes are involved in the vascular reconnection processes between the
two tissues. Despite many transcriptional similarities between separated tops
and grafted tissues, tissues had to be attached for at least three days for phloem
connections to form, regardless of when cutting occurred (Figure S10). Thus, it
appears that RNAs expressed in the separated top or separated bottom are
insufficient to drive graft formation. Instead attachment and the genes activated
by this recognition process including grafting-specific RNA changes (Figure 7A)
or RNAs expressed in the grafted bottom, grafted top and separated top (Figure
6C) are those that contribute to a response that distinguishes attached from
separated plant tissues. Future work should focus on these RNAs to identify the
pathways required for grafting that could be modified to improve graft
formation, wound healing and vascular regeneration. Likewise, the rapid
transcriptional changes below the graft indicate a recognition system that
promotes tissue regeneration. Identifying the exogenous cue that triggers
recognition and understanding how it is perceived should be priorities, as should
understanding whether this phenomenon applies more broadly to inter-tissue
communication, tissue regeneration or tissue fusion events, such as parasitic
plant infections (Musselman, 1980), epidermal fusions (Becraft et al., 1996; Lolle
et al., 1998) or petal fusions (Zhong & Preston, 2015).
624
METHODS
Plant material and grafting
Arabidopsis thaliana accession Columbia was used throughout. The *p35S::GFP-ER*
(Nelson et al., 2007), *pSUC2::GFP* (Imlau et al., 1999), *pUBQ10::PM-tdTomato*
(Segonzac et al., 2012), *pARR5::GFP* (Yanai et al., 2005), *pANT::H2B-YFP* (Randall

et al., 2015), *pLOG4::n3GFP* (De Rybel et al., 2014), *pCASP1::NLS-GFP* (Roppolo et

al., 2011), *p35S::DII-Venus* (Brunoud et al., 2012) lines have been previously

published. For the construction of *pHCA2::RFP*, a 2.9kb 5' upstream region of the

HCA2 gene was cloned into pDONRp4-p1R donor vector, and recombined with

tagRFPer into a destination vector by the Multisite Gateway system (Siligato et

al., 2016). The following primers were used for *pHCA2* cloning: attB4_HCA2(-

)2958 GGGACAACTTTGTATAGAAAGTTGtcgatacgcgggacagatatac

attB1_HCA2ProEnd ggggACTGCTTTTTTGTACAAACTTGttttgtgttctgtatgtttg.

Arabidopsis thaliana micrografting was performed according to a previously

published protocol (Melnyk, 2017a). Briefly, seven day old *Arabidopsis* seedlings

were grown vertically on Murashige and Skoog (MS) medium + 1% bacto agar

(pH5.7; no sucrose) in short day conditions (8 hours of 80-100 μmol m^{-2} s^{-1} light)

at 20°C. Seedlings were placed on one layer of 2.5x4cm sterile Hybond N

membrane (GE Healthcare) on top of two 8.5cm circles of sterile 3 Chr Whatman

paper (Scientific Laboratory Supplies) moisten with sterile distilled water in a

9cm petri dish. In a laminar flow hood using a dissecting microscope, one

cotyledon was removed and a transvers cut through the hypocotyl was made

with a vascular dissecting knife (Ultra Fine Micro Knife; Fine Science Tools).

Grafts were assembled by aligning the two cut halves and joining them together,

after which, the petri dishes was sealed with parafilm and placed vertically

under short day conditions at 20°C. For grafting on a microscope coverslip to

image the graft junction, a 10cm square Petri dish modified by gluing a

microscope coverslip in place of a section of plastic from the back. On top of the

microscope cover slip was placed a 2.5x4 cm rectangle of Hybond N membrane.

At the edges and base of the Petri dish three 3 x 8cm strips of Whatman paper

were placed. Sterile water moistened both Whatman paper and Hybond N. After

which, roots were placed on the Hybond N membrane and hypocotyls on the

coverslip. Grafting then proceeded as above. Graft junctions were imaged

through the coverslip with a Plan-Apochromat 20X/0.8 objective on a Zeiss LSM-

700 or LSM-780 confocal microscope.

Fluorescent assays and microscopy

To test the effect of sugars on grafting, *Arabidopsis thaliana* Col-0 plants were

grown on 1/2MS+ 1% bacto agar (pH5.7; no sucrose) for seven days in short day

conditions. Wild type roots were grafted to scions expressing *pSUC2::GFP* using

the protocol described above but either water or water containing 0.25%, 0.5%

1% or 2% sucrose was added to the grafted plates. Roots were observed for

fluorescence 2-7 days after grafting with a Zeiss V12 dissecting microscope

equipped with a GFP filter. Roots were scored daily and the same plants were

observed during the 7-day assay as previously described (Melnyk, 2017b).

Fluorescent images were taken on a Zeiss LSM-700 or LSM-780 confocal

microscope with a Zeiss Plan-Apochromat 20X/0.8 dry objective. A 488nm argon

laser (Zeiss 780) or 488nm solid-state laser (Zeiss 700) was used for excitation

of GFP and YFP. A 561nm solid-state laser was used for excitation of the

tdTomato fluorescent protein. A T-PMT detector obtained bright-field

transmitted light. Black and white fluorescent images of graft junctions were

taken on a Zeiss V12 dissecting microscope fitted with a Hamamatsu EM-CCD

camera and RFP and YFP filters. FIJI software (Fiji.sc) was used to process

images. Image contrast and brightness were adjusted for controls and samples

equally. For longitudinal images of the graft junction, z-stack projections are

shown and made with the average intensity function in FIJI from stacks

containing the hypocotyl vascular tissues, mesophyll and epidermis.

RNAseq sample and library preparation

Wild type *Arabidopsis thaliana* accession Col-0 were grafted as above taking care

to switch shoot and root between different plants. All grafting and cutting was

performed in the morning to minimize circadian effects. For the 0 hour time

points, plants were transferred from 1/2MS plates to the grafting plates and

immediately harvested. Only intact plants were harvested at the 0 hour time

since at this point, there would be insufficient time to reasonably expect the

separated or grafted samples to be transcriptionally different (the time between

cutting and freezing is less than two minutes). For all other time points, after

cutting plants were left on grafting plates for the respective amount of time.

Tissues were harvested and care taken to separate grafts by gently pulling plants

apart. Approximately 0.5mm of tissue was taken above or below each cut site

and kept separate. Intact plants had 1mm of tissue taken in a similar location on

the hypocotyl as separated or grafted plants. Grafted, separated or intact tissues

were pooled into groups of approximately 24 tissues (1 plates with 24 plants)

which were immediately placed in 96% ethanol on dry ice. After harvesting,

microcentrifuge tubes were briefly centrifuged and the ethanol removed before

storing at -80°C. Plants were grafted over two months to get sufficient material.

Tissues were ground in the microcentrifuge tube using a microcentrifuge pestle

frozen in liquid nitrogen. RNA was extracted using an RNeasy Kit (Qiagen, UK)

following the manufacturer's instructions including on column DNase digestion.

RNA was eluted from the column with 50ul of sterile water. Quality and quantity

of RNA was checked using an Agilent 2200 TapeStation and High Sensitivity (HS)

RNA screentapes (Agilent, UK). After RNA extraction, two to four biological

replicates were combined (50-100 plants) to get sufficient RNA. 90-100ng of

RNA was used to prepare RNAseq libraries using the TruSeq® Stranded mRNA

LT kit (Illumina, UK) according to the manufacturer's instructions. The final PCR

was for 15 cycles and samples were resuspended in 23ul of distilled water.

Quantity and quality of DNA libraries was checked on the Agilent 2200

TapeStation using D1000 screentapes (Agilent, UK). Each sample had two

libraries prepared from grafted tissues or separated tissues at different times so

that independent biological replicates were made. Samples were diluted to 10nM

and 11-12 barcoded samples randomly mixed to make a total of 7 mixes for 7

flow lanes, one mix per lane. Samples were sequenced on the HiSeq 4000

platform (Illumina, UK) with Paired End 75bp transcriptome sequencing (BGI

Tech Solutions, Shenzhen, China).

Iodine staining

Arabidopsis seedlings were placed in a fixation solution (3.7% formaldehyde,

50% ethanol, 5% acetic acid) for 1 hour at room temperature, then transferred

to 70% ethanol for 10 minutes. Afterwards, plants were transferred to 96%

ethanol and stored at -20°C for up to a week. Samples were rehydrated in 50%

ethanol for 1 hour at room temperature, then transferred to distilled water for

30 minutes. Samples were then transferred to a solution of Lugol solution

(Sigma) and stained for 10 minutes at room temperature. Plants were rinsed

with water, then mounted on microscope slides. Images were taken on a Zeiss

Axioimager.M2 microscope with a PlanApochromat 20x objective and SPOT Flex

camera (Imsol, UK).

Pairwise and Bayseq analyses

The reads acquired through high-throughput sequencing were quality trimmed

with sickle (Joshi & Fass, 2011) to increase the read quality before mapping.

Reads were aligned to protein-coding gene sequences acquired from TAIR10

using Bowtie2. Read assignment was performed using the eXpress tool, which

also provided effective gene lengths for use in normalisation. Library scaling

factors were inferred from the sum of the number of reads assigned to the genes

in the lowest seventy-five percentiles of expressed genes for each library

(Hardcastle et al., 2012).

Analyses of the data were carried out using the R package baySeq (Hardcastle &

Kelly, 2010)and clustering based on the posterior probabilities acquired from

this package. For each timepoint, all possible patterns of differential expression

between the graft types were considered, where a 'pattern' defines similarity and

difference between different experimental conditions. For example,

'{Col_cut_bottomGenes=Col:Col_bottomGenes=ungraftedGenes},{Col_cut_topGenes=Col:Col_topGenes}'

defines a pattern in which gene expression is equivalent in the separated

bottoms, the grafted bottoms and the intact plant, but different to the

equivalently expressed separated top and grafted top. The total number of

possible patterns for five experimental conditions (as in this analysis) is fifty-

two.

For a given timepoint, posterior likelihoods on the likelihood of each pattern of

expression are calculated for every gene with greater than ten reads across all

experimental conditions. The patterns were then modified to include orderings

(denoted by < or >), for example, the pattern described would lead to the

ordered pattern

'{Col_cut_bottomGenes=Col:Col_bottomGenes=ungraftedGenes}>{Col_cut_topGenes=Col:Col_topGenes}'

in which gene expression is equivalent in the separated bottoms, the grafted

bottoms and the intact plant and greater than the equivalently expressed

separated top and grafted top. In total, 541 ordered patterns exist in this data

set. Posterior likelihoods for an ordered pattern were assigned to that of the

unordered pattern for genes in which the (normalised) mean expressions within

the equivalently expressed groups conformed to the ordering, and to zero

otherwise.

Based on the posterior likelihoods for the ordered patterns, a similarity score s_{ij}

was established between two genes i and j as the sum over the products of their

likelihoods of each ordered pattern. A single-link agglomerative clustering of

genes, in which a gene will join a cluster if it has a greater than 50% similarity to

any gene within that cluster was then performed based on these similarity

scores. We label each cluster according to the predominant ordered pattern with

high likelihood amongst the genes that comprise it. The change in size of these

clusters over time is shown for the major clusterings in Figure 6.

We can also find likelihoods on comparisons between pairs of experimental

conditions by summing the likelihoods over combinations of patterns. Figure 1B

shows the number of genes identified at each time point in a pairwise analysis

between the grafted top and grafted bottom samples. The likelihood of

symmetric expression (i.e., expression which is equivalent across the graft

junction) is calculated as the sum of the likelihoods of all patterns in which the

grafted top and grafted bottom samples are equivalent. Conversely, asymmetric

expression is calculated as the sum of the likelihoods of all patterns in which the

grafted top and grafted bottom samples are not equivalent. Additional sets can

be formed by considering the ordering of the grafted top and grafted bottoms

samples. Sets of genes are identified at each time point with an FDR of less than

0.05 and a likelihood of symmetric/asymmetric expression greater than 50%.

Genes in this analysis were only included if they were differentially expressed

relative to intact samples.

Gene overlap analyses of up and down regulated genes

To measure if the ratio of up- and down- regulated genes from a previously published dataset is significantly different to the ratio of up- and down-regulated genes in our grafting dataset we only took into account genes that are differentially expressed at a certain time point. A gene was called differentially expressed at a certain time point if the marginal likelihood, calculated by BaySeq, was greater than 0.9 and if the absolute $\log_2$-foldchange was greater than 1. Hence, we only consider genes that are significantly two-fold up- or down-regulated. This definition of differentially expressed genes was also used to filter the published datasets according to the expression values in our transcriptome dataset. Hence, some genes were filtered out from the original published datasets because they did not show a significant up- or down- regulation during a certain time point in our expression data based on our criteria. The histograms (Figure 2,4, 5, S4, S7, S8, S9) show the relative number of up- and down-regulated genes from the published datasets during a certain time point and a certain condition (separated top, separated bottom, grafted top, grafted bottom) based on the number of genes in the published dataset after filtering. To calculate the significance of the difference of the ratios between the published DEGs and all up- and down- regulated genes, we used a two-sided Fisher's exact test. To correct for multiple testing we used the Benjamini-Yekutieli (BY) correction method. Hence, the asterisks in the barplots highlight that the corrected p-value is below 0.05.

Dealing with probe ids from microarray datasets

Due to the fact that some published datasets only used probe ids instead of gene ids to represent their differentially expressed genes we first had to match these probe ids to their corresponding gene ids. This step was done with the R package biomartr (Drost & Paszkowski, 2017). If one probe id matched more than one gene id we used all the corresponding gene ids and tested afterwards if these genes were actually differentially expressed in our dataset. In some cases, one probe id was represented by more than one gene id. Hence, some gene sets contained slightly more gene ids than published probe ids. In contrast, some

probe ids did not match to currently existing gene ids. Hence, some gene sets

contained slightly fewer gene ids than published probe ids.

Gene overlap analyses of gene sets involved in graft formation

For this analysis we used the previously calculated gene sets from `baySeq` of

differentially expressed genes clustered for each time point into grafted bottom,

grafted top, and grafted bottom and top and calculated their overlap to

previously published data sets. The significance of the number of overlapping

genes between the grafted samples and the published datasets was determined

by a one-sided Fisher's exact test, to prove if the overlap is greater than

expected. The resulting p-values were corrected for multiple testing by using the

Benjamini-Yekutieli method.

This procedure was also applied to generate Table 1 to study the overlaps of

symmetrically and asymmetrically expressed genes in the grafting dataset with

previously published sugar-responsive genes.

GO enrichment analysis

The gene ontology analysis (GO) enrichment analysis on grafting-specific genes

was done with a customized R script using the package GOstats (Falcon &

Gentleman, 2007). Gene ontology annotation was used from the Bioconductor

package org.At.tair.db (Carlson, 2017). The p-values calculated by a

hypergeometric test were corrected for multiple testing with the Bonferroni

correction. A GO category was called enriched if the corrected p-value was below

0.05.

A detailed description, the required data and R scripts to reproduce the

clustering (hierarchical clustering and PCA), the statistical analyses regarding

the overlap studies, and the GO enrichment analysis are available via the GitHub

repository (https://github.com/AlexGa/GraftingScripts).

Breaking force measurements

A micro-extensometer was used for all breaking force graft measurements

according to a previously published protocol (Robinson et al., 2017). Briefly,

grafted Col-0 plants were attached to a moving plate using tough tags (0.94X0.50

inches, white, 679 catalog no. TTSW-1000, DiversifiedBiotech) and a

cyanoacrylate glue. The plates were moved apart and the force was measured

using a force sensor (Futek LSB200 10g load cell, Futek Inc.). Upon breaking the

force dropped. The maximum force measured before breaking was recorded.

Images were captured using a Leica SP5 confocal microscope. During the

experiment single z-place images were captured and made into a movie in

ImageJ.

To measure levels of tissue contamination between grafted top and bottom,

grafted green fluorescent protein(*p35S::GFP*) and red fluorescent protein

(*pUBQ10::PM-tdTomato*) expressing plants were pulled apart manually and

imaged on a Zeiss LSM 700 confocal microscope. Z-stacks comprising the

majority of the hypocotyl were made and regions approximately 0.5mm above

and 0.5mm below the graft junction were made into projections using the

average intensity function on FIJI software, and then the mean intensity

quantified for red and green channels on FIJI. The mean intensity of one colour in

the tissue tested for contamination was divided by the mean intensity of the

same colour in the tissue expressing that transgenes to get a percentage of

contamination. A region away from the cut site was also quantified to get a

percentage of spectral overlap between red and green channels. The percent

spectral overlap was then subtracted from the percentage contamination to get

an overall percentage for how much contamination was present.

ACKNOWLEDGEMENTS

We thank Niko Geldner, Dolf Weijers, Paul Tarr, Yka Helariutta, Ruth Stadler and

The Arabidopsis Information Resource for providing seeds. Funding for this

work was provided by Gatsby Charitable Trust grants GAT3272/C and GAT3273-

PR1, by a Knut and Alice Wallenberg Academy Fellowship KAW2016.0274 (to

C.W.M), and by the Howard Hughes Medical Institute and Gordon and Betty

Moore Foundation grant GBMF3406 (to E.M.M.)

REFERENCES

Abel S, Nguyen MD, Theologis A. (1995). The PS-IAA4/5-like family of early auxin-inducible mRNAs in Arabidopsis thaliana. *J Mol Biol* **251**(4): 533-549.

Aloni R. (1980). Role of auxin and sucrose in the differentiation of sieve and tracheary elements in plant tissue cultures. *Planta* **150**(3): 255-263.

Asahina M, Azuma K, Pitaksaringkarn W, Yamazaki T, Mitsuda N, Ohme-Takagi M, Yamaguchi S, Kamiya Y, Okada K, Nishimura T, et al. (2011). Spatially selective hormonal control of RAP2.6L and ANAC071 transcription factors involved in tissue reunion in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **108**(38): 16128-16132.

Becraft PW, Stinard PS, McCarty DR. (1996). CRINKLY4: A TNFR-like receptor kinase involved in maize epidermal differentiation. *Science* **273**(5280): 1406-1409.

Bonke M, Thitamadee S, Mahonen AP, Hauser MT, Helariutta Y. (2003). APL regulates vascular tissue identity in Arabidopsis. *Nature* **426**(6963): 181-186.

Brady SM, Orlando DA, Lee JY, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN. (2007). A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* **318**(5851): 801-806.

Brunoud G, Wells DM, Oliva M, Larrieu A, Mirabet V, Burrow AH, Beeckman T, Kepinski S, Traas J, Bennett MJ, et al. (2012). A novel sensor to map auxin response and distribution at high spatio-temporal resolution. *Nature* **482**(7383): 103-106.

Carlson M 2017. org.At.tair.db: Genome wide annotation for Arabidopsis.

Chen Z, Zhao J, Hu F, Qin Y, Wang X, Hu G. (2017). Transcriptome changes between compatible and incompatible graft combination of Litchi chinensis by digital gene expression profile. *Scientific reports* **7**(1): 3954.

Cheong YH, Chang HS, Gupta R, Wang X, Zhu T, Luan S. (2002). Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in Arabidopsis. *Plant physiology* **129**(2): 661-677.

Chitwood DH, Guo M, Nogueira FT, Timmermans MC. (2007). Establishing leaf polarity: the role of small RNAs and positional signals in the shoot apex. *Development* **134**(5): 813-823.

Cookson SJ, Clemente Moreno MJ, Hevin C, Nyamba Mendome LZ, Delrot S, Magnin N, Trossat-Magnin C, Ollat N. (2014). Heterografting with nonself rootstocks induces genes involved in stress responses at the graft interface when compared with autografted controls. *Journal of experimental botany* **65**(9): 2473-2481.

Cookson SJ, Clemente Moreno MJ, Hevin C, Nyamba Mendome LZ, Delrot S, Trossat-Magnin C, Ollat N. (2013). Graft union formation in grapevine induces transcriptional changes related to cell wall modification, wounding, hormone signalling, and secondary metabolism. *Journal of experimental botany* **64**(10): 2997-3008.

Cordoba E, Aceves-Zamudio DL, Hernandez-Bernal AF, Ramos-Vega M, Leon P. (2015). Sugar regulation of SUGAR TRANSPORTER PROTEIN 1 (STP1)

expression in Arabidopsis thaliana. *Journal of experimental botany* **66**(1):
147-159.
Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR. (2005).
Genome-wide identification and testing of superior reference genes for
transcript normalization in Arabidopsis. *Plant physiology* **139**(1): 5-17.
De Rybel B, Adibi M, Breda AS, Wendrich JR, Smit ME, Novak O, Yamaguchi
N, Yoshida S, Van Isterdael G, Palovaara J, et al. (2014). Plant
development. Integration of growth and patterning during vascular tissue
formation in Arabidopsis. *Science* **345**(6197): 1255215.
De Rybel B, Mahonen AP, Helariutta Y, Weijers D. (2016). Plant vascular
development: from early specification to differentiation. *Nat Rev Mol Cell*
Biol **17**(1): 30-40.
Drost HG, Paszkowski J. (2017). Biomartr: genomic data retrieval with R.
Bioinformatics **33**(8): 1216-1217.
Falcon S, Gentleman R. (2007). Using GOstats to test gene lists for GO term
association. *Bioinformatics* **23**(2): 257-258.
Furuta KM, Yadav SR, Lehesranta S, Belevich I, Miyashima S, Heo JO, Vaten
A, Lindgren O, De Rybel B, Van Isterdael G, et al. (2014). Plant
development. Arabidopsis NAC45/86 direct sieve element morphogenesis
culminating in enucleation. *Science* **345**(6199): 933-937.
Goldschmidt EE. (2014). Plant grafting: new mechanisms, evolutionary
implications. *Frontiers in plant science* **5**: 727.
Hardcastle TJ, Kelly KA. (2010). baySeq: empirical Bayesian methods for
identifying differential expression in sequence count data. *BMC*
Bioinformatics **11**: 422.
Hardcastle TJ, Kelly KA, Baulcombe DC. (2012). Identifying small interfering
RNA loci from high-throughput sequencing data. *Bioinformatics* **28**(4):
457-463.
Hardcastle TJ, Papatheodorou I. (2017). clusterSeq: methods for identifying
co-expression in high-throughput sequencing data. *bioRxiv* **188581**.
Ikeuchi M, Iwase A, Rymen B, Lambolez A, Kojima M, Takebayashi Y,
Heyman J, Watanabe S, Seo M, De Veylder L, et al. (2017). Wounding
triggers callus formation via dynamic hormonal and transcriptional
changes. *Plant physiology*.
Imlau A, Truernit E, Sauer N. (1999). Cell-to-cell and long-distance trafficking
of the green fluorescent protein in the phloem and symplastic unloading
of the protein into sink tissues. *The Plant cell* **11**(3): 309-322.
Ito Y, Nakanomyo I, Motose H, Iwamoto K, Sawa S, Dohmae N, Fukuda H.
(2006). Dodeca-CLE peptides as suppressors of plant stem cell
differentiation. *Science* **313**(5788): 842-845.
Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M,
Sakakibara H, Sugimoto K, et al. (2011). The AP2/ERF transcription
factor WIND1 controls cell dedifferentiation in Arabidopsis. *Current*
Biology **21**(6): 508-514.
Joshi NA, Fass JN 2011. Sickle: A sliding-window, adaptive, quality-based
trimming tool for FastQ files (Version 1.33).
Kondo Y, Nurani AM, Saito C, Ichihashi Y, Saito M, Yamazaki K, Mitsuda N,
Ohme-Takagi M, Fukuda H. (2016). Vascular Cell Induction Culture
System Using Arabidopsis Leaves (VISUAL) Reveals the Sequential

989   Differentiation of Sieve Element-Like Cells. *The Plant cell* **28**(6): 1250-
990   1262.
991 **Lee J, Kubota C, Tsao SJ, Bie Z, Echevarria PH, Morra L, Oda M. (2010).**
992   Current status of vegetable grafting: Diffusion, grafting techniques,
993   automation. *Scientia Horticulturae* **127**: 93-105.
994 **Leyser O. (2011).** Auxin, self-organisation, and the colonial nature of plants.
995   *Current Biology* **21**(9): R331-337.
996 **Lindsay DW, Yeoman MM, Brown R. (1974).** An analysis of the development of
997   the graft union in Lycopersicon esculentum. *Annals of Botany* **38**: 639-
998   646.
999 **Lolle SJ, Hsu W, Pruitt RE. (1998).** Genetic analysis of organ fusion in
1000   Arabidopsis thaliana. *Genetics* **149**(2): 607-619.
1001 **Lough TJ, Lucas WJ. (2006).** Integrative plant biology: Role of phloem long-
1002   distance macromolecular trafficking. *Annual review of plant biology* **57**:
1003   203-232.
1004 **Marsch-Martinez N, Franken J, Gonzalez-Aguilera KL, de Folter S, Angenent**
1005   **G, Alvarez-Buylla ER. (2013).** An efficient flat-surface collar-free
1006   grafting method for Arabidopsis thaliana seedlings. *Plant methods* **9**(1):
1007   14.
1008 **Matsumoto-Kitano M, Kusumoto T, Tarkowski P, Kinoshita-Tsujimura K,**
1009   **Vaclavikova K, Miyawaki K, Kakimoto T. (2008).** Cytokinins are
1010   central regulators of cambial activity. *Proceedings of the National Academy*
1011   *of Sciences of the United States of America* **105**(50): 20027-20031.
1012 **Matsuoka K, Sugawara E, Aoki R, Takuma K, Terao-Morita M, Satoh S,**
1013   **Asahina M. (2016).** Differential Cellular Control by Cotyledon-Derived
1014   Phytohormones Involved in Graft Reunion of Arabidopsis Hypocotyls.
1015   *Plant & cell physiology* **57**(12): 2620-2631.
1016 **Melnyk CW. (2017a).** Grafting with Arabidopsis thaliana. *Methods in molecular*
1017   *biology* **1497**: 9-18.
1018 **Melnyk CW. (2017b).** Monitoring Vascular Regeneration and Xylem
1019   Connectivity in Arabidopsis thaliana. *Methods in molecular biology* **1544**:
1020   91-102.
1021 **Melnyk CW. (2017c).** Plant grafting: insights into tissue regeneration.
1022   *Regeneration* **4**(1): 3-14.
1023 **Melnyk CW, Schuster C, Leyser O, Meyerowitz EM. (2015).** A developmental
1024   framework for graft formation and vascular reconnection in Arabidopsis
1025   thaliana. *Current Biology* **25**(10): 1306-1318.
1026 **Moore R. (1984).** Graft formation in Solanum pennellii (Solanaceae). *Plant Cell*
1027   *Reports* **3**(5): 172-175.
1028 **Musselman LJ. (1980).** The biology of Striga, Orobanche, and other root-
1029   parasitic weeds. *Annual Review of Phytopathology* **18**: 463-489.
1030 **Nelson BK, Cai X, Nebenfuhr A. (2007).** A multicolored set of in vivo organelle
1031   markers for co-localization studies in Arabidopsis and other plants. *The*
1032   *Plant Journal* **51**(6): 1126-1136.
1033 **Nemhauser JL, Hong F, Chory J. (2006).** Different plant hormones regulate
1034   similar processes through largely nonoverlapping transcriptional
1035   responses. *Cell* **126**(3): 467-475.
1036 **Osuna D, Usadel B, Morcuende R, Gibon Y, Blasing OE, Hohne M, Gunter M,**
1037   **Kamlage B, Trethewey R, Scheible WR, et al. (2007).** Temporal

responses of transcripts, enzyme activities and metabolites after adding
sucrose to carbon-deprived Arabidopsis seedlings. *The Plant journal : for*
cell and molecular biology **49**(3): 463-491.
Pitaksaringkarn W, Ishiguro S, Asahina M, Satoh S. (2014). ARF6 and ARF8
contribute to tissue reunion in incised Arabidopsis inflorescence stems.
Plant Biotechnology **31**(1): 49-53.
Randall RS, Miyashima S, Blomster T, Zhang J, Elo A, Karlberg A, Immanen J,
Nieminen K, Lee JY, Kakimoto T, et al. (2015). AINTEGUMENTA and
the D-type cyclin CYCD3;1 regulate root secondary growth and respond to
cytokinins. *Biology open*.
Robinson S, Huflejt M, Barbier de Reuille P, Braybrook SA, Schorderet M,
Reinhardt D, Kuhlemeier C. (2017). An automated confocal micro-
extensometer enables in vivo quantification of mechanical properties
with cellular resolution. *bioRxiv* **183533**.
Roppolo D, De Rybel B, Tendon VD, Pfister A, Alassimone J, Vermeer JE,
Yamazaki M, Stierhof YD, Beeckman T, Geldner N. (2011). A novel
protein family mediates Casparian strip formation in the endodermis.
Nature **473**(7347): 380-383.
Sauer M, Balla J, Luschnig C, Wisniewska J, Reinohl V, Friml J, Benkova E.
(2006). Canalization of auxin flow by Aux/IAA-ARF-dependent feedback
regulation of PIN polarity. *Genes & development* **20**(20): 2902-2911.
Segonzac C, Nimchuk ZL, Beck M, Tarr PT, Robatzek S, Meyerowitz EM,
Zipfel C. (2012). The shoot apical meristem regulatory peptide CLV3
does not activate innate immunity. *The Plant cell* **24**(8): 3186-3192.
Siligato R, Wang X, Yadav SR, Lehesranta S, Ma G, Ursache R, Sevilem I,
Zhang J, Gorte M, Prasad K, et al. (2016). MultiSite Gateway-Compatible
Cell Type-Specific Gene-Inducible System for Plants. *Plant physiology*
170(2): 627-641.
Skylar A, Sung F, Hong F, Chory J, Wu X. (2011). Metabolic sugar signal
promotes Arabidopsis meristematic proliferation via G2. *Dev Biol* **351**(1):
82-89.
Souza CA, Li S, Lin AZ, Boutrot F, Grossmann G, Zipfel C, Somerville SC.
(2017). Cellulose-Derived Oligomers Act as Damage-Associated
Molecular Patterns and Trigger Defense-Like Responses. *Plant physiology*
173(4): 2383-2398.
Sugimoto K, Jiao Y, Meyerowitz EM. (2010). Arabidopsis regeneration from
multiple tissues occurs via a root development pathway. *Developmental*
cell **18**(3): 463-471.
Thum KE, Shin MJ, Palenchar PM, Kouranov A, Coruzzi GM. (2004). Genome-
wide investigation of light and carbon signaling interactions in
Arabidopsis. *Genome biology* **5**(2): R10.
Villadsen D, Smith SM. (2004). Identification of more than 200 glucose-
responsive Arabidopsis genes none of which responds to 3-O-
methylglucose or 6-deoxyglucose. *Plant molecular biology* **55**(4): 467-
477.
Wang L, Ruan YL. (2013). Regulation of cell division and expansion by sugar
and auxin signaling. *Frontiers in plant science* **4**: 163.
Wetmore RH, Rier JP. (1963). Experimental induction of vascular tissues in
callus of angiosperms. *American Journal of Botany* **50**(5): 418-430.

Yanai O, Shani E, Dolezal K, Tarkowski P, Sablowski R, Sandberg G, Samach
A, Ori N. (2005). Arabidopsis KNOXI proteins activate cytokinin
biosynthesis. *Current Biology* **15**(17): 1566-1571.
Yin H, Yan B, Sun J, Jia P, Zhang Z, Yan X, Chai J, Ren Z, Zheng G, Liu H.
(2012). Graft-union development: a delicate process that involves cell-
cell communication between scion and stock for local auxin accumulation.
Journal of experimental botany **63**(11): 4219-4232.
Zheng BS, Chu HL, Jin SH, Huang YJ, Wang ZJ, Chen M, Huang JQ. (2010).
cDNA-AFLP analysis of gene expression in hickory (Carya cathayensis)
during graft process. *Tree physiology* **30**(2): 297-303.
Zhong J, Preston JC. (2015). Bridging the gaps: evolution and development of
perianth fusion. *New Phytologist* **208**(2): 330-335.

1099
1100
1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

SUPPLEMENTARY MATERIAL

Movie S1. Breaking weight measurements at the graft junction, related to
figure 1. A micro-extensometer pulled apart an Arabidopsis plant 9 days after
grafting while force was measured using a force sensor and images captured
using a Zeiss Sp5 confocal microscope. Here, the transmitted light field is shown.
Shoot is on top and root towards the bottom.
1131

Movie S2. Breaking weight measurements at the graft junction, related to
figure 1. A micro-extensometer pulled apart an *Arabidopsis* plant 9 days after
grafting while force was measured using a force sensor and images captured
using a Zeiss Sp5 confocal microscope. The same experiment is shown in Movie
S1, but here, the GFP and RFP channels are shown. Shoot is on top (*p35S::GFP*
expressing) and root towards the bottom (*pUBQ10::PM-tdTomato* expressing).
1138

1139 **Table S1.** Details of previously published datasets used to compare to the
1140 grafting datasets.
1141

1142 **Table S2.** Numbers and categories associated with the Bayseq analysis.
1143 Categories are defined as grafted (Col:Col), separated (Col_cut) or intact
1144 (ungrafted).
1145

1146 **Table S3.** GO analysis for biological process (BP). Shown are the top 20 BP GO
1147 terms for the grafting-specific genes. Time point selected are those when there
1148 are the most genes in grafted bottom samples (48hrs) or grafted top + bottom
1149 samples (120 hrs).
1150

1151 **Table S4.** Normalised reads for all the protein-coding genes in the datasets. By
1152 entering the ATG number of interest, a plot is made which shows a differential
1153 gene expression profile for the gene of interest.
1154

1155 **Dataset S1.** Details of the Bayesian segmentation analysis providing files
1156 containing the ATGs for every cluster for every time point.
1157

1158

Figure S1. Experimental approach, related to figure 1. (A) The breaking
weight required to pull apart grafted plants measured by a micro-extensometer.
(B-C) Contamination of top in bottom or bottom in top was less than 4% as
measured by grafting green fluorescent protein-expressing plants (*p35S::GFP*) to
tomato fluorescent protein-expressing plants (*pUBQ10::PM-tdTomato*) and
measuring amounts of fluorescence at the different wavelengths of emission in
the top segments relative to the bottom. Images show different plants prior to
and after pulling with the percent contamination indicated.
1167

1168

Figure S2. Comparison between RNAseq transcriptome expression profiles and transcriptional fluorescent reporters, related to figure 1. RNAseq expression profiles for various genes upregulated during graft formation were plotted for intact, separated and grafted samples (left panels). Transcriptional reporter-expressing plants were cut and separated, cut and grafted, or left intact. After cutting, plants were imaged and z-projections made at various time points (right panels). For *pCASP1::GFP*, we did not observe a signal in intact plants grafted on glass slides (see Materials and Methods), but observed a signal with 3/5 plants 7 days after grafting on Whatman-nylon membrane, the same condition used for transcriptome library preparation. HAG, hours after grafting. HAS, hours after separation. Dashed lines denote the graft junction.

1180

1181

1182

Figure S3. Hierarchical cluster analysis and principle component analysis
(PCA) for the transcriptome libraries, related to figure 1. (A) Hierarchical
clustering of samples based on log10 transformed TPM values. Similarity
between samples was measured by 1 – spearman correlation coefficient. (B)
PCA of expression data shows clustering of similar samples. Especially the
grafted top and grafted bottom samples are very similar from 120 hours
onwards.

1190

Figure S4. Asymmetry is a feature observed with sugar response and starch accumulation, related to figure 2. (A) Expression profiles for RNAs coded by sugar-repressed genes (*DIN6*, *STP1*) or photosynthetic gene (*RBSC3B*) were plotted for intact, separated and grafted samples. (B) Transcriptional overlap between previously published mannitol-induced, mannitol-repressed, sugar-induced or sugar-repressed genes and our datasets. The numbers in brackets represents the number of glucose or mannitol-responsive genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentially expressed genes (DEG) up or down regulated in our dataset relative to intact samples. Asterisks represent a significant difference (p < 0.05) between the ratio of up- and down- regulated genes in a previously published transcriptome dataset compared to the ratio of all up- and down- regulated genes in our grafting dataset at a certain time point. (C) Lugol staining of grafted plants at various time points reveals dark brown staining associated with starch accumulation. Upper grafted panels are the same as those present in Figure 2 and shown here to compare with controls. HAG, hours after grafting.

1207

1208 **Figure S5. Transcriptional dynamics of genes associated with cambium,**
1209 **phloem, xylem and cell division, related to figure 3.** Expression profiles in
1210 grafted tops or grafted bottoms for various genes of interest were plotted,
1211 normalised to intact samples and plotted on a $\log_2$ scale.

1212

1213

1214

Figure S6. Accumulation of RNA from genes associated with provasculature and phloem, and of RNAs from WOX transcription factors, related to figure 3. (A) RNA accumulation profiles for various genes of interest were plotted over time as measured in grafted tops or grafted bottoms, normalised to intact samples and plotted on a log$_2$ scale. The grafting datasets could also be used to investigate the transcriptional dynamics of related genes, such as the sequential activation of WOX transcription factors at the graft junction. (B) Expression levels of a primary root specific transcript (of the *WOX5* gene) or a lateral root specific transcript (from the *LBD18* gene) were plotted over time for intact, separated and grafted samples.

1225

1226

Figure S7. Transcriptional overlap between previously published datasets
and the grafting datasets, related to Figure 4. Genes whose transcripts are
associated with various cell types or biological processes were taken from
previously published datasets (see Table S1) and compared to the datasets
generated here. The number in brackets represents the number of cell type-
specific or process-specific genes identified in the previous dataset, and overlap
is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or
down-regulated in our dataset relative to intact samples. Asterisks represent a
significant difference (p < 0.05) between the ratio of up- and down- regulated
genes in a previously published transcriptome dataset compared to the ratio of
all up- and down- regulated genes in our grafting dataset at a certain time point.

1238

Figure S8. Transcriptional overlap between previously published root transcriptomic datasets and the grafting RNA-seq dataset, related to Figure 4. Genes whose transcripts are associated with various root regions were taken from previously published datasets (see Table S1) and compared to the datasets generated here. Root layer is indicated in the bottom right of each histogram with reference to the cartoon. The number in brackets represents the number of root layer-specific genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or down-regulated in our dataset relative to intact samples. Asterisks represent a significant difference (p < 0.05) between the ratio of up- and down- regulated genes in a previously published transcriptome dataset compared to the ratio of all up- and down- regulated genes in our grafting dataset at a certain time point.

1251

1252

1253

Figure S9. Transcriptional overlap between previously published hormone responsive RNA datasets and the grafting RNA datasets, related to Figure 5.
Genes whose differential expression is associated with various hormone responses were taken from previously published datasets (see Table S1) and compared to the genes represented in the RNA-seq datasets generated. The number in brackets represents the number of cell type-specific or process-specific genes identified in the previous dataset, and overlap is presented as a ratio out of 1.0 for differentiation expressed genes (DEG) up- or down-regulated in our dataset relative to intact samples. Asterisks represent a significant difference ($p < 0.05$) between the ratio of up- and down- regulated genes in a previously published transcriptome dataset compared to the ratio of all up- and down- regulated genes in our grafting dataset at a certain time point.

1266

1267

1268

Figure S10. A subset of genes is differentially expressed only during graft formation, related to Figure 7. (A) *pSUC2::GFP* expressing *Arabidopsis* shoots were grafted to Col-0 wild type roots 0-5 days after transferring to grafting plates and kept intact until grafting (intact treatment). GFP movement to the roots for phloem connection assays was monitored over 7 days. (B) Plants were cut at day 0 and grafted at days 0-5 after cutting. (C) Shoots were cut at day 0 and grafted at days 0-5 after cutting. Roots were kept intact until immediately before grafting. (D) Roots were cut at day 0 and grafted at days 0-5 after cutting. Shoots were kept intact until immediately before grafting. (E) Genes differentially expressed in grafted tops and grafted bottoms show overlap with previously published genes whose transcripts are associated with the endodermis, vascular induction and grafting (see Table S1 for treatment information). Asterisks represent significant high overlap (p <0.05) of previously published gene sets that are also differentially expressed in the grafted samples at a certain time point.

1284