Molecular function limits divergent protein evolution on planetary timescales

Mariam M. Konaté[a,b*], Germán Plata[a*], Jimin Park[a,c], Harris H. Wang[a,c], and Dennis Vitkup[a,d†]

[a] Department of Systems Biology, Columbia University, 1130 Saint Nicholas Ave., New York, NY, 10032,

USA. [b] Division of Cancer Treatment and Diagnosis, National Cancer Institute, 31 Center Dr., Bethesda,

MD, 20892, USA. [c] Department of Pathology and Cell Biology, Columbia University, 3960 Broadway, New

York, NY, 10032, USA. [d] Department of Biomedical Informatics, Columbia University, 1130 Saint Nicholas

Ave., New York, NY, 10032, USA.

[*]These authors contributed equally to this work.

[†]Correspondence to Dennis Vitkup:

1130 Saint Nicholas Ave. RM 803, New York, NY 10032

Tel: 212 851 5151

E-mail: dv2121@columbia.edu

Abstract

Sequence analyses demonstrate that even after billions of years of independent evolution, protein

orthologs are still diverging from each other. Although the sequence and functional evolution of proteins

have been studied in detail, it is currently unknown whether the requirement to continuously maintain

the same molecular function imposes an effective limit on protein divergence. Here we investigate this

fundamental question using a combined computational and experimental analysis of orthologous

enzymes with the same molecular function. Interestingly, our results demonstrate that the mutual

divergence rates of orthologous enzymes substantially decrease after ~1-2 billion years of independent

evolution. As a result of this slowdown, which is observed well above the levels of detectible sequence

homology, the sequence and structural similarities between ancient orthologs have not substantially

decreased during the last billion years. We find that conservation of molecular function constrains the

maximum divergence of all protein regions, including those distant from the active site residues. By

experimentally characterizing the growth rates of all amino acid substitutions in the *E. coli* FolA protein,

we delineate two stages of long-term sequence divergence with different contributions from sites with

small and large fitness effects. Overall, our analysis suggests that the vast majority of ancient enzyme

orthologs are unlikely to diverge, at least on planetary timescales, beyond ~25% sequence identity.

Introduction

As proteins evolve from a common ancestor, their sequences and structures diverge from each other[1, 2]. Multiple previous studies have investigated the relationship between the conservation of protein molecular function and sequence identity[3-5]. Specifically, characterizing the probability that two sequences share the same molecular function given their sequence similarity[4] has been important for understanding the emergence of new functions within protein families[6, 7], and for the functional annotation of protein sequences[3, 8]. In this work, we focused on a different and currently unaddressed question. Namely, how far can two sequences typically diverge in natural populations while continuously maintaining the same molecular function, for example the ability to catalyze the same enzymatic reaction. We note that the requirement for continuous conservation of molecular function is crucial in this context, as multiple examples of convergent evolution and protein engineering demonstrate that catalysis of the same chemical reactions can be accomplished by proteins with unrelated sequences and different folds[9-11].

It was previously demonstrated that proteins with the same fold frequently diverge to very low (~10%) levels of sequence identity[12]. Because only a relatively small fraction of protein residues (~3-5%) are usually directly involved in catalysis[13], it is plausible that divergent evolution, even when constrained by the same molecular function, will eventually lead to orthologs with essentially random levels of sequence identity. However, recent analyses have shown that even sites located far from catalytic residues may constrain protein evolution, as substitutions at these sites can have substantial effects on enzymatic activity[14]. In this study, we explore the long-term divergence limit of protein orthologs using several models of molecular evolution. Importantly, a divergence limit does not imply that the rate of amino acid substitutions slows down in evolution, but suggests that subsequent substitutions do not substantially further decrease the similarity between orthologous sequences and structures. We also investigate the correlation between experimentally measured fitness effects of amino acid substitutions and the contribution of corresponding protein sites to long-term sequence divergence.

Results

To study the evolution of proteins with the same molecular function, we used orthologous enzymes annotated in UniProt[15] with the same enzymatic activity (see Methods). We focused on enzymes because their molecular function is usually well defined: the Enzyme Commission (EC) classifies enzymatic functions using a hierarchical 4-digit code[16], such that two enzymes that share all four EC digits catalyze the same biochemical reaction. In the present analysis, we used protein sequences representing 64 EC numbers from 22 diverse model organisms across the three domains of life (Table S1).

The considered activities include members of all 6 major enzyme classes: oxidoreductases, transferases,

hydrolases, lyases, isomerases and ligases.

To investigate whether the conservation of enzymatic function limits the divergence between

orthologous sequences, we first considered two simple models of long-term protein evolution, one

without a limit of sequence divergence and the other with an explicit divergence limit. In the first model,

all protein sites have equal and independent substitution rates; see Equation 1, where y represents global

sequence identity, t represents divergence time, and R_0 represents the average substitution rate. In the

second model, in addition to sites with equal and independent substitution rates, there is a minimum

fraction of identical sites at long divergence times, represented by Y_0 in Equation 2.

$$y = 100 * e^{-R_0 * t} \qquad (1)$$

$$y = Y_0 + (100 - Y_0) * e^{-R_0 * t} \qquad (2)$$

We applied the two models to fit sequence divergence for each of the considered enzymatic

functions. Species' divergence times were used to estimate the times of divergence between

corresponding orthologous proteins. For each enzymatic activity, phylogenetic trees based on the

orthologous protein sequences were constructed and compared to species' trees; proteins that showed

clear differences in the tree topologies, suggesting horizontal gene transfer, were excluded from the

analysis (see Methods).

The best model fits for four representative metabolic activities are shown in Figure 1 (black for

the first model and red for the second); see Figure S1 for the other metabolic activities. In 62 of the 64

cases, the second model fits the divergence data significantly better than the first model (F-test P-value

<0.05, Table S2). Moreover, in 61 of the 64 cases, the maximum likelihood value of the parameter Y_0 is

significantly higher than the average sequence identity of the optimal global alignment between random

protein sequences (13.5%, shown in Figure 1 and Figure S1 by dashed black lines) (Wald test P-value

<0.05). The distribution of the optimized parameter Y_0 suggests a long-term sequence identity higher than

25% between considered orthologs (Figure S2a), supporting the idea that conservation of enzymatic

function significantly limits protein divergence.

Figure 1. Sequence divergence of enzyme orthologs as a function of time. The global pairwise sequence identities between pairs of orthologous enzymes are shown as a function of divergence times between the corresponding species. Three models of amino acid substitution were used to fit the data. Model 1 (black line) assumes independent and equal substitution rates across all protein sites. Model 2 (red line) assumes that a given fraction of protein sites remains identical at long divergence times. Model 3 (blue line) assumes a gamma distribution of substitution rates across sites. Best fits of the models are shown for 4 representative EC numbers: **a.** 1.5.1.3, **b.** 2.7.4.3, **c.** 4.2.1.2, **d.** 6.3.4.2. The dashed black lines represent the average sequence identity for the alignment of unrelated protein sequences. The data and corresponding model fits for the other EC numbers considered in the analysis are given in Figure S1 and Table S2.

The two aforementioned models simplify the process of sequence divergence by considering similar substitution rates across protein sites. A more realistic and commonly used model of protein evolution assumes a gamma distribution[17-19] of substitution rates across sites; see Equation 3, where α represents the shape parameter of the gamma distribution. The best fits of such a variable-rate model (blue in Figure 1 and Figure S1) show that the rates of mutual protein sequence divergence between orthologous enzymes have decreased by more than 10 times during ~4 billion years of evolution (see

Methods and Table S3). Across enzymes, the sharpest decrease in divergence rates occurred within 1-2

billion years of divergence from a common ancestor (Figure 1 and Figure S1). Notably, the observed

decrease in divergence rates is substantially higher (by >5 fold) than the one expected simply due to

progressively fewer non-mutated sites in models with equal and independent substitution rates. The

slowdown in divergence does not imply that mutation rates have decreased in molecular evolution.

Instead, a substantial level of conservation is likely necessary for the continuous preservation of molecular

function, leading to amino acid substitutions that do not further decrease sequence identity[20]. The third

model does not explicitly consider a long-term divergence limit. Nevertheless, the projected sequence

identity also shows that the vast majority of orthologous enzymes will remain above 20-25% sequence

identity on the timescales in which Earth environments will remain hospitable to life (estimated at 1-2.8

billion years from the present[21]) (Figure S2b).

$$y = 100 * \left(\frac{R_0 * t}{\alpha} + 1 \right)^{-\alpha}$$ (3)

Interestingly, following the previously introduced metaphor of the expanding protein universe[2],

we can use the third model to express the divergence rate between orthologs as a function of protein

distance ($D = 1 - y$, where y is sequence identity), see Equation 4. This equation, similarly to Hubble's law

of universe expansion[22], describes how the divergence rate changes with increasing distance between

protein orthologs.

$$\frac{\partial D}{\partial t} = 100 * R_0 * (1 - D)^{(\alpha + 1/\alpha)}$$ (4)

We note that the observed divergence limit is not due to an inability to detect remote protein

homologs, as it occurs at relatively high sequence identities (Figure 1 and Figure S1) for which

corresponding orthologs can be easily identified by computational sequence comparison methods.

Furthermore, the results remained similar when we restricted the analysis to orthologous enzyme pairs

with experimentally validated molecular functions (Figure S3), using publications referenced in the

BRENDA database[23].

Next, to understand how different protein sites contribute to long-term sequence divergence

constraints, we experimentally measured the fitness effects of all possible single amino acid substitutions

in a representative enzyme, the *Escherichia coli* dihydrofolate reductase (FolA, EC 1.5.1.3). We chose FolA

for the experiments due to its small size (159 amino acids) and essential role in the *E. coli* metabolism[24].

Also, the long-term protein sequence identity between FolA orthologs (~32%, see Figure 1a) is similar to

other analyzed enzymes (Figure S2a). Following a recently described strategy[25], we used the Multiplex

Automated Genome Engineering (MAGE) approach[26], to introduce every possible amino acid

substitution at each FolA site in *E. coli*. We then measured fitness effects (relative growth rates) of each

variant by performing growth competition experiments, followed by next generation sequencing (see

Methods, Dataset S1).

Figure 2. Sequence divergence of protein sites with different mutant fitness effects in *E. coli*. a. Sequence
identity between pairs of FolA orthologs as a function of divergence time. Results are shown for subsets
of protein sites (represented by different colors) with different average fitness effects of mutations in *E.
coli*. The average sequence identity values were calculated using ~300 bacterial FolA orthologs in the COG
database[27]; divergence times were estimated using bacterial 16S rRNA sequences (see Methods). **b.**
Similar to panel a, but for the sequence divergence of *E. coli* translation initiation factor InfA orthologs
(~500 sequences). Error bars represent the standard deviation of sequence identity in each bin.

Using the growth measurements in *E. coli*, we investigated the long-term sequence divergence of

protein sites with different contribution to fitness. To that end, we sorted FolA protein sites into several

groups according to the experimentally measured average fitness effects. We then used a large multiple

sequence alignment of FolA orthologs (~300 sequences) from the COG database[27] (see Methods) to

estimate sequence identity as a function of corresponding species' divergence times for sites with

different average fitness effects (Figure 2a, different colors). Similar to the global sequence identity, after

~1.5 billion years of evolution, we observed a significant decrease in mutual divergence rates for sites in

the least deleterious fitness bin (Figure 2a, blue). In contrast, sites with the most deleterious mutations in

E. coli (Figure 2a, black) display approximately constant divergence rate throughout evolutionary history.

To understand the generality of the FolA results, we used a similarly obtained fitness dataset[25] for the

E. coli translation initiation factor InfA (Figure 2b). Using the InfA data we observed, on average, lower

mutant growth rates and lower rates of sequence divergence than for FolA. Nevertheless, the observed

long-term divergence patterns were qualitatively similar (Figure 2b). For sites in the least deleterious

fitness bin (Figure 2b, blue) we observed a substantial decrease in divergence after ~2 billon years of

evolution, while sites with most deleterious mutations in *E. coli* (Figure 2b, pink) display an approximately

constant but slow divergence rate.

The results described above suggest a correlation between substitution fitness effects at a protein

site and the site's long-term sequence identity. To further quantify this relationship, we calculated the

probability that each site is occupied by the same amino acid in pairs of orthologs at large phylogenetic

distances, when mutual divergence rates have significantly slowed down (~2 billion years for FolA and

~2.5 billion years for InfA). To this end, we used the TARGETING package[28] to identify phylogenetically

independent pairs of species with FolA and InfA orthologs (Figure 3a), and calculated the fraction of

independent pairs where a site was identical (Figure 3b). Notably, the probability that a protein site is

identical, and thus contributes to the divergence limit, first increases linearly with increasing fitness

effects at the site, and then begins to saturate for sites with high (>30%) fitness effects (Figure 3b).

192
193

Figure 3. Probability of protein sites conservation in phylogenetically independent lineages. a.
Illustration of species pairs representing phylogenetically independent lineages. In the figure, A-B and D-
H are pairs of species that diverged within the specified time window (illustrated by the shaded region),
and do not share more recent edges in the phylogenetic tree. **b.** The probability that protein sites in FolA
(upper panel) or InfA (lower panel) are occupied by identical amino acids as a function of the average
mutant fitness at the sites in *E. coli*. The probability represents the fraction of phylogenetically
independent pairs of orthologs in which sites are identical at long divergence distances (2±0.25 billion
years for FolA, and 2.5±0.25 billion years for InfA). Error bars represent the S.E.M. across sites with a given
fitness effect.

Given that some protein sites are more likely to be conserved between ancient pairs of orthologs,

it is interesting to understand to what extent the same sites contribute to the divergence limit in different

lineages. Towards that end, we calculated the distribution quantifying how frequently protein sites are

occupied by identical amino acids in phylogenetically independent linages (Figure 4). The results, for both

FolA and InfA orthologs, demonstrate that only a relatively small fraction of protein sites (10-20%) are

identical in a majority (>90%) of independent lineages. Therefore, the observed divergence limit is not

due to a uniform set of conserved protein sites between all pairs of orthologs. Instead, heterogeneous

sets of sites contribute to the limit in independent lineages. Analysis of the sites that contribute to

sequence identity between orthologs at two consecutive divergence times (black and gray in Figure 4),

revealed only minor differences in the corresponding probability distributions. Therefore, the observed

slow-down of sequence divergence after 1.5-2 billion years of evolution is accompanied by the

convergence of the probability distribution characterizing the relative contribution of different protein

sites to the limit.

218
219

Figure 4. Contribution of different protein sites to the observed divergence limit. a. Distribution of FolA
protein sites according to the probabilities of being identical in phylogenetically independent lineages.
Probabilities were calculated as the proportion of phylogenetically independent pairs of FolA orthologs
with identical amino acids. Results are shown for divergence times of 2 (black) and 2.5±0.25 (grey) billion
years between pairs of orthologs. **b.** Similar to panel a. but for InfA orthologs. Results are shown for
divergence times of 2.5 (black) and 3±0.25 (grey) billion years. Error bars represent the standard error of
the calculated fractions across sites in each probability bin.

Conservation of molecular function should also significantly constrain protein structural

evolution[8, 29]. To investigate the long-term structural divergence of proteins with the same molecular

function, we structurally aligned orthologous enzymes using the TM-align algorithm[30]. In this analysis,

we used more than 1000 orthologous pairs of enzymes (sharing all 4 EC digits) with known 3D structures

(see Methods). The average root mean square deviation (RMSD) between C-alpha atoms of the

orthologous enzymes increases (Spearman's r=0.44, P-value<1e-20) with divergence time between the

corresponding species (Figure 4a). Nevertheless, the C-alpha RMSD rarely diverges beyond 3 Å even at

long evolutionary distances. Consistent with sequence evolution (Figure 1), we observed a substantial

decrease in the rate of structural divergence after ~1.5 billion years of evolution.

238
239

Figure 5. Long-term structural evolution of proteins with the same molecular function. a. The pairwise
C-alpha root mean square deviation (RMSD) as a function of the divergence time between pairs of
orthologs annotated with the same EC number and experimentally determined structures (gray dots).
RMSD values were calculated based on structural alignments using the TM-align algorithm[30]. Boxes
indicate the median and 25-75 percentiles at the corresponding divergence times, the vertical lines
indicate the 5-95 percentile range, and the red line shows a moving average of the data. **b.** Divergence of
sequence identity at different distances to enzymes' active sites. In this analysis we considered the same
species and enzymatic activities used to explore the global sequence divergence (Figure 1 and Figure S1);
the average sequence identities at each distance shell were calculated across all pairs of orthologs
annotated with the same EC number (see Methods). Error bars show the S.E.M. across ortholog pairs.
250

Only a small fraction of all protein residues usually forms an enzyme active site. Therefore, it is

interesting to investigate how the sequence divergence limit depends on the spatial proximity to active

site residues. While it was recently demonstrated that evolutionary rates correlate with the sites' spatial

distance to catalytic residues[31], it is not clear whether conservation of molecular function would limit

the maximum possible divergence of corresponding protein sites. To address this question, we considered

catalytic site annotations available from the Protein Data Bank[32], UniProt-KB[15] and the Catalytic Site

Atlas[33] and quantified the average divergence of sequence identity at various distances from catalytic

residues (see Methods, Figure 5b). For this analysis, we considered the same set of enzymatic activities

used to study global sequence divergence (Figure 1 and Figure S1). Although, as expected, residues close

to the active site are most highly conserved[31, 34], even residues distant from the active site display an

effective divergence limit at long evolutionary distances. This result suggests that the spatial constraints

required to conserve molecular function usually propagate throughout the entire protein structure.

Discussion

The presented results demonstrate that, in contrast to proteins with the same fold[12], the

requirement to maintain the same molecular function significantly constrains the long-term divergence

of protein orthologs. Moreover, our analysis delineates two different stages of divergence for ancient

orthologs with the same function. During the first stage, substitutions at sites with mild fitness effects

lead to a relatively fast initial divergence. However, after 1-2 billion years of independent evolution,

divergence at these sites effectively saturates. Notably, the long-term sequence identity between these

sites at saturation (30-40%) is still substantially higher than the expected identity between two random

protein sequences (~10%). This suggests that, even at sites with small fitness effects, not all combinations

of amino acids are compatible with maintaining the same molecular function. In contrast, the divergence

rate at sites with large fitness effects is slow but approximately constant throughout ~4 billion years of

evolution. The saturation at sites with small fitness effects and the slow divergence at sites with large

effects, lead to a substantially slower overall sequence and structural divergence during the second stage,

i.e. after ~1-2 billon years of independent evolution. As a result of this slowdown, for the past billion years

there has not been a substantial divergence in sequence or structure between ancient orthologs with the

same molecular function, i.e. no significant further decrease in their sequence and structural similarity.

Our analysis also demonstrates that different protein sites are usually conserved between

orthologs in independent lineages. This result is likely due to the epistatic nature of protein sequence

landscapes, where mutations neutral in one lineage are often prohibitively deleterious in another[35, 36].

Despite epistatic interactions, our experimental results indicate that there is a correlation between the

average fitness effect of mutations at a protein site, measured in a single species, and the long-term

conservation of the site across multiple species. This result is consistent with the recent study describing

correlated fitness landscapes between distantly related TIM barrel proteins with the same enzymatic

activity[37].

A previous analysis by Povolotskaya *et al.*[2] demonstrated that ancient protein orthologs are still

diverging from each other, but that study did not explicitly considered the effect of conservation of

molecular function. Our analysis shows that the requirement to conserve a specific molecular function

significantly constraints sequence and structural divergence of orthologs. For example, enzymes sharing

only the first three digits of the EC annotation, i.e. only a general class of substrates or cofactors[16],

display a significantly lower sequence identity at large phylogenetic distances compared orthologs sharing

all four EC digits (Figure S4). Similarly, significant sequence identity between homologous restriction

endonucleases is usually limited to isoschizomers, i.e. proteins specific to the same target sequence[38].

These patterns suggest that the effective divergence limit is likely a consequence of functional pressure

to closely preserve the relative spatial positions and dynamical properties of protein residues participating

in ligand binding and catalysis[13]. The conservation of the chemical structures of enzyme substrates,

coupled with the requirement for efficient catalysis, likely makes the long-term evolutionary drift in

sequence space increasingly slow. Further analyses of the biochemical and biophysical constraints

affecting orthologs' evolution will improve our understanding of what structural protein features are

essential for catalysis, and how catalytic efficiency may be compromised by deleterious mutations[39].

Methods

Considered enzyme activities and corresponding protein orthologs. We selected for analysis the

sequences annotated in UniProt[15] with EC numbers associated to the following metabolic pathways (as

defined in the KEGG database[40]): Glycolysis and gluconeogenesis, pentose phosphate pathway, TCA

cycle, purine metabolism, and pyrimidine metabolism. Using protein sequences from 22 diverse

organisms (Table S1) we identified clusters of orthologous groups (COGs) using the EdgeSearch

algorithm[41]. Following previous studies, COGs were defined so that any two proteins from different

species and members of the same COG were considered orthologs[42]. COGs were obtained using the

COGsoft software[41]. Specifically, starting from an all-against-all psi-blast[43] search, setting the

database size at 10^8, and using a maximum considered E-value of 0.1. In order to obtain the largest

number of likely orthologs we did not apply a filter on low complexity or composition-based statistics.

Only proteins sharing the same EC number and assigned to the same COG were compared, and only COGs

with sequences in 10 or more of the 22 species were used.

In order to exclude proteins showing evidence of Horizontal Gene Transfer (HGT), we constructed

a maximum likelihood phylogenetic tree of the 12 prokaryotes used in our analysis using a concatenated

alignment of marker genes[44]. The tree was manually compared to the individual trees of the prokaryotic

sequences sharing EC annotations within each COG; COG-specific trees were built using the GAMMA

model of amino-acid substitution implemented in the RAxML software[45]. Proteins that showed clear

differences in tree topologies, suggesting HGT, were excluded from the analysis.

Models of long-term protein sequence evolution. Divergence times between organisms were obtained from the TimeTree database[46]. Specifically, we used the mean divergence times between pairs of species across published studies (November, 2015). Global sequence identities for pairs of proteins annotated with the same molecular function in the same COG were calculated from pairwise alignments with ClustalW2[47]. Sequence identity was computed as the number of identical sites at aligned positions, divided by the total number of aligned sites, i.e. excluding gaps. In the analysis we only considered EC numbers with pairs of orthologs showing at least 5 different divergence times distributed across 4 billion years.

Models were fitted to equations 1 to 3 using the least-squares minimization algorithm implemented in the MATLAB R2017a fitnlm function (The MathWorks, Inc., Natick, MA). The best fits of the model 1 and model 2 were compared using the F-test. The minimum sequence identity parameter in the model 2 (Y_0, from equation 2) was compared, for each enzymatic activity, to the average global sequence identity between unrelated protein pairs using the Wald test.

To assess the effect of computational annotations on the observed divergence results, we repeated the analysis using only sequences with experimentally validated molecular functions. To keep only sequences with validated molecular functions, we manually reviewed published references for enzyme annotations in the BRENDA database[23], and discarded any annotations that were based exclusively on computational methods or high-throughput studies. After filtering the annotations, we only considered EC numbers with pairs of orthologs showing at least 4 different divergence times distributed across 4 billion years.

Calculation of the divergence rate. Based on Model 3, we determined the divergence rate at a given divergence time t by solving for the derivative of Equation 3 with respect to time:

$$\frac{dy}{dt} = \frac{d\left(100*\left(\frac{R_0*t}{\alpha}+1\right)^{-\alpha}\right)}{dt} = -100 * R_0 \left(\frac{R_0*t}{\alpha} + 1\right)^{-\alpha-1} \tag{5}$$

where y represents global sequence identity, t represents divergence time, R_0 represents the average substitution rate, and α represents the shape parameter of the gamma distribution.

FolA competition experiment in *E. coli*. To perform competition experiments we used the EcNR2 strain derived from *E. coli* K12 MG1655. Mutagenesis was performed using Multiplex Automated Genomic

Engineering (MAGE), as previously described[26]. 90 bp DNA oligomers were designed around each folA

codon using the MG1655 wild type sequence as reference. For each codon, all possible nucleotide variants

were synthesized. To avoid simultaneous mutations of multiple codons, cells were transformed targeting

ten consecutive codons at a time. After four rounds of electroporation, cells were recovered and pooled

together at approximately the same concentration based on cell counts. Two competition growth

experiments were carried out, one for each half of the protein. For the competition experiments, cells

were grown in LB media in a turbidostat while maintaining constant volume and cell density. Samples

were taken every 2 hours for a period of 16 hours, spun down, washed in PBS, spun down again and stored

at -20°C until all samples were collected. For each competition, the corresponding FolA region was

amplified through PCR while assigning a specific DNA barcode for each time point. PCR products were

then pooled and paired-end sequenced using the MiSeq Reagent Kit 2 from Illumina. Sequence reads were

deposited to the SRA database with accession number: SRP152339.

To determine, at each time point, the abundance of each mutant relative to wild type, we joined

paired-end reads using SeqPrep (v 1.1) and aligned the joined reads to the folA gene sequence using

Bowtie2[48]. We then counted the number of reads per mutant using a custom script. Reads with more

than a single mutated codon were discarded. Counts were median-normalized to control for noise due to

mutagenesis performed in batches of 10 codons. At each time point we calculated the ratio R_t of mutant

to wild type (WT) reads. In exponential growth, the growth rate difference between a given mutant and

WT was calculated based on the slope of $\ln(R_\tau)$ as a function of time:

$$\ln(R_t) = (m_i - m_{wt}) * t + \ln(R_0)$$

where m_i and m_{wt} represent the mutant and WT growth rates, respectively. Growth rate differences were

calculated only for mutants with at least 5 time points with 20 or more reads. Relative growth rates were

calculated by dividing the slopes in the equation above by the number of e-fold increases given the

average dilution rate of the turbidostat (1.37/h).

To obtain a single value characterizing the effect of all possible mutations at a protein site, we

first averaged the relative growth rates of mutants resulting in the same amino acid change. We then

calculated the average fitness effect of mutations at each protein site by averaging across 20 possible

amino acids substitutions (Dataset S1).

Divergence of sites with different fitness effects. To investigate divergence of sites with different fitness

effects, we obtained sequences of FolA and InfA bacterial orthologs from the COG database[27]. The FolA

orthologs annotated with the same EC number in UniProt (n=311) and the InfA orthologs annotated with

the same KO number in KEGG (n=514) were used to build multiple sequence alignments with

ClustalW2[47]. We then calculated the genetic distance between all pairs of species with aligned

sequences based on the corresponding 16S rRNA genes obtained from the GreenGenes database[49]

(October, 2016); the F84 model of nucleotide substitution implemented in the phylip package[50] was

used to compute the genetic distances. Divergence times were estimated from the 16S genetic distances

based on the linear regression of literature reported divergence times[46] (Table S1). In each divergence

bin (Figure 2), sequence identities of sites with different average fitness effects (represented by different

colors) were averaged across all pairs of orthologs.

To find phylogenetically independent pairs of species with specific divergence times, we used the

16S rRNA gene alignments for species with FolA and InfA orthologs. We built maximum likelihood

phylogenetic trees based on the 16S alignments using RAxML[45]. We then applied the Maximum Pairing

Problem approach by Arnold *et al*.[28] to find, at specific divergence times, the largest number of edge-

disjoint pairs of species.

Analysis of global protein structural evolution. To study the divergence of protein structures as a function

of time, we obtained PDB codes for all proteins associated with EC numbers in the BRENDA database[23].

We then selected for the analysis species with solved enzyme structures for at least 10 different EC

numbers. Psi-blast searches with a conservative E-value cutoff of 10^{-6} were used to identify orthologs

(defined as bi-directional best hits) in the selected species. The 3D structures of orthologous pairs,

annotated with the same EC number, were aligned using the TM-align program[30] to obtain the C-alpha

RMSD values. Pairs of proteins were not considered if more than 70% of the residues of the shortest

protein could not be structurally aligned. We also removed from the analysis pairs of structures with

flexibility between domains, as they could result in large RMSD values despite significant structural

similarity. To identify such proteins we used the FATCAT[51] software to perform flexible structural

alignments of all structure pairs. We then specifically filtered the pairs that were split into two or more

domains by the FATCAT alignments.

Analysis of the enzyme active sites. To analyze divergence as a function of active site distance we used

protein sequences associated with the 64 EC numbers and 22 species considered in Figure 1 and Figure

S1. To that end, PDB[32] was searched for homologous sequences annotated with the same enzymatic activities and with known 3D structures. Annotations of active site residues for the corresponding structures were obtained from the Catalytic Site Atlas[33], PDB and UniProt-KB[15]. For each PDB structure with available active site information, protein sites were then stratified into different layers according to the distance between their alpha carbons and the centroid of the active site residues. Each pair of orthologs was then aligned using ClustalW2[47] with a homolog in PDB annotated with the same activity and with defined distance layers around the active site; the PDB sequence with the highest sequence identity to either member of the pair was used for the alignment. Sequence identities for different layers were calculated based on the structural positions in the corresponding PDB reference sequences.

References

1. Chothia, C., and Lesk, A.M. (1986). The relation between the divergence of sequence and structure in proteins. The EMBO journal *5*, 823-826.
2. Povolotskaya, I.S., and Kondrashov, F.A. (2010). Sequence space and the ongoing expansion of the protein universe. Nature *465*, 922-926.
3. Lee, D., Redfern, O., and Orengo, C. (2007). Predicting protein function from sequence and structure. Nature reviews. Molecular cell biology *8*, 995-1005.
4. Tian, W., and Skolnick, J. (2003). How well is enzyme function conserved as a function of pairwise sequence identity? Journal of molecular biology *333*, 863-882.
5. Worth, C.L., Gong, S., and Blundell, T.L. (2009). Structural and functional constraints in the evolution of protein families. Nature reviews. Molecular cell biology *10*, 709-720.
6. Rost, B. (2002). Enzyme function less conserved than anticipated. Journal of molecular biology *318*, 595-608.
7. Conant, G.C., and Wolfe, K.H. (2008). Turning a hobby into a job: how duplicated genes find new functions. Nat Rev Genet *9*, 938-950.
8. Wilson, C.A., Kreychman, J., and Gerstein, M. (2000). Assessing annotation transfer for genomics: quantifying the relations between protein sequence, structure and function through traditional and probabilistic scores. Journal of molecular biology *297*, 233-249.
9. Bork, P., Sander, C., and Valencia, A. (1993). Convergent evolution of similar enzymatic function on different protein folds: the hexokinase, ribokinase, and galactokinase families of sugar kinases. Protein science : a publication of the Protein Society *2*, 31-40.
10. Galperin, M.Y., Walker, D.R., and Koonin, E.V. (1998). Analogous enzymes: independent inventions in enzyme evolution. Genome Res *8*, 779-790.
11. Omelchenko, M.V., Galperin, M.Y., Wolf, Y.I., and Koonin, E.V. (2010). Non-homologous isofunctional enzymes: a systematic analysis of alternative solutions in enzyme evolution. Biol Direct *5*, 31.
12. Rost, B. (1997). Protein structures sustain evolutionary drift. Folding & design *2*, S19-24.
13. Lehninger, A.L., Nelson, D.L., and Cox, M.M. (2013). Lehninger principles of biochemistry, 6th Edition, (New York: W.H. Freeman).
14. Firnberg, E., Labonte, J.W., Gray, J.J., and Ostermeier, M. (2016). A Comprehensive, High-Resolution Map of a Gene's Fitness Landscape. Molecular biology and evolution *33*, 1378.

15. UniProt, C. (2015). UniProt: a hub for protein information. Nucleic acids research *43*, D204-212.

16. Bairoch, A. (1999). The ENZYME data bank in 1999. Nucleic acids research *27*, 310-311.

17. Yang, Z., Nielsen, R., Goldman, N., and Pedersen, A.M. (2000). Codon-substitution models for
heterogeneous selection pressure at amino acid sites. Genetics *155*, 431-449.

18. Fares, M.A., Byrne, K.P., and Wolfe, K.H. (2006). Rate asymmetry after genome duplication causes
substantial long-branch attraction artifacts in the phylogeny of Saccharomyces species. Molecular
biology and evolution *23*, 245-253.

19. Fletcher, W., and Yang, Z. (2009). INDELible: a flexible simulator of biological sequence evolution.
Molecular biology and evolution *26*, 1879-1888.

20. Meyer, T.E., Cusanovich, M.A., and Kamen, M.D. (1986). Evidence against Use of Bacterial Amino-
Acid-Sequence Data for Construction of All-Inclusive Phylogenetic Trees. Proceedings of the
National Academy of Sciences of the United States of America *83*, 217-220.

21. O'Malley-James, J.T., Cockell, C.S., Greaves, J.S., and Raven, J.A. (2014). Swansong biospheres II:
the final signs of life on terrestrial planets near the end of their habitable lifetimes. Int J Astrobiol
13, 229-243.

22. Hubble, E. (1929). A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae.
Proceedings of the National Academy of Sciences of the United States of America *15*, 168-173.

23. Chang, A., Schomburg, I., Placzek, S., Jeske, L., Ulbrich, M., Xiao, M., Sensen, C.W., and Schomburg,
D. (2015). BRENDA in 2015: exciting developments in its 25th year of existence. Nucleic acids
research *43*, D439-446.

24. Benkovic, S.J., Fierke, C.A., and Naylor, A.M. (1988). Insights into Enzyme Function from Studies
on Mutants of Dihydrofolate-Reductase. Science *239*, 1105-1110.

25. Kelsic, E.D., Chung, H., Cohen, N., Park, J., Wang, H.H., and Kishony, R. (2016). RNA Structural
Determinants of Optimal Codons Revealed by MAGE-Seq. Cell Syst *3*, 563-571 e566.

26. Wang, H.H., Isaacs, F.J., Carr, P.A., Sun, Z.Z., Xu, G., Forest, C.R., and Church, G.M. (2009).
Programming cells by multiplex genome engineering and accelerated evolution. Nature *460*, 894-
898.

27. Galperin, M.Y., Makarova, K.S., Wolf, Y.I., and Koonin, E.V. (2015). Expanded microbial genome
coverage and improved protein family annotation in the COG database. Nucleic acids research *43*,
D261-269.

28. Arnold, C., and Stadler, P.F. (2010). Polynomial algorithms for the Maximal Pairing Problem:
efficient phylogenetic targeting on arbitrary trees. Algorithms Mol Biol *5*, 25.

29. Watson, J.D., Laskowski, R.A., and Thornton, J.M. (2005). Predicting protein function from
sequence and structural data. Current opinion in structural biology *15*, 275-284.

30. Zhang, Y., and Skolnick, J. (2005). TM-align: a protein structure alignment algorithm based on the
TM-score. Nucleic acids research *33*, 2302-2309.

31. Jack, B.R., Meyer, A.G., Echave, J., and Wilke, C.O. (2016). Functional Sites Induce Long-Range
Evolutionary Constraints in Enzymes. PLoS Biol *14*, e1002452.

32. Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N., and
Bourne, P.E. (2000). The Protein Data Bank. Nucleic acids research *28*, 235-242.

33. Porter, C.T., Bartlett, G.J., and Thornton, J.M. (2004). The Catalytic Site Atlas: a resource of
catalytic sites and residues identified in enzymes using structural data. Nucleic acids research *32*,
D129-133.

34. Halabi, N., Rivoire, O., Leibler, S., and Ranganathan, R. (2009). Protein sectors: evolutionary units
of three-dimensional structure. Cell *138*, 774-786.

35. Breen, M.S., Kemena, C., Vlasov, P.K., Notredame, C., and Kondrashov, F.A. (2012). Epistasis as
the primary factor in molecular evolution. Nature *490*, 535-538.

36. Lunzer, M., Golding, G.B., and Dean, A.M. (2010). Pervasive cryptic epistasis in molecular
evolution. PLoS Genet *6*, e1001162.
37. Chan, Y.H., Venev, S.V., Zeldovich, K.B., and Matthews, C.R. (2017). Correlation of fitness
landscapes from three orthologous TIM barrels originates from sequence and structure
constraints. Nat Commun *8*, 14614.
38. Pingoud, A., Wilson, G.G., and Wende, W. (2014). Type II restriction endonucleases--a historical
perspective and more. Nucleic acids research *42*, 7489-7527.
39. Shendure, J., and Akey, J.M. (2015). The origins, determinants, and consequences of human
mutations. Science *349*, 1478-1483.
40. Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2016). KEGG as a reference
resource for gene and protein annotation. Nucleic acids research *44*, D457-462.
41. Kristensen, D.M., Kannan, L., Coleman, M.K., Wolf, Y.I., Sorokin, A., Koonin, E.V., and Mushegian,
A. (2010). A low-polynomial algorithm for assembling clusters of orthologous groups from
intergenomic symmetric best matches. Bioinformatics *26*, 1481-1487.
42. Tatusov, R.L., Koonin, E.V., and Lipman, D.J. (1997). A genomic perspective on protein families.
Science *278*, 631-637.
43. Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. (1997).
Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic
acids research *25*, 3389-3402.
44. Wu, M., and Scott, A.J. (2012). Phylogenomic analysis of bacterial and archaeal sequences with
AMPHORA2. Bioinformatics *28*, 1033-1034.
45. Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics *30*, 1312-1313.
46. Hedges, S.B., Dudley, J., and Kumar, S. (2006). TimeTree: a public knowledge-base of divergence
times among organisms. Bioinformatics *22*, 2971-2972.
47. Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin,
F., Wallace, I.M., Wilm, A., Lopez, R., et al. (2007). Clustal W and Clustal X version 2.0.
Bioinformatics *23*, 2947-2948.
48. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods
9, 357-359.
49. DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D.,
Hu, P., and Andersen, G.L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and
workbench compatible with ARB. Appl Environ Microbiol *72*, 5069-5072.
50. Felsenstein, J. (2005). PHYLIP (Phylogeny Inference Package) version 3.6. Distributed by the
author: Department of Genome Sciences, University of Washington, Seattle.
51. Ye, Y., and Godzik, A. (2003). Flexible structure alignment by chaining aligned fragment pairs
allowing twists. Bioinformatics *19 Suppl 2*, ii246-255.

Table S1. Model species selected and pairwise average divergence times [a] (in billions of years)[46]

	Eukarya										Bacteria									Archaea		
	Homo sapiens	*Rattus norvegicus*	*Xenopus laevis*	*Xenopus laevis / Danio rerio*	*Gallus gallus*	*Drosophila melanogaster*	*Saccharomyces cerevisiae*	*Caenorhabditis elegans*	*Arabidopsis thaliana*	*Oryza sativa*	*Bacillus subtilis*	*Thermotoga maritima*	*Pseudomonas aeruginosa*	*Mycobacterium tuberculosis*	*Yersinia pestis*	*Haemophilus influenzae*	*Salmonella enterica*	*Escherichia coli*	*Agrobacterium fabrum*	*Pyrococcus horikoshii*	*Sulfolobus solfataricus*	*Haloferax volcanii*
H.sapiens	0.0																					
R. norvegicus	0.1	0.0																				
X. laevis	0.4	0.4	0.0																			
D. rerio	0.4	0.4	0.4	0.0																		
G. gallus	0.3	0.3	0.4	0.4	0.0																	
D. melanogaster	0.8	0.8	0.8	0.8	0.8	0.0																
S. cerevisiae	1.3	1.3	1.3	1.3	1.3	1.3	0.0															
C. elegans	0.8	0.8	0.8	0.8	0.8	0.7	1.3	0.0														
A. thaliana	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	0.0													
O. sativa	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	0.2	0.0												
B. subtilis	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	0.0											
T. maritima	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	0.0										
P. aeruginosa	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	0.0									
M. tuberculosis	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	3.1	0.0								
Y. pestis	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	1.4	3.1	0.0							
H. influenzae	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	1.4	3.1	0.7	0.0						
S. enterica	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	1.4	3.1	0.4	0.7	0.0					
E. coli	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	1.4	3.1	0.4	0.7	0.1	0.0				
A. fabrum	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.1	4.0	2.5	3.1	2.5	2.5	2.5	2.5	0.0			
P. horikoshii	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	0.0		
S. solfataricus	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.8	0.0	
H. volcanii	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.7	3.8	0.0

[a] Divergence times were obtained from literature references in the TimeTree database[46]

Figure S1. The global pairwise sequence identities between pairs of orthologous enzymes are shown as a function of divergence times between the corresponding species. The colored lines indicate fits to the data using the three models of amino acid substitution (Equations 1-3). The dashed black lines represent the average sequence identity for the alignment of unrelated protein sequences. Dot colors indicate the psi-blast E-value of the pairwise alignments for a database size of 3.8×10^5 sequences. Fit parameters are presented in Table S2.

Table S2. Fitted parameters and test results for 64 EC numbers.

EC number	Model1		Model2			Model3			Model 1 Vs. Model 2	Y_0 Wald test
	R_0	R^2	Y_0	R_0	R^2	R_0	α	R^2	F test P-value	P-value[a]
1.1.1.37	0.25	0.30	40.95	0.97	0.51	0.96	0.35	0.51	6.0E-15	<1.0E-20
1.1.1.42	0.20	-2.41	63.12	2.70	0.46	5.32	0.09	0.48	9.2E-97	<1.0E-20
1.1.1.44	0.22	-0.68	49.99	1.59	0.71	2.12	0.19	0.62	3.4E-72	<1.0E-20
1.1.1.49	0.31	-0.36	37.81	1.52	0.58	3.51	0.24	0.76	1.6E-68	<1.0E-20
1.1.1.205	0.27	0.50	37.59	0.93	0.79	0.80	0.46	0.78	2.0E-76	<1.0E-20
1.2.1.12	0.15	-0.47	60.46	1.20	0.68	1.06	0.15	0.68	6.1E-57	<1.0E-20
1.2.4.1	0.46	-0.25	41.05	1.83	0.58	2.60	0.27	0.54	3.9E-49	<1.0E-20
1.2.4.2	0.27	0.09	40.46	1.43	0.92	1.55	0.30	0.86	1.5E-29	<1.0E-20
1.3.5.1	0.20	0.48	47.46	0.77	0.73	0.58	0.34	0.73	1.4E-28	<1.0E-20
1.3.5.2	0.26	-0.74	43.15	1.87	0.83	2.85	0.22	0.82	1.8E-53	<1.0E-20
1.5.1.3	0.33	0.24	33.99	1.69	0.88	1.91	0.36	0.84	2.1E-27	<1.0E-20
1.8.1.4	0.27	0.41	38.56	1.01	0.83	0.91	0.42	0.78	2.8E-68	<1.0E-20
1.8.1.9	0.23	-0.26	43.23	1.15	0.17	5.61	0.17	0.34	2.4E-16	<1.0E-20
2.1.1.45	0.18	0.50	49.39	0.77	0.63	0.62	0.30	0.65	7.2E-06	<1.0E-20
2.1.2.2	0.33	0.38	34.56	1.24	0.82	1.54	0.38	0.87	1.4E-13	<1.0E-20
2.2.1.1	0.29	0.12	36.80	1.38	0.44	1.67	0.34	0.43	4.8E-18	<1.0E-20
2.2.1.2	0.21	0.13	46.70	1.19	0.50	1.12	0.27	0.48	5.4E-14	<1.0E-20
2.3.1.12	0.38	0.23	30.07	1.19	0.76	1.68	0.43	0.79	7.1E-49	<1.0E-20
2.3.1.61	0.24	-0.05	44.35	1.35	0.81	1.53	0.26	0.76	1.9E-29	<1.0E-20
2.4.2.1	0.32	-1.75	36.56	2.77	0.43	9.70	0.20	0.34	8.6E-63	<1.0E-20
2.4.2.7	0.26	-1.45	42.23	2.12	0.62	5.11	0.19	0.48	7.0E-49	<1.0E-20
2.4.2.9	0.27	0.58	28.88	0.56	0.62	0.46	0.94	0.62	3.9E-03	6.50E-04
2.4.2.14	0.29	-1.00	39.38	1.82	0.60	3.90	0.23	0.51	1.1E-113	<1.0E-20
2.7.1.12	0.32	-2.50	38.83	3.73	0.81	10.78	0.18	0.61	2.6E-35	<1.0E-20
2.7.1.15	0.37	-2.29	33.36	2.52	0.71	7.85	0.23	0.50	9.7E-112	<1.0E-20
2.7.1.21	0.36	0.58	27.70	1.23	0.81	1.27	0.56	0.82	1.6E-19	<1.0E-20
2.7.1.25	0.20	-0.71	49.51	1.88	0.85	3.05	0.17	0.73	3.6E-22	<1.0E-20
2.7.1.40	0.34	-0.43	35.98	1.49	0.66	2.65	0.29	0.66	8.1E-179	<1.0E-20
2.7.2.3	0.27	0.50	38.49	0.94	0.82	0.82	0.44	0.79	1.3E-70	<1.0E-20
2.7.4.3	0.24	0.16	42.88	1.16	0.82	1.27	0.29	0.81	1.6E-77	<1.0E-20
2.7.4.6	0.25	-1.11	47.99	2.58	0.40	5.26	0.16	0.57	2.2E-123	<1.0E-20
2.7.4.9	0.36	0.27	26.95	1.19	0.68	1.50	0.52	0.66	9.4E-26	<1.0E-20
2.7.4.22	0.28	0.93	0.00	0.28	0.92	0.24	10.0	0.85	N.S.	N.S
2.7.6.1	0.25	0.13	42.55	1.15	0.60	1.11	0.32	0.54	2.8E-55	<1.0E-20
2.7.7.8	0.23	-0.18	44.23	1.70	0.24	4.87	0.18	0.40	3.5E-10	<1.0E-20
3.1.1.31	0.57	-3.32	31.23	5.33	0.67	15.70	0.23	0.67	1.6E-63	<1.0E-20
3.1.3.11	0.28	0.31	37.89	1.11	0.79	1.28	0.35	0.79	7.7E-32	<1.0E-20
3.5.4.5	0.30	-4.58	39.97	3.18	0.62	29.28	0.14	0.34	9.0E-52	<1.0E-20
3.6.1.23	0.30	0.77	27.28	0.64	0.85	0.55	0.92	0.85	2.1E-10	1.59E-12
4.1.2.4	0.28	0.32	36.16	1.55	0.63	1.56	0.36	0.63	9.6E-09	<1.0E-20
4.1.2.13	0.41	-0.69	52.02	1.80	0.68	1.79	0.20	0.66	1.6E-63	<1.0E-20
4.2.1.2	0.18	-0.50	53.09	1.35	0.47	1.85	0.17	0.43	4.1E-48	<1.0E-20
4.2.1.3	0.22	-0.26	52.94	1.35	0.92	1.32	0.20	0.90	3.7E-71	<1.0E-20
4.2.1.11	0.19	0.23	50.69	0.99	0.73	0.91	0.24	0.72	4.6E-103	<1.0E-20
4.3.2.2	0.40	0.21	26.73	1.25	0.53	1.38	0.57	0.48	7.8E-28	<1.0E-20
5.1.3.1	0.26	-0.16	41.62	1.33	0.56	2.00	0.25	0.57	1.2E-40	<1.0E-20
5.1.3.3	0.54	-2.23	35.37	4.01	0.35	12.77	0.21	0.45	9.3E-95	<1.0E-20
5.3.1.1	0.29	0.53	33.27	0.84	0.70	0.73	0.58	0.68	2.7E-27	<1.0E-20
5.3.1.6	0.32	-0.28	34.56	1.40	0.67	1.94	0.34	0.59	3.2E-31	<1.0E-20
5.3.1.9	0.30	-0.33	38.80	1.66	0.25	3.41	0.24	0.22	4.2E-38	<1.0E-20
5.4.2.2	0.40	0.94	13.27	0.56	0.96	0.52	2.30	0.96	1.7E-07	N.S
5.4.99.18	0.18	0.54	49.56	0.80	0.72	0.65	0.29	0.75	1.8E-07	<1.0E-20
6.2.1.1	0.25	-1.88	47.10	2.58	0.44	8.88	0.14	0.35	1.6E-82	<1.0E-20
6.2.1.5	0.24	0.09	43.31	1.28	0.74	1.37	0.28	0.70	4.3E-43	<1.0E-20
6.3.2.6	0.32	0.53	26.81	0.74	0.60	0.62	0.91	0.58	1.7E-04	4.25E-05
6.3.3.1	0.24	0.83	24.72	0.41	0.84	0.35	1.23	0.85	5.7E-03	2.34E-02
6.3.4.2	0.22	-0.22	48.23	1.35	0.81	1.65	0.22	0.81	1.1E-183	<1.0E-20
6.3.4.4	0.26	0.31	41.12	1.26	0.87	1.23	0.33	0.84	3.4E-156	<1.0E-20
6.3.4.13	0.25	0.77	0.00	0.25	0.76	0.26	20.84	0.76	N.S.	N.S
6.3.4.18	0.33	0.24	32.78	1.44	0.91	1.69	0.40	0.85	2.2E-21	<1.0E-20
6.3.5.2	0.21	0.32	44.93	1.06	0.59	0.95	0.31	0.57	9.7E-15	<1.0E-20
6.3.5.3	0.34	0.24	29.22	1.09	0.56	1.20	0.53	0.53	3.5E-18	<1.0E-20
6.3.5.5	0.23	0.40	40.73	0.92	0.63	0.94	0.36	0.63	4.3E-11	<1.0E-20
6.4.1.1	0.24	-0.18	47.69	1.34	0.96	1.28	0.25	0.88	9.1E-47	<1.0E-20

[a] P-values <0.05 indicate Y_0 is significantly higher than 13.5%, the average identity between unrelated sequences

Figure S2. a. The distribution of Y_0 parameter values across 64 EC numbers for Model 2 fits (Equation 2). The Y_0 parameter represents the minimum fraction of protein sites that remain identical at long divergence times. **b.** The distribution across 64 EC numbers of the projected global sequence identity after 7 billion years of divergence according to Model 3 (Equation 3). In both panels the dashed red lines represent the corresponding median values (39% and 33% for a. and b., respectively).

Table S3. Estimated evolutionary rates for pairs of proteins according to Model 3 fits.

EC number	Initial rate (R_0)	Rate at 4 Byr.	Fold-reduction in rate
1.1.1.37	0.96	0.03	28.3
1.1.1.42	5.32	0.01	377.0
1.1.1.44	2.12	0.02	94.5
1.1.1.49	3.51	0.02	158.5
1.1.1.205	0.80	0.04	20.7
1.2.1.12	1.06	0.02	48.9
1.2.4.1	2.60	0.02	106.2
1.2.4.2	1.55	0.03	54.5
1.3.5.1	0.58	0.04	15.6
1.3.5.2	2.85	0.02	126.3
1.5.1.3	1.91	0.03	67.7
1.8.1.4	0.91	0.04	25.1
1.8.1.9	5.61	0.02	307.5
2.1.1.45	0.62	0.03	18.3
2.1.2.2	1.54	0.03	50.7
2.2.1.1	1.67	0.03	58.0
2.2.1.2	1.12	0.03	38.3
2.3.1.12	1.68	0.03	55.6
2.3.1.61	1.53	0.03	56.5
2.4.2.1	9.70	0.02	561.5
2.4.2.7	5.11	0.02	264.6
2.4.2.9	0.46	0.06	8.3
2.4.2.14	3.90	0.02	182.9
2.7.1.12	10.78	0.02	643.2
2.7.1.15	7.85	0.02	426.1
2.7.1.21	1.27	0.03	36.9
2.7.1.25	3.05	0.02	151.4
2.7.1.40	2.65	0.02	107.4
2.7.2.3	0.82	0.04	21.5
2.7.4.3	1.27	0.03	43.2
2.7.4.6	5.26	0.02	289.7
2.7.4.9	1.50	0.03	46.7
2.7.4.22	0.24	0.09	2.6
2.7.6.1	1.11	0.03	35.3
2.7.7.8	4.87	0.02	256.4
3.1.1.31	15.70	0.02	996.1
3.1.3.11	1.28	0.03	40.8
3.5.4.5	29.28	0.01	2149.6
3.6.1.23	0.55	0.05	10.3
4.1.2.4	1.56	0.03	52.5
4.1.2.13	1.79	0.02	75.4
4.2.1.2	1.85	0.02	85.3
4.2.1.3	1.32	0.02	53.2
4.2.1.11	0.91	0.03	31.6
4.3.2.2	1.38	0.03	41.3
5.1.3.1	2.00	0.03	78.7
5.1.3.3	12.77	0.02	774.3
5.3.1.1	0.73	0.04	17.1
5.3.1.6	1.94	0.03	69.9
5.3.1.9	3.41	0.02	153.5
5.4.2.2	0.52	0.06	8.4
5.4.99.18	0.65	0.03	19.4
6.2.1.1	8.88	0.02	548.7
6.2.1.5	1.37	0.03	47.5
6.3.2.6	0.62	0.05	12.4
6.3.3.1	0.35	0.06	5.3
6.3.4.2	1.65	0.02	66.5
6.3.4.4	1.23	0.03	39.6
6.3.4.13	0.26	0.09	2.9
6.3.4.18	1.69	0.03	56.7
6.3.5.2	0.95	0.03	29.4
6.3.5.3	1.20	0.04	34.4
6.3.5.5	0.94	0.03	27.8
6.4.1.1	1.28	0.03	46.4

Figure S3. Sequence identity as a function of divergence time for orthologous enzymes annotated with the same EC number and experimentally validated molecular functions. The color lines indicate fits to the data for the three models of amino acid substitution (Equations 1-3). Results are shown for 12 different EC numbers. For all EC numbers, Model 2 (red) gives a better fit to the data compared to Model 1 (black, F-test P-value < 0.05). Also, for all EC numbers the minimum sequence identity at long divergence times (Y_0, from equation 2) is higher than the average sequence identity between unrelated protein pairs (~13.5%, dashed black lines) (Wald-test P-value < 0.05).

Figure S4. The distribution of sequence identities at long evolutionary distances between orthologous pairs of enzymes annotated (based on UniProt) with the same EC number (red), and only sharing the first three digits of an EC number (black); points in different sequence identity bins are connected using a spline function (solid lines). The distributions of sequence identities were obtained using pairs of orthologs from species that diverged over two billion years ago. Results are based on all enzyme COGs present in the 22 species used to analyze global sequence divergence (Table S1). To avoid functional biases, orthologs with the same EC number (red, n=8646 pairs) were restricted to the same set of COGs as the ones used for orthologs sharing the first three digits of an EC number (black, n=9123 pairs). The two distributions are significantly different (mean: 37.5 vs. 27.8%, Mann-Whitney P-value <10^{-20}).