Therapeutic importance of timely immunophenotyping of breast cancer in a resource-

constrained setting: a retrospective hospital-based cohort study

Arturo López Pineda, PhD[1,§,*], Mario Felipe Rodríguez Moran, MD[2,§], Cleto Álvarez Aguilar,

MD, MS[2,3], Sarah Marlenne Fuentes Valle, MD[3], Román Acosta Rosales, MD[3], Carlos D.

Bustamante, PhD[1]

[1] Stanford University. Department of Biomedical Data Science. Stanford, California, USA

[2] Universidad Michoacana de San Nicolas de Hidalgo. Faculty of Medical and Biological

Sciences "Dr. Ignacio Chavez". Morelia, Michoacan, Mexico.

[3] Mexican Institute of Social Security. Michoacan Delegation. Morelia, Michoacan, Mexico.

[§] Authors contributed equally to this work.

[*] Corresponding author

Address correspondence to:

Arturo Lopez Pineda, PhD

Department of Biomedical Data Science

Stanford University School of Medicine

365 Lasuen Street, Stanford, CA, 94305, USA

Email: arturolp@stanford.edu

Abstract

Background. The Mexican Consensus of 2015 on diagnosis and treatment of breast cancer

proposes the use of surrogate classification based on immunohistochemistry (IHC). In this study,

we assessed the therapeutic value of this consensus recommendation, in the absence of gene

expression data, as a tool for providing opportune and precise care.

Methods and Findings. In this retrospective hospital-based cohort study, we reviewed medical

records of all suspected breast cancer patients in 2014 treated at the General Regional Hospital

No. 1, of the Mexican Institute of Social Security (IMSS), located in Charo, Michoacan (western

Mexico). Then, we followed the medical history of those patients with IHC testing until March

2017. The trajectory of patients was recorded to discover common patterns, and turnaround time

for IHC testing. There were 402 suspected breast cancer patients in 2014, of which 30 had been

tested for some IHC biomarkers (ER, PR, HER2). The surrogate subtyping allowed doctors to

adjust (56.7 %) or confirm (43.3 %) the initial therapeutic regimen. Opportune IHC testing was

found to be beneficial when it was available before or during the first rounds of chemotherapy.

Conclusions. The use of opportune immunohistochemistry provides improved outcomes for

breast cancer patients. This result has the potential to become a roadmap for the implementation

of precision medicine in Mexico.

Keywords: Breast Neoplasms; Immunohistochemistry; Tumor Biomarkers; Medical Records;

Clinical Coding; Mexico

Abbreviations: IMSS, Mexican Institute of Social Security; HGR1, General Regional Hospital

No. 1 in Charo, Michoacan, Mexico.

1. Introduction

Molecular subtyping for breast cancer tumors based on gene expression patterns have been previously recognized to have prognostic value in the management and treatment of breast cancer patients [1]. During the last decade, there have been several commercially accepted multi-gene prognostic tests. Genomic prognostic tests have been associated with decreased adjuvant chemotherapy use among patients, and better outcomes with cost-effective treatments [2]. However, the availability and expense of these genomic tests remain largely prohibitive for constrain-resourced hospital settings, regardless of country income. For example, in the United States, the use of the 21-gene recurrence score Oncotype DX has increased significantly over the last decade due to insurance coverage, but it is estimated that only a quarter of eligible patients have been tested [3].

In resource-constrained settings, the use of surrogate immunohistochemistry (IHC)-based breast cancer subtypes presents as an alternative to molecular subtyping when there is no access to gene expression data or high-throughput sequencing data. Despite the lowering costs of genomic tests, these technologies are only available to researchers in select locations in the world, and they are inaccessible to most patients worldwide. Alternatively, a panel of three IHC markers—estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2)—demonstrated its predictive value for chemotherapy response in breast cancer [4]. There are other IHC markers that could be included for the prediction of treatment response and prognosis; however, further studies are needed to select an optimized surrogate panel [5].

The 2015 Mexican Consensus on diagnosis and treatment of breast cancer [6] is the main working document for oncologists in the country. It recommends routine use of a 3-marker panel

(ER, PR, HER2) IHC approximation to identify four intrinsic breast cancer phenotypes (luminal

A, luminal B, HER-2, and triple negative). In the Mexican population, the estimated frequency

of ER/PR positive is 60%, HER2 positive 20.4%, and triple negative is 23.1% [7,8]. There is a

troubling onset of disease among younger breast cancer patients in Mexico (aged <40), with a

high prevalence of triple-negative breast cancers [9]. Furthermore, there is an increased mortality

trend associated with breast cancer in the country [10].

The Mexican clinical practice guidelines for screening, management, and treatment of

breast cancer [11-13] also recommends the use of IHC testing as part of the histopathology

studies. Despite these recommendations, the widespread use of genetic testing in Mexico remains

elusive due to constrained resources. However, IHC testing is widely available through hospital-

owned pathology laboratories, as well as through outsourced commercial laboratories across the

country.

In this study, we aimed to assess the timeliness of surrogate IHC biomarker testing in the

context of a resource-constrained hospital setting. Evaluating the therapeutic value for patients is

of critical importance, since the demand for more IHC testing will create new logistical

challenges in the healthcare system. The impact of acceptable or critical time delays is ultimately

not well defined, and it would be of great importance for hospital administrators to have a

reference of the value of timely IHC testing. As precision medicine continues to provide better

prognostic biomarkers, translating these benefits into clinical practice increasingly requires a

time-efficient approach from the entire healthcare system.

2. Methods

Ethical approval

This research was reviewed and approved by the Mexican Institute of Social Security (IMSS)'s National Scientific Research Committee and its National Bioethics Committee, under protocol number R-2017-785-010, and locally by IMSS' Local Research and Ethics Committee, under specific protocol number R-2017-1602-14 HGR1 Charo, Michoacan. Stanford's Institutional Review Board provided a non-human subject determination under eProtocol: 40704. Researchers only received access to non-identifiable data under the rigorous considerations of the applicable Mexican bylaws on health research, patient privacy, and electronic medical records. Consent was not required.

Hospital setting

The Mexican Institute of Social Security is a hybrid single-payer system with an integrated network of hospitals nationwide. In Michoacan (a state located in western Mexico), the General Regional Hospital No. 1 (HGR1) is the designated secondary-care facility for the IMSS-insured population, estimated at 1,288,695 people (28% of the state's population), out of 4,584,471 according to the 2015 census [14].

This hospital is in Charo, a suburban community adjacent to the Michoacan capital (Morelia), and it was built in 2012 to replace the old general hospital. HGR1 is the reference secondary-care facility, which serves as reference for seven General Zone Hospitals and 45 Family Medicine Units within the state of Michoacan. HGR1 has an oncology unit with a full complement of fixed staff and facilities available to all patients. It also offers pathology services, diagnostic imaging, and therapeutic capabilities with access to all approved drugs. Patients with

breast cancer who require chemotherapy are sent to the outpatient medical unit, and patients with

breast cancer who require radiation therapy are provided the service through an outsourced

private service in the same city. The personnel include gynecologists, medical oncologist,

surgical oncologists with significant breast cancer training, adequate numbers of nursing and

pharmacy staff, surgeons with significant training, and cancer pathologists. A self-assessment of

the services and facilities provided by these hospitals is S1 Text, which is based upon a modified

version of the oncological service self-assessment levels for low- and middle-income countries

developed by The International Society of Pediatric Oncology (SIOP) committee on Pediatric

Oncology in Developing Countries (PODC) [15].

Patients

The study population for this research was all cumulative breast cancer patients seen in

2014 by medical staff at the IMSS's General Regional Hospital No 1 in Charo, Michoacan. The

institution's breast cancer census was used to select patients that were seen in 2014 at HGR1. To

avoid selection bias in our study, we did not restrict the selection of patients to members of the

female sex or any age group. The inclusion criteria can be seen in Figure 1.

Figure 1. Inclusion criteria.

A patient with suspected breast cancer is referred from a primary-care facility to HGR1

after being examined by his/her family physicians. If the physician suspects the existence of a

breast lesion, the patient undergoes an imaging assessment using the breast imaging-reporting

and data system (BIRADS) score. Upon arriving at HGR1, the patient was either seen by the

breast cancer clinic (BIRADS 0 and 3) for further imaging studies and/or fine-needle biopsy

assessments, or directly referred to the medical oncology service (BIRADS 4, 5 and 6), at which

point they would undergo surgical biopsies for pathology analysis. The gold-standard diagnosis

of breast cancer would be provided by a board-certified pathologist, who would assign one of the

C50 codes (malignant neoplasms of breast) from the international code of diseases version 10

(ICD10).

We reviewed the medical records (both electronic and paper) of all breast cancer patients

at HGR1, and selected those whose biopsies had also been tested with IHC to detect ER, PR,

HER2 antibodies; then, we followed the medical histories of those patients, ending with March

2017, making note of multiple medical visits to the breast cancer clinic, medical oncology

service, and/or surgical oncology service. From their medical records, we extracted information

about IHC testing (antibody ordered, date of ordering, date of results being obtained),

chemotherapy and hormonal drugs administered, and radiation sessions and surgical procedures

undertaken.

Study design

This was a retrospective hospital-based cohort study, using medical records collected

routinely as part of clinical care. The objective was to understand the therapeutic impact on

breast cancer patients of the time taken to test surrogate immunohistochemistry biomarkers (ER,

PR, HER2) in a resource-constrained hospital in western Mexico. We were interested in

reporting the frequency of treatment selection or treatment adjustment frequencies depending on

breast cancer subtyping and the turnaround times for surrogate immunohistochemistry biomarker

testing. The analysis included timeline trajectories and association rules. The STROBE checklist

is provided in S2 Text. All analyses were performed in R version 3.3.2.

Timeline trajectories. We explored the paper- and electronic-based medical records for

all patients with IHC testing, from the patient's first day at the hospital to the last follow-up

pertaining to this study (occurring before March 2017). First, author MFRM manually reviewed

the patient's medical records and curated a set of possible events (transactions) that occurred to

the patients of this study, annotating the time points for each event for each patient. Then, author

ALP annotated those events with standardized coding from the Unified Medical Language

System (UMLS). When disagreement occurred, both authors discussed the translation to assign a

UMLS code to the medical records in Spanish. Finally, similar events were grouped together in

fewer categories. The UMLS codes, UMLS descriptions, Spanish description, and groupings can

be seen in Table 1.

Table 1. List of medical events with UMLS codes and descriptions in English and Spanish.

UMLS code	UMLS description	Description in Spanish
Visit		
C0008952	Clinic Visits	*Atención inicial en la unidad de medicina familiar (primer nivel de atención).*
C2153644	Visit for: gynecological exam	*Atención inicial en la consulta de ginecología (segundo nivel de atención).*
Oncology		
C1620996	Oncology; primary focus of visit; work-up, evaluation, or staging at the time of cancer diagnosis or recurrence (for use in a medicare-approved demonstration project)	*Atención inicial en la consulta de oncología médica.*
C1617848	Oncology; primary focus of visit; expectant management of patient with evidence of cancer for whom no cancer-directed therapy is being administered or arranged at present; cancer-directed therapy might be considered in the future (for use in a medicare-approved demonstration project)	*Atención subsecuente en la consulta de oncología médica.*
Request lab		
C2186763	Request lab results from pathology	*Solicitud de estudios de patología.*
C2186756	Request lab results from hematology	*Solicitud de estudios básicos de laboratorio clínico.*
C2186777	Request lab results from x-ray	*Solicitud de estudios de imagen (tele de tórax, ultrasonido, mastografía).*
C2186774	Request lab results from CT	*Realización de estudios especiales de imagen (tomografía).*
C2186775	Request lab results from MRI	*Realización de estudios especiales de imagen (resonancia magnética).*
Biopsy / Pathology		
C0177666	Needle biopsy of breast	*Realización de biopsia con aguja fina o gruesa.*
C0585992	Surgical biopsy of breast	*Realización de biopsia por escisión quirúrgica.*
C0807321	Pathology report	*Reporte de estudio histopatológico.*
Chemotherapy		
C0086965	Selection for Treatment	*Selección de tipo de tratamiento (quimioterapia, radioterapia, cirugía).*
C4302504	Chemotherapy started	*Inicio de quimioterapia.*
Surgery / Radiation		
C0436382	Radiotherapy started	*Inicio de radioterapia.*

C0024881	Mastectomy	*Realización de mastectomía radical.*
C0851238	Lumpectomy of breast	*Realización de lumpectomía*

IHC results

C3248285	Quantitative HER2 immunohistochemistry (IHC) evaluation of breast cancer consistent with the scoring system defined in the ASCO/CAP guidelines (PATH)	*Solicitud de estudios de inmunohistoquímica (Receptor de HER2).*
C3248286	Quantitative non-HER2 immunohistochemistry (IHC) evaluation of breast cancer (eg, testing for estrogen or progesterone receptors [ER/PR]) performed (PATH)	*Solicitud de estudios de inmunohistoquímica (Receptores de estrógeno y progesterona).*

Treatment adjusted

C1627778	Treatment adjusted per protocol	*Ajuste de tratamiento de primera línea (posterior al primer ciclo de quimioterapia).*
C0419989	Hormone replacement therapy started	*Inicio de tratamiento de reemplazo hormonal.*

Hospitalization / Metastasis

C0019993	Hospitalization	*Complicación de proceso oncológico (internamiento a causa de cáncer)*
C1458156	Recurrent Malignant Neoplasm	*Recaída de proceso oncológico (mismo sitio tumor primario).*
C3694291	Metastasis from malignant neoplasm of breast	*Reconocimiento de metástasis (tumor secundario).*

Remission

C0687702	Cancer Remission	*Evolución favorable de paciente, mejoría clínica; remisión.*

Missing data. Missing data was assumed to be non-missing at random (NMAR), and the

events from Table 1 were considered missing for a specific reason. For example, if the code

C2186775 (request lab result from MRI) was missing from a patient's medical record, we left

that value unassigned, since it was assumed that an MRI was not requested for that patient. There

is a small chance that this assumption might be violated (and introduce some bias) if the

physician did request the exam/procedure but forgot to write it down in the medical record.

However, we did not impute any missing data, since that might be a larger source of bias for this

EHR-based data.

Outliers. We assessed the time-to-event distribution for each event in Table 1, and

reported the mean time using boxplots with whiskers. The outliers were identified using the 1.5

interquartile rule, but were not removed from the downstream analysis since undefined time-to-

event standards exist for these events.

Clustering. A heatmap with all the patients and events was generated. We used

hierarchical clustering to create dendograms. The distance matrix was generated with the time-

to-event matrix, filling all missing events with a negative number (–1000). Distance was

calculated with Pearson's correlation, while linkage was computed as complete.

Association rules. The Apriori algorithm [16] was used to discover association rules that

are common across multiple patients. Apriori first counts event occurrences to determine the

largest 1-item sets. Then, in subsequent steps, it iteratively tries to generate larger candidate item

sets, based on an incremental one-step from the previous step, and assign the strength of

evidence for that set of events based on the number of occurrences in the database. The Apriori

algorithm uses a bottom-up approach. The algorithm finalizes when no candidate list hast larger

item sets. The candidate generation step uses a breadth-first search to efficiently count candidate

item sets, and prune candidates with infrequent patterns. The algorithm does not consider the

time-to-event but only whether the event occurred and in what order it occurred for that patient.

3. Results

The epidemiological information obtained pertaining to this one IMSS hospital can be

seen in Table 2. The results show the patients with breast cancer in 2014 (N=402), which

correlate with national estimates for the country [10]. In this cohort, all patients were female. The

highest prevalence of cancer was within the 50-59 year old age group. The majority of patients

were diagnosed with early stages of cancer (stages I and II). However, only a few number of

patients (N=30) had undergone subtyping with IHC testing in accordance with the Mexican

Consensus. Even in this small cohort, we can still see the same higher prevalence of triple

negative breast cancer cases that was previously reported [17]. It is troubling to see that there

were 18 (4% of) patients younger than 40 years old.

At IMSS, patients with Breast Imaging Reporting and Data System (BIRADS) reports

scoring 1 or 2 are usually deemed low-risk for breast cancer and they have regular annual

screenings at their Family Medicine Unit (UMF). Meanwhile, patients with BIRADS reports 0

and 3 are sent to IMSS's secondary care breast cancer clinic for a closer screening (usually fine-

needle biopsy and 6-month screening); while patients with BIRADS reports 4, 5 or 6 are sent

directly to the secondary or tertiary care oncology services. At the IMSS-HGR1 hospital that we

investigated, 23 patients (6%) had a BIRADS score of 2 or lower, which were not supposed to be

sent to the secondary care facility. In these cases, patients had external biopsy investigations in

non-IMSS hospitals (usually private), and decided to continue their care for follow-up treatment

at the IMSS oncology services.

Table 2. Clinical characteristics of the cohort and receptor status for patients with IHC testing

Characteristic	Total (%)
Overall	**N = 402**
Sex	
Female	402
Male	0
Age at diagnosis	
<40	18 (4%)
40-49	62 (15%)
50-59	174 (43%)
60-69	127 (32%)
70+	21 (5%)
BIRADS	
0,1,2	23 (6%)
3	203 (50%)
4,5,6	162 (40%)
unknown	14 (3%)
Tumor stage	
I	85 (21%)

II	122 (30%)
III	84 (21%)
IV	19 (5%)
unknown	92 (23%)
Patients with IHC testing	**N = 30**
ER testing	
positive	17 (57%)
negative	11 (37%)
unknown	2 (7%)
PR testing	
positive	16 (53%)
negative	12 (40%)
unknown	2 (7%)
HER2 testing	
positive	5 (17%)
negative	23 (77%)
unknown	2 (7%)
Additional IHC testing	
Ki67, P53	1 (3%)
unknown	29 (97%)
Subtype according to the Consensus	14 (47%)
Luminal A	3 (10%)
Luminal B	9 (30%)
Basal-like / Triple negative	2 (7%)
HER2	2 (7%)
not enough information to assign	

Clustering analysis

A hierarchical clustering and heatmap of time-to-events is shown in Figure 2. The time in

which each event occurred for each patient is measured using information found in their medical

record (in days, positive). To be able to create a dissimilarity matrix to calculate distance

between patients, all missing values were assigned a negative value of -1000. It is important to

note that this figure does not provide a context for the ordering in which events occurred, but it

can still help to provide information about patients with similar trajectories. Three main groups

of patients (or clusters) were identified in this graph.

Figure 2. Heatmap of time-to-event with patient clustering.

Patient trajectories

Medical trajectories for patients in the cohort with IHC testing can be seen in Figure 3, clustered according to the groupings shown in Figure 2. Each patient is represented in a panel (rectangle) with colored bars, indicating the events that a patient experienced in the IMSS hospital. Each row represents one year of follow-up treatment for that patient, which can be from one to four rows (because the time maximum time span is three years and five months). The length of each colored bar represents the time between the occurrence of an event and the event that preceded it. Although events do not occur continuously, but, rather, happen at single time points (e.g. a patient visit, obtaining the results of an x-ray), the visual representation shown in Figure 3 provides a sense of how much time was required for each event to occur, assuming nothing else happened at the same time.

Figure 3. Medical trajectories of the cohort with IHC testing. Each patient is represented by a

rectangle, with rows representing each year of follow-up treatment. Events are color-coded

according to the type of event, and the length of the bar represents the duration between the

occurrence of an event and the event that preceded it.

Turnaround time

The time-to-event for all events is shown in Figure 4. It is measured as the number of

days that it took for an event to occur, from the initial visit to the hospital stay (day 0). From this

Figure, it is easy to see that visiting the hospital or beginning treatment occurred relatively at the

beginning of the patients' trajectories, while events that typically take place later did in fact

occur at a posterior time, such as remission or metastasis. Also, one can see there is a wide

distribution of time, when requesting laboratory/imaging testing, which coincides with the

continuity of care.

Figure 4. Boxplots for time-to-event

The turnaround time for immunohistochemistry (IHC) testing can be seen in Figure 5. Only 20 patients had information in their medical records relating to the timing of IHC testing. Although this service is referred to in the IMSS medical records, it is subcontracted to a private laboratory. There was large variation surrounding two key variables: a) the time needed to request IHC testing relative to the first day of follow-up treatment at the hospital, and b) the turnaround time to obtain those results. For the first one, the average time of request was 117 days (95% C.I. 80 – 154 days). For the second one, the average turnaround time for obtaining those results was 56 days (95% C.I. 36 – 77 days).

There were 15 out of 20 patients whose treatments were adjusted, in accordance with IMSS protocol, after obtaining the results of the IHC testing. Overall, for the 30 patients with

IHC testing, 17 patients (56.7%) had their treatments adjusted, and 13 patients (43.3%) had their

treatments confirmed.

Figure 5. Turnaround times for IHC testing. The size of the bar indicates time in days from

requesting results for IHC testing to obtaining them. For each patient, the initial time is the first

day of follow-up treatment at the hospital. The color of the bar indicates the quartile within

which the data falls in the distribution. The diamond mark indicates that the treatment was

adjusted, according to protocols, after obtaining IHC results. The blue dashed line indicates the

1st quartile for the time it took before an IHC request was submitted. Violin plots with whiskers

are correspondingly provided for the time to IHC request (from start to request), time to IHC

results (from start to the obtaining of results), and turnaround time of IHC testing (from request

to results).

Association rules

Association rules were mined from the events of this cohort using the Apriori algorithm.

The five top-ranking association rules are shown in Figure 6. These rules have a support of

63.3% of cases, with a confidence of 90.5% and a rule performance (lift, or precision) of 1.3. For

rules with lift larger than 1, it can be assumed that the right-hand side of the rule is a

consequence of the left-hand side. In our mined rules, we can observe that, in general, IHC

testing, particularly ER/PR testing, results in adjusting treatment plans.

Figure 6. Top-5 association rules mined with the Apriori algorithm

4. Discussion

Key findings

Breast cancer in Mexico (and worldwide) continues to be a public health concern, and

increasing research is being done on the characterization of better genomic variants. Although

our study investigated a small number of patients, the prevalence and epidemiological

characteristics among our cohort are similar to what has previously been estimated for the

country (i.e. in regards to age group and staging) [10]. Our study further investigates the

timeliness of IHC testing, so our conclusions can then be used to inform hospital administrators

and public health officials around the world about the importance of timely IHC testing.

On the therapeutic value. The therapeutic importance of immunophenotyping (the use of

IHC testing for subtyping of cancer patients) needs to be carefully considered and addressed in

treatment plans by oncologists. IMSS protocol states that estrogen positive (ER+) and/or

progesterone positive (PR+) patients could benefit from hormone replacement therapy to prevent

recurrence. For the patient, that usually means that the traditional FEC chemotherapy regimen

composed by 5-fluorouracil, epirubicin, and cyclophosphamide, would be adjusted to include

some hormone replacement therapy, either aromatase inhibitors to block estrogen production

(e.g. anastrozole, letrozole, or exemestane), or some drug that could interfere with the ability of

estrogene to stimulate the growth of breast cells (e.g. tamoxifene or toremifene).

In contrast, patients with HER2 positive (HER2+), which tumors tend to grow and spread

more aggressively, could benefit from a targeted therapy that could prescribe drugs to block

HER2 (e.g. trastuzumab or pertuzumab). However, the use of these drugs can have different side

effects than traditional chemotherapy. The use of in situ hybridization (ISH), instead of IHC, can

be used to determine HER2 status with an overall concordance with IHC, and it may be more

beneficial to use both [18]. There is a need to balance sensitivity of the study with hospital

workload and expenses. Furthermore, HER2 status has been successfully incorporated into

medical practice to guide treatment decisions for breast cancer patients [19]. In fact, the

American Society of Clinical Oncology/College of American Pathologists (ASCO/CAP) updated

their 2013 guidelines to designate more patients as eligible for trastuzumab therapy, in

accordance with ISH and IHC testing [20].

On the timeliness of testing. Opportune IHC testing, understood as the availability of IHC

results before the beginning of any treatment, is of critical importance. For IMSS, the timeliness

of IHC results were typically not opportune, but whenever they were available, they triggered a

response from the clinical oncology team to adjust patient treatments. The widespread

distribution of time to request and turnaround time demonstrates the lack of standardization of

this process. Many factors could have contributed to the delay in IHC testing, including: patient

non-adherence to appointments; hospital overflow, resulting in longer appointments; logistical

problems relating to IHC testing's status an external service to the IMSS hospital of this study;

and administrative problems in the hospital resulting in inadequate tracking of tests and results.

<u>On the need for improved clinical guidelines</u>. It is important to establish an appropriate

timeline for opportune IHC testing. For example, In the United States, the College of American

Pathologists' (CAP) guideline on turnaround time for standard biopsies is around two days for

90% of cases [21]. In fact, the joint guideline on HER2 testing from the American Society of

Clinical Oncology (ASCO) and CAP recommends informing the patients about the expected

turnaround time [22]. In Australia, a study with more than 78,000 patients tested for HER2

status, with both ISH and IHC, revealed that the average turnaround time was between 4-5 days,

which decreased due to the increased volume of patients being tested [23]. In Saudi Arabia, a

study in 2015 showed that 24% of cases fall outside the recommended CAP turnaround time

[24]. The European Society for Medical Oncology (ESMO) published a survey on 24 European

countries were the turnaround time was 10 days or less for 89% of laboratories [25]. In the

United Kingdom, the Royal College of Pathologists recommend the use of key performance

indicators to have a histopathology diagnosis within seven days of biopsy in 90% of cases [26].

Given their resource-available settings, clinical guidelines in developed countries do not

have a recommendation regarding the maximum turnaround time that could be effective in a

patient's treatment trajectory. More importantly, they do not address the time between initial

diagnosis, molecular testing (either via IHC or genetics), and the selection of treatment. In

Mexico, clinical guidelines for breast cancer recommend testing for ER, PR, and HER2 as part of

the histopathology study, but fail to provide guidance regarding turnaround time [13]. IMSS

maintains a medical procedures manual, which includes ten key indicators for breast cancer

screening, diagnosis, and treatment [27]. This manual measures time-to-diagnosis in a 30-day

period, including imaging through mammography and results of histopathology report. In

addition, the institution also measures time-to-treatment measured from the date of diagnosis,

which should be achieved within a 21-day period. Recognizing the resource constraints of IMSS,

the implementation of a key indicator policy related to IHC testing might significantly reduce

turnaround times.

Limitations

On the need for accurate electronic health records (EHR). Missing information was

common across the paper and electronic records in our study. Clinicians and administrative staff

at the IMSS hospital are still getting used to the novel implementation of the EHR, which had an

impact on our ability to better characterize this cohort. In the absence of a cancer registry, our

best estimate of the disease is the institutional census. As the hospitals in Mexico, and elsewhere,

continue to become more and more electronic, there is a need to develop better software tools to

analyze the information obtained, including medical natural language processing and machine

learning applications.

On the visual representation. The medical trajectories shown in Figure 1 are useful in

quickly providing a patient history overview of hospital care. With more work on the user

interface, we envision this tool could eventually represent a valuable visual aid in which a patient

and their companions might be able to better communicate with their clinicians about the

management of their disease (it could probably also appear in a printed version). Currently, some

of the challenges of this tool include the missing identification of events that are overlapping,

and the impossibility of further elaborating on the details of each event. For a hospital

administrator, Figure 1 can provide a quick overview of the patients seen in their hospital, which

could be used as a decision-making tool, with the proper validation. This visual aid creation tool

can be obtained from the repository at https://github.com/arturolp/patientTrajectory.git

Healthcare system implications

Adverse events in the trajectory of an oncological patient, which might include

hospitalization related to cancer, recurrence of tumor, or metastasis, are extremely costly for the

healthcare system. The use of IHC testing was shown in our study to help with the selection of

precise treatment for patients (either by adjusting the treatment or confirming it). The association

rules found in our study confirm that ER and PR testing are associated with the adjustment of

treatment. However, the time in which IHC testing is performed is of critical importance if we

want to influence improved prognosis.

IHC testing is being carried out as an external service at the IMSS hospital in our study.

As the Mexican healthcare system continues to transition from reactive to preventative care, the

need for more IHC testing in breast cancer and other diseases will certainly allow for the further

development of its own testing facilities, therefore allowing for some economy of scale.

We have shown that the use of surrogate immunohistochemistry, as described by the

Mexican Consensus and previous literature worldwide, can have a beneficial therapeutic effect

on breast cancer patients. The aims of any healthcare system should be the identification of

earlier events that can have an impact on downstream events in the trajectory of an oncological

patient's treatment. In resource-constrained settings, it is important not only to consider surrogate

alternatives to more costly diagnostics (e.g. genomic testing), but also to incorporate the

regulatory and logistical aspects of implementing these surrogate IHC tests. IMSS must face the

important challenge of continuing to improve their turnaround times, which should have a

positive impact on the prognosis of their patients. Finally, other healthcare systems would also

benefit from determining the maximum turnaround time in which IHC testing can still be

opportune and have a therapeutic beneficial effect.

Acknowledgements

The authors wish to acknowledge Prof. Ami S. Bhatt, MD, PhD, for her valuable

contribution and comments regarding this project. Also to Katie Kanagawa, PhD, for her

valuable support in editing this manuscript.

Authors' contributions

Conceptualization: MFRM, ALP, and CAA

Data curation: MFRM

Formal analysis: MFRM and ALP

Funding acquisition: CDB

Investigation: MRFM, ALP, SMFV, and CAA

Methodology: ALP

Resources: RAR and CDB

Software: ALP

Supervision: CDB, RAR, and CAA

Visualization: MFRM, ALP

Writing – original draft: ALP

Writing – review & editing: MFRM, CAA, SMFV, RAR, CDB, ALP

Data Availability

Data belongs to the Mexican Institute of Social Security (IMSS), through its state

Delegation of Michoacan. IMSS's Coordination of Health Research may grant access to this data

on a case-by-case basis to researchers who obtain the necessary approvals from IMSS's National

Scientific Research Committee and National Bioethics Committee.

Funding

Research reported in this publication was supported by grant 8316sc-7 from the joint

Center of Excellence in Regulatory Science and Innovation (CERSI) at the University of

California, San Francisco and Stanford University. The funder had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests

RAR declares to be IMSS's state delegate in Michoacan, overseeing the hospital mentioned in this study. The remaining authors declare no conflicts of interest.

References

1.   Sørlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. Proc Natl Acad Sci U S A. National Acad Sciences; 2001;98: 10869–10874. doi:10.1073/pnas.191367098

2.   Abu-Khalf M, Pusztai L. Influence of genomics on adjuvant treatments for pre-invasive and invasive breast cancer. Breast. 2013;22 Suppl 2: S83–7. doi:10.1016/j.breast.2013.07.015

3.   Enewold L, Geiger AM, Zujewski J, Harlan LC. Oncotype Dx assay and breast cancer in the United States: usage and concordance with chemotherapy. Breast Cancer Res Treat. Springer US; 2015;151: 149–156. doi:10.1007/s10549-015-3366-7

4.   Lips EH, Mulder L, de Ronde JJ, Mandjes IAM, Koolen BB, Wessels LFA, et al. Breast cancer subtyping by immunohistochemistry and histological grade outperforms breast cancer intrinsic subtypes in predicting neoadjuvant chemotherapy response. Breast Cancer Res Treat. Springer US; 2013;140: 63–71. doi:10.1007/s10549-013-2620-0

5.   Won JR, Gao D, Chow C, Cheng J, Lau SYH, Ellis MJ, et al. A survey of immunohistochemical biomarkers for basal-like breast cancer against a gene expression

profile gold standard. Mod Pathol. 2013;26: 1438–1450. doi:10.1038/modpathol.2013.97

6.   Cardenas Sanchez J, Bargallo Rocha JE, Erazo Valle A, Poitevin Chacon A, Valero

Castillo V, Perez-Sanchez V. [Consenso Mexicano sobre diagnóstico y tratamiento del

cáncer mamario]. 6 ed. 2015. pp. 1–149.

7.   Lara-Medina F, Perez-Sanchez V, Saavedra-Perez D, Blake-Cerda M, Arce C, Motola-

Kuba D, et al. Triple-negative breast cancer in Hispanic patients: high prevalence, poor

prognosis, and association with menopausal status, body mass index, and parity. Cancer.

2011;117: 3658–3669. doi:10.1002/cncr.25961

8.   Robles-Castillo J, Ruvalcaba-Limón E. [Cáncer de mama en mujeres mexicanas menores

de 40 años]. Ginecologia y Obstetricia de Mexico. 2011;79: 482–488.

9.   Villarreal-Garza C, Alvarez-Gomez RM, Perez-Plasencia C, Herrera LA, Herzog J,

Castillo D, et al. Significant clinical impact of recurrent BRCA1 and BRCA2 mutations in

Mexico. Cancer. 2015;121: 372–378. doi:10.1002/cncr.29058

10.   Mohar-Betancourt A, Reynoso-Noveron N, Armas-Texta D, Gutierrez-Delgado C, Torres-

Dominguez JA. Cancer Trends in Mexico: Essential Data for the Creation and Follow-Up

of Public Policies. Journal of Global Oncology. 2017;: JGO.2016.007476–9.

doi:10.1200/JGO.2016.007476

11.   Secretaria de Salud de Mexico. [Diagnóstico y Tratamiento de la Patología Mamaria

Benigna en Primer y Segundo Nivel de Atención]. 2009;: 1–66.

12.   Secretaria de Salud de Mexico. [Prevention, Screening, and Timely Referral of Suspected

Breast Cancer in the First Level of Care]. 2011 pp. 1–74.

13.   Secretaria de Salud de Mexico. [Diagnóstico y Tratamiento del Cáncer de Mama en

Segundo y Tercer Nivel de Atención]. 2012;: 1–102.

14.   Instituto Nacional de Estadística y Geografía. [Encuesta Intercensal 2015]. 2015.

15.   Parkes J, Hendricks M, Ssenyonga P, Mugamba J, Molyneux E, Schouten van Meeteren

A, et al. SIOP PODC adapted treatment recommendations for standard-risk

medulloblastoma in low and middle income settings. Pediatric Blood & Cancer.

2015;62: 553–564. doi:10.1002/pbc.25313

16.   Agrawal R, Srikant R. Fast algorithms for mining association rules. Proc 20th int conf

very large data bases. 1994.

17.   Villarreal-Garza C, Weitzel JN, Llacuachaqui M, Sifuentes E, Magallanes-Hoyos MC,

Gallardo L, et al. The prevalence of BRCA1 and BRCA2 mutations among young

Mexican women with triple-negative breast cancer. Breast Cancer Res Treat. Springer US;

2015;150: 389–394. doi:10.1007/s10549-015-3312-8

18.   Solomon JP, Dell'Aquila M, Fadare O, Hasteh F. Her2/neu Status Determination in Breast

Cancer: A Single Institutional Experience Using a Dual-Testing Approach With

Immunohistochemistry and Fluorescence In Situ Hybridization. Am J Clin Pathol.

2017;147: 432–437. doi:10.1093/ajcp/aqw224

19.   Goddard KAB, Bowles EJA, Feigelson HS, Habel LA, Alford SH, McCarty CA, et al.

Utilization of HER2 genetic testing in a multi-institutional observational study. Am J

Manag Care. NIH Public Access; 2012;18: 704–712.

20.   Lim TH, Lim AST, Thike AA, Tien SL, Tan PH. Implications of the Updated 2013

American Society of Clinical Oncology/College of American Pathologists Guideline

Recommendations on Human Epidermal Growth Factor Receptor 2 Gene Testing Using

Immunohistochemistry and Fluorescence In Situ Hybridization for Breast Cancer. Arch

Pathol Lab Med. 2016;140: 140–147. doi:10.5858/arpa.2015-0108-OA

21.   Novis DA, Zarbo RJ, Saladino AJ. Interinstitutional comparison of surgical biopsy

505   diagnosis turnaround time: a College of American Pathologists Q-Probes study of 5384

506   surgical biopsies in 157 small hospitals. Arch Pathol Lab Med. 1998;122: 951–956.

507   22.   Wolff AC, Hammond MEH, Hicks DG, Dowsett M, McShane LM, Allison KH, et al.

508   Recommendations for Human Epidermal Growth Factor Receptor 2 Testing in Breast

509   Cancer: American Society of Clinical Oncology/College of American Pathologists

510   Clinical Practice Guideline Update. J Clin Oncol. 2013;31: 3997–4013.

511   doi:10.1200/JCO.2013.50.9984

512   23.   Morey AL, Brown B, Farshid G, Fox SB, Francis GD, McCue G, et al. Determining

513   HER2 (ERBB2) amplification status in women with breast cancer: final results from the

514   Australian in situ hybridisation program. Pathology. Elsevier; 2016;48: 535–542.

515   doi:10.1016/j.pathol.2016.05.007

516   24.   Alshieban S, Al-Surimi K. Reducing turnaround time of surgical pathology reports in

517   pathology and laboratory medicine departments. BMJ Qual Improv Rep. BMJ Open

518   Quality; 2015;4: u209223.w3773. doi:10.1136/bmjquality.u209223.w3773

519   25.   Boleij A, Tembuyser L, Taylor A, Kafatos G, Jenkins-Anderson S, Tack V, et al. PD-

520   015A survey on current RAS-mutation testing practices in Europe. Ann Oncol. Oxford

521   University Press; 2015;26: iv105–iv105. doi:10.1093/annonc/mdv234.14

522   26.   Pathologists TRCO. Key Performance Indicators in Pathology. 2013;: 1–23.

523   27.   Instituto Mexicano del Seguro Social. [Manual Metodológico de Indicadores Médicos

524   2016]. 2016;: 1–452.

```
┌─────────────────────────────┐
│  Cummulative patients with  │
│  confirmed breast cancer    │
│  with visits to HGR1 in 2014│
│  (Institutional census)     │
│  N = 402                    │
└─────────────────────────────┘

                    ┌─────────────────────────────┐
                    │  without IHC testing        │
                    │  (Jan 2014 – Mar 2017)      │
                    │  N = 372                    │
                    └─────────────────────────────┘

                    ┌─────────────────────────────┐
                    │  Deceased                   │
                    │  (Jan 2014 – Mar 2017)      │
                    │  N = 0                      │
                    └─────────────────────────────┘

┌─────────────────────────────┐
│  with IHC testing           │
│  (Jan 2014 – Mar 2017)      │
│  N = 30                     │
└─────────────────────────────┘
```

	C3248286 ER/PR testing	**C1627778** Treatment adjusted
C3248285 HER2 testing	**C3248286** ER/PR testing	**C1627778** Treatment adjusted
C0807321 Pathology report	**C3248286** ER/PR testing	**C1627778** Treatment adjusted
C0086965 Selection for treatment	**C3248286** ER/PR testing	**C1627778** Treatment adjusted
C1617848 Oncology visit (no therapy)	**C3248286** ER/PR testing	**C1627778** Treatment adjusted