In the Absence of Apoptosis, Myeloid Cells Arrest When Deprived of Growth Factor,

But Remain Viable by Consuming Extracellular Glucose

Li Dong[1], Boris Reljic[1], Jen G. Cheung[1], Elizabeth S. Ng[2], Lisa M. Lindqvist[1],

Andrew G. Elefanty[2,3], David L. Vaux[1] and Hoanh Tran[1#]

[1]The Walter and Eliza Hall Institute of Medical Research, and the Department of

Medical Biology, University of Melbourne, Victoria, Australia.

[2]Murdoch Childrens Research Institute, The Royal Children's Hospital, Parkville,

Victoria, Australia.

[3]Department of Anatomy and Developmental Biology, Faculty of Medicine, Nursing

and Health Sciences, Monash University, Clayton, Victoria, Australia.

Running Head: Cell Survival Without ATG5-dependent Autophagy

[#]Address correspondence to Hoanh Tran, tran.h@wehi.edu.au

Abstract

Withdrawal of the growth factor interleukin 3 from IL3-dependent myeloid cells

causes them to undergo Bax/Bak1-dependent apoptosis, whereas factor-deprived $Bax^{-/-}$

$Bak1^{-/-}$ cells remain viable, but arrest and shrink. It was reported that withdrawal of

IL3 from $Bax^{-/-}Bak1^{-/-}$ cells caused decreased expression of the glucose transporter

Glut1, leading to reduced glucose uptake, so that arrested cells required Atg5-

dependent autophagy for long-term survival. In other cell types, a decrease in Glut1 is

mediated by the thioredoxin-interacting protein Txnip, which is induced in IL3-

dependent myeloid cells when growth factor is removed. We mutated *Atg5* and *Txnip*

by CRISPR/Cas9 and found that Atg5-dependent autophagy was not necessary for the

long-term viability of cycling or arrested $Bax^{-/-}Bak1^{-/-}$ cells, and that Txnip was not

required for the decrease in Glut1 expression in response to IL3 withdrawal.

Surprisingly, Atg5-deficient *Bax/Bak1* double mutant cells survived for several weeks

in medium supplemented with 10% fetal bovine serum (FBS), without high

concentrations of added glucose or glutamine. When serum was withdrawn, the

provision of an equivalent amount of glucose present in 10% FBS (~0.5 mM) was

sufficient to support cell survival for more than a week, in the presence or absence of

IL3. Thus, $Bax^{-/-}Bak1^{-/-}$ myeloid cells deprived of growth factor consume extracellular

glucose to maintain long-term viability, without a requirement for Atg5-dependent

autophagy.

Introduction

Mammalian cells import a variety of extracellular nutrients to support their survival, growth and proliferation. *In vitro*, cells are typically propagated in medium supplemented with glucose, glutamine and fetal bovine serum. To generate chemical energy, glucose can be metabolized by oxidative phosphorylation in the mitochondria or by anaerobic glycolysis. Glutamine can be used as an energy source through glutaminolysis, and can also be used, together with a multitude of factors present in serum, to allow the synthesis of essential macromolecules for cell division.

Factor dependent myeloid (FDM) progenitor cell lines[1, 2] require the hematopoietic growth factor interleukin 3 (IL3) to survive and multiply in culture. When IL3 is not provided, FDM cells undergo Bax/Bak1-dependent apoptosis. FDM cells derived from *Bax⁻/⁻Bak1⁻/⁻* mice survive in the absence of IL3, but they undergo cell cycle arrest, similar to those over-expressing Bcl-2[3, 4]. It has been reported that to survive longer than 4 days, IL3-deprived *Bax⁻/⁻Bak1⁻/⁻* cells required Atg5-dependent autophagy[4], an intracellular catabolic process that can support the survival of starved or arrested cells[5].

Cells import glucose when it is available and growth factors can stimulate the amount of glucose that is consumed. *Bax⁻/⁻Bak1⁻/⁻* FDM cells arrested by IL3 withdrawal decrease expression of the glucose transporter Glut1, which was proposed to reduce glucose uptake and metabolism[4]. Several mechanisms have been put forward to explain the regulation of expression and plasma membrane translocation of Glut1. Activation of autophagy was shown to be required for the efficient surface expression of Glut1 in mouse embryonic fibroblasts cultured under hypoxic conditions[6]. In

human hepatocytes subjected to glucose starvation, a mechanism involving

proteasomal degradation of the thioredoxin-interacting protein Txnip was required for

Glut1 stabilization[7]. Moreover, it was shown that Txnip directly binds to Glut1 and

promotes its internalization, thus reducing Glut1 on the plasma membrane and the

ability of cells to take up glucose[7].

In this study, we employed $Bax^{-/-}Bak1^{-/-}$ FDM cells to genetically test whether Atg5-

dependent autophagy is required for the sustained survival of arrested cells deprived

of growth factor. We also asked whether autophagy and/or Txnip are required for the

reduction in Glut1 expression and glucose uptake by arrested myeloid cells.

Results

$Bax^{-/-}Bak1^{-/-}$ myeloid cells shrink and arrest when deprived of IL3, but survive.

Factor dependent myeloid (FDM) cells require interleukin 3 (IL3) to survive and

proliferate[2, 4, 8]. When deprived of IL3 for 4 days, >80% of wild-type $Bax^{+/+}Bak1^{+/+}$

FDM cells died, as indicated by loss of plasma membrane integrity and uptake of

propidium iodide (Figs. 1A and Supplementary S1A). By contrast, and consistent with

the findings of Lum *et al.*[4], $Bax^{-/-}Bak1^{-/-}$ FDM cells survived in the absence of IL3 for

more than 14 days. After 48 h IL3 withdrawal, these cells shrank in size and arrested

(Fig. 1, B and C). This quiescent state was reversible, because when IL3 was returned,

the cells increased in size and resumed cycling. Thus, in the absence of Bax/Bak1-

mediated apoptosis, FDM cells can survive for weeks without IL3, but they shrink

and arrest.

Bax$^{-/-}$Bak1$^{-/-}$ myeloid cells arrested by deprivation of IL3 do not require Atg5-dependent autophagy to survive.

During macroautophagy, LC3B-I is lipidated by a multiprotein complex containing Atg5 to form LC3B-II, which is incorporated into the growing autophagosome, and is commonly used as a marker of autophagy[9]. Lysates of *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells, whether they were cultured in the presence or absence of IL3, contained similar amounts LC3B-II, suggesting that autophagy is occurring in both cycling and arrested cells (Fig. 2A; lanes 1, 3 and 4). Notably, the low levels of both LC3B isoforms I and II in *Bax$^{-/-}$Bak1$^{-/-}$* cells cultured in serum, amino acid and glucose-free Hank's balanced salt solution (HBSS; Fig. 2A, lane 2) is indicative of a strong increase in autophagic flux, which causes the coincident degradation of LC3B proteins after autophagosomes fuse with lysosomes[9].

To determine if Atg5-dependent autophagy was needed to support the sustained survival of IL3-starved cells, we used CRISPR/Cas9 to delete *Atg5* in *Bax$^{-/-}$Bak1$^{-/-}$* cells, generating *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* triple knockout FDM cell clones. Western blotting confirmed in three *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones the absence of the LC3B-II isoform, whether they were cultured with or without IL3, or in HBSS (Fig. 2A). Moreover, levels of LC3B-I in *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones remained similar across the different culture conditions. These results indicate that Atg5-dependent autophagy and autophagic flux are blocked in the *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* lines. After 14 days cultured without IL3, all *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones remained ~80% viable (Fig. 2B). Similar to *Bax$^{-/-}$Bak1$^{-/-}$* FDMs, *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* cells decreased in size and arrested within 48 h of growth factor withdrawal (Fig. 2, C and D).

These results confirm that IL3 receptor signals are needed for $Bax^{-/-}Bak1^{-/-}$ FDM cells

to maintain their normal size and divide, but show that Atg5-dependent autophagy is

not required for their survival, whether they are cycling or are arrested by IL3

withdrawal. Furthermore, because culturing without IL3 caused a similar decrease in

size of the arrested $Bax^{-/-}Bak1^{-/-}$ to that of the $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells, this shrinkage is

not due to Atg5-dependent autophagy.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells arrested by IL3 deprivation downregulate Glut1

independently of Txnip or Atg5 and survive in the absence of added glucose.

Lum *et al*. reported that when IL3 growth factor was removed, $Bax^{-/-}Bak1^{-/-}$ FDM

cells decreased expression of the glucose transporter Glut1[4]. A possible mechanism

for this was proposed by Wu *et al*., who found that thioredoxin-interacting protein

(Txnip) could not only bind to Glut1 and induce its internalization through clathrin-

coated pits, thus removing it from the plasma membrane, but could also reduce the

level of Glut1 messenger mRNA[7]. Consistent with Lum *et al*. and Wu *et al*., levels of

Txnip increased, and those of Glut1 decreased, in lysates of $Bax^{-/-}Bak1^{-/-}$ cells 48 h

following removal of IL3 (Fig. 3A, lane 2).

To determine whether expression of Txnip caused or correlated with changes in Glut1

levels, we deleted *Txnip* in $Bax^{-/-}Bak1^{-/-}$ FDM cells using CRISPR/Cas9. Western

blotting confirmed the absence of Txnip protein in three $Bax^{-/-}Bak1^{-/-}\Delta Txnip$ clonal

lines (Fig. 3A). We found that Glut1 levels were unaffected by deletion of *Txnip*,

whether IL3 was added or not, indicating that the increase in Txnip was not required

for the decrease in Glut1 when cells were cultured without IL3. Furthermore, similar

to $Bax^{-/-}Bak1^{-/-}$ cells, withdrawal of IL3 caused G_1 arrest in $Bax^{-/-}Bak1^{-/-}\Delta Txnip$ cells

within 48 h (Supplementary Fig. S1A). Because induced over-expression of FLAG-

Txnip in $Bax^{-/-}Bak1^{-/-}$ cells had no impact on levels of Glut1 protein or cell cycle

progression (Figs. 3B and Supplementary S1B), an increase in Txnip is not sufficient

to reduce Glut1. Therefore, in FDM cells, the abundance of Txnip has no effect on

levels of Glut1, whether they are treated with IL3 or not. Of note, IL3 deprivation

caused specific downregulation of Glut1, but not Glut4, protein in $Bax^{-/-}Bak1^{-/-}$ and

$Bax^{-/-}Bak1^{-/-}\Delta Atg5$ clones; and levels of Glut1 were similar among proliferating

Bax/Bak1-null cells of all genotypes (Figs. 3A, B and Supplementary Fig. S1C).

Together, these results indicate that although IL3 can promote the expression of

Glut1, neither Txnip nor Atg5-dependent autophagy is required for the decrease in

Glut1 when growth factor is removed from myeloid cells, and neither absence nor

overexpression of Txnip affects the levels of Glut1.

Because Glut1 levels are reduced in IL3-starved $Bax^{-/-}Bak1^{-/-}$ FDM cells, we asked

whether glucose utilization was reduced. Proliferating $Bax^{-/-}Bak1^{-/-}$ cells in the

presence of IL3 responded strongly to the addition of glucose, as indicated by the

increased extracellular acidification rate (ECAR, Fig. 3C), signifying that they readily

consumed and metabolized the added glucose. Following 3 days cultured without IL3,

the rate of glucose metabolism in $Bax^{-/-}Bak1^{-/-}$ cells was reduced, but they

nevertheless still responded to the addition of glucose. With or without IL3, the rates

of glucose metabolism in $Bax^{-/-}Bak1^{-/-}$ lines that also lacked Txnip or Atg5 were

similar to the rates of the parental Bax/Bak1-null cells (Figs. 3C and Supplementary

S1D). When cells were treated with oligomycin to block oxidative phosphorylation

shortly after they reached their maximal glycolytic rate, the steady decline in ECAR

continued relatively unchanged until glycolysis was inhibited with 2-deoxyglucose (2-

DG), causing ECAR values to fall rapidly to baseline. These results indicate that

FDM cells rely predominantly on glycolysis for energy production, and when

deprived of IL3, they retain the capacity to import and metabolise glucose, albeit at a

reduced level.

Because IL3-starved $Bax^{-/-}Bak1^{-/-}$ FDMs took up and metabolized glucose, we

hypothesised that this was sufficient to support their sustained survival despite the

decrease in Glut1. For FDM cell culture, we used DMEM medium supplemented with

4 mM glutamine, 5.5 mM glucose and 10% fetal bovine serum (FBS), estimated to

contain ~0.5 mM glucose (Supplementary Fig. S1E). To test their requirement for

glucose, the cells were cultured in medium supplemented with 4 mM glutamine and

10% FBS, but lacking the added 5.5 mM glucose. Surprisingly, > 80% of IL3-starved

$Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Txnip$ cells remained viable for 3 weeks in culture

without added glucose (Fig. 3D). Furthermore, when IL3 was present, these cells were

able to proliferate (Supplementary Fig. S1F). These results suggest that FDM cells do

not require high concentrations of glucose to maintain long-term viability, whether

IL3 is present or not, and that they consume the glucose (and other nutrients) present

in FBS and/or the added glutamine to support survival or to multiply when stimulated

with growth factor.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to proliferate but can survive in its

absence for weeks.

When glucose is limiting, lymphoid cells metabolise glutamine to survive and

proliferate[10]. We therefore asked whether IL3-dependent myeloid cells exhibited

similar requirements for glutamine. In medium supplemented with 10% FBS but

without added glucose or glutamine, with or without IL3, >95% of wild-type

$Bax^{+/+}Bak1^{+/+}$ FDM cells died as indicated by uptake of PI by day 4 (Supplementary

Fig. S2A), and consistent with their death being due to apoptosis, the nuclei of the

cells had sub-2n DNA (Fig. 4A). Apoptosis was Bax/Bak1-dependent, because

Bax/Bak1-null cells, including clones also lacking Atg5, did not die when cultured in

medium without added glucose and glutamine, but they arrested instead, even when

IL3 was provided (Fig. 4A). When cultured with neither added glucose nor glutamine,

in the presence or absence of IL3, >80% of the arrested $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}$

$\Delta Atg5$ cells survived for more than 12 days (Figs. 4B and Supplementary S2B).

When $Bax^{-/-}Bak1^{-/-}$ cells were starved of glucose and glutamine, most arrested in G_1

with 2n DNA, but some arrested in G_2/M with 4n DNA (Fig. 4A). When glutamine

was added back to $Bax^{-/-}Bak1^{-/-}$ cells that had been cultured for 7 days in glucose- and

glutamine-free medium, the arrested cells resumed cycling within 48 h in the presence

of IL3, whereas adding back glucose had no effect (Supplementary Fig. S2C). These

results indicate that IL3-treated $Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to

proliferate, but can survive for weeks in its absence. Because $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells

behaved similarly to $Bax^{-/-}Bak1^{-/-}$ cells, Atg5-dependent autophagy is not required for

the sustained survival of growth factor-deprived $Bax^{-/-}Bak1^{-/-}$ FDMs, even in cultures

without high levels of glucose and glutamine.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells arrested by glutamine deprivation shrink in a manner

independent of Atg5-mediated autophagy.

Because $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells arrested by IL3 withdrawal

decreased in size (Figs. 1B and 2C), we asked whether cells arrested by glutamine

withdrawal also shrink. In medium containing 10% FBS but lacking added glucose

and glutamine, IL3-treated $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells showed a gradual

decrease in size to almost 1/3 of their original volume over a 20-day period (Fig. 5).

Without IL3, the shrinkage was more acute and cells reached ~1/3 of their original

volume within 7 days. These results suggest that IL3 provided a signal to $Bax^{-/-}Bak1^{-/-}$

myeloid cells arrested by glutamine deprivation to maintain a large size, and confirm

that the shrinkage of these cells upon IL3 and/or nutrient withdrawal do not require

Atg5-dependent autophagy.

Withdrawal of IL3 or glutamine caused reduced expression of Mcl-1 and key

regulators of cell cycle progression.

To elucidate how growth factor and nutrient deprivation impacted FDM cell survival

and proliferation, we performed Western blotting to determine protein levels of

apoptotic and cell cycle regulators. We found that levels of the pro-survival protein

Mcl-1 were markedly reduced in $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells after 3 days

in culture without added glucose and glutamine (Fig. 6A), suggesting that the

decrease in normal levels of Mcl-1 allowed activation of Bax/Bak1 and the apoptosis

of wild-type FDM cells observed in Fig. 4A. Consistent with this notion, addition of a

specific Mcl-1 inhibitor S63845[11] to $Bax^{+/+}Bak1^{+/+}$ cells was sufficient to cause

significant apoptosis within 24 h, even when the cells were cultured with IL3 in

complete medium containing 10% FBS and high concentrations of glucose and

glutamine (Fig. 6B).

As well as causing arrest, IL3 withdrawal reduced c-Myc protein to barely detectable

levels in both $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells (Fig. 6A). Cdk4 and cyclin D3

are key drivers of cell cycle progression in haematopoietic cell lines[12, 13], and these

proteins were also markedly reduced in the IL3-starved, Bax/Bak1-deficient myeloid

cells, suggesting a mechanism for their arrest (Figs. 1C and 2D). Withdrawal of

glutamine from these cells also caused cell cycle arrest, even in the presence of IL3

(Fig. 4A). Under these conditions, IL3 was able to stimulate the expression of c-Myc

and Cdk4, but the levels of these proteins were significantly lower than in the cells

cultured with both glutamine and IL3 (Fig. 6A). The transcription factor Atf4

accumulates in response to a variety of cellular stresses including amino acid

starvation[14]. In glutamine-starved $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells, Atf4

levels increase strongly, but intriguingly only in the presence of IL3. These results

indicate that the signal transduction pathway activated by ligation of the IL3 receptors

remains largely intact in cells cultured without supplemented glutamine, and can

stimulate the cells to remain large, but that glutamine is essential for high levels of c-

Myc and Cdk4/cyclin D3 expression, and for the cells to divide.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells consume extracellular glucose to maintain long-term

viability.

The Bax/Bak1-null cells cultured without added glucose and glutamine would need a

source of energy to survive (Figs. 4B and S2B), so we tested whether they were

obtaining it from the supplemented FBS. In IL3-containing medium lacking added

glucose, glutamine and FBS, all $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ died after 2 days

(Fig. 7A). In the absence of serum, we asked if adding back glutamine or glucose

alone was sufficient to support cell survival. While <40% of cells cultured in medium

supplemented with 4mM glutamine remained viable after 2 days (and all died within

4 days, Supplementary Fig. S3A), >90% of cells cultured with 5.5 mM glucose

survived (for more than 6 days, Supplementary Fig. S3B). In IL3-starved cultures, we

observed a similar dependence on glucose, but not glutamine, for cell viability (Fig.

7B). Strikingly, we found that 0.5 mM glucose only (an equivalent amount present in

10% FBS) was sufficient to support the survival of >80% of IL3-deprived *Bax*[-/-]*Bak1*[-]

[/-] and *Bax*[-/-]*Bak1*[-/-]Δ*Atg5* cells (Fig. 7B) for at least 1 week (Fig. 7C). Collectively,

these results suggest that in the absence of growth factor, *Bax*[-/-]*Bak1*[-/-] myeloid cells

consume extracellular glucose to maintain long-term viability, without the need to

invoke Atg5-dependent autophagy.

Discussion

IL3-dependent myeloid cell lines have provided many insights into the regulation of

fundamental processes such as cell death, cell division, and autophagy, as well as the

signaling pathways that regulate them. For example, it was experiments on an IL3-

dependent myeloid cell line that first revealed that the function of Bcl-2 was to inhibit

cell death[3]. When genes for Bax and Bak1 are deleted in factor-dependent myeloid

(FDM) cells, they fail to undergo apoptosis when IL3 is removed, but nevertheless

will die if starved of nutrients needed to maintain their metabolism. When *Bax*[-/-]*Bak1*[-]

[/-] FDMs are cultured without IL3, they decrease expression of the glucose transporter

Glut1, shrink and arrest, but do not die[4]. This led to the proposal that the cells could

not take up glucose, so they needed autophagy to maintain their metabolism to

survive. Consistent with this idea, Lum *et al.* found that inhibition of autophagy either

by 3-methyladenine, chloroquine or shRNA knockdown of Atg5 caused the cells to

die[4]. However, we also used chloroquine to inhibit autophagy and shRNA to reduce

Atg5 levels, and found that Atg5-dependent autophagy was not required for the

survival *Bax*[-/-]*Bak1*[-/-] myeloid cells deprived of IL3[15]. It is conceivable that the two

Atg5-targeting shRNA sequences used by Lum *et al.*, which is different to our

shRNA, had off-target effects (such as induction of interferons) that sensitized their

cells to autophagy inhibition. Notwithstanding, it is unlikely that off-target effects of

chloroquine caused such contrasting survival outcomes in similar IL3-dependent

lines.

In this report, we have genetically eliminated *Atg5* expression in IL3-dependent *Bax*[-/-]

Bak1[-/-] cells using CRISPR/Cas9. Unlike Lum *et al.*, who concluded that autophagy

was necessary to sustain the survival of IL3-deprived *Bax*[-/-]*Bak1*[-/-] FDM cells beyond

4 days[4], our results demonstrate that canonical Atg5-dependent autophagy pathways

are not required to support the long-term viability of these cells (Fig. 2). We generated

IL3-dependent FDM lines by co-culturing E14.5 fetal liver single-cell suspensions

from *Bax*[-/-]*Bak1*[-/-] embryos with fibroblasts expressing a *HoxB8* retrovirus (to

immortalize the cells) in the presence of IL3[16]. Lum *et al*. obtained IL3-dependent

myeloid progenitors from the bone marrow of adult *Bax*[-/-]*Bak1*[-/-] mice, but did not

specify how these cells were immortalized[4]. It is plausible that differences in the age,

organ of origin, or immortalization protocols used to generate IL3-dependent myeloid

progenitors caused the discrepancy in results between our study and those reported by

Lum and colleagues. An experiment to rule out these differences would be to mutate

Bax, *Bak1* and *Atg5* genes by CRISPR/Cas9 in IL3-dependent myeloid lines isolated

from the fetal livers and bone marrow of wild-type mice, then determine whether

these cells survive long-term in the absence of IL3.

We were able to confirm the observation that culture of FDM cells without IL3

reduced the levels of the glucose transporter Glut1[4], and wondered if the mechanism

for reduction of Glut1 would be similar to that described by Wu *et al*. who found that

thioredoxin interacting protein (Txnip) could reduce the levels of Glut1 in human

hepatocytes[7]. Although we found that Txnip increased in FDM cells when IL3 was

removed, levels of Glut1 decreased by the same amount when IL3 was removed from

Txnip-null cells, and over-expression of exogenous Txnip did not reduce the levels of

Glut1 (Fig. 3). Although we confirmed the reported inverse correlation between levels

of Txnip and Glut1, Txnip is neither required nor sufficient for IL3 to regulate levels

of Glut1 in FDMs. It remains possible that the Txnip-mediated downregulation of

Glut1 is a mechanism specific to hepatocytes and other cell types in response to

glucose availability[7].

Canonical autophagy was shown to be required for the efficient surface expression of

Glut1[6]. Our Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ FDM cells had similar levels of Glut1 to

parental $Bax^{-/-}Bak1^{-/-}$ cells, and they decreased Glut1 to a similar extent when cultured

without IL3. Furthermore, the $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells showed similar levels of

glucose uptake to $Bax^{-/-}Bak1^{-/-}$ cells when cultured with IL3 and without. Therefore, in

FDM cells, Atg5-dependent autophagy is not needed to maintain normal levels of

Glut1, and does not affect the ability of IL3-starved cells to take up and metabolize

glucose (Figs. 3C and S1D).

Consistent with the requirement of glutamine for lymphoid cell survival[10], we found

that wild-type factor dependent myeloid cells died following glutamine deprivation

(Fig. 4). Surprisingly, a high concentration of glutamine was not required to sustain

the survival of Bax/Bak1-null FDM cells in serum-containing cultures not

supplemented with extra glucose, and that these glutamine-deprived cells maintained

a large size in the presence of IL3, but arrested. This suggests that glutamine provided

the source of amino acids needed for cell division, but that in its absence, these cells

import the glucose present in serum to maintain enough metabolic activity to survive

(Fig. 7). Hecht *et al.*[17] reported that the acute shrinkage of B an T lymphocytes

following growth factor withdrawal occurred before the induction of autophagy.

Consistent with Hecht *et al.*, the shrinkage of FDM cells following growth factor or

glutamine withdrawal also occurred independently of Atg5-mediated autophagy (Fig.

5).

In the absence of high levels of glutamine, pathways activated by IL3 could still

converge to induce the expression of c-Myc in Bax/Bak1-null FDM cells, and

although this was response was blunted, the arrested cells maintained a relatively

large size for at least 1 week (Fig. 5). We found that the transcription factor Atf4 was

strongly induced by IL3 in glutamine-starved cells, but at present, it is unclear how or

why signaling from the IL3 receptor leads to increased Atf4 production. Although

Atf4 was shown to activate the expression of several autophagy genes to promote cell

survival[18], we have not been able to confirm that its accumulation induced LC3B

lipidation in the glutamine-starved FDM cells. That IL3 also induced Atf4 expression

in Atg5-deficient *Bax*[-/-]*Bak1*[-/-] cells following glutamine deprivation (Fig. 6), suggests

that Atf4 does not promote canonical autophagy to maintain cell viability under these

conditions. Moreover, cells starved of both IL3 and glutamine showed no detectable

expression of Atf4, yet survived for weeks.

Several pathways have been proposed to explain how engagement of IL3 receptors

prevents activation of Bax and Bak1. For example, it was proposed that IL3 receptors

signaled via Akt to phosphorylate Bad, preventing it from binding to Bcl-x, to thereby

free Bcl-x to inhibit apoptosis[19, 20]. However, experiments in which genes for Bad or

Akt have been mutated have shown that unlike Bax/Bak1, they play relatively minor

roles in the regulation of apoptosis by IL3[16, 21]. These experiments, and those where

genes for all the Bcl-2 family member were deleted[22] suggest that Bax/Bak1

dependent apoptosis is not always signaled via BH3-only proteins, but more often

regulated by controlling the abundance of anti-apoptotic Bcl-2 family members, such

as Bcl-2, Bcl-x, and Mcl-1[23]. The results presented here are consistent with this, and

show that apoptosis of FDMs following the removal of IL3, or culture in the absence

of glutamine, is due to the decrease in abundance of Mcl-1. Cytokines including IL3

keeps cells alive by driving *Mcl-1* transcription, most likely via Jak2 and Stat5[24, 25, 26],

and glutamine is required to maintain the production of Mcl-1 to prevent default

apoptosis. Consistent with a requirement for Mcl-1 to prevent spontaneous activation

of Bax/Bak1, a specific inhibitor of Mcl-1 was able to induce apoptosis of wild-type

$Bax^{+/+}Bak1^{+/+}$ FDM cells cultured with IL3 and glutamine (Fig. 6).

In the absence of growth factor and high concentrations of added glucose and

glutamine, Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ cells remained viable for weeks in serum-

containing medium, and their rapid death in glucose-free cultures was consistent with

the hypothesis that they took up and used the glucose present in serum to survive (Fig.

7). That these cells predominantly generate chemical energy through glycolysis (Fig.

3) was also consistent with their ability to consume only diluted concentrations of

glucose to remain viable for at least a week. Serum-starved Bax/Bak1-null cells

supplemented with high concentrations of glucose and glutamine survived for several

weeks, but arrested, with or without IL3 (data not shown). Collectively, our work

suggests that in the absence of growth factor, *Bax^{-/-}Bak1^{-/-}* FDM cells, including

mutant lines lacking Atg5-dependent autophagy, import extracellular glucose and

amino acids to maintain long-term viability. Their metabolic requirement for glucose

and amino acids is a mechanism to support survival even in cultures where levels of

these nutrients are not optimal for cell proliferation, and the reduced levels of c-Myc,

Cdk4 and cyclin D3, combined with possibly altered levels of other cell cycle

regulators, could explain why growth factor and/or nutrient-starved *Bax^{-/-}Bak1^{-/-}* cells

arrested (Fig. 8)

Methods & Materials

Cell culture

Wild-type *Bax^{+/+}Bak1^{+/+}* and *Bax^{-/-}Bak1^{-/-}* IL3-dependent mouse myeloid progenitor

(FDM) cell lines have been described[16]. We routinely test our cells for mycoplasma

contamination. Cells were cultured in Dulbecco's Modified Eagle Medium (DMEM,

ThermoFisher Scientific Cat# 11885-084; containing 5.55 mM D-glucose and 4 mM

L-glutamine) supplemented with 10% fetal bovine serum (FBS, Sigma Cat# F9423)

and 0.25 ng/mL recombinant murine interleukin 3 (IL3, PeproTech, Cat# 213-13) in a

37°C humidified incubator with 10% CO_2. Cells were deprived of IL3, glucose,

glucose and glutamine, or varying concentrations of FBS, by washing 3 times with

medium that lacked those nutrients, then resuspended in the same medium for

analysis at the indicated time points. Unless indicated otherwise, 1 x 10^5 cells were

seeded in 1 mL of the specified medium per well of a 24-well tissue culture plate

(Corning, Cat# 353047), and one well was analysed for each timepoint. For

experiments without FBS, plastic tubes and tissue culture dishes were pre-coated with

20 μg/mL bovine serum albumin (Sigma Cat# A1470) and rinsed with serum-free

medium before cell seeding. DMEM lacking D-Glucose and sodium pyruvate but

containing 4 mM L-Glutamine (Cat# 11966-025) and DMEM lacking D-Glucose,

sodium pyruvate, L-Glutamine and phenol red (Cat# A14430-01) were purchased

from ThermoFisher Scientific. D-Glucose was purchased from Sigma (Cat# G7528).

The glucose concentration in FBS was determined using an Accu-Chek Performa

glucose meter, according to the manufacturer's instructions.

Lentiviral constructs for CRISPR/Cas9 and inducible overexpression

An inducible lentiviral CRISPR/Cas9 system[27] was used to mutate *Txnip* in *Bax*[-/-]

Bak1[-/-] FDM cells. Guide RNA oligos targeting mouse *Txnip* (guide 1: CCA GCA

GGT GAG AAC GAG A; guide 2: TAT GGT TGC GTA GAC TAC T) were cloned

into the *Bsm*BI site of the pFgH1tUTG GFP lentiviral vector. Guide RNA oligo

targeting mouse *Atg5* (AAG ATG TGC TTC GAG ATG TG) was cloned into the

*Bsm*BI site of the lentiviral vector lentiCRISPR v2 (Addgene #52961[28]). For

overexpression studies, the coding sequence of mouse *Txnip* was PCR amplified from

pEGFP-C1-Txnip (Addgene #18758[29]) with the incorporation of an N-terminal

FLAG-tag. The PCR product was cloned into the *Eco*RI/*Nhe*I sites of a doxycycline

inducible pFTREtight MCS rtTAadvanced GFP lentiviral vector[21].

Antibodies and small molecules

Antibodies: LC3B (D11, Cell Signaling Technology #3868), Atg5 (D5F5U, Cell

Signaling Technology #12994); β-actin (AC-15, Sigma #A1978), Glut1 (Abcam

ab15730), Txnip (D5F3E, Cell Signaling Technology #14715S), Glut4 (1F8, Cell

Signaling Technology #2213), Mcl-1 (Rockland 600-401-394S), c-Myc (Abcam

ab32072), Cdk4 (Santa Cruz Biotechnology sc-260), Cyclin D3 (Cell Signaling

Technology #2936), Atf4 (D4B8, Cell Signaling Technology #11815). Small

molecules: Mcl-1 inhibitor S63845 was synthesized according to Kotschy *et al.*[11] by

SYNthesis Med Chem Pty Ltd (Parkville, Australia) or by Active Biochem (Catalog#

A-6044 – a kind gift from David Huang); DMSO (dimethyl sulfoxide; Cat# 472301)

and propidium iodide (Cat# P4864) were purchased from Sigma.

Measurements of cell death, cell cycle and cell size

Cell death and cell cycle were assayed using propidium iodide (PI) and flow

cytometry. For death assays, cells were harvested and resuspended in PBS containing

1 μg/mL PI. For cell cycle analysis, cells were harvested and resuspended in PBS

containing 0.1% Triton X-100, 5 μg/mL PI and 10 μg/mL RNaseA (Thermo

Scientific), then incubated at 37 °C for 15 min. Samples were processed on

FACSCalibur or LSRFortessa X-20 flow cytometers (BD Biosciences) and the

acquired data analyzed with FlowJo software. Cell volume in Figure 5 was measured

on a CASY instrument (Roche Innovatis AG), a Coulter-based cell counter operating

on the Electrical Current Exclusion principle. In brief, 200 μL aliquots of cells

cultured under the specified conditions were mixed with 5 mL of CASY electrolyte

buffer and analyzed in a 60 μm CASY capillary. Multiple parameters including the

peak cell volume were collected.

Glucose uptake assays

Extracellular acidification (ECAR) rates were measured in live cells using a Seahorse

Bioscience XFe96 Analyzer (Agilent Technologies). In brief, cells were resuspended

in non-buffered DMEM (Seahorse Biosciences) and 200×10^3 cells were seeded per

well in Seahorse Bioscience 96-well plates. The cellular ECAR was analyzed in non-

buffered DMEM containing 2 mM glutamine with the following compounds: 5 mM

glucose, 1 μM oligomycin and 50 mM 2-deoxy-glucose (2DG). For each assay cycle,

measurement time points of 2 min mix, 2 min wait and 5 min measure were collected.

Data reproducibility

Unless indicated otherwise, all experiments shown have been reproduced at least 3

independent times with similar results. While not all repeats were performed wholly

or in exact detail, we ensured that our experimental design allowed us to interpret and

draw conclusions from multiple independent experiments. This is aided by the fact

that our results are unambiguous: the majority of cells are either alive or dead, cycling

or arrested, large or small; and immunoblots show clear changes in protein levels or

modification. We have endeavored to perform experiments that provide consistent

evidence in mechanistically independent ways, rather than repeating the same

experiment many times until statistical significance is reached. The experimental

triangulation of our evidence[30] is best exemplified by multiple independent assays

confirming that autophagy-deficient FDM cells survived for weeks without growth

factor under a variety of different culture conditions (Figs. 2B, 4B, 5, 7C, S3B).

Acknowledgments

We thank Gabriela Brumatti, Rebecca Feltham, James Vince, Simon Monard, David

Huang and Andre Samson for discussion, technical advice and reagents. Funding for

this project was provided by the Australian National Health and Medical Research

Council (NHMRC) Program Grant #1113133 and NHMRC fellowship 1020136 to

DLV; and the Leukemia and Lymphoma Society SCOR grant #7001-13. Work in the

authors' laboratory is made possible by operational infrastructure grants through the

References

1. Perkins AC, Cory S. Conditional immortalization of mouse myelomonocytic,
megakaryocytic and mast cell progenitors by the Hox-2.4 homeobox gene.
Embo J 1993, **12**(10)**:** 3835-3846.
507

508 2. Ekert PG, Read SH, Silke J, Marsden VS, Kaufmann H, Hawkins CJ*, et al.*
509   Apaf-1 and caspase-9 accelerate apoptosis, but do not determine whether
510   factor-deprived or drug-treated cells die. *J Cell Biol* 2004, **165**(6)**:** 835-842.
511

512 3. Vaux DL, Cory S, Adams JM. Bcl-2 gene promotes haemopoietic cell survival
513   and cooperates with c-myc to immortalize pre-B cells. *Nature* 1988,
514   **335**(6189)**:** 440-442.
515

516 4. Lum JJ, Bauer DE, Kong M, Harris MH, Li C, Lindsten T*, et al.* Growth
517   Factor Regulation of Autophagy and Cell Survival in the Absence of
518   Apoptosis. *Cell* 2005, **120**(2)**:** 237-248.
519

520 5. Mizushima N, Komatsu M. Autophagy: renovation of cells and tissues. *Cell*
521   2011, **147**(4)**:** 728-741. doi: 710.1016/j.cell.2011.1010.1026.
522

523 6. Roy S, Leidal AM, Ye J, Ronen SM, Debnath J. Autophagy-Dependent
524   Shuttling of TBC1D5 Controls Plasma Membrane Translocation of GLUT1
525   and Glucose Uptake. *Mol Cell* 2017, **6**(17)**:** 020.
526

527 7. Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G*, et al.* AMPK-
528   dependent degradation of TXNIP upon energy stress leads to enhanced
529   glucose uptake via GLUT1. *Mol Cell* 2013, **49**(6)**:** 1167-1175. doi:
530   1110.1016/j.molcel.2013.1101.1035. Epub 2013 Feb 1128.
531

532 8. Dexter TM, Garland J, Scott D, Scolnick E, Metcalf D. Growth factor-
533   dependent hemopoietic precursor cell lines. *J Exp Med* 1980, **152:** 1036-1047.
534

535 9. Loos B, du Toit A, Hofmeyr JH. Defining and measuring autophagosome
536   flux-concept and reality. *Autophagy* 2014, **10**(11)**:** 2087-2096.
537

538 10. Le A, Lane AN, Hamaker M, Bose S, Gouw A, Barbi J*, et al.* Glucose-
539   independent glutamine metabolism via TCA cycling for proliferation and
540   survival in B cells. *Cell Metab* 2012, **15**(1)**:** 110-121. doi:
541   110.1016/j.cmet.2011.1012.1009.
542

543 11. Kotschy A, Szlavik Z, Murray J, Davidson J, Maragno AL, Le Toumelin-
544   Braizat G*, et al.* The MCL1 inhibitor S63845 is tolerable and effective in
545   diverse cancer models. *Nature* 2016, **538**(7626)**:** 477-482.

547   12.   Dolezal E, Infantino S, Drepper F, Borsig T, Singh A, Wossning T*, et al.* The
548   BTG2-PRMT1 module limits pre-B cell expansion by regulating the CDK4-
549   Cyclin-D3 complex. *Nat Immunol* 2017, **18**(8)**:** 911-920.

551   13.   Sawai CM, Freund J, Oh P, Ndiaye-Lobry D, Bretz JC, Strikoudis A*, et al.*
552   Therapeutic targeting of the cyclin D3:CDK4/6 complex in T cell leukemia.
553   *Cancer Cell* 2012, **22**(4)**:** 452-465.

555   14.   Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M*, et al.* An
556   integrated stress response regulates amino acid metabolism and resistance to
557   oxidative stress. *Mol Cell* 2003, **11**(3)**:** 619-633.

559   15.   Lindqvist LM, Heinlein M, Huang DC, Vaux DL. Prosurvival Bcl-2 family
560   members affect autophagy only indirectly, by inhibiting Bax and Bak. *Proc*
561   *Natl Acad Sci U S A* 2014, **111**(23)**:** 8512-8517. doi:
562   8510.1073/pnas.1406425111. Epub 1406422014 May 1406425127.

564   16.   Ekert PG, Jabbour AM, Manoharan A, Heraud JE, Yu J, Pakusch M*, et al.*
565   Cell death provoked by loss of interleukin-3 signaling is independent of Bad,
566   Bim, and PI3 kinase, but depends in part on Puma. *Blood* 2006, **108**(5)**:** 1461-
567   1468.

569   17.   Hecht VC, Sullivan LB, Kimmerling RJ, Kim DH, Hosios AM, Stockslager
570   MA*, et al.* Biophysical changes reduce energetic demand in growth factor-
571   deprived lymphocytes. *J Cell Biol* 2016, **212**(4)**:** 439-447.

573   18.   B'Chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi Y*, et al.*
574   The eIF2alpha/ATF4 pathway is essential for stress-induced autophagy gene
575   expression. *Nucleic Acids Res* 2013, **41**(16)**:** 7683-7699.

577   19.   Datta SR, Dudek H, Tao X, Masters S, Fu HA, Gotoh Y*, et al.* Akt
578   phosphorylation of bad couples survival signals to the cell-intrinsic death
579   machinery. *Cell* 1997, **91**(2)**:** 231-241.

581   20.   Zha JP, Harada H, Osipov K, Jockel J, Waksman G, Korsmeyer SJ. Bh3
582   domain of bad is required for heterodimerization with bcl-x-l and pro-
583   apoptotic activity. *Journal of Biological Chemistry* 1997, **272**(39)**:** 24101-
584   24104.

586   21.   Green BD, Jabbour AM, Sandow JJ, Riffkin CD, Masouras D, Daunt CP*, et*
587   *al.* Akt1 is the principal Akt isoform regulating apoptosis in limiting cytokine
588   concentrations. *Cell Death Differ* 2013, **20**(10)**:** 1341-1349. doi:
589   1310.1038/cdd.2013.1363. Epub 2013 Jun 1321.

591   22.   O'Neill KL, Huang K, Zhang J, Chen Y, Luo X. Inactivation of prosurvival
592   Bcl-2 proteins activates Bax/Bak through the outer mitochondrial membrane.
593   *Genes Dev* 2016, **30**(8)**:** 973-988. doi: 910.1101/gad.276725.276115. Epub
594   272016 Apr 276727.

596   23.   Goodall KJ, Finch-Edmondson ML, van Vuuren J, Yeoh GC, Gentle IE, Vince
597   JE, *et al.* Cycloheximide Can Induce Bax/Bak Dependent Myeloid Cell Death
598   Independently of Multiple BH3-Only Proteins. *PLoS One* 2016, **11**(11)**:**
599   e0164003.   doi: 0164010.0161371/journal.pone.0164003.   eCollection
600   0162016.
601
602   24.   Callus BA, Mathey-Prevot B. Interleukin-3-induced activation of the
603   JAK/STAT pathway is prolonged by proteasome inhibitors. *Blood* 1998,
604   **91**(9)**:** 3182-3192.
605
606   25.   Chao JR, Wang JM, Lee SF, Peng HW, Lin YH, Chou CH, *et al.* mcl-1 is an
607   immediate-early gene activated by the granulocyte-macrophage colony-
608   stimulating factor (GM-CSF) signaling pathway and is one component of the
609   GM-CSF viability response. *Mol Cell Biol* 1998, **18**(8)**:** 4883-4898.
610
611   26.   Moulding DA, Quayle JA, Hart CA, Edwards SW. Mcl-1 expression in human
612   neutrophils: regulation by cytokines and correlation with cell survival. *Blood*
613   1998, **92**(7)**:** 2495-2502.
614
615   27.   Aubrey BJ, Kelly GL, Kueh AJ, Brennan MS, O'Connor L, Milla L, *et al.* An
616   inducible lentiviral guide RNA platform enables the identification of tumor-
617   essential genes and tumor-promoting mutations in vivo. *Cell reports* 2015,
618   **10**(8)**:** 1422-1432.
619
620   28.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries
621   for CRISPR screening. *Nat Methods* 2014, **11**(8)**:** 783-784.
622
623   29.   Wang Z, Rong YP, Malone MH, Davis MC, Zhong F, Distelhorst CW.
624   Thioredoxin-interacting protein (txnip) is a glucocorticoid-regulated primary
625   response gene involved in mediating glucocorticoid-induced apoptosis.
626   *Oncogene* 2006, **25**(13)**:** 1903-1913.
627
628   30.   Munafo MR, Davey Smith G. Robust research needs many lines of evidence.
629   *Nature* 2018, **553**(7689)**:** 399-401.

Figure Legends

Figure 1. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells shrink and arrest but remain viable when cultured

without IL3.

A. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells survive long-term in the absence of IL3. *Bax$^{+/+}$Bak1$^{+/+}$*

wild-type (black circles) and *Bax$^{-/-}$Bak1$^{-/-}$* (green squares) FDM cells were cultured

without IL3. At the indicated times, cells were harvested, resuspended in PBS

containing propidium iodide (PI) and analyzed by flow cytometry.

B. IL3 withdrawal caused cells to shrink. PI negative populations of *Bax$^{-/-}$Bak1$^{-/-}$* cells

cultured for 48 h in the presence (light gray) or absence (dark gray) of IL3, and the

latter that had IL3 returned for 72 h (green), were compared by forward scatter as an

indicator of cell size.

C. IL3 withdrawal caused cell cycle arrest, which was reversed by re-addition of IL3.

Nuclei from *Bax$^{-/-}$Bak1$^{-/-}$* cells cultured as described in (B) were stained with PI to

indicate DNA content by flow cytometry.

Figure 2. *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* FDM cells shrink and arrest, but remain viable when IL3

is withdrawn.

A. Whole cell lysates from *Bax$^{-/-}$Bak1$^{-/-}$* cells and 3 *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clonal lines

cultured in the presence of IL3, in Hank's Buffered Saline Solution (HBSS) for 2 h, or

in the absence of IL3 for 2 or 48 h, were subjected to Western blotting. The Atg5

immunoblot demonstrates successful CRISPR/Cas9-mediated deletion of *Atg5*. The

LC3B immunoblot demonstrates lack of LC3B lipidation (isoform II) and autophagic

flux (HBSS treatment) in *ΔAtg5* cell clones.

B. *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* FDM cells survive long-term in the absence of IL3. The

indicated *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clones were cultured without IL3 and viability was

determined by PI exclusion and flow cytometry at the indicated times.

C. *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* cells (clone 1) were cultured for 48 h in the presence or absence

of IL3 and PI negative populations were compared by forward scatter to indicate

differences in cell size.

D. Cells from *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clone 1 were cultured for 48 h in the presence or

absence of IL3 then processed in PI buffer to measure DNA content by flow

cytometry.

Figure 3. *Bax⁻/⁻Bak1⁻/⁻* FDM cells arrested by IL3 deprivation survive in the absence

of added glucose.

A. Txnip is not required for the decrease in Glut1 following IL3 withdrawal. Whole

cell lysates from *Bax⁻/⁻Bak1⁻/⁻* and 3 independent *Bax⁻/⁻Bak1⁻/⁻ΔTxnip* FDM cell clones

cultured for 48 h in the presence or absence of IL3 were analyzed by Western blotting

to detect Glut1 and Txnip protein.

B. Forced expression of Txnip is not sufficient to decrease Glut1 protein levels.

Independent *Bax⁻/⁻Bak1⁻/⁻* FDM clones bearing a doxycycline (dox)-inducible FLAG-

Txnip construct or empty vector control were cultured for 72 h in the absence or

presence of dox. Whole cell lysates were prepared and analyzed by Western blotting

to detect Glut1 and Txnip protein.

C. *Bax⁻/⁻Bak1⁻/⁻* and *Bax⁻/⁻Bak1⁻/⁻ΔTxnip* cells can take up and metabolise glucose.

Cells cultured for 72 h in the presence or absence of IL3 were analyzed on the

Seahorse bioanalyzer to measure the extracellular acidification rate (ECAR) as an

indicator of glucose metabolism. The increased ECAR following addition of glucose

signifies increased glycolytic flux in cells. Oligomycin inhibits mitochondrial ATP

synthase and the change in ECAR upon its addition represents the glycolytic reserve

capacity of cells. Glycolysis is inhibited by 2-deoxyglucose (2-DG) so upon its

addition, the ECAR values are produced from non-glycolytic sources. Equal numbers

of cells were seeded for analysis. To correct for the reduced size of IL3-starved cells,

protein concentration was assayed and the ECAR values were normalized to 1 μg

total protein per sample. Curves represent the mean of 3 replicate samples depicted as

solid symbols for each time point.

D. *Bax*$^{-/-}$*Bak1*$^{-/-}$ and *Bax*$^{-/-}$*Bak1*$^{-/-}$*ΔTxnip* cells deprived of IL3 survive long-term in the

absence of added glucose. Cells were cultured without or with IL3 in medium

supplemented with 10% FBS, 4 mM glutamine and 5.5 mM glucose (medium with

added glucose) or in medium supplemented with 10% FBS and 4 mM glutamine

(medium without added glucose). Cell viability was monitored by PI exclusion and

flow cytometry at the indicated times.

Figure 4. When starved of glutamine, *Bax*$^{+/+}$*Bak1*$^{+/+}$ FDM cells undergo apoptosis,

whereas *Bax*$^{-/-}$*Bak1*$^{-/-}$ cells arrest, but can survive for weeks.

A. *Bax*$^{+/+}$*Bak1*$^{+/+}$ cells undergo apoptosis in the absence of glutamine (upper panels),

whereas *Bax*$^{-/-}$*Bak1*$^{-/-}$ and *Bax*$^{-/-}$*Bak1*$^{-/-}$*ΔAtg5* cells survive in the absence of glutamine,

but arrest even when IL3 is present (lower panels). Cells were cultured in medium

supplemented with 10% FBS, with or without IL3 and 4 mM glutamine, in the

absence of added glucose, for 4 days. Cells were processed in PI buffer to assess

DNA content by flow cytometry. Note the absence of cells in S phase when

Bax/Bak1-null cells were cultured without added glutamine.

B. *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* cells survive for weeks in the absence of glutamine. Three

independent *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones were cultured with or without IL3, in medium

supplemented with 10% FBS, without added glucose or glutamine. At the indicated

times, cells were assayed for viability by PI exclusion and flow cytometry.

Figure 5. Withdrawal of IL3 or glutamine caused cell shrinkage independent of Atg5-

mediated autophagy.

Bax$^{-/-}$Bak1$^{-/-}$ and *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* cells were cultured in medium supplemented with

10% FBS, with or without IL3, in the absence of added glucose and glutamine. At the

indicated times, cell volume was measured in a Coulter-based CASY cell counter.

Figure 6. Withdrawal of IL3 or glutamine reduced the expression of Mcl-1 and

several key regulators of cell cycle progression.

A. *Bax$^{-/-}$Bak1$^{-/-}$* cells and *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clone 1 were cultured for 72 h in medium

supplemented with (+) or without (–) added glucose (Glc) and glutamine (Q), as

indicated, in the presence or absence of IL3 (+/–). Whole cell lysates were prepared

and analyzed by Western blotting to detect the indicated proteins. Three independent

membranes were used for immunoblotting and β-actin loading control blots for each

are shown. Membrane #1: anti-Mcl-1 and anti-c-Myc; membrane #2: anti-Cdk4;

membrane #3: anti-cyclin D3 and anti-Atf4.

B. *Bax$^{+/+}$Bak1$^{+/+}$* and *Bax$^{-/-}$Bak1$^{-/-}$* cells cultured in complete medium with IL3 were

treated with DMSO (vehicle control) or 10 μM of the Mcl-1 inhibitor S63845 for 24

h. Cells were harvested and viability determined by PI exclusion and flow cytometry.

Bars depict the mean of 3 independent experiments represented by the 3 different

symbols, performed in triplicate. The experiment represented by the solid circles was

performed with S63845 synthesized by SYNthesis Med Chem. Experiments

represented by the solid triangles and open hexagons were performed with S63845

synthesized by Active Biochem (see Materials and Methods).

Figure 7. In the absence of Atg5-dependent autophagy, IL3-deprived $Bax^{-/-}Bak1^{-/-}$

myeloid cells consume extracellular glucose to maintain long-term viability.

A. $Bax^{-/-}Bak1^{-/-}$ cells and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ clone 1 were cultured in the absence of

FBS for 48 h in medium without or with the specified concentration of glutamine

and/or glucose, with IL3. Cells were assayed for viability by PI exclusion and flow

cytometry. Bars represent the average of 3 independent experiments (depicted by the

3 different symbols).

B. $Bax^{-/-}Bak1^{-/-}$ cells and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ clone 1 were cultured in the absence of

FBS for 48 h in medium with the specified concentration of glutamine or glucose,

without IL3. Cells were assayed for viability by PI exclusion and flow cytometry.

Bars represent the average of 2 independent experiments (depicted by the 2 different

symbols) performed with technical replicates as indicated.

C. $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells were cultured in medium supplemented

with 0.5 mM glucose; without IL3, glutamine or FBS. At the indicated times, cells

were assayed for viability by PI exclusion and flow cytometry. The curves connect the

average values of data points derived from 2 independent experiments (depicted by

the 2 different symbols) performed with technical replicates or without (dotted

symbols) as indicated.

Figure 8.

The growth factor IL3 stimulates the expression of Mcl-1 that inhibits activation of

Bax/Bak1, and stimulates expression of c-Myc, Cdk4 and cyclin D3 to promote cell

growth and division. Optimal concentrations of glucose and glutamine are required to

maintain levels of these proteins needed to prevent activation of Bax/Bak1 (Mcl-1), as

well as for cell proliferation (c-Myc, Cdk4, cyclin D3 and possibly others). In the

absence of growth factor, or when nutrient levels are low, sufficient levels of glucose

transporters (Glut1, Glut4 and possibly others) remain to allow $Bax^{-/-}Bak1^{-/-}$ myeloid

cells to consume extracellular glucose to survive, even in cells lacking Atg5-

dependent autophagy.

A

B   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻*

C   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻*

A

B

C *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clone 1

D *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clone 1

A

A

B

A   +IL3, 0% FBS, 48 h

B   −IL3, 0% FBS, 48 h

C

Bax$^{-/-}$Bak1$^{-/-}$ IL3-dependent myeloid cells