In the Absence of Apoptosis, Myeloid Cells Arrest When Deprived of Growth Factor,

But Remain Viable Without ATG5-dependent Autophagy or Uptake of Added

Glucose or Glutamine.

Li Dong[1], Boris Reljic[1], Jen G. Cheung[1], Elizabeth S. Ng[2], Lisa M. Lindqvist[1],

Andrew G. Elefanty[2,3], David L. Vaux[1] and Hoanh Tran[1#]

[1]The Walter and Eliza Hall Institute of Medical Research, and the Department of

Medical Biology, University of Melbourne, Victoria, Australia.

[2]Murdoch Childrens Research Institute, The Royal Children's Hospital, Parkville,

Victoria, Australia.

[3]Department of Anatomy and Developmental Biology, Faculty of Medicine, Nursing

and Health Sciences, Monash University, Clayton, Victoria, Australia.

Running Head: Cell Survival Without ATG5-dependent Autophagy

[#]Address correspondence to Hoanh Tran, tran.h@wehi.edu.au

Abstract

Withdrawal of growth factor interleukin 3 from IL3-dependent myeloid cells causes

them to undergo Bax/Bak1-dependent apoptosis, whereas factor-deprived $Bax^{-/-}Bak1^{-/-}$

cells remain viable, but arrest. It was reported that withdrawal of IL3 from $Bax^{-/-}Bak1^{-/-}$

$^{/-}$ cells caused decreased expression of the glucose transporter Glut1, leading to

reduced glucose uptake, so that arrested cells required Atg5-dependent autophagy for

long-term survival. In other cell types, a decrease in Glut1 is mediated by thioredoxin-

interacting protein Txnip, which is induced in IL3-dependent myeloid cells when

growth factor is removed. We mutated *Atg5* and *Txnip* by CRISPR/Cas9 and found

that Atg5-dependent autophagy was not necessary for the long-term viability of

cycling or arrested $Bax^{-/-}Bak1^{-/-}$ cells, and that Txnip was not required for the decrease

in Glut1 expression in response to IL3 withdrawal. Surprisingly, in glucose-free

medium, wild-type and *Bax/Bak1* double mutant cells not only survived, but

continued to cycle when IL3 was present. In contrast, $Bax^{+/+}Bak1^{+/+}$ cells died by

apoptosis when deprived of glutamine, and while $Bax^{-/-}Bak1^{-/-}$ cells deprived of

glutamine survived, they arrested even when IL3 was present. Thus, glutamine, but

neither Atg5-dependent autophagy nor glucose uptake, is necessary for cell division,

and none are needed for arrested $Bax^{-/-}Bak1^{-/-}$ myeloid cells to survive for several

weeks.

Introduction

Mammalian cells import a variety of extracellular nutrients to support their survival and growth. *In vitro*, cells are typically propagated in medium supplemented with glucose, glutamine and fetal bovine serum. To generate chemical energy, glucose can be metabolized by oxidative phosphorylation in the mitochondria or by anaerobic glycolysis. Glutamine can be used as an energy source through glutaminolysis, and can also be used, together with a multitude of factors present in serum, to allow the synthesis of essential macromolecules.

Factor dependent myeloid (FDM) progenitor cell lines (1, 2) require the hematopoietic growth factor interleukin 3 (IL3) to survive and multiply in culture. When IL3 is not provided, FDM cells undergo Bax/Bak1-dependent apoptosis. FDM cells derived from *Bax*[-/-]*Bak1*[-/-] mice survive in the absence of IL3, but they undergo cell cycle arrest, similar to those over-expressing Bcl-2 (3, 4). Previously it was reported that to survive longer than 4 days, IL3-deprived *Bax*[-/-]*Bak1*[-/-] cells required Atg5-dependent autophagy (4), an intracellular catabolic process that can support the survival of starved or arrested cells (5).

Cells import glucose when it is available and growth factors can stimulate the amount of glucose that is consumed. *Bax*[-/-]*Bak1*[-/-] FDM cells arrested by IL3 withdrawal decrease expression of the glucose transporter Glut1, which was proposed to reduce glucose uptake and metabolism (4). Several mechanisms have been put forward to explain the regulation of expression and plasma membrane translocation of Glut1. Activation of autophagy was shown to be required for the efficient surface expression of Glut1 in mouse embryonic fibroblasts cultured under hypoxic conditions (6). In

human hepatocytes subjected to glucose starvation, a mechanism involving

proteasomal degradation of the thioredoxin-interacting protein Txnip was required for

Glut1 stabilization (7). Moreover, it was shown that Txnip directly binds to Glut1 and

promotes its internalization, thus reducing Glut1 on the plasma membrane and the

ability of cells to take up glucose (7).

In this study, we employed $Bax^{-/-}Bak1^{-/-}$ FDM cells to genetically test whether Atg5-

dependent autophagy is required for the sustained survival of arrested cells deprived

of growth factor. We also asked whether autophagy and/or Txnip are required for the

reduction in Glut1 expression and glucose uptake by arrested myeloid cells, and

whether cells required glucose uptake in order to cycle.

Results

$Bax^{-/-}Bak1^{-/-}$ myeloid cells shrink and arrest when deprived of IL3, but survive.

Factor dependent myeloid (FDM) cells require interleukin 3 (IL3) to survive and

proliferate (2, 4, 8). When deprived of IL3 for 4 days, >80% of wild-type

$Bax^{+/+}Bak1^{+/+}$ FDM cells died, as indicated by loss of plasma membrane integrity and

uptake of propidium iodide (Figs. 1A and Supplemental S1A). By contrast, and

consistent with the findings of Lum *et al*. (4), $Bax^{-/-}Bak1^{-/-}$ FDM cells survived in the

absence of IL3 for more than 14 days. After 48 h IL3 withdrawal, these cells shrank in

size and arrested (Fig. 1, B and C). This quiescent state was reversible, because when

IL3 was returned, the cells increased in size and resumed cycling. Thus, in the

absence of Bax/Bak1-mediated apoptosis, FDM cells can survive for weeks without

IL3, but they shrink and arrest.

Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ myeloid cells arrested by deprivation of IL3 do not require Atg5-

dependent autophagy to survive.

During macroautophagy, LC3B-I is lipidated by a multiprotein complex containing

Atg5 to form LC3B-II, which can incorporate into the growing autophagosome, and is

a common marker of autophagy (9). Lysates of *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* FDM cells, whether they

were cultured in the presence or absence of IL3, contained similar amounts LC3B-II,

suggesting that autophagy is occurring in both cycling and arrested cells (Fig. 2A;

lanes 1, 3 and 4). Notably, the degradation of both LC3B isoforms I and II in *Bax⁻ᐟ⁻*

Bak1⁻ᐟ⁻ cells under starvation in HBSS (Fig. 2A, lane 2) is indicative of a strong

increase in autophagic flux, which causes the coincident turnover of LC3B proteins

when autophagosomes fuse with lysosomes (9).

To determine if Atg5-dependent autophagy was needed to support the sustained

survival of IL3-starved cells, we used CRISPR/Cas9 to delete *Atg5* in *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻*

cells, generating *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* triple knockout FDM cell clones. Western

blotting confirmed in three *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clones the absence of the LC3B-II

isoform, whether they were cultured with or without IL3, or with HBSS (Fig. 2A).

Moreover, levels of LC3B-I in *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clones remained similar across the

different culture conditions. These results indicate that Atg5-dependent autophagy and

autophagic flux are blocked in the *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* lines. After 14 days in culture

without IL3, all *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clones remained ~80% viable (Fig. 2B). Similar to

Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ FDMs, *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* cells decreased in size and arrested within 48 h

of growth factor withdrawal (Fig. 2, C and D).

These results confirm that IL3 receptor signals are needed for *Bax^-/-^Bak1^-/-* FDM cells

to maintain their normal size, but show that Atg5-dependent autophagy is not required

for their survival, whether they are cycling or are arrested by IL3 withdrawal.

Furthermore, because culturing without IL3 caused a similar decrease in size of the

arrested *Bax^-/-^Bak1^-/-* as of the *Bax^-/-^Bak1^-/-ΔAtg5* cells, this shrinkage is not due to

Atg5-dependent autophagy.

Bax^-/-^Bak1^-/- myeloid cells arrested by IL3 deprivation survive in the absence of

glucose.

Lum *et al*. reported that when growth factor was removed, *Bax^-/-^Bak1^-/-* FDM cells

decreased expression of the glucose transporter Glut1 (4). A possible mechanism for

this was proposed by Wu *et al*., who found that thioredoxin-interacting protein

(Txnip) could not only bind to Glut1 and induce its internalization through clathrin-

coated pits, thus removing it from the plasma membrane, but could also reduce the

level of Glut1 messenger mRNA (7). Consistent with Lum *et al*. and Wu *et al*., Txnip

was increased, and Glut1 was decreased, in lysates of *Bax^-/-^Bak1^-/-* cells 48 h

following removal of IL3 (Fig. 3A, lane 2).

To determine whether expression of Txnip caused or correlated with changes in Glut1

levels, we deleted *Txnip* in *Bax^-/-^Bak1^-/-* FDM cells using CRISPR/Cas9. Western

blotting confirmed the absence of Txnip protein in three *Bax^-/-^Bak1^-/-ΔTxnip* clonal

lines (Fig. 3A). We found that Glut1 levels were unaffected by deletion of *Txnip*,

whether IL3 was added or not, indicating that the increase in Txnip was not required

for the decrease in Glut1 when cells were cultured without IL3. Furthermore,

withdrawal of IL3 caused G_1 arrest in *Bax^-/-^Bak1^-/-ΔTxnip* cells within 48 h, similar to

Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ cells (Figs. 1C and Supplemental S1A). Because induced over-

expression of FLAG-Txnip in *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* cells had no impact on levels of Glut1

protein or cell cycle progression (Figs. 3B and Supplemental S1B), an increase in

Txnip is not sufficient to reduce Glut1. Therefore, in FDM cells, the abundance of

Txnip has no effect on levels of Glut1, whether they are treated with IL3 or not.

Levels of Glut1 were similar among proliferating Bax/Bak1-null cells of all genotypes

and, compared to *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* cells, the decrease in Glut1 expression following IL3

withdrawal in *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clones was unaffected (Supplemental Fig. S1C). Of

note, IL3 deprivation caused specific downregulation of Glut1, but not Glut4, protein

in FDMs (Supplemental Fig. S1C).

Together, these results indicate that although IL3 can promote the expression of

Glut1, neither Txnip nor Atg5-dependent autophagy is required for the decrease in

Glut1 when growth factor is removed from myeloid cells, and neither absence nor

overexpression of Txnip affects the levels of Glut1.

Because Glut1 levels are reduced in IL3-starved *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* FDM cells, we asked

whether glucose utilization was reduced. The rate of glucose metabolism in *Bax⁻ᐟ⁻*

Bak1⁻ᐟ⁻ FDMs was reduced after IL3 withdrawal, but they nevertheless still responded

to the addition of glucose, as indicated by the increased extracellular acidification rate

(ECAR, Fig. 3C). In the absence of IL3, *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* FDM lines that also lacked

Txnip or Atg5 had similar rates of glucose metabolism to the parental Bax/Bak1-null

cells (Figs. 3C and Supplemental S1D). These results suggest that arrested FDM cells

retain the capacity to import and metabolise glucose, albeit at a reduced level.

Because the arrested IL3-starved cells were able to take up glucose, we hypothesised

that this was sufficient to support their sustained survival despite the decrease in

Glut1. To test their requirement for glucose, the cells were cultured in medium that

lacked glucose. Surprisingly, $\geq$ 80% of IL3-starved $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}$

$\Delta Txnip$ cells remained viable at 24 days without glucose (Fig. 3D). Furthermore,

when IL3 was present, these cells were able to proliferate (Supplemental Fig. S1E).

These results indicate that FDM cells will import glucose whether IL3 is present or

not, and that they can still proliferate even in the absence of added glucose.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to proliferate but can survive in its

absence for weeks.

When glucose is limiting, lymphoid cells metabolise glutamine to survive and

proliferate (10). We therefore asked whether FDM cells exhibited similar

requirements for glutamine when cultured in the absence of glucose. In medium that

lacked both glucose and glutamine, with or without IL3, >95% of wild-type

$Bax^{+/+}Bak1^{+/+}$ FDM cells died as indicated by uptake of PI by day 4 (Supplemental

Fig. S2A), and consistent with their death being due to apoptosis, the nuclei of the

cells had sub-2n DNA (Fig. 4A). Apoptosis was Bax/Bak1-dependent, because

Bax/Bak1-null cells did not die when cultured in medium lacking glucose and

glutamine, but they arrested instead, even when IL3 was provided (Figs. 4A and

Supplemental S2B). When cultured with neither glucose nor glutamine, in the

presence or absence of IL3, >80% of $Bax^{-/-}Bak1^{-/-}$ cells survived for 16 days and

beyond (Fig. 4B). Similar findings were observed for $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells

(Supplemental Fig. S2C).

When $Bax^{-/-}Bak1^{-/-}$ FDM cells were starved of glucose and glutamine, most arrested in

G_1 with 2n DNA, but some arrested in G_2/M with 4n DNA (Fig. 4, A and C). When

glutamine was added back to $Bax^{-/-}Bak1^{-/-}$ cells that had been cultured for 7 days

without glucose or glutamine, the arrested cells resumed cycling within 48 h in the

presence of IL3, whereas adding back glucose had no effect (Fig. 4C). These results

indicate that IL3-treated $Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to proliferate,

but can survive up to 4 weeks or more in its absence. Because $Bax^{-/-}Bak1^{-/-}\Delta Atg5$

FDMs behaved similarly to $Bax^{-/-}Bak1^{-/-}$ cells, Atg5-dependent autophagy is not

required for sustained cell survival, even in the absence of glucose and glutamine.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells arrested by glutamine withdrawal shrink in a manner

independent of Atg5-mediated autophagy.

Because $Bax^{-/-}Bak1^{-/-}$ myeloid cells arrested by IL3 deprivation decreased in size (Fig.

1B), we asked whether cells arrested by glutamine withdrawal also shrink. When

cultured in medium lacking glucose and glutamine, but with added IL3, $Bax^{-/-}Bak1^{-/-}$

and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells showed a gradual decrease in size to almost 1/3 of their

original volume over a 20-day period (Fig. 5, A and B). Without IL3, the shrinkage

was more acute and cells reached ~1/3 of their original volume within 7 days. These

results suggest that IL3 provided a signal to cells arrested by glutamine deprivation to

maintain a large size. They also confirm that the shrinkage of $Bax^{-/-}Bak1^{-/-}$ myeloid

cells upon IL3 and/or glutamine withdrawal do not require Atg5-dependent

autophagy.

 Withdrawal of IL3 or glutamine caused reduced expression of Mcl-1 and key

regulators of cell cycle progression.

To elucidate how growth factor and nutrient deprivation impacted FDM cell survival

and proliferation, we performed Western blotting to determine protein levels of

apoptotic and cell cycle regulators. We found that levels of the pro-survival protein

Mcl-1 were markedly reduced in *Bax^{-/-}Bak1^{-/-}* and *Bax^{-/-}Bak1^{-/-}ΔAtg5* cells after 3 days

in culture without glucose and glutamine (Fig. 6A), suggesting that the decrease in

normal levels of Mcl-1 allowed activation of Bax/Bak1 and the apoptosis of wild-type

FDM cells observed in Fig. 4A. Consistent with this notion, addition of a specific

Mcl-1 inhibitor S63845 (11) to *Bax^{+/+}Bak1^{+/+}* cells was sufficient to cause significant

apoptosis within 24 h, even when the cells were cultured in complete medium

containing glutamine and IL3 (Fig. 6B).

As well as causing arrest, IL3 withdrawal reduced c-Myc protein to barely detectable

levels in both *Bax^{-/-}Bak1^{-/-}* and *Bax^{-/-}Bak1^{-/-}ΔAtg5* cells (Fig. 6A). Cdk4 and cyclin D3

are key drivers of cell cycle progression in haematopoietic cell lines (12, 13), and

these proteins were also markedly reduced in the IL3-starved, Bax/Bak1-deficient

myeloid cells, suggesting a mechanism for their arrest (Figs. 1C and 2D). Withdrawal

of glutamine from these cells also caused cell cycle arrest, even in the presence of IL3

(Figs. 4A and Supplemental S2B). Under these conditions, IL3 was able to stimulate

the expression of c-Myc and Cdk4, but the levels of these proteins were significantly

lower than in the cells cultured with both glutamine and IL3 (Fig. 6A). The

transcription factor Atf4 accumulates in response to a variety of cellular stresses

including amino acid starvation (14). In glutamine-starved *Bax^{-/-}Bak1^{-/-}* and *Bax^{-/-}*

Bak1^{-/-}ΔAtg5 cells, Atf4 levels increase strongly, but intriguingly only in the presence

of IL3. These results indicate that the signal transduction pathway activated by

ligation of the IL3 receptors remains largely intact in cells cultured without

supplemented glutamine, and can stimulate the cells to remain large, but that

glutamine is essential for high levels of c-Myc and Cdk4/cyclin D3 expression, and

for the cells to divide.

Discussion

IL3-dependent myeloid cell lines have provided many insights into the regulation of

fundamental processes such as cell death, cell division, and autophagy, as well as the

signaling pathways that regulate them. For example, it was experiments on an IL3-

dependent myeloid cell line that first revealed that the function of Bcl-2 was to inhibit

cell death (3). When genes for Bax and Bak1 are deleted in factor-dependent myeloid

(FDM) cells, they fail to undergo apoptosis when IL3 is removed, but nevertheless

will die if starved of nutrients needed to maintain their metabolism. When $Bax^{-/-}Bak1^{-/-}$

FDMs are cultured without IL3, they decrease expression of the glucose transporter

Glut1, shrink and arrest, but do not die (4). This led to the proposal that the cells could

not take up glucose, so they needed autophagy to maintain their metabolism to

survive. Consistent with this idea, Lum *et al*. found that inhibition of autophagy either

by 3-methyladenine or shRNA knockdown of Atg5 caused the cells to die (4).

However, we also used shRNA to reduce Atg5 levels and found that Atg5-dependent

autophagy was not required for the survival $Bax^{-/-}Bak1^{-/-}$ myeloid cells deprived of

IL3 (16). In this report, we have genetically eliminated *Atg5* expression in IL3-

dependent $Bax^{-/-}Bak1^{-/-}$ cells using CRISPR/Cas9. Unlike Lum *et al*., who concluded

that autophagy was necessary to sustain the survival of IL3-deprived $Bax^{-/-}Bak1^{-/-}$

FDM cells beyond 4 days (4), our results demonstrate that canonical Atg5-dependent

autophagy pathways are not required to support the long-term viability of these cells

(Fig. 2).

We were able to confirm the observation that culture of FDM cells without IL3

reduced the levels of the glucose transporter Glut1 (4), and wondered if the

mechanism for reduction of Glut1 would be similar to that described by Wu *et al.* who

found that thioredoxin interacting protein (Txnip) could reduce the levels of Glut1 in

human hepatocytes (7). Although we found that Txnip increased in FDM cells when

IL3 was removed, levels of Glut1 decreased by the same amount when IL3 was

removed from Txnip-null cells, and over-expression of exogenous Txnip did not

reduce the levels of Glut1 (Fig. 3). Although we confirmed the reported inverse

correlation between levels of Txnip and Glut1, Txnip is neither required nor sufficient

for IL3 to regulate levels of Glut1 in FDMs. It remains possible that the Txnip-

mediated downregulation of Glut1 is a process specific to human hepatocytes (7).

Canonical autophagy was shown to be required for the efficient surface expression of

Glut1 (6). Our Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ FDM cells had similar levels of Glut1 to

parental $Bax^{-/-}Bak1^{-/-}$ cells, and they decreased Glut1 to a similar extent when cultured

without IL3. Furthermore, the $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells showed similar levels of

glucose uptake to $Bax^{-/-}Bak1^{-/-}$ cells when cultured with IL3 and without. Therefore, in

FDM cells, Atg5-dependent autophagy is not needed to maintain normal levels of

Glut1, and does not affect the ability of cells to take up and metabolize glucose.

We were surprised to find that Bax/Bak1-null FDM cells of all genotypes, when

cultured in the absence of glucose and arrested due to withdrawal of IL3, not only

survived, but that these cells, and even wild-type $Bax^{+/+}Bak1^{+/+}$ cells (data not shown),

proliferated in the absence of glucose when IL3 was present. Therefore, even though

FDM cells will take up and metabolise glucose when it is available, even if they are

arrested and their Glut1 levels are decreased, they have no obligatory requirement for

glucose, either to survive or to proliferate. Our data demonstrates that reduced uptake

of glucose is not responsible for the arrest of $Bax^{-/-}Bak1^{-/-}$ FDM cells upon IL3

withdrawal.

Consistent with the requirement of glutamine for lymphoid cell survival (10), we

found that wild-type factor dependent myeloid cells died following glutamine

deprivation (Fig. 4). It was surprising that glutamine was not required to sustain the

survival of Bax/Bak1-null FDM cells in the absence of glucose or Atg5-dependent

autophagy, and that these glutamine-deprived cells maintained a large size in the

presence of IL3, but arrested. This suggests that glutamine provided the source of

amino acids needed for cell division, but that these cells can maintain enough

metabolic activity to survive even in its absence. Hecht *et al*. (17) reported that the

acute shrinkage of B an T lymphocytes following growth factor withdrawal occurred

before the induction of autophagy. Consistent with Hecht *et al*., the shrinkage of FDM

cells following growth factor or glutamine withdrawal also occurred independently of

Atg5-mediated autophagy (Fig. 5).

In the absence of glutamine, pathways activated by IL3 could still converge to induce

the expression of c-Myc in Bax/Bak1-null FDM cells, and although this was response

was blunted, the arrested cells maintained a relatively large size for at least 1 week

(Fig. 5). We found that the transcription factor Atf4 was strongly induced by IL3 in

glutamine-starved cells, but at present, it is unclear how or why signaling from the

IL3 receptor leads to increased Atf4 production. Given that IL3 can stimulate the

expression of ER chaperones in FDM cells (18) and that Atf4 translation increased in

cells starved of amino acids or exposed to stimuli that induced ER stress (14), it is

plausible that IL3 activated the unfolded protein response in the glutamine-deprived

FDMs. Atf4 was also shown to activate the expression of several autophagy genes to

promote cell survival (19), but we have not been able to confirm that its accumulation

induced LC3B lipidation in the glutamine-starved FDM cells. That IL3 also induced

Atf4 expression in Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ cells cultured in glutamine-free media

(Fig. 6), suggests that Atf4 does not promote canonical autophagy to maintain cell

viability under these conditions. Moreover, cells starved of both IL3 and glutamine

showed no detectable expression of Atf4, yet survived for weeks. It would be

interesting to determine if Atf4 plays a role in maintaining the relatively large size of

glutamine-starved $Bax^{-/-}Bak1^{-/-}$ cells cultured with IL3.

Several pathways have been proposed to explain how engagement of IL3 receptors

prevents activation of Bax and Bak1. For example, it was proposed that IL3 receptors

signaled via Akt to phosphorylate Bad, preventing it from binding to Bcl-x, to thereby

free Bcl-x to inhibit apoptosis (20, 21). However, experiments in which genes for Bad

or Akt have been mutated have shown that unlike Bax/Bak1, they play relatively

minor roles in the regulation of apoptosis by IL3 (22, 23). These experiments, and

those where genes for all the Bcl-2 family member were deleted (24) suggest that

Bax/Bak1 dependent apoptosis is not always signaled via BH3-only proteins, but

more often regulated by controlling the abundance of anti-apoptotic Bcl-2 family

members, such as Bcl-2, Bcl-x, and Mcl-1 (25). The results presented here are

consistent with this, and show that apoptosis of FDMs following the removal of IL3,

or culture in the absence of glutamine, is due to the decrease in abundance of Mcl-1.

Cytokines including IL3 keeps cells alive by driving *Mcl-1* transcription, most likely

via Jak2 and Stat5 (26-28), and glutamine is required to maintain the production of

Mcl-1 to prevent default apoptosis. Consistent with a requirement for Mcl-1 to

prevent spontaneous activation of Bax/Bak1, a specific inhibitor of Mcl-1 was able to

induce apoptosis of wild-type $Bax^{+/+}Bak1^{+/+}$ FDM cells cultured with IL3 and

glutamine (Fig. 6).

In the absence of growth factor, glucose, glutamine and Atg5-dependent autophagy,

$Bax^{-/-}Bak1^{-/-}$ FDM cells remained viable for weeks, and their sensitivity to low

concentrations of fetal bovine serum (Supplemental Fig. S3) are consistent with the

hypothesis that they took up and used components present in serum to maintain

enough metabolism in order to survive. Serum factors are likely scavenged by

endocytic processes such as macropinocytosis, which can be stimulated by growth

factors (29). Growth factor-induced endocytic uptake of nutrients could account for

the large size of glutamine-starved cells cultured with IL3. Moreover, the reduced

levels of Cdk4 and cyclin D3, combined with possibly altered levels of other cell

cycle regulators, could explain why glutamine-starved cells arrested, even in the

presence of IL3 that stimulated them to maintain a large size (Fig. 7).

Materials & Methods

Cell culture

Wild-type $Bax^{+/+}Bak1^{+/+}$ and $Bax^{-/-}Bak1^{-/-}$ IL3-dependent mouse myeloid progenitor

(FDM) cell lines have been described (23). Cells were cultured in Dulbecco's

Modified Eagle Medium (DMEM, ThermoFisher Scientific Cat# 11885-084; containing 5.55 mM D-glucose and 4 mM L-glutamine) supplemented with 10% fetal bovine serum (FBS, ThermoFisher Scientific Cat# 16000-044) and 0.25 ng/mL recombinant murine interleukin 3 (IL3, PeproTech, Cat# 213-13) in a 37ºC humidified incubator with 10% CO_2. Cells were deprived of IL3, glucose, glucose and glutamine, or varying concentrations of FBS, by washing 3 times with medium that lacked those nutrients, then resuspended in the same medium for analysis at the indicated time points. Unless indicated otherwise, 1 x 10^5 cells were seeded in 1 mL of the specified medium per well of a 24-well tissue culture plate (Corning, Cat# 353047) for analysis per timepoint. DMEM lacking D-Glucose and sodium pyruvate but containing 4 mM L-Glutamine (Cat# 11966-025) and DMEM lacking D-Glucose, sodium pyruvate, L-Glutamine and phenol red (Cat# A14430-01) were purchased from ThermoFisher Scientific. D-Glucose was purchased from Sigma (Cat# G7528).

Lentiviral constructs for CRISPR/Cas9 and inducible overexpression

An inducible lentiviral CRISPR/Cas9 system (30) was used to mutate *Txnip* in *Bax*$^{-/-}$ *Bak1*$^{-/-}$ FDM cells. Guide RNA oligos targeting mouse *Txnip* (guide 1: CCA GCA GGT GAG AAC GAG A; guide 2: TAT GGT TGC GTA GAC TAC T) were cloned into the *Bsm*BI site of the pFgH1tUTG GFP lentiviral vector. Guide RNA oligo targeting mouse *Atg5* (AAG ATG TGC TTC GAG ATG TG) was cloned into the *Bsm*BI site of the lentiviral vector lentiCRISPR v2 (Addgene #52961; (31)). For overexpression studies, the coding sequence of mouse *Txnip* was PCR amplified from pEGFP-C1-Txnip (Addgene #18758; (32)) with the incorporation of an N-terminal FLAG-tag. The PCR product was cloned into the *Eco*RI/*Nhe*I sites of a doxycycline inducible pFTREtight MCS rtTAadvanced GFP lentiviral vector (33).

Antibodies and small molecules

Antibodies: LC3B (D11, Cell Signaling Technology #3868), Atg5 (D5F5U, Cell Signaling Technology #12994); β-actin (AC-15, Sigma #A1978), Glut1 (Abcam ab15730), Txnip (D5F3E, Cell Signaling Technology #14715S), Glut4 (1F8, Cell Signaling Technology #2213), Mcl-1 (Rockland 600-401-394S), c-Myc (Abcam ab32072), Cdk4 (Santa Cruz Biotechnology sc-260), Cyclin D3 (Cell Signaling Technology #2936), Atf4 (D4B8, Cell Signaling Technology #11815). Small molecules: Mcl-1 inhibitor S63845 was synthesized according to Kotschy et al. (11) by SYNthesis Med Chem Pty Ltd (Parkville, Australia) or by Active Biochem (Catalog# A-6044 – a kind gift from David Huang); DMSO (dimethyl sulfoxide; Cat# 472301) and propidium iodide (Cat# P4864) were purchased from Sigma.

Measurements of cell death, cell cycle and cell size

Cell death and cell cycle were assayed using propidium iodide (PI) and flow cytometry. For death assays, cells were harvested and resuspended in PBS containing 1 μg/mL PI. For cell cycle analysis, cells were harvested and resuspended in PBS containing 0.1% Triton X-100, 5 μg/mL PI and 10 μg/mL RNaseA (Thermo Scientific), then incubated at 37 °C for 15 min. Samples were processed on FACSCalibur or LSRFortessa X-20 flow cytometers (BD Biosciences) and the acquired data analyzed with FlowJo software. Cell volume in Figure 5 was measured on a CASY instrument (Roche Innovatis AG), a Coulter-based cell counter operating on the Electrical Current Exclusion principle. In brief, 200 μL aliquots of cells cultured under the specified conditions were mixed with 5 mL of CASY electrolyte

buffer and analyzed in a 60 μm CASY capillary. Multiple parameters including the

peak cell volume were collected.

Glucose uptake assays

Extracellular acidification (ECAR) rates were measured in live cells using a Seahorse

Bioscience XFe96 Analyzer (Agilent Technologies). In brief, cells were resuspended

in non-buffered DMEM (Seahorse Biosciences) and 200 x 10^3 cells were seeded per

well in Seahorse Bioscience 96-well plates. The cellular ECAR was analyzed in non-

buffered DMEM containing 2 mM glutamine with the following compounds: 5 mM

glucose, 1 μM oligomycin and 50 mM 2-deoxy-glucose (2DG). For each assay cycle,

measurement time points of 2 min mix, 2 min wait and 5 min measure were collected.

Data reproducibility

All experiments shown have been reproduced at least 3 independent times with

similar results. While not all repeats were performed wholly or in exact detail, we

ensured that the experimental design allowed us to interpret and draw conclusions

from multiple independent experiments. This is aided by the fact that our results are

unambiguous: the majority of cells are either alive or dead, cycling or arrested, large

or small; and immunoblots show clear changes in protein levels or modification. We

have endeavored to perform experiments that provide consistent evidence in

mechanistically independent ways, rather than repeating the same experiment many

times until statistical significance is reached.

Acknowledgments

Australian National Health and Medical Research Council (NHMRC) Program Grant

#1113133 and NHMRC fellowship 1020136 to DLV; the Leukemia and Lymphoma

Society SCOR grant #7001-13. Work in the authors' laboratory is made possible by

operational infrastructure grants through the Australian Government Independent

Research Institutes Infrastructure Support (IRISS) and the Victorian State

Government OIS. We thank Drs. Gabriela Brumatti, Rebecca Feltham, James Vince

and Andre Samson for discussion, technical advice and reagents.

References

1. Perkins AC, Cory S. 1993. Conditional immortalization of mouse

myelomonocytic, megakaryocytic and mast cell progenitors by the Hox-2.4

homeobox gene. EMBO Journal 12:3835-46.

2. Ekert PG, Read SH, Silke J, Marsden VS, Kaufmann H, Hawkins CJ, Gerl R,

Kumar S, Vaux DL. 2004. Apaf-1 and caspase-9 accelerate apoptosis, but do not

determine whether factor-deprived or drug-treated cells die. J Cell Biol 165:835-42.

3. Vaux DL, Cory S, Adams JM. 1988. Bcl-2 gene promotes haemopoietic cell

survival and cooperates with c-myc to immortalize pre-B cells. Nature 335:440-442.

4. Lum JJ, Bauer DE, Kong M, Harris MH, Li C, Lindsten T, Thompson CB. 2005.

Growth Factor Regulation of Autophagy and Cell Survival in the Absence of

Apoptosis. Cell 120:237-248.

5. Mizushima N, Komatsu M. 2011. Autophagy: renovation of cells and tissues. Cell

147:728-41. doi: 10.1016/j.cell.2011.10.026.

6. Roy S, Leidal AM, Ye J, Ronen SM, Debnath J. 2017. Autophagy-Dependent

Shuttling of TBC1D5 Controls Plasma Membrane Translocation of GLUT1 and

Glucose Uptake. Mol Cell 6:020.

7. Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, Shen CH, Wen J,

Asara J, McGraw TE, Kahn BB, Cantley LC. 2013. AMPK-dependent degradation of

TXNIP upon energy stress leads to enhanced glucose uptake via GLUT1. Mol Cell

49:1167-75. doi: 10.1016/j.molcel.2013.01.035. Epub 2013 Feb 28.

8. Dexter TM, Garland J, Scott D, Scolnick E, Metcalf D. 1980. Growth factor-

dependent hemopoietic precursor cell lines. J Exp Med 152:1036-1047.

9. Loos B, du Toit A, Hofmeyr JH. 2014. Defining and measuring autophagosome

flux-concept and reality. Autophagy 10:2087-96.

10. Le A, Lane AN, Hamaker M, Bose S, Gouw A, Barbi J, Tsukamoto T, Rojas

CJ, Slusher BS, Zhang H, Zimmerman LJ, Liebler DC, Slebos RJ, Lorkiewicz PK,

Higashi RM, Fan TW, Dang CV. 2012. Glucose-independent glutamine metabolism

via TCA cycling for proliferation and survival in B cells. Cell Metab 15:110-21. doi:

10.1016/j.cmet.2011.12.009.

11. Kotschy A, Szlavik Z, Murray J, Davidson J, Maragno AL, Le Toumelin-

Braizat G, Chanrion M, Kelly GL, Gong JN, Moujalled DM, Bruno A, Csekei M,

Paczal A, Szabo ZB, Sipos S, Radics G, Proszenyak A, Balint B, Ondi L, Blasko G,

Robertson A, Surgenor A, Dokurno P, Chen I, Matassova N, Smith J, Pedder C,

Graham C, Studeny A, Lysiak-Auvity G, Girard AM, Grave F, Segal D, Riffkin CD,

Pomilio G, Galbraith LC, Aubrey BJ, Brennan MS, Herold MJ, Chang C, Guasconi

G, Cauquil N, Melchiore F, Guigal-Stephan N, Lockhart B, Colland F, Hickman JA,

Roberts AW, Huang DC, Wei AH, et al. 2016. The MCL1 inhibitor S63845 is

tolerable and effective in diverse cancer models. Nature 538:477-482.

12. Dolezal E, Infantino S, Drepper F, Borsig T, Singh A, Wossning T, Fiala GJ,

Minguet S, Warscheid B, Tarlinton DM, Jumaa H, Medgyesi D, Reth M. 2017. The

BTG2-PRMT1 module limits pre-B cell expansion by regulating the CDK4-Cyclin-

D3 complex. Nat Immunol 18:911-920.

13.   Sawai CM, Freund J, Oh P, Ndiaye-Lobry D, Bretz JC, Strikoudis A, Genesca

L, Trimarchi T, Kelliher MA, Clark M, Soulier J, Chen-Kiang S, Aifantis I. 2012.

Therapeutic targeting of the cyclin D3:CDK4/6 complex in T cell leukemia. Cancer

Cell 22:452-65.

14.   Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C,

Popko B, Paules R, Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. 2003. An

integrated stress response regulates amino acid metabolism and resistance to oxidative

stress. Mol Cell 11:619-33.

15.   Ng ES, Davis R, Stanley EG, Elefanty AG. 2008. A protocol describing the use

of a recombinant protein-based, animal product-free medium (APEL) for human

embryonic stem cell differentiation as spin embryoid bodies. Nat Protoc 3:768-76.

16.   Lindqvist LM, Heinlein M, Huang DC, Vaux DL. 2014. Prosurvival Bcl-2

family members affect autophagy only indirectly, by inhibiting Bax and Bak. Proc

Natl Acad Sci U S A 111:8512-7. doi: 10.1073/pnas.1406425111. Epub 2014 May 27.

17.   Hecht VC, Sullivan LB, Kimmerling RJ, Kim DH, Hosios AM, Stockslager

MA, Stevens MM, Kang JH, Wirtz D, Vander Heiden MG, Manalis SR. 2016.

Biophysical changes reduce energetic demand in growth factor-deprived

lymphocytes. J Cell Biol 212:439-47.

18.   Brewer JW, Cleveland JL, Hendershot LM. 1997. A pathway distinct from the

mammalian unfolded protein response regulates expression of endoplasmic reticulum

chaperones in non-stressed cells. EMBO J 16:7207-16.

19. B'Chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi Y, Parry L,

Stepien G, Fafournoux P, Bruhat A. 2013. The eIF2alpha/ATF4 pathway is essential

for stress-induced autophagy gene expression. Nucleic Acids Res 41:7683-99.

20. Datta SR, Dudek H, Tao X, Masters S, Fu HA, Gotoh Y, Greenberg ME. 1997.

Akt phosphorylation of bad couples survival signals to the cell-intrinsic death

machinery. Cell 91:231-241.

21. Zha JP, Harada H, Osipov K, Jockel J, Waksman G, Korsmeyer SJ. 1997. Bh3

domain of bad is required for heterodimerization with bcl-x-l and pro-apoptotic

activity. Journal of Biological Chemistry 272:24101-24104.

22. Green BD, Jabbour AM, Sandow JJ, Riffkin CD, Masouras D, Daunt CP,

Salmanidis M, Brumatti G, Hemmings BA, Guthridge MA, Pearson RB, Ekert PG.

2013. Akt1 is the principal Akt isoform regulating apoptosis in limiting cytokine

concentrations. Cell Death Differ 20:1341-9. doi: 10.1038/cdd.2013.63. Epub 2013

Jun 21.

23. Ekert PG, Jabbour AM, Manoharan A, Heraud JE, Yu J, Pakusch M, Michalak

EM, Kelly PN, Callus B, Kiefer T, Verhagen A, Silke J, Strasser A, Borner C, Vaux

DL. 2006. Cell death provoked by loss of interleukin-3 signaling is independent of

Bad, Bim, and PI3 kinase, but depends in part on Puma. Blood 108:1461-8.

24. O'Neill KL, Huang K, Zhang J, Chen Y, Luo X. 2016. Inactivation of

prosurvival Bcl-2 proteins activates Bax/Bak through the outer mitochondrial

membrane. Genes Dev 30:973-88. doi: 10.1101/gad.276725.115. Epub 2016 Apr 7.

25. Goodall KJ, Finch-Edmondson ML, van Vuuren J, Yeoh GC, Gentle IE, Vince

JE, Ekert PG, Vaux DL, Callus BA. 2016. Cycloheximide Can Induce Bax/Bak

Dependent Myeloid Cell Death Independently of Multiple BH3-Only Proteins. PLoS

One 11:e0164003. doi: 10.1371/journal.pone.0164003. eCollection 2016.

26.   Callus BA, Mathey-Prevot B. 1998. Interleukin-3-induced activation of the

JAK/STAT pathway is prolonged by proteasome inhibitors. Blood 91:3182-92.

27.   Chao JR, Wang JM, Lee SF, Peng HW, Lin YH, Chou CH, Li JC, Huang HM,

Chou CK, Kuo ML, Yen JJ, Yang-Yen HF. 1998. mcl-1 is an immediate-early gene

activated by the granulocyte-macrophage colony-stimulating factor (GM-CSF)

signaling pathway and is one component of the GM-CSF viability response. Mol Cell

Biol 18:4883-98.

28.   Moulding DA, Quayle JA, Hart CA, Edwards SW. 1998. Mcl-1 expression in

human neutrophils: regulation by cytokines and correlation with cell survival. Blood

92:2495-502.

29.   Lim JP, Gosavi P, Mintern JD, Ross EM, Gleeson PA. 2015. Sorting nexin 5

selectively regulates dorsal-ruffle-mediated macropinocytosis in primary

macrophages. J Cell Sci 128:4407-19.

30.   Aubrey BJ, Kelly GL, Kueh AJ, Brennan MS, O'Connor L, Milla L, Wilcox S,

Tai L, Strasser A, Herold MJ. 2015. An inducible lentiviral guide RNA platform

enables the identification of tumor-essential genes and tumor-promoting mutations in

vivo. Cell Rep 10:1422-32.

31.   Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide

libraries for CRISPR screening. Nat Methods 11:783-784.

32.   Wang Z, Rong YP, Malone MH, Davis MC, Zhong F, Distelhorst CW. 2006.

Thioredoxin-interacting protein (txnip) is a glucocorticoid-regulated primary response

gene involved in mediating glucocorticoid-induced apoptosis. Oncogene 25:1903-13.

33.   Brumatti G, Salmanidis M, Kok CH, Bilardi RA, Sandow JJ, Silke N, Mason K,

Visser J, Jabbour AM, Glaser SP, Okamoto T, Bouillet P, D'Andrea RJ, Ekert PG.

2013. HoxA9 regulated Bcl-2 expression mediates survival of myeloid progenitors

and the severity of HoxA9-dependent leukemia. Oncotarget 4:1933-47.

Figure Legends

Figure 1. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells shrink and arrest but remain viable when cultured

without IL3.

A. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells survive long-term in the absence of IL3. *Bax$^{+/+}$Bak1$^{+/+}$*

wild-type (black circles) and *Bax$^{-/-}$Bak1$^{-/-}$* (green squares) FDM cells were cultured

without IL3. At the indicated times, cells were harvested, resuspended in PBS

containing propidium iodide (PI) and analyzed by flow cytometry.

B. IL3 withdrawal caused cells to shrink. PI negative populations of *Bax$^{-/-}$Bak1$^{-/-}$* cells

cultured for 48 h in the presence (light gray) or absence (dark gray) of IL3, and the

latter that had IL3 returned for 72 h (green), were compared by forward scatter as an

indicator of cell size.

C. IL3 withdrawal caused cell cycle arrest, which was reversed by re-addition of IL3.

Nuclei from *Bax$^{-/-}$Bak1$^{-/-}$* cells cultured as described in (B) were stained with PI to

indicate DNA content by flow cytometry.

Figure 2. *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* FDM cells shrink and arrest, but remain viable when IL3

is withdrawn.

A. Whole cell lysates from *Bax$^{-/-}$Bak1$^{-/-}$* cells and 3 *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clonal lines

cultured in the presence of IL3, in Hank's Buffered Saline Solution (HBSS) for 2 h, or

in the absence of IL3 for 2 or 48 h, were subjected to Western blotting. The Atg5

immunoblot demonstrates successful CRISPR/Cas9-mediated deletion of *Atg5*. The

LC3B immunoblot demonstrates lack of LC3B lipidation (isoform II) and autophagic

flux (HBSS treatment) in Δ*Atg5* cell clones.

B. *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Atg5* FDM cells survive long-term in the absence of IL3. The

indicated *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Atg5* clones were cultured without IL3 and viability was

determined by PI uptake and flow cytometry at the indicated times.

C. *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Atg5* cells (clone 1) were cultured for 48 h in the presence or absence

of IL3 and PI negative populations were compared by forward scatter to indicate

differences in cell size.

D. Cells from *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Atg5* clone 1 were cultured for 48 h in the presence or

absence of IL3 then processed in PI buffer to measure DNA content by flow

cytometry.

Figure 3. *Bax*$^{-/-}$*Bak1*$^{-/-}$ FDM cells arrested by IL3 deprivation survive in the absence

of added glucose.

A. Txnip is not required for the decrease in Glut1 following IL3 withdrawal. Whole

cell lysates from *Bax*$^{-/-}$*Bak1*$^{-/-}$ and 3 independent *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Txnip* FDM cell clones

cultured for 48 h in the presence or absence of IL3 were analyzed by Western blotting

to detect Glut1 and Txnip protein.

B. Forced expression of Txnip is not sufficient to decrease Glut1 protein levels.

Independent *Bax*$^{-/-}$*Bak1*$^{-/-}$ FDM clones bearing a doxycycline (dox)-inducible FLAG-

Txnip construct or empty vector control were cultured for 72 h in the absence or

presence of dox. Whole cell lysates were prepared and analyzed by Western blotting

to detect Glut1 and Txnip protein.

C. *Bax*$^{-/-}$*Bak1*$^{-/-}$ and *Bax*$^{-/-}$*Bak1*$^{-/-}$Δ*Txnip* cells can take up and metabolise glucose.

Cells cultured for 72 h in the presence or absence of IL3 were analyzed on the

Seahorse bioanalyzer to measure the extracellular acidification rate (ECAR) as an

indicator of glucose metabolism. The increased ECAR following addition of glucose

signifies increased glycolytic flux in cells. Oligomycin inhibits mitochondrial ATP

synthase and the change in ECAR upon its addition represents the glycolytic reserve

capacity of cells. Glycolysis is inhibited by 2-deoxyglucose (2-DG) so upon its

addition, the ECAR values are produced from non-glycolytic sources. Equal numbers

of cells were seeded for analysis. To correct for the reduced size of IL3-starved cells,

protein concentration was assayed and the ECAR values were normalized to 1 µg

total protein per sample. Curves represent the mean of 3 replicate samples depicted as

solid symbols for each time point.

D. *Bax$^{-/-}$Bak1$^{-/-}$* and *Bax$^{-/-}$Bak1$^{-/-}$ΔTxnip* cells deprived of IL3 survive long-term in the

absence of added glucose. Cells were cultured without or with IL3 in complete or

glucose-free medium and analyzed for PI uptake by flow cytometry at the indicated

times.

Figure 4. When starved of glutamine, *Bax$^{+/+}$Bak1$^{+/+}$* FDM cells undergo apoptosis,

whereas *Bax$^{-/-}$Bak1$^{-/-}$* cells arrest, but can survive for weeks.

A. *Bax$^{+/+}$Bak1$^{+/+}$* cells undergo apoptosis in the absence of glutamine (upper panels),

whereas *Bax$^{-/-}$Bak1$^{-/-}$* cells survive in the absence of glutamine, but arrest even when

IL3 is present (lower panels). Cells were cultured with or without IL3 in glucose- and

glutamine-free medium for 4 days then processed in PI buffer to assess DNA content

by flow cytometry. Note the absence of cells in S phase when *Bax$^{-/-}$Bak1$^{-/-}$* cells were

cultured without glutamine.

B. *Bax^{-/-}Bak1^{-/-}* cells survive long-term in the absence of glutamine. Cells were

cultured with or without IL3 in glucose- and glutamine-free medium. At the indicated

times, cells were assayed for viability by PI uptake and flow cytometry.

C. *Bax^{-/-}Bak1^{-/-}* cells require glutamine to divide. Cells were cultured with or without

IL3 for 7 days in glucose- and glutamine-free medium. They were then split in three,

and resuspended in medium without glucose or glutamine, or containing only glucose

(5.55 mM), or only glutamine (4 mM), for a further 2 days. Cells were harvested and

resuspended in PI buffer to assess DNA content by flow cytometry.

Figure 5. Withdrawal of IL3 or glutamine caused cell shrinkage independent of Atg5-

mediated autophagy.

A. *Bax^{-/-}Bak1^{-/-}* cells were cultured in glucose- and glutamine-free medium, with or

without IL3. At the indicated times, cell volume was measured using a Coulter-based

CASY cell counter.

B. *Bax^{-/-}Bak1^{-/-}ΔAtg5* clone 1 was cultured and analyzed as described in **A**.

Figure 6. Withdrawal of IL3 or glutamine reduced the expression of Mcl-1 and

several key regulators of cell cycle progression.

A. *Bax^{-/-}Bak1^{-/-}* cells and *Bax^{-/-}Bak1^{-/-}ΔAtg5* clone 1 were cultured for 72 h with or

without IL3 (+/–) in complete (+) or glucose (Glc)- and glutamine (Q)-free (–)

medium. Whole cell lysates were prepared and analyzed by Western blotting to detect

the indicated proteins. Three independent membranes were used for immunoblotting

and β-actin loading control blots for each are shown. Membrane #1: anti-Mcl-1 and

anti-c-Myc; membrane #2: anti-Cdk4; membrane #3: anti-cyclin D3 and anti-Atf4.

B. *Bax$^{+/+}$Bak1$^{+/+}$* and *Bax$^{-/-}$Bak1$^{-/-}$* cells cultured in complete medium with IL3 were

treated with DMSO (vehicle control) or 10 μM of the Mcl-1 inhibitor S63845 for 24

h. Cells were harvested and viability determined by PI exclusion and flow cytometry.

Bars depict the mean of 3 independent experiments represented by the 3 different

symbols, performed in triplicate. The experiment represented by the solid circles was

performed with S63845 synthesized by SYNthesis Med Chem. Experiments

represented by the solid triangles and open hexagons were performed with S63845

synthesized by Active Biochem (see Materials and Methods).

Figure 7.

The growth factor IL3 stimulates the expression of Mcl-1 that inhibits activation of

Bax/Bak1, and stimulates expression of c-Myc, Cdk4 and cyclin D3 to promote cell

growth and division. Glutamine is required to maintain levels of these proteins needed

to prevent activation of Bax/Bak1 (Mcl-1), as well as for cell proliferation (c-Myc,

Cdk4, cyclin D3 and possibly others). In the absence of growth factor, glucose,

glutamine and Atg5-dependent autophagy, we hypothesise that *Bax$^{-/-}$Bak1$^{-/-}$* myeloid

cells survive by consuming nutrients in fetal bovine serum.

A

B *Bax⁻/⁻Bak1⁻/⁻*

C *Bax⁻/⁻Bak1⁻/⁻*

A

B

C *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clone 1

D *Bax⁻/⁻Bak1⁻/⁻ΔAtg5* clone 1

A

B   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* cells

C   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* cells

A   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* cells

B   *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clone 1

A

B

FIGURE 7   Dong et al.