In the Absence of Apoptosis, Myeloid Cells Arrest When Deprived of Growth Factor,

But Remain Viable Without ATG5-dependent Autophagy or Uptake of Glucose or

Glutamine.

Li Dong, Boris Reljic, Lisa M. Lindqvist, David L. Vaux and Hoanh Tran#

The Walter and Eliza Hall Institute of Medical Research, and the Department of

Medical Biology, University of Melbourne, Victoria, Australia.

Running Head: Cell Survival Without Glucose, Glutamine, or Autophagy

#Address correspondence to Hoanh Tran, tran.h@wehi.edu.au

Abstract

Withdrawal of growth factor interleukin 3 from IL3-dependent myeloid cells causes

them to undergo Bax/Bak1-dependent apoptosis, whereas factor-deprived $Bax^{-/-}Bak1^{-/-}$

cells remain viable, but arrest. It was reported that withdrawal of IL3 from $Bax^{-/-}Bak1^{-/-}$

$^{-/-}$ cells caused decreased expression of the glucose transporter Glut1, leading to

reduced glucose uptake, so that arrested cells required Atg5-dependent autophagy for

long-term survival. In other cell types, a decrease in Glut1 is mediated by thioredoxin-

interacting protein Txnip, which is induced in IL3-dependent myeloid cells when

growth factor is removed. We mutated *Atg5* and *Txnip* by CRISPR/Cas9 and found

that Atg5-dependent autophagy was not necessary for the long-term viability of

cycling or arrested $Bax^{-/-}Bak1^{-/-}$ cells, and that Txnip was not required for the decrease

in Glut1 expression in response to IL3 withdrawal. Surprisingly, in glucose-free

medium, wild-type and *Bax*/*Bak1* double mutant cells not only survived, but

continued to cycle when IL3 was present. In contrast, $Bax^{+/+}Bak1^{+/+}$ cells died by

apoptosis when deprived of glutamine, and while $Bax^{-/-}Bak1^{-/-}$ cells deprived of

glutamine survived, they arrested even when IL3 was present. Thus, glutamine, but

neither Atg5-dependent autophagy nor glucose uptake, is necessary for cell division,

and none are needed for arrested $Bax^{-/-}Bak1^{-/-}$ myeloid cells to survive for several

weeks.

Introduction

Mammalian cells import a variety of extracellular nutrients to support their survival and growth. *In vitro*, cells are typically propagated in medium supplemented with glucose, glutamine and fetal bovine serum. To generate chemical energy, glucose can be metabolized by oxidative phosphorylation in the mitochondria or by anaerobic glycolysis. Glutamine can be used as an energy source through glutaminolysis, and can also be used, together with a multitude of factors present in serum, to allow the synthesis of essential macromolecules.

Factor dependent myeloid (FDM) progenitor cell lines (1, 2) require the haematopoietic growth factor interleukin 3 (IL3) to survive and multiply in culture. When IL3 is not provided, FDM cells undergo Bax/Bak1-dependent apoptosis. FDM cells derived from $Bax^{-/-}Bak1^{-/-}$ mice survive in the absence of IL3, but they undergo cell cycle arrest, similar to those over-expressing Bcl2 (3, 4). Previously it was reported that to survive longer than 4 days, IL3-deprived $Bax^{-/-}Bak1^{-/-}$ cells required Atg5-dependent autophagy (4), an intracellular catabolic process that can support the survival of starved or arrested cells (5).

Cells import glucose when it is available and growth factors can stimulate the amount of glucose that is consumed. $Bax^{-/-}Bak1^{-/-}$ FDM cells arrested by IL3 withdrawal decrease expression of the glucose transporter Glut1, which was proposed to reduce glucose uptake and metabolism (4). Several mechanisms have been put forward to explain the regulation of expression and plasma membrane translocation of Glut1. Activation of autophagy was shown to be required for the efficient surface expression of Glut1 in mouse embryonic fibroblasts cultured under hypoxic conditions (6). In

human hepatocytes subjected to glucose starvation, a mechanism involving

proteasomal degradation of the thioredoxin-interacting protein Txnip was required for

Glut1 stabilization (7). Moreover, it was shown that Txnip directly binds to Glut1 and

promotes its internalization, thus reducing Glut1 on the plasma membrane and the

ability of cells to take up glucose (7).

In this study, we employed $Bax^{-/-}Bak1^{-/-}$ FDM cells to genetically test whether Atg5-

dependent autophagy is required for the sustained survival of arrested cells deprived

of growth factor. We also asked whether autophagy and/or Txnip are required for the

reduction in Glut1 expression and glucose uptake by arrested myeloid cells, and

whether cells required glucose uptake in order to cycle.

Results

$Bax^{-/-}Bak1^{-/-}$ myeloid cells shrink and arrest when deprived of IL3, but survive.

Factor dependent myeloid (FDM) cells require interleukin 3 (IL3) to survive and

proliferate (2, 4, 8). When deprived of IL3 for 4 days, >80 % of wild-type

$Bax^{+/+}Bak1^{+/+}$ FDM cells died, as indicated by loss of plasma membrane integrity and

uptake of propidium iodide (Figs. 1A and Supplemental S1A). By contrast, and

consistent with the findings of Lum *et al*. (4), $Bax^{-/-}Bak1^{-/-}$ FDM cells survived in the

absence of IL3 for more than 14 days. After 48 h IL3 withdrawal, these cells shrank in

size and arrested (Fig. 1, B and C). This quiescent state was reversible, because when

IL3 was returned, the cells increased in size and resumed cycling. Thus, in the

absence of Bax/Bak1-mediated apoptosis, FDM cells can survive for weeks without

IL3, but they shrink and arrest.

Bax$^{-/-}$Bak1$^{-/-}$ myeloid cells arrested by deprivation of IL3 do not require autophagy to

survive.

During macroautophagy, LC3B-I is lipidated by a multiprotein complex containing

Atg5 to form LC3B-II, which can incorporate into the growing autophagosome, and is

a common marker of autophagy. Lysates of *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells contained similar

amounts LC3B-II, whether they were cultured in the presence or absence of IL3 (Fig.

2A, lanes 1 and 2). This suggests autophagy is occurring in both cycling and arrested

cells. To determine if autophagy was needed to support the sustained survival of

arrested IL3-starved cells, we used CRISPR/Cas9 to delete *Atg5* in *Bax$^{-/-}$Bak1$^{-/-}$* cells,

generating *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* triple knockout FDM cell clones. Western blotting

confirmed the three *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones had much less LC3B-II than the *Bax$^{-/-}$*

Bak1$^{-/-}$ cells, whether they were cultured with IL3 or not (Fig. 2A). After 14 days in

culture without IL3, all *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* clones remained ~80% viable (Figs. 2B

and Supplemental S1B). Similar to *Bax$^{-/-}$Bak1$^{-/-}$* FDMs, *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* cells

arrested and shrank in size within 48 h of growth factor withdrawal (Fig. 2, C and D).

These results confirm that IL3 receptor signals are needed for *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells

to maintain their normal size, but show that Atg5-dependent autophagy is not required

for their survival, whether they are cycling or are arrested by IL3 withdrawal.

Furthermore, because culturing without IL3 caused a similar decrease in size of the

arrested *Bax$^{-/-}$Bak1$^{-/-}$* as of the *Bax$^{-/-}$Bak1$^{-/-}$ΔAtg5* cells, this shrinkage is not due to

Atg5-dependent autophagy.

Bax$^{-/-}$Bak1$^{-/-}$ myeloid cells arrested by IL3 deprivation survive in the absence of

glucose.

Lum *et al*. reported that when growth factor was removed, *Bax^-/-Bak1^-/-* FDM cells

decreased expression of the glucose transporter Glut1 (4). A possible mechanism for

this was proposed by Wu *et al*., who found that thioredoxin-interacting protein

(Txnip) could not only bind to Glut1 and induce its internalization through clathrin-

coated pits, thus removing it from the plasma membrane, but could also reduce the

level of Glut1 messenger mRNA (7). Consistent with Lum *et al*. and Wu *et al*., Txnip

was increased, and Glut1 was decreased, in lysates of factor-starved *Bax^-/-Bak1^-/-* cells

48 h following removal of IL3 (Fig. 3A, lanes 1 and 2).

To determine whether expression of Txnip caused or correlated with changes in Glut1

levels, we deleted *Txnip* in *Bax^-/-Bak1^-/-* FDM cells using CRISPR/Cas9. Western

blotting confirmed the absence of Txnip protein in three *Bax^-/-Bak1^-/-ΔTxnip* clones

(Fig. 3A). We found that Glut1 levels were unaffected by deletion of *Txnip*, whether

IL3 was added or not, indicating that the increase in Txnip was not required for the

decrease in Glut1 when cells were cultured without IL3. Furthermore, withdrawal of

IL3 caused G1 arrest in *Bax^-/-Bak1^-/-ΔTxnip* cells within 48 h, similar to *Bax^-/-Bak1^-/-*

cells (Figs. 1A and Supplemental S2A). Because induced over-expression of FLAG-

Txnip in *Bax^-/-Bak1^-/-* cells had no impact on levels of Glut1 protein or cell cycle

progression (Figs. 3B and Supplemental S2B), an increase in Txnip is not sufficient to

reduce Glut1. Therefore, in FDM cells, the abundance of Txnip has no effect on levels

of Glut1, whether they are treated with IL3 or not.

Levels of Glut1 were similar among proliferating Bax/Bak1-null cells of all genotypes

and the decrease in Glut1 expression following IL3 withdrawal in *Bax^-/-Bak1^-/-ΔAtg5*

clones compared to *Bax^-/-Bak1^-/-* cells was unaffected (Supplemental Fig. S2C).

Together, these results indicate that although IL3 can promote expression of Glut1,

neither Txnip nor Atg5-dependent autophagy is required for the decrease in Glut1

when growth factor is removed from myeloid cells, and neither absence nor

overexpression of Txnip affects the levels of Glut1.

Because Glut1 levels are reduced in IL3-starved $Bax^{-/-}Bak1^{-/-}$ FDM cells, we asked

whether glucose utilization was reduced. The rate of glucose metabolism in $Bax^{-/-}$

$Bak1^{-/-}$ FDMs was reduced after IL3 withdrawal, but they nevertheless still responded

to the addition of glucose, as indicated by the increased extracellular acidification rate

(ECAR, Fig. 3C). In the absence of IL3, $Bax^{-/-}Bak1^{-/-}$ FDM lines that also lacked

Txnip or Atg5 had similar rates of glucose metabolism to the parental Bax/Bak1-null

cells (Figs. 3C and Supplemental S2D). Thus, arrested FDM cells retain the capacity

to import and metabolise glucose, albeit at a reduced level.

Because the arrested IL3-starved cells were able to take up glucose, we hypothesised

that this was sufficient to support their sustained survival despite the decrease in

Glut1. To test their requirement for glucose, the cells were cultured in medium that

lacked glucose. Surprisingly, $\geq$ 80% of IL3-starved $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}$

$\Delta Txnip$ cells remained viable at 24 days without glucose (Fig. 3D). Furthermore,

when IL3 was present, these cells were able to proliferate (Supplemental Fig. S2E).

These results indicate that FDM cells will import glucose whether IL3 is present or

not, and that they can still proliferate even in the absence of glucose.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to proliferate, but can survive in its

absence for weeks.

When glucose is limiting, lymphoid cells can metabolise glutamine to survive and

proliferate (9). We therefore asked whether FDM cells exhibited similar requirements

for glutamine when cultured in the absence of glucose. In medium that lacked both

glucose and glutamine, with or without IL3, >95% of wild-type $Bax^{+/+}Bak1^{+/+}$ FDM

cells died as indicated by uptake of PI by day 4 (Supplemental Fig. S3A), and

consistent with their death being due to apoptosis, the nuclei of the cells had sub-2n

DNA (Fig. 4A). Apoptosis was Bax/Bak1 dependent, because Bax/Bak-null cells did

not die when cultured in medium lacking glucose and glutamine, but they arrested

instead (Figs. 4A and Supplemental S3B). When IL3 was present, they arrested as

large cells, whereas when cultured without IL3, they arrested as small cells (Figs. 4B

and Supplemental S3C).

In medium without glucose, glutamine, or IL3, $Bax^{-/-}Bak1^{-/-}$ FDM cells exhibited a

slight decline in viability at day 16 and beyond compared to cultures with IL3 (Fig.

4C). Interestingly, in the presence of IL3, >90% of cells cultured with neither glucose

nor glutamine continued to survive beyond 28 days. $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ clones

cultured in medium lacking glucose and glutamine also maintained ~80% or greater

viability at day 16, with or without IL3 (Supplemental Fig. S3D).

When $Bax^{-/-}Bak1^{-/-}$ FDM cells were starved of glucose and glutamine, most arrested in

G_1 with 2n DNA, but some arrested in G_2/M with 4n DNA (Fig. 4, A and D). When

glutamine was added back to $Bax^{-/-}Bak1^{-/-}$ cells that had been cultured for 7 days

without glucose or glutamine, the arrested cells resumed cycling within 48 h in the

presence of IL3, whereas adding back glucose had no effect (Fig. 4D). These results

indicate that IL3-treated $Bax^{-/-}Bak1^{-/-}$ myeloid cells require glutamine to proliferate,

but can survive up to 4 weeks or more in its absence. Because $Bax^{-/-}Bak1^{-/-}\Delta Atg5$

FDMs behaved similarly to $Bax^{-/-}Bak1^{-/-}$ cells (Supplemental Fig. S3B), Atg5-

dependent autophagy is not required for sustained cell survival, even in the absence of

glucose and glutamine.

Withdrawal of IL3 or glutamine caused reduced expression of Mcl-1 and key

regulators of cell cycle progression.

To elucidate how growth factor and nutrient deprivation impacted FDM cell survival

and proliferation, we performed Western blotting to determine protein levels of

apoptotic and cell cycle regulators. We found that levels of the pro-survival protein

Mcl-1 were markedly reduced in $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells after 3 days

in culture without glucose and glutamine (Fig. 5A), suggesting that the decrease in

normal levels of Mcl-1 allowed activation of Bax/Bak1 and the apoptosis of wild-type

FDM cells observed in Fig. 4A. Consistent with this notion, addition of a specific

Mcl-1 inhibitor S63845 (10) to $Bax^{+/+}Bak1^{+/+}$ cells was sufficient to cause significant

apoptosis within 24 h, even when the cells were cultured in complete medium

containing glutamine and IL3 (Fig. 5B).

As well as causing arrest, IL3 withdrawal reduced c-Myc protein to barely detectable

levels in both $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells (Fig. 5A). Cdk4 and cyclin D3

are key drivers of cell cycle progression in haematopoietic cell lines (11, 12), and

these proteins were also markedly reduced in the IL3-starved, Bax/Bak1-deficient

myeloid cells (Fig. 5A), suggesting a mechanism for their arrest (Figs. 1C and 2C).

Withdrawal of glutamine from these cells also caused cell cycle arrest, even in the

presence of IL3 (Figs. 4A and Supplemental S3B). Under these conditions, IL3 was

able to stimulate the expression of c-Myc and Cdk4, but the levels of these proteins

were significantly lower than in the cells cultured with both glutamine and IL3 (Fig.

5A). The transcription factor Atf4 accumulates in response to a variety of cellular

stresses including amino acid depletion (13). In glutamine-starved $Bax^{-/-}Bak1^{-/-}$ and

$Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells, Atf4 levels increase strongly, but intriguingly only in the

presence of IL3. These results indicate that the signal transduction pathway activated

by ligation of the IL3 receptors remains largely intact in cells lacking glutamine, and

can stimulate the cells to remain large, but that glutamine is essential for high levels

of c-Myc and Cdk4/cyclin D3 expression, and for the cells to divide.

$Bax^{-/-}Bak1^{-/-}$ myeloid cells require serum factors to survive long-term in the absence

of glutamine.

The cells cultured without glutamine or glucose would need a source of energy to

survive, so we tested whether they were obtaining it from fetal bovine serum (FBS) in

the medium. When cultured in 2% FBS, all $Bax^{-/-}Bak1^{-/-}$ cells died after 4 days,

whereas at the same time ~20% of cells cultured with 4% FBS were viable (Fig. 6A).

More than 80% of cells remained viable for 16 days and beyond when cultured with

6%, 8% or 10% FBS. Similar findings were observed for $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells

(Fig. 6B). These results suggest that $Bax^{-/-}Bak1^{-/-}$ myeloid cells survive for several

weeks without glucose and glutamine by consuming nutrients in the FBS, rather than

by using autophagy.

Discussion

IL3-dependent myeloid cell lines have provided many insights into the regulation of

fundamental processes such as cell death, cell division, and autophagy, as well as the

signaling pathways that regulate them. For example, it was experiments on an IL3-

dependent myeloid cell line that first revealed that the function of Bcl-2 was to inhibit

cell death (3). When genes for Bax and Bak1 are deleted in factor-dependent myeloid

(FDM) cells, they fail to undergo apoptosis when IL3 is removed, but nevertheless

will die if starved of nutrients needed to maintain their metabolism. When $Bax^{-/-}Bak1^{-}$

$^{/-}$ FDMs are cultured without IL3, they decrease expression of the glucose transporter

Glut1, shrink and arrest, but do not die (4). This led to the proposal that the cells could

not take up glucose, so they needed autophagy to maintain their metabolism to

survive. Consistent with this idea, Lum et $al.$ found that inhibition of autophagy either

by 3-methyladenine or shRNA knockdown of Atg5 caused the cells to die (4).

However, we also used shRNA to reduce Atg5 levels and found that Atg5-dependent

autophagy was not required for the survival $Bax^{-/-}Bak1^{-/-}$ myeloid cells deprived of

IL3 (14). In this report, we have genetically eliminated $Atg5$ expression in IL3-

dependent $Bax^{-/-}Bak1^{-/-}$ cells using CRISPR/Cas9. Unlike Lum et $al.$, who concluded

that autophagy was necessary to sustain the survival of IL3-deprived $Bax^{-/-}Bak1^{-/-}$

FDM cells beyond 4 days (4), our results demonstrate that canonical Atg5-dependent

autophagy pathways are not required to support the long-term viability of these cells

(Fig. 2).

We were able to confirm the observation that culture of FDM cells without IL3

reduced the levels of the glucose transporter Glut1 (4) (Fig. 3A), and wondered if the

mechanism for reduction of Glut1 would be similar to that described by Wu et $al.$ who

found that thioredoxin interacting protein (Txnip) could reduce the levels of Glut1 in

human hepatocytes (7). Although we found that Txnip increased in FDM cells when

IL3 was removed, levels of Glut1 decreased by the same amount when IL3 was

removed from Txnip-null cells, and over-expression of exogenous Txnip did not

reduce the levels of Glut1 (Fig. 3, A and B). Although we confirmed the reported

inverse correlation between levels of Txnip and Glut1, Txnip is neither required nor

sufficient for IL3 to regulate levels of Glut1 in FDMs.

Canonical autophagy was shown to be required for the efficient surface expression of

Glut1 (6). Our Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ FDM cells had similar levels of Glut1 to

parental $Bax^{-/-}Bak1^{-/-}$ cells, and they decreased Glut1 to a similar extent when cultured

without IL3. Furthermore, the $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ cells showed similar levels of

glucose uptake to $Bax^{-/-}Bak1^{-/-}$ cells when cultured with IL3 and without. Therefore, in

FDM cells, Atg5-dependent autophagy is not needed to maintain normal levels of

Glut1, and does not affect the ability of cells to take up and metabolize glucose (Fig.

S2, C and D).

We were surprised to find that Bax/Bak1-null FDM cells of all genotypes, when

cultured in the absence of glucose and arrested due to withdrawal of IL3, not only

survived, but that these cells, and even wild-type $Bax^{+/+}Bak1^{+/+}$ cells, proliferated in

the absence of glucose when IL3 was present (Figs. 3D and S2E; data not shown).

Therefore, even though FDM cells will take up and metabolise glucose when it is

available, even if they are arrested and their Glut1 levels are decreased, they have no

obligatory requirement for glucose, either to survive or to proliferate. Our work

demonstrates that reduced uptake of glucose is not responsible for the arrest of $Bax^{-/-}$

$Bak1^{-/-}$ FDM cells upon IL3 withdrawal.

Consistent with the requirement of glutamine for lymphoid cell survival (9), we found

that wild-type factor dependent myeloid cells died following glutamine deprivation

(Fig. 4). It was nevertheless surprising that glutamine was not required to sustain the

survival of $Bax^{-/-}Bak1^{-/-}$ FDM cells in the absence of glucose or Atg5-dependent

autophagy, and that the glutamine-deprived cells maintained a large size in the

presence of IL3, but arrested. This suggests that glutamine provided the source of

amino acids needed for cell division, but that these cells can maintain enough

metabolic activity to survive even in its absence.

Several pathways have been proposed to explain how engagement of IL3 receptors

prevents activation of Bax and Bak1. For example, it was proposed that IL3 receptors

signaled via Akt to phosphorylate Bad, preventing it from binding to Bcl-x, to thereby

free Bcl-x to inhibit apoptosis (15, 16). However, experiments in which genes for Bad

or Akt have been mutated have shown that unlike Bax/Bak1, they play relatively

minor roles in the regulation of apoptosis by IL3 (17, 18). These experiments, and

those where genes for all the Bcl-2 family member were deleted (19) suggest that

Bax/Bak1 dependent apoptosis is not always signaled via BH3-only proteins, but

more often regulated by controlling the abundance of anti-apoptotic Bcl-2 family

members, such as Bcl-2, Bcl-x, and Mcl-1 (20). The results presented here are

consistent with this, and show that apoptosis of FDMs following the removal of IL3,

or culture in the absence of glutamine, is due to the decrease in abundance of Mcl-1

(Fig. 5A). Cytokines including IL3 keeps cells alive by driving *Mcl-1* transcription,

most likely via Jak2 and Stat5 (21-23), and glutamine is required to maintain the

production of Mcl-1 to prevent default apoptosis. Consistent with a requirement for

Mcl-1 to prevent spontaneous activation of Bax/Bak1, a specific inhibitor of Mcl-1

was able to induce apoptosis of wild-type $Bax^{+/+}Bak1^{+/+}$ FDM cells cultured with IL3

and glutamine (Fig. 5B).

In the absence of glutamine, pathways activated by IL3 could still converge to induce

the expression of c-Myc in $Bax^{-/-}Bak1^{-/-}$ FDM cells, and although this was response

was blunted, the arrested cells maintained their large size (Figs. 4B and 5A). We

found that the transcription factor Atf4 was strongly induced by IL3 in glutamine-

starved cells (Fig. 5A), but at present, it is unclear how or why signaling from the IL3

receptor leads to increased Atf4 production. Given that IL3 can stimulate the

expression of ER chaperones in FDM cells (24) and that Atf4 translation increased in

cells starved of amino acids or exposed to stimuli that induced ER stress (13), it is

plausible that IL3 activated the unfolded protein response in the glutamine-deprived

FDMs. Atf4 has also been shown to promote the expression of several autophagy

genes (25), but we have not been able to confirm that its accumulation induced LC3B

lipidation in the glutamine-starved FDM cells. Nevertheless, IL3 also induced Atf4

expression in Atg5-deficient $Bax^{-/-}Bak1^{-/-}$ cells cultured in glutamine-free media (Fig.

5A), suggesting that Atf4 does not promote canonical autophagy to maintain cell

viability under these conditions. The importance of Atf4 for the sustained survival of

$Bax^{-/-}Bak1^{-/-}$ FDM cells arrested by glutamine withdrawal will be determined when

we genetically delete $Atf4$ in these cells.

In the absence of glucose, glutamine and Atg5-dependent autophagy, $Bax^{-/-}Bak1^{-/-}$

FDM cells remained viable for weeks, and their sensitivity to low concentrations of

fetal bovine serum (Fig. 6) are consistent with the hypothesis that they took up and

used components present in serum to maintain enough metabolism in order to survive.

Serum factors are likely scavenged by endocytic processes such as macropinocytosis,

which can be stimulated by growth factors (26). Increased endocytic uptake of serum

proteins could account for the observation that at ~2 weeks and beyond in culture

without glutamine, more Bax/Bak1-null cells survived when growth factor was

provided (Figs. 4C and S3E).

Bax$^{-/-}$Bak1$^{-/-}$ FDM cells cultured in the absence of glutamine express low levels of the

cell cycle regulators Cdk4 and cyclin D3 (Fig. 5A). Reduced levels of these proteins,

combined with possibly altered levels of other cell cycle regulators, could explain

why glutamine-starved cells arrested, even in the presence of IL3 that stimulated them

to maintain a large size (Fig. 7). In the absence of growth factor, glucose, glutamine

and Atg5-dependent autophagy, we hypothesise that *Bax$^{-/-}$Bak1$^{-/-}$* myeloid cells

survive by consuming proteins present in bovine serum, and degrading these in

lysosomes and/or proteasomes to maintain a sufficient pool of amino acids. Future

studies will address this hypothesis and determine how cells generate the energy they

need to stay viable under conditions of limited nutrients.

Materials & Methods

Cell culture

Wild-type *Bax$^{+/+}$Bak1$^{+/+}$* and *Bax$^{-/-}$Bak1$^{-/-}$* IL3-dependent mouse myeloid progenitor

cell lines (FDM) have been described (18). Cells were cultured in Dulbecco's

Modified Eagle Medium (DMEM, ThermoFisher Scientific Cat# 11885-084;

containing 1 g/L glucose and 4 mM glutamine) supplemented with 10% fetal bovine

serum (FBS, ThermoFisher Scientific Cat# 16000-044) and 0.25 ng/ml recombinant

murine interleukin 3 (IL3, PeproTech, Cat# 213-13) in a 37ºC humidified incubator

with 10% CO_2. Cells were deprived of IL3, glucose, or glucose and glutamine by

washing 3 times with medium (+10% FBS) that lacked those nutrients, then

resuspended in the same medium for analysis at the indicated time points. DMEM

lacking D-Glucose and sodium pyruvate (Cat# 11966-025) and DMEM lacking D-

Glucose, sodium pyruvate, L-Glutamine and phenol red (Cat# A14430-01) were

purchased from ThermoFisher Scientific. D-Glucose was purchased from Sigma

(Cat# G7528).

Lentiviral constructs for CRISPR/Cas9 and inducible overexpression

An inducible lentiviral CRISPR/Cas9 system (27) was used to mutate *Txnip* in *Bax*$^{-/-}$

Bak1$^{-/-}$ FDM cells. Guide RNA oligos targeting mouse *Txnip* (guide 1: CCA GCA

GGT GAG AAC GAG A; guide 2: TAT GGT TGC GTA GAC TAC T) were cloned

into the *Bsm*BI site of the pFgH1tUTG GFP lentiviral vector. Guide RNA oligo

targeting mouse *Atg5* (AAG ATG TGC TTC GAG ATG TG) was cloned into the

*Bsm*BI site of the lentiviral vector lentiCRISPR v2 (Addgene #52961; (28)). For

overexpression studies, the coding sequence of mouse *Txnip* was PCR amplified from

pEGFP-C1-Txnip (Addgene #18758; (29)) with the incorporation of an N-terminal

FLAG-tag. The PCR product was cloned into the *Eco*RI/*Nhe*I sites of a doxycycline

inducible pFTREtight MCS rtTAadvanced GFP lentiviral vector (30).

Antibodies and compounds

Antibodies: LC3B (D11, Cell Signaling Technology #3868), Atg5 (D5F5U, Cell

Signaling Technology #12994); β-actin (AC-15, Sigma #A1978), Glut1 (Abcam

ab15730), Txnip (D5F3E, Cell Signaling Technology #14715S), Glut4 (1F8, Cell

Signaling Technology #2213), Mcl-1 (Rockland 600-401-394S), c-Myc (Abcam

ab32072), Cdk4 (Santa Cruz Biotechnology sc-260), Cyclin D3 (Cell Signaling

Technology #2936), Atf4 (D4B8, Cell Signaling Technology #11815). Compounds:

Mcl-1 inhibitor S63845 was synthesized according to Kotschy et al. (10) by

SYNthesis Med Chem Pty Ltd (Parkville, Australia); DMSO (dimethyl sulfoxide) was

purchased from Sigma (Cat# 472301).

Cell death and cell cycle assays

Cell death and cell cycle were assayed using propidium iodide (PI, Sigma) and flow

cytometry. For death assays, cells were harvested and resuspended in PBS containing

1 μg/ml PI. For cell cycle analysis, cells were harvested and resuspended in PBS

containing 0.1% Triton X-100, 5 μg/ml PI and 10 μg/ml RNaseA (Thermo Scientific),

then incubated at 37 °C for 15 min. Samples were processed on FACSCalibur or

LSRFortessa X-20 flow cytometers (BD Biosciences) and the acquired data analysed

with FlowJo software.

Glucose uptake assays

Extracellular acidification (ECAR) rates were measured in live cells using a Seahorse

Bioscience XFe96 Analyzer (Agilent Technologies). In brief, cells were resuspended

in non-buffered DMEM (Seahorse Biosciences) and 200×10^3 cells were seeded per

well in Seahorse Bioscience 96-well plates. The cellular ECAR was analyzed in non-

buffered DMEM containing 2 mM glutamine with the following compounds: 5 mM

glucose, 1 μM oligomycin and 50 mM 2-deoxy-glucose (2DG). For each assay cycle,

measurement time points of 2 min mix, 2 min wait and 5 min measure were collected.

Data shown for each cell line are representative of at least 2 independent experiments

performed in triplicate.

Acknowledgments

Australian National Health and Medical Research Council (NHMRC) Program Grant

#1113133 and NHMRC fellowship 1020136 to DLV; the Leukemia and Lymphoma

Society SCOR grant #7001-13. Work in the authors' laboratory is made possible by

operational infrastructure grants through the Australian Government Independent

Research Institutes Infrastructure Support (IRISS) and the Victorian State

Government OIS.

References

1.   Perkins AC, Cory S. 1993. Conditional immortalization of mouse

myelomonocytic, megakaryocytic and mast cell progenitors by the Hox-2.4

homeobox gene. EMBO Journal 12:3835-46.

2.   Ekert PG, Read SH, Silke J, Marsden VS, Kaufmann H, Hawkins CJ, Gerl R,

Kumar S, Vaux DL. 2004. Apaf-1 and caspase-9 accelerate apoptosis, but do not

determine whether factor-deprived or drug-treated cells die. J Cell Biol 165:835-42.

3.   Vaux DL, Cory S, Adams JM. 1988. Bcl-2 gene promotes haemopoietic cell

survival and cooperates with c-myc to immortalize pre-B cells. Nature 335:440-442.

4.   Lum JJ, Bauer DE, Kong M, Harris MH, Li C, Lindsten T, Thompson CB.

2005. Growth Factor Regulation of Autophagy and Cell Survival in the Absence of

Apoptosis. Cell 120:237-248.

5.   Mizushima N, Komatsu M. 2011. Autophagy: renovation of cells and tissues.

Cell 147:728-41. doi: 10.1016/j.cell.2011.10.026.

6.   Roy S, Leidal AM, Ye J, Ronen SM, Debnath J. 2017. Autophagy-Dependent

Shuttling of TBC1D5 Controls Plasma Membrane Translocation of GLUT1 and

Glucose Uptake. Mol Cell 6:020.

7.   Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, Shen CH, Wen

J, Asara J, McGraw TE, Kahn BB, Cantley LC. 2013. AMPK-dependent degradation

of TXNIP upon energy stress leads to enhanced glucose uptake via GLUT1. Mol Cell

49:1167-75. doi: 10.1016/j.molcel.2013.01.035. Epub 2013 Feb 28.

8.   Dexter TM, Garland J, Scott D, Scolnick E, Metcalf D. 1980. Growth factor-

dependent hemopoietic precursor cell lines. J Exp Med 152:1036-1047.

9.   Le A, Lane AN, Hamaker M, Bose S, Gouw A, Barbi J, Tsukamoto T, Rojas

CJ, Slusher BS, Zhang H, Zimmerman LJ, Liebler DC, Slebos RJ, Lorkiewicz PK,

Higashi RM, Fan TW, Dang CV. 2012. Glucose-independent glutamine metabolism

via TCA cycling for proliferation and survival in B cells. Cell Metab 15:110-21. doi:

10.1016/j.cmet.2011.12.009.

10.   Kotschy A, Szlavik Z, Murray J, Davidson J, Maragno AL, Le Toumelin-

Braizat G, Chanrion M, Kelly GL, Gong JN, Moujalled DM, Bruno A, Csekei M,

Paczal A, Szabo ZB, Sipos S, Radics G, Proszenyak A, Balint B, Ondi L, Blasko G,

Robertson A, Surgenor A, Dokurno P, Chen I, Matassova N, Smith J, Pedder C,

Graham C, Studeny A, Lysiak-Auvity G, Girard AM, Grave F, Segal D, Riffkin CD,

Pomilio G, Galbraith LC, Aubrey BJ, Brennan MS, Herold MJ, Chang C, Guasconi

G, Cauquil N, Melchiore F, Guigal-Stephan N, Lockhart B, Colland F, Hickman JA,

Roberts AW, Huang DC, Wei AH, et al. 2016. The MCL1 inhibitor S63845 is

tolerable and effective in diverse cancer models. Nature 538:477-482.

11.   Dolezal E, Infantino S, Drepper F, Borsig T, Singh A, Wossning T, Fiala GJ,

Minguet S, Warscheid B, Tarlinton DM, Jumaa H, Medgyesi D, Reth M. 2017. The

BTG2-PRMT1 module limits pre-B cell expansion by regulating the CDK4-Cyclin-

D3 complex. Nat Immunol 18:911-920.

12.   Sawai CM, Freund J, Oh P, Ndiaye-Lobry D, Bretz JC, Strikoudis A, Genesca

L, Trimarchi T, Kelliher MA, Clark M, Soulier J, Chen-Kiang S, Aifantis I. 2012.

Therapeutic targeting of the cyclin D3:CDK4/6 complex in T cell leukemia. Cancer

Cell 22:452-65.

13.   Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C,

Popko B, Paules R, Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. 2003. An

integrated stress response regulates amino acid metabolism and resistance to oxidative

stress. Mol Cell 11:619-33.

14.   Lindqvist LM, Heinlein M, Huang DC, Vaux DL. 2014. Prosurvival Bcl-2

family members affect autophagy only indirectly, by inhibiting Bax and Bak. Proc

Natl Acad Sci U S A 111:8512-7. doi: 10.1073/pnas.1406425111. Epub 2014 May 27.

15.   Datta SR, Dudek H, Tao X, Masters S, Fu HA, Gotoh Y, Greenberg ME.

1997. Akt phosphorylation of bad couples survival signals to the cell-intrinsic death

machinery. Cell 91:231-241.

16.   Zha JP, Harada H, Osipov K, Jockel J, Waksman G, Korsmeyer SJ. 1997. Bh3

domain of bad is required for heterodimerization with bcl-x-l and pro-apoptotic

activity. Journal of Biological Chemistry 272:24101-24104.

17.   Green BD, Jabbour AM, Sandow JJ, Riffkin CD, Masouras D, Daunt CP,

Salmanidis M, Brumatti G, Hemmings BA, Guthridge MA, Pearson RB, Ekert PG.

2013. Akt1 is the principal Akt isoform regulating apoptosis in limiting cytokine

concentrations. Cell Death Differ 20:1341-9. doi: 10.1038/cdd.2013.63. Epub 2013

Jun 21.

18.   Ekert PG, Jabbour AM, Manoharan A, Heraud JE, Yu J, Pakusch M,

Michalak EM, Kelly PN, Callus B, Kiefer T, Verhagen A, Silke J, Strasser A, Borner

C, Vaux DL. 2006. Cell death provoked by loss of interleukin-3 signaling is

independent of Bad, Bim, and PI3 kinase, but depends in part on Puma. Blood

108:1461-8.

19.   O'Neill KL, Huang K, Zhang J, Chen Y, Luo X. 2016. Inactivation of

prosurvival Bcl-2 proteins activates Bax/Bak through the outer mitochondrial

membrane. Genes Dev 30:973-88. doi: 10.1101/gad.276725.115. Epub 2016 Apr 7.

20.   Goodall KJ, Finch-Edmondson ML, van Vuuren J, Yeoh GC, Gentle IE,

Vince JE, Ekert PG, Vaux DL, Callus BA. 2016. Cycloheximide Can Induce Bax/Bak

Dependent Myeloid Cell Death Independently of Multiple BH3-Only Proteins. PLoS

One 11:e0164003. doi: 10.1371/journal.pone.0164003. eCollection 2016.

21.   Callus BA, Mathey-Prevot B. 1998. Interleukin-3-induced activation of the

JAK/STAT pathway is prolonged by proteasome inhibitors. Blood 91:3182-92.

22.   Chao JR, Wang JM, Lee SF, Peng HW, Lin YH, Chou CH, Li JC, Huang HM,

Chou CK, Kuo ML, Yen JJ, Yang-Yen HF. 1998. mcl-1 is an immediate-early gene

activated by the granulocyte-macrophage colony-stimulating factor (GM-CSF)

signaling pathway and is one component of the GM-CSF viability response. Mol Cell

Biol 18:4883-98.

23.   Moulding DA, Quayle JA, Hart CA, Edwards SW. 1998. Mcl-1 expression in

human neutrophils: regulation by cytokines and correlation with cell survival. Blood

92:2495-502.

24.   Brewer JW, Cleveland JL, Hendershot LM. 1997. A pathway distinct from the

mammalian unfolded protein response regulates expression of endoplasmic reticulum

chaperones in non-stressed cells. EMBO J 16:7207-16.

25.   B'Chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi Y, Parry L,

Stepien G, Fafournoux P, Bruhat A. 2013. The eIF2alpha/ATF4 pathway is essential

for stress-induced autophagy gene expression. Nucleic Acids Res 41:7683-99.

26.   Lim JP, Gosavi P, Mintern JD, Ross EM, Gleeson PA. 2015. Sorting nexin 5

selectively regulates dorsal-ruffle-mediated macropinocytosis in primary

macrophages. J Cell Sci 128:4407-19.

27.   Aubrey BJ, Kelly GL, Kueh AJ, Brennan MS, O'Connor L, Milla L, Wilcox S,

Tai L, Strasser A, Herold MJ. 2015. An inducible lentiviral guide RNA platform

enables the identification of tumor-essential genes and tumor-promoting mutations in

vivo. Cell Rep 10:1422-32.

28.   Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide

libraries for CRISPR screening. Nat Methods 11:783-784.

29.   Wang Z, Rong YP, Malone MH, Davis MC, Zhong F, Distelhorst CW. 2006.

Thioredoxin-interacting protein (txnip) is a glucocorticoid-regulated primary response

gene involved in mediating glucocorticoid-induced apoptosis. Oncogene 25:1903-13.

30.   Brumatti G, Salmanidis M, Kok CH, Bilardi RA, Sandow JJ, Silke N, Mason

K, Visser J, Jabbour AM, Glaser SP, Okamoto T, Bouillet P, D'Andrea RJ, Ekert PG.

2013. HoxA9 regulated Bcl-2 expression mediates survival of myeloid progenitors

and the severity of HoxA9-dependent leukemia. Oncotarget 4:1933-47.

Figure Legends

Figure 1. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells shrink and arrest but remain viable when cultured

without IL3.

A. *Bax$^{-/-}$Bak1$^{-/-}$* FDM cells survive long-term in the absence of IL3. *Bax$^{+/+}$Bak1$^{+/+}$*

wild-type (black circles) and *Bax$^{-/-}$Bak1$^{-/-}$* (green squares) FDM cells were cultured

without IL3. At the indicated times, cells were harvested, resuspended in PBS

containing propidium iodide (PI) and analysed by flow cytometry. See Supplemental

Fig. S1A for a representative independent repeat of this experiment.

B. IL3 withdrawal caused cells to shrink. PI negative populations of $Bax^{-/-}Bak1^{-/-}$ cells

cultured for 48 h in the presence (light gray) or absence (dark gray) of IL3, and the

latter that had IL3 returned for 72 h (green), were compared by forward scatter as an

indicator of cell size. Data shown are representative of 2 independent experiments.

C. IL3 withdrawal caused cell cycle arrest, which was reversed by re-addition of IL3.

Nuclei from $Bax^{-/-}Bak1^{-/-}$ cells cultured as described in (B) were stained with PI to

indicate DNA content by flow cytometry. Data shown are representative of 2

independent experiments.

Figure 2. $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ FDM cells shrink and arrest but remain viable when IL3

is withdrawn.

A. Whole cell lysates from $Bax^{-/-}Bak1^{-/-}$ cells and 3 $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ FDM clonal

lines cultured for 48 h in the presence or absence of IL3 were subjected to Western

blotting. The Atg5 immunoblot demonstrates successful CRISPR/Cas9-mediated

deletion of $Atg5$. The LC3B immunoblot indicates greatly reduced LC3B lipidation

(isoform II) in $\Delta Atg5$ cell clones. Data shown are representative of 2 independent

experiments.

B. $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ FDM cells survive long-term in the absence of IL3. The

indicated $Bax^{-/-}Bak1^{-/-}\Delta Atg5$ clones were cultured without IL3 and viability was

determined by PI uptake at the indicated times. See Supplemental Fig. S1B for a

representative independent repeat of this experiment.

C. Cells from *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clone 1 were cultured for 48 h in the presence or

absence of IL3 then processed in PI buffer to measure DNA content by flow

cytometry. Data shown are representative of 2 independent experiments.

D. *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔAtg5* clone 1 cells were cultured for 48 h in the presence or absence

of IL3 and PI negative populations were compared by forward scatter to indicate

differences in cell size. Data shown are representative of 2 independent experiments.

Figure 3. *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* FDM cells arrested by IL3 deprivation survive in the absence

of glucose.

A. Txnip is not required for the decrease in Glut1 following IL3 withdrawal. Whole

cell lysates from *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* and 3 independent *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔTxnip* FDM cell clones

cultured for 48 h in the presence or absence of IL3 were analysed by Western blotting

to detect Glut1 and Txnip protein. Data shown are representative of 2 independent

experiments.

B. Forced expression of Txnip is not sufficient to decrease Glut1 protein levels.

Independent *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* FDM clones bearing a doxycycline (dox)-inducible FLAG-

Txnip construct or empty vector control were cultured for 72 h in the absence or

presence of dox. Whole cell lysates were prepared and analysed by Western blotting

to detect Glut1 and Txnip protein. The data shown represent 1 experiment.

C. *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻* and *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻ΔTxnip* cells can take up and metabolise glucose.

Cells cultured for 72 h in the presence or absence of IL3 were analysed on the

Seahorse bioanalyzer to measure the extracellular acidification rate (ECAR) as an

indicator of glucose metabolism. The increased ECAR following addition of glucose

signifies increased glycolytic flux in cells. Oligomycin inhibits mitochondrial ATP

synthase and the change in ECAR upon its addition represents the glycolytic reserve

capacity of cells. Glycolysis is inhibited by 2-deoxyglucose (2-DG) so upon its

addition, the ECAR values are produced from non-glycolytic sources. Equal numbers

of cells were seeded for analysis. To correct for the reduced size of IL3-starved cells,

protein concentration was assayed and the ECAR values were normalized to 1 μg

total protein per sample. Curves represent the mean of 3 replicate samples depicted as

solid symbols for each time point. Data shown are representative of 2 independent

experiments.

D. $Bax^{-/-}Bak1^{-/-}$ and $Bax^{-/-}Bak1^{-/-}\Delta Txnip$ cells deprived of IL3 survive long-term in the

absence of glucose. Cells were cultured without or with IL3 in complete or glucose-

free medium and analysed for PI uptake by flow cytometry at the indicated times.

Figure 4. When starved of glutamine, $Bax^{+/+}Bak1^{+/+}$ FDM cells undergo apoptosis,

whereas $Bax^{-/-}Bak1^{-/-}$ cells arrest, but can survive for weeks.

A. $Bax^{+/+}Bak1^{+/+}$ cells undergo apoptosis in the absence of glutamine (upper panels),

whereas $Bax^{-/-}Bak1^{-/-}$ cells survive in the absence of glutamine, but arrest even when

IL3 is present (lower panels). Cells were cultured with or without IL3 in glucose- and

glutamine-free medium for 4 days then processed in PI buffer to assess DNA content

by flow cytometry. Note the absence of cells in S phase when $Bax^{-/-}Bak1^{-/-}$ cells were

cultured without glutamine. Data shown are representative of 2 independent

experiments.

B. PI negative populations of $Bax^{-/-}Bak1^{-/-}$ cells cultured with or without IL3 in

medium lacking glucose or glutamine for 4 days (D4) were compared to cells at day 0

(D0) by forward scatter as an indicator of cell size. Data shown are representative of 2

independent experiments.

C. *Bax^{-/-}Bak1^{-/-}* cells survive long-term in the absence of glutamine. Cells were

cultured with or without IL3 in glucose- and glutamine-free medium. At the indicated

times, cells were assayed for viability by PI uptake. See Supplemental Fig. S3E for an

independent repeat of this experiment.

D. *Bax^{-/-}Bak1^{-/-}* cells require glutamine to divide. Cells were cultured with or without

IL3 for 7 days in glucose- and glutamine-free medium. They were then split in three,

and resuspended in medium without glucose and glutamine, or containing only

glucose (1 g/L), or only glutamine (4 mM), for a further 2 days. Cells were harvested

and resuspended in PI buffer to assess DNA content by flow cytometry. Data shown

are representative of 2 independent experiments.

Figure 5. Withdrawal of IL3 or glutamine reduced the expression of Mcl-1 and

several key regulators of cell cycle progression.

A. *Bax^{-/-}Bak1^{-/-}* cells and *Bax^{-/-}Bak1^{-/-}ΔAtg5* clone 1 were cultured for 72 h with or

without IL3 in complete or glucose (Glc)- and glutamine (Gln)-free medium. Whole

cell lysates were prepared and analysed by Western blotting to detect the indicated

proteins. Three independent membranes were used for immunoblotting and β-actin

loading control blots for each are shown. Membrane #1: anti-Mcl-1 and anti-c-Myc;

membrane #2: anti-Cdk4; membrane #3: anti-cyclin D3 and anti-Atf4. Data shown are

representative of 2 independent experiments.

B. *Bax^{+/+}Bak1^{+/+}* and *Bax^{-/-}Bak1^{-/-}* cells cultured in complete medium with IL3 were

treated with DMSO (vehicle control) or the 10 μM of the Mcl-1 inhibitor S63845 for

24 h. Cells were harvested and viability determined by PI uptake and flow cytometry.

Bars depict the mean of 3 replicate cultures represented by the solid symbols. The

data shown represent 1 experiment.

Figure 6. *Bax^{-/-}Bak1^{-/-}* FDM cells require serum and proteasome activity to survive in the absence of glutamine.

A. Even though they were incapable of undergoing Bax/Bak1-dependent apoptosis, *Bax^{-/-}Bak^{-/-}* FDM cells starved of glucose and glutamine died rapidly when cultured in medium with less than 4% fetal bovine serum. *Bax^{-/-}Bak1^{-/-}* cells were cultured with the indicated concentrations of FBS in medium with IL3 but without supplemented glucose and glutamine. Viability was determined by failure to take up PI at the indicated time points as assessed by flow cytometry. The data shown represent 1 experiment.

B. *Bax^{-/-}Bak1^{-/-}ΔAtg5* FDM cells starved of glucose and glutamine died rapidly when cultured in medium with less than 4% fetal bovine serum. *Bax^{-/-}Bak1^{-/-}ΔAtg5* cells were cultured and analyzed for viability as described above in (A). The data shown represent 1 experiment.

Figure 7.

The growth factor IL3 stimulates the expression of Mcl-1 that inhibits activation of Bax/Bak1, and stimulates expression of c-Myc, Cdk4 and cyclin D3 to promote cell growth and proliferation. Glutamine is required to maintain levels of these proteins needed to prevent activation of Bax/Bak1 (Mcl-1), as well as for cell proliferation (c-Myc, Cdk4, cyclin D3 and possibly others). In the absence of growth factor, glucose glutamine and Atg5-dependent autophagy, we hypothesise that *Bax^{-/-}Bak1^{-/-}* myeloid cells survive by consuming proteins present in fetal bovine serum, and degrading these in lysosomes and proteasomes to maintain a sufficient pool of amino acids for energy production.

A

B *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻*

C *Bax⁻ᐟ⁻Bak1⁻ᐟ⁻*

A

B