Title: Auranofin inhibits virulence in *Pseudomonas aeruginosa*

Authors: Leon Zhen Wei Tan[1,2,6], Ziyan Hong[2,6], Joey Kuok Hoong Yam[2,3], May Margarette

Santillan Salido[2], Bau Yi Woo[1], Sam Fong Yau Li[1], Liang Yang[2,4], Michael Givskov[2,5*], Shu-

Sin Chng[1,2*]

Affiliations:

[1]Department of Chemistry, National University of Singapore, Singapore 117543.

[2]Singapore Center for Environmental Life Sciences Engineering (SCELSE), Singapore.

[3]Interdisciplinary Graduate School, Nanyang Technological University, Singapore 637551,

Singapore.

[4]School of Biological Sciences, Nanyang Technological University, Singapore 639798,

Singapore

[5]Costerton Biofilm Center, Department of Immunology and Microbiology, University of

Copenhagen, DK- 2200, Denmark

[6]These authors contributed equally to this work

*To whom correspondence should be addressed.

E-mail: (S.-S.C) chmchngs@nus.edu.sg, (M.G) mgivskov@sund.ku.dk

Abstract

Pseudomonas aeruginosa is widely attributed as the leading cause of hospital-acquired infections. Due to intrinsic antibiotic resistance mechanisms, as well as the ability to form biofilms, *P. aeruginosa* infections are notoriously difficult to treat. *P. aeruginosa* employs multiple virulence mechanisms to establish infections, and the global virulence regulator, Vfr, plays an important role in modulating these mechanisms. An attractive strategy to combat *P. aeruginosa* infections is the use of anti-virulence compounds. Here, we report the identification of auranofin, an FDA-approved drug, that attenuates virulence in *P. aeruginosa*. We present evidence that auranofin exerts its action by targeting Vfr, which in turn leads to inhibition of quorum sensing (QS), Type III secretion (T3SS) and Type IV pili (TFP). Consistent with potent inhibition of QS and TFP expression, we show that auranofin attenuates biofilm maturation, and when used in combination with colistin, displays strong synergy in eradicating *P. aeruginosa* biofilms both in flow chambers and in a murine implant model. Auranofin may have immediate applications as an anti-virulence drug against *P. aeruginosa* infections.

Introduction

Pseudomonas aeruginosa is a prevalent opportunistic pathogen known for its potent virulence, environmental versatility and multi-drug resistance. Its role as the leading cause of hospital-acquired infections (HAIs) among immuno-compromised patients is well documented[1]. Notably, it is one of the main causes of pneumonia in cystic fibrosis (CF) patients[2], and wound infections in surgical patients or burn victims[3]. Furthermore, *P. aeruginosa* has been known to thrive in a variety of ecological niches[4], allowing colonization on abiotic surfaces like catheters and other implanted medical devices[5,6]. Increasing incidences of multi-drug resistant (MDR) *P. aeruginosa* infections[7,8] in the clinic pose a serious threat to global health, underscoring the need to develop new strategies to combat these pathogens.

P. aeruginosa carries several intrinsic antibiotic resistance mechanisms involving the collaboration of restricted uptake through the outer membrane[9], the use of energy-dependent efflux pumps[10,11], and expression of β-lactamases[12,13]. In addition, *P. aeruginosa*, like many other bacteria, exhibits a form of social behavior known as biofilm formation[14], which further increases its tolerance to antibiotics[15]. A biofilm is a community of bacteria encased in a polymeric mesh consisting of extracellular DNA (eDNA), proteins and polysaccharides. In the clinic, significantly higher patient mortality has been associated with the formation of biofilms in the lungs of CF patients[16]. Specifically, the presence of biofilms increases the complexity of treatment due to the function of the extracellular matrix as a physical barrier against, and sponge for, antimicrobial drugs[17]. The small pore size of the biofilm mesh effectively reduces penetration of lysozymes and macromolecular drugs, presumably decreasing the immediate concentrations of these agents surrounding the cells. Additionally, the biofilm matrix is also highly negatively charged due to the presence of eDNA and polysaccharides, which protect the colony from positively-charged aminoglycosides[18]. Consequently, the inhibition of biofilm

formation, maturation and/or function remains a principal consideration in the search for new

treatment options against *P. aeruginosa*.

An attractive alternative strategy to control bacterial infections has been the use of small

molecules that block virulence. Such anti-virulence compounds may reduce the ability of

bacteria to infect the host[19,20], and/or attenuate bacterial defenses against host immune cells[21],

thereby facilitating clearance of the infection site[22]. The virulence of *P. aeruginosa* is multi-

faceted but largely controlled by the global virulence factor regulator, Vfr[23]. Vfr is a cyclic

adenosine monophosphate (cAMP)-dependent transcription factor that coordinates expression

of roughly 200 genes[24,25]. It regulates virulence directly by activating expression of secretion

systems[26] (e.g. Type III) and the pilus formation machinery[27] (Type IV). The Type III secretion

system (T3SS) in *P. aeruginosa* mediates cytotoxicity against host cells through the delivery

of molecular effectors that can disrupt host processes[28]. The Type IV Pilus (TFP) is important

for twitching motility[29], and often associated with the establishment and maturation of *P.*

aeruginosa biofilms. Vfr also controls quorum sensing (QS)[30], a cell-cell communication

network involving the *las-*, *rhl-* and *pqs*-encoded systems, which plays a role in the production

of several virulence factors including extracellular proteases, biosurfactants and toxins[21], and

is important for biofilm maturation[31] and immune evasion. While many small molecules that

inhibit QS and T3SS have been reported[32–34], no inhibitors of Vfr and TFP have been

discovered to date.

In this work, we identified auranofin, an FDA-approved, gold(I)-based anti-

inflammatory drug, as a potent inhibitor of virulence in *P. aeruginosa*. We show that auranofin

inhibits QS, T3SS and TFP formation at low micromolar concentrations. It does this in part by

targeting cysteines in Vfr and abolishing the ability of Vfr to bind DNA. Using flow cell biofilm

assays, we demonstrate that auranofin is able to attenuate biofilm maturation. Furthermore,

auranofin displays synergistic effects in eradicating *P. aeruginosa* biofilms *in vitro*, when used

in combination with colistin. We propose that auranofin can potentially be re-purposed as an

anti-virulent drug to treat *P. aeruginosa* infections.

Results

Auranofin inhibits QS in *Pseudomonas aeruginosa*

We initially discovered auranofin in a screen to identify small molecules that inhibit QS in *P. aeruginosa*. We screened ~6,000 compounds from various libraries using a *lasB-gfp*(ASV) strain[35], where expression of a fusion construct between LasB N-terminus and an unstable GFP variant (GFP(ASV)) is controlled by constitutively expressed LasR, the cognate transcriptional regulator of the *las* pathway. Inhibitors of this pathway should reduce GFP fluorescence in this assay. In addition, since the *las* pathway controls both the *rhl* and *pqs* pathways, compounds identified in this screen could potentially shut down the entire QS network. We found that auranofin exhibits sub-micromolar (IC_{50}) inhibition of LasB-GFP expression with minimal effects on cell growth (Fig. 1a, S1). Auranofin also strongly inhibits fluorescence in two other reporter strains, *rhlA-gfp*(ASV) and *pqsA-gfp*(ASV)[36], both under the control of their respective transcriptional activators, RhlR and PqsR (Fig. 1b&c). To validate our findings, we examined the effects of auranofin on the levels of RNA transcripts for key QS-regulated genes, as well as on the production of various QS-controlled virulence factors. Auranofin results in the down-regulation of *lasA* and *rhlA* transcripts (Fig. S2), and significantly reduces the levels of elastase (LasB), rhamnolipids and pyocyanin (Fig. 2). Taken together, these results establish that auranofin is a potent QS inhibitor.

Auranofin inhibits expression of Type III secretion system (T3SS) and Type IV pili (TFP) in *P. aeruginosa*

To gain insights into the mechanism(s)-of-action of auranofin, we compared the protein profiles of *P. aeruginosa* in the presence and absence of auranofin. Consistent with it being a QS inhibitor, treatment with auranofin caused reduction in the levels of proteins involved in or regulated by QS (Table S1). Interestingly, we also observed down-regulation of proteins

involved in T3SS and TFP formation. We confirmed that T3SS was down-regulated using

immunoblot analyses (Fig. S4). We also showed that auranofin is able to inhibit TFP-dependent

motility, including swarming and twitching[37,38] (Fig. 3). Our observations here are, in part, due

to transcriptional down-regulation of these systems (Fig. S2). These results imply that

auranofin either has multiple targets, and/or it targets a transcriptional regulator upstream of

QS, T3SS and TFP.

Auranofin inhibits Vfr *in vitro* but has other targets in cells

QS, T3SS and TFP formation are positively regulated by Vfr[25]. We asked if auranofin

directly targets Vfr function. To test this, we examined the ability of auranofin to inhibit the

function of Vfr in binding to promoter sequences *in vitro* using electrophoretic mobility shift

assays (EMSA). We first showed that purified Vfr is able to bind DNA fragments containing

the promoter regions of representative QS (*lasR*) and virulence (*toxA*, *regA*) genes (Fig. S3),

as previously reported[39]. We further demonstrated that auranofin effectively prevents Vfr

binding to these promoters in a dose-dependent manner, indicating that it inhibits Vfr function

in vitro (Fig. 4a, c & d). Gold(I)-based compounds are thiophilic, and auranofin has previously

been shown to react covalently with cysteine residues within target proteins[40,41]. To ascertain

if its inhibition of Vfr proceeded via covalent modification of cysteines, we generated a

cysteine-less Vfr variant (VfrΔ5) in which all five cysteines were replaced by serine residues.

This variant is functional as judged by its ability to bind the *lasR, toxA and regA* promoters in

vitro (Fig. S3). Remarkably, auranofin is no longer able to prevent binding of VfrΔ5 to these

same promoters (Fig. 4b, c & d). These results establish that auranofin inhibits Vfr *in vitro* via

its cysteine residues.

To determine if Vfr is the main target of auranofin in cells, we complemented the Δ*vfr*

strain with wild-type Vfr and the VfrΔ5 variant, and examined the levels of elastase, T3SS

expression and twitching motility in the presence and absence of auranofin. Both wild-type Vfr

and the VfrΔ5 variant were able to complement the Δ*vfr* strain in these assays (Fig. S4). We

showed that auranofin inhibits all three phenotypes in the wild-type Vfr complemented strain,

as expected. Surprisingly, auranofin was still able to inhibit these phenotypes in the VfrΔ5

complemented strain, unlike what we have observed *in vitro*. While we have demonstrated that

auranofin does inhibit Vfr via its cysteine residues *in vitro*, our data suggests auranofin may

have other targets, in addition to Vfr.

Auranofin impedes the development of *P. aeruginosa* biofilms

Both TFP and QS are required during biofilm development[42,43]. TFP are important for

initial surface attachment of cells, as well as for the organization of cells into the typical

"mushroom" architecture of mature *P. aeruginosa* biofilms. QS is also critical for biofilm

maturation, although the mechanisms here are less understood. Since auranofin inhibits both

TFP formation and QS networks, we expect it to also have an impact on the development of *P.*

aeruginosa biofilms. To test this, we monitored biofilm formation and maturation in the

presence and absence of auranofin over a period of 72 h in continuous flow chambers. The

formation of fully matured, canonical mushroom-shaped biofilm structures for wild-type *P.*

aeruginosa was readily observed within 72 h (Fig. 5). Interestingly, while auranofin did not

have obvious effects on the initial surface attachment of cells, the formation of mushroom-

shaped structures was less pronounced; biofilms formed in the presence of auranofin appeared

more flat, similar to biofilms of QS or TFP-null strains[44]. Our data clearly indicate that despite

permitting biofilm attachment and growth, auranofin attenuates the progression of biofilm

maturation, possibly by inhibiting QS and TFP.

Auranofin exhibits synergistic effects with colistin in killing *P. aeruginosa* biofilms

Since auranofin affects the maturation of the biofilm, we asked whether it may display synergy with colistin, a last resort drug commonly used to treat *P. aeruginosa* infections. We evaluated the combined effects of auranofin and colistin by monitoring the ratio of live versus dead cells within *P. aeruginosa* flow cell biofilms after treatment. As previously reported, colistin treatment alone results in a sharp decline in cell viability across a 2-day old biofilm, with the exception of several surviving colistin-tolerant subpopulations[44] (Fig. 6a). Remarkably, adding auranofin prior to biofilm formation allows subsequent colistin additions to completely eradicate the *P. aeruginosa* biofilm. In fact, co-treatment of a 2-day biofilm with auranofin and colistin also results in complete killing. These results demonstrate synergy between auranofin and colistin in a flow cell biofilm.

To test whether auranofin and colistin can work together *in vivo*, we surgically inserted silicone implants coated with 1-day old *P. aeruginosa* biofilms into the peritoneal cavity of mice, and subsequently treated them locally with colistin, auranofin, or a combination of both drugs. Remarkably, combination therapy of auranofin and colistin resulted in significantly lower bacterial titer counts from the implant than either auranofin or colistin alone (Fig. 6b). This confirms the synergy between auranofin and colistin in a murine model.

Discussion

Auranofin is approved by the FDA for the treatment of rheumatoid arthritis. It has been shown to reduce inflammation of joints in rheumatic patients[45]; however, the precise mechanism of how this occurs is not clear. Auranofin also exhibits anti-cancer and anti-parasitic properties[40,46,47]. Furthermore, auranofin is able to kill a broad spectrum of Gram-positive bacteria, and also mycobacteria[41]. All of these activities may be attributed to auranofin's ability to disrupt the thiol-redox balance in cells, in part by targeting the thioredoxin reductases in each of these systems[48]. Interestingly, auranofin is largely ineffective against most Gram-negative bacteria, presumably because of lack of penetration of the bacterial outer membrane[49]. Here, we have demonstrated that auranofin is additionally able to inhibit virulence in *P. aeruginosa*. Despite not being able to accumulate at high concentrations in cells, auranofin causes downregulation of QS, T3SS and TFP formation. One possible target is Vfr, which controls all of these phenotypes. Consistently, we have shown that auranofin inhibits the ability of Vfr to bind to target promoter sequences in vitro. The fact that a cysteine-less Vfr variant is not inhibited in vitro indicates that auranofin also targets cysteines within this protein. Interestingly, *P. aeruginosa* does not gain resistance against auranofin even when expressing cysteine-less Vfr, suggesting that it has multiple targets in cells. This is probable, given that auranofin is a general thiol-targeting reagent[40]. Understanding the mechanism of action of auranofin would require identifying these unknown targets.

Current clinical treatment options of *P. aeruginosa* infections involve the usage of carbapenems, aminoglycosides and ciprofloxacin, to name a few[16]. These drugs are often used in combination or singularly in treatment. Unfortunately, many strains of *P. aeruginosa* have developed resistance to such compounds and drastically reduce the pool of available treatment options for clinicians[7]. Colistin remains one of the few drugs effective against most strains of *P. aeruginosa*. Its usage, however, remains limited and is confined to last resort situations due

to its toxicity[50]. We have shown that auranofin and colistin combine synergistically to

completely eradicate a *P. aeruginosa* biofilm, something not observed with colistin treatment

alone. This synergy is also observed in a murine implant model. We believe that the synergy

between auranofin and colistin works in both directions. It has been shown that disrupting outer

membrane permeability (here with the use of colistin) allows auranofin to penetrate the cell

envelope and kill Gram-negative bacteria, through its potential action on thioredoxin reductase.

At the same time, the inhibition of TFP and QS (by auranofin) during biofilm development

may prevent the formation of colistin-tolerant subpopulations within the biofilm, thereby

allowing complete eradication. The unique synergistic effects of auranofin and colistin may be

exploited as a new treatment regimen for *P. aeruginosa* infections. Given that auranofin is

already an FDA-approved drug, it may not be in the very far future that this approach can be

tested.

Materials and Methods

Detailed methods can be found in SI Appendix, Materials and Methods. All bacterial strains

and plasmids are summarized in Table S2 and S3. Oligonucleotides for PCR are listed in Table

S4.

Acknowledgements

We thank Young-Tae Chang (POSTECH) for allowing access to small molecule libraries used

in this study. We acknowledge Song Lin Chua (SCELSE) for his help in processing and

submitting our samples for proteomics analysis. We thank Yingying Li (SCELSE) for her

guidance in RT-PCR experiments. We are grateful to Arne Rietsch (Case Western Reserve

University) and Katrina Forest (University of Wisconsin-Madison) for their generous gifts of

the α−PcrV and α−Vfr antibodies, respectively. This research was supported by the National

Research Foundation and the Ministry of Education of Singapore under its Research Centre of

Excellence Programme.

Contributions

L.Y., M.G., and S.-S.C. conceived the project and directed the study. Z.H. conducted the QS inhibitor screening assay. L.Z.W.T. conducted all biochemical experiments. J.K.H.Y. and M.M.S.S. performed the mouse model and the flow cell biofilm experiments. B.Y.W. conducted the rhamnolipid analysis with supervision from S.F.Y.L.. L.Z.W.T. and S.-S.C. wrote the manuscript with input from all authors.

Competing financial interests

M.G., S.-S.C., L.Y., L.Z.W.T., and Z.H. have filed an international patent application (PCT/SG2016/050436, filed 6th September 2016) on the work described in this manuscript.

References

1.   Vincent, J.-L. The Prevalence of Nosocomial Infection in Intensive Care Units in Europe. *JAMA* **274,** 639 (1995).

2.   Palmer, K. L., Mashburn, L. M., Singh, P. K. & Whiteley, M. Cystic fibrosis sputum supports growth and cues key aspects of Pseudomonas aeruginosa physiology. *J. Bacteriol.* **187,** 5267–77 (2005).

3.   Church, D., Elsayed, S., Reid, O., Winston, B. & Lindsay, R. Burn wound infections. *Clin. Microbiol. Rev.* **19,** 403–34 (2006).

4.   L M Ringen, C. H. D. A study of the incidence of Pseudomonas aeruginosa from various natural sources. *J. Bacteriol.* **XVI,** 841–845 (1952).

5.   Cole, S. J., Records, A. R., Orr, M. W., Linden, S. B. & Lee, V. T. Catheter-associated urinary tract infection by Pseudomonas aeruginosa is mediated by exopolysaccharide-independent biofilms. *Infect. Immun.* **82,** 2048–58 (2014).

6.   Mittal, R. *et al.* Urinary tract infections caused by Pseudomonas aeruginosa: a

minireview. *J. Infect. Public Health* **2,** 101–11 (2009).

7.   Hirsch, E. B. & Tam, V. H. Impact of multidrug-resistant Pseudomonas aeruginosa

infection on patient outcomes. *Expert Rev. Pharmacoecon. Outcomes Res.* **10,** 441–51

(2010).

8.   Dilip Nathwani, Gowri Raman, Katherine Sulham, M. G. and V. M. Clinical and

economic consequences of hospital-acquired resistant and multidrug-resistant

Pseudomonas aeruginosa infections: a systematic review and meta-analysis. *Antimicrob.*

Resist. Infect. Control **3,** R97 (2014).

9.   Nikaido, H. Molecular basis of bacterial outer membrane permeability revisited.

Microbiol. Mol. Biol. Rev. **67,** 593–656 (2003).

10. Li, X. Z., Livermore, D. M. & Nikaido, H. Role of efflux pump(s) in intrinsic resistance

of Pseudomonas aeruginosa: resistance to tetracycline, chloramphenicol, and

norfloxacin. *Antimicrob. Agents Chemother.* **38,** 1732–1741 (1994).

11. Poole K Krebes K, M. C. N. S. Multiple antibiotic resistance in Pseudomonas aeruginosa:

evidence for involvement of an efflux operon. *J. Bacteriol.* **175,** 7363–7372 (1993).

12. Sanders, C. C. & Sanders, W. E. B-Lactam Resistance in Gram-Negative Bacteria:

Global Trends and Clinical Impact. *Clin. Infect. Dis.* **15,** 824–39 (1992).

13. Okamoto, K., Gotoh, N. & Nishino, T. Pseudomonas aeruginosa reveals high intrinsic

resistance to penem antibiotics: penem resistance mechanisms and their interplay.

Antimicrob. Agents Chemother. **45,** 1964–71 (2001).

14. Costerton, J. W. Bacterial Biofilms: A Common Cause of Persistent Infections. *Science*

(80-.). **284,** 1318–1322 (1999).

15. Høiby, N., Bjarnsholt, T., Givskov, M., Molin, S. & Ciofu, O. Antibiotic resistance of

bacterial biofilms. *Int. J. Antimicrob. Agents* **35,** 322–332 (2010).

16. Driscoll, J. A., Brody, S. L. & Kollef, M. H. The epidemiology, pathogenesis and

treatment of Pseudomonas aeruginosa infections. *Drugs* **67,** 351–68 (2007).

17.   Lewis, K. Riddle of biofilm resistance. *Antimicrob. Agents Chemother.* **45,** 999–1007

(2001).

18.   Chiang, W.-C. *et al.* Extracellular DNA shields against aminoglycosides in

Pseudomonas aeruginosa biofilms. *Antimicrob. Agents Chemother.* **57,** 2352–61 (2013).

19.   Imperi, F., Leoni, L. & Visca, P. Antivirulence activity of azithromycin in Pseudomonas

aeruginosa. *Front. Microbiol.* **5,** 178 (2014).

20.   Maisuria, V. B., Los Santos, Y. L., Tufenkji, N., Déziel, E. & Lazdunski, A. Cranberry-

derived proanthocyanidins impair virulence and inhibit quorum sensing of Pseudomonas

aeruginosa. *Sci. Rep.* **6,** 30169 (2016).

21.   Alhede, M. *et al.* Pseudomonas aeruginosa recognizes and responds aggressively to the

presence of polymorphonuclear leukocytes. *Microbiology* **155,** 3500–3508 (2009).

22.   Hentzer, M. *et al.* Attenuation of Pseudomonas aeruginosa virulence by quorum sensing

inhibitors. *EMBO J.* **22,** 3803–3815 (2003).

23.   Suh, S.-J. *et al.* Effect of vfr mutation on global gene expression and catabolite

repression control of Pseudomonas aeruginosa. *Microbiology* **148,** 1561–9 (2002).

24.   Wolfgang, M. C., Lee, V. T., Gilmore, M. E. & Lory, S. Coordinate regulation of

bacterial virulence genes by a novel adenylate cyclase-dependent signaling pathway.

Dev. Cell **4,** 253–63 (2003).

25.   Coggan, K. a. & Wolfgang, M. C. Global regulatory pathways and cross-talk control

pseudomonas aeruginosa environmental lifestyle and virulence phenotype. *Curr. Issues*

Mol. Biol. **14,** 47–70 (2012).

26.   Davinic, M., Carty, N. L., Colmer-Hamood, J. A., San Francisco, M. & Hamood, A. N.

Role of Vfr in regulating exotoxin A production by Pseudomonas aeruginosa.

Microbiology **155,** 2265–73 (2009).

27. Beatson, S. A., Whitchurch, C. B., Sargent, J. L., Levesque, R. C. & Mattick, J. S. Differential regulation of twitching motility and elastase production by Vfr in Pseudomonas aeruginosa. *J. Bacteriol.* **184,** 3605–13 (2002).

28. Wanyin Deng, Natalie C. Marshall, Jennifer L. Rowland, James M. McCoy, Liam J. Worrall, Andrew S. Santos, N. C. J. S. & B. B. F. Assembly, structure, function and regulation of type III secretion systems. *Nat. Rev. Microbiol.* **15,** 323–337 (2017).

29. Mattick, J. S. Type IV Pili and Twitching Motility. *Annu. Rev. Microbiol.* **56,** 289–314 (2002).

30. Albus, A. M., Pesci, E. C., Runyen-Janecky, L. J., West, S. E. & Iglewski, B. H. Vfr controls quorum sensing in Pseudomonas aeruginosa. *J. Bacteriol.* **179,** 3928–35 (1997).

31. Rutherford, S. T. & Bassler, B. L. Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb. Perspect. Med.* **2,** a012427 (2012).

32. Mcshan, A. C. & De Guzman, R. N. The Bacterial Type III Secretion System as a Target for Developing New Antibiotics. doi:10.1111/cbdd.12422

33. Jakobsen, T. H. *et al.* Food as a source for quorum sensing inhibitors: Iberin from horseradish revealed as a quorum sensing inhibitor of Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* **78,** 2410–2421 (2012).

34. Duncan, M. C., Linington, R. G. & Auerbuch, V. Chemical inhibitors of the type three secretion system: disarming bacterial pathogens. *Antimicrob. Agents Chemother.* **56,** 5433–41 (2012).

35. Hentzer, M. *et al.* Inhibition of quorum sensing in Pseudomonas aeruginosa biofilm bacteria by a halogenated furanone compound. *Microbiology* **148,** 87–102 (2002).

36. Yang, L. *et al.* Computer-aided identification of recognized drugs as Pseudomonas aeruginosa quorum-sensing inhibitors. *Antimicrob. Agents Chemother.* **53,** 2432–2443 (2009).

333   37.   Anyan, M. E. *et al.* Type IV pili interactions promote intercellular association and

334   moderate swarming of Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **111,**

335   18013–8 (2014).

336   38.   Burrows, L. L. Twitching Motility: Type IV Pili in Action. *Annu. Rev. Microbiol.* **66,**

337   493–520 (2012).

338   39.   Fuchs, E. L. *et al.* The Pseudomonas aeruginosa Vfr regulator controls global virulence

339   factor expression through cyclic AMP-dependent and -independent mechanisms. *J.*

340   *Bacteriol.* **192,** 3553–64 (2010).

341   40.   Di Sarra, F., Fresch, B., Bini, R., Saielli, G. & Bagno, A. Reactivity of Auranofin with

342   Selenols and Thiols - Implications for the Anticancer Activity of Gold(I) Compounds.

343   *Eur. J. Inorg. Chem.* **2013,** 2718–2727 (2013).

344   41.   Harbut, M. B. *et al.* Auranofin exerts broad-spectrum bactericidal activities by targeting

345   thiol-redox homeostasis. *Proc. Natl. Acad. Sci.* **112,** 4453–4458 (2015).

346   42.   Rodesney, C. A. *et al.* Mechanosensing of shear by Pseudomonas aeruginosa leads to

347   increased levels of the cyclic-di-GMP signal initiating biofilm development. *Proc. Natl.*

348   *Acad. Sci. U. S. A.* 201703255 (2017). doi:10.1073/pnas.1703255114

349   43.   Solano, C., Echeverz, M. & Lasa, I. Biofilm dispersion and quorum sensing. *Curr. Opin.*

350   *Microbiol.* **18,** 96–104 (2014).

351   44.   Chua, S. L. *et al.* Selective labelling and eradication of antibiotic-tolerant bacterial

352   populations in Pseudomonas aeruginosa biofilms. *Nat. Commun.* **7,** 10750 (2016).

353   45.   Finkelstein, A. E. *et al.* Auranofin. New oral gold compound for treatment of rheumatoid

354   arthritis. *Ann. Rheum. Dis.* **35,** 251–7 (1976).

355   46.   Debnath, A. *et al.* A high-throughput drug screen for Entamoeba histolytica identifies a

356   new lead and target. *Nat. Med.* **18,** 956–60 (2012).

47.   Park, S.-H., Lee, J., Berek, J. & Hu, M. Auranofin displays anticancer activity against

ovarian cancer cells through FOXO3 activation independent of p53. *Int. J. Oncol.* **45,**

1691–8 (2014).

48. Becker, K., Gromer, S., Schirmer, R. H., Mu, S. & Ller, È. Thioredoxin reductase as a

pathophysiological factor and drug target. *Eur. J. Biochem.* **267,** 6118–6125 (2000).

49. Thangamani, S. *et al.* Antibacterial activity and mechanism of action of auranofin

against multi-drug resistant bacterial pathogens. *Sci. Rep.* **6,** 22571 (2016).

50. Ordooei Javan, A., Shokouhi, S. & Sahraei, Z. A review on colistin nephrotoxicity. *Eur.*

J. Clin. Pharmacol. **71,** 801–810 (2015).

Figure 1. Auranofin inhibits all levels of QS with minimal effects on cell growth.

Representative fluorescence induction profiles (relative fluorescence units normalized to cell

density, over time) for (a) LasR-mediated expression of LasB-GFP, (b) RhlR-mediated

expression of P_{rhlA}-*gfp*, and (c) PqsR-mediated expression of P_{pqsA}-*gfp*, in the presence of

indicated concentrations of auranofin, whose chemical structure is shown on the top. Dose-

dependent inhibition curves for each assay are plotted on the right. Error bars represent

standard deviation from technical triplicates.

Figure 2. The production of key QS-controlled virulence factors is reduced in the presence of

auranofin. (a) Elastase activity in the supernatants of PAO1 and Δ*lasIrhlI* cultures treated with

indicated concentrations of auranofin. (b) Relative abundance of rhamnolipid fragments

isolated from PAO1 cells treated with 20 μM auranofin, as quantified using multiple reaction

monitoring mass spectrometry. The chemical structure of rhamnolipid and its measured

fragments are shown on the right. (c) Relative levels of pyocyanin in supernatants of PAO1

and Δ*lasIrhlI* cultures treated with indicated concentrations of auranofin, as measured at

absorbance at 691 nm. Error bars represent standard deviation from biological triplicates.

Figure 3. Motility phenotypes that depend on TFP (i.e. swarming and twitching) are reduced

in the presence of auranofin. (A) Swarming (0.5% LB agar), (b) swimming (0.3% LB agar),

and twitching (1% LB agar) motilities of PAO1 and PAO1 Δ*vfr* strains in the presence or

absence of 20 μM auranofin. For twitching, cells at the base of the plates were stained with

Coomassie blue.

Figure 4. Auranofin inhibits Vfr binding to target promoter sequences in vitro by reacting with

cysteine residues. Polyacrylamide gels showing specific binding of (a) Vfr, (b) VfrΔ5 to P_{lasR},

or both proteins to (c) P_{toxA} and (d) P_{regA}, in the presence of indicated concentrations of

auranofin. The coding region of the gene *rsaL* is used as a probe to control for non-specific

binding.

Figure 5. Auranofin attenuates the progression of *P. aeruginosa* biofilms. Confocal

fluorescence microscopy images illustrating the formation and maturation of PAO1-*gfp*

biofilms over three days in the presence or absence of 1 μM under continuous flow conditions.

Scale bar = 20 μm

Figure 6. Auranofin displays strong synergy in eradicating *P. aeruginosa* biofilms when used

in combination with colistin. (a) Confocal fluorescence microscopy images of two-day old

PAO1-*gfp* biofilms before and after treatment with 10 μg/ml colistin. Auranofin was not added

(top panels), added prior to biofilm formation (middle panels), or added together with colistin

(bottom panels). Dead cells were revealed by staining with propidium iodide. Each panel

represents a square surface area of 212 x 212 μm. (b) Mean c.f.u per ml of *in vivo* PAO1

biofilms obtained from silicone implants in the peritoneal space of mice, treated with auranofin,

colistin or a combination of both drugs. *$P<0.01$, Student's *t*-test.

Figure 1

Figure 2

Figure 3

   Figure 4

a

→Vfr/P*lasR* complex

rsaL
←P*lasR*

P*lasR*	+	+	+	+	+	+	+	+	+	+
rsaL	-	+	+	+	+	+	+	+	+	+
Vfr, µM	-	-	1	1	1	1	1	1	1	1
Auf, µM	-	-	-	100	50	25	12.5	6.25	3.13	1.56

b

→VfrΔ5/P*lasR* complex

rsaL
←P*lasR*

P*lasR*	+	+	+	+	+	+	+	+	+	+
rsaL	-	+	+	+	+	+	+	+	+	+
Vfr, µM	-	-	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1
Auf, µM	-	-	-	100	50	25	12.5	6.25	3.13	1.56

c

→Vfr(Δ5)/P*toxA* complex

rsaL
←P*toxA*

P*toxA*	+	+	+	+	+	+	+	+	+	+
rsaL	-	+	+	+	+	+	+	+	+	+
Vfr, µM	-	-	1	1	1	1	3.1	3.1	3.1	3.1
Auf, µM	-	-	25	12.5	6.25	-	25	12.5	6.25	

d

→Vfr(Δ5)/P*regA* complex

P*regA*
←*rsaL*

P*regA*	+	+	+	+	+	+	+	+	+	+
rsaL	-	+	+	+	+	+	+	+	+	+
Vfr, µM	-	-	1	1	1	1	3.1	3.1	3.1	3.1
Auf, µM	-	-	25	12.5	6.25	-	25	12.5	6.25	

419   **Figure 5**

421 **Figure 6**

422