A hybrid versatile method for state estimation and feature extraction from the trajectory

of animal behavior

Authors: Shuhei J. Yamazaki[1,2], Kazuya Ohara[3], Kentaro Ito[4], Nobuo Kokubun[4,5], Takuma

Kitanishi[6,7,8], Daisuke Takaichi[9], Yasufumi Yamada[10], Yosuke Ikejiri[1,2], Fumie Hiramatsu[1],

Kosuke Fujita[1,14], Yuki Tanimoto[1,15], Akiko Yamazoe-Umemoto[1], Koichi Hashimoto[11],

Katsufumi Sato[12], Ken Yoda[13], Akinori Takahashi[4,5], Yuki Ishikawa[9], Azusa Kamikouchi[9],

Shizuko Hiryu[10], Takuya Maekawa[3], Koutarou D. Kimura[1,2*]

Affiliations

[1]Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

[2]Graduate School of Natural Sciences, Nagoya City University, Nagoya, Aichi, 467-8501, Japan

[3]Graduate School of Information Science and Technology, Osaka University, Suita, Osaka 565-

0871, Japan

[4]Department of Polar Science, SOKENDAI, Tachikawa, Tokyo 190-8518, Japan

[5]National Institute of Polar Research, Tachikawa, Tokyo 190-8518, Japan

[6]Department of Physiology, Osaka City University Graduate School of Medicine, Osaka 545-

8585, Japan

[7]Center for Brain Science, Osaka City University Graduate School of Medicine, Osaka 545-

8585, Japan

[8]PRESTO, Japan Science and Technology Agency (JST), Kawaguchi, Saitama 332-0012, Japan.

[9]Graduate School of Science, Nagoya University, Chikusa, Nagoya, Aichi, 464-8602, Japan

[10]Faculty of Life and Medical Sciences, Doshisha University, Kyotanabe, Kyoto 6100321, Japan

[11]Graduate School of Information Sciences, Tohoku University, Sendai, Miyagi 980-8579, Japan

[12]Atmosphere and Ocean Research Institute, The University of Tokyo, 5-1-5 Kashiwanoha,

Kashiwa, Chiba 277-8564, Japan

[13]Graduate School of Environmental Studies, Nagoya University, Nagoya, Aichi 464-8601,

Japan

[14]Present address: Department of Ophthalmology, Graduate School of Medicine, Tohoku

University, Sendai, Miyagi 980-8574, Japan

[15]Present address: RIKEN Center for Brain Science, Wako, Saitama 351-0198, Japan

*Correspondence: kokimura-lab@umin.ac.jp or kokimura@nsc.nagoya-cu.ac.jp

Manuscript information:

Abstract: 224 words

Main body: 8995 words

Figures & Tables: 12

Supplementary file: 1

ABSTRACT

Animal behavior is the final and integrated output of the brain activity. Thus, recording and analyzing behavior is critical to understand the underlying brain function. While recording animal behavior has become easier than ever with the development of compact and inexpensive devices, detailed behavioral data analysis requires sufficient previous knowledge and/or high content data such as video images of animal postures, which makes it difficult for most of the animal behavioral data to be efficiently analyzed to understand brain function. Here, we report a versatile method using a hybrid supervised/unsupervised machine learning approach to efficiently estimate behavioral states and to extract important behavioral features only from low-content animal trajectory data. As proof of principle experiments, we analyzed trajectory data of worms, fruit flies, rats, and bats in the laboratories, and penguins and flying seabirds in the wild, which were recorded with various methods and span a wide range of spatiotemporal scales—from mm to 1000 km in space and from sub-seconds to days in time. We estimated several states during behavior and comprehensively extracted characteristic features from a behavioral state and/or a specific experimental condition. Physiological and genetic experiments in worms revealed that the extracted behavioral features reflected specific neural or gene activities. Thus, our method provides a versatile and unbiased way to extract behavioral features from simple trajectory data to understand brain function.

INTRODUCTION

The brain receives, integrates, and processes a range of ever-changing environmental information to produce relevant behavioral outputs. Therefore, understanding salient behavioral features can augment our understanding of important aspects of environmental information as well as of brain activity, which links the environmental information to behavior. Recent technological development of compact and inexpensive cameras and/or global positioning system (GPS) devices has facilitated convenient monitoring and recording of animal behavior (Brown and de Bivort, 2018; Dell et al., 2014; Egnor and Branson, 2016). However, the behavioral data generated through these approaches are frequently represented as a few simple measures, such as velocity, migratory distance, or the probability of reaching a particular goal, due to the challenges related to identification of specific aspects of behavior to be analyzed; in other words, it is still difficult to figure out how we can describe an animal behavior meaningfully (Berman, 2018). Owing to poor description of behavior, dynamic neural activity, for example, is not sufficiently interpreted even though simultaneous optical monitoring can measure a large number of time-series neural activities (Alivisatos et al., 2012; Landhuis, 2017). This large asymmetry in data richness between neural activity and behavior has emerged as one of the most significant issues in modern neuroscience (Anderson and Perona, 2014; Gomez-Marin et al., 2014; Krakauer et al., 2017).

One way to overcome the challenges in the appropriate descriptions of behavior is to describe its salient features via comprehensive analysis through an approach such as machine learning. Machine learning involves extracting latent patterns and uncovering knowledge from a large amount of data (Bishop, 2006). In fact, multiple behavioral analysis methods based on machine learning have been reported in the last decade (Baek et al., 2002; Branson et al., 2009; Brown et al., 2013; Dankert et al., 2009; Kabra et al., 2013; Mathis et al., 2018; Robie et al., 2017; Stephens et al., 2008; Vogelstein et al., 2014; Wiltschko et al., 2015). Most of these studies

have classified behavioral states based on detailed analyses of animal postures as observed in

video images (Dell et al., 2014); the classification of behavioral states into classes, such as

foraging, sleeping, chasing, or fighting, is considered to be critical for efficient behavioral

analysis, as each of behavioral features varies differently across different behavioral states

(Egnor and Branson, 2016; Jonsen et al., 2013; Patterson et al., 2008). Although these methods

have worked successfully for the analysis of behavioral videos of worms, fruit flies, and rodents

in laboratories, they have some limitations. First, these methods are not suitable for analyzing

relatively long-distance navigation given their requirement of recording reasonably large and

detailed images of animals in the video frame. Second, the extraction of behavioral features

from a state, as opposed to just state classification, is more critical in understanding how

environmental information and/or brain activities trigger transitions among states for behavioral

response.

To analyze relatively long-distance navigation behavior comprehensively, we developed a

method for the estimation of behavioral states and extraction of relevant behavioral features

based only on the trajectories of animals. For estimating behavioral states, we used an

unsupervised learning method involving the expectation maximization (EM) algorithm

(Dempster et al., 1977) because it is difficult for the human eye to classify behavior into distinct

states without using posture images. For extracting salient behavioral features, we used

information gain, an index used for a supervised learning method (the decision tree analysis)

(Quinlan, 1992), and compared the features between two different experimental conditions (e.g.,

with or without certain stimulus). It is because supervised learning is considered advantageous

in the extraction of characteristic behavioral features and comparing them among multiple

conditions. We named this hybrid supervised/unsupervised machine learning approach as the

*st*ate *e*stimation and *f*eature ex*tr*action (STEFTR) method (Fig. 1).

Because the STEFTR method only uses trajectory information for the analysis, it becomes

possible to analyze movement behavior of various animals regardless of the spatiotemporal

scale of movement. As proof-of-principle experiments, we analyzed the trajectories of worms,

flies, rats, and bats in laboratories and those of penguins and flying seabirds in the wild; these

experiments involved a spatiotemporal scale ranging from mm to 1000 km in space and from

sub-seconds to days in time. The behavioral states of worms and penguins estimated by the

STEFTR method were in reasonable conformation with the ones described in previous

literature, supporting the reliability of our method. We further extracted learning-dependent

behavioral features from a behavioral state of worms, in which one of the behavioral features is

correlated with learning-dependent changes in neural activities. We also analyzed the behavioral

features of mutant strains of worms and found that the patterns of features are correlated with

gene function, suggesting that comprehensive feature extraction may enable us to estimate

unknown functions of a gene product. We were also able to extract learning-dependent features

from bats and pheromone-dependent features from fruit flies. Taken together, our findings

indicate that the STEFTR method allows us to estimate internal state, neural activity, and gene

function related to animal behavior only from movement trajectories, regardless of the recording

method or the spatiotemporal scales.

MATERIAL AND METHODS

Estimation of behavioral states

For the analysis of trajectory information of an animal obtained from video images or from the

GPS device attached to an animal, approximately 1/1,000 and 1/100 of the median recording

time across animals were used as an unit for time frame and the time window for moving

average, respectively (Table 1). These values were used to draw the 8 histograms of the

averages (*Ave*) and the variances (*Var*) of velocity (*V*), bearing (*B*), time-differential of *V* (*dV*)

and *B* (*dB*) as the basic behavioral features. The time window for moving average was critical to

reduce noise and to detect relatively long trends of behavior. From the histograms, a basic

behavioral feature appeared to include multiple normal distributions was selected, and the

number of clusters and boundaries of each cluster were automatically determined by EM

algorithm (see below). In the case of worms, the cluster analysis was performed with specifying

the maximum number cluster of 20. In other cases, maximum cluster number 5 was

predetermined based on the knowledge that the number of basic behavioral states are several in

general (Patterson et al., 2008).

The EM algorithm assigns a cluster label to each time frame although the clustering results,

which reflect behavioral states of an animal, should be smooth in time. To smooth out the

clustering results and removing outlying results, moving average was again applied to the

cluster labels, which resulted in clusters resemble to the human-labeled behavioral states.

When the value of a behavioral feature changes suddenly and largely, the influence of the

change may extend over a wide range. For example, if an animal moving straightly initiates

local search suddenly, dB value will be 0°, 0°, 0°, 0°, 0°, 0°, 180°, 0°, 90°, 0°, 270°, etc. If

moving average with ±5 time frame was applied, the value change occur from -5 time frame of

the sudden value change, which should be compensated. Because worm's clusters 0 and 1

corresponded to this case, the beginning and the end of each cluster 0 was extended by the half

of time window.

The cluster labels obtained as described above were mapped to the corresponding trajectory

position with colors. We used a custom-made python program for calculating basic behavioral

features, Weka data mining software (the University of Waikato, New Zealand) (Frank et al.,

2016) for EM calculation, and Excel (Microsoft) for others.

EM algorithm for cluster analysis

A set of values of the ith basic behavioral feature F_i, which were extracted from trajectories of interest, and the number of clusters N were given. We employed the EM algorithm to cluster F_i into N clusters, i.e., a mixture of N Gaussians. The probability distribution of the Gaussian mixtures M_N is represented as follows:

$$p(f_i|M_N) = \sum_{n=1}^{N} \pi_n N(f_i, \mu_n, \sigma_n),$$

where f_i is a feature value of the ith feature, π_n is the mixture weight of the nth Gaussian, μ_n is the mean of the nth Gaussian, and σ_n is the standard deviation of the nth Gaussian. Thus, the EM algorithm was used to estimate the cluster parameters: π_n, μ_n, and σ_n.

Determination of cluster number using information criteria

The method evaluates a set of clusters (model M_N) obtained by the EM algorithm using information criteria, and finds the best N using the following criteria.

$$AIC(M_N) = -2 \ln L\big(p(F_i|M_N)\big) + 2N,$$

where $L\big(p(F_i|M_N)\big)$ is the maximum likelihood of F_i under model M_N and N is the penalty term that measures model complexity.

Feature extraction with information gain

We leverage information gain to evaluate the classification ability of each feature, i.e., its ability to identify a characteristic of a state (cluster). Information entropy is used to compute the ambiguity of a set of data points according to the following formula:

$$H = -\sum_{n=1}^{N} p_n \log_2 p_n,$$

where p_n is the number of data points belonging to the nth class (cluster). Given that we classify all the data points into two groups (i.e. two experimental conditions) using a particular threshold related to a specific feature, the feature is considered to be a characteristic feature (in that it classifies the data points well) if the ambiguity within the two groups is lower than that of

the original data set. Thus, we evaluated features in terms of their ability in ambiguity reduction

upon classification (information gain).

For worm odor avoidance behavior, we extracted behavioral features that have positive

information gain in naive versus pre-exposed worms or in mock-treated versus pre-exposed

worms. Next, we chose the extracted features that were common for both comparisons; these

features were termed as "features modulated in a learning-dependent manner." For flies,

behavioral features were compared between with or without pheromone tapping. For bats,

behavioral features were compared between unfamiliar flights (1^{st}-3^{rd}) and familiar (10^{th}-12^{th})

flights. The Weka software was used for these calculations.

Behavioral parameters included in a feature vector

For the machine learning analysis of worm' odor avoidance behavior, the following behavioral

features were calculated for each cluster 0 segment from the coordinates of the centroid of the

trajectory: velocity (V), bearing (B), odor concentration the worm experienced during the run

(C), the time differential values for these (dV, dB, and dC), directedness (Dir) (Gorelik and

Gautreau, 2014), curvature (called weathervane; WV) (Iino and Yoshida, 2009), and durations

of cluster 0 and 1 (*Clst0Dur* and *Clst1Dur*, respectively). For V, dV, B, dB, C, dC, and Dir, the

average (*Ave*) and median (*Med*) values for at the initiation (*Ini*), middle (*Mid*), termination

(*Ter*), and all (*All*) periods of a cluster 0 segment were calculated. A total of 333 features was

calculated by combining all these features.

For analyzing changes in the flight of bats, the following behavioral features in each flight were

calculated from the coordinates of the bats and obstacles: three-dimensional flight velocity (V),

horizontal and vertical bearings of the flight (*B_hori* and *B_vert*, respectively), distance (*R_obs*)

and bearing (*B_obs*) of the bat to the nearest edge point of the obstacle chain array, longitudinal

directional distance to the frontal chain array (R_x), and lateral directional distance to the inside

pitch of the chain array (R_y). Time-differential values were calculated for V (dV), B (dB), dB

(ddB), and the flight height (dH), which were calculated with frame units of the high-speed

video cameras (1/125 s). All flight trajectories were divided into three segments: early, middle,

and late terms. The time window for the analysis of each behavioral feature was 0.1 s, 0.2 s, or

0.3 s before or while ($t = 0$) passing through the chain array. A total of 42 features was

calculated by combining all these features.

Excel and Visual C# (Microsoft) were used for the calculations, while the Beeswarm package

for R (The R Project) was used to obtain a scatter plot of the data.

Worms

The culture and handling of *Caenorhabditis elegans* strains were performed according to

techniques described previously (Brenner, 1974). Wild-type Bristol strain RRID:WB-

STRAIN:N2_Male and mutant strains RRID:WB-STRAIN:MT1219 *egl-3(n589)*, RRID:WB-

STRAIN:VC671 *egl-3(ok979)*, RRID:WB-STRAIN:CX4544 *ocr-2(ak47),* RRID:WB-

STRAIN:JC1636 *osm-9(ky10)*, and RRID:WB-STRAIN:FK127 *tax-4(p678)*, RRID:WB-

STRAIN:MT6308 *eat-4(ky5)*, and RRID:WB-STRAIN:IK105 *pkc-1(nj1)* were obtained from

the Caenorhabditis Genetics Center at the University of Minnesota, USA. The RRID:WB-

STRAIN:KDK1 *dop-3(tm1356)* strain was originally obtained from the National BioResource

Project (Japan) and back-crossed with the wild-type N2 strain five times.

A 2-nonanone avoidance assay was performed according to the protocol described previously

(Kimura et al., 2010; Yamazoe-Umemoto et al., 2015). Briefly, 2-3 young adult hermaphrodite

worms grown synchronously were placed in the center of a 9-cm nematode growth media

(NGM) plate. Worm behavior was recorded for 12 min after 2-μL of 30% 2-nonanone (cat. no.

132-04173; Wako, Japan) diluted in 99.5% ethanol (cat. no. 0057-00456; Wako, Japan) were

placed at two spots on the surface of the NGM plate. This assay was performed under the

following three conditions: (1) Naive—the worms cultivated on 6-cm NGM plates with the

RRID:WB-STRAIN:OP-50 bacteria as food were briefly washed with NGM buffer and

subjected to the assay; (2) Pre-exposed—the worms were subjected to the assay after being pre-

exposed to 0.6 µL of 15% 2-nonanone spotted on the lid of a 6-cm NGM plate for 1 h without

food; and (3) Mock—the worms were subjected to the assay after being pre-exposed to ethanol

similarly to the pre-exposed condition. We added the mock-treated control group to ensure that

the starvation itself did not affect the odor avoidance behavior of worms and to extract

behavioral features modulated by odor pre-exposure compared to the naive and mock-treated

control groups. Images of worms on the 9-cm NGM plate during the odor avoidance assay were

acquired by a high-resolution USB camera (DMK 72AUC02; The Imaging Source, USA) with a

lens (LM16JC5MW; Kowa, Japan) at 1 Hz for 12 min. The coordinates of individual animals'

centroids were acquired from the recorded images using the Move-tr/2D software (Library Co.,

Ltd., Tokyo, Japan) and used for the STEFTR analysis.

Similar to the other sensory behaviors of the worms, trajectories in the 2-nonanone avoidance

behavior can be divided into two states: (1) "run"—a relatively long period of straight

movement, and (2) "pirouette"—a period of short movements interrupted by frequent reversals

and turns (Kimura et al., 2010; Pierce-Shimomura et al., 1999). The angular change per second

was calculated from the centroid coordinates, and movements of 1 s with angular changes larger

than 90° were classified as a turn. The histogram of turn intervals could be fitted to two

exponentials, suggesting that the turn intervals are regulated by two probabilistic mechanisms

(Pierce-Shimomura et al., 1999; Yamazoe-Umemoto et al., 2015). The time point at which the

two exponentials intersected was defined as t_{crit}, and turn intervals longer or shorter than the t_{crit}

were classified as runs or included in pirouettes, respectively. The t_{crit} was calculated for the

control (i.e., naive and mock-treated) condition for wild-type and mutant strains. Excel

(Microsoft) was used for the above calculations. The odor concentrations that the worms

experienced at specific spatiotemporal points were calculated according to the dynamic odor

gradient model based on the measured odor concentration (Tanimoto et al., 2017; Yamazoe-

Umemoto et al., 2018).

Statistical analyses were performed with Prism ver. 5.0 for Mac OSX (GraphPad Software, CA,

USA) and R (The R Project). Sample size was determined based on the previous report

(Yamazoe-Umemoto et al., 2015). A part of the original data used in this study had already been

analyzed and published previously (Yamazoe-Umemoto et al., 2015), and re-analyzed with the

STEFTR method.

Penguins

Fieldwork was performed on chick-rearing Adélie penguins at Hukuro Cove colony (69°13′ S,

39°38′ E) in Lützow-Holm Bay, East Antarctica. GPS-depth loggers (GPL380-DT or GPL400-

D3GT, weighing 55-85 g; Little Leonardo, Japan) were deployed among 11 penguins during the

period between 27 December 2016 and 10 January 2017 and recovered from all the birds after

1-2 days. While the loggers were set to record GPS positions and depth every second, they

could not record GPS positions when the penguins were diving. Therefore, we linearly

interpolated the data, when necessary, to obtain GPS positions every 1 minute before further

analysis. See Kokubun et al. for methodological details (Kokubun et al., 2015). This fieldwork

was carried out in accordance with the recommendations of the Law relating to Protection of the

Environment in Antarctica. The protocol was approved by the Ministry of the Environment,

Government of Japan. Sample size was not predetermined.

Flying seabirds

Fieldwork was performed on *Calonectris leucomelas* at Funakoshi-Ohshima Island (39°24'N,

141°59'E) between August and September in 2011, 2012, 2013, and 2015. We attached GPS

loggers (GiPSy-2, 37 × 16 × 4 mm or GiPSy-4, 37 × 19 × 6 mm; TechnoSmArt, Roma, Italy) to

the back feathers of chick-rearing Streaked Shearwaters with Tesa$^{®}$ tape (Beiersdorf AG;

GmbH, Hamburg, Germany) and cyanoacrylate glue (Loctite$^{®}$401; Henkel Ltd., Hatfield, UK).

The loggers were housed in waterproof heat-shrink tubing and set to record one fix per minute.

The total weight of the unit was 25 g, which was less than 5% of the mean mass of the birds in

accordance with the suggested load limit for flying seabirds. After approximately two weeks of

deployment, we recaptured and retrieved the loggers. See Yoda et al. for methodological details

(Yoda et al., 2014). The study was carried out in accordance with the recommendations of the

guidelines of the Animal Experimental Committee of Nagoya University. The protocol was

approved by the Animal Experimental Committee of Nagoya University. Sample size was not

predetermined. A part of the original data used in this study had already been analyzed and

published previously (Yoda et al., 2014), and re-analyzed with the STEFTR method.

Rats

Locomotion data of an adult male Long Evans rat were obtained from the Collaborative

Research in Computational Neuroscience (CRCNS; RRID:SCR_005608) data sharing website

(https://crcns.org, hc-3 dataset, ec013 rat) (Mizuseki et al., 2014). The rat foraged for randomly

dispersed water or foods on an elevated open field (180 cm × 180 cm) for 17 - 53 min. The rat's

position was tracked by monitoring two light-emitting diodes mounted above the head with an

overhead video camera at 30 Hz. The 30-Hz tracking data were resampled to 39.0625 Hz for

offline processing. This study was carried out in accordance with the recommendations of the

Regulations on Animal Experiments in Osaka City University. The protocol was approved by

the Animal Care and Ethics Committee of Osaka City University. Sample size was not

predetermined. The original data used in this study had already been analyzed and published

previously (Diba and Buzsáki, 2008; Mizuseki et al., 2009; 2014), and re-analyzed with the

STEFTR method.

Flies

Fruit flies *D. melanogaster* were raised on standard yeast-based media at 25ºC and 40 - 60%

relative humidity under a 12-hour light/dark cycle. *Canton-S* flies aged between 6-8 days after

eclosion were used as a wild-type. After eclosion, the males were housed singly, while females

were housed in groups until the experiment.

The locomotion measurement was performed as described previously with minor modifications

(Kohatsu and Yamamoto, 2015; Kohatsu et al., 2011). Briefly, a male fly was tethered with a

metal wire on its dorsal thorax and positioned over an air-supported Styrofoam ball (diameter, c.

a. 6 mm). The locomotion trajectory of the fly was recorded by monitoring the rotations of the

Styrofoam ball using an optical computer mouse sensor (BSMRU21BK; BUFFALO INC.,

Nagoya, Japan). The sensor detected the movements of the ball in the horizontal (Δx) and

vertical (Δy) directions, which correspond to lateral and forward movements of the male fly,

respectively. The Δx and Δy values, together with timestamps, were sent to a computer at 60 Hz

via an Arduino Due microcontroller (Switch Science, Japan) with a custom sketch program. The

60-Hz data were down-sampled to 4-Hz data for the information gain analysis. The

measurements were obtained at 25±1°C and 50±10% relative humidity and within 4 hours after

light onset.

Female pheromones were applied to the male fly by placing the female's abdomen in contact

with the male's foreleg at the onset of the measurement. A manipulator (M-3333, Narishige,

Tokyo, Japan) actuated a pipette with a volume of 200 µL (FUKAEKASEI Co., Ltd., China), in

which a live female with her abdomen exposed toward a male fly was captured. We manually

controlled the position of the manipulator to contact the female's abdomen to the male's foreleg.

This contact procedure was omitted in the control experiments.

Visual stimulus was applied directly after pheromone application by starting horizontal

movements of the female fly in front of the male fly as described (Kohatsu et al., 2011). The

visual stimulus consisted of ten left-right horizontal movements of the female that lasted for 40

s. Each movement started with the female fly in the front of the male fly (i.e., center) and

continued as the female fly moved left until it reached the left end of the rail (i.e., 5 mm away

from the center), then moved right until it reached the right end of the rail (i.e., 5 mm away from

the center), and ended when it came back to the center with a constant velocity of 5 mm/s. This

movement was driven by a stepper motor (42BYG Stepper Motor, Makeblock Co., Ltd.,

Shenzhen, China) controlled by a custom sketch program (processing software version 3.3.7).

We defined one round (4 s in total) as the movement related to the female starting to moving

away from the center, reaching the left end of the rail, passing the center, moving away to reach

the right end of the rail, and coming back to the center again. Each of the "moving away" and

"coming back" periods lasted for 1 s.

The Δx and Δy values with timestamps obtained in the final eight rounds were used for the

analysis. To detect the characteristic parameters for the chasing behavior, we used Δx and Δy

values during the period when the female was moving away from the male (2 s/round). We set

the angle of the chasing behavior as 0 degree when the male moved forward. Angles between 0

and 90 degrees indicate that the male fly is moving towards the female that was moving away

from the male.

As the parameters (velocity, bearing, and their time-differential values) were not normally

distributed (Shapiro-Wilk test; see Supplementary Table 13), their values were compared

between conditions (with/without pheromone) using the Mann-Whitney U test followed by

Bonferroni correction for multiple comparison. We used the Steel-Dwass test to compare values

of the parameters between rounds. Statistical analyses were conducted using R software version

3.4.4. No statistical methods were used to pre-determine sample sizes, but our sample sizes are

similar to those in previous studies (Kohatsu and Yamamoto, 2015; Kohatsu et al., 2011).

Bats

Three adult Japanese horseshoe bats (*Rhinolophus ferrumequinum nippon*, body length: 6–8 cm,

body mass: 20–30 g) were captured from natural caves in the Hyogo and Osaka prefectures in

Japan as previously described (Yamada et al., 2016). The bats were housed in a temperature-

and humidity-controlled colony room [4 m (L) × 3 m (W) × 2 m (H)] with a 12-h light/dark

cycle at Doshisha University in Kyoto, Japan, and were allowed to fly freely and given access to

mealworms and water. Captures were conducted under license and in compliance with current

Japanese law. This study was carried out in accordance with the recommendations of Principles

of Animal Care (publication no. 86-23 [revised 1985]) of the National Institutes of Health) and

all Japanese laws. The protocol was approved by the Animal Experiment Committee of

Doshisha University.

Methods for acoustic navigation measurement in bats have been described elsewhere (Yamada

et al., in revision). Briefly, the experiments were conducted in a flight chamber constructed

using steel plates [9 (length) × 4.5 (width) × 2.5 m (height)] under lighting with red filters (>650

nm) to avoid visual effects on the bats. An obstacle environment was constructed using plastic

chains (diameter: 4 cm) suspended from the ceiling of the chamber. The chains were arranged at

15-cm intervals along the *x*-axis and at 22-cm intervals along the *y*-axis so that the bat was

forced to fly in an S-shaped pattern without passing between chains. Three naive bats were

observed for 12 continuous repeated flights so that their echolocation behavior in unfamiliar and

familiar spaces could be compared. In this study, the first three flights were defined as

unfamiliar flights, while the last three flights were defined as familiar flights.

The flight behavior of the bats was recorded at 125 frames/s using two digital high-speed video

cameras (MotionPro X3; IDT Japan, Inc., Japan) placed in the left and right corners of the flight

chamber. Based on a direct linear transformation technique, the successive 3D positions of the

flying bats, as well as the locations of other objects, were reconstructed using motion analysis

software (DIPPMotionPro ver. 2.2.1.0; Ditect Corp., Japan). The statistical calculations were

performed with SPSS version 23 (IBM Corp.).

Calcium imaging of worm's neurons

Calcium imaging of the worms' ASH neurons was performed according to the previous method

with some modifications (Tanimoto et al., 2017). Briefly, transgenic strains expressing GCaMP3

(Tian et al., 2009) and mCherry (Shaner et al., 2004) in ASH sensory neurons under the *sra-6*

promoter (KDK70034 and KDK70072; 20 ng/µl of *sra-6p::GCaMP3*, 20 ng/µl of *sra-*

6p::mCherry, 10 ng/µl of *lin-44p::GFP*, 50 ng/µl of PvuII-cut N2 genomic DNA as a carrier in

N2 background) were placed on an NGM agar plate on a robotic microscope system, OSB2

(Tanimoto et al., 2017). Although these transgenic worms were immobilized with the

acetylcholine receptor agonist levamisole (Lewis et al., 1980) for high-throughput data

acquisition through simultaneous imaging of multiple worms, the previous study revealed that

the ASH activity is essentially unaffected by levamisole-treatment (Tanimoto et al., 2017). A

constant gas flow of 8 mm/min was delivered, in which the mixture rate of 2-nonanone gas with

air was changed to create a temporal gradient of odor concentration. The temporal change in

odor concentration was measured by a custom-made semiconductor sensor before and after the

series of calcium imaging experiments on each day. The fluorescence signals of GCaMP3 and

mCherry in ASH neurons were divided into two channels using W-View (Hamamatsu, Japan),

an image splitting optic, and captured by an electron multiplying charge-coupled detector (EM-

CCD) camera (ImagEM; Hamamatsu, Japan) at 1 Hz. The intensities of fluorescence signals

from cell bodies were extracted and quantified by ImageJ (NIH) after background subtraction.

The average ratio over 30 s prior to the odor increase was used as a baseline (F_0), and the

difference from F_0 (ΔF) was used to calculate the fluorescence intensities of GCaMP3 and

mCherry ($F = \Delta F/F_0$). The ratio between florescence intensities of GCaMP and mCherry

(GCaMP/mCherry) was used in the figure.

RESULTS

Estimation of behavioral states

As the first part of the analysis, we classified the trajectory into several behavioral states based

on the distribution of a basic behavioral feature. Behavior of animals consists of several states

(Jonsen et al., 2013; Patterson et al., 2008), where basic behavioral features such as speed and

direction change are likely distributed probabilistically with a center value that is optimal for

each state. Thus, behavior can be more easily characterized when the behavioral features are

analyzed for each state rather than for the entire behavior as a whole. In fact, classifying the

trajectory into several states is one of the essential preprocessing steps in trajectory mining of

people and vehicles in data science (Zheng, 2015).

For the state classification, we calculated the averages (*Ave*) and variances (*Var*) of four basic

behavioral features: velocity (*V*), temporal changes in velocity (i.e. acceleration, *dV*), bearing

(*B*), and temporal changes in bearing (*dB*). These 8 features were represented in the form of

histograms. Based on the hypothesis that basic behavioral features are likely to be distributed

probabilistically for each state, we chose one of the features whose histogram likely consists of

multiple probabilistic distributions, and classified the feature values into several clusters with

the EM algorithm (Dempster et al., 1977). The EM algorithm, which is an iterative method to

estimate model parameters that maximize the likelihood of the model for data, was used to

estimate parameters such as the average and variance of each cluster. Each cluster was then

regarded as a behavioral state. As proof-of-principle experiments, we analyzed the trajectories

of worms and penguins, whose behavioral states have been studied previously using other

methods (Pierce-Shimomura et al., 1999; Yoda et al., 2001).

The roundworm *Caenorhabditis elegans* has been used as a model animal for quantitative

behavioral analysis owing to the ease of tracking behavior (movement for a few cm on agar

surface can be easily recorded with an inexpensive high-resolution camera), optical monitoring

neural activities, and genetic analyses and manipulations (De Bono and Maricq, 2005). Further,

the neuronal wiring in *C. elegans* has been described in complete detail (White et al., 1986). In

this study, we focused on the avoidance behavior to the repulsive odor of 2-nonanone (Fig. 2A,

left) (Bargmann et al., 1993; Kimura et al., 2010). We chose this behavioral paradigm for the

proof-of-principle experiment because the odor avoidance behavior has been quantitatively,

although not fully, analyzed previously (Kimura et al., 2010; Yamazoe-Umemoto et al., 2015).

The behavior of the worms was recorded with an USB camera for 12 minutes, and the position

of each worm's centroid was extracted every second.

We observed multiple peaks in the histogram corresponding to the variance values for bearing

change (*dB_Var*, Fig. 2B). These values were automatically classified into 5 clusters by the EM

algorithm (Fig. 2C). Upon mapping the clusters on to the trajectories, we found that cluster 0

corresponds to relatively straight part, while the other clusters correspond to more complex

parts (Fig. 2D, upper panel).

Cluster 0 and clusters 1-4 mainly corresponded to "run" and "pirouette," respectively, which are

the two classical behavioral states of worms in various types of sensory behavior (Fig. 2D,

lower panel) (Lockery, 2011; Pierce-Shimomura et al., 1999). "Run" constitutes relatively

straight movement, while "pirouette" is characterized by short, straight movement divided by

frequent large changes in angle (turns and reversals). Runs and pirouettes are usually classified

based on a threshold value for the duration between consecutive large angle changes (Pierce-

Shimomura et al., 1999), unlike in this method. We collectively regarded clusters 1-4 as "cluster

1" because the positional information of worms does not appropriately reflect their actual

locations during pirouettes due to insufficient spatiotemporal resolution of the recording system

for relatively long-distance navigation, such as odor avoidance behavior (Yamazoe-Umemoto et

al., 2015). We found that more than 90% of the cluster 0 and 1 corresponded with the run and

pirouette, respectively (Fig. 2E and F). Therefore, we concluded that the STEFTR method

properly classified the odor avoidance behavior into distinct behavioral states.

Next, we applied the same process to the trajectories of penguins obtained using GPS devices.

Penguins are good model wild animals for studying long-distance navigation given their

relatively large body size for attaching GPS devices and their habit of returning to a colony,

which makes the attachment and the recovery of GPS data easy (Yoda, 2018; Yoda et al., 2001).

In this study, GPS and depth sensors were attached to 11 penguins from a colony in the

Antarctic Continent; the depth sensors were used to evaluate the accuracy of state estimation

(see below). The penguins moved by walking and swimming about 10 km for feeding, and each

dataset contained up to 2 days of data recordings (Table 1 and Fig. 3A). Like in the case of

worms, 8 basic behavioral features were extracted from the penguin trajectory data and

represented as histograms. We chose the average velocity (V_Ave) as it showed multiple peaks

in the histogram. The EM algorithm classified it into 5 clusters (Fig. 3B lower panel, C and D

upper panel).

Interestingly, the clusters exhibited significantly different distributions in multiple behavioral

features. For example, the values for the duration of each bout were much longer in cluster 0

than in clusters 1, 2, and 3 (Fig. 3D, middle panel). In addition, although the clusters were

classified only based on the horizontal velocities, the depths for cluster 0 and 1 were

significantly closer to zero than those for clusters 2, 3, and 4 (Fig. 3D, lower panel). These

results are consistent with the idea that each cluster reflects a behavioral state that is a complex

function of multiple behavioral features.

To evaluate whether the clusters actually reflect different behavioral states, we manually

classified the penguin behavior into four states (resting, transit by walking, transit by

swimming, and diving) based on diving depth (from depth sensor), movement velocity and

distance from the colony (both calculated from the GPS positional information) (Yoda et al.,

2001). Penguins stayed and rested at the colony (location does not change much; depth is zero),

moved on land and ice mainly by walking (location changes relatively slowly; depth is near-

zero), swim in the sea to go to the foraging area (location changes quickly; depth is relatively

shallow in general and sometimes increases when they are moving towards the feeding area),

dive deeply at the foraging area (location does not change much; dives occur continuously in

bouts), and then come back to the colony by swimming and walking. The resting at the colony

and swimming correlated with clusters 0 and 4, respectively (Fig. 3E and F). In addition, most

of clusters 1 and 2 correlated with walking, while about 50% of cluster 3 corresponded to diving

(Fig. 3F). Thus, when a behavioral state is classified to a cluster other than cluster 3, the

penguin is likely to be resting, walking, or swimming. If a behavioral state is classified to

cluster 3, which is ~10% of all the behavior recorded, the penguin is either diving or swimming.

Remarkably, although the clustering is only based on the trajectories of 11 penguins for a few

days, the false positive rates were less than 10% and the sensitivity of the analysis was greater

than 90% in all the cases (Fig. 3G). Thus, we concluded that the STEFTR method can

reasonably estimate different behavioral states only based on trajectory data.

The STEFTR method was also applied to the trajectories of flying seabirds in the wild and rats

in the laboratory. The seabirds, *Calonectris leucomelas,* traveled ~100 times longer distances

(up to 1,000 km) with ~10 times the speeds compared to penguins (Matsumoto et al., 2017;

Yoda et al., 2014). Like in the case of penguins, the average velocity (V_Ave) showed multiple

peaks and was classified into 5 clusters (Fig. 4A and B) with significantly different profiles for

duration and directedness (Fig. 4C). In the case of rats, the variance of velocity change

(dV_Var) was chosen, and the EM algorithm classified the behavior into 4 clusters (Fig. 4D and

E). Interestingly, significant differences were observed in several features such as duration and

speed (Fig. 4F). Such information can help ecologists estimate the candidates for feeding areas

where fishes may be more densely distributed and discover biologically important marine areas.

It can also help neuroscientists in estimating candidate conditions to further explore specific

neural activities.

Comprehensive extraction of behavioral features modulated by learning

As a second part of the STEFTR method, comprehensive feature extraction was performed by

comparing a specific behavioral state in two different conditions, such as cluster 0 of worms

before and after learning. Comprehensive semi-automated analysis can be very helpful to

compare behavioral features in two conditions. This is because even when the overall result of

behavioral responses are different in two conditions, it is still difficult to quantitatively

determine which part of the trajectories are different (Fig. 2A left and right, for example).

Furthermore, even if several behavioral features are found to be different, it is possible that

other more prominent feature differences may exist. We considered that learning-dependent

changes in behavior should be one of the best models for comprehensive feature extraction

because the differences in behavioral features should reflect learning-dependent changes in

neural/brain activities.

As an useful index for feature extraction, we chose information gain, the index for decision tree

analysis (Quinlan, 1986). Binary decision tree analysis is for splitting a dataset into two sub-

groups by automatically selecting the best feature and its parameter showing the largest

information gain (i.e., difference of uncertainty, or "information entropy", between before and

after division). Each data point is then classified into one of the sub-classes based on whether it

has a larger or smaller value than an automatically determined threshold. When applied for

binary classification, decision tree analysis automatically evaluates the classification

performance of a large number of features as designed by the researchers. This analysis results

in the extraction of certain features, allowing us to easily understand the utility of particular

features in the classification. This approach is substantially different from those that employ

support vector machines and/or deep neural networks, wherein the relationships between the

classification and the features of the data cannot be easily discerned.

We first analyzed learning-dependent changes in worm odor avoidance behavior. Worm odor

avoidance behavior is enhanced by pre-exposure to the odor as a type of non-associative

learning, and pre-exposed worms migrate significantly longer distances from the odor source

than control worms do during the same period (Fig. 2A) (Kimura et al., 2010). This

phenomenon is interesting because prior exposure to a stimulus generally causes reduction,

instead of enhancement, of the response to the stimulus through adaptation or habituation.

Although this is a simple form of learning, this odor learning is modulated by multiple

neuromodulators, including dopamine, octopamine (the worm counterpart of mammalian

noradrenaline), and neuropeptides (Kimura et al., 2010; Yamazoe-Umemoto et al., 2015).

Previous quantitative analyses have shown that the enhanced odor avoidance behavior is not

caused by changes in speed, but mostly by increases in run duration (Yamazoe-Umemoto et al.,

2015). However, this did not rule out the possibility that other behavioral features play more

profound effects.

As an example of comprehensive feature extraction from a behavioral state, we focused on

learning-dependent changes in cluster 0 (run) because the values of their centroid migration are

quantitatively more reliable than cluster 1 (pirouette) as mentioned above. In addition to the

basic behavioral features used for the estimation of behavioral states (V, dV, B, and dB), we

also calculated directedness (Dir) (Gorelik and Gautreau, 2014), and the odor concentration (C)

and temporal change in odor concentration (dC) that each worm experienced during the odor

avoidance behavior; C and dC were calculated based on actual measurements of the dynamic

odor gradient (Tanimoto et al., 2017; Yamazoe-Umemoto et al., 2018). For these, we calculated

the initiation (Ini), middle (Mid), termination (Ter), and all (All) values of a cluster 0 segment

(Fig. 5A). In addition, different time windows (1-6 s in this case) were used to calculate these

values because a behavioral feature could be apparent only within a specific temporal window

(for example, velocity of run (i.e. cluster 0) starts decreasing 2 s before the end of a run (Pierce-

Shimomura et al., 1999)). We also calculated durations (Dur) of cluster 0 and 1, and the

weathervane index (WV) (Iino and Yoshida, 2009). Information gain for each of these features

were compared between naive/mock and pre-exposed conditions (Fig. 5B, for example). The

information gain values for each of the features have been summarized in Table 2, and the

details are described in Supplementary Tables 2-11.

Through this analysis, we were able to find new as well as previously known behavioral

features that exhibited learning-dependent changes. First, we found that the duration of each

cluster 0 (*Clst0Dur*) exhibited higher information gain (Table 2), which corresponded to

significantly increased cluster 0 duration (Fig. 6B). This result is consistent with the findings

from previous reports (Kimura et al., 2010; Yamazoe-Umemoto et al., 2015), highlighting the

reliability of this method. We also found that the velocity at the beginning of each cluster 0

(*V_Ini*) consistently exhibited higher information gain in the average and median values in

multiple time windows (Table 2); these values were also significantly different in the pre-

exposed worms as compared to the control worms (Fig. 6C). The previous study has not

identified this difference as only average values per run (*i.e.* cluster 0) have been calculated in

the study (Yamazoe-Umemoto et al., 2015). Although the contribution of this behavioral feature

to enhanced odor avoidance is unclear at present, our results indicate that the STEFTR method

can reveal characteristic feature(s) under specific conditions, which is difficult for human

analyses to accomplish.

Odor stimuli during runs, which likely drive the worms' odor avoidance behavior, were also

found to be consistently modulated (Table 2). In fact, odor concentration (C) was significantly

lower, and the temporal change in odor concentration (dC) was significantly closer to zero (i.e.

shallower) in a learning-dependent manner (Fig. 6D and E). Because the previous study

demonstrated that worm odor avoidance behavior depends on dC rather than C at least in the

naive condition (Tanimoto et al., 2017), one possibility is that the changes in the responsiveness

of worms to dC is the underlying reason for the enhanced odor avoidance. However, it is also

possible that the odor-experienced worms were somehow located farther away from the odor

source than the unexperienced worms, and hence, sensed lower odor concentrations and

shallower odor concentration change than the latter.

Responsiveness of sensory neurons to odor increase was modulated by the odor learning

If the change in sensitivity to dC/dt is the reason underlying enhanced odor avoidance behavior,

it should be associated with changes in neural activity. Thus, we analyzed the responsiveness of

a likely candidate, ASH nociceptive neurons (Bargmann, 2006; Kaplan, 1996). Previously, we

have established the OSB2 microscope system that allows for *in vivo* calcium imaging of *C.*

elegans neurons in the presence of odor stimuli resembling those that the worms experience

during the odor avoidance behavior in the plates (Fig. 7A) (Tanimoto et al., 2017). Using the

OSB2 system, we found that ASH neurons are the major sensory neurons to cause pirouettes

upon increases in 2-nonanone concentration (Tanimoto et al., 2017). However, whether the

ASH response is modulated by 2-nonanone experience has not yet been studied.

We found that ASH responses were indeed modulated by prior odor experience. When the

worms were stimulated with a 5 nM/s odor increase rate, which is the lowest rate of change to

cause the threshold-level behavioral response in the previous study (Tanimoto et al., 2017),

ASH neurons in naive as well as mock-treated worms exhibited robust responses (Fig. 7B).

However, the ASH responses were significantly reduced in the pre-exposed worms (Fig. 7B and

C). This suggests that prior odor experience causes a reduction in the neuronal response to a

slight increase in odor concentration, subsequently causing longer run durations and enhanced

odor avoidance behavior (Fig. 7D).

Extracted behavioral features of mutant strains correspond to gene function

Next, we comprehensively analyzed learning-dependent behavioral changes in the mutant *C.*

elegans strains. Many mutant strains of *C. elegans* showing impaired learning have already

been isolated and characterized (Bargmann, 2006; Sasakura and Mori, 2013), and the behavioral

abnormalities observed in these mutants should reflect the role of the causal genes in neural

function. In fact, we have previously shown that two different genes involved in the enhanced

odor avoidance behavior cause different abnormalities in behavioral features when mutated

(Yamazoe-Umemoto et al., 2015). However, as the behavioral features exhibited by a mutant

strain could be different from one another, identification of abnormal behavioral features is

often laborious and time-consuming.

In addition to studying the previously described mutants with defective enhanced odor

avoidance behavior (*egl-3* and *egl-21* for neuropeptide biosynthesis, and *dop-3* for dopamine

receptor) (Kass et al., 2001; Suo et al., 2004; Yamazoe-Umemoto et al., 2015), we also analyzed

mutant strains found to be involved in the phenomenon in this study (*ocr-2* and *osm-9* for TRP

channels; *tax-4* for CNG channel; *eat-4* for vesicular glutamate transporter; *pkc-1* for protein

kinase) (Colbert et al., 1997; Komatsu et al., 1996; Land et al., 1994; Lee et al., 1999; Tobin et

al., 2002) (Table 3).

Neuropeptide mutant strains did not exhibit learning-dependent changes in behavioral features,

except for the velocity of *egl-3(ok979)*. This result is consistent with the previous finding that

neuropeptide signaling is required for the acquisition of odor memory (Yamazoe-Umemoto et

al., 2015). *egl-3(ok979)* may have exhibited stronger phenotypes than *egl-3(n589)* because they

are nonsense and missense mutants, respectively. Also consistent with the previous report

(Yamazoe-Umemoto et al., 2015), the *dop-3* mutants exhibited abnormalities in direction-

related behavioral features (*B* and *Dir*) while the changes in cluster 0 durations and velocities

are similar to those of wild-type worms (Table 3). Furthermore, with respect to the newly added

mutant strain, similar patterns are observed in *ocr-2* and *osm-9* mutants of the TRP channel

involved in sensory perception. On the other hand, *tax-4*, which is also involved in sensory

perception but expressed in a different set of sensory neurons (Komatsu et al., 1996; Tobin et

al., 2002), and *eat-4* and *pkc-1* mutants showed different patterns of abnormalities. Taken

together, our results suggest that the patterns of features extracted from mutant strains may

reflect functional groupings of the mutated genes. Thus, profiling and classification of extracted

mutant features of unknown genes may be useful in the estimation of their physiological

functions.

Feature extraction of fly sexual behavior

Next, we applied the technique to comprehensive feature extraction of animal behavior under

specific situations in two different conditions—heterosexual chasing behavior of *Drosophila*

melanogaster with or without pheromone sensation. On an experimental tracking system (Fig.

8A), male flies chased the target female flies' abdomens after tapping them with their forelegs

to sense the cuticular pheromone, although males do not show such chasing behavior before

tapping (Kohatsu et al., 2011). While this pheromone-driven behavior has been generally used

for the observation of neural activity during courtship behavior in fruit flies, the behavioral

features have not yet been elucidated in great detail.

In this study, we used a tracking system as described in previous studies, where a male was

exhibited the moving female abdomen with 8 time left-right round trip after a pheromone

sensation (Fig. 8A) (Kohatsu et al., 2011). We detected positive information gains in the

velocity, changes in velocity, bearing, and changes in bearing in a pheromone sensation-

dependent manner (Fig. 8B). Unexpectedly, the information gains of velocity were higher in the

earlier round trips and decreased over the trips (Fig. 8B). This suggests that the pheromonal

effect promoting chasing behavior decreases over time. To confirm the result of the STEFTR

method, we re-analyzed the speed of the male locomotion along with time-series. Consistent

with the STEFTR result, the velocity of males that had tapped the female significantly

decreased over the trips (orange group in Fig. 8C; significant differences between rounds 1, 4,

and 8), whereas that of control flies (gray group in Fig. 8C) remained mostly unchanged

(significant difference only between rounds 1 and 5). Thus, the STEFTR method can even

uncover behavioral features that fluctuate over time. The decreased tracking velocity may

reflect a decrease in motivation in the fly brain, which can be assessed directly by observing the

temporal changes in neuronal activity related to the courtship-motivation circuit in the fly brain

(Yamamoto and Koganezawa, 2013; Zhang et al., 2018).

Feature extraction of learning-dependent modulation of acoustic navigation in bats

To further demonstrate the general applicability of the method, we examined features of

acoustic navigation in bats. We have previously reported that bats improve their flight trajectory

in an indoor space with obstacles in a learning-dependent manner (Yamada et al., in revision).

Here, we optimized features such as velocity (V), distance to the obstacle chain array (R_obs

and R_x), and horizontal bearing of the flight (B_hori) for the experimental paradigm (Fig. 9A

and B). Interestingly, although the velocity (V) itself was modulated by flight experience, the

change in velocity (dV) was not (Fig. 9C and D), suggesting that bats determine flight speed

before initiating navigation, but not during navigation, at least in this experimental conditions.

As the vocalizations of bats reflect their attention or decisions (Moss and Surlykke, 2010), our

results suggest that the STEFTR method can be used to elucidate such higher brain functions

during navigation.

DISCUSSION

Measuring and analyzing behavior is one of the most prominent steps in understanding brain

function. In order to utilize "behavioral big data", we developed a hybrid

supervised/unsupervised technique, the STEFTR method, to estimate behavioral states and to

efficiently extract behavioral features solely from the trajectories of animal movement. The

behavioral states of worms and penguins estimated with the STEFTR method were in

reasonable agreement with the ones based on previous knowledge, highlighting the validity of

our method. In addition, one of the learning-dependent behavioral features extracted from

worms corresponded to a change in neural activity. Furthermore, we were able identify

temporally dynamic changes through feature extraction from fly courtship behavioral data.

One of the advantages of the STEFTR method is its versatility. Multiple methods have been

reported for the behavioral analysis of specific animals under specific conditions (Baek et al.,

2002; Branson et al., 2009; Brown et al., 2013; Dankert et al., 2009; Kabra et al., 2013; Mathis

et al., 2018; Robie et al., 2017; Stephens et al., 2008; Vogelstein et al., 2014; Wiltschko et al.,

2015). However, the animals and experimental conditions for which each of these methods can

be applied are rather limited. For example, it is still not easy to effectively and robustly extract

an animal's posture from a video image, especially in the wild. Even in laboratories, the

parameters generally need to be adjusted again when imaging conditions changed (Dell et al.,

2014; Egnor and Branson, 2016). In contrast to these methods, our method allows for

behavioral analysis based on positional information that can be extracted from animal video

images as well as from different methods such as GPS device.

As a first step of the STEFTR method, we estimated behavioral states from animal trajectories.

State estimation is one of the critical processes of movement analysis of animals in the wild as

well as of cars and people in ecology and data science, respectively (Egnor and Branson, 2016;

Jonsen et al., 2013; Patterson et al., 2008; Zheng, 2015). However, the analytical methods that

can be applied to the analysis of various types of animals (and cars and people) are still in

debate (Gurarie et al., 2016; Zheng, 2015). In the STEFTR method, we aimed to analyze

behavior without previous knowledge of the animal and/or experimental condition and

independent of the spatiotemporal scale of the behavior. We analyzed migratory velocity and

direction, the most fundamental elements of moving objects, with appropriate moving-averaged

data. Interestingly, in case of both worms and penguins, the moving average with a temporal

window that is about 1% of median length of the behavioral record provided results that were

reasonably consistent with behavioral states categorized by manual labeling (Fig. 2D-F and 3E-

G). This result suggests that the ~1% temporal window works as a low-pass filter for properly

estimating behavioral states in a scale-free manner. We estimated the number of behavioral

states to be equal to or less than 5 (Jonsen et al., 2013), which worked well in the above-

mentioned cases.

State estimation based only on trajectory analysis using the STEFTR method is not perfect, as

shown in the case of penguins (Fig. 3E). However, the estimated behavioral states likely

provide us with important information for further experiments, such as when and where in the

spatiotemporal behavioral profile of the animal behavior should be analyzed in detail, especially

in the case where the behavior has not been studied intensively in quantitative manner. For

example, relatively small movements at places distant from their nest in the wild may

correspond to the feeding area. For neurobiological/physiological analysis, the transition from

one state to the other could be triggered by a specific change in the sensory stimulus and

associated with specific neural activities. It should also be noted that the STEFTR method

allows semi-automatic state estimation and feature extraction, which is suitable for large-scale

behavioral analysis of mutant strains of laboratory model animals.

The Hidden Markov Model (HMM) has been used for the estimation of behavioral states

(Gurarie et al., 2016; Jonsen et al., 2013). However, the HMM depends on the idea of the

Markov process, wherein the state transition is probabilistic. In general, state transition in

animal behavior is triggered by external as well as internal stimuli, which are not always

probabilistic. While the STEFTR method assumes a probabilistic normal distribution of a

behavioral feature in a state, it does not assume probabilistic transition between states.

For comprehensive feature extraction, we used information gain, an index used in decision tree

analysis. Decision tree analysis is one of the machine learning techniques used for classification.

Classification analysis involves classifying new, unlabeled data into appropriate classes using

characteristic features and the parameters that have been extracted from the known class-labeled

data. In the present study, however, the classification itself was not meaningful because the data

were already classified (with/without learning or with/without sex pheromone). Instead, we

focused on the procedure in the classification that identifies features useful for distinguishing

between the two classes. In other words, behavioral features that are different between two

classes (*i.e.,* conditions) should be able to effectively classify the behavioral data of animals in

two conditions.

Although the STEFTR method does not directly provide information about brain/neural activity

underlying animal behavior, it provides us with clues required to formulate hypotheses related

to the experimental investigation of the neural activity, as shown in the case of learning-

dependent changes in behavioral features and neural activity (Fig. 7). For example, animals in

the wild experience continuously changing visual, auditory, and olfactory stimuli, each of which

contain multi-dimensional information (color, shape, tone, different chemical compounds, etc.).

Therefore, it is difficult to identify which aspect(s) of the particular stimulus actually triggers a

change in animal behavior. Estimation of behavioral states using the STEFTR method will

allow us to identify the behavior-triggering stimulus by focusing on the timing and/or place of

the behavioral transition. Similarly, large-scale recording of neural activities from moving

animals in the laboratory itself is difficult to interpret. However, state estimation and feature

extraction of the behavior will greatly facilitate the identification of neural activities that are

associated with behavioral transitions and/or specific behavioral features.

FIGURE LEGENDS

Figure 1. A workflow of the STEFTR method. Trajectory data of animals are used to calculate 8

basic behavioral features, and one of them are analyzed by the EM algorithm to estimate

behavioral states (upper panels). From a behavioral state, behavioral features are

comprehensively evaluated by using information gain (lower panels).

Figure 2. State estimation of worms. (A) Examples of the trajectories of 3 worms before (left) or

after odor learning (right) in 12 min of 2-nonanone avoidance assay, overlaid on a schematic

drawing of a 9 cm plate. One of the trajectories is magnified below. (B) The histograms of 8

basic behavioral features. Horizontal and vertical axes indicate the values and the density of

each feature. *dB_Var* was chosen (red square) for the EM analysis. (C) Classification of *dB_Var*

by the EM algorithm. Each cluster distribution (cluster 0, 1, 2, 3, and 4 are indicated by purple,

blue, green, orange, and red dashed lines, respectively, similarly to the following figures) and

the sum of clusters (red solid line) are shown. Inset is a magnified view. (D) Comparison of the

cluster 0 and 1 (upper panel; purple and green, respectively) with run and pirouette (lower

panel; blue and red, respectively) on a trajectory. The initial 2 minutes (gray in both panels)

were excluded from the analyses because worms do not avoid the odor during the period

(Kimura et al., 2010). (E) Event numbers of cluster 0 (left) and 1 (right) in run and pirouette. (F)

Matching matrix of the state estimation.

- 33 -

Figure 3. State estimation of penguins. (A) The trajectories of 11 penguins (black lines) on the

Antarctic Continent (gray area; white area is the sea). Horizontal and vertical axes indicate

longitude and latitude, respectively. (B) The histograms of 8 basic behavioral features (upper

panels) and the classification by the EM algorithm. *V_Ave* was chosen (red square) and

classified into 5 clusters (lower panels). (C) Mapping of the five clusters on the trajectory. (D)

Differences in velocity, duration, and depth among the clusters. Each dot represents a cluster

bout, and the bars represent the median and the first and third quartiles. Significant differences

among clusters suggest that the clusters correspond to different behavioral states. Statistical

values were calculated using Kruskal-Wallis test with *post hoc* Dunn's test. **$p < 0.01$, ***$p <$

0.001. (E) An example of comparison of the clusters from the STEFTR analysis with the

behavioral states by manually classified labels, which is based on diving depth, movement

speed recorded from GPS data, and distance from the colony. (F) Event numbers of each

clusters. (G) Matching matrix of the state estimation. The statistical details are described in

Supplementary Table 1.

Figure 4. State estimation of flying seabirds in the Pacific Ocean (A-C) and rats in the open

maze (D-F). (A) From the histograms of 8 basic features of flying seabirds (left panels), the

average of velocity (*V_Ave*) was chosen and classified into 5 clusters (right panel). (B) Mapping

of the clusters on the trajectory. The gray region is the northern part of Japan (Tohoku and

Hokkaido area), while the white region is the sea. (C) Significant differences were observed in

duration (left) and directedness (right). (D) From the histograms of 8 basic features of rats (left

panels), the variance of velocity change (d*V_Var*) was chosen and classified into 4 clusters

(right panel). (E) An example of trajectories of one rat. (F) Significant differences were

observed in duration (left) and velocity (right). Each dot represents a cluster bout, and the bars

represent the median and the first and third quartiles. Statistical values were calculated using

Kruskal-Wallis test with *post hoc* Dunn's test. ***p < 0.001. The statistical details are described

in Supplementary Table 1.

Figure 5. Feature extraction of worm behavior. (A) Schematic drawing of the behavioral

features. (B) One example (*V_Ini*; average of time window 2) of calculation of information

gain.

Figure 6. Extracted features modulated by the odor learning. (A) Enhanced odor avoidance

behavior in worms caused by odor pre-exposure. Left: End-points of 25 worms in each

condition plotted on a schematic representation of the assay plate. Right: Avoidance distance

(distance between the center line of the plate and the end-point of the behavior) of each worm.

Each dot represents a worm. Significant differences were observed between the pre-exposed

worms and the naive and mock-treated worms (***p < 0.001, Kruskal-Wallis test with *post-hoc*

Dunn's test). (B, C, D and E) Distributions of extracted features. Duration (B), the initial value

of velocity (C; average of time window 2), the average odor concentration (D), and the average

odor concentration change (E) of each run (***p < 0.001, Kruskal-Wallis test with *post-hoc*

Dunn's test). Each dot represents a cluster bout, and the bars represent the median and the first

and third quartiles. The statistical details are described in Supplementary Table 1.

Figure 7. Sensory responses to slight increases in odor concentration were reduced by pre-

exposure to the odor. (A) A schematic drawing of calcium imaging of neural activity of worms

under odor stimuli. Several immobilized worms were simultaneously exposed to an odor flow

whose concentration was changed by controlling syringe pumps. (B) Responses

(GCaMP/mCherry) of ASH neurons in naive (n = 25), mock-treated (n = 29), and pre-exposed

(n = 26) worms. Thick lines with gray shadows indicate mean ± standard error of mean, while

thin lines indicate individual responses. (C) Distributions of peak values during the odor-

increasing phase (t = 40-80 s) shown in panel B. The bars represent the median. (***$p < 0.001$,

Kruskal-Wallis test with *post-hoc* Dunn's test). (D) A model relationship between odor

concentration change and behavioral response during navigation along the odor gradient. When

naive and mock-treated worms sensed a slight increase in the odor concentration, which is a

sign of migrating in the wrong direction, they stopped a run and started a pirouette to search for

a new direction. In contrast, the pre-exposed worms did not respond to a slight increase in odor

concentration, leading to longer run durations (and shorter pirouette durations in total as a

consequence), which likely contribute to the enhanced avoidance distance. The statistical details

are described in Supplementary Table 1.

Figure 8. Pheromone-driven responses of male fruit flies decreased over time. (A) A schematic

drawing of the experimental setup. A female fly was actuated leftward and rightward in front of

the male fly. The locomotion of the male fly was monitored by an optical sensor, which

recorded lateral (Δx) and forward (Δy) movements at 4 Hz. (B) Information gain. Darker and

lighter colors mean larger and smaller values, respectively. (C) Distribution of velocity in the

chasing behavior of male flies. Control (without female tapping, gray dots) and experimental

(with female tapping, orange dots) groups are shown. Solid and dotted lines represent the

median and the first and third quartiles, respectively. Asterisks indicate the statistical

significance between the control and test groups (Mann-Whitney U test followed by Bonferroni

test for multiple comparison correction, p < 0.05). Different characters in each group indicate

statistical significance among rounds (Steel-Dwass test, $p < 0.05$). The statistical details are

described in Supplementary Tables 12 and 13.

Figure 9. Learning-dependent changes in bat acoustic navigation. (A) The experimental setup

for monitoring the 3D flight trajectory of a bat during obstacle avoidance flight in a chamber.

(B) Representative flight trajectories of a bat in the horizontal plane during repeated flights in

the obstacle course. The figure on the top combines the first three (red) and last three (blue)

flight trajectories. Each behavioral feature was collected in three segments: earlier, middle, and

later terms. The figure on the bottom shows an expanded view of the earlier term in the first

flight. Definition of the horizontal bearing of the flight (*B_hori*), distance (*R_obs*), and bearing

(*B_obs*) of the bat to the nearest edge point of the obstacle chain array, longitudinal directional

distance to the frontal chain array (*R_x*), and lateral directional distance to the inside pitch of the

chains array (*R_y*) are indicated. Time windows for the analysis of each behavioral feature were

0.1, 0.2, or 0.3 s before or while ($t = 0$) passing through the chain array. (C) A list of extracted

features of bat acoustic navigation modulated by flight experience. (D) Distributions of *V(-0.3)*

and *dV(-0.3)* are plotted. The bars represent the median and the first and third quartiles. (*$p <$

0.05, Kruskal-Wallis test with *post-hoc* Dunn's test). The statistical details are described in

Supplementary Table 1.

ACKNOWLEDGEMENTS

We thank Drs. André Brown, Katsuyoshi Matsushita, Ken-ichi Hironaka, Takuma Degawa,

Daisuke Yamamoto, Soh Kohatsu and the Kimura laboratory members for suggestions and

comments on this work. Nematode strains were provided by the Caenorhabditis Genetics Center

(funded by the NIH Office of Research Infrastructure Programs P40 OD010440) and by

National Bioresource Project funded by the Ministry of Education, Culture, Sports, Science and

Technology (MEXT), Japan. This work was supported by Interdisciplinary graduate school

program for systematic understanding of health and disease (for S.J.Y.), by the Tohoku

Ecosystem-Associated Marine Science (for K. S.), by KAKENHI 25249020 and JP16H06536

(for K.H.), 24681006, JP16H01769, JP16H06541 (for K.Y.), 16H04655, 18H05069,

and17K19450 (for A.K.), JP16H06542 (for S.H.), JP16H06539 (for T.M.), JP16H06545 (for

K.D.K.) from the MEXT, by PRESTO JPMJPR14D8 (for S.H.) by JST, and by Osaka

University Co-Creation Program (for T.M. and K.D.K.).

AUTHOR CONTRIBUTIONS

S.J.Y, T.M. and K.D.K. designed the experiments, S.J.Y, K.O., K.I., N.K., T.K., D. T., Y.Y.,

Y.I., F.H., K.F., Y.T., A.Y.-U., K.H., K.S., K.Y., A.T., Y.I., A.K., S.H. and K.D.K. performed

the experiments, S.J.Y. and T.M. analyzed the data, and S.J.Y. and K.D.K. wrote the

manuscript. All authors reviewed the manuscript.

REFERENCES

Alivisatos, A. P., Chun, M., Church, G. M., Greenspan, R. J., Roukes, M. L., and Yuste, R.

(2012). The Brain Activity Map Project and the Challenge of Functional Connectomics.

Neuron 74, 970–974. doi:10.1016/j.neuron.2012.06.006.

Anderson, D. J., and Perona, P. (2014). Toward a Science of Computational Ethology. *Neuron*

84, 18–31. doi:10.1016/j.neuron.2014.09.005.

Baek, J.-H., Cosman, P., Feng, Z., Silver, J., and Schafer, W. R. (2002). Using machine vision

to analyze and classify Caenorhabditis elegans behavioral phenotypes quantitatively. *J*

Neurosci Methods 118, 9–21.

Bargmann, C. I. (2006). Chemosensation in C. elegans. *WormBook*, 1–29.

doi:10.1895/wormbook.1.123.1.

Bargmann, C. I., Hartwieg, E., and Horvitz, H. R. (1993). Odorant-selective genes and neurons

mediate olfaction in C. elegans. *Cell* 74, 515–527.

Berman, G. J. (2018). Measuring behavior across scales. *BMC Biol* 16, 23. doi:10.1186/s12915-

018-0494-7.

Bishop, C. M. (2006). *Pattern Recognition and Machine Learning*. Springer Verlag.

Branson, K., Robie, A. A., Bender, J., Perona, P., and Dickinson, M. H. (2009). High-

throughput ethomics in large groups of Drosophila. *Nat Meth* 6, 451–457.

doi:10.1038/nmeth.1328.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. *Genetics* 77, 71–94.

Brown, A. E. X., and de Bivort, B. (2018). Ethology as a physical science. *Nat Phys* 14, 653–

657. doi:10.1038/s41567-018-0093-0.

Brown, A. E. X., Yemini, E. I., Grundy, L. J., Jucikas, T., and Schafer, W. R. (2013). A

dictionary of behavioral motifs reveals clusters of genes affecting Caenorhabditis elegans

locomotion. *Proc Natl Acad Sci USA* 110, 791–796. doi:10.1073/pnas.1211447110.

Colbert, H. A., Smith, T. L., and Bargmann, C. I. (1997). OSM-9, a novel protein with

structural similarity to channels, is required for olfaction, mechanosensation, and olfactory

adaptation in Caenorhabditis elegans. *J Neurosci* 17, 8259–8269.

Dankert, H., Wang, L., Hoopfer, E. D., Anderson, D. J., and Perona, P. (2009). Automated

monitoring and analysis of social behavior in Drosophila. *Nat Meth* 6, 297–303.

doi:10.1038/nmeth.1310.

De Bono, M., and Maricq, A. V. (2005). Neuronal substrates of complex behaviors in C.

elegans. *Annu Rev Neurosci* 28, 451–501. doi:10.1146/annurev.neuro.27.070203.144259.

Dell, A. I., Bender, J. A., Branson, K., Couzin, I. D., de Polavieja, G. G., Noldus, L. P. J. J., et
al. (2014). Automated image-based tracking and its application in ecology. *Trends in*
Ecology & Evolution 29, 417–428. doi:10.1016/j.tree.2014.05.004.

Dempster, A. P., Laird, N. M., Rubin, D. B., 1977 (1977). Maximum likelihood from
incomplete data via the EM algorithm. *J Royal Stat Soc B* 39, 1–38. doi:10.2307/3315718.

Diba, K., and Buzsáki, G. (2008). Hippocampal network dynamics constrain the time lag
between pyramidal cells across modified environments. *J Neurosci* 28, 13448–13456.
doi:10.1523/JNEUROSCI.3824-08.2008.

Egnor, S. E. R., and Branson, K. (2016). Computational Analysis of Behavior. *Annu Rev*
Neurosci 39, 217–236. doi:10.1146/annurev-neuro-070815-013845.

Frank, E., Hall, M. A., and Witten, I. H. (2016). "The WEKA Workbench (Online Appendix),"
in *Data Mining* (Morgan Kaufmann).

Gomez-Marin, A., Paton, J. J., Kampff, A. R., Costa, R. M., and Mainen, Z. F. (2014). Big
behavioral data: psychology, ethology and the foundations of neuroscience. *Nat Neurosci*
17, 1455–1462. doi:10.1038/nn.3812.

Gorelik, R., and Gautreau, A. (2014). Quantitative and unbiased analysis of directional
persistence in cell migration. *Nat Protoc* 9, 1931–1943. doi:10.1038/nprot.2014.131.

Gurarie, E., Bracis, C., Delgado, M., Meckley, T. D., Kojola, I., and Wagner, C. M. (2016).
What is the animal doing? Tools for exploring behavioural structure in animal movements.
J Anim Ecol 85, 69–84. doi:10.1111/1365-2656.12379.

Iino, Y., and Yoshida, K. (2009). Parallel use of two behavioral mechanisms for chemotaxis in
Caenorhabditis elegans. *J Neurosci* 29, 5370–5380. doi:10.1523/JNEUROSCI.3633-
08.2009.

Jonsen, I. D., Basson, M., Bestley, S., Bravington, M. V., Patterson, T. A., Pedersen, M. W., et
al. (2013). State-space models for bio-loggers: A methodological road map. *Deep Sea*
Research Part II: Topical Studies in Oceanography 88–89, 34–46.
doi:10.1016/j.dsr2.2012.07.008.

Kabra, M., Robie, A. A., Rivera-Alba, M., Branson, S., and Branson, K. (2013). JAABA:

Interactive machine learning for automatic annotation of animal behavior. *Nat Meth* 10,

64–67. doi:10.1038/nmeth.2281.

Kaplan, J. M. (1996). Sensory signaling in Caenorhabditis elegans. *Curr Opin Neurobiol* 6,

494–499.

Kass, J., Jacob, T. C., Kim, P., and Kaplan, J. M. (2001). The EGL-3 proprotein convertase

regulates mechanosensory responses of Caenorhabditis elegans. *J Neurosci* 21, 9265–9272.

Kimura, K. D., Fujita, K., and Katsura, I. (2010). Enhancement of odor avoidance regulated by

dopamine signaling in Caenorhabditis elegans. *J Neurosci* 30, 16365–16375.

doi:10.1523/JNEUROSCI.6023-09.2010.

Kohatsu, S., and Yamamoto, D. (2015). Visually induced initiation of Drosophila innate

courtship-like following pursuit is mediated by central excitatory state. *Nature*

Communications 6, 6457. doi:10.1038/ncomms7457.

Kohatsu, S., Koganezawa, M., and Yamamoto, D. (2011). Female contact activates male-

specific interneurons that trigger stereotypic courtship behavior in Drosophila. *Neuron* 69,

498–508. doi:10.1016/j.neuron.2010.12.017.

Kokubun, N., Lee, W. Y., Kim, J.-H., and Takahashi, A. (2015). Chinstrap penguin foraging

area associated with a seamount in Bransfield Strait, Antarctica. *Polar Science* 9, 393–400.

doi:10.1016/j.polar.2015.10.001.

Komatsu, H., Mori, I., Rhee, J. S., Akaike, N., and Ohshima, Y. (1996). Mutations in a cyclic

nucleotide-gated channel lead to abnormal thermosensation and chemosensation in C.

elegans. *Neuron* 17, 707–718.

Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A., and Poeppel, D. (2017).

Neuroscience Needs Behavior: Correcting a Reductionist Bias. *Neuron* 93, 480–490.

doi:10.1016/j.neuron.2016.12.041.

Land, M., Islas-Trejo, A., and Rubin, C. S. (1994). Origin, properties, and regulated expression

of multiple mRNAs encoded by the protein kinase C1 gene of Caenorhabditis elegans. *J.*

Biol. Chem. 269, 14820–14827.

Landhuis, E. (2017). Neuroscience: Big brain, big data. *Nature* 541, 559–561.

doi:10.1038/541559a.

Lee, R. Y., Sawin, E. R., Chalfie, M., Horvitz, H. R., and Avery, L. (1999). EAT-4, a homolog

of a mammalian sodium-dependent inorganic phosphate cotransporter, is necessary for

glutamatergic neurotransmission in caenorhabditis elegans. *J Neurosci* 19, 159–167.

Lewis, J. A., Wu, C. H., Berg, H., and Levine, J. H. (1980). The genetics of levamisole

resistance in the nematode Caenorhabditis elegans. *Genetics* 95, 905–928.

Lockery, S. R. (2011). The computational worm: spatial orientation and its neuronal basis in C.

elegans. *Curr Opin Neurobiol* 21, 782–790. doi:10.1016/j.conb.2011.06.009.

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., et al. (2018).

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.

Nat Neurosci, 1–12. doi:10.1038/s41593-018-0209-y.

Matsumoto, S., Yamamoto, T., Yamamoto, M., Zavalaga, C. B., and Yoda, K. (2017). Sex-

related differences in the foraging movement of streaked shearwaters calonectris

leucomelas breeding on Awashima Island in the sea of Japan. *Ornithological Science* 16,

23–32. doi:10.2326/osj.16.23.

Mizuseki, K., Diba, K., Pastalkova, E., Teeters, J., Sirota, A., and Buzsáki, G. (2014).

Neurosharing: large-scale data sets (spike, LFP) recorded from the hippocampal-entorhinal

system in behaving rats. *F1000Res* 3, 98. doi:10.12688/f1000research.3895.1.

Mizuseki, K., Sirota, A., Pastalkova, E., and Buzsáki, G. (2009). Theta Oscillations Provide

Temporal Windows for Local Circuit Computation in the Entorhinal-Hippocampal Loop.

Neuron 64, 267–280. doi:10.1016/j.neuron.2009.08.037.

Moss, C. F., and Surlykke, A. (2010). Probing the natural scene by echolocation in bats. *Front*

Behav Neurosci 4. doi:10.3389/fnbeh.2010.00033.

Patterson, T. A., Thomas, L., Wilcox, C., Ovaskainen, O., and Matthiopoulos, J. (2008). State-

space models of individual animal movement. *Trends in Ecology & Evolution* 23, 87–94.

doi:10.1016/j.tree.2007.10.009.

Pierce-Shimomura, J. T., Morse, T. M., and Lockery, S. R. (1999). The fundamental role of
pirouettes in Caenorhabditis elegans chemotaxis. *J Neurosci* 19, 9557–9569.

Quinlan, J. R. (1986). Induction of Decision Trees. *Mach Learn* 1, 81–106.
doi:10.1023/A:1022643204877.

Quinlan, J. R. (1992). *C4.5*. San Mateo: Morgan Kaufmann.

Robie, A. A., Hirokawa, J., Edwards, A. W., Umayam, L. A., Lee, A., Phillips, M. L., et al.
(2017). Mapping the Neural Substrates of Behavior. *Cell* 170, 393–406.e28.
doi:10.1016/j.cell.2017.06.032.

Sasakura, H., and Mori, I. (2013). Behavioral plasticity, learning, and memory in C. elegans.
Curr Opin Neurobiol 23, 92–99. doi:10.1016/j.conb.2012.09.005.

Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N. G., Palmer, A. E., and Tsien,
R. Y. (2004). Improved monomeric red, orange and yellow fluorescent proteins derived
from Discosoma sp. red fluorescent protein. *Nat Biotech* 22, 1567–1572.
doi:10.1038/nbt1037.

Stephens, G. J., Johnson-Kerner, B., Bialek, W., and Ryu, W. S. (2008). Dimensionality and
Dynamics in the Behavior of C. elegans. *PLoS Comput Biol* 4, e1000028.
doi:10.1371/journal.pcbi.1000028.

Suo, S., Ishiura, S., and Van Tol, H. H. M. (2004). Dopamine receptors in C. elegans. *Eur. J.*
Pharmacol. 500, 159–166. doi:10.1016/j.ejphar.2004.07.021.

Tanimoto, Y., Yamazoe-Umemoto, A., Fujita, K., Kawazoe, Y., Miyanishi, Y., Yamazaki, S. J.,
et al. (2017). Calcium dynamics regulating the timing of decision-making in C. elegans.
eLife 6, e21629. doi:10.7554/eLife.21629.

Tian, L., Hires, S. A., Mao, T., Huber, D., Chiappe, M. E., Chalasani, S. H., et al. (2009).
Imaging neural activity in worms, flies and mice with improved GCaMP calcium
indicators. *Nat Meth* 6, 875–881. doi:10.1038/nmeth.1398.

Tobin, D., Madsen, D., Kahn-Kirby, A., Peckol, E., Moulder, G., Barstead, R., et al. (2002).
Combinatorial expression of TRPV channel proteins defines their sensory functions and
subcellular localization in C. elegans neurons. *Neuron* 35, 307–318.

Vogelstein, J. T., Park, Y., Ohyama, T., Kerr, R. A., Truman, J. W., Priebe, C. E., et al. (2014).
Discovery of Brainwide Neural-Behavioral Maps via Multiscale Unsupervised Structure
Learning. *Science* 344, 386–392. doi:10.1126/science.1250298.

White, J. G., Southgate, E., Thomson, J. N., and Brenner, S. (1986). The Structure of the
Nervous System of the Nematode Caenorhabditis elegans. *Philosophical Transactions of*
the Royal Society B: Biological Sciences 314, 1–340. doi:10.1098/rstb.1986.0056.

Wiltschko, A. B., Johnson, M. J., Iurilli, G., Peterson, R. E., Katon, J. M., Pashkovski, S. L., et
al. (2015). Mapping Sub-Second Structure in Mouse Behavior. *Neuron* 88, 1121–1135.
doi:10.1016/j.neuron.2015.11.031.

Yamada, Y., Hiryu, S., and Watanabe, Y. (2016). Species-specific control of acoustic gaze by
echolocating bats, Rhinolophus ferrumequinum nippon and Pipistrellus abramus, during
flight. *J Comp Physiol A* 202, 791–801. doi:10.1007/s00359-016-1121-0.

Yamamoto, D., and Koganezawa, M. (2013). Genes and circuits of courtship behaviour in
drosophila males. *Nat Rev Neurosci* 14, 681–692. doi:10.1038/nrn3567.

Yamazoe-Umemoto, A., Fujita, K., Iino, Y., Iwasaki, Y., and Kimura, K. D. (2015). Modulation
of different behavioral components by neuropeptide and dopamine signalings in non-
associative odor learning of Caenorhabditis elegans. *Neurosci Res* 99, 22–33.
doi:10.1016/j.neures.2015.05.009.

Yamazoe-Umemoto, A., Iwasaki, Y., and Kimura, K. (2018). Measuring Spatiotemporal
Dynamics of Odor Gradient for   Small Animals by Gas Chromatography. *BIO-*
PROTOCOL 8. doi:10.21769/bioprotoc.2797.

Yoda, K. (2018). Advances in bio-logging techniques and their application to study navigation
in wild seabirds. *Advanced Robotics* 58, 1–10. doi:10.1080/01691864.2018.1553686.

Yoda, K., Naito, Y., Sato, K., Takahashi, A., Nishikawa, J., Ropert-Coudert, Y., et al. (2001). A

1102  new technique for monitoring the behaviour of free-ranging Adélie penguins. *Journal of*

1103  *Experimental Biology* 204, 685–690.

1104 Yoda, K., Shiomi, K., and Sato, K. (2014). Foraging spots of streaked shearwaters in relation to

1105  ocean surface currents as identified using their drift movements. *Progress in Oceanography*

1106  122, 54–64. doi:10.1016/j.pocean.2013.12.002.

1107 Zhang, S. X., Miner, L. E., Boutros, C. L., Rogulja, D., and Crickmore, M. A. (2018).

1108  Motivation, Perception, and Chance Converge to Make a Binary Decision. *Neuron* 99,

1109  376–388.e6. doi:10.1016/j.neuron.2018.06.014.

1110 Zheng, Y. (2015). Trajectory Data Mining. *ACM Trans. Intell. Syst. Technol.* 6, 1–41.

1111  doi:10.1145/2743025.

1112

Table 1. Summary of recording and analysis conditions of animal behavior.

Animal	worms	penguins	flying seabirds	rats	flies	bats
Condition	naive, mock, preexposed	1	1	1	with or without pheromone	with or without learning
Animal # / condition	50 each	11	54	1	27, 30	3 each
Trajectory # / condition	50 each	11	54	45	27, 30	9 each
Recording time (min-max)	600 s	1626-2787 min	484-8293 min	1031-3194 s	16 s	1.67-2.53 s
Recording time (median)	600 s	2279 min	972 min	1822 min	16 s	1.95 s
Time unit	1 s	1 min	1 min	1 s	0.25 s	0.008 s
Time window for moving average	12 s	20 min	10 min	20 s	-	-
Feature used for EM	dB_Var	V_Ave	V_Ave	dV_Var	-	-
Length unit	mm	m	km	cm	a.u.	m
Approx. distance (min-max)	5-30 mm	1-10 km	100-2000 km	180 × 180 cm	1 - 500 mm	6 m

1113

Table 2. Learning-dependent features extracted from cluster 0 in odor avoidance behavior of wild-type worms.

time	value	Clst0 Dur	Clst1 Dur	V				dV				B				dB				Dir				C				dC				WV
		All	All	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All
-	Ave	0.098	0.057	-	-	-	-	-	-	-	-	-	-	-	-	0.072	-	-	-	-	-	-	-	0.163	-	-	-	0.088	-	-	-	
-	Med	-	-	0.040	-	-	-	-	-	-	-	-	-	-	-	0.102	-	-	-	-	-	-	-	0.160	-	-	-	0.088	-	-	-	
-	Q1st	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.065	-	-	-	-	-	-	-	0.164	-	-	-	0.107	-	-	-	
-	Q3rd	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.083	-	-	-	-	-	-	-	0.161	-	-	-	0.080	-	-	-	
-	Var	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		-	-	-	0.080	-	-	-	
-	1 %ile	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.163	-	-	-	0.090	-	-	-	
-	5 %ile	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.165	-	-	-	0.100	-	-	-	
-	95 %ile	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.153	-	-	-	0.108	-	-	-	
-	99 %ile	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.148	-	-	-	0.060	-	-	-	
1	Ave	-	-	-	0.114			-				-				-				-	-	-	-	-	0.147	0.162	0.162	-	0.116	0.093	0.104	-
2	Ave	-	-	-	0.120		0.049	-				-				-		0.048		-				-	0.148	0.166	0.163	-	0.111	0.105	0.122	-
3	Ave	-	-	-	0.109			-				-				-		0.049		-				-	0.151	0.163	0.162	-	0.110	0.099	0.128	-
3	Med	-	-	-	0.100		0.039	-				-				-				-	-	-	-	-	0.149	0.162	0.162	-	0.114	0.089	0.134	-
3	Var	-	-	-				-				-				-				-	-	-	-	-	0.111	0.087	0.041	-		0.060	0.050	-
4	Ave	-	-	-	0.099			-				-				-		0.050		-				-	0.147	0.166	0.165	-	0.082	0.098	0.122	-
4	Med	-	-	-	0.100			-				-				-				-	-	-	-	-	0.149	0.166	0.165	-	0.085	0.097	0.145	-
4	Var	-	-	-				-				-				-				-				-	0.102	0.083	0.040	-	0.057	0.082	0.082	-
5	Ave	-	-	-				-				-				-		0.048		-				-	0.146	0.163	0.166	-	0.090	0.112	0.148	-
5	Med	-	-	-	0.078			-				-				-				-	-	-	-	-	0.148	0.162	0.165	-	0.093	0.110	0.125	-
5	Var	-	-	-				-				-				-				-				-	0.097	0.092	0.054	-	0.061	0.112	0.078	-
6	Ave	-	-	-	0.072			-				-				-		0.056		-				-	0.148	0.164	0.146	-	0.083	0.112	0.169	-
6	Med	-	-	-	0.075			-				-				-				-	-	-	-	-	0.148	0.166	0.147	-	0.085	0.111	0.149	-
6	Var	-	-	-				-				-				-				-				-	0.089	0.095	0.060	-		0.116	0.081	-

Numbers are the sum of information gains obtained from the comparisons of naive vs. preexp. and mock. vs. preexp. Darker and lighter colors correspond to larger and smaller values, respectively. Blank cells, no value for information gain was obtained. "-", not calculated.

1114

1115

Table 3. Patterns of learning-dependent behavioral features of cluster 0 in odor avoidance behavior of mutant worms.

Strain	Clst0 Dur	Clst1 Dur	V				dV				B				dB				Dir				C				dC			
	All	All	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter	All	Ini	Mid	Ter
wild-type	Up	D/U	Down	Down		Down									Down		Down						Down	Down	Down	Down	Up	Up	Up	Up
egl-3(n589)																							Up	Up	Up	Up				
egl-3(ok979)			Down	Down	Down	Down																	Up	Up	Up	Up				
egl-21																														
dop-3	Up		Down		Down	Down					Up		Up	Up				Up	Down		Down						Up		Up	Up
ocr-2			Up			Up																								
osm-9	Up		Up	Up	Up	Up																								
tax-4						Down									Down	U/D							D/U	D/U	D/U	D/U				
eat-4			Down	Down		Down													Down											
pkc-1															Down		Down						Up				Down			

Darker and lighter colors mean larger and smaller information gain values, respectively. "Up" and "Down" indicates the value changes of pre-exposed worms compared to naive and mock-treated worms. "U/D" or "D/U" indicates the value of pre-exposed worms is higher than naive and lower than mock-treated worms, and *vice versa*. Blank cells, no value for information gain was obtained. The details of information gain are described in Supplementary Tables 2-11.

1116

Trajectory data
(x, y)

Histograms for
basic behavioral
features

Clustering with
the EM algorithm

Mapping the cluster
labels

Calculating behavioral
features of a cluster

Calculating information gain

Plotting the result

a reference
direction

A

Worms
（*Caenorhabditis elegans*）

before learning

after learning

odor
source

start

10 mm

B

V   *dV*   *B*   *dB*

Ave.

Var.

C

Density

dB_Var [(° / s)²]

D

Mapped cluster labels

"Run and pirouette"

E

Events in cluster 0

run   pir.

Events in cluster 1

run   pir.

F

Actual	run	pirouette
Predicted	cluster 0	cluster 1
Sensitivity	0.906	0.955
False positive	0.045	0.094
False negative	0.094	0.045
Specificity	0.955	0.906

Figure 4

A

All

Ini

Mid

Ter

$$Directedness = \frac{\text{-----}}{\sim}$$

B

Parameter	naive		preexp		entropy	Info gain
	Low	High	Low	High		
Original	347		349		1[a]	-
V_Ini	234	113	108	241	0.9017[b]	0.0983[c]

(c = a − b)

A

B

C

D

A

B

	V	*dV*	*B*	*dB*
Overall	0.279	0.260	0.146	0.176
Round 1	0.254	0.200	0.038	0.201
Round 2	0.268	0.218	0.177	0.277
Round 3	0.222	0.171	0.070	0.214
Round 4	0.175	0.235	0.184	0.204
Round 5	0.206	0.229	0.216	0.224
Round 6	0.168	0.221	0.118	0.240
Round 7	0.165	0.188	0.103	0.158
Round 8	0.027	0.087	0	0.062

C

Figure 9

A

High speed video cameras

Obstacle chains

2.5 m

9 m

4.5 m

Z
Y
X

B

Earlier term | $t_e = 0$ | Middle term | $t_m = 0$ | Later term | $t_l = 0$

start

Y
X

—— Initial 3 flights
—— Last 3 flights

0.5 m

R_x (-0.3)

B_obs (-0.3)

X axis (0°)

B_hori (-0.3)

R_y (-0.3)

R_obs (-0.3)

$t_e = -0.3$ s

$t_e = -0.2$ s

$t_e = -0.1$ s

$t_e = 0$ s

0.3 m

C

Features	Info Gain
R_x (-0.3)	0.297
B_hori (-0.3)	0.285
R_obs (-0.3)	0.269
V (-0.3)	0.216
B_hori (-0.2)	0.213
ddB_hori (-0.3)	0.207
V (-0.2)	0.191
R_x (-0.2)	0.191
R_obs (-0.2)	0.191

D

$V (-0.3)$ [m/s]

*

initial   last

$dV (-0.3)$ [m/s²]

initial   last