# Probabilistic data integration identifies reliable gametocyte-specific proteins and transcripts in malaria parasites

Lisette Meerstein-Kessel[1,2], Robin van der Lee[1#a], Will Stone[2,3], Kjerstin Lanke[2], David A Baker[4], Pietro Alano[5], Francesco Silvestrini[5], Chris J Janse[6], Shahid M Khan[6], Marga van de Vegte-Bolmer[2], Wouter Graumans[2], Rianne Siebelink-Stoter[2], Taco WA Kooij[2], Matthias Marti[7], Chris Drakeley[3], Joseph J. Campo[8], Teunis JP van Dam[1#b], Robert Sauerwein[2], Teun Bousema[2¶], Martijn A Huynen[1¶]*

[1] Centre for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboud university medical center, Nijmegen, The Netherlands
[2] Department of Medical Microbiology, Radboud university medical center, Nijmegen, The Netherlands
[3] Department of Immunology and Infection, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom
[4] Department of Pathogen Molecular Biology, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom
[5] Dipartimento Malattie Infettive, Istituto Superiore di Sanità, Rome, Italy
[6] Department of Parasitology, Leiden University Medical Center, Leiden, The Netherlands
[7] Wellcome Trust Center for Molecular Parasitology, Institute of Infection, Immunity and Inflammation, College of Medical Veterinary & Life Sciences, University of Glasgow, Glasgow, Scotland, United Kingdom
[8] Antigen Discovery Inc., Irvine, California, USA
[#a] Current Address: Centre for Molecular Medicine and Therapeutics, Department of Medical Genetics, BC Children's Hospital Research Institute, University of British Columbia, Vancouver, BC, Canada
[#b] Current Address: Theoretical Biology and Bioinformatics, Department of Biology, Utrecht University, Utrecht, The Netherlands

* Corresponding author
Email Martijn.Huijnen@radboudumc.nl (MAH)

[¶] These authors contributed equally to this work.

## Abstract

Plasmodium gametocytes are the sexual forms of the malaria parasite essential for transmission to mosquitoes. To better understand how gametocytes differ from asexual blood stage parasites we performed a systematic analysis of available 'omics data for *P. falciparum* and other *Plasmodium* species. 18 transcriptomic and proteomic data sets were evaluated for the presence of curated "gold standards" of 41 gametocyte-specific versus 46 non-gametocyte genes and integrated using Bayesian statistics, resulting in integrated gametocyte-specificity scores for all *P. falciparum* genes.

To illustrate the utility of the gametocyte score, we explored newly predicted gametocyte-specific genes as potential biomarkers of gametocyte carriage and exposure. We analyzed the humoral immune response in field samples against 34 novel gametocyte-specific antigens and found five antigens to be differentially recognized by gametocyte carriers as compared to malaria-infected individuals without detectable gametocytes. We also validated the gametocyte-specificity of 15 identified gametocyte transcripts on culture material and samples from naturally infected individuals, resulting in eight transcripts that were >1000-fold higher expressed in gametocytes compared to asexual parasites and whose transcript abundance allowed gametocyte detection in naturally infected individuals.

Our integrated genome-wide gametocyte-specificity scores provide a unique resource to identify targets and monitor *P. falciparum* gametocytemia.

## Introduction

Despite a decrease in malaria incidence and mortality over the past two decades, malaria remains a major global health challenge [1,2]. Furthermore, the emergence and spread of insecticide resistance in mosquitoes [3] and artemisinin resistance in *Plasmodium falciparum* (*Pf*) [4–6] threaten recent gains in malaria control. The decline in malaria burden and the necessity to contain artemisinin-resistance have increased interest in malaria elimination that may require interventions that specifically aim to prevent malaria transmission. Malaria transmission depends on male and female gametocytes, the sexually reproducing forms of the *Plasmodium* parasite that are ingested by blood-feeding *Anopheles* mosquitoes. In the mosquito gut, gametocytes may complete the parasite's reproductive cycle and, following sporogonic development, render the mosquito infectious. Factors that govern gametocyte production and infectivity remain poorly understood. Whilst recent studies have shed light on the processes controlling gametocyte commitment [7,8], commitment and maturation of gametocytes may differ between infections and over the course of infections, under influence of environmental and host factors [9,10]. A better understanding of gametocyte dynamics during infections, as well as the development of tools to monitor or target gametocytes, may be informed by high-throughput protein and transcriptome studies [11,12]. In the past 15 years, a number of large-scale studies on *Plasmodium* gametocytes have been reported: The proteome of *Pf* and the rodent malaria parasite *Plasmodium berghei* (*Pb)* have been examined by mass spectrometry [13–21], and the transcriptome of both species by microarray and RNA sequencing [15,20,22–28]. These studies differed in their focus and resolution in examining (sexual) developmental stages and each faced challenges in detecting low abundance proteins [29] and by the purity of parasite populations [13,14]. The use of fluorescent parasites and fluorescence-assisted sorting of staged parasites have recently permitted a better discrimination of proteins in either male or female gametocytes [16,17,20] and have allowed more detailed comparisons of *Plasmodium* life-stages. However, individual studies are still vulnerable to imperfect sample purity, and other sources of uncertainty such as correct gene identification for accurate peptide assignment. These technical and methodological challenges lead to discrepancies between individual studies and hamper firm conclusions about gametocyte-specificity of proteins and transcripts. We utilized the numerous published proteomics and transcriptomics *Plasmodium* data sets in a comprehensive data integration framework to obtain a consensus of gametocyte-specific transcripts and proteins. Our data integration approach, based on a naïve Bayesian classifier, allows: (i) inherent weighting of each contributing data set based on the presence

of expert curated gold standard lists of gametocyte and asexual proteins, (ii) resolution of

conflicting results without disregarding data and (iii) construction of a transparent scoring

system in which the relative contribution of each data set is directly visible. Using this

approach, we propose a robust gametocyte-specificity score for all *Pf* genes that allows a

consensus list of gametocyte-specific genes at protein and transcript level. We illustrate the

utility of our findings by examining naturally acquired responses to newly identified

gametocyte-specific proteins in gametocyte-carriers and non-carriers by protein microarray.

In addition, we confirmed gametocyte-specificity for a selection of gametocyte-specific

transcripts using culture material from geographically distinct *Pf* strains and samples from

naturally infected malaria patients.

## Results

**Weighted integration of proteomics and gene expression data**

Using Bayesian statistics, we integrated *Plasmodium* mass spectrometry and transcript

datasets from 17 different studies on *P. falciparum (Pf;* n=14*)*, *P. berghei (Pb;* n=3*)* and *P.*

*vivax (Pv*; n=1*)*. Since gametocyte biology differs between *Plasmodium* species, scores

were calculated for the total set of *Plasmodium* studies and for *Pf* only. Unsupervised

clustering of genes based on peptide counts or mRNA expression resulted in grouping

according to data acquisition method rather than parasite stage (Fig 1), illustrating the

necessity of a supervised approach to discriminate between gametocyte-specific and non-

gametocyte genes. To objectively assess the value of individual data sets and allow their

assembly into a gametocyte-specificity score, we created a gold standard that served as a

benchmark for every sample. This gold standard was collected from literature review and

comprises two lists; one of asexually expressed proteins, mainly blood stage but also

sporozoite and liver stage, and one of known gametocyte proteins (Supplementary Table

S1). A gametocyte-specificity score was then derived for each gene by comparing its

expression in all studies to the relative expression of the gametocyte and asexual gold

standards in those samples (Supplementary Fig S1-2). Proteins or RNAs detected in a study

with high discriminative power for gametocyte and asexual gold standard genes (ratio of

gametocyte to asexual gold standard genes) received higher gametocyte-specificity scores

than those detected in a study with lower discriminative power. The individual log-

transformed scores per gene were combined for proteomics and transcriptomics data

separately. Scores for *Pf*-only studies (Supplementary Table S2) and all combined data sets

were highly correlated (r=0.9867 and r=0.9514 for proteomics and transcriptomics,

respectively) and the latter were used in the remainder of the manuscript.

The distribution of scores for proteins and transcripts are presented for the two sets of genes

of the gold standard as well as for all other genes (Fig 2A). As expected, the gametocyte

and asexual gold standard set of genes are perfectly separated by their respective

proteomics-derived scores and show only little overlap in their transcriptomics-derived

scores (Fig 2B).  The shift in the density peak of the proteomics compared to transcriptomics

is due to an inherent property of the method that gives a negative score for gametocyte-

specificity to all proteins that were not detected in the proteomics studies (n=1583). The

highest scoring 100 genes for proteomics and transcriptomics contained 26 (63.4 %) and 15

(36.6 %) of the 41 gold standard gametocyte genes, respectively (Fig 2C), indicating both a

powerful gametocyte gold standard and that many other genes are equally specific for

gametocytes.

Translationally repressed genes are common in late stage female gametocytes [30,31] and

are detectable by high transcriptomics and low proteomics score. In our analysis 461 genes

have this profile (Supplementary Table S3), including genes that are known to be

translationally repressed like Pfs28 [30,32] and 186 genes with a previously reported bias

towards expression in female gametocytes [20].

**Cross-validation illustrates the improved predictive power of the integrated data.** Ten-

fold cross-validation was performed using random subsamples of the gold standard lists to

predict the ranks of left-out genes. The resulting proteomics ranking shows near perfect

sensitivity, with all but two gold standard gametocyte genes ranking higher than the gold

standard asexual genes (Fig 3). The added value of our integrated approach is illustrated by

the receiver operating characteristic curve where the integration of data sets gave higher

sensitivity and areas under the curve for both proteomics and transcriptomics than any

individual study (Fig 3). Using the Bayesian integration based on the complete gold

standards, we ranked all *Pf* proteins by giving them a gametocyte-specificity score

(Supplementary Table S2). All proteins with a score >5 (n=602) were considered

gametocyte-specific. Most of these have not consistently been described as "specific" or

"enriched" in gametocytes in the original data sets (Fig 4A and Supplementary Table S4).

Previous studies defined 315 [13] to 1725 [20] proteins as gametocyte-specific for *Pf*. Not

only did our integrated approach lead to a better recovery of gold standard listed known

gametocyte proteins, we also identified 178 genes with undescribed function as gametocyte-

specific (Supplementary Table S2). We further identify a number of proteins as gametocyte-

specific even though they had been reported as asexual by previous studies (Fig 4B).

A recent proteomics study of male and female *Pf* gametocytes [33], not included in our

original analysis, was used to test the robustness of our scores. When we included this data

set in our final Bayesian proteomics scores, both gametocyte scores and gene ranks before

and after addition of this data set were highly correlated (Pearson's r=0.997 and Spearman's

rho=0.995, respectively). Furthermore, the top 100 gametocyte proteins did not change and

the top 602 proteins were 96% identical (578 of 602). Taken together, cross-validation and

independent data suggest that the integrated gametocyte-specificity score is robust and

contains potential novel gametocyte markers.

**Predicted gametocyte-specific proteins are recognized by gametocyte-carriers.**

As an illustration of the utility of gametocyte-specific proteins as markers of gametocyte

exposure, we utilized protein microarray data from a study that aimed to characterize the

immune profile associated with transmission-reducing immunity in naturally infected

gametocyte carriers (Stone, Campo *et al.* 2017 under review). For the current study, we

compared responses to our gold standard gametocyte genes (n=41) and novel gametocyte

genes from our 100 highest scoring proteins that were on the array (n=34). Antibody

prevalences for these genes were compared between Gambian gametocyte carriers and

Gambians who carried asexual parasites but not gametocytes by microscopy. Antibody

responses to these gametocyte-specific genes were generally higher in gametocyte carriers

(Fig 5A). A significantly higher antibody prevalence in gametocyte carriers was detected for

five of the novel gametocyte antigens (Fig 5B, p<0.05 in Fisher's exact, corrected for

multiple testing, Supplementary Table S5, Supplementary Fig S3). Only two of these five

have an assigned function – a DNA ligase, and Gamete egress and sporozoite-traversal

protein (GEST). For two of the three remaining *Plasmodium* proteins, we were able to

predict a function based on homology, using the sensitive homology detection tool HHpred

[34]. PF3D7_1251000 is homologous to the co-chaperone HSP20 heat shock protein and

PF3D7_1439600 is homologous to the MLRQ subunit of complex IV of the oxidative

phosphorylation, underlining the enrichment in mitochondrial proteins as discussed below

(Supplementary Table S6 includes homology predictions for all conserved, highly

gametocyte-specific *Plasmodium* proteins).

**Gametocyte-specific RNA transcripts detect (submicroscopic) gametocyte carriage**

Of the 100 highest-scoring transcripts, 15 non-gold standard candidates were randomly

selected for qRT-PCR validation (Table 2). Mature gametocytes of four *Pf* strains from

different geographical origins were compared to asexual blood stage parasites. The

minimum transcript abundance difference ($Ct_{Asexuals} - Ct_{Gametocytes}$) ranged from 4.76 to 14.95

(Fig 6A and Supplementary Table S7 qPCR & primers), reflecting 27.1 to 31,500-fold higher

transcript numbers in gametocytes compared to asexual parasites and confirming

pronounced upregulation of all selected targets in gametocytes. With a very conservative

threshold of 1,000-fold enrichment in three of the four strains tested (Fig 6A), eight of the 15

tested transcripts were highly specific to gametocytes. Transcript abundance in ring stage

parasites was assessed and compared to Pfs25 mRNA, an established and highly abundant

yet intron-less female gametocyte specific transcript [11,35]. Five out of eight gametocyte

specific transcripts were undetectable in asexual ring stage at ≤$10^5$ parasites/mL, similar in

specificity to Pfs25 (Fig 6B); the five most sensitive gametocyte markers detected

gametocytes across the range of $10^2 - 10^6$ gametocytes/mL (Fig 6C). In RNA samples from

a previously reported clinical trial conducted in Kenya [36], all eight gametocyte markers

detected gametocytes at densities below $10^3$/mL (Fig 6D).

**Gametocyte-specific proteins are enriched for cytoskeletal movement and**

**metabolism functions**

To uncover novel characteristics underlying gametocyte function, we analyzed over

represented gene ontology (GO) terms in our integrated consensus gametocyte proteins.

The 100 highest ranked proteins were examined for enrichment of GO terms that reflect

specific biological functions. Microtubule based movement, metabolism of carboxylic acids

and metabolism of nucleic acids were highly enriched among gametocyte proteins

(Supplementary Fig S4). Four out of the six putative *Pf* dynein heavy chain proteins are

found back among the 100 most gametocyte-specific proteins, alongside a tubulin gamma

chain and a tubulin chaperone. The importance of DNA elongation and ligation processes is

reflected in GO term associations as well as in antibody response to a DNA ligase (Fig 5B).

The "classic" Go term enrichment calculation was complemented by a rank-based gene set

enrichment analysis (GSEA). GSEA uses all *Pf* genes and their respective (proteomics

derived) gametocyte-specificity score and thus does not include an arbitrary cutoff of the

proteins that are or are not gametocyte-specific. It confirmed the above-mentioned results,

and in addition to those the terms "mitochondrial protein complex" (GO:0098798) and "TCA"

(GO:0006099), were observed to be enriched, stressing both mitochondrial location and

processes.

Although male gametocytes carry pre-synthesized proteins to rapidly form eight motile

gametes upon activation in the mosquito midgut, we did not observe flagellum associated

terms in the GSEA. The reason for this is that current GO term annotation for *Pf* has only

one gene with the "cilium" GO term for cellular component (GO:0005929; PF3D7_1025500),

one for "axoneme" (GO:0005930; PF3D7_0828700) and none with the biological-process

terms "cilium or flagellum-dependent cell motility" (GO:0001539) or "axoneme assembly"

(GO:0035082). We supplemented the GO annotation with a list of 28 *Pf* cilium genes
(Methods). The newly assembled "cilium" GO term now acquired the highest enrichment
score in the GSEA (Supplementary Fig S5). This may reflect the formation of the flagella of
the microgamete but may also (partially) reflect intracellular trafficking and or be associated
with genome replication as the term has overlap with the genes annotated for microtubule
processes (Supplementary Table S5 GO terms).
219

## Discussion

220
221
Combining proteomic and transcriptomic data from 17 sources, we present an integrated
consensus score for gametocyte-specific proteins and transcripts. We predict 602
gametocyte-enriched proteins of which 186 are currently without ascribed function. We
illustrate the potential utility of our gametocyte score by providing evidence for differential
recognition of gametocyte proteins by naturally infected gametocyte carriers and the
sensitive detection of mRNA of novel gametocyte transcripts in field samples.
228
The gametocyte proteome of *P. falciparum (Pf)* has been assessed repeatedly. Individual
lists of gametocyte-specific proteins [13,14,16,17,19,20] have unavoidable limitations related
to comparator (asexual) parasite stages, sample purity, assay sensitivity and arbitrary cut-
offs used to define gametocyte-specificity and show only partial agreement. To acquire a
more robust gametocyte specificity score, we integrated data from these individual studies,
along with studies of purified asexual parasites and related *Plasmodium* species. Including
gene expression data from multiple species generally increases the likelihood that the
combined gene expression data reflect underlying biology, as observed in the Apicomplexa
[37,38]. We applied a Bayesian classifier first applied to 'omics data by Jansen and
colleagues [39] and adapted by Van der Lee and colleagues to identify genes involved in
anti-viral immune responses [40]. The probabilistic approach combines the evidence from all
studies in an unbiased way, without giving *a priori* preference of one study over another.
Instead, the measurements of all studies were weighted inherently during the scoring
process by assessing the retrieval of a gold standard set of genes. As these gametocyte and
asexual gold standard sets are of central importance to the study, they have undergone
expert curation (see Methods and Acknowledgments). The power of this integrative
approach lies not only in weighting data sets by the retrieval of gold standard genes but also
in the opportunity to exclude proteins from the gametocyte-specific list by appreciating their
presence in (other) asexual samples.  A further strength of the approach is that it allows the
ranking of gametocyte proteins that have only been reported in a subset of studies. Our

integration of data sets reveals that 602 proteins are likely to be specific to gametocytes

although very few gametocyte-specific proteins were detected in every underlying dataset

and seven proteins had never before been reported as gametocyte-specific.

A general limitation of all mass spectrometry (MS) studies is their bias toward highly

abundant proteins. Proteins with low-level expression may be missed in a bulk proteome

analysis. After integration of the MS studies listed in Table 1, 1583 *Pf* proteins were never

detected, representing approximately 28% of all proteins encoded by *Pf*. Some of these

might be of too low abundance or expressed during sporozoite or gamete, ookinete and liver

stage, which are underrepresented or not included in our data, respectively. New advances

in MS that include the sensitive detection of peptides from currently understudied

*Plasmodium* life stages may shed light on these currently uncharacterized genes. In addition,

approaches that focus specifically on post translational modifications like phosphorylation of

proteins as has been done for asexual parasites [41–44] may add new lines of evidence

towards gametocyte-specific functions of proteins. Our approach suggests that the currently

available MS data is sufficiently comprehensive to identify stage specific proteins when

analysed in an integrative approach. We examined this directly by incorporating a new *Pf*

gametocyte MS study [33] in our scoring. The authors reported 44 new gametocyte-specific

proteins that were not reported by earlier studies. We compared this data set to our

integrated data set and found 24 of the 44 had been detected in one or more erythrocytic

stages or sporozoites [14,18,41,45,46] while 11 others had been identified in a (single)

gametocyte sample before (Supplementary Fig S6). Importantly, the scores and top 100

gametocyte genes remained unaltered by integrating this new dataset.

The ranking of gametocyte-specificity that we provide here can i) aid in understanding the

biology of this life stage and ii) improve diagnostics related to gametocyte exposure and

carriage. Regarding gametocyte biology, our high-ranking gametocyte-specific genes are

enriched for mitochondrial, metabolism and microtubule processes and DNA replication. The

enrichment of mitochondrial localization and process is consistent with what we know about

the enlarged mitochondrion of gametocyte stages [47] and increased activity of the citric acid

cycle [48]. DNA replication terms are highly enriched which is consistent with what happens

in the subsequent life stage in which the (micro)gamete rapidly duplicates its genomic DNA

three times. Regarding the use of the gametocyte score to inform gametocyte diagnostics,

diagnostics can directly detect nucleic acids specific to gametocytes [11] or detect antibody

responses reflecting past/recent exposure as is increasingly used for asexual *P. falciparum*

and *P. vivax* parasites [49,50]. We use our integrated gametocyte list to explore its utility for

both approaches. We validated 15 transcript targets in four different *Pf* strains, comparing

transcript abundance in gametocytes and asexual parasites. All tested targets were enriched

in gametocytes. Five targets were tested for their sensitivity and can recognize 100

gametocytes/mL, while the signal is undetectable when fewer than $10^5$-$10^6$ ring stage

parasites/mL are present. In practical terms, these markers may be used to reliably detect

gametocytes at densities well below the microscopic threshold of detection in samples

without high-densities of asexual parasites, similar to the gametocyte marker that is currently

most widely used, the female gametocyte-specific Pfs25 [35].

As an alternative approach to the detection of gametocyte carriage in populations, we

utilized a gametocyte-enriched protein microarray (Stone, Campo et al. under review) to

determine antibody responses to genes that we here describe as highly gametocyte-specific.

The expression system used for the array has known limitations with the expression of

conformational proteins [51] and should thus be considered a 'rule in' rather than 'rule out'

approach to immune recognition. Moreover, the array was constructed with the aim of

detecting surface proteins or exported proteins whilst our list does not require these

characteristics. Only 34 of our top 100 novel gametocyte antigens were thus printed on this

array. Antibody responses to five gametocyte proteins were significantly more prevalent in

gametocyte carriers than in carriers of the asexual blood stage only. This is the first

evidence that antibody responses may be indicative of current gametocyte carriage. Refined

studies with longitudinal sampling and gametocyte detection by sensitive qRT-PCR

methodologies are needed to formally assess antibody kinetics in relation to gametocyte

exposure and determine whether recent markers of exposure to blood stage antigens [49]

can be complemented by a set of markers for recent or long-term gametocyte exposure.

We described the assembly of a curated gold standard set of gametocyte and asexual

proteins and used this new resource to rank the likelihood of all *Pf* proteins and transcripts

being specific to the gametocyte stage. Data from 17 publicly available studies were

integrated to resolve partially conflicting evidence. The resulting consensus lists can be used

for guidance of future investigations as we have shown the value of our predictions by in

vitro validation.

## Materials and Methods
316

**Assembly of a gold standard for gametocyte and asexual proteins to weigh whole**

**proteome/transcriptome data sets**

To build a gold standard against which the performance of individual data sets could be

assessed, we identified proteins that are known to be expressed in either asexual parasites

(mostly blood stage, also including sporozoites and liver stage) or gametocytes. This list was

initially informed by literature review (Supplementary table S1) for expression in the

respective stages, supplemented with a number of blood stage or transmission blocking

vaccine candidates, and additionally reviewed and updated by an international group of

gametocyte experts (including the authors DAB, PA, FS, CJJ, SMK, TWAK, MM, CD, RS

and TB). The asexual gold standard list contains 46 proteins; the gametocyte list contains 41

proteins. These gold standard lists (Supplementary Table 1) represent the balance between

very strict inclusion criteria and sufficient set size to evaluate the quality of all data sets

integrated. We tested for the detection of these proteins or transcripts in the respective

samples, using a Bayesian statistics approach that we have successfully applied previously

for genes involved in anti-viral immune defense [40].

**Data selection and integration**

Data sets that measured protein and transcript abundance in *Pf* gametocytes were balanced

with data from other life stages and supplemented with studies of the rodent malaria parasite

*P. berghei*. One MS study on *P. vivax* was included as it is based on of *ex vivo* blood

material as opposed to all other studies that used *in vitro* cultivated parasites. Unique

peptide counts were retrieved from plasmoDB (version 28) in which sequenced peptides

from the published studies are always mapped to the most recent genome annotation, or

supplementary material of the respective studies. Many aspects determine how well a

protein is represented in proteomics data that are obtained via MS, like its length or

posttranslational modifications. Remapping original MS data to newly annotated genomes

improves the quality of the predicted proteins [19]. We were however not able to retrieve

those data from the studies [13,14,16] and therefore decided to take those proteins at face

value. Notice that also these early studies contribute significantly to our integrated lists.

Expression percentiles were retrieved from plasmoDB (version 28) or calculated from raw

data in the respective supplementary material. Gametocyte samples were summarized if

applicable (using the maximum peptide count/expression percentile of different stages or

male and female gametocytes) as were asexual samples, only considering the highest

expression in any sample or time point.

For MS and transcriptomics data sets, separate scores for gametocyte-specificity of any *Pf*

gene have been calculated. In brief, protein or transcript expression has been categorized

from absent to high expression levels as given by number of unique peptides or expression

percentiles, respectively. For each of the respective bins, a score was calculated depending

on the relative retrieval of gametocyte and asexual gold standard genes. The log ratio of

these retrieved genes defined the score for all other genes within the same bin. The final

gametocyte score calculates as the prior probability of a gene being gametocyte-specific that

is updated using the contributions of the data sets:

$$GametocyteScore = log_2\left(\frac{P_{gct}}{P_{\sim gct}}\right) + \sum_{i=1}^{n} log_2\left(\frac{P(data_i|Gct)}{P(data_i|\sim Gct)}\right) \qquad (1)$$

$$with\ \frac{P(data_i|Gct)}{P(data_i|\sim Gct)} = \frac{gametocyteGS_i}{asexualGS_i} \qquad (2)$$

where gametocyteGS and asexualGS are the fractions of retrieved gametocyte and asexual

gold standard genes in sample i, respectively. We used a pseudocount of 1 if necessary to

prevent division by zero if none of the gold standard genes was retrieved in this specific

sample and bin. It was assumed that the likelihood of a gene to be either gametocyte or

asexual specific is equally high, thus the (log-transformed) prior equals 0 and the final score

depends solely on the integrated data. In the selection of a set of proteins that we assigned

to be gametocyte-specific we chose a cutoff score of 5.0 (proteomics-derived). The cutoff

score of 5.0 can be interpreted as: a gene has to be $2^5$ = 32 times more likely to be

gametocyte-specific than asexual specific. The score of 5.0 was based on the behavior of

the gold standard genes. Out of the 41 gametocyte gold standard genes, 37 have a score

higher than 5.0, while none of the asexual gold standard genes do.

When applicable, genes from *Pb* and *Pv* were treated as their respective *Pf* orthologs as

retrieved from plasmoDB [52], to be able to integrate all data sets. When no ortholog is

known, the respective non-*Pf* data sets did not contribute to the score of this particular gene.

Scores using *Pf* data exclusively were also calculated (Supplementary Table S2 includes all

scores and rankings with expression information from all integrated studies).

**Cross-validation of the scoring method**

We performed a ten-fold cross-validation to assess the predictive performance of the

integrated gametocyte-specificity score (i.e. its ability to discriminate known gametocyte vs.

asexual genes). For that, we subsampled both gold standard gene sets ten times (folds),

without replacement (i.e. each gene is selected exactly once). Then for each fold we re-

weighed and integrated the data sets based on nine-tenth of gold standard genes, and

collected the ranks of the one-tenth of genes that were left out in that particular fold. A ROC

curve was constructed based on those ranks. Using the same strategy, ROC curves for

individual data sets that comprised both gametocyte and asexual samples were constructed

for comparison.

**Protein microarray to measure humoral immune responses**

A protein microarray that was enriched for gametocyte proteins was produced and probed
as described earlier for a study aiming to unravel the immune signature of naturally acquired
transmission-reducing immune responses in gametocyte carriers (Stone, Campo et al. 2017
under review). As a control group, Gambian asexual parasite carriers without gametocytes
by microscopy were included in the probing. For the current study array data from this
control group (n=63) and Gambian gametocyte carriers were used (n=164) [53–58]. For
these populations, responses to 34 newly defined highly gametocyte-specific antigens (from
24 genes) were compared between gametocyte carriers and non-carriers. Seropositivity for
each of the antigens was determined using a mixture model-based cutoff and related to
gametocyte carriage using Fisher's exact test, corrected for multiple testing (Benjamini-
Hochberg) of a total of 75 antigens (including 41 antigens from the gametocyte gold
standard).
403
**Transcript abundance in different life stages and strains**
The abundance of 15 predicted gametocyte-specific targets was measured in asexual
parasites and gametocytes of four different *Pf* strains from *in vitro* culture. The *Pf* strains
used are of West African (NF54, NF166, NF175) and Southeast Asian origin (NF135). All
strains were cultured and synchronized as described previously [20]. Using established
standard curves, the same concentrations of parasites were compared for Ct values in qRT-
PCR (for primers, see Supplementary Table S5). Extracted nucleic acids were DNase-
treated before reverse transcription when introns were absent from the targets. Initial
comparison was between mixed asexual blood stage parasites (considering the lowest Ct
measured in any strain and replicate) and stage V gametocyte (highest Ct measured per
strain and replicate). Promising targets with a high $Ct_{Asexuals}$ - $Ct_{Gametocytes}$ were further
examined in serial dilutions of stage V gametocytes and synchronized asexual material of
the strain NF54 (10, 20, 30, 40 hours post invasion, resembling early rings, late rings,
trophozoites and schizonts, respectively). All qRT-PCR reactions were analyzed in technical
triplicates, from biological triplicates (NF54) or duplicates (remaining strains).
RNA samples were used from a clinical malaria trial conducted in Western Kenya [36].
Samples from days 3 and 7 after treatment were selected to ensure a range of (low-density)
gametocyte carriage to test qRT-PCR sensitivity.
422
**Go term enrichment in top 100 proteins or rank-based enrichment**
Current GO term annotation for *Pf* was retrieved from plasmoDB (release 30) and analyzed
using the topGO R package [59] for the enrichment of terms in the 100 highest scoring
proteins versus all *Pf* proteins. Semantic clustering of the significant GO terms in Biological
Process ontology was done with the Revigo webtool for *Pf* [60]. Second, gene set

enrichment analysis (GSEA) based on all *Pf* proteins and their ranks and scores was

performed using the software available at

http://software.broadinstitute.org/gsea/downloads.jsp. Based on the cilium genes reported by

the Syscilia consortium [61], we assembled a "cilium" GO term of mixed ontology

(GO:9999999) for *Pf* with 28 predicted orthologs (Supplementary Table S6).

**Data availability**

All data generated or analysed during this study are included in this published article (and its

Supplementary Information files).

-    The gold standard lists (Supplementary Table S1)

-    Bayesian gametocyte scoring for proteomics and transcriptomics data. Includes *Pf*

439              only-scores and expression values for any gene and individual data set

440              (Supplementary Table S2)

441         -    Potential translationally repressed genes with high transcriptomics score (>7) and

442              low proteomics score (<-10, Supplementary Table S3)

443         -    Overview over previously reported gametocyte-specificity per study (Supplementary

444              Table S4)

445         -    Seroprevalence in The Gambia in gametocyte carriers and non-carriers

446              (Supplementary Table S5)

447         -    Function predictions for highly gametocyte-specific proteins with lacking annotation

448              (Supplementary Table S6)

449         -    Transcript validation in 15 targets, including primer sequences for qRT-PCR

450              (Supplementary Table S7)

451         -    GO term analyses with cilium genes (Supplementary Table S8)

# References

1.  WHO. Reversing the Incidence of Malaria 2000–2015. In 2015.
2.  Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature. 2015;526(7572):207–11.
3.  Ranson H, Lissenden N. Insecticide Resistance in African Anopheles Mosquitoes: A Worsening Situation that Needs Urgent Action to Maintain Malaria Control. Trends Parasitol. Elsevier Ltd; 2016;32(3):187–96.
4.  Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, et al. Spread of Artemisinin Resistance in *Plasmodium falciparum* Malaria. N Engl J Med. 2014;371(5):411–23.
5.  Takala-Harrison S, Jacob CG, Arze C, Cummings MP, Silva JC, Dondorp AM, et al. Independent emergence of artemisinin resistance mutations among Plasmodium falciparum in Southeast Asia. J Infect Dis. 2015;211(5):670–9.
6.  Ménard D, Khim N, Beghain J, Adegnika AA, Shafiul-Alam M, Amodu O, et al. A Worldwide Map of Plasmodium falciparum K13-Propeller Polymorphisms. N Engl J Med. 2016;374(25):2453–64.
7.  Kafsack BFC, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et al. A transcriptional switch underlies commitment to sexual development in malaria parasites. Nature. 2014;507(7491):248–52.
8.  Pelle KG, Oh K, Buchholz K, Narasimhan V, Joice R, Milner DA, et al. Transcriptional profiling defines dynamics of parasite tissue sequestration during malaria infection. Genome Med. 2015;7(1):1–20.
9.  Johnston GL, Smith DL, Fidock DA. Malaria's Missing Number: Calculating the Human Component of R0 by a Within-Host Mechanistic Model of Plasmodium falciparum Infection and Transmission. PLoS Comput Biol. 2013;
10. Bousema T, Drakeley C. Epidemiology and infectivity of Plasmodium falciparum and Plasmodium vivax gametocytes in relation to malaria control and elimination. Clinical Microbiology Reviews. 2011. p. 377–410.
11. Joice R, Narasimhan V, Montgomery J, Sidhu AB, Oh K, Meyer E, et al. Inferring Developmental Stage Composition from Gene Expression in Human Malaria. PLoS Comput Biol. 2013;9(12):1–13.
12. Proietti C, Doolan DL. The case for a rational genome-based vaccine against malaria. Frontiers in Microbiology. 2015.
13. Lasonder E, Ishihama Y, Andersen JS, Vermunt AMW, Pain A, Sauerwein RW, et al. Analysis of the Plasmodium falciparum proteome by high-accuracy mass spectrometry. Nature. 2002;419(6906):537–42.
14. Florens L, Washburn MP, Raine JD, Anthony RM, Grainger M, Haynes JD, et al. A proteomic view of the Plasmodium falciparum life cycle. Nature. 2002;419(6906):520–6.
15. Hall N, Karras M, Raine JD. A Comprehensive Survey of the Plasmodium Life Cycle by Genomic, Transcriptomic, and Proteomic Analyses. Science (80- ). 2005;307(5706):82–6.
16. Khan SM, Franke-Fayard B, Mair GR, Lasonder E, Janse CJ, Mann M, et al. Proteome Analysis of Separated Male and Female Gametocytes Reveals Novel Sex-Specific Plasmodium Biology. Cell. 2005;121(5):675–87.
17. Silvestrini F, Lasonder E, Olivieri A, Camarda G, van Schaijk B, Sanchez M, et al. Protein Export Marks the Early Phase of Gametocytogenesis of the Human Malaria Parasite Plasmodium falciparum. Mol Cell Proteomics. 2010;9(7):1437–48.
18. Oehring SC, Woodcroft BJ, Moes S, Wetzel J, Dietz O, Pulfer A, et al. Organellar proteomics reveals hundreds of novel nuclear proteins in the malaria parasite Plasmodium falciparum. Genome Biol. BioMed Central Ltd; 2012;13(11):R108.
19. Tao D, Ubaida-Mohien C, Mathias DK, King JG, Pastrana-Mena R, Tripathi A, et al. Sex-

partitioning of the Plasmodium falciparum Stage V Gametocyte Proteome Provides Insight
into falciparum-specific Cell Biology. Mol Cell Proteomics. 2014;13(10):2705–24.

20. Lasonder E, Rijpma SR, van Schaijk BCL, Hoeijmakers WAM, Kensche PR, Gresnigt MS, et al.
Integrated transcriptomic and proteomic analyses of P. falciparum gametocytes: molecular
insight into sex-specific processes and translational repression. Nucleic Acids Res.
2016;44(13):gkw536.

21. Suárez-Cortes P, Sharma V, Bertuccini L, Costa G, Bannerman N-L, Sannella AR, et al.
Comparative proteomics and functional analysis reveal a role of P. falciparum osmiophilic
bodies in malaria parasite transmission. Mol Cell Proteomics. 2016 Jul 18;

22. Bozdech Z, Llinás M, Pulliam BL, Wong ED, Zhu J, DeRisi JL. The transcriptome of the
intraerythrocytic developmental cycle of Plasmodium falciparum. PLoS Biol. 2003;1(1):85–
100.

23. Le Roch KG, Zhou Y, Blair PL, Grainger M, Moch JK, Haynes JD, et al. Discovery of gene
function by expression profiling of the malaria parasite life cycle. Science (80- ).
2003;301(September):1503–8.

24. Young JA, Fivelman QL, Blair PL, de la Vega P, Le Roch KG, Zhou Y, et al. The Plasmodium
falciparum sexual development transcriptome: A microarray analysis using ontology-based
pattern identification. Mol Biochem Parasitol. 2005;143(1):67–79.

25. Llinas M. Comparative whole genome transcriptome analysis of three Plasmodium falciparum
strains. Nucleic Acids Res. 2006;34(4):1166–73.

26. Otto TD, Wilinski D, Assefa S, Keane TM, Sarry LR, Böhme U, et al. New insights into the
blood-stage transcriptome of Plasmodium falciparum using RNA-Seq. Mol Microbiol.
2010;76(1):12–24.

27. López-Barragán MJ, Lemieux J, Quiñones M, Williamson KC, Molina-Cruz A, Cui K, et al.
Directional gene expression and antisense transcripts in sexual and asexual stages of
Plasmodium falciparum. BMC Genomics. 2011;12(1):587.

28. Otto TD, Böhme U, Jackson AP, Hunt M, Franke-Fayard B, Hoeijmakers WAM, et al. A
comprehensive evaluation of rodent malaria parasite genomes and gene expression. BMC
Biol. 2014;12(1):86.

29. Wasinger VC, Zeng M, Yau Y. Current status and advances in quantitative proteomic mass
spectrometry. Int J Proteomics. 2013;2013:180605.

30. Mair GR, Braks JAM, Garver LS, Wiegant JCAG, Hall N, Dirks RW, et al. Regulation of Sexual
Development of Plasmodium by Translational Repression. Science (80- ). 2006;313:667–9.

31. Mair GR, Lasonder E, Garver LS, Franke-Fayard BMD, Carret CK, Wiegant JCAG, et al. Universal
features of post-transcriptional gene regulation are critical for Plasmodium zygote
development. PLoS Pathog. 2010;6(2).

32. Miao J, Fan Q, Parker D, Li X, Li J, Cui L. Puf Mediates Translation Repression of Transmission-
Blocking Vaccine Candidates in Malaria Parasites. PLoS Pathog. 2013;

33. Miao J, Chen Z, Wang Z, Shrestha S, Li X, Li R, et al. Sex-Specific Biology of the Human Malaria
Parasite Revealed from the Proteomes of Mature Male and Female Gametocytes. Mol Cell
Proteomics. 2017;

34. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection
and structure prediction. Nucleic Acids Res. 2005;33(SUPPL. 2):244–8.

35. Stone W, Sawa P, Lanke K, Rijpma S, Oriango R, Nyaurah M, et al. A Molecular Assay to
Quantify Male and Female Plasmodium falciparum Gametocytes: Results From 2 Randomized
Controlled Trials Using Primaquine for Gametocyte Clearance. J Infect Dis. 2017;

36. Graves PM, Gelband H, Garner P. Primaquine or other 8-aminoquinoline for reducing
Plasmodium falciparum transmission. Cochrane database Syst Rev. 2015;

37. Huynen MA, Snel B, Van Noort V. Comparative genomics for reliable protein-function
prediction from genomic data. Trends in Genetics. 2004.

38. Butler CL, Lucas O, Wuchty S, Xue B, Uversky VN, White M. Identifying novel cell cycle

proteins in apicomplexa parasites through co-expression decision analysis. PLoS One. 2014;

39.  Jansen R, Yu HY, Greenbaum D, Kluger Y, Krogan NJ, Chung SB, et al. A Bayesian networks
approach for predicting protein-protein interactions from genomic data. Science (80- ). 2003;

40.  van der Lee R, Feng Q, Langereis MA, ter Horst R, Szklarczyk R, Netea MG, et al. Integrative
Genomics-Based Discovery of Novel Regulators of the Innate Antiviral Response. PLoS
Comput Biol. 2015;11(10).

41.  Treeck M, Sanders JLL, Elias JEE, Boothroyd JCC. The Phosphoproteomes of Plasmodium
falciparum and Toxoplasma gondii Reveal Unusual Adaptations Within and Beyond the
Parasites' Boundaries. Cell Host Microbe. Elsevier; 2011;10(4):410–9.

42.  Lasonder E, Green JL, Camarda G, Talabani H, Holder A a., Langsley G, et al. The plasmodium
falciparum schizont phosphoproteome reveals extensive phosphatidylinositol and cAMP-
protein kinase A signaling. J Proteome Res. 2012;11(11):5323–37.

43.  Pease BN, Huttlin EL, Jedrychowski MP, Talevich E, Harmon J, Dillman T, et al. Global analysis
of protein expression and phosphorylation of three stages of plasmodium falciparum
intraerythrocytic development. J Proteome Res. 2013;12(9):4028–45.

44.  Lasonder E, Green JL, Grainger M, Langsley G, Holder A a. Extensive differential protein
phosphorylation as intraerythrocytic *Plasmodium falciparum* schizonts develop into
extracellular invasive merozoites. Proteomics. 2015;n/a-n/a.

45.  Bowyer PW, Simon GM, Cravatt BF, Bogyo M. Global profiling of proteolysis during rupture of
Plasmodium falciparum from the host erythrocyte. Mol Cell Proteomics. 2011;10(5):M110
001636.

46.  Lindner SE, Swearingen KE, Harupa A, Vaughan AM, Sinnis P, Moritz RL, et al. Total and
putative surface proteomics of malaria parasite salivary gland sporozoites. Mol Cell
Proteomics. 2013;12(5):1127–43.

47.  Okamoto N, Spurck TP, Goodman CD, McFadden GI. Apicoplast and mitochondrion in
gametocytogenesis of Plasmodium falciparum. Eukaryot Cell. 2009;8(1):128–32.

48.  MacRae JI, Dixon MW, Dearnley MK, Chua HH, Chambers JM, Kenny S, et al. Mitochondrial
metabolism of sexual and asexual blood stages of the malaria parasite Plasmodium
falciparum. BMC Biol. 2013;11(1):67.

49.  Helb DA, Tetteh KKA, Felgner PL, Skinner J, Hubbard A, Arinaitwe E, et al. Novel serologic
biomarkers provide accurate estimates of recent Plasmodium falciparum exposure for
individuals and communities. Proc Natl Acad Sci U S A. 2015;1501705112-.

50.  Cutts JC, Powell R, Agius P a, Beeson JG, Simpson J a, Fowkes FJI. Immunological markers of
Plasmodium vivax exposure and immunity: a systematic review and meta-analysis. BMC Med.
2014;

51.  Crompton PD, Kayala MA, Traore B, Kayentao K, Ongoiba A, Weiss GE, et al. A prospective
analysis of the Ab response to Plasmodium falciparum before and after a malaria season by
protein microarray. Proc Natl Acad Sci U S A. 2010;107(15):6958–63.

52.  Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, et al. PlasmoDB: A
functional genomic database for malaria parasites. Nucleic Acids Res. 2009;

53.  Drakeley CJ, Secka I, Correa S, Greenwood BM, Targett GAT. Host haematological factors
influencing the transmission of Plasmodium falciparum gametocytes to Anopheles gambiae
s.s. mosquitoes. Trop Med Int Heal. 1999;

54.  Targett G, Drakeley C, Jawara M, Von Seidlein L, Coleman R, Deen J, et al. Artesunate reduces
but does not prevent posttreatment transmission of Plasmodium falciparum to Anopheles
gambiae. J Infect Dis. 2001;

55.  Drakeley CJ, Eling W, Teelen K, Bousema JT, Sauerwein R, Greenwood BM, et al. Parasite
infectivity and immunity to Plasmodium falciparum gametocytes in Gambian children.
Parasite Immunol. 2004;

56.  Sutherland CJ, Ord R, Dunyo S, Jawara M, Drakeley CJ, Alexander N, et al. Reduction of
malaria transmission to Anopheles mosquitoes with a six-dose regimen of co-artemether.

PLoS Med. 2005;2(4):0338–46.

57.   Dunyo S, Milligan P, Edwards T, Sutherland C, Targett G, Pinder M. Gametocytaemia after
drug treatment of asymptomatic Plasmodium falciparum. PLoS.Clin.Trials. 2006.

58.   Hallett RL, Dunyo S, Ord R, Jawara M, Pinder M, Randall A, et al. Chloroquine/Sulphadoxine-
Pyrimethamine for Gambian Children with Malaria: Transmission to Mosquitoes of
Multidrug-Resistant Plasmodium falciparum. PLoS Clin Trials. 2006;

59.   Alexa A and Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology. R package
version 2.24.0. 2016.

60.   Supek F, Bošnjak M, Škunca N, Šmuc T. Revigo summarizes and visualizes long lists of gene
ontology terms. PLoS One. 2011;6(7).

61.   van Dam TJ, Wheway G, Slaats GG, Huynen MA, Giles RH. The SYSCILIA gold standard
(SCGSv1) of known ciliary components and its applications within a systems biology
consortium. Cilia. Cilia; 2013;2(1):7.

62.   Moreno-Pérez DA, Dégano R, Ibarrola N, Muro A, Patarroyo MA. Determining the
Plasmodium vivax VCG-1 strain blood stage proteome. J Proteomics. Elsevier B.V.;
2015;113:268–80.

## Acknowledgments

We are thankful to Colin Sutherland (LSHTM London, UK) for commenting on the manuscript and providing serum samples that we used to study antibody responses in Gambian individuals. David Conway and Johannes Dessens (LSHTM London, UK) gave helpful feedback on the initial gold standard lists. We further thank Adam D. Shandling, Jozelyn V. Pablo and Andy A. Teng from Antigen Discovery Inc. (Irvine, CA) for their work on the protein microarray.

TB was supported by the Netherlands Organization for Scientific Research (nwo.nl), through a VIDI fellowship 016.158.306. The Radboud Institute for Health Sciences (rihs.nl), supported LM through grant R-2765. The Virgo consortium (virgo.nl), grant FES0908, supported the work of RvdL and TJPvD. TWAK is supported by the Netherlands Organization for Scientific Research (NWO-VIDI 864.13.009). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Author Contributions Statement

MAH, TB, RvdL and LMK conceptualised the work. LMK, RvdL, TJPvD, MAH and TB analysed the data. LMK, TB, DAB, PA, FS, CJJ, SMK, TWAK, MM and RS assembled the Gold Standard. KL, MvdVB, WG, RSS, LMK and WS conducted and analysed experiments, LMK wrote the first draft of the manuscript, all authors reviewed the manuscript.

**Competing financial interests**

JJC is employed by Antigen Discovery, Inc. The authors declare no further competing interests.

**Figure Legends**

*Figure 1* **Figure 1**. *Clustered data sets used in this study with genes ranked according to their protein or transcript expression*. *Level of expression as detected in the respective samples with unique peptide counts for MS data and percentiles for transcriptomics. See Table 1 for study keys. Distribution of asexual(a)/gametocyte(g) samples (red/blue) is shown in top bar, proteomics (P) and transcriptomics (T) (dark/light grey) in lower bar.*

*Figure 2* **Figure 2**. *Gametocyte-specificity scores for P. falciparum genes derived from proteomics (P) and transcriptomics (T) data sets*. *(A) Boxplot for integrated scores for the two gold standard sets and all other Pf genes, derived from proteomics, transcriptomics or all data sets (combined). (B) Density of P and T gametocyte scores, individual gold standard genes and their scores are indicated at the bottom (red, asexual, blue gametocyte). (C) 100 highest ranking proteins and transcripts, gametocyte gold standard in blue. (D) Correlation of the gametocyte-specificity scores derived from all integrated MS studies and* Pf *MS studies only.*

*Figure 3* **Figure 3**. *Validation of Bayesian gametocyte scoring with area under the curve (AUC) values*. *Integrated data and individual data sets are compared by 10-fold cross-validation (subsampling of gametocyte and asexual gold standard sets). Integrated proteomics (P) and transcriptomics (T) scores in bold lines. P. berghei data sets in shades of red, individual proteomics and transcriptomics studies with short and long dashes, respectively. See Table 1 for study keys.*

*Figure 4* **Figure 4. Comparison of reported gametocyte-specific proteins in mass spectrometry studies.** *(A) Proteins reported as gametocyte-specific by six individual studies, agreements on gametocyte-specificity are summarized in the table. Bayesian: gametocyte-specific proteins (n=602) that have a score > 5 after data integration. The overlap with previously published data sets is shown, but not to scale. Overlap between the individual studies is not shown for better visibility. Note that the Lasonder 2002 study includes proteins that were found in gametocytes or gametocytes and gametes. (B) Proteins that were reported as non-gametocytic and are (partially) included after data integration*

*Figure 5* **Figure 5**. *Seroprevalence in two cohorts of parasite carriers in The Gambia*. *(A+B) Antibodies against the highest scoring gametocyte-specific proteins were measured on protein microarrays. Comparison of positivity (mixture-model cutoff) in gametocyte carriers (n=164) and non-carriers (n=63). Gametocyte presence determined by microscopy. All individuals were positive for asexual parasites.  (A) Prevalence of antigens from the gold standard and predicted gametocyte-specific proteins (B) Antigens of five predicted gametocyte-specific proteins are preferentially recognized by gametocyte carriers. Error bars indicate the upper limit of the 95% confidence interval around the proportion.  p<0.05*

698    *Fisher's exact test, corrected for multiple testing of a total of 75 antigens (Benjamini-*
699    *Hochberg)*

700
<sub>Figure 6</sub> **Figure 6**. **Validation of gametocyte-specific targets in qRT-PCR** *Targets are sorted*
*for decreasing gametocyte-specificity in all panels, see Table 2. (A) Minimum transcript*
*abundance in blood stage versus gametocytes in different Pf strains. 1000-fold enrichment*
*of transcript in gametocytes over asexuals was assumed when delta-Ct was 10 or higher*
*(dashed line), considering the lowest Ct value detected in any asexual concentration-*
*matched sample. This threshold was not met by the transcripts with gene IDs in grey. (B)*
*Detection limit of eight validated targets alongside Pfs25 in serial dilutions of Pf NF54*
*asexual stage parasites (ring stage parasites 10-20 hours post invasion). (C) Detection limit*
*of the most sensitive targets in serial dilutions of stage V gametocytes. (A-C) For Pf NF54,*
*all n=3, other strains n=2 biological replicates, all measurements in triplicates. (D) Sensitivity*
*of eight validated targets in Kenyan blood samples of varying gametocyte densities.*

**713**  **Table 1. Data sets used for integration**

**714**  Life stages Spz sporozoites, Ri rings, Troph trohozoites, Schiz schizonts, Mer

**715**  merozoites, Gct gametocytes. MS mass spectrometry (highest unique peptide count

**716**  in any of the samples), T transcriptomics (highest percentile in any of the samples).

**717**  Data from Miao et al. 2017 was integrated after analyses of high scoring proteins,

**718**  ranks and scores are included in Supplementary Table 2.

**719**  *Asexual microarray data by Llinas and others retrieved from plasmoDB version 28

**720**  (Data set "Pfal3D7 real-time transcription and decay"), no accompanying publication.

| Study | Reference | Species | Life Stage | Integrated Data |
|-------|-----------|---------|------------|-----------------|
| Fl02 | Florens et al. 2002 [14] | *Pf* | Spz, Troph, Mer, Gct | MS asexual/Gct |
| La02 | Lasonder et al. 2002 [13] | *Pf* | Troph, Schiz, Gct | MS asexual/Gct |
| Le03 | LeRoch et al. 2003 [23] | *Pf* | Spz, Ri, Troph, Schiz, Mer, Gct | T asexual/Gct |
| Ha05 | Hall et al. 2005 [15] | *Pb* | Ri, Troph, Schiz, Gct | MS + T asexual/Gct |
| Kh05 | Khan et al. 2005 [16] | *Pb* | Mixed blood stage, Gct | MS asexual/Gct |
| Yo05 | Young et al. 2005 [24] | *Pf* | Gct | T Gct |
| Ll06 | Llinas et al. 2006 [25] | *Pf* | All blood stages, synchronized | T asexual |
| Ot10 | Otto et al. 2010 [26] | *Pf* | All blood stages, synchronized | T asexual |
| Bo11 | Treeck et al. 2011 [41] | *Pf* | Schiz | MS asexual |
| Lo11 | Lopez-Barragan et al. [27] | *Pf* | Troph, Schiz, Gct | T asexual/Gct |
| Oe12 | Oehring et al. 2012 [18] | *Pf* | Ri, Troph, Schiz | MS asexual |
| Mo14 | Moreno-Perez et al. 2014 [62] | *Pv* | Ri, Troph, Schiz | MS asexual |
| Ot14 | Otto et al. 2014 [28] | *Pb* | Ri, Troph, Schiz, Gct | T asexual/Gct |
| Si14 | Silvestrini et al. 2010, Tao et al. 2014 (re-analyzed) [17,19] | *Pf* | Troph, Schiz, Gct | MS asexual/Gct |
| Ta14 | Tao et al. 2014 [19] | *Pf* | Gct | MS Gct |
| Ll15 | Llinas et al. 2015* | *Pf* | All blood stages, synchronized | T asexual |
| La16 | Lasonder et al. 2016 [20] | *Pf* | Gct | MS + T Gct |
| Su16 | Suarez-Cortes et al. 2016 [21] | *Pf* | Gct | MS Gct |
| Mi17 | Miao et al. 2017 [33] | *Pf* | Gct | MS Gct |

**721**

**722**

**723**

| Rank | Gene ID | Description | Name | Intron-spanning | Least Ct difference |
|---|---|---|---|---|---|
| 3 | PF3D7_1143600 | conserved Plasmodium protein, unknown function | — | no | 9.37 |
| 5 | PF3D7_1147200 | tubulin--tyrosine ligase, putative | — | no | 8.52 |
| 6 | PF3D7_1026100 | conserved Plasmodium protein, unknown function | — | yes | 12.94 |
| 7 | PF3D7_1438800 | conserved Plasmodium protein, unknown function | — | yes | 8.57 |
| 8 | PF3D7_0625100 | sphingomyelin synthase 2, putative | SMS2 | no | 11.04 |
| 12 | PF3D7_0930000 | procollagen lysine 5-dioxygenase, putative | — | no | 11.96 |
| 14 | PF3D7_0518700 | mRNA-binding protein PUF1 | PUF1 | yes | 8.31 |
| 15 | PF3D7_0303900 | phosphatidylethanolamine-binding protein, putative | — | yes | 10.87 |
| 16 | PF3D7_1466600 | conserved Plasmodium protein, unknown function | — | no | 5.67 |
| 17 | PF3D7_1107900 | mechanosensitive ion channel protein, putative | MSCS | no | 6.33 |
| 18 | PF3D7_1214500 | conserved Plasmodium protein, unknown function | — | yes | 10.61 |
| 24 | PF3D7_1131500 | conserved Plasmodium protein, unknown function | — | no | 10.41 |
| 51 | PF3D7_0929600 | G2 protein, putative | — | yes | 8.37 |
| 61 | PF3D7_0816800 | meiotic recombination protein DMC1, putative | DMC1 | yes | 13.29 |
| 75 | PF3D7_0903800 | LCCL domain-containing protein | CCp4 | yes | 12.70 |

**Table 2: Properties of putative gametocyte-specific targets.**
Rank in transcriptomics (all data sets) for specificity in gametocytes. Random sample of top
100, excluding the gold standard. If primers are not intron-spanning, samples were DNase I
treated. Ct difference is the difference between the lowest Ct detected in asexual samples
and the highest Ct in concentration-matched stage V gametocytes, averaged across strains
*Pf* NF54, NF135, NF166 and NF175.

**A**

Gametocyte specifictiy over asexual (log2)

Proteomics    Transcriptomics

Gametocyte Gold standard n=41
Asexual Gold standard n=46
Other Pf genes

**B**

Density

**C**

Top 100 P

26 of 41 genes from gametocyte gold standard

25 of predicted gametocyte proteins are of *unknown function*

53 of predicted transcripts are of *unknown function*

15 of 41 genes from gametocyte gold standard

Top 100 T

**D**

Proteomics scores (Pf only)

Proteomics scores (all data sets)

r=0.9867477

| studies reporting as gametocyte-specific or enriched | number of proteins |
|---|---|
| 0 | 7 |
| 1 | 48 |
| 2 | 156 |
| 3 | 209 |
| 4 | 117 |
| 5 | 51 |
| 6 | 14 |
| total | 602 |