Genome analysis reveals evolutionary mechanisms of adaptation in systemic dimorphic fungi

José F. Muñoz[1], Juan G. McEwen[2,3], Oliver K. Clay[3,4] , Christina A. Cuomo[1]*

[1]Broad Institute of MIT and Harvard, Cambridge, MA, United States.

[2] Cellular and Molecular Biology Unit, Corporación para Investigaciones Biológicas, Medellín, Colombia.

[3] School of Medicine, Universidad de Antioquia, Medellín, Colombia.

[4] School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia

* cuomo@broadinstitute.org

ABSTRACT

Dimorphic fungal pathogens cause a significant human disease burden and unlike most fungal
pathogens affect immunocompetent hosts. Most dimorphic fungi are found in the family
Ajellomycetaceae, including the genera *Histoplasma, Blastomyces, Paracoccidioides*, and the recently
described *Emergomyces*. To examine the origin of virulence and host adaptation in these fungal
pathogens, we compared the gene content of classic systemic, opportunistic, and non-pathogenic
species, including new genomes for *Emmonsia* species and two closely non-pathogenic species,
Helicocarpus griseus and *Polytolypa hystricis.* We examined differences in gene content between
pathogens and environmental fungi, and found that gene families related to plant degradation,
synthesis of secondary metabolites, and amino acid and lipid metabolism are retained in *H. griseus* and
P. hystricis. Genes of central importance in dimorphic pathogenic fungi such as heat shock response
proteins, morphological switch related proteins are conserved in non-pathogenic species. However, we
found that changes in the copy number of proteases, phosphotransferases and transcription factors in
systemic dimorphic relative to non-dimorphic species may have aided the evolution of specialized gene
regulatory programs to rapidly adapt to higher temperatures and new nutritional environments. Notably,
we found that both of the basal branching, non-pathogenic species appear homothallic, with both
mating type locus idiomorphs fused at a single locus, whereas all related pathogenic species are
heterothallic. These differences revealed that independent changes in nutrient acquisition capacity but
similar host/substrate shift from plants to animals have occurred in the Onygenaceae and
Ajellomycetaceae, and highlighted how the dimorphic pathogens from Ajellomycetaceae have adapted
to the host immune response and decreased their capacity for life stages in environmental niches.

INTRODUCTION

Among millions of ubiquitous fungal species that pose no threat to humans, a small number of species cause devastating diseases in immunocompetent individuals. Notably, many human pathogenic species are ascomycetes in the order Onygenales, which comprises dermatophytes (e.g. *Trichophyton* spp.), the spherule-forming dimorphic pathogen *Coccidioides*, and yeast-forming dimorphic pathogens *Histoplasma*, *Paracoccidioides*, and *Blastomyces*. Dimorphism is a specialized morphogenetic adaptation allowing both growth in the environment and colonization of a host and is critical for the lifecycle of dimorphic fungal pathogens [1]. In yeast-forming pathogens, dimorphism has been broadly defined as the ability to generate both yeast (e.g. blastoconidia) in the host or at 37°C, and mycelia (e.g. conidia-producing or other hyphae) in the environment [1]. During fungal evolution, dimorphism has arisen independently multiple times in saprophytic fungi, as is manifested in the wide distribution of dimorphic species across the Ascomycota, Basidiomycota and Zygomycota phyla [2]. Notably, the dimorphic fungi in the order Onygenales are primary pathogens causing systemic mycosis in healthy humans, and collectively cause over 650,000 new infections a year in the United States alone [3]. Dimorphic fungi can persist as latent infections in tens of millions of people worldwide and may reactivate when the host becomes immune-deficient [1].

The largest cluster of thermally dimorphic human pathogenic fungi belongs to the Onygenales family Ajellomycetaceae, which includes *Histoplasma*, *Paracoccidioides*, and *Blastomyces*. In addition to these medically important genera, the Ajellomycetaceae family also includes more rarely observed opportunistic pathogenic species including *Emmonsia parva* and *Emmonsia crescens*, which undergo a thermal dimorphic transition to produce adiaconidia rather than yeast and cause adiospiromycosis sporadically in humans [4,5]. This group also includes *Lacazia loboi*, the yeast-like etiological agent of lobomycosis [6]. Of public health concern, recent reports have documented the worldwide emergence of yeast-forming Ajellomycetaceae that cause systemic mycoses predominantly in immunocompromised patients, often with high case-fatality rates [7,8]. Morphological and phylogenetic analyses were combined to demarcate species boundaries within the Ajellomycetaceae with clinical significance, which led to the description of the novel dimorphic species *Emergomyces africanus*, *E. pasteurianus*, and *Blastomyces percursus* [8,9]. Non-dimorphic species have also been documented as early diverging Ajellomycetaceae, including *Helicocarpus griseus* and *Polytolypa hystricis*, neither of which has been associated with disease in mammals. These species are found in environmental samples and do not appear to undergo a thermally regulated dimorphic transition [10–12].

We hypothesized that such phenotypic differences within the Ajellomycetaceae could be associated
with gene family expansions or contractions, as a consequence of gene duplication and gene loss
events, and that species comparison might reveal evolutionary mechanisms of adaptation in the
systemic dimorphic fungi. Previous comparative and population genomic studies within
Ajellomycetaceae had found evidence of gene family contractions and expansions associated with
virulence [5,13,14]. Other Onygenales genera outside the Ajellomycetaceae represented by published
genome sequences include *Coccidioides* and the non-pathogenic related species *Uncinocarpus reesii*
[15,16], as well as diverse dermatophytes including species that cause skin infections [17,18]. Recently, the
genomes of four non-pathogenic Onygenaceae species closely related to *Coccidioides* were described,
providing additional resolution into changes in gene content between pathogenic and non-pathogenic
Onygenales [19]. Together these studies helped establish how gene content is related to the life cycle of
different dimorphic pathogenic fungi and dermatophytes; for example, gene family contractions in
cellulases and other plant metabolism genes, and gene family expansions in proteases, keratinases
and other animal tissue metabolism genes, indicated that dimorphic fungi switched from a nutrition
system based on plants to one system based on animals, though mostly relative to relative to
outgroups to the Onygenales such as *Aspergillus* [13,16–18]. In the Onygenales, this hypothesis was tested
experimentally; comparing growth on a wide variety of compounds revealed that in *Uncinocarpus reesii*
hyphal growth was restricted on carbohydrates and considerably improved on proteins [13].

To examine key transition points in evolution of virulence and host adaptation in the dimorphic fungi, we
increased the phylogenetic density within the Ajellomycetaceae by sequencing the genomes of non-
pathogenic and emerging species and then performed comparative genomic analyses of the systemic,
opportunistic, and non-pathogenic species. Although representative genomes for most of the described
pathogenic Onygenales genera have been sequenced, the only strictly non-pathogenic species that
have been sequenced are closely related to *Coccidioides* in the family Onygenaceae. Here, we have
sequenced four additional Onygenales species from the family Ajellomycetaceae; this includes *H.*
griseus and *P. hystricis*, neither of which has been associated with disease in mammals, and two
additional adiaspore-forming strains of *E. parva* and *E. crescens* with distinct phenotypic properties.
Leveraging our previous genomic studies and these additional genome sequences, we have
characterized the extent of gene family expansions and contractions to provide a better understanding
of the evolutionary mechanisms of adaptation in systemic dimorphic fungi in the Ajellomycetaceae.

RESULTS

Genome comparisons of *Ajellomycetaceae*, the largest group of dimorphic human pathogenic
102

fungi
103

Building on previous genomic analyses of the members from the Ajellomycetaceae [5,9,13,14,20], we
104

sequenced using Illumina technology the genomes of four species within Ajellomycetaceae with
105

demarcated phenotypic differences. In addition to *Emmonsia parva* UAMH139 and *Emmonsia crescens*
106

UAMH3008, we sequenced the genomes of one strain of *Emmonsia parva* UAMH130 (type strain) and
107

one additional strain of *E. crescens* UAMH4076. *E. parva* UAMH130 is from lungs of a rodent in USA,
108

and *E. crescens* UAMH4076 is from a greenhouse source in Canada [21]. We also sequenced two
109

species that live in soil, where they are found in animal excrement but not associated with disease;
110

neither species appears to undergo a thermal dimorphic switch to a parasitic phase, as there is no
111

observation of a yeast-like form [12,21]. This includes one strain of *Helicocarpus griseus*, UAMH5409
112

(Type strain; synonym *Spiromastix grisea*), isolated from gazelle dung in Algeria, and one strain of
113

Polytolypa hystricis, UAMH7299, isolated from porcupine dung in Canada.
114

The *de novo* genome assemblies of these four species have similar sizes to those of other genera
116

within the Ajellomycetaceae including *Histoplasma*, *Emergomyces* and *Paracoccidioides* [5,9,13,14], and
117

have high representation of conserved eukaryotic genes (**Figure S1, S2**). The assembly sizes were
118

27.8 Mb in *E. parva* UAMH130, 33.8 Mb in *E. crescens* UAMH4076, 32.0 Mb in *P. hystricis*, and 34.7
119

Mb in *H. griseus* (**Table 1**). In addition, using both the CEGMA [22] and BUSCO [23] conserved eukaryotic
120

gene sets, we found that the predicted genes for each sequenced species are nearly complete; all of
121

these genomes include 96-98% of the core genes (**Figure S2**). The total numbers of predicted genes in
122

E. parva UAMH130 (n = 8,202), and *E. crescens* UAMH4076 (n = 8,909) were similar to those found in
123

annotated genome assemblies of other Ajellomycetaceae (n = 9,041 on average); however, relative to
124

other Ajellomycetaceae *P. hystricis*, and *H. griseus* have a modest increase in gene content (n = 9,935
125

and n = 10,225, respectively; **Table 1**; **Figure S1**).
126

Examining the guanine-cytosine (GC) content in the genomes *E. parva* UAMH130, *E. crescens*
128

UAMH4076, *H. griseus* UAMH5409 and *P. hystricis* UAMH7299 revealed that none of these genome
129

assemblies showed evidence of the pronounced bimodality of GC-content observed in *B. gilchristii* and
130

B. dermatitidis (**Figure S3**; **Table 1**; [5]). This highlights that the acquisition of repetitive elements
131

contributing to bimodal GC distribution was a unique phenomena during the evolution of the *B. gilchristii*
132

and *B. dermatitidis* within the Onygenales. In addition, UAMH139 showed a small peak of low GC that
133

is largely absent in UAHM130 (44.4 % and 47.5 % GC, respectively; **Table 1**), which is consistent with
134

the phylogenetic divergence between these two strains, and highlights an independent gain of repetitive sequences (10.4 % and 2.6 % repetitive bases in UAMH139 and UAMH130, respectively; **Table 1**). Similarly, *E. crescens* UAMH3008 and UAHM4076 have demarcated differences in GC and repetitive content; UAHM4076 has a higher peak of low GC sequences (42.9 %GC, 15.0 % repetitive bases in UAMH4076; 45.2 %GC, 5.1 % repetitive bases in UAMH3008; **Table 1**). Additionally, the more basal strain more basal *P. hystricis* UAMH7299 has a representation of low GC sequences (45.6 %GC, 12.8 % repeat), but in *H. griseus* UAMH5409 they are almost absent (48.0 %GC, 1.8 % repeats; **Figure S3**). These results highlight the dynamic and independent expansion of repetitive elements during Ajellomycetaceae evolution.

Phylogenomic analysis revealed multiple transitions within dimorphic pathogens to enable human infection

We estimated a strongly supported phylogeny of the newly sequenced *E. parva* UAMH130*, E. crescens* UAMH4075, *H. griseus* and *P. hystricis* strains relative to other dimorphic fungi using maximum likelihood and 2,505 single copy core genes (**Figure 1**). Branch length and topology supported the placement and relationships of *P. hystricis* and *H. griseus* within the Ajellomycetaceae as basally branching. The two *E. parva* isolates (UAMH130 and UAMH139) do not form a single well-defined clade, confirming that *E. parva* may not be a single species. In addition to longer branch lengths suggesting higher divergence of these isolates, the average genome-wide identity between UAMH130 and UAMH139 is much lower even comparing across groups that represent recently subdivided species, for example, within *B. gilchristii* and *B. dermatitidis* or *P. lutzii* and *P. brasiliensis*, with an average identity of 88.6% versus 95.8% and 91.1 respectively (**Figure S4**). *E. parva* (UAMH130-Type) is the most basal member of the clade including *E. parva* UAMH139 and *Blastomyces* species *B. percursus, B. gilchristii, and B. dermatitidis*. Previous molecular approaches [11,24] and recent multi-locus approaches [8,9] also support the polyphyletic nature of *E. parva* isolates. Although both isolates of *E. crescens* (UAMH3008 and UAMH4076) form a single clade, the genetic variation between them is also greater than that observed within other species from the same genus in the Ajellomycetaceae, with long branch lengths and slightly lower average genome identity (91.8%) between these isolates than seen in other intraspecies comparisons within the family (**Figure S4**). The *E. crescens* clade is closely related to the recently proposed genus *Emergomyces* including the *E. pasteurianus/africanus* species [9]. The *E. crescens – Emergomyces* clade is a sister group of the clade including *Histoplasma* and *Blastomyces*, and *Paracoccidioides* is in a basal position for the dimorphic pathogens of the Ajellomycetaceae (**Figure 1**).

The intercalated clades of primary pathogens (e.g. *H. capsulatum*, *P. brasiliensis* and *B. dermatitidis*),
opportunistic pathogens (e.g. *E. pasteurianus*, *E. africanus*) producing a parasitic yeast phase, and
non-pathogens producing adiaconidia-type cells rather than yeast cells (e.g. *E. parva* and *E. crescens*)
suggests that the dimorphic fungi in the Ajellomycetaceae have undergone numerous evolutionary
transitions allowing adaptation, infection and virulence to humans and other mammals, and that
interactions with other eukaryotes in the environment may help maintain the capacity for pathogenic
growth in a mammalian host (**Figure 1**; **Table 2**). The non-thermally-dimorphic species *H. griseus* and
P. hystricis are highly supported as early diverging lineages of Ajellomycetaceae as previously
suggested [12]; genomic analysis strongly supports *P. hystricis* as the earliest diverging lineage within the
sequenced Ajellomycetaceae, and large genetic variation between these early saprophytes. The basal
position of both *H. griseus* and *P. hystricis* in the family Ajellomycetaceae and the large divergence
between them suggests that the history of this family could involve evolution from saprophytic non-
thermotolerant and non-yeast-forming ancestral species to dimorphic human pathogenic species after
the divergence with *H. griseus*, including the adaptation to higher temperatures and the appearance of
a dimorphic switch and a yeast parasitic phase (**Figure 1**).

Substantial gene family contractions underlie the shift from growth on plant to animals

By comparing the gene conservation and functional annotation of primary and opportunistic
dimorphic pathogens with their saprophytic relatives *H. griseus* and *P. hystricis,* we identified significant
decreases in gene family size (PFAM domain and GO-term counts, corrected p-value < 0.05)
associated with a host/substrate shift from plants to animals in the pathogenic Ajellomycetaceae
(**Figure 2**). Plant cell wall degrading enzymes are reduced in number or totally absent in dimorphic
pathogens compared with *H. griseus* and *P. hystricis*, including several glycosyl hydrolases, and fungal
cellulose binding domain-containing proteins (**Figure 2; Table S2**). These gene family changes suggest
that dimorphic fungi from Ajellomycetaceae are not typical soil fungi in that they might maintain a close
association with living mammals or other organisms such as protozoa or insects. This transition
appears to have occurred within the family Ajellomycetaceae after divergence from *Helicocarpus* and
Polytolypa (**Figure 1**). Our analysis also recapitulates previous reports comparing smaller numbers of
genomes from the order Onygenales [5,13,16,19] that showed that genes coding for enzymes involved in the
deconstruction of plant cell walls were absent from all the human pathogens (e.g. *Paracoccidioides*,
Blastomyces, *Histoplasma*, *Coccidioides*) and non-pathogenic Onygenaceae (e.g. *Uncinocarpus reesii*,
Byssoonygena ceratinophila, *Amauroascus mutatus*, *Amauroascus niger*, *Chrysosporium*
queenslandicum) whereas these enzymes were commonly conserved in the non-pathogens outside the
order Onygenales (including *Aspergillus, Fusarium*, *Neurospora*; **Figure 2; Table S2**). Notably,

increasing the phylogenetic density within the order Onygenales in this study revealed that the loss of plant cell wall degradative enzymes occurred independently in two families within the Onygenales, and this highlights transitions between saprophytic and pathogenic species, where *Helicocarpus* and *Polytolypa* are have retained enzymes required to live on plant material in the soil, whereas non-pathogenic species from the Onygenaceae, which are not adapted to survive in mammals, have also lost capacity to degrade plant material.

Other notable changes in gene content included loss of protein families associated with secondary metabolite biosynthesis such as polyketide synthase dehydratases, and beta-ketoacyl synthases domains in dimorphic Ajellomycetaceae relative to *H. griseus* and *P. hystricis* (**Figure 2**; **Table S3**). We identified predicted biosynthetic clusters using the program antiSMASH (antibiotics and Secondary Metabolism Analysis Shell [25]). Overall, dimorphic pathogenic fungi from Ajellomycetaceae have fewer polyketide synthase (PKs) gene clusters than the saprophytic *H. griseus* and *P. hystricis* (**Table S3; Figure S5**). Dimorphic pathogens had lost essential genes or complete clusters of the type 1 and type 3 PKs, and terpene clusters (**Table S3**). Type 3 PKs are not common in fungi, and were not previously reported in the Ajellomycetaceae. Furthermore, *H. griseus* and *P. hystricis* but not dimorphic fungi have genes related to those involved in antibiotic biosynthetic pathways (ko01055; **Table S2**). Previous studies had shown that *P. hystricis* secretes a pentacyclic triterpenoid exhibiting antifungal and antibiotic activity, denominated Polytolypin [26], suggesting this terpene cluster is intact and may produce this molecule. Loss of such secondary metabolic pathways may weaken responses to other microbes in the environment. In addition to secondary metabolite biosynthesis, Ajellomycetaceae dimorphic fungi showed significant decreases in pathways related with the production of sphingolipids, chloroalkane and chloroalkene degradation, linoleic acid metabolism, and bisphenol degradation (KEGG-EC counts, corrected p-value < 0.05; **Table S2**). Loss of degradative pathways for bisphenol and chloralkanes in particular likely reflect encountering these molecules in the environment but not during pathogenic growth. Altogether, these shifts in gene content highlight how the Ajellomycetaceae dimorphic pathogenic fungi became less adapted to survive in the soil than the more basally diverging saprophytic species *H. griseus* and *P. hystricis*.

Transition among carbohydrate metabolism and protein catabolism

We found that the evolution of carbohydrate metabolism and protein catabolism also occurred independently within the Ajellomycetaceae. We annotated carbohydrate active enzymes (CAZy) and peptidases (MEROPS; **Methods**), and using enrichment analysis found substantial shifts in the relative proportion of these groups among dimorphic and saprophytic species. Notably, the pathogenic

Ajellomycetaceae species, but not the saprophytic species *H. griseus* and *P. hystricis*, showed a
dramatic reduction of carbohydrate active enzymes, including 44 categories that were totally absent
and 29 that were significantly depleted; these included glycoside hydrolases, glycosyltransferases,
carbohydrate esterases, and polysaccharide lyases (corrected $p < 0.05$; **Figure 3**). The fact that these
enzymes are reduced in dimorphic pathogenic Ajellomycetaceae but not in *H. griseus* and *P. hystricis*
confirms that reduced carbohydrate metabolism occurred independently Ajellomycetaceae from the
reduction in the Onygenaceae, originally noted in the spherule-dimorphic pathogen *Coccidioides* [16]. In
contrast to carbohydrate active enzymes, we found that the total number of peptidases is relatively
similar in dimorphic and saprophytic species within the Ajellomycetaceae. Significant expansions were
found in specific families in *Blastomyces*, *Histoplasma* and Onygenaceae and Arthrodermataceae
species relative to *P. hystricis* and *H. griseus*, including the A11, S12, and S08A peptidases (corrected
$p < 0.05$; **Figure 3**). The peptidase family A11 contains endopeptidases involved in the processing of
polyproteins encoded by retrotransposons, and the increase in copy number is correlated with the
genome expansion due to the proliferation of repetitive elements in *Blastomyces* [5]. The peptidase
family S12 contains serine-type D-Ala-D-Ala carboxypeptidases, and S8 contains the serine
endopeptidase subtilisin. Two families of fungal metalloendopeptidases (S8 and M35) were previously
reported to be expanded in *Coccidioides* relative to saprophytic species outside the order Onygenales
[16]. In addition, it has been reported that the Ajellomycetaceae includes fewer copies of multiple classes
of peptidases (M36, M43, S8) as well as an associated inhibitor (I9, inhibitor of S8 protease) relative to
other Onygenales [5]. Comparing across all the Onygenales, we found that all pathogenic Onygenales,
as well as saprophytes related to *Coccidioides* such as *U. reesii*, had a higher ratio of proteases to
carbohydrate active enzymes than the saprophytic Ajellomycetaceae *H. griseus* and *P. hystricis*
(**Figure 3**). This strengthens the evidence that the adaptation and specialization of dimorphic
Ajellomycetaceae occurred independently and that non-pathogenic Onygenaceae may represent an
intermediate state in showing gene content changes associated with pathogens, which allowed these
fungi to use proteins as a main source of energy in nutrient-limited environments.
263
264

Loss of transporters in Ajellomycetaceae dimorphic fungi
In addition to major shifts in enzymes that support nutrient acquisition, the dimorphic
Ajellomycetaceae have undergone losses of families of transporters. These include the large classes of
major facilitator superfamily transporters (PF07690) and sugar transporters (PF00083) (**Figure 2)**. The
number of transporters correlates with the numbers of genes predicted to have transmembrane helices
(1,848 and 1,745 proteins have predicted transmembrane domains in *H. griseus* and *P. hystricis,*

respectively, relative to an average of 1,490 in dimorphic Ajellomycetaceae pathogens, **Figure 4**)**.** To
characterize gene gain and losses in transporters, we annotated transporter families (**Methods,** [27]), and
found significant expansion of 18 classes of transporters encompassing 9 families in *H. griseus* and *P.*
hystricis relative to dimorphic Ajellomycetaceae pathogens (corrected *p-value* < 0.05; **Figure 4**).
Transporter families that were totally absent or significantly reduced in pathogenic species included the
sugar porter, anion:cation symporter, AAA-ATPase, ammonium transporter and carnitine O-Acyl
transferase. The substrates of these transporters include galacturonate, Rip1, hexoses, dipeptides,
allantoate, ureidosuccinate, allantoin, glycerol, sulfate, sulfite, thiosulfate, sulfonates, tartrate, thiamine,
cellobiose, cellodextrin, ammonia, methylamine, phenylacetate, beta-lactams, trichothecene, and
carnitine O-octanoyl (**Figure 4**). Most of these compounds are only present in the plant cell wall,
including the cellulose, hemicellulose, or pectin types, supporting that dimorphic pathogens have
evolved to have more limited compounds from plant cell wall as source of energy.
283
Pathogenesis-related genes are conserved in both saprophytic and pathogenic lifestyles
Genes known to be important in the response to stress imposed by the host, including
virulence-associated or yeast-phase specific genes of central importance in dimorphic fungi in the
Onygenales, are conserved in *H. griseus* and *P. hystricis*. These include heat shock response proteins
(*HSF*, *HSP90*, *HSP70*), dimorphic switch related proteins (*RYP1, RYP2, RYP3, AGS1, FKS1*),
oxidative stress response proteins (*CATB*, *CATP*, *SOD3*), as well as antigens (*PbGP43*, *PbP27*, *BAD1*;
Table S4;[20]). In dimorphic pathogenic fungi both morphological transitions and growth temperature are
linked, and loss of genes necessary for high-temperature growth and morphological transition in these
pathogens results in attenuated virulence, which highlights that they are essential for pathogenesis [28–
30]. Some of these genes such as the *RYP*1-3 transcriptional regulators and the dimorphism-regulating
histidine kinase *DRK1* have been connected with morphogenetic adaptation in response to
environmental stimuli, and with key determinants of the mycelia to yeast transition in dimorphic human
pathogenic fungi in the Ajellomycetaceae [28,29]. Although *H. griseus* and *P. hystricis* are not dimorphic
and cannot grow at 37°C, the heat shock response has been shown to be highly evolutionarily
conserved among eukaryotes. The fact that genes of central importance for high-temperature growth
and dimorphic switch are conserved in *H. griseus* and *P. hystricis* suggests that these saprophytes may
sense different gradations of temperature, or that the mechanism that triggers a rapid response to the
adaptation may not be rapid enough to enable these fungi to grow at 37°C.
302
Dimorphic Ajellomycetaceae may have evolved specialized mechanisms to regulate their transcriptional
responses; these include expansion of proteins related to the regulation of transcription, and protein

kinase activity (GO term enrichment analysis; *q*-value < 0.05; **Table S2**), which are all expanded in

dimorphic pathogens relative to *H. griseus* and *P. hystricis.* Thus, the gain of transcription factors,

transcriptional regulators and phosphotransferases suggests that rapid evolution of transcriptional

mechanisms may underlie the adaptation to both high temperature and the dimorphic transition. As one

of the largest categories enriched in dimorphic Ajellomycetaceae but not in saprophytic species was the

protein kinase activity (**Table S2**), we further classified protein kinases using Kinannote [31], including the

divergent fungal-specific protein kinase (*FunK1*; ref; **Methods**). We found an expansion of the *FunK1*

family, in dimorphic Ajellomycetaceae as previously reported [5,13]. As we increased the phylogenetic

density in the Ajellomycetaceae, including species and strains with high genetic and phenotypic

variation this allowed resolution of the evolutionary history of this family of kinases (**Figure 5**). *FunK1*

family is enriched in the systemic pathogens *B. dermatitidis/gilchristii*, *H. capsulatum*, *Paracoccidioides*,

but limited in saprophytic species, *H. griseus* and *P.* hystricis. Notably, this family is also limited in the

adiaspore-forming *E. parva* (UAMH130, UAMH139), *E. crescens* (UAMH4076) but not in *E. crescens*

(UAMH3008). In addition, we found that the recently described primary pathogen *B. percursus* includes

a large expansion of FunK1 kinases, with almost twice the number as compared with its closest relative

B. dermatitidis (**Figure 5**). Phylogenetic analysis of *FunK1* genes revealed that this family undergone

independent lineage-specific expansions in primary dimorphic pathogens of both Ajellomycetaceae and

Onygenaceae, but not in non-pathogenic or opportunistic dimorphic species (**Figure 5**).

To identify specific genes that evolved within the Ajellomycetaceae to enable the dimorphic transition

and infection of mammals, we identified and examined ortholog clusters that were unique to these

pathogens and absent in both *H. griseus* and *P. hystricis*. We identified 75 ortholog clusters that were

present in dimorphic Ajellomycetaceae and dimorphic Onygenaceae species but absent in *H. griseus*

and *P. hystricis* (i.e. 'sporophyte gene loss events'), and 212 ortholog clusters that were present in

dimorphic Ajellomycetaceae, but absent in the dimorphic Onygenaceae species, *H. griseus* and *P.*

hystricis (i.e. 'dimorphic Ajellomycetaceae gene gain events'; **Table S5**). The gene loss events include

a mold specific protein (CIMG_00805), a phenol 2-monooxygenase (BDFG_05966), as well as several

protein kinases and transporters. The gene gain events include several protein kinases, transporters,

transcription factors, and peptidases (**Table S5**). In addition, we annotated genes that have been

broadly associated with host-pathogen interactions (Pathogen Host interaction database PHI), and

genes that have been reported as virulence factors in fungal species (Database of Fungal Virulence

Factors DFVF) [32,33]. We found that many genes that have been associated with host-pathogen

interactions are present in the saprophytic *P. hystricis* and *H. griseus,* and in fact some appear at

higher copy number counts (**Table S5**). This emphasizes that host-pathogen interaction and virulence

factors are not only important for pathogenicity in mammals, but also for saprophytic species to survive
and adapt to stress in the environment, including for example the ability to evade the engulfment by
amoebae, or the interaction with insects [34,35].

Conservation of genes induced during *in vivo* infection

We also hypothesized that genes induced during mammalian infection in dimorphic
Ajellomycetaceae pathogens might have different gene conservation patterns among *Blastomyces*,
Histoplasma, *Paracoccidioides*, and the novel species and saprophytes. Transcriptional responses
during an *in vivo* mouse model of infection had been studied in *B. dermatitidis* (strain ATCC26189; [5])
and *P. brasiliensis* (strain Pb18; [36]). In both *B. dermatitidis* and *P. brasiliensis* the high-affinity zinc
transporter, *ZRT1* (BDFG_09159; PADG_06417) and the flavodoxin-like protein *PST2* (BDFG_08006;
PADG_07749) were highly induced during infection; these genes are conserved in all Ajellomycetaceae
included in this analysis (**Table S6**). Of the 72 *in vivo*-induced genes in *B. dermatitidis* ATCC26188,
22% are absent from the *P. hystricis* genome and 11% are absent from the *H. griseus* genome. The
genes absent in *P. hystricis* include the cysteine dioxygenase (BDFG_08059), an acetyltransferase
(BDFG_07903), a thioesterase (BDFG_02348), a superoxide dismutase (BDFG_07895), a
sodium/hydrogen exchanger (BDFG_05427), and a response regulator protein (BDFG_01066). In
addition, we identified three secreted proteins that were found induced during the interaction of *B.*
dermatitidis with macrophages (BDFG_08689, BDFG_06057, BDFG_03876) [5] that are found
conserved only in the dimorphic Ajellomycetaceae pathogens but not in *H. griseus, P. hystricis,* or
outgroup species. These are strong candidates for effector proteins involved in host infection. Similar
numbers of genes found induced during *in vivo* infection in *P. brasiliensis* were absent in *P. hystricis*
and *H. griseus*, (23% and 17.5%, respectively; **Table S6**). Genes absent in both species include Gpr1
family protein (PADG_08695), isovaleryl-CoA dehydrogenase (PADG_05046), glutathione-dependent
formaldehyde-activating (PADG_05832), predicted transmembrane protein (PADG_05761), and ergot
alkaloid biosynthetic protein A (PADG_07739). Many of the genes that were absent in *P. hystricis* and
H. griseus, were also absent in at least one of the dimorphic pathogens in the Ajellomycetaceae,
highlighting species-specific gene gain events that may confer virulence specialization.

Evolution of the mating type locus in the Onygenales family Ajellomycetaceae

To elucidate the evolution of the mating system in the Ajellomycetaceae, we identified and
characterized the mating locus of *P. hystricis*, and *H. griseus*. Notably, these two species have both
mating type idiomorphs *HMG* box (*MAT 1-2*) and alpha box (*MAT1-1*); these are therefore the only
homothallic species identified so far within the Ajellomycetaceae (**Figure 6**). Comparative analysis

showed that the locus is not expanded relative to other Ajellomycetaceae species (21 kb between the
flanking genes *SLA2* to *APN2/COX13*), unlike the expansion that observed in the larger genomes of *B.*
dermatitidis or *B. gilchristii* (~60 kb; [37]). One difference is that a sequence inversion between *APN2* and
COX13 is uniquely observed in *P. hystricis* relative to all other sequenced Ajellomycetaceae species,
including *H. griseus* and previously reported species [5,14,37,38]. This inversion observed in *P. hystricis*
(mating type idiomorph flanked by *SLA2* and *APN2/COX13*) is similar to the gene order in the
Onygenaceae species *Coccidioides* spp. but not in the Arthrodermataceae species *M. gypseum* or *T.*
rubrum (**Figure 6; Table S7**) suggesting that independent inversion events within the locus. In addition,
we identified the mating type locus in non-pathogenic Onygenaceae, and found that *Amauroascus*
mutatus and *Byssoonygena ceratinophila* have both mating type idiomorphs (**Table S7**). In *A. mutatus*,
HMG box (*MAT 1-2*) and alpha box (*MAT1-1*) are linked in the same scaffold, however they are
separated by 184 kb, including 65 protein-coding genes (**Figure 6**). The identification of homothallic
mating loci non-pathogenic species within both the Ajellomycetaceae and the Onygenaceae suggests
that there have been multiple transitions between homothallic and heterothallic sexual states in
dimorphic pathogens and non-pathogenic species in the Onygenales.
388

DISCUSSION

The evolution of the dimorphic Onygenales is characterized by multiple transitions between saprophytic and human pathogenic growth. By sequencing the genomes of two additional dimorphic adiaspore-forming species (*Emmonsia parva*, *E. crescens*) and two early diverging non-pathogenic, non-dimorphic species (*Helicocarpus griseus* and *Polytolypa hystricis)*, we analyzed how changes in gene content correlate with transitions to pathogenesis. Our results establish an outer bound on the timing of two key events in the evolution of the Ajellomycetaceae; both the loss of genes involved in digesting plant cell walls and the expansion of proteases, transcriptional regulators, and protein kinases, occurred after divergence with *H. griseus*. The shift in these families was previously observed in both *Paracoccidioides* within this group and in the related *Coccidioides–Uncinocarpus* clade (Desjardins et al. 2011; Sharpton et al. 2009); more recently the comparison with non-pathogenic Onygenaceae species supported previous reports of the contraction of gene families involved in digesting plant cell walls throughout the Onygenales, and of the expansion of gene families involved in digesting animal protein (Whiston et al., 2016). Whereas the studies on contraction of gene families involved in digesting plant cell walls addressed other fungi outside the order Onygenales, our analysis shows that *H. griseus* and *P. hystricis* did not experience such a contraction, which suggests that contractions in gene families involved in digesting plant cell walls throughout the Onygenales have occurred multiple times independently and more recently in Onygenaceae and Ajellomycetaceae families.

Since many virulence factors and dimorphic switch related proteins were also conserved in the non-dimorphic Ajellomycetaceae species, this suggests that these proteins could be needed not only for survival in the host, and that presence of virulence factors alone does not specify pathogenicity. In dimorphic Ajellomycetaceae other mechanisms such as gene regulation or selection may play a role in how these genes respond during distinct stimuli. In addition, it is possible that those virulence factors in mammals were also needed for the survival in the soil, for example to survive interaction with amoebae or in insects [39]. However, there are some virulence factors that are needed for virulence in mammals that are not necessary for virulence in other eukaryotes. For example, the alpha mating factor locus in *Cryptococcus neoformans* contributes to virulence in mice, but no difference was observed in the interaction of congenic *MAT*α and *MAT***a** strains with amoebae (Nielsen et al. 2005). While the human host is a very different habitat in comparison with soil or excrements of animals that are the most common natural niche for this group of fungi, interactions with other eukaryotes in the environment likely selected for some properties that also predisposed species to become pathogenic to humans. Selection pressures in the environment are responsible for the emergence and maintenance

of traits that confer upon some soil microbes the capacity for survival in animal hosts. This is important to understand the evolution and emergence of novel pathogens. For example, *Emergomyces africanus* has recently emerged and numerous cases of novel yeast-forming species have been reported [7–9]. However, *Emergomyces* species have not been found in other mammals like rodents or in the soil. In addition, *E. africanus* has a very restricted geographic distribution. Across the Onygenales, nearly all species have a high lower ratio of proteases to carbohydrate active enzymes; the exceptions are the two saprophytes *P. hystricis* and *H. griseus*, but not the saprophytes related to *Coccidioides* such as *Uncinocarpus reesii*. This genomic profile matches the results of growth assays for the non-pathogenic species *U. reesii*; this species can grow on a wide range of proteinaceous substrates but only a very limited range of carbohydrates, namely cellulose and its component glucose[13]. Together, these findings suggest that fungi in the Onygenales transitioned from saprophytes that digest plant materials to saprophytes that digest more limited plant materials and a wide range of animal proteins, and finally those that are major animal pathogens.

We found many of the genes with roles in the dimorphic transition and virulence are conserved in the saprophytic species *H. griseus* and *P. hystricis*, and we did not find larger patterns of gain of functional classes in pathogenic species. However, we found smaller changes in gene content related to the regulation of gene expression and signaling that may account for the ability to grow at high temperature, species specific morphological transitions, and therefore their differences in virulence, since rapid thermo-tolerance and morphological adaptation is essential for pathogenesis [28,29,40]. Overall, we found significant expansions of the number of the protein kinase *FunK1* family, transcription factors and other genes associated with the regulation of gene expression in yeast-forming Ajellomycetaceae relative to adiaspore-forming and saprophytic species. Our comparison with saprophytic Ajellomycetaceae highlights dramatic decreases and complete absences of several classes of gene families that are important for survival and growth in a soil environment.

Intriguingly, we found that both more basal non-pathogenic Ajellomycetaceae species *H. griseus* and *P. hystricis* appear homothallic, in striking contrast to all the previously described largely pathogenic species in this group that are all heterothallic. *P. hystricis* has a fused mating type locus, with both idiomorphs, *HMG* box (*MAT 1-2*) and alpha box (*MAT1-1*), fused at a single locus, and the same is likely true of *H. griseus*, as genes are located at the ends of two scaffolds in this assembly. In addition, we found that this is also true in the non-pathogenic Onygenaceae species *Amauroascus mutatus* and *Byssoonygena ceratinophila* have both mating type idiomorphs. *A. mutatus* also has both mating type idiomorphs located on the same scaffold; however, these are separated by a much larger distance than

is typically observed. This suggests that this locus has been subject to novel recombination events
either to bring the two idiomorphs nearby on the same scaffold or alternatively in rearranging a fused
locus. While it has been previously suggested that the ancestor of the Onygenales very likely was
heterothallic [38], the placement of multiple lineages of homothallic species in this group raises the
possibility of a heterothallic ancestor, with more recent transitions in pathogenic species to be
homothallic and rearrangements of the mating locus in others such as *P. hystricis* and *A. mutatus*. Loss
of the capacity for homothallic self-mating could increase the frequency of out-crossing and this may be
beneficial to some species under pressure from their environment or host [41] such as the pathogenic
species in this group. Examining the mating type loci of additional non-pathogenic species in the
Onygenales would further reveal the if there are novel configurations of homothallic or heterothallic loci,
which may help infer the mating ability of ancestral species and the timing of transitions in pathogens
and non-pathogens.
469

Our phylogenomic analysis revealed that dimorphic pathogens in the Onygenales have undergone
multiple evolutionary transitions that enabled infection of humans and other mammals. Some such as
the loss of plant cell wall degrading enzymes are convergent for pathogens in the two families of
Onygenales. Our comparisons also revealed major differences in non-pathogenic species between
these two families, which suggested that some species appear intermediate between saprophytes and
pathogens in terms of having a shift in metabolic capacity without associated pathogenic growth.
Targeting additional non-pathogenic species within the Ajellomycetaceae (e.g. *Emmonsiellopsis*) for
genome sequencing and comparison would help further refine the picture of the overall evolution of this
important group of fungi.
479

480 **MATERIALS AND METHODS**

482 **Genome sequencing**

483  Species were selected for sequencing based on previous estimates of high genetic variation

484 among the members of the genus *Emmonsia*, especially in the *E. parva* species, and based on

485 observations of the emergence of emmonsia-like species in the Ajellomycetaceae causing systemic

486 human mycoses worldwide [8,9]. We selected one strain of *Emmonsia parva* (UAMH 130; CBS 139881;

487 type strain) and one additional strain of *Emmonsia crescens* (UAMH 4076; CBS 139868). *E. parva*

488 UAMH130 is from the lungs of a rodent in the USA, and *E. crescens* UAMH4076 is from a greenhouse

489 source in Canada, and mated with *E. crescens* UAMH129, 349 [21]. In addition, the first strictly non-

490 pathogenic non-dimorphic species from the Ajellomycetaceae, one strain of *Helicocarpus griseus*

491 (UAMH 5409; a.k.a. *Spiromastix grisea*) and *Polytolypa hystricis* (UAMH 7299) were selected for whole

492 genome sequencing. Genomic DNA was isolated from mycelial phase at 22 ºC. We constructed one

493 library with 180-base inserts and sequenced on the Illumina HiSeq 2000 platform to generate 101 bp

494 paired-end reads with high coverage (173X UAMH 4076; 199X UAMH 130; 205X UAMH 5409; 163X

495 UAMH 7299).

497 **Genome assembly, gene prediction and annotation**

498  The 101-bp Illumina reads of *E. parva* UAMH130, *E. crescens* UAMH4076, *P. hystricis* and *H.*

499 *griseus* were assembled using ALLPATHS-LG [42] with default parameters. All four *de novo* assemblies

500 were evaluated using the GAEMR package (http://software.broadinstitute.org/software/gaemr/), which

501 revealed no aberrant regions of coverage, GC content or contigs with sequence similarity suggestive of

502 contamination. Scaffolds representing the mitochondrial genome were separated out from the nuclear

503 assembly.

505 Genes were predicted and annotated by combining calls from multiple methods to obtain the best

506 consensus model for a given locus. These included *ab initio* predictions (GlimmerHMM, Augustus,

507 Snap, GeneMark-ES), homologous inference (Genewise, TBlastN), and gene model consolidation

508 programs (EvidenceModeler) [43,44]. For the protein coding-gene name assignment we combined

509 HMMER PFAM/TIGRFAM, Swissprot and Kegg products. Kinannote was used to annotate protein

510 kinases [31]. To evaluate the completeness of predicted gene sets, the representation of core eukaryotic

511 genes was analyzed using CEGMA genes [22] and BUSCO [23].

513 **Identification of orthologs and phylogenomic analysis**

To compare gene content and conservation, we identified orthologous gene clusters in these

sequenced genomes and in other Onygenales genomes using OrthoMCL (version 1.4) with a Markov

in a on index of 1.5 and a maximum e-value of 1e-5 [45]. We included the genomes of the classical

dimorphic pathogenic species (*Blastomyces dermatitidis, B. gilchristii, Histoplasma capsulatum,*

Paracoccidioides brasiliensis, P. lutzii, Coccidioides immitis, and C. posadasii), novel dimorphic human

pathogenic species (*Blastomyces percursus*, *Emergomyces africanus*, *E. pasteurianus*), and two

dimorphic non-human pathogenic species *Emmonsia parva* UAMH139 and *Emmonsia crescens*, along

with other species, including three *Aspergillus* (**Table S1**). In addition, we annotated and included in the

compare gene content and conservation analysis the available genomes of saprophytic species from

the family Onygenaceae *Byssoonygena ceratinophila* (UAMH 5669), *Amauroascus mutatus* (UAMH

3576), *Amauroascus niger* (UAMH 3544), and *Chrysosporium queenslandicum* (CBS 280.77), and the

genome of the yeast-forming dimorphic pathogen *Emergomyces orientalis* (5Z489, **Table S1**).

To examine the phylogenetic relationship of the newly sequenced *E. parva* and *E. crescens*, *P. hystricis*

and *H. griseus* relative to other dimorphic fungi we used single copy orthologs determined and

clustered using OrthoMCL. A total of 31 genomes from the Onygenales order and three *Aspergillus*

genomes were chosen to estimate the species phylogeny (**Table S1**). These include the four newly

strains described here: *E. parva* UAMH 130, *E. crescens* UAMH 4076, *P. hystricis* UAMH 7299 and *H.*

griseus UAMH 5409, as well as the following: three *Blastomyces dermatitidis* (ATCC26199,

ATCC18188, ER-3), one *Blastomyces gilchristii* (SLH14081), one *Emmonsia parva* (UAMH 139), one

Emmonsia crescens (UAMH 3008), two *Histoplasma* (WU24, G186AR)*,* five *Paracoccidioides* (Pb01,

Pb03, Pb18, PbCnh, Pb300), two *Coccidioides* (RS, Silveira), *Uncinocarpus reesii* (UAMH 1704),

Microsporum gypseum (CBS 118893), *Trichophyton rubrum* (CBS 118892), *Aspergillus nidulans*

(FGSC A4), *A. flavus* (NRRL3357) and *A. fumigatus* (Af293). Protein sequences were aligned using

MUSCLE, and a phylogeny was estimated from the concatenated alignments using RAxML v7.7.8 [46]

with model PROTCATWAG with a total of 1,000 bootstrap replicates.

541   Gene family and protein domain analysis

To identify gene content changes that could play a role in the evolution of the dimorphism and

pathogenesis within Ajellomycetaceae, we searched for expansions or contractions in functionally

classified genes compared to the other fungi from the order Onygenales. Genes were functionally

annotated by assigning PFAM domains, GO terms, and KEGG classification. HMMER3 [47] was used to

identify PFAM domains using release 27. GO terms were assigned using Blast2GO [48], with a minimum

e-value of 1×10^{-10}. Protein kinases were identified using Kinannote [31] and divergent FunK1 kinases

were further identified using HMMER3. We further annotated carbohydrate active enzymes, peptidases,

and transporter families using the CAZY version 07-15-2016 [49], MEROPS version 9.12 [50], and TCDB

version 01-05-2017 [27] databases, respectively. Proteins were searched against their corresponding

databases using BLAST, with minimum e-values of 1×10^{-80} for CAZY, 1×10^{-20} for MEROPS, and

1×10^{-20} for TCDB.

To identify functional enrichments in the compared genomes, we used four gene classifications:

OrthoMCL similarity clusters, PFAM domains, KEGG pathways, and Gene Ontology (GO), including

different hierarchy levels, MEROPS and CAZY categories, and transporter families. Using a matrix of

gene class counts for each classification type, we identified enrichment comparing two subsets of

queried genomes using Fisher's exact test. Fisher's exact test was used to detect enrichment of PFAM,

KEGG, GO terms, CAZY, MEROPS, and transporter families between groups of interest, and p-values

were corrected for multiple comparisons [51]. Significant (corrected p-value < 0.05) gene class

expansions or depletions were examined for different comparisons.

Data Availability Statement

The assemblies and annotations were deposited in GenBank under the following BioProject accession

numbers: *Emmonsia parva* strain UAMH 130 (PRJNA252752), *Emmonsia crescens* strain UAMH 4076

(PRJNA252751), *Polytolypa hystricis* strain UAMH 7299 (PRJNA234736), and *Helicocarpus griseus*

strain UAMH 5409 (PRJNA234735).

ACKNOWLEDGEMENTS

We thank Lynne Sigler for her help in selecting strains for sequencing. We thank the Broad Genomics

Platform for generating Illumina sequence for this study, Sarah Young and Margaret Priest for their

assistance in assembly and annotation, and Christopher Desjardins for helpful comments on the

manuscript. This project has been funded in whole or in part with Federal funds from the National

Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and

Human Services, under award N°: U19AI110818. This work was partly supported by Colciencias via the

grants under Contract N°: 122256934875 and N°: 221365842971, and by the Universidad de Antioquia

via a "Sostenibilidad 2016" grant.

Author contributions

Conceived and designed the experiments: JFM JGM OKC CAC. Performed the assembly and

annotation: JFM CAC. Analyzed the data: JFM CAC. Contributed reagents/materials/analysis tools:

JGM OKC. Wrote the paper: JFM CAC.

 Tables and Figures

586 **Table 1.** Statistics of the annotated genome assemblies of *Emmonsia*, *Helicocarpus* and *Polytolypa*

Species	Strain	Assembly (Mb)	Scaffolds (count)	N50 (kb)	GC (%)	Repeat (%)	Genes (count)
Emmonsia parva	UAMH 130	27.8	383	173.2	47.5	2.6	8,202
Emmonsia crescens	UAMH 4076	33.8	1,288	96.6	42.9	15.0	8,909
Helicocarpus griseus	UAMH 5409	34.7	641	135.7	48.0	1.8	10,225
Polytolypa hystricis	UAMH 7299	32.0	546	124.6	45.6	12.8	9,935

588 **Table 2.** Thermo-dimorphic pathogenic fungi, expansion and diversity within Ajellomycetaceae

Species	Family	Dimorphism	Parasitic phase	Virulence in huma
Blastomyces gilchristii	Ajellomycetaceae	Yes	Intermediate yeast	Primary pathogen
Blastomyces dermatitidis	Ajellomycetaceae	Yes	Intermediate yeast	Primary pathogen
Blastomyces percursus	Ajellomycetaceae	Yes	Large yeast	Primary pathogen
Emmonsia parva	Ajellomycetaceae	Yes	Small adiaconidia	Opportunistic
Histoplasma capsulatum	Ajellomycetaceae	Yes	Small yeast	Primary pathogen
Emmonsia crescens	Ajellomycetaceae	Yes	Large adiaconidia	Opportunistic
Emergomyces pasteurianus	Ajellomycetaceae	Yes	Small yeast	Opportunistic
Emergomyces africanus	Ajellomycetaceae	Yes	Small yeast	Opportunistic
Paracoccidioides brasiliensis	Ajellomycetaceae	Yes	Multibudding yeast	Primary pathogen
Paracoccidioides lutzii	Ajellomycetaceae	Yes	Multibudding yeast	Primary pathogen
Lacazia loboi	Ajellomycetaceae	ND	Multibudding yeast	Primary pathogen
Helicocarpus griseus	Ajellomycetaceae	No	No	Non-pathogen
Polytolypa hystricis	Ajellomycetaceae	No	No	Non-pathogen
Coccidioides immitis	Onygenaceae	Yes	Spherules	Primary pathogen
Coccidioides posadasii	Onygenaceae	Yes	Spherules	Primary pathogen
Sporothrix schenckii	Ophiostomataceae	Yes	Small yeast	Primary pathogen
Talaromyces marneffei	Trichocomaceae	Yes	Small septate yeast	Primary pathogen

Figure 1. Phylogenomic relationships and diversity in the family Ajellomycetaceae. Maximum likelihood phylogeny (2,505 core genes based on 1,000 replicates) including 20 annotated genome assemblies of species within the family Ajellomycetaceae, five other Onygenales and three *Aspergilli*. All nodes were supported by 100% of bootstrap replicates. Color bars indicate the genus and icons represent most common source or host. Illustrations of the most common morphology at 37 °C or in the host are depicted for selected species highlighting the phenotypic variation within the Ajellomycetaceae and the transition to human hosts represented by pathogenicity traits. * Formerly *Emmonsia parva*.

Figure 2. Gene family changes across the Ajellomycetaceae. Heatmap depicting results of a protein family enrichment analysis (PFAM domains; corrected *p-value* < 0.05) comparing the gene content of the non-dimorphic/non-pathogenic *Polytolypa hystricis* and *Helicocarpus griseus* with all other depicted members of the family Ajellomycetaceae including the dimorphic pathogens *Histoplasma*, *Emmonsia*, *Blastomyces*, *Emergomyces* and *Paracoccidioides*. Values are colored along a blue (low counts) to red (high counts) color scale, with color scaling relative to the low and high values of each row. Each protein family domain has a color code (right) indicating a global functional category. *Es*: *Emergomyces*; *Ea*: *Emmonsia*.

609
Figure 3. Shifts in carbohydrate metabolism and peptidase activity in Ajellomycetaceae.
Heatmaps depicting for each taxon carbohydrate metabolism categories (CAZy) and peptidase families
(MEROPs) across the Ajellomycetaceae species included in this study and other compared genomes.
These categories were found significantly enriched (corrected *p-value* < 0.05) in the non-dimorphic
systemic fungi *P. hystricis* and *H. griseus* relative to Ajellomycetaceae dimorphic fungi, or within
Ajellomycetaceae species and other compared genomes from Onygenaceae and Arthrodermataceae
families, or Eurotiales order. The ratio of MEROPS genes to CAZY genes for each genome across the
Ajellomycetaceae species included in this study and other compared genomes is shown at right. *Es*:
Emergomyces; *Ea*: *Emmonsia*.
619

Figure 4. Shifts in metabolism matched substrate preference within Ajellomycetaceae contraction of transporters within the Ajellomycetaceae. (right) Heatmap depicting the number of family transporters (TCDB; color-code blue: low, red: high) for each taxon across the Ajellomycetaceae species and other compared genomes. These families were found significantly enriched (test, corrected *p-value* < 0.05) in the non-dimorphic *P. hystricis* and *H. griseus* relative to dimorphic Ajellomycetaceae fungi. (left) Total number of transporters identified in the genomes for each taxon included in this study. *Es*: *Emergomyces*; *Ea*: *Emmonsia*.

Figure 5. Expansion and contraction of protein kinases class _FunK1_ across the Ajellomycetaceae. (top) _FunK1 A_ and _FunK1 B_ counts for each taxon across the Ajellomycetaceae species and other compared genomes. The tree shown corresponds to the maximum likelihood phylogenetic tree of the family Ajellomycetaceae from Fig. 1. (bottom) Maximum likelihood tree of _FunK1_ family showing expansion in Ajellomycetaceae yeast-forming (blue) pathogenic species. Each species (top) or gene (bottom) has a color code indicating the parasitic phase at 37 °C (yeast, adiaspore, spherule, or non-pathogenic).

638
Figure 6. Mating type evolution within the Onygenales family Ajellomycetaceae. (top) Synteny
schema depicting orientation and conservation of the genes adjacent to the mating type locus
idiomorphs HMG box (*MAT 1-2*, red) and alpha box (*MAT1-1*, dark red) in the Ajellomycetaceae. The
more basal species, *P. hystricis*, has both mating type idiomorphs intact, being the only homothallic
species identified so far within the Ajellomycetaceae. The locus has similar size to other
Ajellomycetaceae species with 21 kb average from *SLA2* (blue) to *APN2*/*COX13* (green/orange), but
not *Blastomyces dermatitidis/gilchristii* (60 kb; Li W et al., 2013). A synteny inversion encompassing
APN2 and *COX13* is uniquely observed in this ancestor relative to the rest of Ajellomycetaceae species
included in this study, including *H. griseus*. Grey genes are mating locus hypothetical proteins (MLHP).
(bottom) Schema depicting orientation and conservation of the genes adjacent to the mating type locus
idiomorphs HMG box (*MAT 1-2*, red) and alpha box (*MAT1-1*, dark red) in non-Ajellomycetaceae
Amaurascus mutatus, *C. immitis*, *U. reesii*, and *M. gypseum*. In *A. mutatus*, both mating type
idiomorphs are linked in the same scaffold, however they are separated by 184 kb, including 65 protein-
coding genes (gray).
653

References

1.   Klein, B. S. & Tebbets, B. Dimorphism and virulence in fungi. *Curr Opin Microbiol* **10,** 314–9
(2007).
2.   Sil, A. & Andrianopoulos, A. Thermally Dimorphic Human Fungal Pathogens--Polyphyletic
Pathogens with a Convergent Pathogenicity Trait. *Cold Spring Harb Perspect Med* **5,** a019794
(2015).
3.   Gauthier, G. M. Dimorphism in fungal pathogens of mammals, plants, and insects. *PLoS Pathog*
11, e1004608 (2015).
4.   England, D. M. & Hochholzer, L. Adiaspiromycosis: an unusual fungal infection of the lung.
Report of 11 cases. *Am J Surg Pathol* **17,** 876–86 (1993).
5.   Munoz, J. F. *et al.* The Dynamic Genome and Transcriptome of the Human Fungal Pathogen
Blastomyces and Close Relative *Emmonsia. PLoS Genet* **11,** e1005493 (2015).
6.   Herr, R. A. *et al.* Phylogenetic analysis of *Lacazia loboi* places this previously uncharacterized
pathogen within the dimorphic Onygenales. *J Clin Microbiol* **39,** 309–14 (2001).
7.   Kenyon, C. *et al.* A dimorphic fungus causing disseminated infection in South Africa. *N Engl J*
Med **369,** 1416–1424 (2013).
8.   Schwartz, I. S. *et al.* 50 Years of *Emmonsia* Disease in Humans: The Dramatic Emergence of a
Cluster of Novel Fungal Pathogens. *PLoS Pathog* **11,** e1005198 (2015).
9.   Dukik, K. *et al.* Novel taxa of thermally dimorphic systemic pathogens in the Ajellomycetaceae
(Onygenales). *Mycoses* **60,** 296–309 (2017).
10.   Marin-Felix, Y. *et al. Emmonsiellopsis*, a new genus related to the thermally dimorphic fungi of
the family Ajellomycetaceae. *Mycoses* **58,** 451–60 (2015).
11.   Untereiner, W. A. *et al.* The Ajellomycetaceae, a new family of vertebrate-associated
Onygenales. *Mycologia* **96,** 812–821 (2004).
12.   Untereiner, W. A., Scott, J. A., Naveau, F. A., Currah, R. S. & Bachewich, J. Phylogeny of
Ajellomyces, *Polytolypa* and *Spiromastix* (Onygenaceae) inferred from rDNA sequence and
non-molecular data. *Stud Mycol* **47,** 25–35 (2002).
13.   Desjardins, C. A. *et al.* Comparative genomic analysis of human fungal pathogens causing
paracoccidioidomycosis. *PLoS Genet* **7,** e1002345 (2011).
14.   Munoz, J. F. *et al.* Genome Diversity, Recombination, and Virulence across the Major Lineages
of Paracoccidioides. *mSphere* **1,** (2016).
15.   Neafsey, D. E. *et al.* Population genomic sequencing of Coccidioides fungi reveals recent
hybridization and transposon control. *Genome Res* **20,** 938–46 (2010).
16.   Sharpton, T. J. *et al.* Comparative genomic analyses of the human fungal pathogens
Coccidioides and their relatives. *Genome Res* **19,** 1722–31 (2009).
17.   Burmester, A. *et al.* Comparative and functional genomics provide insights into the pathogenicity
of dermatophytic fungi. *Genome Biol* **12,** R7 (2011).
18.   Martinez, D. A. *et al.* Comparative genome analysis of *Trichophyton rubrum* and related
dermatophytes reveals candidate genes involved in infection. *MBio* **3,** e00259-12 (2012).
19.   Whiston, E. & Taylor, J. W. Comparative Phylogenomics of Pathogenic and Nonpathogenic
Species. *G3 Bethesda* **6,** 235–44 (2015).
20.   Munoz, J. F. *et al.* Genome update of the dimorphic human pathogenic fungi causing
paracoccidioidomycosis. *PLoS Negl Trop Dis* **8,** e3348 (2014).
21.   Sigler, L. Ajellomyces crescens sp. nov., taxonomy of *Emmonsia* spp., and relatedness with
Blastomyces dermatitidis (teleomorph *Ajellomyces dermatitidis*). *J Med Vet Mycol* **34,** 303–14
(1996).
22.   Parra, G., Bradnam, K. & Korf, I. CEGMA: a pipeline to accurately annotate core genes in
eukaryotic genomes. *Bioinformatics* **23,** 1061–7 (2007).

23.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31,** 3210–3212 (2015).
24.   Peterson, S. W. & Sigler, L. Molecular genetic variation in *Emmonsia crescens* and *Emmonsia*
parva, etiologic agents of adiaspiromycosis, and their phylogenetic relationship to *Blastomyces*
dermatitidis (*Ajellomyces dermatitidis*) and other systemic fungal pathogens. *J Clin Microbiol* **36,**
2918–25 (1998).
25.   Blin, K. *et al.* antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary
identification. *Nucleic Acids Res.* **45,** W36–W41 (2017).
26.   Gamble, W. R., Gloer, J. B., Scott, J. A. & Malloch, D. Polytolypin, a new antifungal triterpenoid
from the coprophilous fungus *Polytolypa hystricis*. *J. Nat. Prod.* **58,** 1983–1986 (1995).
27.   Saier, M. H., Jr., Tran, C. V. & Barabote, R. D. TCDB: the Transporter Classification Database
for membrane transport protein analyses and information. *Nucleic Acids Res* **34,** D181-6 (2006).
28.   Beyhan, S., Gutierrez, M., Voorhies, M. & Sil, A. A temperature-responsive network links cell
shape and virulence traits in a primary fungal pathogen. *PLoS Biol* **11,** e1001614 (2013).
29.   Nemecek, J. C., Wuthrich, M. & Klein, B. S. Global control of dimorphism and virulence in fungi.
Science **312,** 583–8 (2006).
30.   Rappleye, C. A., Eissenberg, L. G. & Goldman, W. E. *Histoplasma capsulatum* alpha-(1,3)-
glucan blocks innate immune recognition by the beta-glucan receptor. *Proc Natl Acad Sci USA*
104, 1366–1370 (2007).
31.   Goldberg, J. M. *et al.* Kinannote, a computer program to identify and classify members of the
eukaryotic protein kinase superfamily. *Bioinformatics* **29,** 2387–94 (2013).
32.   Lu, T., Yao, B. & Zhang, C. DFVF: database of fungal virulence factors. *Database J. Biol.*
Databases Curation **2012,** (2012).
33.   Winnenburg, R. *et al.* PHI-base: a new database for pathogen host interactions. *Nucleic Acids*
Res. **34,** D459–D464 (2006).
34.   Casadevall, A., Steenbergen, J. N. & Nosanchuk, J. D. `Ready made' virulence and `dual use'
virulence factors in pathogenic environmental fungi-the Cryptococcus neoformans paradigm.
Curr Opin Microbiol **6,** 332–337 (2003).
35.   Derengowski Lda, S. *et al.* The transcriptional response of *Cryptococcus neoformans* to
ingestion by *Acanthamoeba castellanii* and macrophages provides insights into the evolutionary
adaptation to the mammalian host. *Eukaryot Cell* **12,** 761–74 (2013).
36.   Pigosso, L. L. *et al. Paracoccidioides brasiliensis* presents metabolic reprogramming and
secretes a serine proteinase during murine infection. *Virulence* **0,** 1–18 (2017).
37.   Li, W. *et al.* Identification of the mating-type (MAT) locus that controls sexual reproduction of
Blastomyces dermatitidis. Eukaryot Cell **12,** 109–17 (2013).
38.   Li, W., Metin, B., White, T. C. & Heitman, J. Organization and evolutionary trajectory of the
mating type (MAT) locus in dermatophyte and dimorphic fungal pathogens. *Eukaryot Cell* **9,** 46–
58 (2010).
39.   Steenbergen, J. N., Shuman, H. A. & Casadevall, A. *Cryptococcus neoformans* interactions with
amoebae suggest an explanation for its virulence and intracellular pathogenic strategy in
macrophages. *Proc. Natl. Acad. Sci.* **98,** 15245–15250 (2001).
40.   Rappleye, C. A. & Goldman, W. E. Defining virulence genes in the dimorphic fungi. *Annu Rev*
Microbiol **60,** 281–303 (2006).
41.   Heitman, J., Sun, S. & James, T. Y. Evolution of fungal sexual reproduction. *Mycologia* **105,** 1–
27 (2013).
42.   Gnerre, S. *et al.* High-quality draft assemblies of mammalian genomes from massively parallel
sequence data. *Proc Natl Acad Sci U A* **108,** 1513–8 (2011).
43.   Haas, B. J., Zeng, Q., Pearson, M. D., Cuomo, C. A. & Wortman, J. R. Approaches to Fungal
Genome Annotation. *Mycology* **2,** 118–141 (2011).

44. Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using EVidenceModeler and
the Program to Assemble Spliced Alignments. *Genome Biol* **9,** R7 (2008).
45. Li, L., Stoeckert, C. J., Jr. & Roos, D. S. OrthoMCL: identification of ortholog groups for
eukaryotic genomes. *Genome Res* **13,** 2178–89 (2003).
46. Stamatakis, A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
thousands of taxa and mixed models. *Bioinformatics* **22,** 2688–90 (2006).
47. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7,** e1002195 (2011).
48. Conesa, A. *et al.* Blast2GO: a universal tool for annotation, visualization and analysis in
functional genomics research. *Bioinformatics* **21,** 3674–6 (2005).
49. Cantarel, B. L. *et al.* The Carbohydrate-Active EnZymes database (CAZy): an expert resource
for Glycogenomics. *Nucleic Acids Res* **37,** D233-8 (2009).
50. Rawlings, N. D., Barrett, A. J. & Bateman, A. MEROPS: the peptidase database. *Nucleic Acids
Res* **38,** D227-33 (2010).
51. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc Natl Acad Sci
U A* **100,** 9440–5 (2003).
768
769