A strategy to exploit surrogate sire technology in livestock breeding programs

Paolo Gottardo[1], Gregor Gorjanc[1], Mara Battagin[1], R Chris Gaynor[1], Janez Jenko[1], Roger Ros Freixedes[1], C Bruce A Whitelaw, Alan J Mileham[3], William O Herring[2], John M Hickey[1][§]

[1]The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush, Midlothian, Scotland, UK

[2]The Pig Improvement Company, Genus PLC, 100 Bluegrass Commons Blvd., Ste 2200, Hendersonville, TN 37075, USA

[3]Genus PLC, 1525 River Rd., DeForest, Wisconsin, USA 53532

[§]Corresponding author

Email addresses:

PG: paolo.gottardo@roslin.ed.ac.uk

GG: gregor.gorjanc@roslin.ed.ac.uk

MB: mara.battagin@roslin.ed.ac.uk

RCG: chris.gaynor@roslin.ed.ac.uk

JJ: janez.jenko@roslin.ed.ac.uk

RRF: roger.ros@roslin.ed.ac.uk

CBAW: bruce.whitelaw@roslin.ed.ac.uk

AJM: Alan.Mileham@genusplc.com

WOH: William.Herring@genusplc.com

JMH: john.hickey@roslin.ed.ac.uk

Abstract

Background

In this work, we performed simulations to develop and test a strategy for exploiting surrogate sire technology in livestock breeding programs.

Methods

One hundred replicates of various scenarios were performed. Scenarios followed a common overall structure but differed in the strategy used to identify elite donors and how these donors were used in the product development part.

Results

The results of this study showed that using surrogate sire technology would significantly increase the genetic merit of commercial sires, by as much as 6.5 to 9.2 years' worth of genetic gain compared to a conventional breeding program. The simulations suggested that an identification strategy involving three-stages, the first of which uses genomic selection while the subsequent stages use progeny testing was the most effective strategy of all of those tested.

Conclusions

The use of one or a handful of elite donors to generate the production animals would be very different to current practice. While the results demonstrate the great potential of surrogate sire technology there are considerable risks but also other opportunities. Practical implementation of surrogate sire technology would need to account for these.

Background

In this study, we performed simulations to develop a strategy for exploiting surrogate sire technology [1] in livestock breeding programs. Surrogate sire technology allows the creation of males that lack their own germline cells, but have transplanted spermatogonial stem cells from other "elite donor" males. With this technology, a single male could give rise to huge numbers of progeny, potentially as much as all the production level animals in a particular time period. Using a single male to generate many or even all production animals offers many advantages. Firstly, it would reduce the genetic lag between the elite nucleus animals and the production animals. Secondly, it could enable better matching of specific management plans to the genetics. Thirdly, as we outline in the discussion it could enable elite donors to be selected for combining ability. The latter could increase production on farm and increase investment and innovation in breeding by enabling a greater ability to protect intellectual property.

Typically, livestock breeding programs are implicitly or explicitly organized in pyramid structures (Figure 1). At the top layer of the pyramid is the nucleus population. Nucleus populations are improved using recurrent selection and nowadays most selection decisions are made using genomic based testing rather than traditional phenotype based testing strategies [2–6]. In the middle of the pyramid is the multiplier layer. Multiplier herds multiply the genetics from the nucleus and sometimes cross purebred lines to produce F_1 (two-way) crosses. The base of the pyramid is the commercial layer. It is made up of commercial herds, and contains the majority of animals. The commercial producers often make a final cross between the terminal line sires and the maternal line dams.

The need to generate huge numbers of production animals in the commercial

layer and the limited number of progeny that any male can produce means that large

numbers of nucleus animals must contribute genetics to the subsequent layers and that

one to several generations are required to undertake the multiplication process. These

factors gives rise to a genetic lag, a difference in genetic mean between the nucleus

and commercial layers, equivalent to several years genetic gain [7] (e.g., 4 years in a

conventional pig breeding program). Surrogate sire technology would allow a single

elite nucleus male to produce very large numbers of progeny, by donating

spermatogonial stem cells to its surrogates [1]. This could shorten the lag between the

nucleus, multiplication and commercial layers. It could also increase the genetic merit

in multiplication and commercial layers, because these animals could descend from

the best male or best few males in the nucleus as opposed to many of the nucleus

males.

Using surrogate sire technology in this way would require that animal

breeding programs be restructured so that the best males be identified and tested to

become donors of spermatogonial stem cells. It would require that this process of

identification and testing takes place in a sufficiently small amount of time so that the

extra genetic gain at the commercial level would not be significantly reduced by the

extra time required by the breeders to identify, test donors and generate surrogate

sires.

Plant breeding programs that produce clonally propagated individuals or

inbred or hybrid lines (Figure 1) are conceptually similar to what a restructured

animal breeding program that identifies elite donors might look like. These types of

plant breeding programs seek: (i) to identify the best individual, or a handful of

individuals, from a population of individuals; and (ii) to disseminate this individual

very widely at the commercial layer [8]. To identify the best individual, plant breeders

typically use multiple stage testing and selection. As the breeding program progresses

through these stages the number of individuals being tested is reduced and the

precision of these tests increases. The small numbers of individuals in the final stages

are intensively tested in large replicated experiments that are repeated across several

environments and years. This ensures that the final individual released by the breeding

program is very well characterized and carries a minimal risk of having major

undetected weakness. This is necessary because this individual will have a huge

footprint in the commercial layer. Similar levels of evaluation would be needed in

livestock if elite donors were to have equally large footprints on the commercial layer.

The objective of this study was to develop a strategy for exploiting surrogate

sire technology in livestock breeding programs. This strategy involved a subtle, but

important, reorganisation in the animal breeding programs so that they combine

components of traditional animal and plant breeding programs. The reorganization is

similar to the two-part breeding program design that has recently proposed for the

incorporation of genomic selection into plant breeding programs [9]. The

reorganization involves an explicit partitioning of animal breeding programs into a

population improvement component and a product development component. The

population improvement component would be similar to the currently used recurrent

selection component in many animal breeding nucleus populations that use genomic

selection. The product development component would be similar to traditional plant

breeding programs and would involve a number of stages of testing to identify the

elite donors. The product development component could make use of testing for

combining ability, if that was appropriate for the particular species of interest.

Several alternative versions of the reorganized breeding program were

compared to different variants of a conventional breeding using simulation. The

alternative versions varied: (i) the number of stages of testing; (ii) the accuracy of

genomic prediction at the first stage of testing; (iii) the numbers of individuals tested

at subsequent stages; and (iv) the number of progeny used to test the individuals at

each stage. The results showed that using surrogate sire technology would

significantly increase the genetic merit of commercial sires, by as much as between

6.5 and 9.20 years' worth of genetic gain compared to different variants of a

conventional breeding program. The simulations suggested that an identification

strategy involving three-stages, the first of which uses genomic selection while the

subsequent stages use progeny testing was the most effective strategy of all of those

tested. The use of one or a handful of elite donors to generate the production animals

would be very different to current practice. While the results demonstrate the great

potential of surrogate sire technology there are considerable risks and these are

discussed.

Methods

Simulation was used to evaluate the impact of a strategy for exploiting surrogate sire technology in livestock breeding programs. One hundred replicates of various scenarios were performed. Scenarios followed a common overall structure (Fig. 2) but differed in the strategy used to identify elite donors and how these donors were used in the product development part.

Conceptually, the simulation scheme was divided into historical and future phases. The historical phase represented historical evolution and recent historical animal breeding efforts up to the present day, under the assumption that livestock populations have evolved for tens of thousands of years, followed by 22 recent generations of modern animal breeding with selection on genomic breeding values in a nucleus population. The future phase represented 20 future generations of modern animal breeding, with selection on genomic breeding values in a nucleus population that subsequently supplied genetic improvement to multiplication and commercial layers. The historical animal breeding generations were denoted -21 to 0 and the future animal breeding generations were denoted 1 to 20. The multiplier and commercial layers were not explicitly simulated but were instead represented with the average genetic merit of nucleus males that would give rise to multiplication and commercial animals while accounting for the time lag. Specifically, we only focused on a breeding program that produced terminal males.

Simulations involved the five following steps:

(i)   Generating whole-genome sequence data,

(ii)   Generating quantitative trait variants (**QTV**) affecting phenotypes,

(iii) Generating pedigree structures for a typical livestock population,

(iv) Performing selection and dissemination to the commercial layer, and

(v) Testing the effect of a strategy for exploiting surrogate sire technology to do

the dissemination.

Results are presented as the mean of one hundred replicates for each scenario

and encompass the genetic merit of nucleus males that would give rise to commercial

animals at a given time point.

Whole-genome sequence data and historical evolution

Sequence data were generated using the Markovian Coalescent Simulator

(MaCS) [10] and AlphaSim [11] for 400 base haplotypes for each of ten 10

chromosomes in the genome. Chromosomes (each 100 cM long and comprising 10^8

base pairs) were simulated using a per site mutation rate of 2.5×10^{-8}, a per site

recombination rate of 1.0×10^{-8}, and an effective population size (N_e) that varied over

time in accordance with estimates for the Holstein cattle population [12–15] as

follows: N_e was set to 100 in the final generation of the coalescent simulation, to $N_e =$

1256, 1000 years ago, to $N_e = 4350$, 10,000 years ago, and to $N_e = 43,500$, 100,000 years

ago, with linear changes in between these time-points. The resulting sequences had

approximately 540,000 segregating sites.

Quantitative trait variants

A quantitative trait was simulated by randomly sampling 10,000 QTV from

the segregating sequence sites in the base population, with the restriction that 1000

QTV were sampled from each of the 10 chromosomes. For these QTV, the allele

substitution effect was randomly sampled from a normal distribution with a mean of 0

and standard deviation of 0.01 (1.0 divided by the square root of the number of QTV).

The effects of the QTV were in turn used to compute true breeding values (TBV) for a

trait.

To simulate selection using genomic selection, without the full computational

burden and complexity of simulating training sets and then calculating genomic

breeding values using best linear unbiased prediction, pseudo genomic breeding

values (**GEBV**) were calculated as TBVs with certain levels of noise added to them to

render them representative of genomic breeding values with different levels of

accuracy. Pseudo GEBV with accuracy of 0.50, 0.70 and 0.90 were used.

Pedigree structure, gamete inheritance, and recombination rates

After generating whole-genome sequence data and QTV, a pedigree of 42

generations for the nucleus population was simulated. Each generation included 1,000

(**SmallScenario**) or 5,000 (**BigScenario**) individuals of equal sex ratio. The different

numbers of individuals were used to quantify any impact of nucleus population size

on the benefit of surrogate sire technology. All females (500 for the SmallScenario or

2,500 for the BigScenario) and 50 males, selected by truncation selection on GEBV,

were used as the parents of each generation. In the first generation of the recent

historical animal breeding population (i.e. generation -22), the chromosomes of each

individual were sampled from the 400 base haplotypes. In later generations (i.e.,

generations -21 to 20), the chromosomes of each individual were sampled from

parental chromosomes with possible recombination events. A recombination rate of 1

Morgan per chromosome was used, resulting in a 10 Morgan genome. Crossovers

were simulated to occur without interference.

Scenarios

Two different strategies were used to identify males from the nucleus who,

through the use of conventional multiplication or surrogate sire technology, would

give rise to commercial animals. The first strategy, which represents a conventional

strategy, involved selecting the top 50, 200, or 500 males in both the SmallScenario

and the BigScenario based on GEBV. The second strategy was inspired by the multi-

stage testing strategies that are common in plant breeding. Here two-stage and three-

stage testing strategies were evaluated. In the two-stage strategy a subset of potential

elite donors was first selected based on GEBV (**S0**) and the best of these were then

progeny tested (**S1**). Based on these test results the best individual or set of

individuals was used as elite donors of spermatogonial stem cells to surrogate sires. In

the three-stage strategy a subset of potential elite donors was first selected based on

GEBV (**S0**), the best of these were then progeny tested (**S1**), and finally the best of

these were further progeny tested with even more resources per individual (**S2**). Thus,

just as is the case with plant breeding programs, as the testing progressed through

these stages the number of individuals being tested was reduced and the precision

with which they were tested increased. And as with plant breeding programs, the

small numbers of individuals that were tested in the final stages were very intensively

tested.

To quantify the impact of different amounts of testing resources and different

distributions of these resources across the different stages of testing different

accuracies of GEBV at S0 were simulated, different numbers of elite donor candidates

tested and different numbers of progeny for testing each elite donor candidate at S1

and S2 were simulated. At S0 GEBV with accuracy of 0.50, 0.70 and 0.90 were used.

To ensure that each breeding program had the same costs a total of 14,000 progeny

were assumed to be available for the progeny testing stages. When this involved a

single stage of progeny testing (S1 only) the 14,000 progeny were used to test 14 elite

donor candidates each tested with 1,000 progeny, 28 elite donor candidates each tested with 500 progeny, 56 elite donor candidates each tested with 250 progeny, 112 elite donor candidates each tested with 125 progeny, 224 elite donor candidates each tested with 63 progeny or 448 elite donor candidates each tested with 31 progeny.

When the multi-stage testing strategy involved two-stages of progeny testing (S1 and S2) either 2,000, 4,000 or 6,000 of the progeny were assigned to S1 testing with the remaining 12,000, 10,000 or 8,000 being assigned to S2. At S1 either 100, 200 or 400 elite donor candidates were tested. Thus, when 2,000 progeny were used at S1 the 100 elite donor candidates were each tested with 20 progeny, the 200 elite donor candidates were each tested with 10 progeny, and the 400 elite donor candidates were each tested with 5 progeny. When 4,000 progeny were used at S1 the 100 elite donor candidates were each tested with 40 progeny, the 200 elite donor candidates were each tested with 20 progeny, and the 400 elite donor candidates were each tested with 10 progeny. When 6,000 progeny were used at S1 the 100 elite donor candidates were each tested with 60 progeny, the 200 elite donor candidates were each tested with 30 progeny, and the 400 elite donor candidates were each tested with 15 progeny.

At S2 either 10 or 20 elite donor candidates that had been advanced from S1 were tested. When 12,000 progeny remained to be used at S2 the 10 elite donor candidates were each tested with 1,200 progeny and the 20 elite donor candidates were each tested with 600 progeny. When 10,000 progeny remained to be used at S2 the 10 elite donor candidates were each tested with 1,000 progeny and the 20 elite donor candidates were each tested with 500 progeny. When 8,000 progeny remained

to be used at S2 the 10 elite donor candidates were each tested with 800 progeny and

the 20 elite donor candidates were each tested with 400 progeny.

From each of these testing strategies either 1 or 5 elite donors were to donate

spermatogonial stem cells to surrogate sires used in the commercial layer. It was

assumed that producing a sufficient number of surrogate sires from these donors to

meet the needs of the commercial layer would take one further year. In practice, this

period could be used to perform even more intense testing on the elite donors.

The time taken to transfer germplasm from the nucleus to the commercial

layer was assumed to be 4 years for the conventional breeding program design that

did not involve the use of surrogate sire technology, 5.5 years for the multi-stage

testing strategy with two-stages of testing and 6.5 years for the multi-stage testing

strategy with three-stages of testing. However, in order to enable comparison, the

genetic merit of sires that would be used in the commercial layer but produced by

either the conventional or the surrogate sire technology we needed to adjust the

expected genetic merit of the sires from the conventional strategy to account for the

extra genetic lag invoked by the multiplication process. The elite donors to be used as

surrogates sires would not need to be multiplied and thus would not have their genetic

merit diluted in the way that those who follow a conventional multiplication process

would have, because the females in the multiplier layer would be of lower genetic

merit than that of the sires entering it. To account for this, we calculated a dilution

factor as follows:

$$D = \sum_{i=1}^{k=6} \Delta G_{t-i} * (w)^i$$

Where D is the dilution factor, ΔG is the average genetic gain in generation t

and w is the relationship coefficient between the animal and his maternal male

ancestor in the generation i. k was fixed to 6 meaning that for every correction we

went back 6 generations with w ranging from 0.5 in $t-1$ generation to 0.015625 in

the $t-6$ generation. This dilution factor added an equivalent of ~1.04 years of extra

time to the conventional multiplication process.

All of these different factors gave 96 different scenarios for each level of

GEBV accuracy assumed. The steps and the resources used at each step are shown in

Figure 2.

Comparison of different scenarios

To ensure that sufficient numbers of generations had been traversed to allow

the dilution factor to be calculated we chose to present the results in terms of the

genetic merit of terminal sires used in the commercial layer emerging from

generation 11 and each subsequent generation calculated in units of the standard

deviation of TBV of the nucleus animals in the base generation ($\sigma_{TBV_{base}}$) as

$(\overline{TBV_{curr}} - \overline{TBV_{base}})/\sigma_{TBV_{base}}$, where $\overline{TBV_{curr}}$ is the mean TBV of the nucleus males

chosen n years earlier that gave rise to sires of the commercial population and

$\overline{TBV_{base}}$ is the mean TBV of nucleus animals in the base generation. Generation 0

was used as the base generation to observe the genetic improvement since the start of

the future generations of animal breeding. Calculating the genetic merit in the

commercial layer in this way allowed the different strategies to be compared in

absolute terms and at specific time points in terms of the number of years it takes the

conventional breeding program when selecting the top 50 males to deliver the same

level of genetic merit to the commercial layer.

Results

Using surrogate sire technology was effective for increasing the genetic merit of terminal sires used in the commercial layer. When animals in the commercial layer were progeny of surrogate sires their genetic merit was as much as 6.5 to 9.20 years-worth of genetic gain higher than those that were progeny of sires emerging from different variants of a conventional multiplication process. In both the SmallScenario and the BigScenario the most effective strategy for identifying donors involved three-stages of selection in which genomic selection was used at the first stage, 100 donors were tested each with 60 progeny at the second stage and 20 donors were tested each with 400 progeny at the third stage (see Table1 for details). The benefit of surrogate sire technology was greatest when the accuracy of GEBV was lowest and when the conventional strategy required large proportions of the nucleus males to be used in the commercial layer.

In what follows the results are divided into 4 sub sections for ease of presentation: (i) comparison of conventional and the best performing strategies to identify elite donors for surrogate sires; (ii) comparison of different variants of a two-stage testing strategy to identify elite donors for surrogate sires; (iii) comparison of different variants of a three-stage testing strategy to identify elite donors for surrogate sires; and (iv) comparison of two and three-stage testing strategies to identify elite donors for surrogate sires.

Comparison of conventional and the best performing strategies to identify elite donors for surrogate sires

Figure 4 and Figure S1 shows the average genetic merit of commercial sires

derived from the best performing strategy to identify elite donors and the conventional

strategy that did not use surrogate sires plotted against time, for three different GEBV

accuracies (0.5, 0.7 and 0.9) for both the SmallScenario and the BigScenario

respectively. We compared the surrogate sires strategy with three different

conventional strategies that either selected the top 50, 200, or 500 males to be

advanced to the multiplier layer. At all points in time and for all three GEBV

accuracies commercial sires derived from the best performing strategy to identify elite

donors had a higher genetic merit than those derived from the conventional strategy

and the benefit was greater when more males were advanced to the multiplier layer

for the conventional strategy. The benefit of using surrogate sire technology decreased

as the accuracy of GEBV increased. Across time the difference between using the

conventional strategy and using surrogate sire technology was almost constant. These

trends were common between the SmallScenario and the BigScenario, although with

differences in magnitude.

Table 1, enumerates some of the main results than can be observed in Figure 4

and Figure S1. Across all accuracies and scenarios tested the best performing strategy

to identify elite donors involved a first screen of all candidates based on a GEBV

followed by two-stages of progeny testing on a reduced set of candidates with the

outcome being the release of a single donor. The benefit was greater when more males

were advanced to the multiplier layer for the conventional strategy. When the

accuracy of GEBV was low (0.5) the best strategy involved using 6,000 progenies

distributed across the 100 candidates at the first stage of testing followed by using

8,000 progenies distributed across the 20 candidates at the second stage of testing.

This process of testing and producing a surrogate sire was assumed to take 2.5

additional years compared to the conventional strategy. After accounting for this extra

time and for the dilution factor in the conventional multiplication process, on average

across all time points the use of surrogate sire technology with this strategy delivered

between 6.50 years' worth of extra genetic gain in commercial sires compared to

when 50 sires were advanced to the multiplier layer for the conventional strategy and

9.20 years' worth of extra genetic gain in commercial sires compared to a when 500

sires were advanced to the multiplier layer for the conventional strategy for the

SmallScenario. For the BigScenario the equivalent values were between 2.72 and 4.10

years' worth of extra genetic gain in commercial sires.

When GEBV accuracy was higher the optimal distribution of testing resources

was slightly different. Instead of testing 100 candidates at the first stage of progeny

testing, as was the case when the GEBV accuracy was 0.5, 200 candidates were tested

in the best performing scenario. All other parameters were the same as those for when

the GEBV accuracy was 0.5. The benefit of using surrogate sire technology decreased

with increasing accuracy of GEBV and the magnitude changed significantly between

the SmallScenario and the BigScenario.

On average across all time points the SmallScenario delivered between 6.50

and 9.20 years' worth of extra genetic gain in commercial sires when GEBV accuracy

was 0.5. When GEBV accuracy was 0.7 these values reduced to between 4.52 and

7.20 years' worth of genetic gain in commercial sires and when GEBV accuracy was

0.9 they further reduced to between 2.40 and 5.00 years worth of genetic gain.

On average across all time points the BigScenario delivered between 2.72 and

4.10 years' worth of extra genetic gain at the commercial level when GEBV accuracy

was 0.5. When GEBV accuracy was 0.7 these values reduced to between 2.10 and

3.50 years' worth of genetic gain in commercial sires and when GEBV accuracy was

0.9 the impact of using surrogates sire technology delivered on average to between

1.20 and 2.50 years' worth of genetic gain.

The differences in the SmallScenario and the BigScenario were due to the

different proportions of candidate males that give rise to commercial sires. In the

SmallScenario 10% to 100% contribute while the in the BigScenario 2% to 20%

contribute.

For simplicity of presentation and based on the consistency of trends described

above, in the following sections we only present comparisons to the conventional

strategy in which 50 males were advanced from the nucleus to the multiplier layer.

Comparison of different variants of a two-stage testing strategy

Tables 2 and 3 show the performance of different two-stage testing strategies.

Performance is measured as the average years' worth of extra genetic gain at the

commercial level delivered by surrogate sire technology compared to a conventional

strategy for both the SmallScenario (Table 2) and the BigScenario (Table 3).

Consistent with the results reported in the previous section the benefit of surrogate

sires was always lower when the accuracy of GEBV was higher. In some situations,

the benefit was minimal. In all cases, there was an intermediate optimum for the

numbers of candidates tested. Selecting 5 candidates to become elite donors was

always worse than selecting one candidate. This behaviour was observed in both the

SmallScenario and the BigScenario although with some interesting differences. The

BigScenario showed a general shrinkage of years' worth of genetic gain if compared

to the SmallScenario, resulting in a general increase in the number of scenarios that

showed only a small benefit from use the surrogate sires technology (See tables S1

and S2).

At all levels of GEBV accuracy the best or joint best of the two-stage testing

strategies involved a screen of candidates based on genomic breeding values followed

by a single stage of progeny testing in which 112 candidates were tested, each with

125 progeny, followed by the choice of a single donor. Using this strategy on the

SmallScenario gave 5.26, 3.58, or 2.20 years worth of extra genetic gain in

commercial sires (Table 2) whereas applying it to the BigScenario gave 2.54, 1.97 or

1.07 years worth of extra genetic gain in commercial sires and when GEBV accuracy

was 0.5, 0.7 or 0.9 respectively (Table 3).

Just as for the case of selecting one elite donor of spermatogonial cells for

surrogate sires, when selecting five elite donors of spermatogonial cells for surrogate

sires, testing 112 candidates, each with 125 progeny, gave the highest benefit with

3.62, 2.55 and 1.24 years worth of extra genetic gain in SmallScenario (Table 2) and

2.00, 1.10 and 0.95 in BigScenario (Table 2).

Comparison of different variants of a three-stage testing strategy

Tables 4 and 5 shows the performance of different three-stage testing

strategies when either 1 or 5 sires were selected to become an elite donor. By varying

several parameters we tested 216 (108 for the SmallScenario and 108 for the

BigScenario) different strategies of three-stage testing when the total resources

available for progeny testing were kept constant across strategies and equal to that of

the two-stage testing strategies. The parameters were the split of resources between

the first and second stages of progeny testing, the number of donor candidates and

number of progeny used to test each donor candidate at each of the stages of progeny

testing, the accuracy of GEBV for the first screening of donor candidates and the

number of donor candidates finally chosen for production of surrogate sires.

Three-stage testing gave a greater benefit than two-stage testing. As for two-

stage testing, using one elite donor for surrogate sires gave a greater benefit than using

five elite donors and the benefit of surrogate sire technology was greater when the

accuracy of GEBV was lower. A total of 14,000 progenies were split across the two

stages of progeny testing. Increasing the resources in the first stage of testing gave

increased benefit from using surrogate sire technology. For example, with the

SmallScenario when the accuracy of GEBV was 0.5 and 6,000 progeny were used in

the first progeny test and 8,000 were used in the second progeny test and when one

elite donor was used, the best performing strategy gave a benefit of 6.50 years worth

of extra genetic gain in commercial sires with surrogate sire technology compared to a

conventional strategy where 50 top males were selected using GEBV. This was a

greater benefit than the 5.80 years worth of extra genetic gain for the best performing

strategy when splitting the progeny 4,000 for the first stage and 10,000 for the second

stage which was in turn better than the 5.37 years worth of extra genetic gain for the

best performing strategy when splitting the progeny 2,000 for the first stage and

12,000 for the second stage. This trend of greater benefit when more progeny were

dedicated to the first stage of progeny testing was observed for almost all strategies

that were tested.

For the SmallScenario the difference between testing 100 or 200 donor

candidates at the first stage of progeny testing was not consistent. That said, when the

accuracy of GEBV was 0.5 testing 100 candidates at the first stage of progeny testing

was usually better than testing 200 and testing 200 candidates was usually better than

testing 400. At higher accuracies, there were little differences between testing 100 or

200 candidates at the first stage of progeny testing but both of these sets of strategies

were usually better than strategies that tested 400 candidates at the first stage of

testing.

In the SmallScenario testing 20 elite donor candidates at the second stage of

progeny testing was almost always better than testing 10 candidates. A total of 54

strategies were evaluated for SmallScenario only. In only 6 of these strategies testing

10 candidates was better than testing 20.

Overall for the SmallScenario, when the GEBV accuracy was 0.5, the best

performing of different three-stage testing strategies used 6,000 progenies in the first

stage of progeny testing to test 100 candidates each with 60 progenies, used 8,000

progenies in the second stage of progeny testing to test 20 candidates each with 400

progeny and selected a single donor for surrogate sires. This strategy test gave a

benefit of 6.50 years worth of extra genetic gain in commercial sires compared to a

conventional strategy. The same distribution of testing resources was also the joint

best when 5, instead of 1, donors were chosen for surrogate sires.

Although all the results presented here refer to the SmallScenario the same

trend was observed for the BigScenario with only differences in magnitude of years'

worth of genetic gain, specifically a reduction in the benefit observed for surrogate

sire technology (See table S1 and S2).

Discussion

The results of this paper suggest that surrogate sire technology could be very beneficial for the dissemination of genetic gain from livestock breeding programs. In summary, our results indicate that benefits of the as much as 6.5 years' worth of genetic gain in commercial sires could be realized with surrogate sire technology. It was best to identify donors for surrogate sires via a three-stage testing strategy involving a first screen based on GEBV followed by two stages of progeny testing. The benefits of surrogate sires were greater when GEBV accuracy was lower. To discuss these results we divide the discussion into four sections: (i) Possible explanations for the observed trends; (ii) Justification and impact of assumptions that were made; (iii) The potential impact of surrogate sire technology on the redesign of animal breeding programs; and (iv) Risks and opportunities of using surrogate sire technology.

Possible explanations for the observed trends

That surrogate sire technology generates such a benefit in terms of years' worth of genetic gain can be explained in the context of the breeders' equation. In any cohort the best few individuals will be some number of standard deviations above the average of the cohort. For example, for the scenario where surrogate sires technology delivered 6.5 years' worth of genetic gain the best individual was 2.69 standard deviations above the mean on average across all years simulated. In contrast, the top 50 individuals were 1.95 standard deviations above the mean on average across all years simulated. Given the breeding program, in this scenario, proceeded at a rate of genetic progress of 0.36 standard deviations per year then choosing the best male as a donor for surrogate sires rather than the top 50 males would produce commercial sires

that are better for more than 8 years' worth of genetic gain. However, given the

process of identifying donors for surrogate sires, or the top 50 males in the

conventional scheme, does not have perfect accuracy and that it takes time to identify

donors and produce surrogate sires (we assumed 3 years) the final result reduces to

6.5 years' worth of genetic gain.

The benefit of surrogate sires technology depended on the proportion of male

candidates that the conventional scheme used to give rise to commercial sires. If the

breeding program needed to use a large proportion of its male candidates (e.g., the top

200 or 500) the benefit of surrogate sire technology was greater than if it needed to

use a few. Again, this result is entirely consistent with quantitative genetic theory. The

average genetic merit of the top 100 or more male candidates will be lower than that

of the top 50 candidates. Therefore, if a breeding program needs to deliver 50 sires to

the commercial level the benefit of using surrogate sires compared to a conventional

strategy will be lower than if it needs to deliver 100 sires. This explains why the

benefit of surrogate sires appeared lower for the BigScenario which had 2,500

candidate males per year than for the SmallScenario which had 500 candidate males

per year. If both scenarios deliver the same number of sires to the commercial level

(e.g., 100 sires) the average difference in genetic merit between the top 100 and the

top single individual will be lower for the BigScenario than for the SmallScenario.

The same logic also explains why selecting five as opposed to one donor for surrogate

sires gave a lower benefit.

The trends observed in terms of differences in the benefit accruing from

different strategies to test the benefit of surrogate sires also can be explained in the

context of the breeders' equation. When the accuracy of GEBV, which was used to

perform the first stage of testing, was lower the benefit of surrogate sire technology

was higher. Under the conventional strategy, the average genetic merit of the nucleus

candidates that gave rise to commercial sires will be lower when the GEBV accuracy

from the first-stage is lower than when it is higher. With surrogate sires this reduction

in genetic merit due to the low accuracy of GEBV for the first screening is

compensated by the subsequent progeny testing stages. This is in line with the

analysis of Dickerson and Hazel [16], who compared the use of progeny testing as a

supplement to earlier culling on own or sibling performance. Their conclusion was

that the use of progeny test is warranted when heritability is low in which case

accuracy of GEBV from own or sibling phenotypes (or genomic prediction in our

study) is low. Genomic selection can be thought of as a light touch first screen, the

purpose of which is to identify the top portion of animals which then have a large

amount of progeny testing resources focused upon them. The purpose of the

subsequent progeny testing stages can be thought of as being to search for the best

individual within this group.

534   This same logic also explains why the three-stage strategy was better than the

two-stage strategy. In the three-stage strategy there were two-stages of progeny

testing after the initial stage compared to just one stage of progeny testing in the two-

stage strategy. In the three-stage strategy the first stage of genomic selection can also

be thought of as a light touch second screen, the purpose of which is to identify the

top portion of animals (e.g., the top 50%) which then have a large amount of progeny

testing resources focused upon them. In the three-stage strategy the first stage of

progeny testing serves to use a portion of the overall testing resources to evaluate

many candidates relatively inaccurately in order to discard most candidates. In the

second stage of progeny testing the remaining candidates can be tested more

intensively and the final candidate can be chosen very precisely. In terms of the breeders' equation the first stage of progeny testing maximizes selection intensity while the second stage maximize selection accuracy. The three-stage strategy appears to address both of these parameters more optimally than the two-stage strategy.

While there is substantial body of literature on multi-stage selection [17–21], we are not aware of literature that addresses joint optimisation of proportion of selected individuals, testing resources (i.e., accuracy), and number of selection stages. However, the observed trends in this study have logical explanations. All the performances observed in the two-stages strategy are expected. Although it is well known that increasing the number of progeny per candidate leads to an increasing progeny test's accuracy [22,23], it is equally well known that the number of candidates to be tested is an extremely important variable and the trade-off between the two is crucial to optimize the selection process. In our simulations, the optimum was reached when 112 candidates were progeny tested, given a fixed number of available resources, in our case 14,000 progeny. This behaviour was consistent across the different levels of GEBV accuracy, although the level of accuracy heavily influenced the amount of extra genetic gain because higher GEBV's accuracy directly translates to a better selection of the animals even before performing the progeny test. These trends are consistent with the long-established use of multiple stages of testing in plant breeding programs[8]. Most plant breeding programs that deliver a single genotype (e.g., inbred line) to the commercial layer use multistage testing strategies to identify this individual. Typically, in these multistage testing strategies a very large number of individuals are imprecisely evaluated at the first stage of testing. At each subsequent stage of testing the number of individuals tested is reduced and the precision with which they are evaluated is increased.

Justification and impact of assumptions that were made

There are a huge range of strategies that could be taken for identification of donors for surrogate sires and we only compared a small subset in this study. We choose the tested range of scenarios because we believe they could demonstrate the properties of surrogate sire technology. They show that in some circumstances surrogate sires can deliver a large benefit and in others small benefit. We chose the three levels of GEBV accuracy as these levels reflect what might be possible in breeding programs of various sizes in the next years. To ensure that all strategies used an equal set of resources we set the total number of progeny involved in progeny testing to 14,000. We chose this number as it was divisible in many ways and thereby enabled several strategies to be compared and because this number was similar the 10,000 progeny that would be used by an animal breeding program that each year tested 100 candidates each with 100 progeny, a scale of progeny testing that was not uncommon in some animal breeding programs before the advent of genomic selection.

In the two-stage testing strategy the total testing resources were distributed across many or few candidates. As expected testing an intermediate to higher number of candidates (i.e., 112 to 224) tested using relatively small numbers of progeny (i.e., 125 to 63) gave higher benefits than testing a few candidates (e.g., 14) with many progeny or a very high number of candidates (448) with very few progeny (31). These trends fit within the expectations of the breeders' equation and occur due to the interplay between selection intensity and selection accuracy. However, when the selected elite donors of spermatogonial cells for surrogate sires are to be used to produce huge numbers of progeny at the commercial level, the risk of a donor

carrying some major defect that was not identified by the testing process must also be

minimized. For this reason, it is unlikely that a strategy in which donors are tested

with a single stage of progeny testing using a ~200 or less progeny would ever be

used by a commercial breeding program.

It was this logic that motivated us in our design of the three-stage testing

scheme. Our intuition was that in the first stage of progeny testing many candidates

could be evaluated with relatively low accuracy but in the second stage of progeny

testing a handful of individuals could be tested with many progeny. We evaluated 10

or 20 candidates and tested them using either 800 or 400 progeny as we felt that this

would reflect a level of testing that would give a very high degree of certainty that the

selected donors would not carry any major defects that were undetected by the testing

process.

A major assumption of this study was the amount of time it took to identify

elite donors and then to make surrogate sires. In this study, we assumed that this

would take 1.5 years for the two-stage strategy and 2.5 years for the three-stage

strategy. This was based on the logic that in pigs it would take ~6 months for a male

to reach sexual maturity, a further ~4 months for a successful gestation, and a further

~6 months to collect terminal line phenotypes on progeny. Thus, to perform the first

stage progeny testing on an individual would take ~16 months. We assumed that it

would take another year to perform a second round of progeny testing and equally one

year to produce surrogate sires. In contrast, we assumed that the genetic improvement

from the conventional program could be delivered to the commercial population

within 3.5 to 4 years in a highly optimized breeding program but that this genetic

merit would be diluted to some degree by the genetic merit of females in the

multiplier layer. This resulted in the two-stage strategy taking 5.5 years and the three-

stage strategy taking 6.5 years for using surrogate sire technology compared to about

~5.04 years for the conventional strategy (calculated as 4 years in time plus a further

~1.04 years to account for the dilution of genetic merit). It is likely that this time

could be shortened or lengthened for both the strategies using surrogate sire

technology and the conventional strategy in several ways and depending on the

assumed species. The benefit of using surrogate sire technology would change

accordingly.

The potential impact of surrogate sires on the redesign of animal breeding

programs

Animal breeding programs maximize the genetic merit of commercial animals

within the available financial, physical, technical and physiological constraints.

Implicitly a breeding program has two objectives: (i) improving the mean of the

population; and (ii) developing and delivering a product to the commercial producers.

In dairy cattle for example, before the advent of genomic selection, breeders used

progeny testing schemes that intensively evaluated relatively small numbers of

candidate males and used the best performing of these as both parents to improve the

population and as a breeding product to be used by the commercial layer. In doing so

dairy cattle breeders maximised selection accuracy but were constrained in their

ability to increase selection intensity and decrease generation interval. However, the

products used by commercial producers were well tested and therefore an individual

producer could rely on using relatively few sires, who together could serve entire

geographic regions. The advent of genomic selection changed this paradigm

significantly. Under genomic selection progeny testing of small numbers of

candidates has been replaced with a genomic testing of large numbers of candidates.

Those with best predictions are used as parents to improve the population and as

products sold in teams of sires (i.e., a group of sires sold together rather than a single

sire sold on its own). In doing so dairy cattle breeders increased selection intensity

and reduced generation interval, but are somewhat constrained in their ability to

achieve very high levels of accuracy. Given that each candidate male has not had their

merit assessed based on phenotypes of their progeny there is a risk that certain sires

are not that good or may carry mutations that are highly detrimental (e.g., a *de-novo*

mutation that prevents progeny from lactating) [24–26]. To overcome this risk,

breeders recommend that commercial producers use semen of a larger number (i.e.,

team) of sires and limit their use of any one sire.

A strategy for using surrogate sire technology would need to exploit aspects of

both genomic selection and progeny testing. Genomic selection could be used to drive

the population improvement and, as demonstrated in the present paper, to identify a

subset of candidates that could enter a progeny testing scheme. The role of the

progeny testing component would be to ensure that the surrogate sires that are

released to the commercial sector are not significantly worse than it is predicted to be

based on GEBV and that they do not carry detrimental mutations. As demonstrated by

the results of the present study a multi-stage strategy to progeny testing was a more

efficient use of resources than a single-stage strategy that was commonly used in by

animal breeding programs in the past. Multi-stage testing has a long history of use in

plant breeding programs which also have a long history delivering products to

commercial producers in a way that is highly analogous to what surrogate sires would

enable for animal producers.

The majority of commercial producers for all of the major stable crops (maize,

wheat, rice) use inbred lines or hybrids between a pair of inbred lines at the

commercial level. These inbred or hybrid lines can be grown on huge areas. Plant and

animal breeding program designs have diverged somewhat over the years owing to

differences in biology, economics, and technical possibilities. Surrogate sire

technology, combined with genomic selection, could result in a coalescence of

designs across plant and animal breeding. One such design that could apply to both is

the two-part scheme recently proposed by Gaynor et al. (2017)[9]. In this scheme,

rapid recurrent selection based on genomic breeding values is used to increase the

mean of the population, while a multi-stage testing strategy is used to periodically

extract, test, and develop a product from the population. The population improvement

component resembles the traditional nucleus layer of animal breeding programs, while

the product development component resembles the traditional multi-stage testing to

derive inbred or hybrid lines of plant breeding programs. The latter could also be seen

as an improved multiplication layer of animal breeding programs that exploit breed

complementarity to deliver a commercial product.

In the present work, we focused on the use of surrogate sires to serve as sires

of commercial individuals (e.g., a terminal sire in a pig population). To do this, donors

for surrogate sires were chosen based on their general combining ability. The strategy

could also be extended so that specific combining ability was exploited to produce a

relatively homogenous set of females from a maternal line that are crossed with single

terminal male (via surrogate sires). Use of specific combining ability is widespread in

hybrid crops where it exploits complementarity of pairs of individuals and heterosis

generated by the "nicking together" of specific pairs of individuals. An extension of

the strategy for the use of surrogate technology that was proposed in the present paper

to enable it to exploit specific combining ability could involve extending the multi-

stage testing to include a later set of stages where specific crosses are progeny tested.

Because in livestock the parents of individuals are outbred (compared to crops where

they are inbred) a tiered strategy may be needed in the maternal line(s) which

homogenizes the haplotypes that are present in dams released from the maternal lines.

For example, using a single surrogate sire, grandsire and great-grandsire on the

maternal population would result in a pool of females that carried one of two

haplotypes for 87.5% (0.5 + 0.25 + 0.125) of their genome. The terminal surrogate

sire would be chosen for what could be termed pseudo specific combining ability that

nicked well with these haplotypes.

Risks and opportunities of using surrogate sire technology

Surrogate sires present risk and opportunities to commercial production. The

most obvious risk relates to the genetic homogeneity of commercial animals if a

single surrogate sire, or a set of very closely related surrogate sires were used. If a

disease emerged that this homogenous group of animals was susceptible to it could

have major detrimental impacts on commercial production. Having such large groups

of homogeneous animals would also increase the selection pressure on disease

pathogens to evolve pathogenicity to the group. Plant breeders and commercial crop

growers have extensive experience in managing the potential to have genetic

homogeneity across large segments of the production area. They have developed

strategies to minimize the risk of disease outbreaks and other failures such as crop

rotation, using multiple varieties on a farm, creating varietal blends consisting of

multiple genotypes, and taking holistic strategies to pathogen management [27]. Aside

from rotation, which may be difficult to use in practice, these strategies would likely

have important roles in ensuring the effectiveness of surrogate sire technology in

livestock.

717   The most obvious opportunity emanating from surrogate sire technology also

relates to the genetic homogeneity of commercial animals and can also draw on

practices that are well established in crop production. In crops, management plans

(e.g., https://catalog.extension.oregonstate.edu/em9004) are supplied to a farmer

alongside the seed. These plans are specifically tailored to the genotype of the variety

based on extensive sets of field trials. They include recommendations for target

market, expected performance, optimum sowing date, seeding rate, soil type and

water, fertilizer, pesticide and fungicide requirements. These management plans

complement the genetics of the variety and increase the benefit obtained from the

genetic potential in a generic environment. Similar management plans could be

developed for surrogate sires and the benefits would be similarly expected to exceed

the benefit (e.g., 6.5 year's worth of genetic gain) that was observed in the present

study for the genetics alone. The phenotype data collected to development of the

management plans would also serve to further test and validate a particular surrogate

sire.

732   Another obvious opportunity emanating from surrogate sire technology that

also relates to the genetic homogeneity is the potential for increasing the homogeneity

of product. In livestock production, uniformity of end product is an important topic. In

meat animals, for instance, uniformity has economic benefits because excessive

variability in carcass weight or conformation is penalized by slaughterhouses [28,29].

A genetically homogeneous commercial population, achieved through the use of

surrogate sire technology, could aid product uniformity. If this was to be achieved

most of this increase in uniformity would need to emanate from matching very

specific management plans to the homogenous genetics because homogeneous

genetics in itself has limited ability to increase phenotypic homogeneity. Van Vleck

[30] showed that in the context of cloned animals, if heritability is 25%, then the

phenotypic standard deviation among clones would be 87% of that of uncloned

animals and only if heritability is 100% will clone mates have complete uniformity.

Compared to conventional multiplication surrogate sire technology enables

shorter lag between nucleus and production animals and requires a smaller number of

parents contributing to commercial animals. This offers several advantages including

the ability to rapidly change the entire genetics at the production level. This could be

used to rapidly respond to sudden changes in requirements such as pressure from a

new disease or the emergence of a new market for the product that has specific

requirements (e.g., marbling levels in meat).

Surrogate sire technology would be costly to implement in practice because it

would require capacity in advanced molecular biology and infrastructure for progeny

testing to maintained. However, it presents other opportunities through which costs

can be saved. For example, multiplier populations to produce terminal sires would not

need to be maintained.

Finally, surrogate sire technology either by the release of a handful of

individuals from the nucleus or through the release of individuals selected for pseudo

specific combining ability presents breeding programs with an enhanced opportunity

to protect its intellectual property. This protection of intellectual property would the

breeding companies incentive to invest in breeding programs thus avoiding critical

market failure, which increases benefits to farmers. Typically breeding companies

share the benefits two-thirds to the farmers and one third to the breeding company

which more than offsets the cost of purchase of the genetics by a produced. The most

spectacular example of the benefits of cross-breeding are seen in Maize which has

seen a 6-fold increase in productivity since hybrid breeding was introduced in the

1930's [31].

Conclusions

The results of this study showed that using surrogate sire technology would

significantly increase the genetic merit of commercial sires, by as much as 6.5 to 9.20

years' worth of genetic gain compared to a conventional breeding program. The

simulations suggested that an identification strategy involving three-stages, the first of

which uses genomic selection while the subsequent stages use progeny testing was the

most effective strategy of all of those tested. The use of one or a handful of elite

donors to generate the production animals would be very different to current practice.

While the results demonstrate the great potential of surrogate sire technology there are

considerable risks but also other opportunities. Practical implementation of surrogate

sire technology would need to account for these.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JMH conceived the study. JMH, MB, and PG designed the study. PG performed the

analysis. JMH and PG wrote the first draft of the manuscript. MG, WOH, GG, JJ,

AJM and RCG helped to interpret the results and refine the manuscript. All authors

read and approved the final manuscript.

Acknowledgements

The authors acknowledge the financial support from the BBSRC ISPG to The Roslin

Institute, from Genus PLC, the BBSRC, and Innovate UK.

References

1. Park K-E, Kaucher AV, Powell A, Waqas MS, Sandmaier SES, Oatley MJ, et al. Generation of germline ablated male pigs by CRISPR/Cas9 editing of the NANOS2 gene. 2017;7:40176.

2. De Roos APW. Recent trends in genomic selection in dairy cattle. 62nd Annu. Meet. EAAP. Stavanger, Norway: Wageningen Academic Publishers; 2011. p. Contribution 01–7, p. 4.

3. Pryce JE, Daetwyler HD. Designing dairy cattle breeding schemes under genomic selection: a review of international research. Anim. Prod. Sci. 2012;52:107–14.

4. Meuwissen T, Hayes B, Goddard M. Accelerating Improvement of Livestock with Genomic Selection. Annu. Rev. Anim. Biosci. 2013;1:221–37.

5. Van Eenennaam AL, Weigel KA, Young AE, Cleveland MA, Dekkers JCM. Applied Animal Genomics: Results from the Field. Annu. Rev. Anim. Biosci. 2014;2:105–39.

6. Knol EF, Nielsen B, Knap PW. Genomic selection in commercial pig breeding.
Anim. Front. 2016;6:15.

7. Visscher P, Pong-Wong R, Whittemore C, Haley C. Impact of biotechnology on
(cross)breeding programmes in pigs. Livest. Prod. Sci. 2000;65:57–70.

8. Acquaah G. Principles of Plant Genetics and Breeding. 2nd Edition edition.
Hoboken, NJ: Wiley-Blackwell; 2012.

9. Gaynor RC, Gorjanc G, Bentley AR, Howell P, Jackson R, Woolliams JA, et al. A
two-part strategy for using genomic selection to develop inbred lines. Crop Sci. 2016;

10. Chen GK, Marjoram P, Wall JD. Fast and flexible simulation of DNA sequence
data. Genome Res. 2009;19:136–42.

11. Faux A-M, Gorjanc G, Gaynor RC, Battagin M, Edwards SM, Wilson DL, et al.
AlphaSim: Software for Breeding Program Simulation. Plant Genome. 2016;9.

12. Rodríguez-Ramilo ST, Fernández J, Toro MA, Hernández D, Villanueva B.
Genome-Wide Estimates of Coancestry, Inbreeding and Effective Population Size in
the Spanish Holstein Population. PLoS ONE [Internet]. 2015 [cited 2015 Apr 25];10.
Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4399946/

13. Hill WG. Understanding and using quantitative genetic variation. Philos. Trans. R.
Soc. B Biol. Sci. 2009;365:73.

14. Brotherstone S, Goddard M. Artificial selection and maintenance of genetic
variance in the global dairy cow population. Philos. Trans. R. Soc. B Biol. Sci.
2005;360:1479.

15. Hayes BJ, Visscher PM, McPartlan HC, Goddard ME. Novel Multilocus Measure
of Linkage Disequilibrium to Estimate Past Effective Population Size. Genome Res.
2003;13:635–43.

16. G.E. Dickerson, Hazel LN. Effectiveness of selection on progeny performance as
a supplement to earlier culling in livestock. J.Agr.Res. 69:459.

17. Cunningham EP. Multi-stage index selection. Theor. Appl. Genet. 1975;46:55–61.

18. Saxton AM. A comparison of exact and sequential methods in multi-stage index
selection. Theor. Appl. Genet. 1983;66:23–8.

19. Ducrocq V, Colleau J J. Optimum truncation points for independent culling level
selection on a multivariate normal distribution, with an application to dairy cattle
selection. Genet Sel Evol. 1989;21:185–98.

20. Xie C, Xu S, Mosjidis JA. Multistage selection indices for maximum genetic gain
and economic efficiency in red clover. Euphytica. 1997;98:75–82.

21. Mi X, Utz HF, Technow F, Melchinger AE. Optimizing Resource Allocation for
Multistage Selection in Plant Breeding with R Package Selectiongain. Crop Sci.
2014;54:1413–8.

22. Robertson A. Optimum Group Size in Progeny Testing and Family Selection.
Biometrics. 1957;13:442–50.

23. Meuwissen THE, Goddard ME. Optimization of progeny tests with prior
information on young bulls. Livest. Prod. Sci. 1997;52:57–68.

24. Milan D, Jeon J-T, Looft C, Amarger V, Robic A, Thelander M, et al. A Mutation
in PRKAG3 Associated with Excess Glycogen Content in Pig Skeletal
Muscle. Science. 2000;288:1248.

25. VanRaden PM, Olson KM, Null DJ, Hutchison JL. Harmful recessive effects on
fertility detected by absence of homozygous haplotypes. J. Dairy Sci. 2011;94:6153–
61.

26. Kadri NK, Sahana G, Charlier C, Iso-Touru T, Guldbrandtsen B, Karim L, et al. A
660-Kb Deletion with Antagonistic Effects on Fertility and Milk Production
Segregates at High Frequency in Nordic Red Cattle: Additional Evidence for the
Common Occurrence of Balancing Selection in Livestock. PLOS Genet.
2014;10:e1004049.

27. McDonald BA, Linde C. The population genetics of plant pathogens and breeding
strategies for durable resistance. Euphytica. 2002;124:163–80.

28. Mulder HA, Bijma P, Hill WG. Prediction of Breeding Values and Selection
Responses With Genetic Heterogeneity of Environmental Variance. Genetics.
2007;175:1895.

29. Mulder HA, Bijma P, Hill WG. Selection for uniformity in livestock by exploiting
genetic heterogeneity of residual variance. Genet. Sel. Evol. 2008;40:37.

30. van Vleck LD. Implications of cloning for breed improvement strategies: are
traditional methods of animal improvement obsolete? J. Anim. Sci. 1999;77 Suppl
2:111–21.

31. Duvick DN. Genetic Progress in yield of United States maize (Zea mays L.).
Maydica. 2005;50:193–202.

Figures

Figure 1. Example animal (on the left) and plant (on the right) breeding schemes.

Figure 2. Map of the scenarios used in the study.

Figure 3. Timeline of the different approaches to identify and disseminate genetic

improvement.

Figure 4. Average genetic merit of commercial sires derived from the best performing

strategy to identify elite donors and the conventional strategy (top 50, 200 and 500

sires) that did not use surrogate sire technology for SmallScenario (a and b) and

BigScenario (c and d) plotted against time when accuracies of genomic selection was

0.5 or 0.7.

Tables

Table 1. Best performant scenarios across all the simulations performed for 3

different levels of GEBV (0.5,0.7 and 0.9).

Table 2. Performance of the scenarios tested using the two-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) for SmallScenario.

Table 3. Performance of the scenarios tested using the two-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) for BigScenario.

Table 4. Performance of the scenarios tested using the three-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) when 1 sire was selected to

become surrogate sire for SmallScenario.

Table 5. Performance of the scenarios tested using the three-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) when 5 sires were selected to

become surrogate sires for SmallScenario.

Supplementary material

Figure S1. Average genetic merit of commercial sires derived from the best

performing strategy to identify elite donors and the conventional strategy (top 50, 200

and 500 sires) that did not use surrogate sire technology for SmallScenario (a and b)

and BigScenario (c and d) plotted against time when accuracy of genomic selection

was 0.9.

Table S1. Performance of the scenarios tested using the three-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) when 1 sire was selected to

become surrogate sire for the BigScenario.

Table S2. Performance of the scenarios tested using the three-stage testing strategy

across 3 different levels of GEBV (0.5,0.7 and 0.9) when 5 sires were selected to

become surrogate sires for the BigScenario.

Figure 1 Typical animal (on the left) and plant (on the right) breeding schemes

Figure 2 Mind map of the scenarios used in the study

Figure 3 Timeline of the different approaches to identify and disseminate genetic improvement

Figure 4 Average genetic merit of commercial sires derived from the best performing strategy to identify elite donors and the conventional strategy (top 50, 200 and 500 sires) that did not use surrogate sire technology for SmallScenario (a and b) and BigScenario (c and d) plotted against time

Figure S1 Average genetic merit of commercial sires derived from the best performing strategy to identify elite donors and the conventional strategy (top 50, 200 and 500 sires) that did not use surrogate sire technology for SmallScenario (a) and BigScenario(b) plotted against time

Table 1. Best performing strategies for identifying elite donors for 3 different levels of accuracy

Accuracy GEBV	Sires tested S1	Sires tested S2	Sires selected[*]	Resources[**]	Conv50 $\bar{X}_{YGG}$[***]	Conv200 $\bar{X}_{YGG}$[***]	Conv500 $\bar{X}_{YGG}$[***]
				Small Scenario			
0.5	100	20	1	6000S1/8000S2	6.50	7.50	9.20
0.7	200	20	1	6000S1/8000S2	4.52	6.50	7.20
0.9	200	20	1	6000S1/8000S2	2.40	4.50	5.00
				Big Scenario			
0.5	100	20	1	6000S1/8000S2	2.72	3.50	4.10
0.7	200	20	1	6000S1/8000S2	2.10	2.50	3.50
0.9	200	20	1	6000S1/8000S2	1.20	1.70	2.50

*Number of sires selected to become elite donors
**Total number of progeny allocated in the first progeny test (S1) and in the second progeny test (S2) respectively
***Average Years' worth of genetic gain (YGG) calculated across 20 generations of future breeding

Table 2 Best performing of the two-stage strategies for identifying elite donors for 3 different levels of accuracy for SmallScenario

Sire Tested	Progeny/Sire	Sires selected	$YGG_{0.5}$***	$YGG_{0.7}$***	$YGG_{0.9}$***
14	1000	1	4.12	3.0	1.80
28	500	1	4.72	3.0	1.20
56	250	1	5.11	3.50	2.20
112	125	1	5.26	3.58	2.20
224	63	1	4.80	2.77	1.30
448	31	1	3.81	2.06	1.11
14	1000	5	2.90	1.90	0.2
28	500	5	3.10	2.06	0.5
56	250	5	3.60	2.40	1.10
112	125	5	3.62	2.55	1.24
224	63	5	3.41	1.90	0.33
448	31	5	2.76	1.63	0.23

*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively*

Table 3 Best performing of the two-stage strategies for identifying elite donors for 3 different levels of accuracy for BigScenario

Sire Tested	Progeny/Sire	Sires selected	$YGG_{0.5}$***	$YGG_{0.7}$***	$YGG_{0.9}$***
14	1000	1	2.27	1.74	0.72
28	500	1	2.35	1.9	0.80
56	250	1	2.48	1.95	0.99
112	125	1	2.54	1.97	1.07
224	63	1	2.05	1.82	0.85
448	31	1	1.86	1.50	0.39
14	1000	5	1.73	1.21	0.49
28	500	5	1.73	1.25	0.69
56	250	5	1.94	1.08	0.95
112	125	5	2.00	1.10	0.95
224	63	5	1.79	1.04	0.48
448	31	5	1.02	0.77	0.33

*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively*

Table 4 Best performing of the three-stage strategies for identifying elite donors for 3 different levels of accuracy for SmallScenario when 1 sire was selected to become an elite donor

Resources[*]	Sire Tested S1	Progeny/Sire S1	Sire Tested S2	Progeny/Sire S2	$YGG_{0.5}$[***]	$YGG_{0.7}$[***]	$YGG_{0.9}$[***]
2000S1/12000S2	100	20	10	1200	5.28	3.51	2.20
	100		20	600	5.37	3.62	2.37
	200	10	10	1200	4.91	3.25	2.22
	200		20	600	5.12	3.32	2.12
	400	5	10	1200	4.52	3.68	2.05
	400		20	600	4.72	2.67	1.40
4000S1/10000S2	100	40	10	1000	5.53	3.57	2.18
	100		20	500	5.80	4.01	2.31
	200	20	10	1000	5.30	3.46	2.43
	200		20	500	5.37	3.78	2.32
	400	10	10	1000	4.31	3.29	1.64
	400		20	500	4.52	3.49	1.38
6000S1/8000S2	100	60	10	800	5.92	4.13	2.01
	100		20	400	6.50	4.18	2.16
	200	30	10	800	5.27	4.21	2.06
	200		20	400	5.71	4.50	2.40
	400	15	10	800	5.01	3.40	1.62
	400		20	400	5.78	3.51	1.22

*Number of total progeny allocated in the first progeny test (S1) and in the second progeny test (S2) respectively
*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively

Table 5 Best performing of the three-stage strategies for identifying elite donors for 3 different levels of accuracy for SmallScenario when 5 sires were selected to become an elite donor

Resources*	Sire Tested S1	Progeny/Sire S1	Sire Tested S2	Progeny/Sire S2	$YGG_{0.5}$***	$YGG_{0.7}$***	$YGG_{0.9}$***
2000S1/12000S2	100	20	10	1200	4.14	2.14	1.12
	100		20	600	4.44	2.17	1.20
	200	10	10	1200	3.04	2.11	1.20
	200		20	600	3.68	2.51	1.26
	400	5	10	1200	2.15	1.50	1.00
	400		20	600	2.19	1.43	1.05
4000S1/10000S2	100	40	10	1000	4.41	2.45	1.29
	100		20	500	4.53	2.51	1.23
	200	20	10	1000	4.10	2.24	1.10
	200		20	500	4.12	2.71	1.25
	400	10	10	1000	4.16	1.69	1.05
	400		20	500	4.18	2.02	1.17
6000S1/8000S2	100	60	10	800	4.55	3.09	1.64
	100		20	400	5.05	3.19	1.77
	200	30	10	800	4.60	2.12	1.30
	200		20	400	5.05	2.25	1.38
	400	15	10	800	4.09	1.67	1.16
	400		20	400	4.61	2.23	1.25

Number of total progeny allocated in the first progeny test (S1) and in the second progeny test (S2) respectively
*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively*

Table S1 Best performing of the three-stage strategies for identifying elite donors for 3 different levels of accuracy for BigScenario when 1 sire was selected to become an elite donor.

Resources[*]	Sire Tested S1	Progeny/Sire S1	Sire Tested S2	Progeny/Sire S2	$YGG_{0.5}$ [***]	$YGG_{0.7}$ [***]	$YGG_{0.9}$ [***]
2000S1/12000S2	100	20	10	1200	2.21	1.53	0.90
	100		20	600	2.31	1.70	0.92
	200	10	10	1200	2.11	1.55	1.0
	200		20	600	2.16	1.11	0.88
	400	5	10	1200	2.24	1.31	0.90
	400		20	600	2.28	1.50	0.77
4000S1/10000S2	100	40	10	1000	2.21	1.58	0.83
	100		20	500	2.14	1.65	0.85
	200	20	10	1000	2.11	2.07	0.91
	200		20	500	2.20	2.09	0.97
	400	10	10	1000	2.32	1.67	0.89
	400		20	500	2.33	1.99	0.93
6000S1/8000S2	100	60	10	800	2.53	2.02	0.99
	100		20	400	2.72	2.08	1.11
	200	30	10	800	2.44	1.90	1.12
	200		20	400	2.63	2.10	1.20
	400	15	10	800	2.33	1.95	0.91
	400		20	400	2.39	1.95	0.90

*Number of total progeny allocated in the first progeny test (S1) and in the second progeny test (S2) respectively
*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively

Table S2 Best performing of the three-stage strategies for identifying elite donors for 3 different levels of accuracy for BigScenario when 5 sire was selected to become an elite donor

Resources[*]	Sire Tested S1	Progeny/Sire S1	Sire Tested S2	Progeny/Sire S2	$YGG_{0.5}$[***]	$YGG_{0.7}$[***]	$YGG_{0.9}$[***]
2000S1/12000S2	100	20	10	1200	1.66	1.19	0.65
	100		20	600	1.65	1.19	0.78
	200	10	10	1200	1.56	1.25	0.63
	200		20	600	1.50	0.98	0.34
	400	5	10	1200	1.58	1.12	0.70
	400		20	600	1.64	1.18	0.77
4000S1/10000S2	100	40	10	1000	1.69	1.27	0.66
	100		20	500	1.72	1.22	0.59
	200	20	10	1000	1.55	1.25	0.43
	200		20	500	1.63	1.37	0.57
	400	10	10	1000	1.32	1.25	0.31
	400		20	500	1.45	1.35	0.44
6000S1/8000S2	100	60	10	800	1.80	1.10	0.37
	100		20	400	1.85	1.37	0.67
	200	30	10	800	1.64	1.15	0.46
	200		20	400	1.63	1.19	0.50
	400	15	10	800	1.72	1.19	071
	400		20	400	1.60	1.23	0.88

*Number of total progeny allocated in the first progeny test (S1) and in the second progeny test (S2) respectively

** Values below 0.1 years were set to 0

*** Average years' worth of genetic gain over 20 generations of future selection starting from a GEBV with accuracy 0.5, 0.7 and 0.9 respectively