Vagal Nerve Stimulation Triggers Wide-Spread Evoked Responses and Changes in Large-Scale Functional Connectivity in Rat Brains

Jiayue Cao[1,4], Kun-Han Lu[2,4], Terry Powley[1,3,4], and Zhongming Liu[1,2,4*]

[1]*Weldon School of Biomedical Engineering,*

[2]*School of Electrical and Computer Engineering,*

[3]*Department of Psychological Science,*

[4]*Purdue Institute of Integrative Neuroscience*

Purdue University, West Lafayette, Indiana, United States

*Correspondence

Zhongming Liu, PhD

Assistant Professor of Biomedical Engineering

Assistant Professor of Electrical and Computer Engineering

College of Engineering, Purdue University

206 S. Martin Jischke Dr.

West Lafayette, IN 47907, USA

Phone: +1 765 496 1872

Fax: +1 765 496 1459

Email: zmliu@purdue.edu

Abstract

Background: Vagus nerve stimulation (VNS) is a FDA-approved therapy for epilepsy and depression. However, the efficacy of VNS varies, and its mechanism is unclear. Prior studies have used functional magnetic resonance imaging (fMRI) to map brain activations with VNS in human brains but yielded inconsistent findings. The source of the inconsistency might be attributed to the complex temporal characteristics of VNS-evoked responses that cannot be fully explained by simplified response models.

Objective: We used fMRI to characterize the effects of VNS in rat brains without assuming any *priori* response model. The questions of primary interest were 1) whether and where the VNS evoked responses at the level of networks, and 2) whether and where the VNS altered the patterns of interactions among intrinsic functional networks.

Methods: We performed 7-Tesla blood oxygenation level dependent (BOLD) fMRI on Sprague–Dawley rats receiving simultaneous electrical stimulation at the left cervical vagus nerve. Using independent component analysis, we mapped the patterns of network activations and interactions in contrast to those in the resting state.

Results: Left cervical VNS evoked widespread responses over ~76% of the brain volume with the most notable activations in the brainstem and hypothalamus. The VNS evoked complex and distinctive temporal responses at different brain networks and altered the patterns of interactions among networks, especially for the limbic system.

Conclusion: This paper reports a method for exploring the complex response of VNS in a data-driven manner. The results suggest that VNS evokes wide-spread BOLD responses and alters large-scale functional connectivity.

Keywords: vagal nerve stimulation, functional magnetic resonance imaging, model-free analysis**,** brain-wide response

Abbreviations: blood oxygenation level dependent (BOLD), central neural system (CNS), functional

magnetic resonance imaging (fMRI), general linear model (GLM), hemodynamic response function

(HRF), independent components (ICs), independent component analysis (ICA), vagus nerve stimulation

(VNS)

Acknowledgements: Thank Dr. Nanyin Zhang for sharing the functional atlas of the rat brain. Thank

Elizabeth Baronowsky and Dr. Matthew Ward for their assistance in the initial training in animal surgery.

Funding: This work was supported by the National Institutes of Health [grant numbers, 1OT2TR001965-

01, R01MH104402].

Introduction

Since the 1800s (1, 2), vagus nerve stimulation (VNS) has been studied as a potential way to treat various diseases, including epilepsy, depression, tinnitus, Alzheimer's Disease, and obesity (3-8). The therapeutic benefits apparently depend on the effects of VNS on the central neural system (CNS) mediated through neuroelectrical or neurochemical signaling (9). Studies have been conducted to evaluate the CNS responses to VNS with various techniques. For example, invasive recordings of unit activity or field potentials have shown VNS-evoked neuronal responses in the nucleus of solitary tract (10), the locus coeruleus (11), and the hippocampus (12). These techniques offer high neuronal specificity but only cover spatially confined targets. In contrast, electroencephalogram (EEG) has been used to reveal VNS-induced synchronization or desynchronization of neural oscillations in the macroscopic scale (13, 14), while being severely limited by its spatial resolution and specificity as well as its inability to detect activity from deep brain structures. However, sub-cortical regions are of interest for VNS, because the vagus nerves convey signals to the brain through polysynaptic neural pathways by projecting to the brainstem, then subcortical areas, and lastly the cortex (9, 15).

Complementary to conventional electrophysiological approaches, functional neuroimaging allows characterizing the effects of VNS throughout the brain volume. Using positron emission tomography (PET) or single-photon emission computerized tomography (SPECT), prior studies have reported VNS evoked responses in the thalamus, hippocampus, amygdala, inferior cerebellum, and cingulate cortex (16-19); but these techniques are unable to capture the dynamics of responses due to their poor temporal resolution. In this regard, functional magnetic resonance imaging (fMRI) is more favorable because it offers balanced and higher spatial and temporal resolution. Previous human VNS-fMRI studies have reported VNS-evoked blood oxygenation level dependent (BOLD) responses in the thalamus, hypothalamus, prefrontal cortex, amygdala, and hippocampal formation (20-24). However, the reported activation patterns vary notably across existing studies (25), highlighting different regions responding to the VNS or the same regions with the opposite responses (20, 22). The variation may implicate the diversity of VNS' therapeutic effects in

different clinical subjcts but could also be attributable to the different analysis protocols (25). Therefore,

before moving forward and evaluating VNS' therapeutic effects using fMRI, it is necessary to verify

reliability of the current analysis method or develop a more reliable method for analyzing VNS-evoked

BOLD responses.

Functional MRI not only localizes the CNS responses of VNS (25), it also reveals the patterns and

dynamics of functional networks during VNS. Characterizing the network-level effect of VNS has been

central to understanding the mechanisms of VNS-based therapetutics. For example, given a lesion in the

locus coruleus, VNS fails to suppress epilepsy (26). Given blockade of the muscarinic receptor, VNS fails

to promote perceptual learning (27). These findings support the notion that VNS alters large-scale

functional networks in part through the adrenergic and cholinergic systems. However, how VNS affects the

patterns of temporal interactions among regions or networks (or functional connectivity) has rarely been

addressed (28), while fMRI has become the primary tool for studying functional connectivity in the resting

state (29, 30).

In this study, we aimed to address the BOLD effects of VNS in rat brains. Compared to human

studies, the use of animal models allowed us to minimize the inter-subject variation in genetics, gender,

age, weight, and health conditions. In such well-controlled conditions, the effects of VNS were mapped and

characterized in the whole brain by using BOLD-fMRI at 7-T. Our first goal was to verify the analysis

method as one of the sources of variation in VNS-fMRI studies and to apply a data-driven activation

mapping method, namely independent component analysis (ICA), to avoid the potential limitations or bias.

When studying the VNS effects, the primary effects of interest include not only the locations and dynamics

of VNS-evoked activations but also the changes in large-scale functional connectivity during VNS relative

to the resting state .

Methods and Materials

Subjects

A total of 17 Sprague–Dawley rats (male, weight: 250-350g; Envigo RMS, Indianapolis, IN) were studied according to a protocol approved by the Purdue Animal Care and Use Committee (PACUC) and the Laboratory Animal Program (LAP). Of the 17 animals, seven animals were used for VNS-fMRI experiments; ten animals were used for resting state fMRI. All animals were housed in a strictly controlled environment (temperature 21±1°C and 12 h light-dark cycle, lights on at 6:00 AM, lights off at 6:00 PM).

Animal preparation

For the VNS-fMRI experiments, each animal was initially anesthetized with 5% isoflurane and maintained with continuous administration of 2-3% isoflurane mixed with oxygen and a bolus of Rimadyl (5 mg/Kg, Zoetis) administrated subcutaneously. After a toe-pinch check for adequate anesthesia, a 2-3cm midline incision was made starting at the jawline and moving caudally. The left cervical vagus nerve was exposed and isolated by removing the surrounding tissue. A bipolar cuff electrode (Microprobes, made of platinum, with 1mm distance between contacts) was then wrapped around the exposed vagus nerve. For resting-state fMRI experiments, animals were anesthetized with the same dose of isoflurane without the aforementioned surgical procedures.

Once animals were moved to the MRI scanner, their heads were constrained with customized head restrainer. Then a bolus of dexdomitor (7.5 µg/Kg, Zoetis) was administrated subcutaneously. At about 15-20 mins following the bolus injection, dexdomitor was subcutaneously infused at 15 µg/Kg/h and increase every hour as needed (31). Isoflurane was reduced to be 0.1-0.5% mixed with oxygen. Both the dexdomitor infusion rate and the isoflurane dose were adjusted to maintain the respiration between 40 to 70 times per min and the heart rate between 250 to 350 beats per minute.

Vagus nerve stimulation

The bipolar cuff electrode was connected to a current stimulator (model 2200, A-M system) placed

outside of the MRI room through a 3-m twisted-pair of copper wire. Stimulation current was delivered in

10s-ON-50s-OFF cycles. When it was ON, biphasic square pulses (pulse width: 0.1 ms; amplitude: 1.0 mA;

frequency: 10 Hz) were delivered. For a total of 10 mins, ten cycles of stimulation were delivered in each

imaging session. Resting (stimulation-free) at least one minute long was given between sessions. Up to 4

sessions were scanned for each animal. Fig. 1 illustrates the VNS paradigm.

MRI and fMRI

MRI data were acquired with a 7-T small-animal MRI system (BioSpec 70/30, Bruker) equipped

with a volume transmitter coil (86 mm inner diameter) and 4-channel surface receiver coil. After localizer

scans, T2-weighted anatomical images were acquired with a rapid acquisition with relaxation enhancement

(RARE) sequence (repetition time (TR)=5804.607s, effective echo time (TE)=32.5ms, echo spacing=10.83

ms, voxel size=0.125×0.125×0.5mm^3, RARE factor=8, flip angle (FA)=90°). The BOLD fMRI data were

acquired by using a 2-D single-shot gradient echo echo-planar imaging (EPI) sequence (TR=1 s, TE=15

ms, FA=55°, in-plane resolution about 0.6×0.6 mm^2, slice thickness=1 mm). In the VNS group, the VNS

was delivered with the aforementioned block-design paradigm during continuous EPI.

Throughout the imaging experiment, the animal's body temperature was maintained at 37 ± 0.5 °C

using an animal-heating system. The oxygen saturation level (SpO$_2$) was maintained above 96%. Heart rate

and respiration rate were monitored with a small-animal physiological monitoring system (Kent Scientific).

Data preprocessing

MRI and fMRI data were preprocessed by using Analysis of Functional Neuroimages (AFNI) and

custom-built functions in MATLAB. Within each session, the fMRI data were corrected for motion, by

registering every volume to the first volume using *3dvolreg*. After removing the first ten volumes, *retroicor*

was used to correct for the motion artifacts due to respiratory and cardiac activity (32, 33). Then, *slicetimer*

was used to correct the timing for each slice. For each animal, we first normalized the EPI image to its T_2

weighted structural images and then registered to the template (34) using *flirt*. Motion artifact was further

removed by regressing out the six motion-correction parameters. The fMRI data were then spatially

smoothed with a 3-D Gaussian kernel with 0.5-mm full width at half maximum (FWHM). The fMRI time

series were detrended by regressing out a 2^{rd}-order polynomial function voxel by voxel.

General linear model analysis

We explored using the conventional general linear model (GLM) analysis to map the activations

evoked by VNS as in previous studies (23, 24, 35-37). Specifically, we derived a response model by

convolving the stimulation block (modeled as a box-car function) with a canonical hemodynamic response

function (HRF) (modeled as a gamma function). For each session, the fMRI signal at every voxel was

correlated with this response model. The correlation coefficient was converted to a z-score by using the

Fisher's z-transform. The voxel-wise z-score was averaged across sessions and evaluated for statistical

significance with one-sample t-test ($p < 0.05$, uncorrected).

This analysis revealed the group-level brain activation with VNS given an assumed response model.

Since the validity of this response model for VNS was not established, we intentionally varied the response

model by assuming three different peak delays (3s, 6s, or 9s) in the HRF. We compared the activation maps

obtained with the three different response models, to assess the dependence of the resulting activation map

on the presumed response characteristics.

Independent component analysis

In contrast to the voxel-wise general linear model (GLM) analysis, we used ICA to map networks

and their responses to VNS in a data-driven or model-free manner. For each session and each voxel, the

fMRI signal during VNS was demeaned and divided by its standard deviation. The resulting fMRI data

were then concatenated across all VNS-fMRI sessions. Infomax ICA (38) was used to decompose the

concatenated data into 20 spatially independent components (ICs). Each of these ICs included a spatial map

and a time series, representing a brain network and its temporal dynamics, respectively. In the spatial maps

of individual ICs, the intensities at each given voxel represented the weights by which the time series of

corresponding ICs were combined to explain this voxel's fMRI time series. Such weights were converted

to Z-statistics and then thresholded as described in a previous paper (39). The threshold was selected such

that the false negative rate was three times as large as the false positive rate. To obtain the false negative

and positive rates, the Z-statistics of all voxels in an IC map were modeled as a two-Gaussian mixture

distributions: one representing the noise, the other representing the signal.

Following ICA, we evaluated the VNS-evoked response separately for each IC, instead of each

voxel. To do so, each IC's time series was segmented for every VNS block. Each segment lasted 54 seconds,

starting from 3 seconds before the onset of a VNS block to 41 seconds after the offset of this block, while

the block size was 10 seconds. To address whether an IC responded to the VNS, we treated each time point

as a random variable and each segment as an independent sample. One-way analysis of variance (ANOVA)

test was conducted against a null hypothesis that there was no difference among all the time points (meaning

no response). The ICs that were statistically significant (p<5e-6) were considered as activated by VNS.

For each activated IC, we further characterized its temporal response to VNS. Briefly, we identified

the time points during or after the VNS block, where the signals significantly differed from the pre-stimulus

baseline by using Tukey's honest significant difference (HSD) test as post-hoc analysis following the

previous ANOVA test. Following this statistical test, the VNS-activated ICs were further classified into

three types (i.e. positive, negative, and mixed) of responses by visual inspection.

Functional connectivity analysis

We further addressed whether VNS altered the patterns of temporal interactions among functional

networks identified by ICA. To do so, the voxel time series was demeaned and standardized for each fMRI

session including both VNS and resting conditions. The fMRI data were concatenated across all sessions

and were then decomposed by ICA to yield 20 spatially ICs or networks, along with their corresponding

time series. The first IC was removed because it was identified as the global component. The time series of

the rest ICs were divided into the signals corresponding to the VNS sessions, and those corresponding to

all resting-state sessions.

We defined the functional connectivity between networks as the temporal correlations between the

ICs. The correlations were evaluated separately for the resting and VNS conditions and all pairs of ICs.

Based on their temporal correlations, we grouped the ICs into clusters by applying k-means clustering

method (with squared Euclidean distance) to ICs' temporal correlation matrix. As a result, the correlations

tended to be stronger within clusters than between clusters.

We further evaluated the differences in functional connectivity between the VNS condition and the

resting state. For this purpose, the functional connectivity between ICs was evaluated for each VNS session,

as well as each resting-state session. Their differences between these two conditions were evaluated using

unpaired two-sample t-test (p<0.05, uncorrected). The functional connectivity was plotted in the functional

connectogram.

Results

Model-based VNS activations were sensitive to variation of the response model

Seven rats were scanned for fMRI while their left cervical vagus nerve was electrically stimulated

in a (10s-ON-50s-OFF) block-design paradigm as illustrated in Fig. 1. The BOLD response phase-locked

to VNS block appeared complex and variable across regions of interest (ROIs). For example, the BOLD

responses were notably different across three ROIs, namely the retrosplenial cortex (RSC), brainstem (BS),

and dorsal caudate putamen (Cpu) in a functional atlas (40) of the rat brain (Fig. 2a). These responses were

not readily explainable by a typical response model derived from a canonical HRF (Fig. 2b, top). The GLM

analysis with three different response models (by varying the HRF peak latency with 3s increments) yielded

almost entirely distinctive activation maps (Fig. 2b), each of which was only marginally significant ($p<0.05$,

uncorrected). Therefore, VNS-evoked BOLD responses were too complex and variable to be captured by a

single response model. As such, the conventional GLM analysis likely leads to incomplete and inconsistent

activations with VNS, which possibly accounts for the diverging findings reported in the related literature

(20-24).

VNS induced widespread and complex network responses

We evaluated the VNS-evoked responses using a data-driven analysis method, and highlight the

responses at the level of networks, where the networks were identified as spatially ICs. It turned out that 15

out of the 20 ICs were significantly activated by VNS (one-way ANOVA, p<5e-6, Fig. 3a). Those activated

ICs collectively covered 76.03% of the brain volume (Fig. 3b). Among the activated regions, the brainstem

and the hypothalamus exhibited stronger responses than other areas.

The response time courses were also notably different across ICs. As shown in Fig. 3.a through 2.c.,

the time points, which were different from the baseline, are highlighted in red (p<0.05, Tukey's HSD). The

polarity of the red marked time points further characterized each IC as positive, negative, or biphasic-mixed

response. The negative response group included the amygdala, dorsal striatum, primary motor cortex,

midbrain, left somatosensory cortex, and superior cerebellum. The positive response group mainly included

the brainstem, thalamus, and hypothalamus region. The mixed response group included the hippocampal

formation, cingulate cortex, and prelimbic & infralimbic cortex. The ICs sharing similar responding polarity

were presumably more functionally associated with one another. Besides polarity characteristics, responses'

timing differences could also group ICs into long-lasting and stimulus-locked responses. Long-lasting

responses happened in the hippocampal formation, prefrontal cortex, amygdala. Stimulus-locked responses

showed in the brainstem and hypothalamus. The ICs with long-lasting responses were mostly involved in

high-level functions, such as forming memory, making decisions, and modulating emotion. The ICs with

time-locked responses had the more direct vagal projections involving fewer synapses. Considering both

polarity and timing differences, VNS responses varied across regions and networks and could reflect how

different regions changed their communications with each other. Therefore, studying the time courses

across different ICs could help us understand the changes of region-regions (or IC-IC) interaction altered

by VNS.

VNS altered functional connectivity

We evaluated the IC-IC interactions during VNS and compared with those in the resting state

(control group). Three clusters were obtained with k-means clustering separately for the VNS and resting-

state groups (Fig. 4.a). Cluster 1 was similar in both the VNS and control groups, and represented the

sensorimotor cortex. In contrast, cluster 2 and 3 were notably different in the two groups. For cluster 2,

VNS strengthened the correlations between the hippocampal formation and such regions as the retrosplenial

cortex relative to the resting state. For cluster 3, VNS reduced the correlations between the prefrontal cortex

and such regions as the basal ganglia. We further compared the functional connectivity of each IC-IC pair

to better illustrate the changes in temporal correlation across different regions (ICs) ($P<0.005$, t-test). Fig

4.b shows that VNS manipulated the degree of interactions among different regions in the limbic system.

The cingulate cortex was less correlated with the ventral striatum. The left and right hippocampal formation

formed stronger functional connectivity, and as a whole hippocampal formation formed stronger functional

connectivity with the retrosplenial cortex During VNS, the functional connectivity was not only reorganized

within the limbic system but also between the limbic system and the sensorimotor cortex. VNS strengthened

the interaction across the sensorimotor cortex with the hippocampal formation, retrosplenial cortex, and

dorsal striatum, but weakened the functional connectivity between the sensorimotor cortex and the cingulate

cortex. In summary, VNS reorganized the functional connectivity within the limbic system and altered the

interactions between the limbic system and the sensorimotor cortex.

Discussion

This paper studied VNS induced BOLD responses on seven healthy rats with the same stimulating

protocol. Using data-driven analysis method, our results showed that VNS evoked brain-wide BOLD

responses covering ~76% of the brain regions, but the distinct responding time courses varied across regions

and were different from the canonical HRF, which was commonly used in the previous VNS-fMRI studies.

In addition, VNS also altered the functional networks compared to those observed in the resting state. In

particular, VNS altered functional connectivity within the limbic system and between the limbic system

and the sensorimotor cortex. Such changes in functional connectivity may underlie the differential functions

of VNS in modulating emotion and sensorimotor plasticity.

VNS evoke brain-wide responses

A major finding in this study was that VNS evoked time-locked and widespread BOLD responses

over most parts of the brain. This finding appeared surprising at the first glance, since simulation was

applied to the left cervical vagus – a seemingly narrowly-focused entry of neuromodulation. Nevertheless,

previous studies suggest that neural activity may drive global fluctuations in resting-state fMRI activity

(41), and simple (e.g., checkerboard) visual stimulation may evoke whole-brain fMRI responses (42).

Common to those prior studies and this study may be the notion that the brain is so densely wired and

interconnected that focal modulation may induce a cascade of responses through neuronal circuits. Such

network responses may even have a global reach, if the stimulation innervates sub-cortical structures with

distributed modulatory effects on the brain (43).

In this regard, widespread responses to VNS may be mediated through diffusive neuromodulation

triggered by VNS. Vagal afferents project to the parabrachial nucleus, locus coeruleus, raphe nuclei through

nucleus of solitary tract (44). From the parabrachial nucleus, locus coeruleus, and raphe nuclei, connectivity

extends onto hypothalamus, thalamus, amygdala, anterior insular, infralimbic cortex, and lateral prefrontal

cortex (45-48). In fact, the widespread VNS-evoked activations reported herein are consistent with the full

picture gathered from piecemeal activations observed in prior VNS-fMRI (see reviews in (25)),

transcutaneous VNS-fMRI studies (49), and EEG-fMRI studies (13, 14). In light of those results, the extent

of the VNS effects on the brain has been under-estimated in prior studies, likely due to the use of simplified

response models that fail to capture the complex and variable responses across all activated regions.

Origins and interpretation of different response characteristics

Results in this study suggest that VNS induces a variety of BOLD responses that vary across regions.

In addition to coarse and qualitative classification of various responses as positive, negative, or mixed (first

negative and then positive) (Fig. 3), the responses at various regions or networks also differed in terms of

transient vs. sustained behaviors during and following VNS. For example, the responses at the brainstem

and the hypothalamus showed a very rapid rise around 2s and rapid decay around 5s following the onset of

VNS. Although the generalizable origins of transient BOLD responses are still debatable (50-52), we

interpret the transient responses to VNS as a result of direct neuroelectric signaling through the vagal nerves.

Nuclei in the brainstem, e.g., NTS, contain neurons receiving direct projections from the vagus, and in turn

connect to the hypothalamus. Such brain structures are thus well-positioned to respond rapidly to VNS.

Also, the rapid decay of the BOLD response in the brainstem and hypothalamus may indicate neuronal

adaption – a factor of consideration for designing the duration and duty cycle of VNS. However, such

interpretation should be taken with caution. The neurovascular coupling (modeled as the HRF) behaves as

a low-pass filter through which the BOLD response is generated from local neuronal responses. Although

the peak latency in HRF is 4 to 6s in humans, it is as short as 2s in rodents (53), making it relatively more

suitable for tracking transient neuronal dynamics.

Another intriguing observation was the prolonged BOLD responses that sustained for a long period

following the offset of VNS. In the striatum, hippocampus, as well as the prelimbic and infralimbic cortex,

the VNS-evoked response lasted for 40s or even longer, while the VNS only lasted 10s (Fig. 3). Such

prolonged responses suggest potentially long-lasting effects of VNS. This observation is also in line with

clinical studies showing that the effects of VNS on seizure suppression are not limited to the stimulation,

but sustain during periods in the absence of VNS (54). Moreover, those regions showing prolonged effects

of VNS tended to be higher-level functional areas presumably involved in learning, decision-making,

memory, and emotion-processing. Speculatively, it implies that the VNS-based modulation of cognitive

functions or dysfunctions operates in a relatively longer time scale while imposing potentially therapeutic

effects on a longer term.

VNS alters network-network interactions

This study highlights the importance of evaluating the effects of VNS on functional connectivity,

which measures the degree to which regions or networks interact with each other. It is widely recognized

that brain functions emerge from coordination among regions (55). However, prior studies address the

effects of VNS in focal and target regions (56), whereas the effects of VNS on functional connectivity is

perhaps more functionally relevant. In line with this perspective, a recent study has shown that

transcutaneous VNS modulates the functional connectivity in the default mode network in patients with

major depressive disorders, and the change in functional connectivity is related to the therapeutic efficacy

across individual patients [50]. Thus, VNS may reorganize the patterns of interactions among functional

networks – a plausible network basis underlying VNS-based therapeutics.

Our results show that VNS reorganizes the functional connectivity with respect to the limbic system.

Relative to the intrinsic functional connectivity in the resting state, VNS increases functional connectivity

between the retrosplenial cortex and hippocampal formation, of which the functional roles are presumably

related to memory, learning, and monitoring sensory inputs (57, 58). In addition, VNS increases functional

connectivity between the sensory cortex and the striatum, of which the functional roles are presumably the

integration of sensorimotor, cognitive, and motivation/emotion (59). In contrast, VNS decreases functional

connectivity between the cingulate cortex and the ventral striatum, which likely affect the emotional control

of visceral, skeletal, and endocrine outflow (60). Collectively, these observations lead us to speculate that

VNS biases the limbic system to shift its functional role from emotional processing to perceptual learning.

Such speculation is consistent with the therapeutic effects of VNS in depression patients (61, 62) and the

cortical plasticity of interest to perceptual learning and motor rehabilitation induced by VNS (63, 64).

Model-free activation mapping in the level of networks

Central to this study is the use of model-free and data-driven analysis for mapping activations in

the level of networks, instead of voxels or regions. This is in contrast to conventional GLM analysis used

in previous VNS-fMRI studies, which assumes that neural responses sustain in the entire period of VNS,

and the BOLD effects of neural responses may be modeled with a canonical HRF (23, 24, 35-37). Both of

these assumptions may not be entirely valid. Neural responses may exhibit a range of non-linear

characteristics. The typical HRF model is mostly based on data or findings obtained from the cortex during

sensory stimulation (65), whereas no study has modeled the HRF for VNS. Moreover, the neurovascular

coupling may also vary across regions in the brain, especially between sub-cortical and cortical areas due

to their differences in local vasculatures (66). Thus, the GLM analysis with a single and empirical response

model most likely falls short for explaining the complex and variable responses across all brain regions.

The model-free analysis allowed us to detect the VNS-evoked brain response in a data-driven way

without assuming any prior response model. A similar strategy has been used to test for voxel-wise BOLD

responses to visual stimulation in humans (42). What was perhaps unique in this study was the use of the

model-free analysis on the activity of spatially independent components, rather than that of single voxels.

Each IC contained a set of voxels (or locations) that shared a common pattern of temporal dynamics. ICA

utilized the fact that individual voxels were organized by networks, not in isolation, to extract the network

activity as the time series of each IC, which reflected the (weighted) average activity of all the voxels that

belonged to each IC. As such, the signal-to-noise ratio was higher for IC-wise activity than for voxel-wise

activity, providing better sensitivity for detecting activations at the network level. This model-free analysis

method is thus arguably more favorable than conventional GLM analysis, especially when the response

characteristics are complex and unclear, e.g., given VNS.

Potentially confounding cardiac and respiratory effects

The BOLD signal is an indirect measure of neural activity. Therefore, it may be affected by non-neuronal physiological fluctuations (67). Previous studies have shown that VNS causes cardiorespiratory effects, e.g., variations in the heart rate, respiration rate, and SpO2 (68, 69). Such effects may potentially confound the interpretation of the VNS-induced BOLD responses regarding neuronal activations. Such confounding effects were highlighted in a prior study, in which VNS was found to decrease the heart rate and in turn decrease the BOLD signal throughout the rat brain (70). In this study, we were concerned about this potential confounding effect, and reduced to the pulse width of VNS to 0.1ms, instead of 0.5ms in that study (70). Such shortened pulse width largely mitigated the cardiac effects, as no obvious changes in heart rate were noticeable during experiments.

Also, the cardiac effects would manifest themselves as the common physiological response observable throughout the brain. This was not the case in this study. Despite wide-spread activations with VNS, the responses at individual regions exhibited different temporal characteristics, which could not be readily explained by a common confounding source (e.g., respiratory or cardiac). Instead, the regions with similar response patterns formed well-patterned functional networks, resembling intrinsic resting-state networks previously observed in rats (71). For these reasons, it was unlikely that the VNS-induced activation and functional-connectivity patterns were the result at non-neuronal cardiac or respiratory effects. Similar justifications are also applicable to the confounding respiratory effects. Nevertheless, future studies are desirable to fully disentangle the neuronal vs. non-neuronal effects of VNS. Of particular interest is using multi-echo fMRI to differentiate BOLD vs. non-BOLD effects (67), and combining electrophysiology and fMRI to pinpoint the neural origin of the fMRI response to VNS (72).

References

1.   Gaskell WH. The Electrical changes in the Quiescent Cardiac Muscle which accompany Stimulation of the Vagus Nerve. J Physiol 1886;7(5-6):451-2.

2.   Robinson C, Draper G. Studies with the electrocardiograph on the action of the vagus nerve on the human heart : II. the effects of vagus stimulation on the hearts of children with chronic valvular disease. J Exp Med 1912;15(1):14-48.

3.   Ben-Menachem E. Vagus-nerve stimulation for the treatment of epilepsy. Lancet Neurol 2002;1(8):477-82.

4.   George MS, Sackeim HA, Rush AJ, Marangell LB, Nahas Z, Husain MM, et al. Vagus nerve stimulation: a new tool for brain research and therapy. Biol Psychiatry 2000;47(4):287-95.

5.   De Ridder D, Vanneste S, Engineer ND, Kilgard MP. Safety and efficacy of vagus nerve stimulation paired with tones for the treatment of tinnitus: a case series. Neuromodulation 2014;17(2):170-9.

6.   Peña DF, Engineer ND, McIntyre CK. Rapid remission of conditioned fear expression with extinction training paired with vagus nerve stimulation. Biol Psychiatry 2013;73(11):1071-7.

7.   Sjögren MJ, Hellström PT, Jonsson MA, Runnerstam M, Silander HC, Ben-Menachem E. Cognition-enhancing effect of vagus nerve stimulation in patients with Alzheimer's disease: a pilot study. J Clin Psychiatry 2002;63(11):972-80.

8.   Bodenlos JS, Kose S, Borckardt JJ, Nahas Z, Shaw D, O'Neil PM, et al. Vagus nerve stimulation acutely alters food craving in adults with depression. Appetite 2007;48(2):145-53.

9.   Henry TR. Therapeutic mechanisms of vagus nerve stimulation. Neurology 2002;59(6 suppl 4):S3-S14.

10.   Beaumont E, Campbell RBP, Andresen MC, Scofield SL, Singh K, Libbus I, et al. Clinically-styled vagus nerve stimulation augments spontaneous discharge in second and higher order sensory neurons in rat nucleus of the solitary tract. American Journal of Physiology-Heart and Circulatory Physiology 2017:ajpheart. 00070.2017.

11.   Groves DA, Bowman EM, Brown VJ. Recordings from the rat locus coeruleus during acute vagal nerve stimulation in the anaesthetised rat. Neuroscience letters 2005;379(3):174-9.

12.   Larsen LE, Wadman WJ, Marinazzo D, van Mierlo P, Delbeke J, Daelemans S, et al. Vagus nerve stimulation applied with a rapid cycle has more profound influence on hippocampal electrophysiology than a standard cycle. Neurotherapeutics 2016;13(3):592-602.

13.   Chase MH, Nakamura Y, Clemente CD, Sterman MB. Afferent vagal stimulation: neurographic correlates of induced EEG synchronization and desynchronization. Brain Res 1967;5(2):236-49.

14.   Bartolomei F, Bonini F, Vidal E, Trébuchon A, Lagarde S, Lambert I, et al. How does vagal nerve stimulation (VNS) change EEG brain functional connectivity? Epilepsy Res 2016;126:141-6.

15.   Ressler KJ, Mayberg HS. Targeting abnormal neural circuits in mood and anxiety disorders: from the laboratory to the clinic. Nature neuroscience 2007;10(9):1116.

16.   Garnett ES, Nahmias C, Scheffel A, Firnau G, Upton AR. Regional cerebral blood flow in man manipulated by direct vagal stimulation. Pacing Clin Electrophysiol 1992;15(10 Pt 2):1579-80.

17.   Ko D, Heck C, Grafton S, Apuzzo ML, Couldwell WT, Chen T, et al. Vagus nerve stimulation activates central nervous system structures in epileptic patients during PET H2(15)O blood flow imaging. Neurosurgery 1996;39(2):426-30; discussion 30-1.

18.   Henry TR, Votaw JR, Pennell PB, Epstein CM, Bakay RA, Faber TL, et al. Acute blood flow changes and efficacy of vagus nerve stimulation in partial epilepsy. Neurology 1999;52(6):1166-73.

19. Zobel A, Joe A, Freymann N, Clusmann H, Schramm J, Reinhardt M, et al. Changes in regional
cerebral blood flow by therapeutic vagus nerve stimulation in depression: an exploratory approach.
Psychiatry Res 2005;139(3):165-79.
20. Devous M, Husain M, Harris T, Rush A. Effects of VNS on regional cerebral blood flow in
depressed subjects. European Psychiatry 2002;17:113-4.
21. Sucholeiki R, Alsaadi TM, Morris Iii GL, Ulmer JL, Biswal B, Mueller WM. fMRI in patients
implanted with a vagal nerve stimulator. Seizure 2002;11(3):157-62.
22. Liu W-C, Mosier K, Kalnin A, Marks D. Vagus Nerve Stimulation in patients: a BOLD fMRI study.
NeuroImage 2001;13(6):810.
23. Bohning DE, Lomarev MP, Denslow S, Nahas Z, Shastri A, George MS. Feasibility of vagus nerve
stimulation–synchronized blood oxygenation level–dependent functional MRI. Investigative Radiology
2001;36(8):470-9.
24. Lomarev M, Denslow S, Nahas Z, Chae JH, George MS, Bohning DE. Vagus nerve stimulation
(VNS) synchronized BOLD fMRI suggests that VNS in depressed adults has frequency/dose dependent
effects. J Psychiatr Res 2002;36(4):219-27.
25. Chae JH, Nahas Z, Lomarev M, Denslow S, Lorberbaum JP, Bohning DE, et al. A review of
functional neuroimaging studies of vagus nerve stimulation (VNS). J Psychiatr Res 2003;37(6):443-55.
26. Krahl SE, Clark KB, Smith DC, Browning RA. Locus coeruleus lesions suppress the seizure-
attenuating effects of vagus nerve stimulation. Epilepsia 1998;39(7):709-14.
27. Nichols JA, Nichols AR, Smirnakis SM, Engineer ND, Kilgard MP, Atzori M. Vagus nerve
stimulation modulates cortical synchrony and excitability through the activation of muscarinic receptors.
Neuroscience 2011;189:207-14.
28. Fang J, Rong P, Hong Y, Fan Y, Liu J, Wang H, et al. Transcutaneous vagus nerve stimulation
modulates default mode network in major depressive disorder. Biological psychiatry 2016;79(4):266-73.
29. Biswal B, Zerrin Yetkin F, Haughton VM, Hyde JS. Functional connectivity in the motor cortex of
resting human brain using echo‐planar mri. Magnetic resonance in medicine 1995;34(4):537-41.
30. Van Den Heuvel MP, Pol HEH. Exploring the brain network: a review on resting-state fMRI
functional connectivity. European neuropsychopharmacology 2010;20(8):519-34.
31. Pawela CP, Biswal BB, Hudetz AG, Schulte ML, Li R, Jones SR, et al. A protocol for use of
medetomidine anesthesia in rats for extended studies using task-induced BOLD contrast and resting-
state functional connectivity. Neuroimage 2009;46(4):1137-47.
32. Glover GH, Li TQ, Ress D. Image-based method for retrospective correction of physiological
motion effects in fMRI: RETROICOR. Magn Reson Med 2000;44(1):162-7.
33. Birn RM, Diamond JB, Smith MA, Bandettini PA. Separating respiratory-variation-related
fluctuations from neuronal-activity-related fluctuations in fMRI. Neuroimage 2006;31(4):1536-48.
34. Valdés-Hernández PA, Sumiyoshi A, Nonaka H, Haga R, Aubert-Vásquez E, Ogawa T, et al. An in
vivo MRI template set for morphometry, tissue segmentation, and fMRI localization in rats. Frontiers in
neuroinformatics 2011;5.
35. Nahas Z, Teneback C, Jeong-Ho C, Mu Q, Molnar C, Kozel FA, et al. Serial vagus nerve stimulation
functional MRI in treatment-resistant depression. Neuropsychopharmacology 2007;32(8):1649.
36. Kraus T, Kiess O, Hösl K, Terekhin P, Kornhuber J, Forster C. CNS BOLD fMRI effects of sham-
controlled transcutaneous electrical nerve stimulation in the left outer auditory canal–a pilot study.
Brain stimulation 2013;6(5):798-804.
37. Kraus T, Hösl K, Kiess O, Schanze A, Kornhuber J, Forster C. BOLD fMRI deactivation of limbic and
temporal brain structures and mood enhancing effect by transcutaneous vagus nerve stimulation. J
Neural Transm. 2007;114(11):1485-93.
38. Bell AJ, Sejnowski TJ. An information-maximization approach to blind separation and blind
deconvolution. Neural computation 1995;7(6):1129-59.

39.   Beckmann CF, Smith SM. Probabilistic independent component analysis for functional magnetic
resonance imaging. IEEE transactions on medical imaging 2004;23(2):137-52.
40.   Ma Z, Perez P, Liu Y, Hamilton C, Liang Z, Zhang N. Functional atlas of the awake rat brain: A
neuroimaging study of rat brain specialization and integration. Neuroimage 2016.
41.   Schölvinck ML, Maier A, Frank QY, Duyn JH, Leopold DA. Neural basis of global resting-state fMRI
activity. Proceedings of the National Academy of Sciences 2010;107(22):10238-43.
42.   Gonzalez-Castillo J, Saad ZS, Handwerker DA, Inati SJ, Brenowitz N, Bandettini PA. Whole-brain,
time-locked activation with simple tasks revealed using massive averaging and model-free analysis. Proc
Natl Acad Sci U S A 2012;109(14):5487-92.
43.   Lee JH, Durand R, Gradinaru V, Zhang F, Goshen I, Kim D-S, et al. Global and local fMRI signals
driven by neurons defined optogenetically by type and wiring. Nature 2010;465(7299):788.
44.   Paxinos G. The rat nervous system: Academic press; 2014.
45.   Van Bockstaele EJ, Peoples J, Valentino RJ. A.E. Bennett Research Award. Anatomic basis for
differential regulation of the rostrolateral peri-locus coeruleus region by limbic afferents. Biol Psychiatry
1999;46(10):1352-63.
46.   Dorr AE, Debonnel G. Effect of vagus nerve stimulation on serotonergic and noradrenergic
transmission. J Pharmacol Exp Ther 2006;318(2):890-8.
47.   Manta S, Dong J, Debonnel G, Blier P. Enhancement of the function of rat serotonin and
norepinephrine neurons by sustained vagus nerve stimulation. J Psychiatry Neurosci 2009;34(4):272-80.
48.   Ruffoli R, Giorgi FS, Pizzanelli C, Murri L, Paparelli A, Fornai F. The chemical neuroanatomy of
vagus nerve stimulation. J Chem Neuroanat 2011;42(4):288-96.
49.   Frangos E, Komisaruk BR. Access to vagal projections via cutaneous electrical stimulation of the
neck: fMRI evidence in healthy humans. Brain stimulation 2017;10(1):19-27.
50.   Renvall V, Hari R. Transients may occur in functional magnetic resonance imaging without
physiological basis. Proc Natl Acad Sci U S A 2009;106(48):20510-4.
51.   Tyler CW, Kontsevich LL, Ferree TC. Independent components in stimulus-related BOLD signals
and estimation of the underlying neural responses. Brain Res 2008;1229:72-89.
52.   Obata T, Liu TT, Miller KL, Luh WM, Wong EC, Frank LR, et al. Discrepancies between BOLD and
flow dynamics in primary and supplementary motor areas: application of the balloon model to the
interpretation of BOLD transients. Neuroimage 2004;21(1):144-53.
53.   Ma Y, Shaik MA, Kozberg MG, Kim SH, Portes JP, Timerman D, et al. Resting-state hemodynamics
are spatiotemporally coupled to synchronized and symmetric neural activity in excitatory neurons. Proc
Natl Acad Sci U S A 2016;113(52):E8463-E71.
54.   Zabara J. Inhibition of experimental seizures in canines by repetitive vagal stimulation. Epilepsia
1992;33(6):1005-12.
55.   Anastassiou CA, Shai AS. Psyche, Signals and Systems. Micro-, Meso-and Macro-Dynamics of the
Brain: Springer; 2016. p. 107-56.
56.   Tian P, Teng IC, May LD, Kurz R, Lu K, Scadeng M, et al. Cortical depth-specific microvascular
dilation underlies laminar differences in blood oxygenation level-dependent functional MRI signal.
Proceedings of the National Academy of Sciences 2010;107(34):15246-51.
57.   Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex.
Trends in cognitive sciences 2000;4(6):215-22.
58.   Vogt BA, Finch DM, Olson CR. Functional heterogeneity in cingulate cortex: the anterior
executive and posterior evaluative regions. Cerebral cortex 1992;2(6):435-43.
59.   Balleine BW, Delgado MR, Hikosaka O. The role of the dorsal striatum in reward and decision-
making. Journal of Neuroscience 2007;27(31):8161-5.

60.   Parkinson JA, Willoughby PJ, Robbins TW, Everitt BJ. Disconnection of the anterior cingulate
cortex and nucleus accumbens core impairs Pavlovian approach behavior: Further evidence for limbic
cortical–ventral striatopallidal systems. Behavioral neuroscience 2000;114(1):42.
61.   Sackeim HA, Rush AJ, George MS, Marangell LB, Husain MM, Nahas Z, et al. Vagus nerve
stimulation (VNS (TM)) for treatment-resistant depression: efficacy, side effects, and predictors of
outcome. Neuropsychopharmacology 2001;25(5):713.
62.   Nemeroff CB, Mayberg HS, Krahl SE, McNamara J, Frazer A, Henry TR, et al. VNS therapy in
treatment-resistant depression: clinical evidence and putative neurobiological mechanisms.
Neuropsychopharmacology 2006;31(7):1345.
63.   Schwarz LA, Luo L. Organization of the locus coeruleus-norepinephrine system. Current Biology
2015;25(21):R1051-R6.
64.   Mesulam M, Mufson E, Wainer B, Levey A. Central cholinergic pathways in the rat: an overview
based on an alternative nomenclature (Ch1–Ch6). Neuroscience 1983;10(4):1185-201.
65.   Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg M, Turner R. Event-related fMRI: characterizing
differential responses. Neuroimage 1998;7(1):30-40.
66.   Li B, Freeman RD. Neurometabolic coupling in the lateral geniculate nucleus changes with
extended age. Journal of neurophysiology 2010;104(1):414-25.
67.   Kundu P, Brenowitz ND, Voon V, Worbe Y, Vértes PE, Inati SJ, et al. Integrated strategy for
improving functional connectivity mapping using multiecho fMRI. Proceedings of the National Academy
of Sciences 2013;110(40):16187-92.
68.   Zaaimi B, Grebe R, Wallois F. Animal model of the short-term cardiorespiratory effects of
intermittent vagus nerve stimulation. Auton Neurosci 2008;143(1-2):20-6.
69.   Murray BJ, Matheson JK, Scammell TE. Effects of vagus nerve stimulation on respiration during
sleep. Neurology 2001;57(8):1523-4.
70.   Reyt S, Picq C, Sinniger V, Clarençon D, Bonaz B, David O. Dynamic Causal Modelling and
physiological confounds: a functional MRI study of vagus nerve stimulation. Neuroimage 2010;
52(4):1456-64.
71.   Sierakowiak A, Monnot C, Aski SN, Uppman M, Li T-Q, Damberg P, et al. Default mode network,
motor network, dorsal and ventral basal ganglia networks in the rat brain: comparison to human
networks using resting state-fMRI. PloS one 2015;10(3):e0120345.
72.   Viswanathan A, Freeman RD. Neurometabolic coupling in cerebral cortex reflects synaptic more
than spiking activity. Nature neuroscience 2007;10(10):1308.

Figure 1. Experimental design. Each rat was electrically stimulated at the left cervical vagus through a

cuff electrode implanted in an acute surgery. The stimulation (0.1ms pulse width, 1.0mA amplitude,

10Hz) was delivered in a 10s-ON-50s-OFF cycle, while the rat was scanned for fMRI with 1s TR.

Figure 2. VNS-evoked responses varied across regions. (a) shows the response time series averaged

within each of the three color-coded regions of interest shown: the retrosplenial cortex (RSC) (blue), the

brainstem, and the dorsal caudate putamen (Cpu). (b) shows the highly different activation maps based on

the response models derived with the HRF, for which different peak latencies were assumed to be 3s, 6s,

or 9s.

Figure 3. VNS evoked widespread and complex responses in the brain. (**a**) VNS-evoked responses for

different brain networks derived with ICA. The ICA-defined networks are labeled: amygdala (Amy),

caudate putamen (Cpu), hippocampus, (Hipp), cingulate cortex (Cing), prelimbic cortex (PrL), infralimbic

cortex (IL), brain stem, hypothalamus (HTh), thalamus (Tha), superior colliculus (SC), cerebellum (Cb),

primary and secondary motor cortex (M1, M2), and primary and secondary somatosensory cortex (S1,

S2). For each network, the time points at which the responses were significant are shown in red. (**b**) The

VNS-activated voxels cover 76.03% of the brain volume. The color is in arbitrary units, representing the

standard deviation of the voxel-wise response averaged across repetitions of VNS. The locations with the

greatest responses are highlighted with arrows.

Figure 4. VNS altered the functional connectivity among intrinsic functional networks. (**a**) shows the

network-network correlations between independent components derived with ICA. The left shows the

correlation matrix during the resting state, or the "control" condition. The right shows the correlation

matrix during VNS, or the "VNS" condition. Smaller squares highlight the networks (or ICs) that were

clustered into groups (based on k-means clustering). (**b**) shows the IC-IC functional connectivity that was

significantly different between the VNS and control conditions (t-test, P<0.005). Red lines represent

increases in functional connectivity, and green lines represent decreases in functional. The thickness of

the lines represents the (VNS versus control) change in correlation. The brain maps show the spatial

patterns of individual ICs. Corresponding to the squares in (a), the arc lines illustrate how the ICs were

clusters into groups, for the VNS condition (inner circle) and the control condition (outer circle).