Whole genome sequence and

comparative analysis of *Borrelia burgdorferi* MM1

Neda Jabbari, Gustavo Glusman, Lena M. Joesch-Cohen, Panga Jaipal Reddy, Robert L. Moritz,

Leroy Hood, and Christopher G. Lausted*

Institute for Systems Biology, Seattle, Washington, United States of America

*Corresponding author:

E-mail: clausted@systemsbiology.org (CGL)

Abstract

Lyme disease is caused by spirochaetes of the *Borrelia burgdorferi* sensu lato genospecies. Complete genome assemblies are available for fewer than ten strains of *Borrelia burgdorferi* sensu stricto, the primary cause of Lyme disease in North America. MM1 is a sensu stricto strain originally isolated in the midwestern United States. Aside from a small number of genes, the complete genome sequence of this strain has not been reported. Here we present the complete genome sequence of MM1 in relation to other sensu stricto strains and in terms of its Multi Locus Sequence Typing. Our results indicate that MM1 is a new sequence type which contains a conserved main chromosome and 15 plasmids. Our results include the first contiguous 28.5 kb assembly of lp28-8, a linear plasmid carrying the *vls* antigenic variation system, from a *Borrelia burgdorferi* sensu stricto strain.

Introduction

Lyme disease is the most prevalent tick-borne disease in North America. Resulting from infections by spirochetes of the *Borrelia burgdorferi* sensu lato (s.l.) genospecies, the most frequent initial manifestation is erythema migrans, a characteristic expanding ring-shaped cutaneous lesion[1,2]. While some patients experience erythema migrans with or without flu-like symptoms, others may develop extracutaneous manifestations that can affect the nervous system (neuroborreliosis), the heart (Lyme carditis) or the joints (Lyme arthritis)[3–5]. What role pathogen genetic diversity plays in the producing varying symptoms is unknown. Currently, genomic sequence of forty seven strains of *Borrelia burgdorferi* sensu stricto (s.s.), the primary clinically relevant species in North America[6], are available from NIH GenBank. Of these, fewer than ten are described as complete assemblies. Given the complexity of Lyme manifestations and the scarcity of sequenced genomes, sequencing and characterization of more strains is vital for the understanding of *Borrelia* genetic diversity and disease manifestations.

The MM1 strain was isolated from the kidney of a white-footed mouse, *Peromyscus leucopus*, in Minnesota [7,8]. It was used in early studies of Lyme disease [7,9–12] and it appears in patent applications for the development of *B. burgdorferi* detection assays and recombinant vaccines against Lyme disease [13,14]. However, its whole genome sequence has not yet been reported. We present here the full genome sequence of MM1, and provide comparative analysis, Multi Locus Sequence Typing (MLST), and annotation of its genome. Most notably, we report on the presence of the lp28-8 plasmid which contains its *vls* antigenic variation system.

Materials and methods

Growth conditions and DNA preparation

B. burgdorferi MM1 spirochetes were purchased from ATCC (ATCC 51990), resuspended and maintained in laboratory-produced media that resembles MKP/BSK-II media with 10% heat-inactivated rabbit serum (R7136-60, Sigma-Aldrich, USA), based on recipes provided by Dr. K. Strle[15,16]. ATCC 51990 MM1was first suspended in 10 mL of MKP/BSK-II media, then 1 mL of the diluted cells was inoculated into 13 mL of MKP/BSK-II media in 15 mL culture tubes (Corning tubes, USA) and incubated at 34°C, 5.0% CO_2 with the lids of the tubes tightly sealed to minimize the oxygen level (microaerophilic condition). The seeded density of spirochetes was 5×10^6 cells/mL and cells were grown until they reached mid-log phase (i.e., spirochete count reached $3 – 5 \times 10^7$). MM1 spirochetes four passages away from ATCC lot were harvested and washed with PBS buffer (pH 7.4) four times to remove the media components and pelleted, ready for subsequent DNA processing and genome sequencing.

Total DNA was isolated from pelleted MM1 spirochetes using DNeasy blood and tissue kit (Qiagen, USA). RNase A (Qiagen) was used to digest the RNA content as recommended by the manufacturer's protocol. DNA was purified using the Agencourt AMPure XP method (Beckman Coulter, USA) and quantified with the Quant-iT PicoGreen dsDNA Assay Kit (ThermoFisher Scientific, USA) and a NanoDrop ND-1000 spectrophotometer. Integrity of the isolated DNA was checked on a 0.6 % agarose gel as well as with an Agilent DNA 12000 Kit (Agilent Technologies, Inc., USA).

Genome sequencing

DNA sequence data were obtained primarily using the Pacific Biosciences (PacBio) Single Molecule, Real-Time (SMRT) system with DNA/Polymerase Binding Kit P6 v2, MagBead Kit v1 and DNA Sequencing Reagent Kit 4.0 v2 at the PacBio Sequencing Services of the University of Washington (Seattle, WA, USA). Long DNA reads were collected in one SMRT cell on a PacBio RSII instrument. Additional long reads were obtained using the Oxford Nanopore Technologies (ONT) MinION system. The MinION genomic DNA library was prepared according to the ONT Nanopore Sequencing Kit SQK-MAP006 protocol and quantified with Quant-iT PicoGreen dsDNA Assay Kit (ThermoFisher Scientific, USA). Sequencing reads were generated on Flow Cells (Flo-MAP003) with MinKNOW 0.51.1.62 and base-called with Metrichor 2.39. "2D pass" reads were converted into FASTA format with poretools v. 0.5.1. These reads are the highest quality reads, selected from reads where both strands of a double-stranded fragment were analyzed together in one pore.

Genome assembly

De novo assembly of PacBio reads was performed with Hierarchical Genome Assembly Process (HGAP v2) using a minimum seed read length of 7800 bp. Contigs were merged or circularized with Circlator (v 1.5.0) [17]. In order to acquire the final consensus sequence, raw reads were filtered and mapped to the *B. burgdorferi* MM1 assembly with the resequencing protocol as implemented in SMRT Analysis Software v2.3.0.

Sequence characterization and visualisation

Tandem repeats were defined with Unipro UGENE v1.9.8 as follows: 1–9 bp, 10–100 bp
and >100 bp unit sizes were termed micro-, mini- and macro-satellite, respectively, and
overlapping tandem repeats were excluded [18]. BLAST alignments of the sequences along with
annotations were plotted using Easyfig (v 2.2.2) [19] and the Artemis Comparison Tool (ACT;
Release 13.0.0)[20].

B. burgdorferi MM1 plasmid identification and nomenclature

The plasmid content of MM1 was first evaluated visually with respect to the standard
type strain B31. Reads were mapped to the B31 genome (assembly GCA_000008685.2) with the
"resequencing" protocol included in PacBio's SMRT Analysis software v2.3.0 (S1 Fig). Next,
MM1 plasmid identities were confirmed by the sequence type of the *Borrelia* paralogous family
(PFam) 32 protein encoded by each plasmid as defined by Casjens *et al*. [21]. Specifically, the
protein sequences of MM1 and three fully-sequenced *B. burgdorferi* s.s. strains (B31, N40 and
JD1; assemblies GCA_000008685.2, GCA_000166635.2 and GCA_000166655.2, respectively)
were aligned with BLAST using an E-value cutoff of 0.001. These alignments were then
clustered using Spectral Clustering of Protein Sequences (SCPS) [22] to infer homology relations
between sequences based on pairwise similarity scores. Members of the four protein families,
PFam32, PFam49, PFam50 and PFam57/62, as reasonable candidates for functioning in plasmid
replication and partitioning [23], were identified on each contig (S1 Table). Plasmids were
assigned names according to the PFam32 type matching their encoding PFam32 member (>95%

amino acid sequence identity). Two exceptions were made to this method: cp9 and lp28-8. The

smallest plasmid, cp9, does not encode PFam32, so it was identified by the sequence type of its

PFam49, PFam50 and PFam57/62 members (S1 Table). Other than cp9, short contigs with no

identifiable PFam32 type were not considered. The PFam32 protein from what would eventually

be identified as the MM1 lp28-8 plasmid shared only a 60% amino acid sequence identity with

PFam32 sequence type from B31 and JD1 cp32-3 plasmids. Instead, a protein BLAST against

the NCBI non-redundant protein database, revealed it as fully identical to PFam32 type on *B.*

burgdorferi 94a lp28-8 plasmid.

The MM1 lp54 plasmid subtype was identified following the criteria by Casjens *et*

al.[24]. Dot plot analysis was performed to identify large (>400 bp) insertions, inversions or

deletions in MM1 lp54 plasmid relative to lp54 subtypes. Nucleotide BLAST sequence

comparisons and overall identical plasmid organization were investigated with Artemis

Comparison Tool (ACT; Release 13.0.0) [20].

Correct replicon identification was confirmed visually by creating genome-scale dotplots

of MM1 versus itself and versus B31 (S2 Fig) using GEPARD v1.4[25].

Functional annotation

Genome features were annotated with RAST v2.0 [26] with the initial NCBI Taxonomy ID set to *B. burgdorferi* (139). To create a comparative annotation map of main chromosomes, FASTA sequences of all *B. burgdorferi* strains with complete genome assemblies were downloaded from NCBI GenBank. B31, CA382, JD1, N40, PAbe, PAli, and ZS7 (accession numbers AE000783.1, CP005925.1, CP002312.1, CP002228.1, CP019916.1, CP019844.1, and CP001205.1, respectively). The main chromosomes of B31, CA382, JD1, N40, and ZS7 were re-annotated with RAST annotation server using the same settings used for MM1 genome annotation. GenBank annotations for PAbe and PAli were used without modification.

Phylogenetic analysis

A phylogenetic tree was constructed to evaluate the relationship of MM1 with respect to seven prominent laboratory *B. burgdorferi* s.s. strains. The tree was inferred from the whole main chromosome sequence data of s.s. strains MM1, B31, CA382, JD1, N40, PAbe, PAli, and ZS7 using REALPHY v.1.12[27] with outgroup *B. bissettii* DN127 (CP002746.1). The strains were ordered for plotting based on genomic distance from MM1 as determined by GGDC v.2.1[28] with its default settings for formula two. The tree was plotted using TreeGraph2 v2.13[29].

The eight chromosomal housekeeping genes *clpA*, *clpX*, *nifS*, *pepX*, *pyrG*, *recG*, *rplB* and *uvrA* of the Multi Locus Sequence Typing (MLST) scheme [30] were used to determine MM1 Sequence Type (ST). Corresponding locus sequences in MM1 were queried against the *Borrelia* MLST database [31]. Alleles with perfect matches to the database were assigned type numbers.

Where perfect matches did not exist, single-base mismatches were noted and confirmed using Oxford Nanopore data. For confirmation, the novel alleles were searched against all MinION filtered 2D reads using FASTA v.36[32]. Hits were aligned with Clustal Omega v.1.2.1[33] and visually inspected (S3 Fig).

Results and discussion

Overall description of the *B. burgdorferi* MM1 genome

The MM1 genome, comprising a single large chromosome and fifteen plasmids, was determined to be 1,280,240 bp in size based on an average coverage depth of 737× that ranged from 351× to 2312× across the replicons (Table 1). The main linear chromosome represents 71% of the genome size (908,512 bp), with the plasmid content providing the remaining 29% (371,728 bp). According to RAST analysis, the MM1 genome carries 1338 annotated genes with 865 (65%) on the main chromosome including 36 RNA coding genes and 473 plasmid features (Table 1). Of all the annotated features, 37% are connected to a subsystem, the components of which are mainly (92%) on the chromosome (Table 1).

Table 1. GenBank, Coverage and Annotation Details for *B. burgdorferi* MM1 Genome

Replicon[a]	Accession	Base Coverage	Annotations[o]
Chromosome	MG385646	782	865
lp54	MG385643	804	74
lp17	MG385647	832	17
lp25	MG385648	637	25

lp28-3	MG385649	550	25
lp28-4	MG385650	655	31
lp28-8	MG385657	589	29
lp36	MG385651	789	34
cp26	MG385644	725	27
cp9	MG385645	2312	12
cp32-1	MG385655	555	53
cp32-4	MG385656	435	20
cp32-5	MG385658	376	31
cp32-6	MG385652	741	28
cp32-7	MG385654	351	20
cp32-9	MG385653	662	47

[a] GenBank BioProject PRJNA397229, BioSample SAMN07452586.

[b] Of all annotated features, 456 chromosomal and 39 total plasmid features are linked to a subsystem.

From among the subsystem categories present in the MM1 genome, Protein Metabolism and Sulfur Metabolism have the highest and lowest subsystem feature counts, respectively. Interestingly, all the subsystem categories except for "Phages, Prophages, Transposable elements, Plasmids" category are also present in the s.s. B31 genome (Table 2).

Given that the plasmid profile of *B. burgdorferi* may undergo changes as a result of prolonged *in vitro* cultivation [34–36], our sequenced low-passage culture of MM1 (see Materials and methods), like other sequenced *Borrelia,* may be no longer homogeneous with some plasmids lost during culturing.

Table 2. Subsystem Categories Present in MM1 Genome in Relation to B31

Category	Subsystem feature counts	
	MM1	B31
Protein Metabolism	171	164
RNA Metabolism	68	82
Motility and Chemotaxis	62	58
DNA Metabolism	55	55
Carbohydrates	48	49
Nucleosides and Nucleotides	38	34
Fatty Acids, Lipids, and Isoprenoids	37	35
Cell Wall and Capsule	36	41
Amino Acids and Derivatives	32	32
Cell Division and Cell Cycle	26	26
Virulence, Disease and Defense	22	23
Stress Response	21	15
Membrane Transport	18	15
Cofactors, Vitamins, Prosthetic Groups, Pigments	17	21
Phosphorus Metabolism	10	14
Respiration	9	9
Regulation and Cell signaling	7	12
Potassium metabolism	6	4

Phages, Prophages, Transposable elements, Plasmids	6	_
Miscellaneous	3	10
Dormancy and Sporulation	2	2
Sulfur Metabolism	1	1

The main, linear chromosome of *B. burgdorferi* MM1

B. burgdorferi MM1 carries a main linear chromosome of 908,512 bps in length with a GC content of 28.6%—similar in size (902,191 - 922,801 bp) and GC content (28.5 - 28.6%) to the main chromosomes of other s.s. strains that have a complete genome assembly (Table 3). Consistently, the MM1 chromosome shares a 99% nucleotide sequence identity and a similar tandem repeat profile (e.g., microsatellite, minisatellite and macrosatellite) with these other main chromosomes (Table 3).

Table 3. *B. burgdorferi* MM1 Chromosome in Relation to Representative *B. burgdorferi* Strains

Strain	Length (bp)	GC%	Tandem Repeats		
			Micro-satellite	Mini-satellite	Macro-satellite
B. burgdorferi MM1	908512	28.6	933	1	1
B. burgdorferi JD1	922801	28.6	951	2	0
B. burgdorferi PAbe	910728	28.6	939	2	1
B. burgdorferi B31	910724	28.6	940	2	2
B. burgdorferi CA382	910736	28.6	944	2	1

B. burgdorferi PAli	909921	28.6	940	2	0
B. burgdorferi ZS7	906707	28.5	947	2	0
B. burgdorferi N40	902191	28.6	931	0	0

With the exception of the right end extension, the chromosomal gene contents of sequenced s.s. strains are essentially identical [37,38]. Accordingly, comparative annotation map of MM1 and s.s. strains suggests a consistent overall gene synteny over the entire chromosome with some variation in the right end extension (Fig 1).

Fig 1. Comparative annotation map of the main chromosome of *B. burgdorferi* s.s. strains with complete genome assemblies. Rectangles represent annotated genes, color-coded by type: CDS in light blue, predicted CDS in dark blue, hypothetical CDS in red, rRNA in green and tRNA in black. Rectangle heights denote gene length; genes shorter than 1 kb are displayed enlarged for visibility. Features above or below the midline for each strain represent genes transcribed on the top or bottom strand, respectively. The phylogenetic tree was inferred from the whole main chromosome sequence data and rooted using *B. bissettii* DN127 as outgroup.

B. burgdorferi MM1 plasmid content and variation from *B. burgdorferi* strains

An initial mapping of raw sequence reads from the MM1 genome to the reference strain B31 (see Materials and Methods) indicated less than 26% base calling for B31 linear plasmids (lp) lp21, lp28-1, lp28-2, lp38, lp56 and lp5 suggesting their absence in MM1 genome. However,

the presence/absence of other B31 plasmids and, in particular, the circular plasmid (cp) cp32s

that are homologous nearly throughout their lengths in *B. burgdorferi* [37] remained ambiguous.

The observed depth of coverage across B31 plasmids is depicted in S1 Fig.

According to the PFam32 nomenclature scheme [21], the sequenced culture of MM1

carries fifteen plasmids encoding seven linear and seven circular plasmid PFam32 types. The

small cp9 plasmid that does not encode a PFam32 protein defines an additional plasmid

compatibility type in MM1 genome. While some plasmids might be spontaneously lost between

original isolation and the low passage culture used for sequencing, the number of identified

plasmids in MM1 is within the previously defined range in a set of 14 *B. burgdorferi* strains (e.g.

6-11 and 6-12 circular and linear sequenced plasmids, respectively) [24].

A comparison of variation in plasmid profile, size and GC content among MM1, B31 ,

JD1 and N40 strains, is shown in Table 4. Similar to the type strain B31, MM1 carries linear

plasmids lp54, lp17, lp25, lp28-3, lp28-4, lp36 and circular plasmids cp9, cp26, cp32-1, cp32-4,

cp32-6, cp32-7, cp32-9. Among the non-cp32 plasmids, lp17, lp25, and lp36 appear to have

major structural differences between the two strains (Table 4, Fig 2). The MM1 genome carries

the linear plasmid PFam32 type, lp28-8, which is not present in the sequenced B31. In addition,

MM1 carries cp32-5 PFam32 type, which is not encoded by B31 plasmids, but is encoded by the

circular plasmids of N40 and present in strains JD1 in fusion with cp32-5 as cp32-1+5 (Table 4)

[37].

Table 4. *B. burgdorferi* MM1 plasmid content in relation to *B. burgdorferi* B31, JD1 and N40 strains.

Replicon[a]	Code	MM1		B31		JD1		N40	
		Length	GC%	Length	GC%	Length	GC%	Length	GC%
lp54	A	53.8	28.1	53.7	28.1	52.9	28.2	53.8	28.1
lp17	D	14.4	23.7	16.8	23.1	17.5	23.9	20.7	24.3
lp21	U	_	_	18.8	20.7	_	_	_	_
lp25	E	24.2	25.4	24.2	23.4	22.8	25.4	23.4	23.3
lp28-1	F	_	_	28.2	32.8	24.4	32.7	_	_
lp28-2	G	_	_	29.8	31.6	_	_	29.5	31.7
lp28-3	H	26.8	25.0	28.6	25.0	28.8	25.0	_	_
lp28-4	I	27.5	24.5	27.3	24.5	30.1	24.7	29.4	24.6
lp28-5	Y	_	_	_	_	24.6	23.9	26.6	25.3
lp28-6	Z	_	_	_	_	27.0	32.4	_	_
lp28-7	AA	_	_	_	_	30.0	31.6	_	_
lp28-8º	RC	28.5	33.2	_	_	_	_	_	_
lp36	K	28.0	25.5	36.8	26.9	22.8	25.6	29.2	31.6
lp38	J	_	_	38.8	26.1	27.6	22.9	37.9	26.0
lp5	T	_	_	5.2	23.8	_	_	_	_
lp56	Q	_	_	53.0	27.3	_	_	_	_
cp26	B	26.5	26.3	26.5	26.3	26.5	26.2	26.5	26.3
cp9	C	11.4	24.3	9.4	23.7	_	_	8.7	24.3

cp32-1	P	30.9	29.3	30.8	29.4	60.7[n]	29.2	_	_
cp32-3	S	_	_	30.2	28.9	29.7	28.9	_	_
cp32-4	R	13.3	26.8	30.3	29.3	_	_	15.9	27.0
cp32-5	V	21.9	28.5	_	_	60.7[n]	29.2	29.6	29.2
cp32-6	M	20.0	27.7	29.8	29.3	30.8	29.0	_	_
cp32-7	O	14.0	26.3	30.8	29.1	_	_	17.5	27.2
cp32-8	L	_	_	30.9	29.1	31.1	29.1	_	_
cp32-9	N	30.6	29.1	30.7	29.3	30.7	29.2	30.7	29.0
cp32-10[y]	Q	_	_	_	_	30.3	29.3	30.2	29.2
cp32-11	W	_	_	_	_	29.9	29.0	_	_
cp32-12	X	_	_	_	_	30.0	29.3	27.8	28.1
Total plasmids		15 : 7 (L), 8 (C)		21: 12 (L), 9 (C)		20: 11 (L), 9 (C)		16: 8 (L), 8 (C)	

[a]Length is listed in kbps.

[o]In strains of *B. afzelii*, where lp28-8 is documented (e.g. K78 and PKo), code "AC" is initially given to plasmid lp28-8 and later revised to "RS", whereas, in others (e.g. *B. burgdorferi* strain 94a, *B. valaisiana* VS116, *B. spielmanii* A14S), code N is initially given and later revised to "RS" [24]. Consistently, we designate "RS" as the code for MM1 lp28-8 plasmid.

[n]JD1cp32-1 and cp32-5 are described as fused into 60.7 kbps circular cp32-1+5 plasmid [37]. [y]In B31 cp32-10 is integrated into lp56 plasmid [37]. L, linear; C, Circular.

Fig 2. A diagrammatic depiction of the differences in the organization of the non-cp32 plasmids present in *B. burgdorferi* MM1 and B31 strains. Annotated genes with or without CDSs are represented as grey arrows. PFam organization and synteny is displayed on each plasmid in MM1 (top) and B31

(bottom) strains as red (PFam32), green (PFam49), yellow (PFam50) and blue (PFam57/62) arrows. Among B31the 4 PFams in B31, genes that appear to be disrupted and pseudogene relatives are not color coded [23] and only PFam types with consecutive numbers in MM1 are color coded. BLAST alignments between plasmid pairs (E-value <0.001) are indicated by blocks ranging in color from red (80% sequence identity) to dark blue (100% sequence identity). GC content is calculated in a window of 400 bps and displayed as a blue histogram bar of a GC content lower than 50%. Sequences are shown in full and drawn to scale with scale bars representing 5 kbps.

The plasmids lp54 and cp26 in *B. burgdorferi* MM1

The linear plasmid lp54 and the circular plasmid cp26 have for the most part conserved synteny among the s.l. genomes [39,40]. Among the lp54 plasmid genes with low sequence identity across *Borrelia* species, are the genes encoding decorin binding proteins A (DbpA) and B (DbpB) that are critical for the overall virulence of *B. burgdorferi [41]*. DbpA is highly heterogeneous between and within s.l. species with as low as 70% sequence identity within s.s. strains, whereas DbpB is more conserved (>96% identity) within s.s. [42–44]. Consistent with the substantial difference in their sequence heterogeneity among *Borrelia* isolates, DbpA and DbpB proteins of MM1 (corresponding gene locus tags BbuMM1_A250 and BbuMM1_A260, respectively) share 79% and 100% sequence identity to B31 (Fig 3).

Fig 3. Synteny map of lp54 containing the PFam54 gene array in MM1, JD1, B31 and N40 strains. Left: The lp54 plasmid of MM1 *B. burgdorferi* in comparison to JD1, B31 and N40 strains.The strains are indicated in the left side of the map along with organizational subtypes indicated by Roman numerals in

parenthesis. Plasmid is represented by a horizontal white bar labeled with positions with annotations on light grey bars. BLAST alignments between the lp54 assemblies are indicated by blocks ranging in color from green (95% sequence identity) to dark blue (100% sequence identity) with a minimum Score Cutoff of 28. Annotated features are indicated in dark grey and orange (dbpA and dbpB genes, PFam54 gene array). Right: A zoomed view of PFam54 gene array of lp54 plasmid in JD1, B31, MM1 and N40 strains. PFam54 variable region is sandwiched by *bba64*, *bba65*, and *bba66* on one side, and *bba73* on the other side. Among the 4 strains, MM1 variable region is highly conserved with N40.

Near one terminus of the lp54 plasmid, we identified a PFam54 gene array of close homologs of complement regulator-acquiring surface protein 1, CRASP-1, that has the most variable gene content caused by gene duplication, loss and sequence diversification [45,46]. This variable region is between the genes homologous to *bba66* and *bba73* in a set of ten *Borrelia* lp54 plasmids [45]. The gene order within MM1 lp54 variable region is very different from that in B31; MM1 carries Bbu_MM1A660 located between genes homologous to *bba66* and *bba68*, whereas B31 carries the *bba70* gene that is absent in MM1 (Fig 3). A comparison of the lp54 plasmid variable region in MM1 and N40 strains reveals highly conserved synteny and sequence, with 99% sequence identity between the Bbu_MM1A660 and BbuN40_A67a genes (Fig 3). MM1 and N40 lp54 plasmids have conserved gene order throughout their length. They do not carry insertions, inversions or deletions larger than 400 bps and inter-plasmid DNA exchanges relative to one another [24], therefore, MM1 lp54 plasmid appears to be a type II lp54.

B. burgdorferi cp26 plasmid is present in all natural isolates and encodes functions critical to bacterial viability [47]. The highly diverse outer-surface protein C (OspC) against which mammals develop protective immunity is encoded by the *ospC* gene on the cp26 plasmid

[48]**.** MM1 strain carries a type U *ospC* allele with 100% nucleotide sequence identity to type U

ospC alleles in northern US s.s. strains 94a and CS5[49] 84% sequence identity with type A

ospC allele in B31 strain.

B. burgdorferi MM1 plasmid lp28-8

MM1 plasmid lp28-8 encoding PFam32 protein is 100% and 90% identical in amino acid

sequence to the lp28-8 encoding PFam32 proteins in *B. burgdorferi* 94a and *B. valaisiana*

VS116, respectively. MM1 lp28-8 carries a typical cluster of PFam32, PFam49, PFam50,

PFam57/62 that is 100% identical to the corresponding region of lp28-8 in *B. burgdorferi* 94a

[24] and 86% identical to the corresponding region of lp28-8 in *B. afzelii* strains PKo and K78,

B. valaisiana VS116 and *B. spielmanii* A14S (Fig 4).

Fig 4. A diagrammatic depiction of the differences in the organization of the lp28-8 plasmids in

MM1 and *B. burgdorferi* s.l. species. Annotations are represented as grey arrows. MM1 PFam

organization and synteny are displayed on each alignment as red (PFam32), green (PFam49), yellow

(PFam50) and blue (PFam57/62) arrows. Only PFam types with consecutive numbers are color coded.

BLAST sequence comparison (E-value <0.001) represents a similarity range from 80% (red) to 100%

(blue) among plasmid pairs. GC content is calculated in a window of 400 bps and displayed as a blue

histogram bar of a GC content less or higher than 50%. The size of lp28-8 in *B. burgdorferi* s.s. 94a

(largest contig), *B. afzelii* strain 78, *B. afzelii* strain PKo, *B. valaisiana* VS 116, and *B. spielmanii* A14S is

21,295, 28,638 , 20715, 18022, and 24828 bps respectively. Sequences are shown in full and scale bars

represent 5 kbps.

This assembly of the MM1 lp28-8 plasmid is the most complete for a *B. burgdorferi* s.s. to date. Previous contigs had not approached 28 kb [24,37,50]. In fact, it was recently described as a new PFam32 type present only in one s.s. strain, 94a, from among a set of 14 characterized s.s. isolates. Due to difficulties in assembling long repetitive tracts, the sequence of lp28-8 plasmid of 94a remains unfinished and reported as contigs of 21,295, 4,910 and 393 base pairs [24]. Our sequenced *B. burgdorferi* MM1 lp28-8 is a continuous contig of 28,515 bps in length that is 99% identical to lp28-8 in *B. burgdorferi* 94a over 95% of its sequence, and its leftmost 13 kb shares a 99% identity to plasmid lp38 subtype V. A comparison of sequence and organization of lp28-8 in MM1 to lp28-8 in *B. burgdorferi* 94a and s.l. species such as *B. afzelii [51,52]*, *B. valaisiana* and *B. spielmanii* [53] is depicted in Fig 4.

MM1 lp28-8 plasmid carries a *vls* locus in one end with a characteristic high GC content [54]. The *vls* locus is required for long-term survival of Lyme *Borrelia* in infected mammals and is characterized with a *cis* location of the expression site *vlsE* and a contiguous array of *vls* silent cassettes [54]. Consistently, the average GC content of annotations within the *vls* locus of MM1 is 47% leading to an elevated average GC content of 33.2% for plasmid lp28-8, while the GC content range for the rest of MM1 genome is 23.7% to 29.3% (Table 4, Fig 4). MM1 lp28-8 includes an array of 18 sequences with at least 85% sequence identity to *vlsE* (Fig 5).

Fig 5. The *Vls* antigenic variation system in *B. burgdorferi* MM1. This locus on lp28-8 plasmid contains the expression site *vlsE* as well as a contiguous array of 18 DNA sequence blocks that are each at least 85% identical to *vlsE* (BbuMM1_RS290). BLAST alignments between the MM1 *vlsE* locus and lp28-8 plasmid are indicated by blocks ranging in color from green (85% sequence identity) to dark blue

(100% sequence identity) with a minimum Score Cutoff of 85. The leftmost 13134 bps of lp28-8 plasmid is 99% identical to lp38 subtype V.

The cp32 plasmids of *B. burgdorferi* MM1

The cp32 plasmid family members are described to have 12 PFam32 types in a set of 14 fully analyzed s.s. genome sequences [24]. The *B. burgdorferi* MM1 genome contains six members of the cp32 family, cp32-1, cp32-4, cp32-5, cp32-6, cp32-7, and cp32-9 (Table 4). Based on Spectral Clustering of Protein Sequence analysis, the MM1 cp32 plasmids consistently share an amino acid sequence identity of more than 99% with their PFam32 genes in either B31 or N40 strains (S1 Table). Among the six, cp32-5 plasmid is absent in *B. burgdorferi* B31.

MLST reveals *B. burgdorferi* MM1 strain is a new sequence type

Based on the MLST scheme, no isolate in the *Borrelia* MLST database [31] was found to have the same allelic profile as MM1, therefore, we conclude that MM1 is a unique sequence type (ST). Of the eight loci used in the MLST scheme, six loci (*clpA*, *clpX*, *pyrG*, *recG*, *rplB* and *uvrA* excluding *nifS* and *pepX*) in the MM1 genome were the same allelic type as the corresponding loci in ST37 and ST18 s.s. isolates within the *Borrelia* MLST database (Table 5). *NifS* and *pepX* loci were partially matched (different in only 1 nt) to allelic types 12 and 1, respectively. The closest allelic profile to MM1 was observed in ST37 isolates with 1 bp mismatch in *nifS* loci. Interestingly, ST37 and ST18 isolates belong to Northeastern US or Eastern/East-central Canada, whereas MM1 is a Midwestern USA isolate.

Table 5. MM1 MLST in relation to isolates (ST 37 and ST 18) with 6 out of 8 allelic types identical to MM1

isolate	country[a]	source[o]	allelic type							
			clpA	*clpX*	*nifS*	*pepX*	*pyrG*	*recG*	*rplB*	*uvrA*
MM1[o]	US	M	7	6	_	_	1	5	5	5
ST37	US (19), CA (2)	H (2), T(19)	7	6	12	1	1	5	5	5
ST18	US (20), CA (5)	H (7), T (18)	7	6	6	1	1	5	5	5

[a]All US locations are Northeastern US and Eastern/East-central Canada locations respectively. M, mouse; H, human, T, tick. [o]MM1 *nifS* and *pepX* loci match partially (1 nt difference) to *nifS*-12 and *pepX*-1, respectively.

Conclusions

We report the full genome sequence and organization of *B. burgdorferi* MM1 derived from *in vitro* cultured spirochetes. Like other *B. burgdorferi* s.s. strains, MM1 appears to have a quite evolutionarily stable chromosome with little variation in size and content [21,37,38]. Our MM1 genome assembly produced 15 plasmids, of which seven are linear and eight are circular types. Based on the unique allelic profile of MM1 in MLST analysis, our study identifies MM1 as a new ST among Northern USA *B. burgdorferi* s.s. isolates. Except for the 94a strain, the lp28-8 plasmid had not been previously detected in sequenced cultures of s.s.. MM1 thus represents only the second s.s. strain to contain this plasmid and the first fully assembled. MM1 carries *vls* antigenic variation system on its lp28-8 plasmid which consists of the expression site *vlsE* and an array of 18 cassettes. The characterization of MM1 as a Midwestern USA isolate and

a well-established s.s. strain in Lyme research and vaccine studies, contributes to understanding

Borrelia diversity, and will facilitate the development of more specific diagnostics and vaccines.

Acknowledgments

We thank Barbara Grimes† for initial culturing of the MM1 spirochetes used in this study

and Dr. Qiang Tian for assistance with nanopore sequencing. We also thank Mary Brunkow for

her guidance on the project and Max Robinson for his suggestions for the analysis. We

gratefully acknowledge support from The Wilke Family Foundation, The Steven and Alexandra

Cohen Foundation, Jeff and MacKenzie Bezos, and the National Institutes of Health, National

Institute for General Medical Sciences, National Centers for Systems Biology Grant

2P50GM076547-06A1.

†Passed away June 16, 2017.

References

1. Swedish Council on Health Technology Assessment. Treatment Duration for Lyme Disease. Stockholm: Swedish Council on Health Technology Assessment (SBU); 2016.

2. Lyme disease. Recognising and treating erythema migrans. Prescrire Int. 2015;24: 247–249.

3. Murray TS, Shapiro ED. Lyme disease. Clin Lab Med. 2010;30: 311–328.

4. Hengge UR, Tannapfel A, Tyring SK, Erbel R, Arendt G, Ruzicka T. Lyme borreliosis. Lancet Infect Dis. 2003;3: 489–500.

5. Robinson ML, Kobayashi T, Higgins Y, Calkins H, Melia MT. Lyme carditis. Infect Dis Clin North Am. 2015;29: 255–268.

6. Feder HM Jr, Johnson BJB, O'Connell S, Shapiro ED, Steere AC, Wormser GP, et al. A critical appraisal of "chronic Lyme disease." N Engl J Med. 2007;357: 1422–1430.

7. Xu Y, Johnson RC. Analysis and comparison of plasmid profiles of Borrelia burgdorferi sensu lato strains. J Clin Microbiol. 1995;33: 2679–2685.

8. Loken KI, Wu CC, Johnson RC, Bey RF. Isolation of the Lyme disease spirochete from mammals in Minnesota. Proc Soc Exp Biol Med. 1985;179: 300–302.

9. Lee S-H, Lee J-H, Park H-S, Jang W-J, Koh S-E, Yang Y-M, et al. Differentiation of Borrelia burgdorferi sensu lato through groEL gene analysis. FEMS Microbiol Lett. 2003;222: 51–57.

10. Hughes CA, Johnson RC. Methylated DNA in Borrelia species. J Bacteriol. 1990;172: 6602–6604.

11. Miller JC, Stevenson B. Immunological and genetic characterization of Borrelia burgdorferi BapA and EppA proteins. Microbiology. 2003;149: 1113–1125.

12. Xu Y, Kodner C, Coleman L, Johnson RC. Correlation of plasmids with infectivity of Borrelia burgdorferi sensu stricto type strain B31. Infect Immun. 1996;64: 3870–3876.

13. Anthony C. Caputa, Russell F. Bey, Michael P. Murtaugh. Recombinant vaccine against Lyme disease. US Patent. 5554371 A, 1996.

14. Roger N. Picken HCA. Dna probes and primers for detection of b. burgdorferi using the

polymerase chain reaction. European Patent. 0487709 B1, 1996.

15. Barbour AG. Isolation and cultivation of Lyme disease spirochetes. Yale J Biol Med. 1984;57: 521–525.

16. Ruzić-Sabljić E, Strle F. Comparison of growth of Borrelia afzelii, B. garinii, and B. burgdorferi sensu stricto in MKP and BSK-II medium. Int J Med Microbiol. 2004;294: 407–412.

17. Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. Circlator: automated circularization of genome assemblies using long sequencing reads. Genome Biol. 2015;16: 294.

18. Zhou K, Aertsen A, Michiels CW. The role of variable DNA tandem repeats in bacterial adaptation. FEMS Microbiol Rev. 2014;38: 119–141.

19. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. Bioinformatics. 2011;27: 1009–1010.

20. Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG, Parkhill J. ACT: the Artemis Comparison Tool. Bioinformatics. 2005;21: 3422–3423.

21. Sherwood R. Casjens CHEAIS. Borrelia genomics: chromosome, plasmids, bacteriophages and genetic variation. In: D. Scott Samuels and Justin D. Radolf, editor. Borrelia: molecular biology, host interaction and pathogenesis. Caister Academic Press; pp. 27–54.

22. Nepusz T, Sasidharan R, Paccanaro A. SCPS: a fast implementation of a spectral method for detecting protein families on a genome-wide scale. BMC Bioinformatics. 2010;11: 120.

23. Casjens S, Palmer N, van Vugt R, Huang WM, Stevenson B, Rosa P, et al. A bacterial genome in flux: the twelve linear and nine circular extrachromosomal DNAs in an infectious isolate of the Lyme disease spirochete Borrelia burgdorferi. Mol Microbiol. 2000;35: 490–516.

24. Casjens SR, Gilcrease EB, Vujadinovic M, Mongodin EF, Luft BJ, Schutzer SE, et al. Plasmid diversity and phylogenetic consistency in the Lyme disease agent Borrelia burgdorferi. BMC Genomics. 2017;18: 165.

25. Krumsiek J, Arnold R, Rattei T. Gepard: a rapid and sensitive tool for creating dotplots on genome scale. Bioinformatics. 2007;23: 1026–1028.

26. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid annotations using subsystems technology. BMC Genomics. 2008;9: 75.

27. Bertels F, Silander OK, Pachkov M, Rainey PB, van Nimwegen E. Automated reconstruction of whole-genome phylogenies from short-sequence reads. Mol Biol Evol.

2014;31: 1077–1088.

28. Auch AF, Klenk H-P, Göker M. Standard operating procedure for calculating genome-to-genome distances based on high-scoring segment pairs. Stand Genomic Sci. 2010;2: 142–148.

29. Stöver BC, Müller KF. TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses. BMC Bioinformatics. 2010;11: 7.

30. Margos G, Gatewood AG, Aanensen DM, Hanincová K, Terekhova D, Vollmer SA, et al. MLST of housekeeping genes captures geographic population structure and suggests a European origin of Borrelia burgdorferi. Proc Natl Acad Sci U S A. 2008;105: 8730–8735.

31. Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the population level. BMC Bioinformatics. 2010;11: 595.

32. Pearson WR, Lipman DJ. Improved tools for biological sequence comparison. Proc Natl Acad Sci U S A. 1988;85: 2444–2448.

33. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011;7: 539.

34. Schwan TG, Burgdorfer W, Garon CF. Changes in infectivity and plasmid profile of the Lyme disease spirochete, Borrelia burgdorferi, as a result of in vitro cultivation. Infect Immun. 1988;56: 1831–1836.

35. Grimm D, Elias AF, Tilly K, Rosa PA. Plasmid stability during in vitro propagation of Borrelia burgdorferi assessed at a clonal level. Infect Immun. 2003;71: 3138–3145.

36. Biškup UG, Strle F, Ružić-Sabljić E. Loss of plasmids of Borrelia burgdorferi sensu lato during prolonged in vitro cultivation. Plasmid. 2011;66: 1–6.

37. Casjens SR, Mongodin EF, Qiu W-G, Luft BJ, Schutzer SE, Gilcrease EB, et al. Genome stability of Lyme disease spirochetes: comparative genomics of Borrelia burgdorferi plasmids. PLoS One. 2012;7: e33280.

38. Schutzer SE, Fraser-Liggett CM, Casjens SR, Qiu W-G, Dunn JJ, Mongodin EF, et al. Whole-genome sequences of thirteen isolates of Borrelia burgdorferi. J Bacteriol. 2011;193: 1018–1020.

39. Haven J, Vargas LC, Mongodin EF, Xue V, Hernandez Y, Pagan P, et al. Pervasive recombination and sympatric genome diversification driven by frequency-dependent selection in Borrelia burgdorferi, the Lyme disease bacterium. Genetics. 2011;189: 951–966.

40. Sherwood R. Casjens, Wai Mun Huang, Eddie B. Gilcrease, Weigang Qiu, William D.

Mccaig, Benjamin J. Luft, Steven E. Schutzer, Claire M. Fraser. Comparative Genomics of Borrelia burgdorferi. In: Felipe C. Cabello, Dagmar Hulinska, Henry P. Godfrey, editor. Molecular Biology of Spirochetes. 2006. pp. 79–95.

41. Shi Y, Xu Q, McShan K, Liang FT. Both decorin-binding proteins A and B are critical for the overall virulence of Borrelia burgdorferi. Infect Immun. 2008;76: 1239–1246.

42. Roberts WC, Mullikin BA, Lathigra R, Hanson MS. Molecular analysis of sequence heterogeneity among genes encoding decorin binding proteins A and B of Borrelia burgdorferi sensu lato. Infect Immun. 1998;66: 5275–5285.

43. Heikkilä T, Seppälä I, Saxen H, Panelius J, Peltomaa M, Huppertz H-I, et al. Cloning of the gene encoding the decorin-binding protein B (DbpB) in Borrelia burgdorferi sensu lato and characterisation of the antibody responses to DbpB in Lyme borreliosis. J Med Microbiol. 2002;51: 641–648.

44. Heikkilä T, Seppälä I, Saxen H, Panelius J, Yrjänäinen H, Lahdenne P. Species-specific serodiagnosis of Lyme arthritis and neuroborreliosis due to Borrelia burgdorferi sensu stricto, B. afzelii, and B. garinii by using decorin binding protein A. J Clin Microbiol. 2002;40: 453–460.

45. Wywial E, Haven J, Casjens SR, Hernandez YA, Singh S, Mongodin EF, et al. Fast, adaptive evolution at a bacterial host-resistance locus: the PFam54 gene array in Borrelia burgdorferi. Gene. 2009;445: 26–37.

46. Qiu W-G, Schutzer SE, Bruno JF, Attie O, Xu Y, Dunn JJ, et al. Genetic exchange and plasmid transfers in Borrelia burgdorferi sensu stricto revealed by three-way genome comparisons and multilocus sequence typing. Proc Natl Acad Sci U S A. 2004;101: 14150–14155.

47. Byram R, Stewart PE, Rosa P. The essential nature of the ubiquitous 26-kilobase circular replicon of Borrelia burgdorferi. J Bacteriol. 2004;186: 3561–3569.

48. Barbour AG, Travinsky B. Evolution and distribution of the ospC Gene, a transferable serotype determinant of Borrelia burgdorferi. MBio. 2010;1. doi:10.1128/mBio.00153-10

49. Attie O, Bruno JF, Xu Y, Qiu D, Luft BJ, Qiu W-G. Co-evolution of the outer surface protein C gene (ospC) and intraspecific lineages of Borrelia burgdorferi sensu stricto in the northeastern United States. Infect Genet Evol. 2007;7: 1–12.

50. Margos G, Hepner S, Mang C, Marosevic D, Reynolds SE, Krebs S, et al. Lost in plasmids: next generation sequencing and the complex genome of the tick-borne pathogen Borrelia burgdorferi. BMC Genomics. 2017;18: 422.

51. Casjens SR, Mongodin EF, Qiu W-G, Dunn JJ, Luft BJ, Fraser-Liggett CM, et al. Whole-genome sequences of two Borrelia afzelii and two Borrelia garinii Lyme disease

agent isolates. J Bacteriol. 2011;193: 6995–6996.

52. Schüler W, Bunikis I, Weber-Lehman J, Comstedt P, Kutschan-Bunikis S, Stanek G, et al. Complete genome sequence of Borrelia afzelii K78 and comparative genome analysis. PLoS One. 2015;10: e0120548.

53. Schutzer SE, Fraser-Liggett CM, Qiu W-G, Kraiczy P, Mongodin EF, Dunn JJ, et al. Whole-genome sequences of Borrelia bissettii, Borrelia valaisiana, and Borrelia spielmanii. J Bacteriol. 2012;194: 545–546.

54. Norris SJ. vls Antigenic Variation Systems of Lyme Disease Borrelia: Eluding Host Immunity through both Random, Segmental Gene Conversion and Framework Heterogeneity. Microbiol Spectr. 2014;2. doi:10.1128/microbiolspec.MDNA3-0038-2014

Supporting Information

S1 Fig. Coverage of MM1 sequence reads across *B. burgdorferi* B31 plasmids. An initial mapping of MM1 reads to the type strain B31 suggested lack of plasmids lp21, lp28-1, lp28-2, lp38, lp56 and lp5 in MM1 genome, while the presence/absence status of other plasmids in MM1 remained ambiguous. B31 plasmids are shown in each plot with X-axis and Y-axis representing Reference Start Position and Coverage respectively.

S2 Fig. Genome-scale dotplots of *Bb* MM1 and B31. A) Dotplot of MM1 versus B31 shows a very high degree of similarity. MM1 contains a subset of the plasmids of B31. B) Dotplot of MM1 versus itself illustrates two characteristics of Borrelia genomes. First, the cp32 family of plasmids show great similarity. Second, a large repetitive region, the *vls* locus, appears as a prominent dark square on plasmid lp28-8.

S3 Fig. Nanopore sequencing reads confirm the novel SNPs in *nifS* and *pepX*. Top: The *de novo* assembly of the *nifS* gene matches the MLST *nifS* type 12 except for a C->T polymorphism. Eleven nanopore reads align to this region and all contain the T variant. Bottom: The de novo assembly of the *pepX* gene matches the MLST *pepX* type 1 except for a T->C polymorphism. Nine nanopore reads align to this region and eight contain the C variant. The two SNPs of interest are noted with a red asterisk (*).

S1 Table. MM1 gene members of PFam32, PFam49, PFam50 and PFam57/62 identified by spectral clustering of protein sequences. The table lists the results of SCPS analysis for the 4 PFams 32, 49, 50 and 57/62 (see Materials and methods) in MM1, N40, JD1 and B31 strains. Genes located on replicons of the same type are arranged in nearby rows within each paralogous gene family. Dashes mark missing entries, which may fall below cut-off or may be missing in the respective strain.

Figure 1

Figure 2

Figure 3

Figure 4

B. burgdorferi MM1

B. burgdorferi 94a

B. burgdorferi MM1

B. afzelii K78

B. burgdorferi MM1

B. afzelii PKo

B. burgdorferi MM1

B. valaisiana VS116

B. burgdorferi MM1

B. spielmanii A14S

Figure 5