Energetic costs, precision, and efficiency of a biological motor in cargo transport

Wonseok Hwang[1] and Changbong Hyeon[1, *]

[1]*Korea Institute for Advanced Study, Seoul 02455, Republic of Korea*

Molecular motors play key roles in organizing the interior of cells. An ideal motor would translocate cargos with a high speed and a minimal error in transport distance (or time), while consuming minimal amount of energy. The travel distance and its variance of motor are, however, constrained by energy consumption, the physical principle of which has recently been formulated into the *thermodynamic uncertainty relation*. Here, we reinterpret the thermodynamic uncertainty relation to define the efficiency of a motor in cargo transport. By analyzing the existing motility data of several types of molecular motors, we calculate the energetic cost (or efficiency) of generating time traces of a given precision. The energetic cost for kinesin-1 varies nonmonotonically with ATP concentration and load (f), and is locally minimized at $[\text{ATP}] \approx 200$ μM and $f \approx 4$ pN. Remarkably, this local minimum vanishes and the cost increases significantly for a mutant that has a longer neck-linker, which underscores the importance of molecular structure in transport properties. Calculations show that the energetic cost and precision of motility also vary depending on the motor type. For the biological motors studied here, the efficiency of motor is effectively optimized under cellular condition ($[\text{ATP}] \approx 1$ mM, $f = 0 - 1$ pN). We find that among the motors, kinesin-1 at single molecule level is the most efficient in cargo transport.

Biological systems are in nonequilibrium steady states (NESS) in which the energy and material currents flow constantly in and out of the system. Subjected to incessant thermal and nonequilibrium fluctuations, dynamics of cellular constituents are inherently stochastic and thus error-prone. To minimize errors, biological systems adopt a plethora of energy-consuming strategies out of equilibrium. Trade-off relations between the energetic cost and information processing are ubiquitous in biochemical and biomolecular processes, and have been a recurring theme in biology for many decades [1–3].

A recent study by Barato and Seifert [4] has formulated the trade-off between energy consumption and precision of an observable from dissipative processes in NESS into a concise inequality relation known as the *thermodynamic uncertainty relation*. To be more explicit, the uncertainty measure $\mathcal{Q}$, defined as a product between the energy consumption (heat dissipation, $Q(t)$) from an energy-driven process in steady state and the squared fluctuation of an output observable $X(t)$ relative to its mean $\epsilon_X^2(t) = \langle \delta X^2 \rangle / \langle X \rangle^2$, is constant. It has been further conjectured that for an arbitrary chemical network formulated by Markov jump processes $\mathcal{Q}$ cannot be smaller than $2k_B T$,

$$\mathcal{Q} = Q(t) \times \epsilon_X^2(t) \geq 2k_B T. \qquad (1)$$

The measure $\mathcal{Q}$ quantifies the uncertainty of a dynamic process. The smaller the value of $\mathcal{Q}$, the more regular and predictable is the trajectory generated from the process, rendering the output observable more precise. The proof and physical significance of this inequality have been discussed [4–9]. In the presence of large fluctuations inherent to cellular dynamics, harnessing energy into precise motion is critical for accuracy in cellular computation. The uncertainty measure $\mathcal{Q}$ can be used to assess the efficiency of suppressing the uncertainty in dynamical process via energy consumption.

Historically, the efficiency of heat engine has been discussed in terms of the thermodynamic efficiency, the aim of which is to maximize the amount of work extracted from heat reservoirs with two different temperatures [10]. For a nonequilibrium machine powered by chemical force maintained in the live cell, power production for a given free energy consumption could be a relevant quantity to maximize. Meanwhile, for transport motors in cells, the uncertainty measure $\mathcal{Q}$ can be used to assess the efficiency of a motor (or motors) in cargo transport. Because the displacement (or travel distance) is a natural output observable of interest in the cargo transport, we set $X = l(t)$, which recasts Eq.1 into

$$\mathcal{Q} = \dot{Q} \frac{2D}{V^2} \geq 2k_B T. \qquad (2)$$

$\mathcal{Q}$ is minimized by a motor satisfying the following three conditions: It transports cargos (i) at high speed ($V \sim \langle l(t) \rangle / t$), (ii) with small error ($D \sim \langle \delta l(t)^2 \rangle / t$) in the displacement (or punctual delivery to a target site), and (iii) with small energy consumption ($\dot{Q}$). An efficient cargo transporter is characterized by a small $\mathcal{Q}$ with its minimal bound $2k_B T$.

Here, we calculate the "efficiency" of several biological motors (kinesin-1, KIF17 and KIF3AB in kinesin-2 family, myosin-V, dynein, and F_1-ATPase) in terms of $\mathcal{Q}$, and study how it changes with varying conditions of load

FIG. 1. Analysis of kinesin-1 based on the 6-state double cycle kinetic model. **A.** Schematics of the 6-state double cycle kinetic network for hand-over-hand dynamics of kinesin-1, where T, D, and ϕ denote ATP-, ADP-bound, and apo state, respectively. Through binding $[(1)\rightarrow(2)]$ and hydrolysis of ATP $[(2)\rightarrow(5)]$, and release of ADP $[(5)\rightarrow(6)]$, kinesin moves forward in the $\mathcal{F}$-cycle $[(1) \rightarrow (2) \rightarrow (5) \rightarrow (6) \rightarrow (1)]$, where it takes a backstep in the $\mathcal{B}$-cycle $[(4) \rightarrow (5) \rightarrow (2) \rightarrow (3) \rightarrow (4)]$. The arrows in the figure depict the direction of reaction currents. In both cycles, each chemical step is reversible and the transition rate from the state (i) to (j) is given by k_{ij}. **B.** Reaction current $J_{\mathcal{F}}$, $J_{\mathcal{B}}$, and J as a function of load. The three cartoons illustrate the amount of current along the $\mathcal{F}$ and $\mathcal{B}$ cycles at each value of load. $f < 0$ and $f > 0$ correspond to the assisting and hindering load, respectively. **C.** The ratio between the forward and backward fluxes $J_{\mathcal{F}}/J_{\mathcal{B}}$ as function of f at fixed [ATP]. The stall forces, determined at $J_{\mathcal{F}}/J_{\mathcal{B}} = 1$ (dashed line), are narrowly distributed between $f = 6 - 8$ pN. (**D**–**H**) V, D, $\dot{Q}$, $\dot{W}$, $\dot{E}$ as a function of f and [ATP]. The white dashed lines depict the locus of [ATP]-dependent stall force, and the dashed lines in magenta indicate the condition of [ATP] = 2 mM. **I**. Dependences of $\dot{E}$, $\dot{Q}$, $\dot{W}$ on f at [ATP] = 2 mM.

(f) and [ATP]. Also, of great interest is to identify the optimal condition for motor efficiency, if any, that $\mathcal{Q}$ is minimized. To evaluate $\mathcal{Q}$, one should know $\dot{Q}$, D, and V of the system (see Eq.2), which can be gained by building a proper kinetic network model that can delineate the dynamical characteristics of the system [11]. Our analysis on motors will show that $\mathcal{Q}$ is a complex function of f and [ATP], and that $\mathcal{Q}(f, [\text{ATP}])$ is sensitive to a subtle variation in motor structure.

RESULTS AND DISCUSSION

Chemical driving force, steady state current, and heat dissipation of double cycle kinetic network for kinesin-1

The chemomechanical dynamics of a molecular motor can be mapped on an appropriate kinetic network where nonzero steady state current created by ATP hydrolysis free energy $(-\Delta\mu_{\text{hyd}} > 0)$ acts as the driving force. Given a proper network model, the transport properties of the motor, V and D, can be expressed in terms of a set of rate constants $\{k_{ij}\}$, where k_{ij} is associated with the transition from the i-th to j-th state [4, 12, 13] (see **SI** text). The formal expressions of V and D in terms

of $\{k_{ij}(f, [\text{ATP}])\}$ are used to fit simultaneously the experimental data of V and D obtained under varying conditions of ATP and f [11]. The set of rate constants $\{k_{ij}\}$ decided from the fit further allows us to calculate the affinity ($\mathcal{A}$), and reaction current (J) of the network, and hence the heat dissipation from the motor, $\dot{Q}$.

For kinesin-1, we considered a 6-state network [14], consisting of two cycles, $\mathcal{F}$ and $\mathcal{B}$ (Fig. 1A). This minimal kinetic scheme accommodates 4 different kinetic paths: ATP-hydrolysis induced forward/backward step and ATP-synthesis induced forward/backward step. Although the conventional (N=4)-state unicyclic kinetic scheme confers a similar result with the 6-state double cycle network scheme at small f, it is led to a physically problematic interpretation when the molecular motor is stalled and starts taking backsteps at large hindering load. The backstep in unicyclic reaction schemes, by construction, is produced by a reversal of the forward cycle [15], which implies that the backstep is always realized via the synthesis of ATP from ADP and P_i. More importantly, in calculating $\dot{Q}$ from kinetic network, the unicyclic reaction scheme is led to $\dot{Q} = 0$ under the stall condition, which is not compatible with the physical reality that an idling car still burns fuel and dissipates heat ($\dot{Q} \neq 0$). To build a more physically sensible model that considers the possibility of ATP-induced (fuel-burning)

backstep, we extend the unicyclic reaction scheme into a multi-cyclic scheme which takes into account an ATP-consuming stall, i.e., a futile cycle [14–17].

In the double cycle model, kinesin-1 predominantly moves forward through $\mathcal{F}$-cycle under small hindering ($f > 0$) or assisting load ($f < 0$), whereas it takes a backstep through the $\mathcal{B}$-cycle under large hindering load. In principle, the current within the $\mathcal{F}$-cycle, $J_{\mathcal{F}}$, itself is decomposed into the forward ($J_{\mathcal{F}}^+$) and backward current ($J_{\mathcal{F}}^-$), and $J_{\mathcal{F}} = J_{\mathcal{F}}^+ - J_{\mathcal{F}}^- > 0$ is satisfied in NESS. Although a backstep could be realized through an ATP synthesis [18], corresponding to $J_{\mathcal{F}}^-$, experimental data [15, 19, 20] suggest that such backstep current (ATP synthesis induced backstep, $J_{\mathcal{F}}^-$) is smaller than $J_{\mathcal{B}}^+$ (ATP hydrolysis induced backstep).

The dynamics realized in the double cycle network can be illuminated in terms of variation of $J_{\mathcal{F}}$ and $J_{\mathcal{B}}$ with increasing f (Fig.1B). Without load ($f = 0$), kinesin-1 predominantly moves forward ($J_{\mathcal{F}} \gg J_{\mathcal{B}}$). This imbalance diminishes with increasing f. At stall conditions, the two reaction currents are balanced ($J_{\mathcal{F}} = J_{\mathcal{B}}$), so that the net current J associated with the mechanical stepping defined between the states (2) and (5) vanishes ($J = J_{\mathcal{F}} - J_{\mathcal{B}} = 0$), but nonvanishing current due to chemistry still remains along the cycle of $\rightarrow (2) \rightarrow (3) \rightarrow (4) \rightarrow (5) \rightarrow (6) \rightarrow (1) \rightarrow (2) \rightarrow$. A further increase of f beyond the stall force renders $J_{\mathcal{F}} < J_{\mathcal{B}}$, augmenting the likelihood of backsteps.

For a given set of rate constants, it is straightforward to calculate the rates of heat dissipation ($\dot{Q}$), work production ($\dot{W}$), and total energy supply ($\dot{E}$). The total heat generated from the kinetic cycle depicted in Fig. 1A is decomposed into the heat generated from two subcycles, $\dot{Q}_{\mathcal{F}}$ and $\dot{Q}_{\mathcal{B}}$, each of which is the product of reaction current and affinity [4, 14, 21–24]

$$\dot{Q} = \dot{Q}_{\mathcal{F}} + \dot{Q}_{\mathcal{B}} = J_{\mathcal{F}}\mathcal{A}_{\mathcal{F}} + J_{\mathcal{B}}\mathcal{A}_{\mathcal{B}}. \tag{3}$$

Here, the affinities (driving forces) for the $\mathcal{F}$ and $\mathcal{B}$ cycles are

$$\mathcal{A}_{\mathcal{F}} = k_B T \log\left(\frac{k_{12}k_{25}k_{56}k_{61}}{k_{21}k_{16}k_{65}k_{52}}\right) = (-\Delta\mu_{\text{hyd}}) - fd_0 \tag{4}$$

and

$$\mathcal{A}_{\mathcal{B}} = k_B T \log\left(\frac{k_{23}k_{34}k_{45}k_{51}}{k_{32}k_{25}k_{54}k_{43}}\right) = (-\Delta\mu_{\text{hyd}}) + fd_0. \tag{5}$$

The explicit forms of $J_{\mathcal{F}}$ as a function of $\{k_{ij}\}$ are available (see Eq. S25) but the expression is generally more complicated than the affinity. It is of note that at $f = 0$, the chemical driving forces for $\mathcal{F}$ and $\mathcal{B}$ cycles are identical to be $-\Delta\mu_{\text{hyd}}$. The above decomposition of affinity associated with each cycle into the chemical driving force and the work done by the motor results from the Bell-like expression of transition rate between the states (2)

and (5) as $k_{25} = k_{25}^o e^{-\theta fd_0/k_B T}, k_{52} = k_{52}^o e^{(1-\theta)fd_0/k_B T}$ [14, 25, 26] (see **Materials and Methods**). Thus, Eqs. 3, 4, and 5 lead to

$$\dot{Q} = (J_{\mathcal{F}} + J_{\mathcal{B}})(-\Delta\mu_{\text{hyd}}) - (J_{\mathcal{F}} - J_{\mathcal{B}})fd_0 = \dot{E} - \dot{W} \tag{6}$$

where $\dot{E} = (J_{\mathcal{F}} + J_{\mathcal{B}})(-\Delta\mu_{\text{hyd}})$ is the rate of total chemical free energy supplied to the system via ATP hydrolysis. $\dot{W} = (J_{\mathcal{F}} - J_{\mathcal{B}})fd_0$ is the rate of work production.

Dependences of V, D, $\dot{Q}$, and $\dot{W}$ for kinesin-1 on f and [ATP] make a few points visually clear (see Fig. 1): (i) The stall condition, indicated with a white dashed line in each map, divided all the 2D maps of V, D, $\dot{Q}$, and $\dot{W}$ into two regions; (ii) In contrast to V (Fig. 1D) and D (Fig. 1E), which decrease monotonically with f, $\dot{Q}$ and $\dot{W}$ display non-monotonic dependence on f (Figs. 1F, G). At high [ATP], $\dot{Q}$ is maximized at $f = 0$, 10 pN, whereas $\dot{W}$ is maximized at 5 pN and minimized at 10 pN (see Fig. 1I calculated at [ATP]=2 mM).

At the stall force f_{stall} (≈ 7 pN) (Figs. 1B, 1I, black dashed line), the reaction current of the $\mathcal{F}$-cycle is exactly balanced with that of $\mathcal{B}$-cycle ($J = J_{\mathcal{F}} - J_{\mathcal{B}} = 0$), giving rise to zero work production ($\dot{W} = fV = fd_0 J = 0$). The numbers of forward and backward steps taken by kinesin motors are identical, and hence there is no net directional movement ($V = 0$) [20]. It is important to note that even at the stall condition, kinesin-1 consumes the chemical free energy of ATP hydrolysis, dissipating heat in both forward and backward steps, and hence rendering $\dot{Q} = (J_{\mathcal{F}} + J_{\mathcal{B}})(-\Delta\mu_{\text{hyd}})$ always positive.

Next, branching of reaction current of the $\mathcal{F}$-cycle into $\mathcal{B}$-cycle with increasing f gives rise to nonmonotonic changes of $\dot{Q}$ and $\dot{W}$ with f. At small f, $\dot{Q}$ decreases with f because exertion of load gradually deactivates the $\mathcal{F}$-cycle via the decrease of $\mathcal{A}_{\mathcal{F}}$ (Eq.3, Eq.4, Fig. S2C) and $J_{\mathcal{F}}$ (Fig. S2A). By contrast, $\mathcal{A}_{\mathcal{B}}$ and $J_{\mathcal{B}}$ increase with f (Eq.5, Figs. S2A and D), which leads to an increase of $\dot{Q}_{\mathcal{B}}$. The nonmonotic dependence of $\dot{W}$ on f can be analyzed in a similar way. Because the heat dissipated from the whole network is decomposed into the parts ($\dot{Q} = \dot{Q}_{\mathcal{F}} + \dot{Q}_{\mathcal{B}}$), the activation and deactivation of the cycles $\mathcal{F}$ and $\mathcal{B}$ (i.e., $J_{\mathcal{F}}$ and $J_{\mathcal{B}}$) are negatively correlated (Figs. S2A and B).

Motor head distortion at high external stress hinders the binding and hydrolysis of ATP in the catalytic site [27–29]; $k_{ij} = 0$ when ATP cannot be processed. This effect is modeled into the rate constants such that $k_{ij} = 2k_{ij}^o(1 + e^{\chi_{ij}fd_0/k_B T})^{-1}$ with $\chi_{ij} > 0$ ($k_{ij} \neq k_{25}, k_{52}$) [14]. Thus, it naturally follows that $\dot{Q} = 0$ at f far greater than f_{stall} (Fig.1B, F, I).

FIG. 2. $\mathcal{Q}$ calculated based on kinesin motility data [30] using the 6-state double-cycle model [14] at varying load (f) and [ATP], where $f > 0$ and $f < 0$ signify the hindering assisting load, respectively. **A.** 2-D contour plot of $\mathcal{Q} = \mathcal{Q}(f, [ATP])$. The solid lines in magenta are the loci of minimal $\mathcal{Q}$ at varying [ATP] for a given f (**B**). The dashed lines in magenta are the loci of minimal $\mathcal{Q}$ at varying f for a given [ATP] (**C**). The star symbol marks the cellular condition of [ATP] $\approx$ 1 mM and $f \approx$ 1 pN.

Quantification of $\mathcal{Q}$ for kinesin-1

Unlike V, D, and $\dot{Q}$ that are maximized at large [ATP] and small f (Figs. 1D, E, F), the uncertainty measure $\mathcal{Q}(f, [ATP])$ displays a complex functional dependence (Fig. 2A). (i) Small $\mathcal{Q}$ at low [ATP] is a trivial outcome of the detailed balance condition, when the amount of [ATP] is in balance with [ADP] and [P_i]. The motor, without chemical driving force and only subjected to thermal fluctuations ($\dot{Q} \to 0$), is on average motionless ($V \to 0$); $\mathcal{Q}$ is minimized in this case ($\mathcal{Q} \to 2k_BT$). (ii) In reference to the stall condition ($f = f_{stall}([ATP])$), demarcated by the white dashed lines in Fig.1, $\mathcal{Q}$ is generally smaller for $f < f_{stall}([ATP])$, in which the reaction current along the $\mathcal{F}$-cycle is more dominant than for $f > f_{stall}([ATP])$. At the stall, $\mathcal{Q}$ diverges because of $V \to 0$ and $\dot{Q} \neq 0$. (iii) Notably, a suboptimal value of $\mathcal{Q} \approx 4 \ k_BT$ is identified at [ATP] = 210 μM and $f = 4.1$ pN (Figs. 2).

Comparison of $\mathcal{Q}$ between different types of kinesins

The dynamic property of molecular motor differs from one motor type to another. Effect of modifying motor structure on the transport properties as well as on the directionality and processivity of molecular motor has been of great interest because it provides glimpses into the design principle of a motor at molecular level [29, 32–35]. Here, we explore how modification to motor structure changes the efficiency of motor in terms of $\mathcal{Q}$. To this end, we analyze single-molecule motility data of a mutant of kinesin-1 (Kin6AA), and homodimeric and het-

FIG. 3. $\mathcal{Q}(f, [ATP]$ calculated for (**A**) Kin6AA [17], (**B**) homodirmeric kinesin-2 KIF17 [31], and (**C**) heterotrimeric kinesin-2 KIF3AB [31]. **D**. $\mathcal{Q}([ATP])$ at fixed f (upper panels) and $\mathcal{Q}(f)$ at fixed [ATP] (lower panels) calculated for kinesin-1 (solid magenta lines for 6-state network model and dashed magenta lines for unicyclic model), Kin6AA (orange lines), KIF17 (green lines), and KIF3AB (blue lines). $\mathcal{Q} = 2k_BT$ is marked with the black dashed lines.

erotrimeric kinesin-2 (KIF17 and KIF3AB).

The data of Kin6AA, a mutant of kinesin-1 that has longer neck-linker domain in each monomer, were taken from Ref. [17]. Six amino-acid residues inserted to the neck-linker reduces the internal tension along the neck-linker which plays a critical role for regulating the chemistry of two motor heads and coordinating the hand-over-hand motion [15, 36]. Disturbance to this motif is expected to affect V and D of the wild type. We analyzed the data of Kin6AA again using the 6-state network model (Figs. 1A, S4, S5, Table II. See **SI** for detail), indeed finding reduction of V and D (Fig. S5A) as well as its stall force (Fig. S5A, white dashed line). Of particular note is that the rate constant k_{25} associated with the mechanical stepping process is reduced by two order of magnitude (Table II). In $\mathcal{Q}(f, [ATP])$ (Fig. 3A), the suboptimal point observed in kinesin-1 (Fig. 2A) vanishes (Fig. 3A), and $\mathcal{Q}$ diverges around $\sim$ 4 pN due to the decreased stall force. Finally, overall, the value of $\mathcal{Q}$ has increased dramatically. This means that compared with kinesin-1 ($\mathcal{Q} \approx 7 \ k_BT$), the trajectory of Kin6AA is less regular and unpredictable ($\mathcal{Q} \approx 20 \ k_BT$), and thus Kin6AA is three fold less efficient in cargo transport. Note that $\Delta Q \approx -(2\mathcal{Q}/\epsilon_l^3)\Delta\epsilon_l$, and hence it is energetically demanding to improve the precision when ϵ_l is already small.

Next, the values of $\mathcal{Q}$ were calculated for two active forms of vertebrate kinesin-2 class motors responsible for intraflagellar transport (IFT). KIF17 is a homodimeric form of kinesin-2, and KIF3AB is a heterotrimeric form made of KIF3A, KIF3B, and a nonmotor accessory protein, KAP. To quantify their motility properties, we digitized single-molecule motility data from Ref. [31] and fitted them to the 6-state double-cycle model (Figs. S6, S8) (See **SI** for detail). The calculated $\mathcal{Q}$s for KIF17 and KIF3AB show that $\mathcal{Q}(f, [\text{ATP}])$ is qualitatively similar to the kinesin-1 with some variations. $\mathcal{Q}$ for KIF17 forms a shallow local minimum of $\mathcal{Q} \approx 9.2 \ k_BT$ at $[\text{ATP}] = 200$ μM and $f = 1.5$ pN (Fig. 3B), whereas such suboptimal condition vanishes in KIF3AB (Fig. 3C). However, KIF3AB has a local valley around $f \sim 4$ pN and 1 μM $\lesssim [\text{ATP}] \lesssim 10$ mM in which $\mathcal{Q} \approx 4k_BT$.

The plots of $\mathcal{Q}([\text{ATP}])$ at fixed f and $\mathcal{Q}(f)$ with fixed $[\text{ATP}]$ in Fig. 3D recapitulate the difference between different classes of kinesins more clearly. The following features are noteworthy. (i) An extension of neck-linker domain (Kin6AA, orange lines) dramatically increases $\mathcal{Q}$ compared with the wild type (Kinesin WT, magenta lines). (ii) Non-monotonic behaviors of $\mathcal{Q}([\text{ATP}])$ are qualitatively similar for all kinesins although $\mathcal{Q}$ is, in general, the smallest for kinesin-1. (iii) The movement of KIF3AB (black lines) becomes the most regular at low $[\text{ATP}]$ ($\lesssim 10$ μM). (iv) $\mathcal{Q}$ for the unicyclic model of kinesin-1 (dashed magenta lines) displays only small deviations as long as $0 \lesssim f \lesssim 4$ pN $\ll f_{\text{stall}} \approx 7$ pN.

Comparison of $\mathcal{Q}$ among different types of motors

We further investigate $\mathcal{Q}(f, [\text{ATP}])$ for myosin-V, dynein, and F_1-ATPase using the kinetic network models proposed in the literature [37–39].

Myosin-V: The model studied in Ref. [37] consists of three cycles (Fig. 4A): chemomechanical forward cycle $\mathcal{F}$, dissipative cycle $\mathcal{E}$, and pure mechanical cycle $\mathcal{M}$. In $\mathcal{F}$-cycle, myosin-V either moves forward by hydrolyzing ATP or takes backstep via ATP synthesis. In $\mathcal{M}$-cycle, myosin-V moves backward under load without involving chemical reactions. The $\mathcal{E}$-cycle, consisting of ATP binding $[(2) \rightarrow (5)]$, ATP hydrolysis $[(5) \rightarrow (6)]$, and ADP release $[(6) \rightarrow (2)]$ (Fig. 4B), was originally introduced to connect the two cycles $\mathcal{F}$ and $\mathcal{M}$. The calculation of $J_{\mathcal{E}}$ and $J_{\mathcal{F}}$ reveals that a gradual deactivation of $\mathcal{F}$-cycle with decreasing [ATP] activates the $\mathcal{E}$-cycle (Fig. S11A, 100 μM $\lesssim [\text{ATP}] \lesssim 1$ mM, $f \lesssim 1$ pN). Thus, $\mathcal{E}$-cycle can be deemed as a futile $\mathcal{F}$-cycle, which is activated when chemical driving force is balanced with a load f at low [ATP]. $\mathcal{Q}(f, [\text{ATP}])$ calculated at $[\text{ADP}] = 70$ μM and $[\text{P}_i] = 1$ mM using the rate constants from Ref. [37] (see **SI** for details and Fig. S11) reveals no local minimum

FIG. 4. $\mathcal{Q}$ for various motors as a function of f and [ATP]. **A.** Kinetic model for myosin-V consisting of three cycles $\mathcal{F}$, $\mathcal{E}$, and $\mathcal{M}$ [37]. $\mathcal{Q}(f, [\text{ATP}])$ calculated at $[\text{ADP}] = 70$ μM and $[\text{P}_i] = 1$ mM (See also Fig. S11D for 2-D heat map). **B.** Kinetic model for monomeric dynein [38], and the corresponding $\mathcal{Q}(f, [\text{ATP}])$ calculated at $[\text{ADP}] = 70$ μM and $[\text{P}_i] = 1$ mM (See also Fig.S13D for 2-D heat map). **C.** $\mathcal{Q}(f, [\text{ATP}])$ at $[\text{ADP}] = 70$ μM and $[\text{P}_i] = 1$ mM (See also Fig.S13D for 2-D heat map) using the kinetic model for F_1-ATPase from Ref. [39]. Other quantities such as V, D, and $\mathcal{A}$ as a function of f and [ATP] are provided in Figs. S11, S13, S14. **D.** $\mathcal{Q}([\text{ATP}])$ at fixed f (upper panels) and $\mathcal{Q}(f)$ at fixed [ATP] (lower panels) for kinesin-1 (magenta), myosin-V (orange), and monomeric dynein (green).

in this condition. However, at $[\text{ADP}] = 0.1$ μM and $[\text{P}_i] = 0.1$ μM, which is the condition studied in Ref. [37], a local minimum with $\mathcal{Q} = 6.5$ k_BT emerges at $f = 1.1$ pN and $[\text{ATP}] = 20$ μM (Fig. S12D, Table I). Both values of f and [ATP] at the suboptimal condition of myosin-V are smaller than those of kinesin-1 (Table I). In (N=2)-unicyclic model for myosin-V (Fig. S15) [40], $\mathcal{Q}$ has local valley around $f \sim 2$ pN and $[\text{ATP}] \sim 10$ μM; and $\Delta\mathcal{Q}$ $[\equiv \mathcal{Q} - \mathcal{Q}_b$ where $\mathcal{Q}_b = (\mathcal{A}/N) \coth (\mathcal{A}/2Nk_BT) \geq 2k_BT$ with $N = 2$ is the stronger bound known for unicyclic model [4]] is locally minimized at $f = 0.03$ pN and [ATP] $= 17$ μM. Similar to the result from multi-cyclic model with $[\text{ADP}] = 0.1$ μM and $[\text{P}_i] = 0.1$ μM, the values of f and [ATP] which are associated with the valley of $\mathcal{Q}$ or the local minimum of $\Delta\mathcal{Q}$ are identified over much lower range than those for the kinesin-1 (Table I).

Dynein: $\mathcal{Q}$ for monomeric dynein evaluated using the (N=9)-unicyclic model in Ref. [38] (Fig. 4B) is locally minimized to $\mathcal{Q} \approx 8.1$ $k_B T$ at $f = 2.1$ pN, [ATP] = 420 μM (Fig. 4D, S13D). Compared to kinesin-1, $\mathcal{Q}$ for monomeric dynein is optimized at lower f. $\Delta\mathcal{Q}$ has a local minimum 5.1 $k_B T$ at $f = 2.0$ pN and [ATP] = 480 μM, which are similar conditions that minimize $\mathcal{Q}$ locally (Fig. S13D).

F_1-ATPase: $\mathcal{Q}$ for F_1-ATPase using the (N=6)-state unicyclic model in Ref. [39] reveals that there is a valley around torque $\tau \approx -10$ pN·nm and [ATP] ≈ 10 μM reaching $\mathcal{Q} \approx 4$ $k_B T$ (Fig. 4C). $\Delta\mathcal{Q}$ is locally minimized to 0 $k_B T$ at torque $\tau = -8.6$ pN·nm and [ATP] = 16 μM. In this condition, $\mathcal{Q} = \mathcal{Q}_b = 4.2$ $k_B T$ and $V = 360$ rad/s (Figs. 4C, S14A and D). Notably, $\mathcal{Q}$ for F_1-ATPase is optimized at $\tau < 0$ in which ATP is synthesized, which is consistent with biologically expected role of F_1-ATPase as an ATP synthase *in vivo*.

To compare between the motors, we plot $\mathcal{Q}([\text{ATP}])$ at fixed f and $\mathcal{Q}(f)$ at fixed [ATP] in Fig. 4D. $\mathcal{Q}_{\text{kinesin-1}} < \mathcal{Q}_{\text{monomeric dynein}} < \mathcal{Q}_{\text{myosin-V}}$ over the broad range of f and [ATP]. Remarkably, $\mathcal{Q}_{\text{myosin-V}}$ is smaller than other motors at low [ATP] ($= 1-10$ μM) and at small f ($= 1-2$ pN).

CONCLUDING REMARKS

Biological motors are far superior to macroscopic machines in converting chemical free energy into linear movement. The thermal noise is utilized to rectify the ATP chemistry induced conformational dynamics into unidirectional movement, which conceptualizes the unidirectional movement by the Brownian ratchet [41], but it also comes with a cost of overcoming the thermal noise that makes the movement of biological motors inherently stochastic and error-prone. Mechanism of harnessing energy into faster and more precise motion is critical for the accuracy of cellular computation. The uncertainty measure $\mathcal{Q}$ quantifies the efficiency of improving the speed and regularity of dynamics for a given energetic cost.

Here, we have quantified the efficiency of biological motors using the uncertainty measure $\mathcal{Q}$ and found that the efficiencies of motors are all semi-optimized near the cellular condition (star symbols in Figs. 2, 3, 4). Q versus ϵ_l plots (Fig. 5, ϵ_θ for F_1-ATPase motor) for the various motors calculated at cellular condition, which is reminiscent of the recent study on the free-energy cost of accurate biochemical oscillations [45], sorts the biological motors in the order of growing $\mathcal{Q}$. The plots indicate that kinesin-1 is the best motor whose $\mathcal{Q}(\approx 7.2$ $k_B T)$ approaches the ideal motor limit ($\mathcal{Q} = 2$ $k_B T$). Note that $\mathcal{Q}(\approx 19$ $k_B T)$ for the mutant kinesin-1, KinAA6, is the diametrical opposite of kinesin-1.

FIG. 5. Plots of Q versus ϵ_l for various motors whose efficiency in cargo transport under cellular condition is quantified with $\mathcal{Q}$. The plot for the ideal motor ($\mathcal{Q} = 2$ $k_B T$) is plotted with a dashed line. $\mathcal{Q}/k_B T = 7.2$ (kinesin-1), 7.7 (F_1-ATPase), 9.1 (monomeric dynein), 9.9 (KIF17), 9.9 (KIF3AB), 13 (myosin-V), 19 (Kin6AA). For cellular condition, we assume [ATP] = 1 mM [42] and $f = 1$ pN [43, 44]; zero torque is assumed for F_1-ATPase.

To what extent are our findings on the *in vitro* single motor properties generalized into those in live cells? First, the force hindering the motor movement varies with cargo size and subcellular location; the load or viscoelastic drag exerted against motors inside the cell varies dynamically [46, 47]. Yet, actual forces opposing the cargo movement in cytosolic environment are $\lesssim 1$ pN [43, 44]. Since $\mathcal{Q}$s for microtubule-based motors, kinesin-1, kinesin-2, and dynein, are narrowly tuned, varying only a few $k_B T$ over the range of $0 \leq f \leq 4$ pN at [ATP] = 1 mM (Figs. 3D, 4D), our discussion on the *in vitro* single motor property in terms of $\mathcal{Q}$ can be extended to cargo transport in cytosolic environment. Next, a team of motors is often responsible for cargo transport in the cell [48]. It has, however, been shown that *in vitro* two kinesin motors attached to a cargo do not coordinate under low load, and saturating ATP [49]. Although trajectories generated by multiple motors have not been analyzed here, extension of the present analysis to such cases should be straightforward.

$\mathcal{Q}(f, [\text{ATP}])$ and the suboptimal condition of $\mathcal{Q}$ vary with the type of motor. Here, we note that the optimal condition for $\mathcal{Q}(f, [\text{ATP}])$ is distinct from that of other quantities such as flux $J(f, [\text{ATP}])$ [50] (equivalent to $V(f, [\text{ATP}])$ and power efficiency $\eta(f, [\text{ATP}]) \equiv \dot{W}/\dot{E}$ (see SI Figures). Remarkably, biological motors in native states effectively optimize $\mathcal{Q}$ in accordance with cellular condition, and that a modification of structure disturbs such suboptimality, also leading to dramatic increase of $\mathcal{Q}$. In axonal transport, of particular importance is the timely and efficient delivery of cellular material, the failure of which is linked to neuropathology [51, 52]. Since there are already numerous regulatory control mechanisms as well as other motors, it could be argued that the role played by the optimized single motor properties dictated by $\mathcal{Q}$ is redundant in light of the overall function

of axonal transport. Yet, given that cellular regulations are realized through a multiple layers of checkpoints [53], the optimized efficiency of motors at single molecule level can also be viewed as one of the apparatus for assuring the optimal cargo transport.

METHODS

To study the transport properties of molecular motors, we used experimental data, available in the literature, of V and D under varying conditions of f and [ATP]. Once a set of kinetic rate constants $\{k_{ij}\}$ that defines the network model is determined by fitting the data of $V(\{k_{ij}(f,[\text{ATP}])\})$ and $D(\{k_{ij}(f,[\text{ATP}])\})$, it is straightforward to calculate $\dot{Q}(f,[\text{ATP}])$ [11], and hence $\mathcal{Q}(f,[\text{ATP}])$.

For kinesin-1, we considered a 6-state double-cycle kinetic network (Fig.1A) and modeled the load-dependent rate constants differently from other kinetic steps [14]. The load dependence of kinetic rate was modeled using $k_{25} = k_{25}^o e^{-\theta f d_0/k_B T}$ and $k_{52} = k_{52}^o e^{-(1-\theta)f d_0/k_B T}$ for the mechanical step between the states (2) and (5), and $k_{ij}(f) = 2k_{ij}^o(1 + e^{\chi_{ij} f d_0/k_B T})^{-1}$ for other steps associated with ATP chemistry. The condition of $\chi_{ij} = \chi_{ji}$ makes only the mechanical transition contribute to the work [14]. The rate constants determined in the $\mathcal{F}$-cycle are used for the corresponding chemical transition steps in the $\mathcal{B}$-cycle [14]. For example, ADP dissociation rate constant k_{23} of $\mathcal{B}$-cycle is equal to k_{56} which describes ADP dissociation in $\mathcal{F}$-cycle. Similarly, $k_{32} = k_{65}, k_{34} = k_{61}, k_{43} = k_{16}, k_{45}^{bi} = k_{12}^{bi}, \chi_{23} = \chi_{56}, \chi_{34} = \chi_{61}, \chi_{45} = \chi_{12}$. k_{54} is determined from the equality $(k_{12}k_{25}k_{56}k_{61}/k_{21}k_{52}k_{65}k_{16}) = (k_{23}k_{34}k_{45}k_{52}/k_{32}k_{43}k_{54}k_{25})$ which results from the fact that ATP hydrolysis free energy that drives the $\mathcal{F}$- and $\mathcal{B}$-cycle is identical, and hence $k_{54} = k_{21}(k_{52}/k_{25})^2$ [14]. As the experimental data from Ref. [30] are scarce at high load condition, which activates the $\mathcal{B}$-cycle, it is not easy to determine all the parameters for $\mathcal{F}$ and $\mathcal{B}$ cycles simultaneously using the existing data. To circumvent this difficulty, we fit the data using the following procedure. First, the affinity $\mathcal{A}$ at $f = 0$ was determined from our previous study that employed the (N=4)-state unicyclic model [11]. Even though $\mathcal{B}$-cycle is not considered in Ref. [11], $J_{\mathcal{B}} \approx 0$ at $f \sim 0$, which justifies the use of unicyclic model at $f \ll f_{\text{stall}}$. Next, the range of parameters were constrained during the fitting procedure (Table III) based on the values obtained in [14, 15]). To fit the data globally, we employed the `minimize` function with 'L–BFGS–B' method from the scipy library.

For Kin6AA, KIF17, and KIF3AB, motility data digitized from Ref. [17, 31] were fit to the same 6-state network model used for kinesin-1. For myosin-V, dynein,

and F_1-ATPase, we employed kinetic network models and corresponding rate constants used in Ref. [37–39]. Further details are provided in **SI**.

ACKNOWLEDGEMENTS

We thank Steven P. Gross for insightful comments on cargo transport *in vivo*. This work was performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1607611. We acknowledge the Center for Advanced Computation in KIAS for providing computing resources.

* hyeoncb@kias.re.kr
[1] J. J. Hopfield, Proc. Natl. Acad. Sci. U. S. A. **71**, 4135 (1974).
[2] P. Mehta and D. J. Schwab, Proc. Natl. Acad. Sci. U. S. A. **109**, 17978 (2012).
[3] G. Lan, P. Sartori, S. Neumann, V. Sourjik, and Y. Tu, Nature physics **8**, 422 (2012).
[4] A. C. Barato and U. Seifert, Phys. Rev. Lett. **114**, 158101 (2015).
[5] T. R. Gingrich, J. M. Horowitz, N. Perunov, and J. L. England, Phys. Rev. Lett. **116**, 120601 (2016).
[6] P. Pietzonka, A. C. Barato, and U. Seifert, Phys. Rev. E. **93**, 052145 (2016).
[7] S. Pigolotti, I. Neri, É. Roldán, and F. Jülicher, Phys. Rev. Lett. **119**, 140604 (2017).
[8] C. Hyeon and W. Hwang, Phys. Rev. E. **96**, 012156 (2017).
[9] K. Proesmans and C. Van den Broeck, EPL **119**, 20001 (2017).
[10] J. S. Lee and H. Park, Scientific Reports **7**, 10725 (2017).
[11] W. Hwang and C. Hyeon, J. Phys. Chem. Lett. **8**, 250 (2017).
[12] Z. Koza, J. Phys. A. **32**, 7637 (1999).
[13] J. L. Lebowitz and H. Spohn, Journal of Statistical Physics **95**, 333 (1999).
[14] S. Liepelt and R. Lipowsky, Phys. Rev. Lett. **98**, 258102 (2007).
[15] C. Hyeon, S. Klumpp, and J. N. Onuchic, Phys. Chem. Chem. Phys. **11**, 4899 (2009).
[16] A. Yildiz, M. Tomishige, A. Gennerich, and R. D. Vale, Cell **134**, 1030 (2008).
[17] B. E. Clancy, W. M. Behnke-Parks, J. O. L. Andreasson, S. S. Rosenfeld, and S. M. Block, Nat. Struct. Mol. Biol. **18**, 1020 (2011).
[18] D. D. Hackney, Proc. Natl. Acad. Sci. U. S. A. **102**, 18338 (2005).
[19] M. Nishiyama, H. Higuchi, and T. Yanagida, Nature Cell Biol. **4**, 790 (2002).
[20] N. J. Carter and R. A. Cross, Nature **435**, 308 (2005).
[21] U. Seifert, Rep. Prog. Phys. **75**, 126001 (2012).
[22] H. Qian and D. A. Beard, Biophys. Chem. **114**, 213 (2005).
[23] H. Qian, Phys. Rev. E **69**, 012901 (2004).

[24] H. Ge and H. Qian, Phys. Rev. E. **81**, 051133 (2010).

[25] M. E. Fisher and A. B. Kolomeisky, Proc. Natl. Acad. Sci. **96**, 6597 (1999).

[26] M. E. Fisher and A. B. Kolomeisky, Proc. Natl. Acad. Sci. U. S. A. **98**, 7748 (2001).

[27] S. Uemura and S. Ishiwata, Nature Struct. Biol. **10**, 308 (2003).

[28] C. Hyeon and J. N. Onuchic, Proc. Natl. Acad. Sci. U. S. A. **104**, 2175 (2007).

[29] C. Hyeon and J. N. Onuchic, Biophys. J. **101**, 2749 (2011).

[30] K. Visscher, M. J. Schnitzer, and S. M. Block, Nature **400**, 184 (1999).

[31] B. Milic, J. O. L. Andreasson, D. W. Hogan, and S. M. Block, Proc. Nati. Acad. Sci. **23**, 201708157 (2017).

[32] J.-C. Liao, M. W. Elting, S. L. Delp, J. A. Spudich, and Z. Bryant, J. Mol. Biol. **392**, 862 (2009).

[33] Z. Bryant, D. Altman, and J. A. Spudich, Proc. Natl. Acad. Sci. U. S. A. **104**, 772 (2007).

[34] B. Jana, C. Hyeon, and J. N. Onuchic, PLoS Comp. Biol. **8**, e1002783 (2012).

[35] M. Hinczewski, R. Tehver, and D. Thirumalai, Proc. Natl. Acad. Sci. U. S. A. **110**, E4059 (2013).

[36] C. Hyeon and J. N. Onuchic, Proc. Natl. Acad. Sci. U. S. A. **104**, 17382 (2007).

[37] V. Bierbaum and R. Lipowsky, Biophys. J. **100**, 1747 (2011).

[38] A. Šarlah and A. Vilfan, Biophys. J. **107**, 662 (2014).

[39] E. Gerritsma and P. Gaspard, Biophys. Rev. Lett. **05**, 163 (2010).

[40] A. B. Kolomeisky and M. E. Fisher, Biophys. J. **84**, 1642 (2003).

[41] R. D. Astumian, Science **276**, 917 (1997).

[42] R. Milo and R. Phillips, *Cell Biology by the Numbers* (Garland Science, 2015).

[43] M. A. Welte, S. P. Gross, M. Postner, S. M. Block, and E. F. Wieschaus, Cell **92**, 547 (1998).

[44] G. T. Shubeita, S. L. Tran, J. Xu, M. Vershinin, S. Cermelli, S. L. Cotton, M. A. Welte, and S. P. Gross, Cell **135**, 1098 (2008).

[45] Y. Cao, H. Wang, Q. Ouyang, and Y. Tu, Nature Physics **11**, 772 (2015).

[46] B. R. J. Narayanareddy, S. Vartiainen, N. Hariri, D. K. O'Dowd, and S. P. Gross, Traffic **15**, 762 (2014).

[47] J. C. Wortman, U. M. Shrestha, D. M. Barry, M. L. Garcia, S. P. Gross, and C. C. Yu, Biophys. J. **106**, 813 (2014).

[48] Q. Li, S. J. King, A. Gopinathan, and J. Xu, Biophys. J. **110**, 2720 (2016).

[49] B. C. Carter, M. Vershinin, and S. P. Gross, Biophys. J. **94**, 306 (2008).

[50] A. I. Brown and D. A. Sivak, Proc. Nati. Acad. Sci. (2017).

[51] E. Mandelkow and E.-M. Mandelkow, Trends in Cell Biol. **12**, 585 (2002).

[52] M. Hafezparast, R. Klocke, and C. Ruhrberg et al., Science **300**, 808 (2003).

[53] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, and P. Walter, *Molecular Biology of the Cell*, 5th ed. (Garland Science, 2008).

[54] A. Wachtel, J. Vollmer, and B. Altaner, Phys. Rev. E **92**, 042132 (2015).

[55] B. E. Clancy, W. M. Behnke-Parks, J. O. L. Andreasson, S. S. Rosenfeld, and S. M. Block, Nat. Struct. Mol. Biol..

[56] D. T. Gillespie, J. Phys. Chem..

[57] M. P. Singh, R. Mallik, S. P. Gross, and C. C. Yu, Proc. Natl. Acad. Sci. **102**, 12059 (2005).

[58] J. A. Wagoner and K. A. Dill, J. Phys. Chem. B **120**, 6327 (2016).

[59] U. Seifert, Phys. Rev. Lett. **95**, 040602 (2005).

[60] B. Derrida, J. Stat. Phys. **31**, 433 (1983).

[61] H. Qian, Annu. Rev. Phys. Chem. **58**, 113 (2007).

[62] M. E. Fisher and A. B. Kolomeisky, Proc. Natl. Acad. Sci. U. S. A. **96**, 6597 (1999).

Supplementary Information

CALCULATION OF V AND D OF KINESIN-1 IN 6-STATE MULTI-CYCLIC MODEL

For multicyclic models, V and D can be expressed as functions of a set of rate constants $\{k_{ij}\}$ that defines the master equation by generalizing the method by Koza [12] (Alternatively, technique based on the large deviation theory can be used. See Ref. [13, 54]). For the derivation, a generating function is defined for each cycle $\mathcal{I}$ in the network model (for the 6-state double cycle kinetic network, $\mathcal{I}$ can represent the forward ($\mathcal{F}$-cycle) or backward cycle ($\mathcal{B}$-cycle) (Fig. 1A)). Next,

by considering an asymptotic limit of the generating function at $t \to \infty$, $V(\{k_{ij}\})$ and $D(\{k_{ij}\})$ shall be found.

Master equation

For a system with N chemical states ($\{1, 2, \cdots, N\}$), a generalized state $\mu^{\mathcal{I}}(t)$ is defined by using the chemical state of the motor at time t and the number of completed $\mathcal{I}$-cycles ($n_c^{\mathcal{I}}(t)$). For example, if kinesin is at chemical state i ($\in \{1, 2, \cdots, 6\}$) at time t, the generalized state of $\mathcal{F}$-cycle is $\mu^{\mathcal{F}}(t) = i + N^{\mathcal{F}} \times n_c^{\mathcal{F}}(t)$. $P(\mu^{\mathcal{I}}, t)$, which represents the probability of the system is at $\mu^{\mathcal{I}}$ at time t, satisfies

$$\frac{\partial P(\mu^{\mathcal{I}}, t)}{\partial t} = \sum_{\xi} K_{\mu^{\mathcal{I}}-\xi, \mu^{\mathcal{I}}} P(\mu^{\mathcal{I}} - \xi, t) - K_{\mu^{\mathcal{I}}, \mu^{\mathcal{I}}-\xi} P(\mu^{\mathcal{I}}, t), \tag{S1}$$

where $K_{\mu,\nu} = \sum_{\alpha} k_{\mu,\nu}^{\alpha}$ and k_{ij}^{α} denotes the rate of transition from state i to state j that follows the α-th pathway. Here, the periodicity of network model imposes $k_{\mu+N,\nu+N}^{\alpha} = k_{\mu,\nu}^{\alpha}$, $K_{\nu+N,\mu+N} = K_{\nu,\mu}$, and $k_{\mu,\nu}^{\alpha} = k_{i,j}^{\alpha}$ for $\mu = i$ (mod N) and $\nu = j$ (mod N). The range of (integer) summation index ξ depends on the existing pathways for $\mathcal{I}$-cycle and can be infinity. Hereafter, the superscript $\mathcal{I}$ on μ shall be omitted for the brevity.

Following Ref. [12], we define

$$P_j(\mu, t) \equiv P(\mu, t) \delta_{\mu,j}^N \tag{S2}$$

where,

$$\delta_{\mu,j}^N = \begin{cases} 1, \text{ if } j = \mu \text{ (mod } N) \\ 0, \text{ otherwise} \end{cases} \tag{S3}$$

Here, $j \in \{1, 2, \cdots, N\}$. Multiplying $\delta_{\mu,j}^N$ on both sides of Eq.S1 and using the equality $\delta_{\mu,j}^N = \delta_{\mu-\xi,j-\xi}^N$, we get

$$\frac{\partial P_j(\mu, t)}{\partial t} = \sum_{\xi} K_{j-\xi,j} P_{j-\xi}(\mu - \xi, t) - K_{j,j-\xi} P_j(\mu, t). \tag{S4}$$

Generating function

We define a generation function from which V and D can be derived straightforwardly. The generating function for $\mathcal{I}$-cycle is defined by

$$Q_j^{\mathcal{I}}(z, t) \equiv \sum_{\mu=-\infty}^{\infty} e^{z X_\mu^{\mathcal{I}}} P_j(\mu, t). \tag{S5}$$

where $X_\mu^{\mathcal{I}}$ denotes the generalized coordinate for $\mathcal{I}$-cycle at generalized state μ. Then Eq.(S4) and the equality $(X_\mu^{\mathcal{I}} - X_{\mu-\xi}^{\mathcal{I}}) P_{j-\xi}(\mu - \xi, t) = (X_j^{\mathcal{I}} - X_{j-\xi}^{\mathcal{I}}) P_{j-\xi}(\mu - \xi, t)$ imposed by $\delta_{\mu,j}^N = \delta_{\mu-\xi,j-\xi}^N$ gives

$$\frac{\partial Q_j^{\mathcal{I}}(z, t)}{\partial t} = \sum_{\xi} \left(\sum_{\mu=-\infty}^{\infty} e^{z X_\mu^{\mathcal{I}}} K_{j-\xi,j} P_{j-\xi}(\mu - \xi, t) \right) - \sum_{\xi} K_{j,j-\xi} Q_j^{\mathcal{I}}(z, t)$$
$$= \sum_{\xi} e^{z d_{j-\xi,j}^{\mathcal{I}}} K_{j-\xi,j} Q_{j-\xi}^{\mathcal{I}}(z, t) - \sum_{\xi} K_{j,j-\xi} Q_j^{\mathcal{I}}(z, t) \tag{S6}$$

where $d_{\mu\nu}^{\mathcal{I}} \equiv X_\nu^{\mathcal{I}} - X_\mu^{\mathcal{I}}$. In general, different cycle has different $\{d_{\mu,\nu}^{\mathcal{I}}\}$. For example, for the $\mathcal{F}$-cycle in Fig. 1A,

$$d_{i,j}^{\mathcal{F}} = \begin{cases} 1, \text{ for } i = 6, j = 1 \\ -1, \text{ for } i = 1, j = 6 \\ 0, \text{ otherwise.} \end{cases} \quad (S7)$$

whereas for the $\mathcal{B}$-cycle,

$$d_{i,j}^{\mathcal{B}} = \begin{cases} 1, \text{ for } i = 3, j = 4 \\ -1, \text{ for } i = 4, j = 3 \\ 0, \text{ otherwise.} \end{cases} \quad (S8)$$

Finally, for a $\mathcal{X}$-cycle,

$$d_{i,j}^{\mathcal{X}} = \begin{cases} 1, \text{ for } i = 2, j = 5 \\ -1, \text{ for } i = 5, j = 2 \\ 0, \text{ otherwise.} \end{cases} \quad (S9)$$

where $d_{ij}^{\mathcal{X}}$ denotes the change in travel distance of motor divided by its step size.

Generator

Eq. (S6) can be expressed more succinctly as

$$\partial_t Q_j^{\mathcal{I}}(z,t) = \sum_{i=1}^{N} \Gamma_{ij}^{\mathcal{I}} Q_i^{\mathcal{I}}(z,t) \quad (S10)$$

where

$$\Gamma_{ij}^{\mathcal{I}}(z) = \begin{cases} \sum_\alpha k_{ij}^\alpha e^{z d_{ij}^{\mathcal{I},\alpha}}, \text{ if } i \neq j \\ -\sum_{m=1(\neq i)}^{N} \sum_\alpha k_{im}^\alpha, \text{ if } i = j \end{cases} \quad (S11)$$

With α, an index to discern the pathways, Γ can be written as $N \times N$ matrix.

Asymptotic limit of generating function

Here we consider the asymptotic limit $(t \to \infty)$ in which V and D are well defined for an arbitrary chemical network model. The general solution of Eq.(S10) can be written as [12]

$$Q_j^{\mathcal{I}}(z,t) = \sum_m T_{mj}^{\mathcal{I}}(z,t) e^{\lambda_m^{\mathcal{I}}(z)t} \quad (S12)$$

where $\lambda_m^{\mathcal{I}}(z)$'s $(m = 0, 1, 2, \ldots, N)$ are the eigenvalues of $\Gamma^{\mathcal{I}}(z)$. For a system in (unique) steady state, the eigenvalues satisfy $\lambda_0^{\mathcal{I}}(0) = 0$ and $\lambda_m^{\mathcal{I}}(0) < 0$ for $m \neq 0$. Thus, at $t \to \infty$ and when $z \sim 0$,

$$\lim_{t \to \infty} Q_j^{\mathcal{I}}(z,t) \sim T_{0j}^{\mathcal{I}}(z,t) e^{\lambda_0^{\mathcal{I}}(z)t} \quad (S13)$$

Now, summed over the index j, Eq.(S5) is led to

$$\begin{aligned} \sum_{j=1}^{N} Q_j^{\mathcal{I}}(z,t) &= \sum_{j=1}^{N} \sum_{\mu=-\infty}^{\infty} e^{z X_\mu^{\mathcal{I}}} P_j(\mu, t) \\ &= \sum_{\mu=-\infty}^{\infty} e^{z X_\mu^{\mathcal{I}}} P(\mu, t) \\ &\equiv Q^{\mathcal{I}}(z,t). \end{aligned} \quad (S14)$$

From Eq.(S13), at $t \to \infty$, we have

$$Q^{\mathcal{I}}(z,t) \sim \sum_j T_{0j}^{\mathcal{I}}(z,t) e^{\lambda_0^{\mathcal{I}}(z)t} = h^{\mathcal{I}}(z,t) e^{\lambda_0^{\mathcal{I}}(z)t} \quad (S15)$$

where $h^{\mathcal{I}}(z,t) \equiv \sum_j T_{0j}^{\mathcal{I}}(z,t)$. Since $Q^{\mathcal{I}}(0,t) = 1$ and $\lambda_0^{\mathcal{I}}(z = 0) = 0$, $h^{\mathcal{I}}(0,t) \sim 1$ at $t \to \infty$.

Velocity and Diffusion coefficient

In this section, we first define flux $J^{\mathcal{I}}$ and diffusion coefficient $D^{\mathcal{I}}$ of $\mathcal{I}$-cycle using $X^{\mathcal{I}}(t)$ at $t \to \infty$. Then by using the asymptotic form of the generating function, we will get the relation between $J^{\mathcal{I}}$ and $D^{\mathcal{I}}$, and lowest eigenvalue $\lambda_0^{\mathcal{I}}(z)$.

The average of generalized coordinate $X^{\mathcal{I}}(t)$ can be written as

$$\langle X^{\mathcal{I}}(t) \rangle = \partial_z Q^{\mathcal{I}}(z,t)|_{z=0} \sim (h^{\mathcal{I}})' + t (\lambda_0^{\mathcal{I}})' \quad (S16)$$

where Eq.(S15) was used and the prime denotes a partial derivative along z at $z = 0$. The flux of $\mathcal{I}$-cycle is defined by

$$J_{\mathcal{I}} \equiv \lim_{t \to \infty} \frac{\langle X^{\mathcal{I}}(t) \rangle}{t} = (\lambda_0^{\mathcal{I}})'. \quad (S17)$$

$J_{\mathcal{X}}$ multiplied by the step size d_0 corresponds to the velocity V of motor

Similarly, the diffusion coefficient $D_{\mathcal{I}}$ is obtained by considering the second moment of $X^{\mathcal{I}}$.

$$\begin{aligned} \langle (X^{\mathcal{I}})^2 \rangle &= \partial_z^2 Q^{\mathcal{I}}(z,t) \\ &= (h^{\mathcal{I}})'' + 2t(h^{\mathcal{I}})'(\lambda_0^{\mathcal{I}})' + t(\lambda_0^{\mathcal{I}})'' + ((\lambda_0^{\mathcal{I}})')^2 t^2, \end{aligned} \quad (S18)$$

which gives

$$D_{\mathcal{I}} = \lim_{t \to \infty} \frac{\langle (X^{\mathcal{I}}(t))^2 \rangle - \langle X^{\mathcal{I}}(t) \rangle^2}{2t} = \frac{(\lambda_0^{\mathcal{I}})''}{2} \quad (S19)$$

and the diffusion coefficient of motor is obtained by $D = d_0^2 D_{\mathcal{X}}$.

Characteristic polynomial

To express the derivatives of $\lambda_0^{\mathcal{I}}$ in terms of rates $\{k_{ij}\}$, we use the characteristic polynomial of $\Gamma^{\mathcal{I}}(z)$ [12].

$$\det(\lambda_0^{\mathcal{I}}\mathbb{I} - \Gamma^{\mathcal{I}}(z)) = \sum_{n=0}^{N}(\lambda_0^{\mathcal{I}})^n C_n(z) = 0. \qquad \text{(S20)}$$

By differentiating both side of Eq.S20 with respect to z and setting $z = 0$, we get

$$C_0' + C_1(\lambda_0^{\mathcal{I}})' = 0, \qquad \text{(S21)}$$

and

$$C_0'' + 2C_1'(\lambda_0^{\mathcal{I}})' + C_1(\lambda_0^{\mathcal{I}})'' + 2C_2((\lambda_0^{\mathcal{I}})')^2 = 0. \qquad \text{(S22)}$$

From Eqs.(S21) and (S22), we get

$$J_{\mathcal{I}} = (\lambda_0^{\mathcal{I}})' = -\frac{C_0'}{C_1} \qquad \text{(S23)}$$

$$D_{\mathcal{I}} = \frac{(\lambda_0^{\mathcal{I}})''}{2} = -\frac{C_0'' + 2C_1'(\lambda_0^{\mathcal{I}})' + 2C_2(\lambda_0^{\mathcal{I}})'}{2C_1} = -\frac{C_0'' + 2J_{\mathcal{I}} + 2C_2 J_{\mathcal{I}}}{2C_1} \qquad \text{(S24)}$$

C_n's and their derivatives, which depend on the choice of $X^{\mathcal{I}}$, can be readily found by taking derivative along $\lambda_0^{\mathcal{I}}(z)$ on the characteristic polynomial and using $\lambda_0^{\mathcal{I}}(0) = 0$ [12].

work.

Explicit expression of $J_{\mathcal{F}}$

Here, we provide the expression of $J_{\mathcal{F}}$ in terms of rate constants $\{k_{ij}\}$ for the 6-state double cycle kinetic net-

$J_{\mathcal{F}} = J_{\mathcal{F}}^+ - J_{\mathcal{F}}^-$

$= \Big(k_{12}(k_{25}k_{32}k_{43} + k_{23}k_{34}k_{45} + k_{25}(k_{32} + k_{34})k_{45})k_{56}k_{61} - k_{16}k_{21}(k_{34}k_{45}k_{52} + k_{32}(k_{43} + k_{45})k_{52} + k_{32}k_{43}k_{54})k_{65} \Big) \Big/$

$\Big(((k_{25}k_{32}k_{43} + k_{23}k_{34}k_{45} + k_{25}(k_{32} + k_{34})k_{45})k_{56} + k_{21}(k_{34}k_{45}(k_{52} + k_{56})$

$+ k_{32}(k_{45}(k_{52} + k_{56}) + k_{43}(k_{52} + k_{54} + k_{56}))))k_{61} + k_{21}(k_{34}k_{45}k_{52} + k_{32}(k_{43} + k_{45})k_{52} + k_{32}k_{43}k_{54})k_{65}$

$+ k_{16}((k_{25}k_{32}k_{43} + k_{23}k_{34}k_{45} + k_{25}(k_{32} + k_{34})k_{45})k_{56} + (k_{32}k_{43}k_{52} + k_{32}k_{45}k_{52} + k_{34}k_{45}k_{52} + k_{32}k_{43}k_{54}$

$+ k_{25}(k_{34}k_{45} + (k_{34} + k_{43})k_{54} + k_{32}(k_{43} + k_{45} + k_{54})) + k_{23}((k_{43} + k_{45})k_{52} + k_{43}k_{54} + k_{34}(k_{45} + k_{52} + k_{54})))k_{65}$

$+ k_{21}(k_{34}k_{45}(k_{52} + k_{56}) + k_{43}k_{54}k_{65} + k_{34}(k_{45} + k_{54})k_{65} + k_{32}(k_{54}k_{65} + k_{45}(k_{52} + k_{56} + k_{65})$

$+ k_{43}(k_{52} + k_{54} + k_{56} + k_{65})))) + k_{12}((k_{34}k_{45}(k_{52} + k_{56}) + k_{32}(k_{45}(k_{52} + k_{56}) + k_{43}(k_{52} + k_{54} + k_{56})))k_{61}$

$+ (k_{34}k_{45}k_{52} + k_{32}(k_{43} + k_{45})k_{52} + k_{32}k_{43}k_{54})k_{65} + k_{23}(k_{45}(k_{52} + k_{56})k_{61}$

$+ k_{43}(k_{52} + k_{54} + k_{56})k_{61} + k_{45}k_{52}k_{65} + k_{43}(k_{52} + k_{54})k_{65} + k_{34}((k_{52} + k_{54} + k_{56})k_{61} + (k_{52} + k_{54})k_{65}$

$+ k_{45}(k_{56} + k_{61} + k_{65}))) + k_{25}(k_{43}k_{54}(k_{61} + k_{65}) + k_{34}(k_{54}(k_{61} + k_{65}) + k_{45}(k_{56} + k_{61} + k_{65}))$

$+ k_{32}(k_{54}(k_{61} + k_{65}) + k_{43}(k_{56} + k_{61} + k_{65}) + k_{45}(k_{56} + k_{61} + k_{65})))) \Big)$

Similarly, the expressions of $J_{\mathcal{B}}$ and $D_{\mathcal{I}}$ in terms of $\{k_{ij}\}$ can be obtained.

ANALYSIS OF OTHER TYPES OF KINESINS

kinesin-1 mutant (Kin6AA)

Single molecule motility data digitized from Ref. [55] was fitted to 6-state network model (Figs. 1A, S4) by using the same method employed for the analysis of kinesin-1 data (**Methods**). However, 4 additional initial conditions for k_{25} ({300, 3000, 30000, 3000000}) were explored. Thus, total 245 initial conditions were explored. The rate constants estimated from this procedure are shown in Table. II.

Kinesin-2 (KIF17, KIF3AB)

Single molecule motility data digitized from Ref. [31] was again fitted to the 6-state model (Fig. 1A, S6, S8) following the identical procedure employed in the analysis of kinesin-1 data (**Methods**). However, two additional initial conditions for k_{25} ({30000, 3000000}) were explored, which results in total 147 initial conditions. The rate constants are shown in Table. II.

MYOSIN-V

We summarize the multi-cyclic model for myosin-V [37] which consists of ATP-dependent chemomechanical forward cycle $\mathcal{F}$, dissipative cycle $\mathcal{E}$, and ratcheting cycle (ATP independent stepping cycle) $\mathcal{M}$ (Fig. 4A). We first provide the explanation of how V and D of myosin-V are calculated. Next, the affinity and heat production ($\dot{Q}$) are expressed in terms of a set of rates $\{k_{ij}\}$. Finally, $\mathcal{Q}$ shall be calculated using V, D, and $\dot{Q}$.

Calculation of V and D

The $\mathcal{M}$-cycle consisting of a single state (Fig.4A) prevents the application of Eq.(S11). To circumvent this difficulty, the model with additional state $(5')$ is considered (Fig. S10). The $(5')$-state is chemically equivalent to the state (5), but describes motor in different position on actins, such that $X(5) = X_0$ and $X(5') = X_0 \pm d_0$ where $d_0 = 36$ nm for myosin-V. In this new network, the rate constants κ_{ij}'s are

$$
\begin{aligned}
\kappa_{2,5} = \kappa_{2,5'} &= \frac{k_{25}}{2} \\
\kappa_{6,5} = \kappa_{6,5'} &= \frac{k_{65}}{2} \\
\kappa_{5,2} = \kappa_{5',2} &= k_{52} \\
\kappa^f_{5,5'} = \kappa^f_{5',5} &= k_{55,f} \\
\kappa^b_{5,5'} = \kappa^b_{5',5} &= k_{55,b}
\end{aligned}
\tag{S25}
$$

where the subscripts f and b denote the forward and backward motion, respectively. Other rate constants satisfy $\kappa_{ij} = k_{ij}$. This modification can be justified by considering stochastic movement of myosin-V on the chemical network [56]: $\kappa_{i,5}, \kappa_{i,5'}$ are set to $k_{i5}/2$, such that the outgoing fluxes from the states $i = (2)$, (6) to the state (5) remain identical in the both networks depicted in Fig.4A and Fig.S10. Next, we set $\kappa_{5,i} = \kappa_{5',i}$ to keep the inward fluxes toward (6), (2) idential for the two networks. Finally, $\kappa^f_{5,5'} = \kappa^f_{5',5} = k_{55,f}$ and $\kappa^b_{5,5'} = \kappa^b_{5',5} = k_{55,b}$. These modification of rate constants enable us to describe transitions within the $\mathcal{M}$-cycle.

Now, the distance matrix scaled by d_0 is

$$
\begin{aligned}
d^{\mathcal{X}}_{3,4} &= 1, \\
d^{\mathcal{X}}_{4,3} &= -1, \\
d^{\mathcal{X},f}_{5,5'} &= 1, \\
d^{\mathcal{X},f}_{5',5} &= 1, \\
d^{\mathcal{X},b}_{5,5'} &= -1, \\
d^{\mathcal{X},b}_{5',5} &= -1.
\end{aligned}
\tag{S26}
$$

Other elements $(d^{\mathcal{X}}_{i,j})$ are all zero. Thus, $\Gamma^{\mathcal{X}}_{i,j}$ is written as (with $(7) \equiv (5')$)

$$
\begin{pmatrix}
-\kappa_{12}-\kappa_{14} & \kappa_{12} & 0 & \kappa_{14} & 0 & 0 & 0 \\
\kappa_{21} & -\kappa_{21}-\kappa_{23}-\kappa_{25}-\kappa_{26}-\kappa_{27} & \kappa_{23} & 0 & e^z\kappa_{25} & \kappa_{26} & \kappa_{27} \\
0 & \kappa_{32} & -\kappa_{32}-\kappa_{34} & \kappa_{34} & 0 & 0 & 0 \\
\kappa_{41} & 0 & \kappa_{43} & -\kappa_{41}-\kappa_{43} & 0 & 0 & 0 \\
0 & e^{-z}\kappa_{52} & 0 & 0 & -\kappa_{52}-\kappa_{56}-\kappa_{57,b}-\kappa_{57,f} & \kappa_{56} & \kappa_{57,b}+\kappa_{57,f} \\
0 & \kappa_{62} & 0 & 0 & \kappa_{65} & -\kappa_{62}-\kappa_{65}-\kappa_{67} & \kappa_{67} \\
0 & \kappa_{72} & 0 & 0 & \kappa_{75,b}+\kappa_{75,f} & \kappa_{76} & -\kappa_{72}-\kappa_{75,b}-\kappa_{75,f}-\kappa_{76}
\end{pmatrix}
\tag{S27}
$$

Now, the travel velocity V and the diffusion coefficient D of myosin-V are readily acquired by using Eqs.(S23, S24, and S27).

Affinities and heat production

The affinities of individual cycles are

$$
\begin{aligned}
\mathcal{A}_{\mathcal{F}} &= k_B T \log \left(\frac{k_{12}k_{23}k_{34}k_{41}}{k_{21}k_{32}k_{43}k_{41}} \right), \\
\mathcal{A}_{\mathcal{E}} &= k_B T \log \left(\frac{k_{25}k_{56}k_{62}}{k_{52}k_{65}k_{26}} \right), \\
\mathcal{A}_{\mathcal{M}} &= k_B T \log \left(\frac{k_{55,f}}{k_{55,b}} \right).
\end{aligned}
\tag{S28}
$$

Only the following rate constants depend on the load (f):

$$
\begin{aligned}
k_{34} &= k_{34}^o e^{-\theta d_m f/k_B T} \\
k_{43} &= k_{43}^o e^{-(1-\theta)d_m f/k_B T} \\
k_{56} &= k_{56}^o \frac{1 + e^{-\chi d_m f_c/k_B T}}{1 + e^{\chi d_m (f-f_c)/k_B T}} \\
k_{52} &= k_{52}^o \frac{1 + e^{-\chi d_m f_c/k_B T}}{1 + e^{\chi d_m (f-f_c)/k_B T}} \\
k_{55,b} &= \frac{D'}{k_B T} \frac{f d_m - U}{d_m^2} \frac{1}{1 - e^{(U - f d_m)/k_B T}} \\
k_{55,f} &= k_{55,b} e^{-f d_m/k_B T}
\end{aligned}
\tag{S29}
$$

where

$$
\begin{aligned}
\theta &= 0.65 \\
\chi &= 4 \\
f_c &= 1.6 \text{ pN} \\
U &= 20 \ k_B T \\
D' &= 4.7 \times 10^{-4} \mu m/s^2
\end{aligned}
\tag{S30}
$$

as described in Ref. [37]. Thus, the affinities can be written as

$$
\begin{aligned}
\mathcal{A}_{\mathcal{F}} &= k_B T \log \left(\frac{k_{12}^o k_{23}^o k_{34}^o k_{41}^o}{k_{21}^o k_{32}^o k_{43}^o k_{41}^o} \right) - f d_0 \\
\mathcal{A}_{\mathcal{E}} &= \mathcal{A}_{\mathcal{E},f=0} \\
\mathcal{A}_{\mathcal{M}} &= -f d_0.
\end{aligned}
\tag{S31}
$$

The relation $\mathcal{A}_{\mathcal{M}} = -f d_0$ results from the fact that $\mathcal{M}$-cycle is ATP-independent and activated by the load. Thus, the heat production rate of the system is

$$
\dot{Q} = J_{\mathcal{F}} \mathcal{A}_F + J_{\mathcal{E}} \mathcal{A}_E + J_{\mathcal{M}} \mathcal{A}_M
\tag{S32}
$$

$J_{\mathcal{F}}, J_{\mathcal{E}}$, and $J_{\mathcal{M}}$ can be calculated by using Eqs. (S23) and (S27). Finally, $\mathcal{Q}$ for myosin-V is given by

$$
\mathcal{Q}_{\text{Myosin-V}} = \frac{2D}{V^2} \dot{Q}
\tag{S33}
$$

where $D = D_{\mathcal{X}} d_0^2$ and $V = J_{\mathcal{X}} d_0$.

MODEL FOR DYNEIN

The unicyclic model with 9 chemical states for the movement of monomeric dynein is considered based on the model shown in Fig. 3 of Ref. [38]. Identical rates constants shown in Table 3 of Ref. [38] were employed for the calculation. To describe force-dependence, we model the rate constant for power stroke k_{+PS} and for reverse stroke k_{-PS} as follows.

$$
\begin{aligned}
k_{+PS} &= k_{+PS,f=0} e^{-\theta \frac{f d_0}{k_B T}} \\
k_{-PS} &= k_{-PS,f=0} e^{(1-\theta) \frac{f d_0}{k_B T}}
\end{aligned}
\tag{S34}
$$

where $\theta = 0.3$ is selected based on the previous study [57, 58]. Again, V and D were calculated using Eqs. S23, S24.

Affinity and heat production

The affinity for unicyclic model is written as [11, 22, 59]

$$
\begin{aligned}
\mathcal{A} &= k_B T \log \prod_{i=1}^{N} \frac{k_{i,i+1}}{k_{i+1,i}} \\
&= -\Delta \mu_{\text{hyd}} - f d_0
\end{aligned}
\tag{S35}
$$

where $d_0 = 8.2$ nm. The second term, describing force-dependence, is originated from the use of Eq. S34. Finally, $\mathcal{Q}$ is

$$
\mathcal{Q} = \frac{2D}{V d_0} \mathcal{A}.
\tag{S36}
$$

MODEL FOR F$_1$-ATPASE

Here, we summarize the unicyclic model developed for F$_1$-ATPase in Ref. [39]. The model is $(N = 2)$ unicyclic model (Fig. 4C) where 3 cycles in chemical state space correspond to a single rotation in real space (angle changes by 90° upon transition from the state (1) to state (2) whereas transitions from the state (2) to (1)′

induce 30° rotation (Fig. 4C). The model is valid when the torque applied to F_1-ATPase is small enough ($\tau \lesssim 30$ pN·nm) that the mechanical cycle is tightly coupled to the chemical reaction [39]. The dependences of rate constants on the torque are

$$
\begin{aligned}
k_{12}(\tau, \zeta) &= k_{12}^{\mathrm{bi}}(\tau, \zeta) \times [\mathrm{ATP}] \\
&= \frac{1}{e^{a_{k_{12}}(\tau)} + \zeta e^{b_{k_{12}}(\tau)}} \times [\mathrm{ATP}] \\
k_{21'}(\tau, \zeta) &= \frac{1}{e^{a_{k_{21'}}(\tau)} + \zeta e^{b_{k_{21'}}(\tau)}} \\
k_{1'2}(\tau, \zeta) &= k_{1'2}^{\mathrm{bi}}(\tau, \zeta) \times [\mathrm{ADP}][\mathrm{Pi}] \\
&= \frac{1}{e^{a_{k_{1'2}}(\tau)} + \zeta e^{b_{k_{1'2}}(\tau)}} \times [\mathrm{ADP}][\mathrm{Pi}] \\
k_{21}(\tau, \zeta) &= \frac{k_{12}(\zeta, \tau) u_2(\zeta, \tau)}{k_{21}(\zeta, \tau)} e^{(\Delta\mu_{\mathrm{hyd}}^0 + \frac{2\pi}{3}\tau)/k_B T}
\end{aligned}
\tag{S37}
$$

where $\Delta\mu_{\mathrm{hyd}}^0 = -12.5\ k_B T \approx -50$ pN·nm, ζ is the friction coefficient (for example, if the γ-shaft of F_1-ATPase is attached to a bead of radius r, $\zeta = 2\pi\eta r^3(4+3\sin^2 \pi/6)$ [39] with the water viscosity $\eta = 1$ cP $= 10^{-9}$ pN×s× nm^{-2}. In our calculation, $r = 40$ nm as in Ref. [39]), and $a_i(\tau), b_i(\tau)$ are polynomial function of τ defined in Ref. [39].

V, D, affinities, and heat production

For (N=2)-unicyclic model, the speed of rotation V, diffusion coefficient D, and affinity $\mathcal{A}$ are [11, 12, 21, 22, 25, 60, 61]

$$
\begin{aligned}
V &= d_R \frac{k_{1,2}k_{2,1'} - k_{2,1}k_{1',2}}{k_{1,2} + k_{2,1'} + k_{2,1} + k_{1',2}} \equiv d_R J, \\
D &= \frac{d_R^2}{2} \left[\frac{k_{1,2}k_{2,1'}}{k_{2,1}k_{1',2}} + 1 - 2 \left(\frac{k_{1,2}k_{2,1'}}{k_{2,1}k_{1',2}} - 1 \right)^2 \frac{k_{2,1}k_{1',2}}{\sigma^2} \right] \\
&\quad \times \frac{k_{2,1}k_{1',2}}{\sigma}, \\
\mathcal{A} &= k_B T \log\left(\frac{k_{1,2}k_{2,1'}}{k_{2,1}k_{1',2}} \right) \\
&= \left(-\Delta\mu_{\mathrm{hyd}}^0 + k_B T \log\left(\frac{[ATP]}{[ADP][Pi]} \right) \right) - \frac{2\pi}{3}\tau \\
&= -\Delta\mu_{\mathrm{hyd}} - W,
\end{aligned}
\tag{S38}
$$

where $d_R = \frac{2\pi}{3}$ is the radian distance that motor travels upon ATP hydrolysis, $\sigma = k_{12} + k_{21'} + k_{21} + k_{1'2}$, and $W \equiv \frac{2\pi}{3}\tau$ denotes the work done by the motor. Here, $\tau > 0$ implies the motor performs work against the hindering load. Thus, $\mathcal{Q}$ is given by

$$
\mathcal{Q} = \frac{2D}{V d_R}\mathcal{A}
\tag{S39}
$$

.

UNICYCLIC KINETIC MODEL FOR KINESIN-1

To analyze the kinesin-1 data, we also considered ($N = 4$)-unicyclic model (Fig. S3A) which was used in our previous study [11]. Briefly, the model consists of four forward rates $\{u_1, u_2, u_3, u_4\}$ and four backward rates $\{w_1, w_2, w_3, w_4\}$. Only $u_1(= k^{bi}[\mathrm{ATP}])$ depends on [ATP]. Barometric dependence of the rates on forces are assumed: $u_n = u_n^o e^{-f d_0 \theta_n^+ / k_B T}$ and $w_n = w_n^o e^{f d_0 \theta_n^- / k_B T}$ with $\sum_{n=1}^{N}(\theta_n^+ + \theta_n^-) = 1$ [26, 62]. V, D, A, and a set of rate constants used in the calculation of $\mathcal{Q}$ are available in Ref. [11]

UNICYCLIC KINETIC MODEL FOR MYOSIN-V

For myosin-V, we also considered the ($N = 2$) unicyclic model from Ref. [40]. Briefly, the model consists of two forward rates $\{u_1, u_2\}$ and two backward rates $\{w_1, w_2\}$. Only $u_1(= k[\mathrm{ATP}])$ and $w_2(= k'[\mathrm{ATP}]^\alpha)$ depend on [ATP]. Here, $\alpha = 1/2$. Different choice of α introduces only minor difference in the results as argued in [40]. Barometric dependences of the rates on forces are assumed again: $u_n = u_n^o e^{-f d_0 \theta_n^+ / k_B T}$ and $w_n = w_n^o e^{f d_0 \theta_n^- / k_B T}$ with $\sum_{n=1}^{N}(\theta_n^+ + \theta_n^-) = 1$ [26, 62]. The parameters used in the calculation are available in Eqs. (12), (13) in Ref. [40]. Identical expressions for V, D, and $\mathcal{A}$ from Eq. S38 were used for the calculation except for $W = f d_0$.

FIG. S1. Analysis of experimental data, digitized from Ref. [30], using the 6-state model [14] The solid lines are the fits to the data **A.** V vs [ATP] at f =1.05 pN (red square), 3.59 pN (blue circle), and 5.63 pN (black triangle). **B.** V vs f at [ATP] = 5 μM. **C.** V vs f at [ATP] = 2 mM. **D.** Stall force as a function of [ATP], measured by 'Position clamp' (red square) or 'Fixed trap' (blue circle) methods. **E.** D vs [ATP] at f =1.05 pN (red square), 3.59 pN (blue circle), and 5.63 pN (black triangle). D was estimated from $r = 2D/Vd_0$. **F.** D vs f at [ATP] = 2 mM.

FIG. S2. Various physical properties of kinesin-1 calculated using 6-state network model (Fig. 1A) at varying f and [ATP]. **A**. Transport properties (flux J and D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}(f, [\text{ATP}])$. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$. For $f > f_{\text{st}}$, η is set to zero. The white dashed lines indicate the stall condition.

FIG. S3. Various physical properties of kinesin-1 calculated using (N=4)-unicycle model [11, 26] at varying f and [ATP]. We used the same model parameters from Ref. [11]. **A**. Transport properties (V, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}$, $\mathcal{Q}_b$, and their difference $\Delta\mathcal{Q}$. Identical data of $\mathcal{Q}$ and $\Delta\mathcal{Q}$ are shown again as contour plots for clarity. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\mathrm{st}}$). The white dashed lines indicate the stall condition.

FIG. S4. Motility data of Kin6AA, a mutant made of kinesin-1 to which six additional amino-acids are inserted in the neck-linker domains [55], and the theoretical fits made using 6-state double-cycle kinetic network model (Fig. 1A).

FIG. S5. Various physical properties of kin6AA calculated using 6-state network model (Fig. 1A) at varying f and [ATP]. **A**. Transport properties (J, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}(f, [\text{ATP}])$. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S6. Motility data of homodimeric kinesin-2 (KIF17) [31] and their theoretical fits (solid lines).

FIG. S7. Various physical properties of KIF17 calculated using 6-state network model (Fig. 1A) at varying f and [ATP]. **A**. Transport properties (J, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D.** $\mathcal{Q}(f, [\text{ATP}])$. **E.** Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S8. Motility data of heterotrimeric kinesin-2 (KIF3AB) [31] and their theoretical fits (solid lines).

FIG. S9. Various physical properties of KIF3AB calculated using 6-state network model (Fig. 1A) at varying f and [ATP]. **A**. Transport properties (J, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}(f, [\text{ATP}])$. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S10. Augmented kinetic network model of myosin-V. Each line represents a reversible kinetics.

FIG. S11. Various physical properties of myosin-V calculated using the multi-cyclic model [37] at varying f and [ATP]. [ADP] = 70 μM and [Pi] = 1 mM condition was used for the calculation. **A**. Transport properties (J, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D.** $\mathcal{Q}(f, [\text{ATP}])$. **E.** Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S12. Various physical properties of myosin-V calculated using the multi-cyclic model [37] at varying f and [ATP]. [ADP] = 0.1 μM and [Pi] = 0.1 μM condition was used for the calculation. **A.** Transport properties (J, D). **B.** Heat $\dot{Q}$ and work production $\dot{W}$. **C.** Thermodynamic affinity $\mathcal{A}$. **D.** $\mathcal{Q}(f, [\text{ATP}])$. **E.** Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S13. Various physical properties of dynein monomer calculated using ($N = 9$)-unicyclic model [38] at varying f and [ATP]. [ADP] = 70 μM and [Pi] = 1 mM condition was used for the calculation. **A.** Transport properties (V, D). **B.** Heat $\dot{Q}$ and work production $\dot{W}$. **C.** Thermodynamic affinity $\mathcal{A}$. **D.** $\mathcal{Q}(f, [\text{ATP}])$. **E.** Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

FIG. S14. Various physical properties of F$_1$-ATPase calculated using ($N = 2$)-unicyclic model (Fig. 4C) at varying f and [ATP]. [ADP] = 70 μM and [Pi] = 1 mM condition was used for the calculation. **A**. Transport properties (V, D). **B**. Heat $\dot{Q}$ and work production $\dot{W}$. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}(f,[\text{ATP}])$. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

(N=2)-uni cyclic

FIG. S15. Various physical properties of myosin-V calculated using ($N = 2$)-unicyclic model [40] at varying f and [ATP]. **A**. Transport properties (V, D). Same data are shown twice over the different range of f for clarity. **B**. Heat $\dot{Q}$ and work production $\dot{W}$. Same data are shown twice over the different range of f for clarity. **C**. Thermodynamic affinity $\mathcal{A}$. **D**. $\mathcal{Q}(f,[\text{ATP}])$. $\Delta\mathcal{Q}$ is shown twice by using 2-D heat map and contour plot for clarity. **E**. Power efficiency $\eta \equiv \dot{W}/(\dot{W} + \dot{Q})$ ($\eta = 0$ for $f > f_{\text{st}}$). The white dashed lines indicate the stall condition.

TABLE I. Optimal f and [ATP] that locally minimize $\mathcal{Q}$. For unicyclic models of kinesin-1, myosin-V, and F$_1$-ATPase, also shown are f and [ATP] that minimizes $\Delta\mathcal{Q} = \mathcal{Q} - \mathcal{Q}_b$, where Q_b ($\geq 2k_BT$) refers to the stronger lower bound for unicyclic kinetic scheme [4].

	Kinesin-1 (multi-cycle)	Kinesin-1 (unicycle)	KIF17 (multi-cycle)	Myosin-V (multi-cycle) [ADP]=[P$_i$]=0.1 μM	Myosin-V (unicycle)	Dynein monomer (unicycle)	F$_1$-ATPase (unicyclic)
f (pN)	4.1	3.2 [a]	1.5	1.1	0.03 [a]	2.1	8.6 [a] [b]
[ATP] (μM)	210	460 [a]	200	20	17 [a]	420	16 [a]
$\mathcal{Q}_{\min}$ (k_BT)	4.0	4.5	9.2	6.5	14	8.1	4.2
$\Delta\mathcal{Q}_{\min}$ (k_BT)	n/a	1.6	n/a	n/a	0	5.1	0

[a] Condition for local minimization of $\Delta\mathcal{Q} = \mathcal{Q} - \mathcal{Q}_b$.
[b] For F$_1$-ATPase, we consider a resisting torque (τ with the unit of pN·nm) against the rotation of the motor.

TABLE II. Parameters determined for the 6-state double-cycle model [14]. The unit of rate constants ($\{k_{ij}\}$) is s^{-1} except for k_{ij}^{bi} ($[k_{ij}^{bi}] = \mu M^{-1}s^{-1}$). The rates in the table are determined for $f = 0$.

	Kinesin-1	Kin6AA	KIF17	KIF3AB
k_{12}^{bi}	2.8	10	10	10
k_{21}	4200	92	3600	500
k_{25}	1.6×10^6	1.6×10^4	4.0×10^4	6.2×10^6
k_{52}	1.1	3.4	0.079	7.2
k_{56}	190	680	590	92
k_{65}	10	4.1	13	37
k_{61}	250	58	310	320
k_{16}	230	260	1100	750
k_{54}	2.1×10^{-9}	4.3×10^{-6}	1.4×10^{-8}	6.8×10^{-10}
θ	0.61	0.59	0.34	0.82
χ_{12}	0.15	0.12	0.15	0.09
χ_{56}	0.0015	0.0	0.012	0.021
χ_{61}	0.11	0.18	0.17	0.16

TABLE III. Initial values and constraints applied during the fit of kinesin 6-state double-cyclic model. The units are same with shown in Table II. For k_{65} and k_{16}, we used 7 initial values $(0.001, 0.01, 0.1, 1, 10, 100, 1000)$.

k_{12}^{bi}	$0.5 \leq 1.8 \leq 10$	k_{25}	$10^4 \leq 3 \times 10^5 \leq 10^7$	k_{56}	$10 \leq 200 \leq 10^4$	k_{61}	$10 \leq 200 \leq 10^4$
k_{21}	$10 \leq 100 \leq 10^4$	k_{52}	NA	k_{65}	$10^{-4} \leq 10^{[-3,-2,-1,0,1,2,3]} \leq 10^4$	k_{16}	$0^{-4} \leq 10^{[-3,-2,-1,0,1,2,3]} \leq 10^4$
θ	$0 \leq 0.3 \leq 1$	χ_{12}	$0 \leq 0.25 \leq 1$	χ_{56}	$0 \leq 0.05 \leq 1$	χ_{61}	$0 \leq 0.05 \leq 1$