Identification and characterization of *Gypsophila paniculata* color morphs in Sleeping Bear

Dunes National Lakeshore, MI, USA.

Marisa L. Yang[1], Emma Rice[2], Hailee Leimbach-Maus[2], and Charlyn G. Partridge[2]

[1]Environmental Science Policy, and Management, University of California, Berkeley, 260

Mulford Hall, Berkeley, CA 94720, email: marisayang79@gmail.com

[2]Annis Water Resources Institute, 740 W. Shoreline Dr., Muskegon, MI 49441, email:

riceemm@mail.gvsu.edu; hailee.pavisich@gmail.com

Corresponding author: Charlyn Partridge, Annis Water Resources Institute, Grand Valley State

University, 740 W. Shoreline Dr., Muskegon, MI; 49441, email: partridc@gvsu.edu

Abstract

Background.

Gypsophila paniculata (baby's breath) is an invasive species found throughout much of the

northwest United States and western Canada. Recently, plants exhibiting a different color

morphology were identified within the coastal dunes along eastern Lake Michigan. The common

baby's breath (*G. paniculata*) typically produces stems that are purple in color (purple morph),

while the atypical morph has stems that are green-yellow (green-yellow morph). The purpose of

this study was to characterize these newly identified morphs and determine if they are genetically

distinct species from the common baby's breath in order to assess whether alternative

management strategies should be employed to control these populations.

Methods. We sequenced two chloroplast regions, rbcL and matK, and one nuclear region, ITS2,

from the purple morphs and green-yellow morphs collected from Sleeping Bear Dunes National

Lakeshore, MI, USA (SBDNL). Sequences were aligned to the reference sequences from other

Gypsophila species obtained from the Barcode of Life (BOLD) and GenBank databases. We also

collected seeds from wild purple morph and wild green-yellow morph plants in SBDNL. We

grew the seeds in a common garden setting and characterized the proportion of green-yellow

individuals produced from the two color morphs after five-months of growth.

Results. Phylogenetic analyses based upon rbcL, matK, and ITS2 regions suggest that the two

color morphs are not distinct species and they both belong to *G. paniculata*. Seeds collected from

wild green-yellow morphs produced a significantly higher proportion of green-yellow

individuals compared to the number produced by seeds collected from wild purple morphs.

However, seeds collected from both color morphs produced more purple morphs than green-

yellow morphs.

Discussion. Based upon these results, we propose that the two color morphs are variants of *G.*

paniculata. Given the significant difference in the number of green-yellow morphs produced

from the seeds of each morph type, we also suggest that this color difference has some genetic

basis. We propose that current management continue to treat the two color morphs in a similar

manner in terms of removal to prevent the further spread of this species.

Introduction

The Great Lakes sand dunes comprise the most extensive freshwater dune complex in the world, stretching over 1,000 km^2 in Michigan alone. Within northwest Michigan, the sand dunes ecosystem is vital both environmentally and economically. It is home to a number of threatened and endangered species, including piping plover (*Charadrius melodus*) and Pitcher's thistle (*Cirsium pitcheri)*. Colonization of invasive species in this region has the potential to significantly alter the biological composition of these native communities (Leege and Murphy, 2001; Emery et al. 2013). One invasive species of significant concern is the perennial baby's breath (*Gypsophila paniculata*). In 2015, baby's breath was listed by the Michigan Department of Natural Resources (DNR) as a "priority" invasive species for detection and control in Michigan's northern lower peninsula (DNR, 2015). Since its colonization in the region it has spread along a 260 km stretch of the Michigan shoreline. Baby's breath produces a large taproot system that can extend down to 4 meters in depth, which likely helps it outcompete native vegetation for limited resources (Darwent & Coupland, 1966; Karamanski, 2000). In addition, while many of the vulnerable and endangered plant species in these areas are seed limited, (e.g., Pitcher's thistle produces approximately 50-300 seeds per plant total or 'per lifetime' (Bevill et al., 1999)), baby's breath can produce up to 14,000 seeds per plant annually (Stevens, 1957), effectively outcompeting native species in terms of overall yield. This has led to baby's breath composing approximately 50-80% of the ground cover in some areas (Karamanski, 2000; Emery et al., 2013).

One concern with current management efforts is that anecdotal evidence suggests there may be a new baby's breath variant within the Michigan dune system. In 2011 and 2012 The Nature Conservancy (TNC) removal crews reported baby's breath plants with different character

traits than what is commonly observed (TNC, 2014). The atypical morph has stems and leaves

that are lighter in color and more yellow than the common *G. paniculata* purple morph (Figure

1a-c). The purple morph has a thick taproot (4-7 cm in diameter) just below the caudex that

remains unbranched for approximately 60 – 100 cm (Darwent & Coupland, 1966). Severing just

below the intersection of the caudex and the taproot is where manual removal efforts target to

limit regrowth. However, TNC removal crews suggested that the atypical green-yellow morph's

root system seemed to be more diffuse, making it harder to identify a primary taproot and thus,

harder to sever without the potential for regrowth (TNC, 2014). Currently, these green-yellow

morphs are treated with herbicide application (glyphosate) when observed; however, if this is a

newly invaded baby's breath species and it continues to spread into areas where threatened or

endangered species are present, removal methods will be a primary concern and alternative

management strategies may need to be considered for these populations.

One of the first steps toward adapting current management strategies for this invasive is

to identify whether the green-yellow morph is a genetically distinct species from the purple

morph. While *G. paniculata* is the dominant invasive baby's breath species in northwest

Michigan, a number of other species have been introduced to North America and, specifically,

the Great Lakes region (Pringle, 1976, Voss & Reznicek, 2012). For example, *G. elegans*, *G.*

scorzonerifolia, *G. muralis,* and *G. acutifolia* have been collected within Michigan (Pringle,

1976; Reznicek et al., 2011; Voss & Reznicek, 2012), and *G. perfoliata* is reported to have

become naturalized in the United States (Pringle, 1976). *G. muralis* is an annual and has a very

distinct morphology compared to the other *Gypsophila* species identified around the Great

Lakes. It typically only reaches 5-20 cm in height, has linear leaves, and commonly produces

white to pink flowers (Borkaudah, 1962). *G. elegans*, also an annual, is commonly sold in this

region in commercial wildflower packets. It typically has a smaller taproot compared to *G.*

paniculata, and its coloration can be similar to that observed for the green-yellow morph. *G.*

scorzonerifolia and *G. acutifolia* are perennials and specimens of these species have been

collected in counties within the Great Lakes dune system that also contain *G. paniculata*

infestations (Voss, 1957; Pringle, 1976). Both *G. scorzonerifolia* and *G. acutifolia* have a deep

taproot and are similar in height to *G. paniculata,* and thus, can superficially resemble *G.*

paniculata (Voss, 1957; Pringle, 1976). However, both can be distinguished from *G. paniculata*

in that their leaves tend to be longer and wider, and the pedicels and calyces are glandular as

opposed to glabrous in *G. paniculata* (Voss, 1957; Pringle, 1976). Given the potential for these

other species to invade the fragile habitat of the Michigan dune system, the goal of this work was

to characterize the genetic relationship between the newly recognized green-yellow morph and

the common purple morph to determine if they are the same species.

Methods and materials

DNA Extraction, Amplification, and Sequencing

We collected leaf tissue from 1 green-yellow morph and 16 purple morphs in 2016 and

an additional 15 green-yellow morphs in 2017 from Sleeping Bear Dunes National Lakeshore

(SBDNL), Empire, MI, USA (specifically: 44.884941 N, 86.062111 W and 44.875302 N,

86.056821 W). Plant tissue collections were approved by the National Parks Service (permit ID

SLBE-2015-SCI-0013). Leaf tissue was dried in silica gel until DNA extractions could take

place. DNA was extracted using a Qiagen DNeasy Plant Mini Kit (Qiagen, Hilden, Germany).

After extraction, the DNA samples were placed through Zymo OneStep PCR inhibitor removal

columns (Zymo, Irvine, CA) to remove any secondary metabolites that might inhibit PCR

amplification. The DNA for each sample was then quantified using a NanoDrop 2000

(ThermoFisher, Waltham, MA).

The DNA of green-yellow morphs and purple morphs was amplified at three genetic

regions: large subunit of the ribulose-bisphosphate carboxylase gene (rbcL), maturase K (matK),

and internal transcribed spacer 2 (ITS2). A combination of ITS region and matK have been used

to differentiate between other *Gypsophila* species (specifically, *G. elegans* and *G. repens*) in

previous studies (Fior et al., 2006). The rbcL region was amplified using rbcL 1F and rbcL 724R

primers (Fay et al., 1997), matK was amplified using matK 390F and matK 1440R primers (Fior

et al., 2006), and the ITS2 region was amplified using ITS2 2SF and ITS2 S3R primers (Chen et

al., 2010). PCR reactions for all loci consisted of 1X Taq Buffer, 2.0 mM $MgCl_2$, 0.3 µM dNTP,

0.08 mg/mL BSA, 0.4 µM forward primer, 0.4 µM reverse primer, and 0.5 units of Taq

polymerase in a 20 µL reaction volume. The thermal cycle protocols consisted of the following:

for rbcL, an initial denaturing step of 95ºC for 2 minutes, followed by 35 cycles of 94ºC for 1

minute, 55ºC for 30 seconds, and 72ºC for 1 minute. A final elongation step was performed at

72ºC for 7 minutes. For matK, the thermal profile consisted of 26 cycles of 94ºC for 1 minute,

48ºC for 30 seconds, and 72ºC for 1 minute, followed by a final elongation step at 72ºC for 7

minutes. For ITS2, an initial denaturing step of 95ºC for 2 minutes was applied, followed by 35

cycles of 95ºC for 30 seconds, 50ºC for 30 seconds, 72ºC for 1.5 minutes, and a final elongation

step of 72ºC for 8 minutes. Successful amplification was checked by running the PCR product on

a 2% agarose gel stained with ethidium bromide. PCR reactions were then cleaned using

ExoSAP-IT PCR Product Cleanup Reagent (ThermoFisher, Waltham, MA). Sequencing

reactions were performed with the forward and reverse primers for each of the three regions.

Sequencing reactions were cleaned using a Sephadex column (GE Healthcare Life Science,

Marlborough, MA) and sequenced on an ABI Genetic BioAnalyzer 3130xl (Applied Biosystems,

Foster City, CA). Out of the 16 green-yellow morphs a total of 13 were successfully sequenced

for rbcL, 13 were successfully sequenced for matK, and 14 were successfully sequenced for

ITS2. For the purple morphs a total of 15, 12, and 15 individuals were successfully sequenced

for rbcL, matK and ITS2, respectively.

Reference sequences for rbcL, matK, and ITS2 of other *Gypsophila spp*. were

downloaded from either the Barcode of Life Database (BOLD) (http://www.barcodeoflife.org) or

GenBank (https://www.ncbi.nlm.nih.gov/genbank/). We primarily focused on *Gypsophila*

species with reported occurrences within the United States, but also incorporated other species if

their information was available on BOLD. Sequences of the three regions were not always

available for the same species, thus for rbcL these included *G. paniculata*, *G. elegans*, *G.*

fastigiata, *G. scorzonerifolia*, *G. perfoliata*, and *G. muralis*. For matK the species included *G.*

paniculata, *G. elegans*, *G. fastigiata*, *G. scorzonerifolia*, *G. perfoliata*, *G. muralis*, *G. altissima*,

and *G. repens*. For the IST2 region, the species included *G. paniculata*, *G. elegans, G.*

scorzonerifolia, G. perfoliata, G. repens, and *G. acutifolia.* Because sequences for all three

regions were not available for all species, our merged phylogeny only contained *G. paniculata*,

G. elegans, G. scorzonerifolia, and *G. perfoliata* reference sequences. The accession numbers

and sequences for all reference species are provided in Supplemental Table 1. All FASTA files

corresponding to these data will be deposited in the Dryad database and sequences have been

submitted to GenBank (ITS2: MG385003-385031, matK: MG603322-603346, rbcL:

MG547346-547373).

Alignment and Phylogenetic Analysis

All successful sequences from our field samples, as well as sequences for other

Gyposphila species reference sequences obtained from BOLD or GenBank, were imported into

the program MEGA7 (version 7.0.14) (Kumar et al., 2016) and sequences for each of the three

regions were aligned both individually and with all sequences combined using Muscle (Edgar,

2004). The total number of base pairs aligned and analyzed for each region included: 427 base

pair (bp) for rbcL, 702 bp for matK, 201 bp for ITS2, and 1330 bp for the three regions

combined. All alignment parameters were kept at their default settings. Once aligned, we used

MEGA7 to identify the most appropriate substitution model (rbcL: Jukes-Cantor, matK: Tamura

3-parameter, ITS2: Jukes-Cantor, all genes combined: Tamura 3-parameter with gamma

distribution). We then created phylogenetic trees using a maximum-likelihood (ML) approach

with 500-replicated bootstrap analyses, as well as using neighbor joining, and parsimony models.

We also constructed a TCS haplotype network (Clement et al., 2002) based upon the combined

sequences using the statistical parsimony approach (Templeton et al., 1992) in the program

PopART (v 1.7, http://popart.tago.ac.nz).

Color Morph Germination

On May 8, 2018 we planted a total of 207 seeds collected from mature purple morphs

and 255 seeds collected from mature green-yellow morphs from SBDNL. For the purple morph

seeds, these were collected from a total of 14 plants that were sampled in 2016 (average 15 seeds

per plant) and 7 plants that were sampled in 2017 (average 1.7 seeds per plant). For the green-

yellow morphs, these seeds were collected from a total of 17 plants (15 seeds per plant) in 2017.

Plants were grown in the GVSU – Allendale greenhouse from May until August, 2018. The

greenhouse was on a 17 hour light/ 7 hour dark cycle. The average day temperature was 21ºC

and night temperature was 15 ºC. In August the plants were transported to the greenhouse at

AWRI-GVSU where they were allowed to grow until October 15, 2018. The greenhouse at

AWRI-GVSU has no external lighting source or temperature controls, and thus more closely

resembled seasonal day/night and temperature cycles. Plants were sampled after a decrease in

temperature occurred (from a high of 23 ºC on October 10, 2018 to a high of 11 ºC on October

15, 2018). Previous greenhouse observations have found that the differences between the purple

and green-yellow morphs can be best detected after a sudden drop in temperature (personal

observation, CGP). On October 15, we characterized the color of all individuals that successfully

germinated and survived over the 5-month period. We used a chi-square analysis in the R

statistical package (v3.5.1) to determine if germination success differed between seeds from the

two color morphs and whether the proportion of seeds that developed into green-yellow morphs

significantly differed between seeds collected from mature purple and mature green-yellow

plants.

Results

Our results indicate that the green-yellow morph identified in SBDNL is not a genetically

distinct species from the common purple found throughout SBDNL. The rbcL, matK, ITS2, and

combined dataset showed similar patterns with both the green-yellow morphs and the purple

morphs clustering together. The phylogenies constructed from rbcL and matK independently

show that the two color morphs cluster separately from *G. fastigata*, *G. elegan*, *G. muralis*, and

G. repens. For the rbcL locus, the relationship of the color morphs to *G. paniculata* and *G.*

scorzonerifolia was not resolved. Additionally, when we only examined the matK gene, the color

morphs clustered separately but within a clade that also included *G. altissima*, *G. scorzonerifolia*,

and *G. paniculata*. The ITS2 region was able to provide more resolution between *G. paniculata*,

G. scorzonerifolia, *G. acutifolia*, and the color morphs, with the color morphs clustering with *G.*

paniculata (Figure 2a-d; Supplemental Figure 1-8) and separately from the *G. scorzonerifolia*

and *G. acutifolia* clade. The same pattern was observed when all regions were analyzed together

(Figure 2d), with the exception that this phylogeny did not include *G. acutifolia*. In addition, the

TCS haplotype network shows that the purple and green-yellow morphs have shared haplotypes.

These two haplotypes are only one mutation away from one another and the *G. paniculata*

reference, while the next closest species*, G. scorzonerifolia* is 15 mutations away (Figure 3).

This further suggests that both color morphs are *G. paniculata*.

Of the regions analyzed for the green-yellow morphs, purple morphs, and reference

sequences, rbcL was the most conserved sequence with an overall mean genetic distance (*d*) =

0.004, followed by matK (*d* = 0.015) and ITS2 (*d* = 0.038). For the ITS2 region, there were six

purple morphs and one green-yellow morph that clustered together inside the *G. paniculata*

branch (Figure 2c & 2d). Further examination of the electropherograms for these individuals

show that they are likely heterozygous at position 138 of our aligned sequence and amplification

bias of the 'A' single nucleotide polymorphism (SNP) over the allele containing the 'G' SNP is

driving this pattern.

Color Morph Germination

Out of the 207 seeds that were collected from mature purple morphs and planted in the

greenhouse, 82 successfully germinated and survived over the 5 month period (39.6%). Out of

these 82 plants, only one green-yellow morph was produced (1.2%), while the remaining seeds

all produced purple morphs. Out of the 255 seeds collected from mature green-yellow morphs

and planted in the greenhouse, 105 successfully germinated and survived over the 5 month

period (41.2%). This was not significantly different than the proportion of plants that

successfully germinated from the purple morph seeds ($\chi^2 = 0.06$, df = 1, p = 0.81). Of the 105

successfully germinated seeds from the green-yellow morph plants, 12 developed into green-

yellow morph plants (11.4%), 91 developed into purple morphs plants (86.7%), and two plants

could not be determined (they appeared to be green-yellow morphs but displayed some dark

spots on the stem). The proportion of seeds that produced green-yellow individuals significantly

differed between seeds collected from mature green-yellow morphs and seeds collected from

mature purple morphs ($\chi^2 = 5.9$, df = 1, p = 0.015).

Discussion

Overall, our data suggest that the green-yellow morph is not a genetically distinct species

from the purple morph, and that both morphs are *G. paniculata*. For all molecular markers used,

the green-yellow and the purple color morphs grouped together. RbcL, matK, and ITS2 are

common 'barcode' genes used to delineate plant species (Newmaster et al., 2006; Group et al.,

2009; Chen et al., 2010; Yao et al., 2010; Stoeckle et al., 2011) and when used in combination

they provided adequate resolution to separate out the *Gypsophila* species included in this study.

In our data set, rbcL and matK worked well to separate our color morphs from *G. elegans*, *G.

muralis* and both of these species have been reported to occur in the Great Lakes region

(Reznicek et al., 2011; Voss & Reznicek 2012). While the morphology of *G. muralis* is very

distinct from the color morphs in SBDNL, it was initially thought that *G. elegans* shared some

similar traits to that originally described by the TNC removal crews and was a potential

candidate species for the green-yellow color morph. Based upon these results, this is clearly not

the case.

The phylogeny based on ITS2 region and the combined sequences provided the best

resolution for assigning the relationship of our *Gypsophila* species. Like the rbcL and matK

phylogenies, all the purple and green yellow morphs grouped together. For this region, the color

morphs also grouped within the same clade as the *G. paniculata* reference sequence. While *G.*

scorzonerifolia and *G. acutifolia* have also been recorded in the Great Lakes regions (Pringle,

1976), and have a similar general phenotype as *G. paniculata*, these species were clearly within a

distinct clade that was separate from the two color morphs. Similarly, while *G. perfoliata* has

been reported to be naturalized in North America (Pringle, 1976), it grouped outside of the *G.*

paniculata and color morph cluster.

Our greenhouse germination study showed that seeds collected from mature green-yellow

morphs produced a significantly higher proportion of green-yellow individuals than seeds

collected from mature purple morphs. Of the seeds collected from mature green-yellow morphs

11% resulted in green-yellow morphs, while only 1% of seeds from mature purple morphs

resulted in green-yellow morphs. However, seeds from both color morphs primarily produced

purple morphs. The mechanism driving the color difference between the purple and green-

morphs is currently unknown. Within SBDNL, the purple morph is the most common form, with

green-yellow individuals found interspersed in a couple of locations throughout the dunes

(personal observation, CGP/HLM/ER). The largest observed group of green-yellow morphs

consists of a few hundred plants clumped within approximately an acre-sized area and

interspersed throughout large groups of purple morphs. Based upon the dispersal patterns of the

two morphs throughout the dunes, and our germination results, the color difference observed

does not appear to be solely environmentally driven, and likely has a genetic component.

Potential candidate genes that could be influencing these color differences include those involved

in the anthocyanin pathway, which influences red – purple coloration in a number of plants

(Asen et al., 1972; de Pascual-Teresa et al., 2002; Abdel-Aal et al., 2006). In addition,

anthocyanin can rapidly accumulate in the shoots of plants following cold exposure (Leng et al.,

2000), and we have observed an increase in the amount of purple coloration in the purple morphs

as temperatures decrease (personal observation, CGP). Further work will begin to elucidate the

specific mechanism influencing this color difference in invasive *G. paniculata* populations, as

well as to explore whether this color variation drives functional differences between the morphs.

Taken together, these data show that the purple and green-yellow morph within SBDNL

are the same species, and that species is *G. paniculata*. One concern with the green-yellow

morph initially noted by TNC removal crews was that the taproot tended to be more diffuse than

the purple morph, potentially making manual removal of these plants less effective. However, we

have not noted differences in the taproot structure between these two morphs when grown under

controlled conditions (Figure 4). Additionally, our lab's personal observations (CGP, HLM) in

the field have not found any indication that large differences in root structure occur between

mature plants of the two color morphs. Therefore, current management approaches for these

populations should be maintained to control the further spread of *G. paniculata* throughout the

Michigan coastal dune system.

Conclusions

Our data show that both the purple and green-yellow color morph of baby's breath in

Sleeping Bear Dunes National Lakeshore are *G. paniculata* and the observed color differences

likely have some genetic basis. Based on this current information, we recommend that these

color morphs continue to be managed in a similar manner and that distinct management

strategies do not need to be established at this time.

Acknowledgements

We would like to thank Shaun Howard from The Nature Conservancy for his help in identifying

the green-yellow morph and Benjamin Giffin for his help with sequencing analysis. We would

also like to thank Kurt Thompson and Doug Haywick for assisting with the figure construction,

and Alexis Hoskins and Doug Haywick for helping with planting. Funding for this project was

provided through an EPA-Great Lake Restoration Initiative Grant and an NSF- Research

Experience for Undergraduates Grant.

References

Abdel-Aal ESM, Young JC, Rabalski I. 2006. Anthocyanin composition in black, blue, pink,

purple, and red cereal grains. *Journal of Agriculture Food Chemistry* 54(13): 4696-4704.

Asen S, Stewart RN, Norris KH. 1972. Co-pigmentation of anthocyanins in plant tissues and its

effect on color. *Phytochemistry* 11(3): 1139-1144

Barkoudah YI. 1962. Revision of Gypsophila, Bolanthus, Ankyropetalum and Phryna. *Wentia* 9:

1-203.

Bevill RL, Louda SM, Stanforth LM. 1999. Protection from natural enemies in managing rare

plant species. *Conservation Biology* 13(6): 1323-1331.

Chen S, Yao H, Han J, Liu C, Song J, Shi L, Zhu Y, Ma X, Gao T, Pang X, Luo K, Li X, Jai X,

Lin Y, Leon C. 2010. Validation of the ITS2 region as a novel DNA barcode for

identifying medicinal plant species. *PloS One* **5**(1): e8613. doi:

10.1371/journal.pone.0008613L

Clement M, Snell Q, Walker P, Pondosa D, Crandall KA. 2002. TCS: Estimating gene

genealogies. Parallel and Distributed Processing Symposium, International Proceedings,

2, 184.

de Pascual-Teresa S, Santos-Buelga C, Rivas-Gonzalo JC. 2002. LC–MS analysis of

anthocyanins from purple corn cob. *Journal of the Science of Food and*

Agriculture 82(9): 1003-1006.

Darwent AL, Coupland RT. (1966). Life history of *Gypsophila paniculata*. *Weeds* 14(4): 313-

318.

Department of Natural Resources (DNR). 2015. Michigan Invasive Species Grant Program

Handbook. http://www.michigan.gov/documents/dnr/2015-MISGP-

hndbk_491809_7.pdf?source=govdelivery

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Research* 32(5): 1792-1797.

Emery SM, Doran PJ, Legge JT, Kleitch M, Howard S. 2013. Aboveground and belowground

impacts following removal of the invasive species baby's breath (*Gypsophila paniculata*)

on Lake Michigan sand dunes. *Restoration Ecology* 21(4): 506-514.

Fay MF, Swensen SM, Chase MW. 1997. Taxonomic affinities of *Medusagyne oppositifolia*

(Medusagynaceae). *Kew Bulletin* 52(1): 111-120.

Fior S, Karis PO, Casazza G, Minuto L, Sala F. 2006. Molecular phylogeny of the

Caryophyllaceae (Caryophyllales) inferred from chloroplast matK and nuclear rDNA ITS

sequences. *American Journal of Botany* 93(3): 399-411.

Group CPW, Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, van der

Bank M, Chase MW, Cowan RS, Erickson DL, Fazekas AJ. 2009. A DNA barcode for

land plants. *Proceedings of the National Academy of Sciences 106*(31): 12794-12797.

Jukes TH, Cantor CR. 1969. Evolution of protein molecules. In*:* Munrio HN, ed. *Mammalian*

Protein Metabolism. New York: Academic Press. pp. 21-132

Karamanski TJ. 2000. Nationalized Lakeshore: The Creation and Administration of Sleeping

Bear Dunes National Lakeshore. Online Book. National Park Service, Department of the

Interior. http://www.cr.nps.gov/history/online_books/slbe/

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis

version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33(7): 1870-1874.

Lee SC, Wang CH, Yen CE, Chang C. 2017. DNA barcode and identification of the varieties and

provenances of Taiwan's domestic and imported made teas using ribosomal internal

transcribed spacer 2 sequences. *Journal of Food and Drug Analysis* 25(2): 260-274. doi:

10.1016/j.jfda.2016.06.008

Leege LM, Murphy PG. 2001. Ecological effects of the non-native *Pinus nigra* on sand dune

communities. *Canadian Journal of Botany* 79(4): 429-437.

Leng P, Itamura H, Yamamura H, & Deng XM. 2000. Anthocyanin accumulation in apple and

peach shoots during cold acclimation. *Scientia Horticulturae, 83*(1): 43-50.

Newmaster SG, Fazekas AJ, Ragupathy S. 2006. DNA barcoding in land plants: evaluation of

rbcL in a multigene tiered approach. *Botany 84*(3): 335-341.

Pringle JS. 1976. *Gypsophila scorzonerifolia* (Caryophyllaceae), a naturalized species in the

Great Lakes region. *Michigan Botanist* 15: 215-219.

Reznicek AA, Voss EG, Walters BS. February. 2011. *Michigan Flora Online*. University of

Michigan. Web. March 6, 2019. https://michiganflora.net/genus.aspx?id=Gypsophila.

Stevens OA. 1957. Weights of seeds and numbers per plant. *Weeds* 5: 46-55.

Stoeckle MY, Gamble CC, Kirpekar R, Young G, Ahmed S, Little DP. 2011. Commercial teas

highlight plant DNA barcode identification successes and obstacles. *Scientific reports 1*:

42.

Tamura K. 1992. Estimation of the number of nucleotide substitutions when there are strong

transition-transversion and G + C-content biases. *Molecular Biology and Evolution* 9(4):

678-687.

Templeton AR, Crandall KA, Sing CF. 1992. A cladistics analysis of phenotypic association

with haplotypes inferred from restriction endonuclease mapping and DNA sequence data.

III. Cladogram estimation. *Genetics* 132: 619-633.

The Nature Conservancy (TNC). 2014. Lake Michigan Coastal Dunes Restoration Project. 2013

Field Season Report. The Nature Conservancy and Partners, Lansing, Michigan

Voss, EG. 1957. Observations on the Michigan flora—VI. Distribution records of some

angiosperms new, rare, or misinterpreted in the state. *Brittonia 9*(2): 83-101.

Voss EG, Reznicek, AA. 2012. *Field manual of Michigan flora*. Ann Arbor: University of

Michigan Press.

385
Yao H, Song J, Liu C, Luo K, Han J, Li Y, Pang X, Xu H, Zhu Y, Xiao H, & Chen S. 2010. Use

of ITS2 region as the universal DNA barcode for plants and animals. *PloS One 5*(10):

e13102.

Figure Legends

Figure 1: (A) Green-yellow morph, (B) common purple morph and (C) stem of the green-yellow

and purple baby's breath morph found in Sleeping Bear Dunes National Lakeshore.

Figure 2: Phylogenetic analysis of the purple and green-yellow baby's breath color morphs in

relationship to other *Gypsophila* species. All evolutionary histories were inferred using

maximum likelihood methods. (A) Phylogeny based on the rbcL region, (B) phylogeny based on

the matK region, (C) phylogeny based on the ITS2 region, (D) Phylogeny based on rbcL, matK,

and ITS2 combined. For rbcL and ITS2 we used a Jukes Cantor (JC) model of molecular

evolution (Jukes and Cantor, 1969). For matK we used a Tamura 3-parameter (T92) model of

molecular evolution with uniform distribution, and for the combined data set we used the T92

model of molecular evolution with a gamma distribution (Tamura 1992).

Figure 3: A TCS haplotype network based on rbcL, matK and ITS2 combined for the purple and

green-yellow baby's breath color morphs and the *G. paniculata*, *G. elegans*, *G. perfoliata*, and

G. scorzonerifolia reference sequences. The size of the ovals correspond to the haplotype

frequency. The hash marks represent the number of mutations between each haplotype.

Figure 4: (A) Green-yellow morph, and (B) common purple morph after 5-months in the GVSU

greenhouse. Note the similarity in taproot structure between the two plants.

412
413

A.

B.

C.

A. rbcL

B. matK

C. ITS2

D. All Sequences