1 Identification and characterization of *Gypsophila paniculata* color morphs in Sleeping Bear
2 Dunes National Lakeshore, MI, USA.
3

4 Marisa L. Yang[1], Emma Rice[2], Hailee Leimbach-Maus[2], and Charlyn G. Partridge[2]
5
6 [1]Environmental Science Policy, and Management, University of California, Berkeley, 260
7 Mulford Hall, Berkeley, CA 94720, email: marisayang79@gmail.com
8 [2]Annis Water Resources Institute, 740 W. Shoreline Dr., Muskegon, MI 49441, email:
9 riceemm@mail.gvsu.edu; haileepavischi@gmail.com
10
11 Corresponding author: Charlyn Partridge, Annis Water Resources Institute, Grand Valley
12 State University, 740 W. Shoreline Dr., Muskegon, MI; 49441, email: partridc@gvsu.edu

13
14
15 **Abstract**
16

Gypsophila paniculata (baby's breath) is an invasive species found throughout much of the

northwest United States and western Canada. Two distinct color morphs of baby's breath

were identified in areas within the coastal dunes along eastern Lake Michigan. The more

common morph produces stems that are purple in color (purple morph), while the atypical

morph has stems that are green-yellow (green-yellow morph). The purpose of this study

was to characterize these morphs and determine if they are genetically distinct species in

order to assess whether alternative management strategies should be employed to control

these populations. We sequenced two chloroplast regions, RbcL and MatK, and one nuclear

region, ITS2, from the purple morphs and green-yellow morphs collected from Sleeping

Bear Dunes National Lakeshore, MI, USA. Sequences were aligned with reference *G.*

paniculata and *G. elegans* sequences obtained from GenBank and the Barcode of Life

(BOLD) databases. Phylogenetic analyses suggest that the two color morphs belong *to G.*

paniculata and do not appear to be distinct genetic strains. We propose that current

management practices continue to treat the two color morphs in a similar manner in terms

of removal to prevent the further spread of this species.

Introduction

The Great Lakes sand dunes comprise the most extensive freshwater dune complex in the

world, stretching over 1,000 km^2 in Michigan alone. Within northwest Michigan, the sand dunes

ecosystem is vital both environmentally and economically. It is home to a number of threatened

and endangered species, including piping plover (*Charadrius melodus*) and Pitcher's thistle

(*Cirsium pitcheri*). The colonization of invasive species into this region can significantly alter the

biological composition of native communities (Olson 1958, Leege and Murphy 2001, Emery et

al. 2013). One invasive species of significant concern is the perennial baby's breath (*Gypsophila*

paniculata). In 2015, baby's breath was listed by the Department of Natural Resources (DNR) as

a "priority" invasive species for detection and control in Michigan's northern lower peninsula

(DNR 2015). Since its colonization in the region it has spread along a 260 km stretch of the

Michigan shoreline. Baby's breath produces a large primary taproot that can extend down to 4

meters in depth, which likely helps it outcompete native vegetation for limited resources

(Darwent and Coupland 1966, Karamanski 2000). The presence of baby's breath in this system is

associated with a reduction in the number of threatened Pitcher's thistle and an increase in the

number of other non-native species (Emery et al. 2013). In addition, while many of the

vulnerable and endangered plant species in these areas are seed limited, (e.g., Pitcher's thistle

produces approximately 50-300 seeds per plant (Bevill et al. 1999)), baby's breath can produce

up to 14,000 seeds per plant annually (Stevens 1957), effectively outcompeting native species in

terms of overall yield. This has led to baby's breath composing approximately 50-80% of the

ground cover in some areas (Karamanski 2000, Emery et al. 2013).

One concern with current management efforts is that there is some anecdotal evidence

that there may be a new baby's breath variant (morph) within the Michigan dune system. In 2011

and 2012 The Nature Conservancy (TNC) removal crews reported baby's breath plants with

different character traits than what is commonly observed (TNC 2014). The atypical morph has

stems and leaves that are lighter in color and more yellow than the common purple morph

(Figure 1a-c). The purple morph has a thick taproot (4-7 cm in diameter) just below the caudex

that remains unbranched for approximately 60 – 100 cm (Darwent & Coupland 1966). Severing

just below the intersection of the caudex and the taproot is where manual removal efforts target

to limit regrowth. However, the atypical green-yellow morph's root system was reported by TNC

to be more diffuse, making it harder to identify a primary taproot and thus, harder to sever

without the potential for regrowth (TNC 2014). Currently, these green-yellow morphs are treated

with herbicide application (glyphosate) when observed; however, if this morph continues to

spread into areas where threatened or endangered species are present, removal methods will be a

primary concern. If the green-yellow morph and the purple morph represent two discrete species,

then alternative management strategies may need to be considered for these populations.

One of the first steps toward adapting current management strategies for this population

is to identify whether the green-yellow morph is genetically distinct from the purple morph.

While *G. paniculata* is the dominant invasive baby's breath species in northwest Michigan, there

are a number of distinct *G. paniculata* commercial strains that could potentially invade these

areas considering seeds are commonly sold in stores throughout the region (personal

observation). In addition, other baby's breath species, including *G. elegans* are sold in this region

in commercial wildflower packets. *G. elegans* is an annual species that has a smaller taproot

compared to *G. paniculata,* and its coloration is similar to that of the green-yellow morph. Thus,

the goal of this work was to use traditional 'barcode' genes for plants to characterize the genetic

relationship between the green-yellow morph and the common purple morph.

81
82 **Methods and materials**

83
84 *DNA Extraction, Amplification, and Sequencing*

85
We collected leaf tissue from 1 green-yellow morph and 16 purple morphs in 2016 and

an additional 15 green-yellow morphs in 2017 from Sleeping Bear Dunes National Lakeshore

(SBDNL), Empire, MI, USA (specifically: 44.884941 N, 86.062111 W and 44.875302 N,

86.056821 W). Leaf tissue was dried in silica gel until DNA extractions could take place. DNA

was extracted using Qiagen DNeasy Plant Mini Kits (Qiagen, Hilden, Germany). After

extraction, the DNA samples were placed through Zymo OneStep PCR inhibitor removal

columns (Zymo, Irvine, CA) to remove any secondary metabolites that might inhibit PCR

amplification. The DNA for each sample was then quantified using a NanoDrop 2000

(ThermoFisher, Waltham, MA).

The DNA of both green-yellow morphs and purple morphs was amplified at three

proposed 'barcoding' genes for plants: large subunit of the ribulose-bisphosphate carboxylase

gene (RbcL), maturase K (MatK), and internal transcribed spacer 2 (ITS2). The RbcL region was

amplified using RbcL 1F and RbcL 724R primers (Chen et al. 2010), MatK was amplified using

MatK 390F and MatK 1440R primers (Fior et al. 2006), and the ITS2 region was amplified using

ITS2 2SF and ITS2 S3R primers (Chen et al. 2010). Individual PCR reactions for all loci

consisted of 1X Taq Buffer, 2.0 mM $MgCl_2$, 0.3 μM dNTP, 0.08 mg/mL BSA, 0.4 μM forward

and reverse primers, and 0.5 units of Taq polymerase in a 20 μL reaction volume. The thermal

cycle protocols consisted of the following: for RbcL, an initial denaturing step of 95ºC for 2

minutes, followed by 35 cycles of 94ºC for 1 minute, 55ºC for 30 seconds, and 72ºC for 1

minute. A final elongation step was performed at 72ºC for 7 minutes. For MatK, the thermal

profile consisted of 26 cycles of 94ºC for 1 minute, 48ºC for 30 seconds, and 72ºC for 1 minute,

followed by a final elongation step at 72ºC for 7 minutes. For ITS2, an initial denaturing step of

95ºC for 2 minutes was applied, followed by 35 cycles of 95ºC for 30 seconds, 50ºC for 30

seconds, 72ºC for 1.5 minutes, and a final elongation step of 72ºC for 8 minutes. Successful

amplification was checked by running the PCR product on a 2% agarose gel stained with

ethidium bromide. PCR reactions were then cleaned using ExoSAP-IT PCR Product Cleanup

Reagent (ThermoFisher, Waltham, MA). Sequencing reactions were performed for both forward

and reverse primers for each of the three genes. Sequencing reactions were cleaned using a

Sephadex column (GE Healthcare Life Science, Marlborough, MA) and sequenced on an ABI

Genetic BioAnalyzer 3130xl (Applied Biosystems, Foster City, CA). Out of the 16 green-yellow

morphs a total of 13 were successfully sequenced for RbcL, 13 were successfully sequenced for

MatK, and 14 were successfully sequenced for ITS2. For the purple morphs a total of 15, 12, and

15 individuals were successfully sequenced for RbcL, MatK and ITS2, respectively.

119   *G. paniculata* and *G. elegans* reference sequences for RbcL, MatK, and ITS2 were

120   downloaded either from GenBank (https://www.ncbi.nlm.nih.gov/genbank/) or the Barcode of

121   Life Database (BOLD) (http://www.barcodeoflife.org). Their accession numbers and sequences

122   are provided in Supplemental Table 1. All fasta files corresponding to these data will be

123   deposited in the Dryad database and sequences will be submitted to GenBank.

124   *Alignment and Phylogenetic Analysis*

125   All success sequences from our field samples, as well as sequences for *G. elegans*

and *G. paniculata* obtained from GenBank or BOLD, were imported into the program

MEGA7 (version 7.0.14) (Kumar et al. 2016) and sequences for each of the three genes

were aligned both individually and with all sequences combined using Muscle (Edgar

2004). The total number of base pairs aligned and analyzed for each region included: 516

base pair (bp) for RbcL, 720 bp for MatK, 264 bp for ITS2, and 1498 bp for the three regions

combined. All alignment parameters were kept at their default settings. Once aligned, we used

MEGA7 to identify the most appropriate substitution model (RbcL: Jukes-Cantor, MatK:

Tamura 3-parameter, ITS2: Jukes-Cantor, all genes combined: Tamura 3-parameter). We then

created phylogenetic trees using a maximum-likelihood (ML) approach with 500-replicated

bootstrap analyzes, as well as using neighbor joining, and parsimony models.

Results and Discussion

Our results indicate that the green-yellow morph identified in SBDNL is not a genetically

distinct species from the common purple *G. paniculata* found throughout SBDNL. The RbcL,

MatK, ITS2, and combined dataset showed similar patterns with both the green-yellow morphs

and the purple morphs clustering together with a *G. paniculata* reference sequence, and

separately from *G. elegans* reference sequences (Figure 2a-d; Supplemental Figure 1). This

suggests that both color morphs belong to *G. paniculata*.

Of the regions analyzed for the green-yellow morphs, purple morphs, and reference

sequences, RbcL was the most conserved sequence with an overall mean genetic distance (d) =

0.001, followed by MatK ($d = 0.002$) and ITS2 ($d = 0.018$). For the ITS2 region, there were 6

purple *G. paniculata* morphs and one green-yellow morph that clustered together inside the *G.*

paniculata branch (Figure 2a & 2d). Further examination of the chromatograms for these

individuals show that they are likely heterozygous at position 195 of our aligned sequence and

amplification bias of the 'A' SNP over the allele containing the 'G' SNP is driving this pattern.

The phylogeny constructed from the RbcL sequence showed one *G. paniculata* reference

sequence (SDH831-14) grouping together with *G. elegans*. All reference sequences for RbcL

were obtained from the BOLD database, and it is possible that this specific individual was

misidentified. Further examination of this specimen from the herbarium image posted to BOLD

indicates a small taproot compared to the upper foliage

(http://www.boldsystems.org/pics/SDH/CCDB-24938-B09_h235508%2B1424896806.jpg),

which is one characteristic of the annual *G. elegans*. While there is also sequence data for both

the ITS2 and MatK regions for this specimen in BOLD, these were not used for this dataset due

to higher levels of sequence fragmentation compared to our samples. However, both the ITS2

and MatK sequences from this *G. paniculata* reference sequence aligned most closely to *G.*

elegans in BLAST.

While our data suggest that both the purple and green-yellow morphs are *G. paniculata*,

whether the green-yellow morph is a different commercial genetic strain is still unclear. RbcL,

MatK, and ITS2 are common 'barcode' genes used to delineate plant species, but their ability to

distinguish genetically distinct strains of the same species can be limited. RbcL and MatK are

highly conserved and would likely not distinguish between different cultivars; however, ITS2 is

more variable and has been used to distinguish between different varieties of domestic and

imported teas (*Camellia sinensis*) (Lee et al. 2017). In our dataset, the purple and green-yellow

morphs from SBDNL grouped separately from the *G. paniculata* reference sequence, but within

the same clade. The *G. paniculata* reference sequence for ITS2 was obtained from GenBank and

is part of an unpublished study, so we do not have information regarding where this sample was

collected. However, given the distinct grouping between the *G. paniculata* reference sequence

and our samples, ITS2 may be a promising region to assess the ability to distinguish among

different commercial strains or regional varieties of *G. paniculata*. If this is the case, then our

data would suggest that the purple and yellow-green morphs not only belong to the same species

(*G. paniculata)*, but are likely derived from the same or similar initial source populations.

The mechanism driving the color difference between the purple and green-morphs is

currently unknown. Within SBDNL, the purple morph is the most common form, with green-

yellow individuals found interspersed in a couple of locations throughout the dunes. The largest

observed group of green-yellow morphs consists of a few hundred plants clumped within

approximately an acre-sized area and interspersed throughout large groups of purple morphs.

Occasionally, small pockets of green-yellow individuals have also been found together in other

areas of SBDNL and Zetterberg Preserve (approximately 38 km from SBDNL). Based upon the

dispersal patterns of the two morphs throughout the dunes, the color difference observed does

not appear to be solely environmentally driven, and likely has a genetic component. Potential

candidate genes could include those involved in the anthocyanin pathway, which influences red –

purple coloration in a number of plants (Asen et al. 1972, de Pascual-Teresa et al. 2002, Abdel-

Aal et al. 2006, Knievel et al. 2009). Further work will begin to elucidate the specific mechanism

influencing these color difference in invasive *G. paniculata* populations, as well as to explore

whether this color difference drives functional differences between the morphs.

In terms of management strategies, the purple and green-yellow morph are not distinct

species and are likely derived from the same genetic strain, so there does not appear to be a need

to manage these two morphs differently. One concern with the green-yellow morph initially

noted by TNC removal crews was that the taproot tended to be more diffuse than the purple

morph potentially making manual removal of these plants less effective. We are currently

examining root growth between the green-yellow and purple morphs under controlled conditions,

however, our lab's personal observations in the field have not found any indication that large

differences in root structure occur between mature plants of the two color morphs. Therefore,

current management approaches for these populations should be maintained to control the further

spread of *G. paniculata* throughout the Michigan coastal dune system.

Acknowledgements

This project was funded through an EPA-Great Lakes Restoration Initiative Grant and through

an NSF-REU grant. We would like to thank Shaun Howard from The Nature Conservancy for

his help in identifying the green-yellow morph and Benjamin Giffin for his help with sequencing

analysis. We would also like to thank Kurt Thompson and Brayden Partridge for assisting with

the figure construction.

207
208   **References**
209
210   Abdel-Aal, E.S.M., Young, J.C., and Rabalski, I. 2006. Anthocyanin composition in black, blue,
211   pink, purple, and red cereal grains. J. Agric. Food Chem. **54**(13): 4696-4704.
212
213   Asen, S., Stewart, R.N., and Norris, K.H. 1972. Co-pigmentation of anthocyanins in plant
214   tissues and its effect on color. Phytochemistry **11**(3): 1139-1144.
215
Bevill, R.L., Louda, S.M., and Stanforth, L M. 1999. Protection from Natural Enemies in
managing rare plant species. Conserv. Biol. **13**(6): 1323-1331.

Chen, S., Yao, H., Han, J., Liu, C., Song, J., Shi, L., Zhu, Y., Ma, X., Gao, T., Pang, X., Luo, K., Li, X.,
Jai, X., Lin, Y., and Leon, C. 2010. Validation of the ITS2 region as a novel DNA
barcode for identifying medicinal plant species. PloS One **5**(1): e8613. doi:
10.1371/journal.pone.0008613L

de Pascual-Teresa, S., Santos-Buelga, C., and Rivas-Gonzalo, J.C. 2002. LC–MS analysis of
anthocyanins from purple corn cob. J. Sci. Food Agric. **82**(9): 1003-1006.
224

225   Darwent, A.L., and Coupland, R.T. 1966. Life history of *Gypsophila paniculata*. Weeds **14**(4):
226   313-318.
227
228   Department of Natural Resources (DNR). 2015. Michigan Invasive Species Grant Program
229   Handbook. http://www.michigan.gov/documents/dnr/2015-MISGP-
230   hndbk_491809_7.pdf?source=govdelivery
231
232   Edgar, R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
233   throughput. Nucleic Acids Res. **32**(5): 1792-1797.
234
235   Emery, S.M., Doran, P.J., Legge, J.T., Kleitch, M., and Howard, S. 2013. Aboveground and
236   belowground impacts following removal of the invasive species baby's breath
237   (*Gypsophila paniculata*) on Lake Michigan sand dunes. Restoration Ecol. **21**(4): 506-
238   514.
239
240   Fior, S., Karis, P. O., Casazza, G., Minuto, L., and Sala, F. 2006. Molecular phylogeny of the
241   Caryophyllaceae (Caryophyllales) inferred from chloroplast matK and nuclear rDNA
242   ITS sequences. Am. J. Bot. **93**(3): 399-411.
243
244   Jukes, T.H. and Cantor, C.R. 1969. Evolution of protein molecules. *In* Mammalian Protein
245   Metabolism. Edited by H. N. Munro. Academic Press, New York. pp. 21-132
246
247   Karamanski, T.J. 2000. Nationalized Lakeshore: The Creation and Administration of
248   Sleeping Bear Dunes National Lakeshore. Online Book. National Park Service,
249   Department of the Interior. http://www.cr.nps.gov/history/online_books/slbe/
250
251   Kumar, S., Stecher, G., & Tamura, K. 2016. MEGA7: Molecular Evolutionary Genetics
252   Analysis version 7.0 for bigger datasets. Mol. Biol. Evol. **33**(7): 1870-1874.
253
254   Lee, S. C., Wang, C. H., Yen, C. E., and Chang, C. 2017. DNA barcode and identification of the
255   varieties and provenances of Taiwan's domestic and imported made teas using
256   ribosomal internal transcribed spacer 2 sequences. J. Food Drug Anal. **25**(2): 260-
257   274. doi: 10.1016/j.jfda.2016.06.008
258
259   Leege, L. M., Murphy, P. G. 2001. Ecological effects of the non-native *Pinus nigra* on sand
260   dune communities. Can. J. Bot. **79**(4): 429-437.
261
262   Stevens, O. A. 1957. Weights of seeds and numbers per plant. Weeds **5**: 46-55.
263
264   Tamura K. 1992. Estimation of the number of nucleotide substitutions when there are
265   strong transition-transversion and G + C-content biases. Mol. Biol. Evol. **9**(4): 678-
266   687.
267
268   The Nature Conservancy (TNC). 2014. Lake Michigan Coastal Dunes Restoration Project.
269   2013 Field Season Report. The Nature Conservancy and Partners, Lansing, Michigan
270

271
272   Figure Legends:
273
274   Figure 1: (A) Green-yellow morph, (B) common purple morph and (C) Stem of the green-
275   yellow and purple baby's breath morph found in Sleeping Bear Dunes National Lakeshore.
276
Figure 2: Phylogenetic construction of the purple and green-yellow baby's breath color
morphs in relationship to *G. paniculata* and *G. elegans* reference sequences. All
evolutionary histories were inferred using maximum likelihood methods. (A) Phylogeny
based on RbcL, MatK, and ITS2 combined, (B) phylogeny based on the RbcL region, (C)
phylogeny based on the MatK region, (D) phylogeny based on the ITS2 region. For RbcL and
ITS2 we used a Jukes Cantor model of molecular evolution (Jukes and Cantor, 1969). For
MatK and the combined data set we used a Tamura 3-parameter model of molecular
evolution (Tamura 1992).
285
286
287

288
289
290   Figure 1

292
293   Figure 2

294
295