N^6-methyladenosine modification and the YTHDF2 reader protein play cell type
specific roles in lytic viral gene expression during Kaposi's sarcoma-associated
herpesvirus infection

Charles Hesser[1], John Karijolich[2,3], Dan Dominissini[4,5,6], Chuan He[4,7], and Britt Glaunsinger[1,2,7]

[1]Department of Molecular and Cell Biology, [2]Department of Plant & Microbial Biology,
University of California Berkeley; [3]Department of Pathology, Microbiology, and Immunology,
Vanderbilt University Medical Center, Nashville, TN 37232-2363, [4]Department of Chemistry,
Department of Biochemistry and Molecular Biology, and Institute for Biophysical Dynamics, The
University of Chicago, Chicago, Illinois, USA; [5]Chaim Sheba Medical Center, Tel Hashomer
52621, Israel; [6]Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel, [7]Howard Hughes
Medical Institute

*Corresponding author: glaunsinger@berkeley.edu

Abstract

Methylation at the N^6 position of adenosine (m^{6}A) is a highly prevalent and reversible

modification within eukaryotic mRNAs that has been linked to many stages of RNA processing

and fate. Recent studies suggest that m^{6}A deposition and proteins involved in the m^{6}A pathway

play a diverse set of roles in either restricting or modulating the lifecycles of select viruses.

Here, we report that m^{6}A levels are significantly increased in cells infected with the oncogenic

human DNA virus Kaposi's sarcoma-associated herpesvirus (KSHV). Transcriptome-wide m^{6}A-

sequencing of the KSHV-positive renal carcinoma cell line iSLK.219 during lytic reactivation

revealed the presence of m^{6}A across multiple kinetic classes of viral transcripts, and a

concomitant decrease in m^{6}A levels across much of the host transcriptome. However, we found

that depletion of the m^{6}A machinery had differential pro- and anti-viral impacts on viral gene

expression depending on the cell-type analyzed. In iSLK.219 cells the pathway functioned in a

pro-viral manner, as depletion of the m^{6}A writer METTL3 and the reader YTHDF2 significantly

impaired viral lytic cycle progression and virion production. This defect could be attributed to

their roles in the post-transcriptional accumulation of the major viral lytic transactivator ORF50,

which is m^{6}A modified. In contrast, although the ORF50 mRNA was also m^{6}A modified in KSHV

infected B cells, ORF50 expression was instead increased upon depletion of METTL3, YTHDF1,

or, to a lesser extent, YTHDF2. These results highlight that the m^{6}A pathway is centrally

involved in regulating KSHV gene expression, and underscore how the outcome of this

dynamically regulated modification can vary significantly between cell types.

Author Summary

In addition to its roles in regulating cellular RNA fate, methylation at the N^6 position of

adenosine (m^6A) of mRNA has recently emerged as a mechanism for regulating viral infection.

While it has been known for over 40 years that the mRNA of nuclear replicating DNA viruses

contain m^6A, only recently have studies began to examine the distribution of this modification

across viral transcripts, as well as characterize its functional impact upon viral lifecycles. Here,

we apply m^6A-sequencing to map the location of m^6A modifications throughout the

transcriptome of the oncogenic human DNA virus Kaposi's sarcoma-associated herpesvirus

(KSHV). We show that the m^6A machinery functions in a cell type specific manner to either

promote or inhibit KSHV gene expression. Thus, the KSHV lifecycle is impacted by the m^6A

pathway, but the functional outcome may depend on cell lineage specific differences in m^6A-

based regulation.

Introduction

The addition of chemical modifications is critical to many steps of mRNA processing and

the regulation of mRNA fate. There are more than 100 different RNA modifications, but the

most abundant internal modification of eukaryotic mRNAs is N^6-methyladenosine (m^6A), which

impacts nearly every stage of the posttranscriptional mRNA lifecycle from splicing through

translation and decay [1-6]. The breadth of impacts ascribed to the m^6A mark can be attributed

to its creation of new platforms for protein recognition, in part via local changes to the RNA

structure [4,7-12]. The reversibility of m^{6}A deposition through the activity of demethylases

termed erasers adds a further layer of complexity by enabling dynamic regulation, for example

during developmental transitions and stress [1,4,5,13-15]. Deposition of m^{6}A occurs co- or post-

transcriptionally through a complex of proteins with methyltransferase activity known as

writers, which include the catalytic subunit METTL3 and cofactors such as Mettl14 and WTAP

[1,4,14,16,17]. The modification is then functionally 'interpreted' through the selective binding

of m^{6}A reader proteins, whose interactions with the mRNA promote distinct fates.

The best-characterized m^{6}A readers are the YTH domain proteins. The nuclear YTHDC1

reader promotes exon inclusion [6], whereupon m^{6}A-containing mRNA fate is guided in the

cytoplasm by the YTHDF1-3 readers. Generally speaking, YTHDF1 directs mRNAs with 3' UTR

m^{6}A modifications to promote translation [3], whereas YTHDF2 recruits the CCR4-NOT

deadenylase complex to promote mRNA decay [18]. YTHDF3 has been proposed to serve as a

co-factor to potentiate the effects of YTHDF1 and 2 [3,19,20]. Although the individual effects of

YTHDF1 and 2 seem opposing, the YTHDF proteins may coordinate to promote accelerated

mRNA processing during developmental transitions and cellular stress [1]. YTHDC2, the fifth

member of the YTH family proteins, was recently shown to play critical roles in mammalian

spermatogenesis through regulating translation efficiency of target transcripts [21]. Additional

examples of distinct functions for m^{6}A readers under specific contexts such as heat shock are

rapidly emerging [13].

Given the prevalence of the m^{6}A modification on cellular mRNAs, it is not surprising that

a number of viruses have been shown to contain m^{6}A in their RNA [22-29]. Indeed, a potential

viral benefit could be a less robust innate antiviral immune response, as m^{6}A modification of in

vitro synthesized RNAs diminishes recognition by immune sensors such as TLR3 and RIG-I

[30,31]. That said, the functional consequences of viral mRNA modification appear diverse and

include both pro- and anti-viral roles. In the case of Influenza A, a negative sense ssRNA virus,

m^6A and the reader YTHDF2 have been shown to promote viral replication [32]. Furthermore,

multiple studies have mapped the sites of m^6A modification in the human immunodeficiency

virus (HIV) genome, and shown that it promotes the nuclear export of HIV mRNA as well as viral

protein synthesis and RNA replication [24,26,28]. Roles for the YTHDF proteins in during HIV

infection remain varied however, as Tirumuru and colleagues propose they function in an anti-

viral context by binding viral RNA and inhibiting reverse transcription, while Kennedy and

colleagues observe they enhance HIV replication and viral titers [24,28]. A more consistently

anti-viral role for the m^6A pathway has been described for the *Flaviviridae*, whose (+) RNA

genomes are replicated exclusively in the cytoplasm and contain multiple m^6A sites in their

genomic RNA [23,25]. An elegant study by Horner and colleagues showed that depletion of

m^6A writers and readers or the introduction of m^6A-abrogating mutations in the viral E1 gene

all selectively inhibit hepatitis C virus (HCV) assembly [23]. Similarly, depletion of METTL3 or

Mettl14 enhances Zika virion production [25].

Despite the fact that m^6A modification of DNA viruses was first reported more than 40

years ago for simian virus 40, herpes simplex virus type 1, and adenovirus type 2, roles for the

modification in these and other DNA viruses remain largely unexplored [33-37]. Unlike most

RNA viruses, with few exceptions DNA viruses replicate in the nucleus and rely on the cellular

transcription and RNA processing machinery, indicating their gene expression strategies are

likely interwoven with the m^6A pathway. Indeed, it was recently shown that the nuclear reader

YTHDC1 potentiates viral mRNA splicing during lytic infection with Kaposi's sarcoma-associated

herpesvirus (KSHV) [38]. However, roles for the cytoplasmic m^6A readers during DNA virus

infection, as well as a transcriptome-level understanding of how DNA viruses incorporate m^6A

and influence the spectrum of m^6A deposition on cellular mRNAs remain key open questions.

Here, we sought to address these questions by measuring and mapping the abundance

of m^6A marks across the viral and host transcriptome during lytic KSHV infection. This

gammaherpesvirus remains the leading etiologic agent of cancer in AIDS patients, in addition to

causing the lymphoproliferative disorders multicentric Castleman's disease and primary

effusion lymphoma. The default state for KSHV in cultured cells is latency, although in select

cell types the virus can be reactivated to engage in lytic replication, which involves a temporally

ordered cascade of gene expression. We reveal that m^6A levels are significantly increased upon

KSHV reactivation, which is due to a combination of m^6A deposition across multiple kinetic

classes of viral transcripts and a concomitant decrease in m^6A levels across much of the host

transcriptome. Depletion of m^6A writer and cytoplasmic reader proteins either moderately or

severely impaired viral lytic cycle progression in the KSHV iSLK.129 reactivation model,

suggesting this pathway potentiates the KSHV lytic cycle. Interestingly, however, the roles for

the m^6A writer and readers shifted to instead display neutral or anti-viral activity in the TREX-

BCBL1 reactivation model. These findings thus demonstrate that while KSHV mRNAs are

marked by m^6A, the functional consequences of this mark can vary significantly depending on

cell context, reinforcing both the functional complexity and dynamic influence of m^6A.

Results

KSHV mRNA contains m^{6}A modifications

Epitranscriptome mapping has revealed significant roles for the m^{6}A pathway in the lifecycle and regulation of several RNA viruses, but similar global analyses have yet to be performed for a DNA virus. Given that herpesviral mRNAs are transcribed and processed in the nucleus using the cellular RNA biogenesis machinery, we hypothesized that these viruses would engage the m^{6}A pathway. We therefore first quantified how KSHV reactivation impacted total cellular m^{6}A levels in the KSHV-positive renal carcinoma cell line iSLK.219 (**Fig 1**). These cells are a widely used model for studying viral lytic events, as they stably express the KSHV genome in a tightly latent state but harbor a doxycycline (dox)-inducible version of the major viral lytic transactivator ORF50 (also known as RTA) that enables efficient entry into the lytic cycle [39,40]. Polyadenylated (polyA+) RNA was enriched from untreated (latent) or dox-reactivated iSLK.219 cells and the levels of m^{6}A were quantitatively analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS) (**Fig 1A**). Indeed, we observed a three-fold increase in total m^{6}A levels upon KSHV lytic reactivation, suggesting that m^{6}A deposition significantly increased during viral replication (**Fig 1B**).

We next sought to discern whether the increase in m^{6}A during the KSHV lytic cycle favors host or viral mRNAs using high throughput m^{6}A RNA sequencing (m^{6}Aseq) [41]. This technique can reveal both the relative abundance and general location of m^{6}A in KSHV and cellular mRNA. Total m^{6}A containing RNA was immunoprecipitated from 2 biological replicates of latent or lytically reactivated iSLK.219 cells using an m^{6}A-specific antibody. DNase-treated

total mRNA was fragmented to lengths of 100 nt prior to immunoprecipitation and then

subjected to m^{6}A-seq. Total RNA-seq was run in parallel for each sample, allowing the degree

of m^{6}A modification to be normalized with respect to transcript abundance because the levels

of many transcripts change upon viral lytic reactivation. Peaks with a fold-change four or higher

(FC>4) and a false discovery rate of 5% or lower (FDR>5%) in both replicates were considered

significant, although it is possible that additional transcripts detectably modified to lower levels

or in a more dynamic manner may also be functionally regulated by m^{6}A (complete list of viral

peaks with FC>2 in **S1 Table**). In lytically reactivated samples, 10 transcripts comprising genes of

immediate early, early, and late kinetic classes displayed significant m^{6}A modification in both

replicates (**Fig 2A** and **S1 Fig**). Within these KSHV mRNAs, m^{6}A peaks were detected primarily in

coding regions, although in some cases the location of a peak in a coding region overlaps with a

UTR (**S1 Fig**). Furthermore, all but one peak contains at least one instance of the GG(m^{6}A)C

consensus sequence. While many of the modified viral transcripts contained only one m^{6}A

peak, multiple peaks were found in certain transcripts, including the major lytic transactivator

ORF50 (**Fig 2B**). Of note, exon2 of ORF50 contained one m^{6}A peak of FC>4 in replicate one, and

three m^{6}A peaks in replicate two, each of which have at least one m^{6}A consensus motif, further

increasing confidence that these peaks accurately represent m^{6}A modified sites. Furthermore,

the viral ncRNA PAN, which has been reported to comprise over 80% of nuclear PolyA+ RNA

during lytic reactivation [42], contains FC>4 peaks in both replicates. Modification of PAN likely

accounts for the marked three-fold increase in cellular m^{6}A content observed upon lytic

reactivation (Fig 1B). As anticipated given the restricted viral gene expression profile during

latency, unreactivated samples had many fewer m^{6}A containing viral mRNAs, with the only

FC>4 peaks occurring in both replicates located in ORF4. Although ORF4 is a lytic transcript, its

coding region overlaps with the 3' UTR of K1, which is expressed during both the latent and lytic

phases of the viral lifecycle (**Fig 2A**, **S1 Fig**) [43,44].

To validate the m^6A-seq results, we performed m^6A RNA immunoprecipitation (RIP)

followed by quantitative real-time PCR (RT-qPCR) on six of the viral transcripts predicted to be

m^6A modified from the m^6A-seq data. This technique allows determination of the relative level

of m^6A content in a given transcript compared to an unmodified transcript. As controls, we

included primers for the cellular GAPDH transcript, which is known not to be m^6A modified, and

the Dicer transcript, which is m^6A modified [41]. The m^6A RIP RT-qPCR confirmed modification

of the vIL-6, K1, ORF50, ORF57 and PAN viral transcripts, in agreement with m^6A-seq results (**Fig**

2C). In summary, we found m^6A modification in approximately one third of KSHV transcripts

upon lytic reactivation, consistent with the hypothesis that this pathway contributes to KSHV

gene expression.

We next compared the distribution of m^6A peaks in host mRNAs from unreactivated

versus reactivated cells to assess whether lytic KSHV infection altered the m^6A profile of cellular

transcripts. Analyzing the two independent replicates for each condition, we found an average

of 14,092 m^6A modification sites (FC>4 and FDR>5%) in host transcripts pre-reactivation,

compared to 10,537 peaks post-reactivation (**Fig 2D** and **S2 Table**). We observed that this >25%

decrease in m^6A deposition on cellular mRNA encompassed a wide spectrum of transcripts, and

no notable patterns were apparent by GO term analysis for functional categories enriched in

the altered population. Thus, while the functional impact of the altered host m^6A profile

remains unresolved, the observation that KSHV lytic infection increased the level of m^6A in total

poly A+ RNA despite decreasing its presence in cellular mRNA implies that m^{6}A deposition

during infection favors viral transcripts.

m^{6}A and the reader YTHDF2 mediate viral gene expression and virion production in iSLK.219

cells

Given the significant deposition of m^{6}A across KSHV transcripts, we reasoned that m^{6}A

might play an important role in potentiating the viral lifecycle. We therefore examined the

effect of depleting the m^{6}A writers and readers on KSHV virion production using a supernatant

transfer assay. The KSHV genome in iSLK.219 cells contains a constitutively expressed version

of GFP, which allows for fluorescence-based monitoring of infection by progeny virions. We

performed siRNA-mediated knockdown of METTL3, the catalytic subunit responsible for m^{6}A

deposition, as well as the m^{6}A readers YTHDF 1, 2 and 3 (**Fig 3A**). Cells were then treated with

dox and sodium butyrate to induce lytic reactivation for 72 h, whereupon supernatants were

collected and used to infect 293T recipient cells. The number of GFP positive 293T cells at 24

hpi was measured by flow cytometry (**Fig 3B**). Notably, for virus generated from METTL3

depleted cells, only 7% of recipient cells were infected compared to 82% for virus generated

during treatment with a control siRNA (**Fig 3B**). YTHDF2 depletion caused an even more

pronounced defect, resulting in a near absence of virion production. In contrast, YTHDF3

knockdown resulted in only a modest defect and virion production was unaffected by YTHDF1

knockdown (**Fig 3B**). Thus, the m^{6}A writer METTL3 and the reader YTHDF2 play important roles

in driving KSHV infectious virion production in iSLK.219 cells.

We then sought to determine the stage of the viral lifecycle impacted by the m^{6}A

pathway by measuring the impact of writer and reader depletion on the abundance of viral

mRNAs of different kinetic classes. First, levels of representative immediate early, delayed

early, and late viral mRNAs were measured by RT-qPCR following lytic reactivation for 72 hours.

Of note, all of these transcripts contained at least one m^{6}A peak. METTL3 depletion did not

appear to impact accumulation of the ORF50 immediate early mRNA at this time point, but

resulted in a significant defect in accumulation of the delayed early transcript ORF57 and late

gene transcript K8.1 (**Fig 3C**). Consistent with the virion production data, we observed a striking

defect in the accumulation of each of the viral transcripts upon YTHDF2 depletion, suggesting

that this protein is essential for lytic KSHV gene expression beginning at the immediate early

stage (**Fig 3C**). In contrast, viral gene expression was either unaltered or modestly increased

upon depletion of YTHDF1 or YTHDF3 (**Fig 3C**). These results for ORF50 and ORF59 were

confirmed at the protein level by Western blot, showing that depletion of YTHDF2 but not the

other m^{6}A pathway components resulted in a near total inhibition of viral protein expression at

this late stage in the viral lifecycle (**Fig 3D**). We noted that although depletion of METTL3 did

not significantly decrease the level of ORF50 mRNA, levels of METTL3 knockdown in excess of

80% have only been reported to reduce m^{6}A levels in Poly A RNA by 20-30% [17]. Thus, at least

some fraction of ORF50 (and other) transcripts may still be m^{6}A modified due to residual

enzyme activity upon depletion of METTL3.

In agreement with the above findings, we also observed that iSLK.219 cells depleted of

METTL3 and YTHDF2 displayed a prominent defect in viral reactivation, as measured by

expression of red fluorescent protein (RFP) driven by the PAN lytic cycle promoter from the

viral genome (**Fig 3E**). Similarly, ORF50 protein production was also markedly reduced upon

YTHDF2 depletion at the 24 h time point, which represents the early phase of the lytic cycle for

cells reactivated with dox alone (**Fig 3F**). Collectively, these results suggest that m^{6}A

modification is integral to the KSHV lifecycle, and that YTHDF2 plays a particularly prominent

role in mediating KSHV lytic gene expression in iSLK.219 cells.

The m^{6}A pathway post-transcriptionally controls ORF50 expression in iSLK.219 cells

ORF50 is the major viral transcriptional transactivator, and its expression is essential to

drive the KSHV lytic gene expression cascade [45]. The observations that ORF50 is m^{6}A

modified and that its accumulation is dependent on YTHDF2 indicate that the m^{6}A pathway

plays key roles in ORF50 mRNA biogenesis or fate in iSLK.219 cells, potentially explaining the

lytic cycle progression defect in the knockdown cells. m^{6}A deposition has been reported to

occur both co-transcriptionally and post-transcriptionally [1,16,17,46] . To determine whether

the m^{6}A pathway is important for ORF50 synthesis or mRNA turnover, we measured its

transcription in reactivated iSLK.219 cells upon depletion of METTL3, YTHDF1, 2, or 3 using 4-

thiouridine (4sU) metabolic pulse labeling. 4sU is a uridine derivative that can be used to

isolate newly transcribed RNA, as thiol-specific biotinylation of the 4sU-containing RNA enables

its purification over streptavidin-coated beads [47,48]. At 24 h post reactivation, the siRNA

treated iSLK.219 cells were pulse labeled with 4sU for 30 min, whereupon the labeled nascent

RNA was isolated by biotin-streptavidin purification and viral transcripts were quantified by RT-

qPCR (**Fig 4A**). Despite the defect in ORF50 accumulation observed upon YTHDF2 depletion

(see **Fig 3A**), we observed no decrease in ORF50 transcription upon depletion of any of the m^{6}A

writer or reader proteins (**Fig 4B**). However, in YTHDF2 depleted cells, there was a prominent

defect in the transcription of ORF37, likely because transcription of delayed early and late viral

genes is dependent on the presence of ORF50 protein (**Fig 4C**).

The observation that YTHDF2 was important for mediating the post-transcriptional,

stable accumulation of ORF50 mRNA suggested that the transcriptional context of ORF50

should not impact its regulation by the m^{6}A pathway in iSLK cells. Indeed, depletion of METTL3

or YTHDF2 also reduced the accumulation of ORF50 protein in an uninfected iSLK cell line

containing an integrated, dox-inducible ORF50 gene (iSLK.puro cells) (**Fig 4D and 4E**). Here,

YTHDF3 depletion resulted in an increase in ORF50 expression, which we also observed to a

more modest degree in the iSLK.219 cells (see **Fig 3D**). Overall, these data suggest that YTHDF2

depletion causes post-transcriptional degradation of the ORF50 mRNA, leading to a block in

KSHV lytic cycle progression in infected iSLK.219 cells.

The impact of m^{6}A on KSHV infection is cell type specific

Finally, given the diversity of functions reported for m^{6}A in controlling cellular processes and

RNA virus infections [1,22,27], we sought to evaluate the role of this pathway in another widely

used KSHV infected cell line of distinct origin, the B cell line TREX-BCBL1 [49]. Similar to

iSLK.219 cells, TREX-BCBL1 cells also contain a dox-inducible copy of ORF50 to boost

reactivation. The m^{6}A pathway components were depleted from TREX-BCBL1 cells via siRNA

treatment, whereupon cells were reactivated for 72 hr with dox, TPA, and ionomycin and the

abundance of the ORF50 and ORF59 lytic cycle proteins was monitored by western blot.

Consistent with observations in iSLK.219 cells, ORF50 and ORF59 expression was not affected

by depletion of YTHDF3, although it was enhanced by depletion of YTHDF1 (**Fig 5A and 5B**).

However, knockdown of METTL3 and YTHDF2 resulted in phenotypes clearly distinct from those

observed in iSLK.219 cells. There was a marked increase in ORF50 expression in METTL3

depleted TREX-BCBL1 cells, and near or above WT levels of ORF50 and ORF59 upon YTHDF2

knockdown (**Fig 5B**). We next examined whether the ORF50 mRNA was subject to m^6A

modification in reactivated TREX-BCBL1 cells by m^6A RIP, followed by RT-qPCR using control or

ORF50 specific primers. Indeed, there was a clear enrichment of ORF50 in the reactivated,

m^6A-containing RNA population (**Fig 5C**). As expected, we detected the m^6A modified DICER

transcript in both reactivated and unreactivated cells, whereas the unmodified GAPDH

transcript was present in neither sample (**Fig 5C**). Thus, although it is possible that the

modification state of some viral mRNAs differs between iSLK.219 and TREX-BCBL1 cells, the

ORF50 transcript is modified in both cell types.

288   Finally, to determine whether these differences in the contribution of the m^6A writer

and readers impacted the outcome of the viral lifecycle, we measured the impact of METTL3

and YTHDF1-3 protein knockdown on virion production using a supernatant transfer assay.

TREX-BCBL1 cells lack the viral GFP marker, and thus infection of recipient cells was instead

measured by RT-qPCR for the KSHV latency-associated LANA transcript. Consistent with the

viral protein expression data from METTL3 or YTHDF2-depleted TREX-BCBL1 cells, we observed

a modest, though not statistically significant, increase in the levels of LANA mRNA in the 293T

recipient cells infected with these supernatants (**Fig 5D**). These data confirm that METTL3 and

YTHDF2 are not required for virion production in TREX-BCBL1 cells, and suggest they may

instead decrease the efficiency of viral amplification. In summary, while there are some

similarities in the impact of YTHDF1 and YTHDF3 in iSLK.219 and TREX-BCBL1 cells, METTL3 and

YTHDF2 appear to function in a pro-viral capacity in iSLK.219 cells but in an antiviral capacity in

TREX-BCBL1 cells. These findings highlight how at least a subset of m^{6}A pathway functions and

targets may diverge between cell types.

Discussion

Although m^{6}A modification of viral RNAs has been recognized for more than 40 years,

only recently are the contributions of this epitranscriptomic mark towards viral life cycles

beginning to be revealed. Thus far, global epitranscriptomic analyses have documented m^{6}A

deposition during infections with HIV, Influenza A virus and several members of the *Flaviviridae*,

with a diverse set of resulting pro- and anti-viral roles [23-26,28,32]. The breadth and

occasionally apparently contrasting functions for the m^{6}A pathway during infection are perhaps

unsurprising given the dynamic role for this modification in controlling mRNA fate and its ability

to impact virtually every stage of host gene expression [1,27]. Here, we provide the first global

analysis of the m^{6}A epitranscriptome during lytic infection with a DNA virus, KSHV, revealing the

presence of m^{6}A across multiple kinetic classes of viral transcripts and a general decrease in

m^{6}A deposition on cellular mRNAs. In the widely used KSHV-positive cell line iSLK.219, we

found that depletion of several components of the m^{6}A pathway inhibited the KSHV lytic cycle,

most notably by restricting the post-transcriptional accumulation of the viral lytic transactivator

ORF50. The YTHDF2 reader protein proved particularly important, as its depletion eliminated

lytic entry and virion production. These observations are suggestive of a pro-viral role for m^{6}A

in the iSLK.219 KSHV reactivation model. However, m^{6}A marks on mRNA in a cell are

widespread and contribute to a large variety of cellular and pathogenic processes that likely

occur in a cell type or context-dependent manner. In this regard, it is notable that a distinct set

of phenotypes was observed for m^{6}A pathway components in the B cell line TREX-BCBL1. Here,

YTHDF2 had little effect on ORF50 expression, whereas depletion of METTL3 or YTHDF1

increased ORF50 abundance, more suggestive of an anti-viral role. Thus, although KSHV

engages the m^{6}A pathway in multiple cell types, these findings underscore the importance of

not broadly extrapolating m^{6}A roles from a particular cell type, as this complex regulatory

pathway can functionally vary in a cell type dependent manner.

What might be the basis for these phenotypic differences between cell types in the

context of KSHV infection? While this work was in preparation, Ye and colleagues performed an

RT-qPCR based analysis demonstrating m^{6}A deposition in many KSHV mRNAs in BCBL1 cells,

including ORF50 [38]. We also observed ORF50 modification in TREX-BCBL1 cells. Thus, it is not

the case that the viral mRNAs engage the m^{6}A methyltransferase machinery in one cell type but

not the other, although it is possible that site specificity of m^{6}A deposition varies between cell

lines. The facts that m^{6}A deposition is dynamic and does not strictly occur on consensus motifs

render this possibility challenging to resolve. Indeed, how m^{6}A deposition selectively controls

gene regulation on particular transcripts or under particular stimuli remains a central

unanswered question in the field [1]. We hypothesize that the distinct phenotypes derive either

from how the viral modifications are 'interpreted' in each cell type and/or indirect effects

driven by an altered m^{6}A profile on cellular mRNAs. The recent finding that m^{6}A modification of

ORF50 in BCBL1 cells contributes to efficient splicing through binding of YTHDC1 argues that

modifications can have a direct cis-acting impact on KSHV mRNA fate [38]. However,

herpesviral mRNAs are heavily reliant on host machinery at every stage of their biogenesis.

Given that cellular mRNA fate is significantly altered upon depletion of METTL3 and the YTHDF

reader proteins [1-3,18,50], it is possible that cell type specific changes in the abundance of a

host factor(s) required for viral mRNA stability also contribute to the phenotypic differences.

Furthermore, in HIV infected cells m^{6}A modification and YTHDF proteins have been proposed to

have a combination of pro-viral and anti-viral effects, including negatively impacting reverse

transcription, enhancing mRNA export, and increasing viral protein production [24,26,28].

Therefore, the m^{6}A pathway might similarly facilitate distinct phenotypes at different stages of

the KSHV lifecycle.

As the 'interpreters' of m^{6}A marks, the individual reader proteins play prominent roles

in modulating gene expression. Generally speaking, in HeLa and 293T cells, YTHDF1 binding

correlates with increased translational efficiency, YTHDF2 binding accelerates mRNA decay, and

YTHDF3 may serve as a cofactor to assist the other reader protein function [1-4,19,20,50].

However, other roles for these factors are rapidly emerging, particularly in the context of cell

stress, infection, or in the control of specific transcripts [7,13,15,23-26,28,50,51]. Furthermore,

m^{6}A is enriched in certain tissues, and different m^{6}A patterns have been found depending on

the tissue and developmental stage [41,52]. Intriguingly, a recent study showed that hypoxia

increases global m^{6}A content of mRNA, with many m^{6}A modified RNAs exhibiting increased

stability, raising the possibility that m^{6}A deposition could also stabilize transcripts during other

forms of cellular stress [53]. In KSHV-infected iSLK.219 cells, YTHDF2 appears essential for the

post-transcriptional accumulation of ORF50, a role seemingly at odds with its more canonical

mRNA destabilizing function. Although less dramatic viral gene expression phenotypes were

observed upon Mettl3 and YTHDF1 depletion, they nonetheless were required for WT levels of

progeny virion production. It is perhaps notable that even though YTHDF1 depletion modestly

enhanced the abundance of the ORF50, ORF37, ORF57, and K8.1 mRNAs, this ultimately did not

lead to WT virion production. An important consideration may be that m^{6}A factors

differentially impact specific KSHV transcripts, or play different roles at distinct times during

infection. However, dissecting these possibilities is likely to be complicated by the changes in

ORF50 expression (either positive or negative), which will have ripple effects on the entire lytic

life cycle.

Another relevant question is the extent to which m^{6}A mediates its effects on KSHV gene

expression co-transcriptionally versus post-transcriptionally. A recent report indicated that m^{6}A

is primarily installed in nascent mRNA in exons and affects cytoplasmic stability, but not splicing

[16,46]. It has also been demonstrated that that m^{6}A can be installed co-transcriptionally, and

that slowing the rate of RNA Pol II elongation enhances m^{6}A modification of mRNAs in a

manner that ultimately decreases translation efficiency [16,46]. These add to a growing body

of literature indicating that the position of m^{6}A in a transcript is a key feature impacting the

functional consequence of the modification [1,3,6,7,13,20]. For example, m^{6}A in the 3' UTR has

been shown to recruit YTHDF1 and enhance translation initiation in HeLa cells, while deposition

of m^{6}A in the 5' UTR has been shown to enhance 5' cap independent translation [3,7,13].

Whether these position-linked effects on translation extend to viral transcripts remains to be

tested, although there does not appear to be a consistent enrichment in a particular region of

viral mRNAs for the viruses analyzed thus far. In KSHV, m^6A sites are found throughout viral

ORFs, some of which also overlap with untranslated regions of other viral transcripts. As KSHV

transcription depends on the host RNA Pol II, the speed of transcriptional elongation on viral

mRNAs likely impacts co-transcriptional deposition and positioning of m^6A, and thus may

ultimately regulate translation efficiency of a given mRNA. Thus, in the context of KSHV

reactivation, a wide variety of mechanisms exist through which m^6A modification could impact

the transcription and translation of viral mRNA. Deciphering these remains an important

challenge for future studies, as we are currently in the early stages of understanding how this

and other viruses interface with the m^6A RNA modification pathway.

Materials and Methods

Cell culture. The renal carcinoma cell line iSLK.puro containing a doxycycline-inducible copy of

ORF50, and the KSHV infected renal carcinoma cell line iSLK.219 bearing docycycline-inducible

ORF50 [40] were cultured in Dulbecco's modified Eagle medium (DMEM; Invitrogen) with 10%

fetal bovine serum (FBS; Invitrogen, HyClone) and 100 U/ml penicillin-streptomycin

(Invitrogen). The KSHV-positive B cell line TREX-BCBL-1 containing a doxycycline-inducible

version of ORF50 [49] was cultured in RPMI medium (Invitrogen) supplemented with 20% FBS,

100 U/ml penicillin/streptomycin, and 200 uM L-glutamine (Invitrogen). HEK293T cells (ATCC)

were grown in DMEM (Invitrogen) supplemented with 10% FBS. To induce lytic reactivation of

iSLK.219 cells, $2x10^6$ cells were plated in a 10 cm dish with 1 ug/ml doxycycline (BD Biosciences)

and 1 mM sodium butyrate for 72 hr. Lytic reactivation of TREX-BCBL-1 cells was achieved by

treatment of $7x10^5$ cells/ml with 20 ng/ml 2-O-tetradecanoylphorbol-13-acetate (TPA, Sigma), 1

ug/ml doxycycline (BD Biosciences), and 500 ng/ml ionomycin (Fisher Scientific) for 72 hr

(western Blot blots for viral gene expression), or for 120 hr (supernatant transfer experiments).

siRNA experiments. For iSLK.219 cells, 100 pmol of siRNA was reverse transfected into $5x10^5$

cells plated in a 6-well dish using Lipofectamine RNAimax (Life Technologies). 24 hours post

transfection, cells were trypsinized and re-seeded on a 10 cm plate. The next day, a second

transfection was performed with 400 pmol siRNA and $2x10^6$ cells using the same method. The

following day, cells were lytically reactivated in a 10 cm plate. 24 hours post-reactivation, cells

were lysed in RIPA buffer (10 mM Tris-Cl (pH 8.0), 1 mM EDTA, 1% Triton X-100, 0.1% sodium

deoxycholate, 0.1% SDS, 140 mM NaCl) to evaluate knockdown efficiency.

For TREX-BCBL-1 cells, 200 pmol of siRNA was nucleofected into $2x10^6$ cells using Lonza

Cell Line Nucleofector Kit V and a Lonza Nucleofector 2b set to Program T001. After

nucleofection, cells were immediately resuspended in 2.2 ml of RPMI media in a 12 well plate.

24 hours later, 200 pmol of siRNA was added again to $2x10^6$ cells using the same protocol. 48

hours after the second transfection, cells were lysed in RIPA buffer and knockdown efficiency

was analyzed by Western Blot.

The following Qiagen siRNAs were used: SI00764715 and SI04279121 targeting YTHDF1,

SI04174534 targeting YTHDF2, SI00764778 and SI04205761 targeting YTHDF3, custom siRNA

targeting Mettl3 (sequence targeted: CTGCAAGTATGTTCACTATGA), Qiagen SI03650318

(negative control siRNA). The following Dharmacon siRNAs were used: SMARTpool siGenome

(M-021009-01-0005), targeting YTHDF2, and siGENOME Non-Targeting siRNA Pool #1

(D0012061305).

Supernatant transfer assay and quantification of virion production. Assays were performed as

previously described [54]. Briefly, for iSLK.219 cells, viral supernatant was collected 72 hr post-

reactivation, filtered, and added to uninfected HEK293T cells by spinfection (1500 rpm, 90

minutes at room temperature). 12 hours later, supernatant was removed and replaced with

fresh media, whereupon the cells were assessed for the successful transfer of the GFP-

containing KSHV BAC 24 hr post-infection using a BD Accuri C6 flow cytometer. Briefly, cells

were trypsinized, washed twice in PBS and resuspended in FACS Buffer (PBS with 1% FBS).

Uninfected HEK293T cells were used to define the GFP negative population. The percentage of

GFP expressing cells was quantified using FlowJo Software (FlowJo LLC). For virus produced in

TREX-BCBL-1 cells, supernatant transfers were performed as in iSLK.219 cells, except the virus

was transferred to HEK293T cells at 120 hr post-reactivation. To quantify virus produced in

TREX-BCBL-1 cells, RNA was extracted from HEK293T cells 48 hours post-supernatant transfer,

and viral gene expression was quantified by RT-qPCR using primers specific for LANA.

Affinity purification and Western blotting. Cell lysate was collected and analyzed as previously

described [54]. Briefly, iSLK.219 or TREX-BCBL-1 cells were trypsinized, washed with PBS and

lysed in RIPA buffer with protease inhibitors. After washing, SDS-sample buffer (indicate

components) was added to samples to elute bound proteins. Lysates were resolved by SDS-

PAGE and western blots were carried out with the following antibodies: rabbit ORF50 (gift of

Yoshihiro Izumiya, UC Davis), rabbit α-K8.1 (1:10000, antibody generated for this study), rabbit

α-ORF59 (1:10000, antibody generated for this study), rabbit α-Mettl3 (Bethyl, 1:1000), rabbit

α-YTHDF1 (Proteintech, 1:1000), rabbit α-YTHDF2 (Millipore, 1:1000), rabbit α-YTHDF3 (Sigma,

1:1000), and goat α-mouse and goat α-rabbit HRP secondary antibodies (1:5000; Southern

Biotech).

4sU Labeling. Following siRNA knockdown and 24 hr reactivation, iSLK.219 cells were pulse

labeled with DMEM containing 500 µM 4sU (Sigma) for 30 minutes, followed by PBS wash and

immediate isolation of total RNA with TRIzol. 4SU isolation was performed as previously

described [55]. 4SU isolated RNA was analyzed by RT-qPCR.

RT-qPCR. Total RNA was harvested using TRIzol following the manufacturer's protocol. Samples

were DNase treated using Turbo DNase (Ambion), and cDNA was synthesized from 2 ug of total

RNA using AMV reverse transcriptase (Promega), and used directly for quantitative PCR (qPCR)

analysis with the DyNAmo ColorFlash SYBR green qPCR kit (Thermo Scientific). All qPCR results

were normalized to 18S levels and WT or scramble control set to 1.

LC-MS/MS analysis of m^{6}A. Total RNA was isolated from iSLK219 cells with TRIzol reagent.

Dynabeads mRNA purification kit (Ambion) was used to isolate polyA(+) RNAs from 100ug of

total RNA according to the manufacturer. 100-200 ng of polyadenylated RNA was spiked with

10uM of 5-fluorouridine (Sigma) and digested by nuclease P1 (1 U) in 25 µL of buffer containing

25 mM NaCl and 2.5 mM $ZnCl_2$ at 42 °C for 2-4 h, followed by addition of NH_4HCO_3 (1 M, 3 µL)

and bacterial alkaline phosphatase (1 U) and incubation at 37 °C for 2 h. The sample was then

filtered (Amicon 3K cutoff spin column), and 5 µL of the flow through was analyzed by liquid

chromatography (LC) coupled to an Orbitrap-XL mass spectrometer (MS) equipped with an

electrospray ionization source (QB3 Chemistry facility).

m⁶A-RIP and m⁶A-RIP-sequencing. Total cellular RNA (containing KSHV RNA) was extracted and

purified by TRIzol and then DNAse treated with Turbo DNase (Ambion). 30 ul protein G

magnetic beads (Invitrogen) were blocked in 1% BSA solution for 1 hour, followed by incubation

with 12.5 ug affinity-purified anti-m⁶A polyclonal antibody (Millipore) at 4°C for 2 hr with head-

over-tail rotation. 100 ug purified RNA was added to the antibody-bound beads in IP buffer

(150 mM NaCl, 0.1% NP-40, and 10 mM Tris-HCl [pH 7.4]) containing RNAse inhibitor and

protease inhibitor cocktail and incubated overnight at 4°C with head-over-tail rotation. The

beads were washed three times in IP buffer, and then RNA was competitively eluted with 6.7

mM m⁶A-free nucleotide solution (Sigma Aldrich). RNA in the eluate was phenol chloroform

extracted and then reverse transcribed to cDNA for Real-Time qPCR analysis.

High-throughput sequencing of the KSHV methylome (m⁶Aseq) was carried following the

previously published protocol [56]. In brief, 2.5 mg total cellular RNA was prepared from

iSLK.219 cells that were either unreactivated, or reactivated for five days with doxycycline. RNA

was isolated and DNAse treated as in the m⁶A RIP, except the RNA was first fragmented to

lengths of ~100 nt prior to immunoprecipitation with anti-m⁶A antibody (Synaptic Systems).

Immunoprecipitated RNA fragments and comparable amounts of input were subjected to first-

strand cDNA synthesis using the NEBNext Ultra RNA Library Prep Kit for Illumina (New England

Biolabs). Sequencing was carried out on Illumina HiSeq2500 according to the manufacturer's

instructions, using 10 pM template per sample for cluster generation, TruSeq SR Cluster kit v3

(Illumina), TruSeq SBS Kit v3-HS (Illumina) and TruSeq Multiplex Sequencing primer kit

(Illumina).A reference human transcriptome was prepared based on the University of California,

Santa Cruz (UCSC) and a reference KSHV transcriptome based on KSHV 2.0 annotation [57] .

Analysis of m^{6}A peaks was performed using the model-based analysis of ChIP-seq (MACS) peak-

calling algorithm. Peaks were considered significant if their MACS-assigned fold change was

greater than four and individual FDR value less than 5%. Sequencing data are available on GEO

repository (accession number GSE104621).

Statistical Analysis: All results are expressed as means +/- S.E.M. of experiments independently

repeated at least three times, except where indicated. Unpaired Student's t test was used to

evaluate the statistical difference between samples. Significance was evaluated with P values

<0.05.

Acknowledgements

This research was funded by NIH (http://www.nih.gov/) grants R01AI122528 to B.G. and

HG008688 to C.H. B.G. and C.H. are investigators of the Howard Hughes Medical Institute.

<u>References</u>

1.   Roundtree IA, Evans ME, Pan T, He C. Dynamic RNA Modifications in Gene Expression
Regulation. Cell. 2017;169: 1187–1200. doi:10.1016/j.cell.2017.05.045

2.   Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, et al. N(6)-methyladenosine-dependent
regulation of messenger RNA stability. Nature. 2013;505: 117–120.
doi:10.1038/nature12730

3.   Wang X, Zhao BS, Roundtree IA, Lu Z, Han D, Ma H, et al. N(6)-methyladenosine
Modulates Messenger RNA Translation Efficiency. Cell. 2015;161: 1388–1399.
doi:10.1016/j.cell.2015.05.014

4.   Fu Y, Dominissini D, Rechavi G, He C. Gene expression regulation mediated through

reversible m(6)A RNA methylation. Nat Rev Genet. 2014;15: 293–306.
doi:10.1038/nrg3724

5.   Zhao BS, Wang X, Beadell AV, Lu Z, Shi H, Kuuspalu A, et al. m(6)A-dependent maternal
mRNA clearance facilitates zebrafish maternal-to-zygotic transition. Nature. 2017;542:
475–478. doi:10.1038/nature21355

6.   Xiao W, Adhikari S, Dahal U, Chen Y-S, Hao Y-J, Sun B-F, et al. Nuclear m(6)A Reader
YTHDC1 Regulates mRNA Splicing. Molecular Cell. 2016;61: 507–519.
doi:10.1016/j.molcel.2016.01.012

7.   Meyer KD, Patil DP, Zhou J, Zinoviev A, Skabkin MA, Elemento O, et al. 5' UTR m(6)A
Promotes Cap-Independent Translation. Cell. 2015;163: 999–1010.
doi:10.1016/j.cell.2015.10.012

8.   Liu N, Pan T. N(6)-methyladenosine–encoded epitranscriptomics. Cell Research. 2016;23:
98–102. doi:10.1038/nsmb.3162

9.   Liu N, Dai Q, Zheng G, He C, Parisien M, Pan T. N(6)-methyladenosine-dependent RNA
structural switches regulate RNA-protein interactions. Nature. 2015;518: 560–564.
doi:10.1038/nature14234

10.   Zhou KI, Parisien M, Dai Q, Liu N, Diatchenko L, Sachleben JR, et al. N(6)-
Methyladenosine Modification in a Long Noncoding RNA Hairpin Predisposes Its
Conformation to Protein Binding. J Mol Biol. 2016;428: 822–833.
doi:10.1016/j.jmb.2015.08.021

11.   Liu N, Zhou KI, Parisien M, Dai Q, Diatchenko L, Pan T. N(6)-methyladenosine alters RNA
structure to regulate binding of a low-complexity protein. Nucleic Acids Research.
2017;45: 6051–6063. doi:10.1093/nar/gkx141

12.   Lewis CJT, Pan T, Kalsotra A. RNA modifications and structures cooperate to guide RNA-
protein interactions. Nat Rev Mol Cell Biol. 2017;18: 202–210. doi:10.1038/nrm.2016.163

13.   Zhou J, Wan J, Gao X, Zhang X, Jaffrey SR, Qian S-B. Dynamic m(6)A mRNA methylation
directs translational control of heat shock response. Nature. 2015;526: 591–594.
doi:10.1038/nature15377

14.   Meyer KD, Jaffrey SR. The dynamic epitranscriptome: N(6)-methyladenosine and gene
expression control. Nat Rev Mol Cell Biol. 2014;15: 313–326. doi:10.1038/nrm3785

15.   Xiang Y, Laurent B, Hsu C-H, Nachtergaele S, Lu Z, Sheng W, et al. RNA m(6)A methylation
regulates the ultraviolet-induced DNA damage response. Nature. 2017;543: 573–576.
doi:10.1038/nature21671

16.   Slobodin B, Han R, Calderone V, Vrielink JAFO, Loayza-Puch F, Elkon R, et al. Transcription

Impacts the Efficiency of mRNA Translation via Co-transcriptional N(6)-adenosine
Methylation. Cell. 2017;169: 326–337.e12. doi:10.1016/j.cell.2017.03.031

17.   Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, et al. A METTL3-METTL14 complex mediates
mammalian nuclear RNA N(6)-adenosine methylation. Nat Chem Biol. 2014;10: 93–95.
doi:10.1038/nchembio.1432

18.   Du H, Zhao Y, He J, Zhang Y, Xi H, Liu M, et al. YTHDF2 destabilizes m(6)A-containing RNA
through direct recruitment of the CCR4-NOT deadenylase complex. Nat Commun.
2016;7: 12626. doi:10.1038/ncomms12626

19.   Shi H, Wang X, Lu Z, Zhao BS, Ma H, Hsu PJ, et al. YTHDF3 facilitates translation and decay
of N(6)-methyladenosine-modified RNA. Cell Research. 2017;27: 315–328.
doi:10.1038/cr.2017.15

20.   Li A, Chen Y-S, Ping X-L, Yang X, Xiao W, Yang Y, et al. Cytoplasmic m(6)A reader YTHDF3
promotes mRNA translation. Cell Research. 2017;27: 444–447. doi:10.1038/cr.2017.10

21.   Hsu PJ, Zhu Y, Ma H, Guo Y, Shi X, Liu Y, et al. Ythdc2 is an N(6)-methyladenosine binding
protein that regulates mammalian spermatogenesis. Cell Research. 2017;27: 1115–1127.
doi:10.1038/cr.2017.99

22.   Gokhale NS, Horner SM. RNA modifications go viral. PLoS Pathog. 2017;13: e1006188.
doi:10.1371/journal.ppat.1006188

23.   Gokhale NS, McIntyre ABR, McFadden MJ, Roder AE, Kennedy EM, Gandara JA, et al.
N(6)-methyladenosine in Flaviviridae Viral RNA Genomes Regulates Infection. Cell Host
Microbe. 2016;20: 654–665. doi:10.1016/j.chom.2016.09.015

24.   Kennedy EM, Bogerd HP, Kornepati AVR, Kang D, Ghoshal D, Marshall JB, et al.
Posttranscriptional m(6)A Editing of HIV-1 mRNAs Enhances Viral Gene Expression. Cell
Host Microbe. 2016;19: 675–685. doi:10.1016/j.chom.2016.04.002

25.   Lichinchi G, Zhao BS, Wu Y, Lu Z, Qin Y, He C, et al. Dynamics of Human and Viral RNA
Methylation during Zika Virus Infection. Cell Host Microbe. 2016;20: 666–673.
doi:10.1016/j.chom.2016.10.002

26.   Lichinchi G, Gao S, Saletore Y, Gonzalez GM, Bansal V, Wang Y, et al. Dynamics of the
human and viral m(6)A RNA methylomes during HIV-1 infection of T cells. Nat Microbiol.
2016;1: 16011. doi:10.1038/nmicrobiol.2016.11

27.   Gonzales-van Horn SR, Sarnow P. Making the Mark: The Role of Adenosine Modifications
in the Life Cycle of RNA Viruses. Cell Host Microbe. 2017;21: 661–669.
doi:10.1016/j.chom.2017.05.008

28.   Tirumuru N, Zhao BS, Lu W, Lu Z, He C, Wu L. N(6)-methyladenosine of HIV-1 RNA

regulates viral infection and HIV-1 Gag protein expression. Elife. 2016;5: 165.
doi:10.7554/eLife.15528

29. Martínez-Pérez M, Aparicio F, López-Gresa MP, Bellés JM, Sánchez-Navarro JA, Pallás V.
Arabidopsis m(6)A demethylase activity modulates viral infection of a plant virus and the
m(6)A abundance in its genomic RNAs. Proc Natl Acad Sci USA. 2017.
doi:10.1073/pnas.1703139114

30. Durbin AF, Wang C, Marcotrigiano J, Gehrke L. RNAs Containing Modified Nucleotides
Fail To Trigger RIG-I Conformational Changes for Innate Immune Signaling. MBio. 2016;7:
e00833–16. doi:10.1128/mBio.00833-16

31. Karikó K, Buckstein M, Ni H, Weissman D. Suppression of RNA recognition by Toll-like
receptors: the impact of nucleoside modification and the evolutionary origin of RNA.
Immunity. 2005;23: 165–175. doi:10.1016/j.immuni.2005.06.008

32. Courtney DG, Kennedy EM, Dumm RE, Bogerd HP, Tsai K, Heaton NS, et al.
Epitranscriptomic Enhancement of Influenza A Virus Gene Expression and Replication.
Cell Host Microbe. 2017;22: 377–386.e5. doi:10.1016/j.chom.2017.08.004

33. Lavi S, Shatkin AJ. Methylated simian virus 40-specific RNA from nuclei and cytoplasm of
infected BSC-1 cells. 1975;72: 2012–2016.

34. Krug RM, Morgan MA, Shatkin AJ. Influenza viral mRNA contains internal N(6)-
methyladenosine and 5'-terminal 7-methylguanosine in cap structures. J Virol. 1976;20:
45–53. doi:10.1177/1091581816683642

35. Kahana C, Lavi S, Groner Y. Identification and mapping of N(6)-methyladenosine
containing sequences in simian virus 40 RNA. Nucleic Acids Research. 1979;6: 2879–
2899.

36. Moss B, Gershowitz A, Stringer JR, Holland LE, Wagner EK. 5'-Terminal and internal
methylated nucleosides in herpes simplex virus type 1 mRNA. J Virol. 1977;23: 234–239.

37. Sommer S, Salditt-Georgieff M. The methylation of adenovirus-specific nuclear and
cytoplasmic RNA. Nucleic Acids Research. 1976;3: 749–765.

38. Ye F, Chen ER, Nilsen TW. Kaposi's Sarcoma-Associated Herpesvirus Utilizes and
Manipulates RNA N(6)-Adenosine Methylation To Promote Lytic Replication. J Virol.
2017;91. doi:10.1128/JVI.00466-17

39. Vieira J, O'Hearn PM. Use of the red fluorescent protein as a marker of Kaposi's sarcoma-
associated herpesvirus lytic gene expression. Virology. 2004;325: 225–240.
doi:10.1016/j.virol.2004.03.049

40. Myoung J, Ganem D. Infection of Lymphoblastoid Cell Lines by Kaposi's Sarcoma-

Associated Herpesvirus: Critical Role of Cell-Associated Virus. J Virol. 2011;85: 9767–
9777. doi:10.1128/JVI.05136-11

41. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg
S, et al. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq.
Nature. 2012;485: 201–206. doi:10.1038/nature11112

42. Sun R, Lin SF, Gradoville L, Miller G. Polyadenylylated nuclear RNA encoded by Kaposi
sarcoma-associated herpesvirus. PNAS. 1996;93: 11883–11888.

43. Bai Z, Huang Y, Li W, Zhu Y, Jung JU, Lu C, et al. Genomewide mapping and screening of
Kaposi's sarcoma-associated herpesvirus (KSHV) 3'untranslated regions identify
bicistronic and polycistronic viral transcripts as frequent targets of KSHV microRNAs. J
Virol. 2014;88: 377–392. doi:10.1128/JVI.02689-13

44. Chandriani S, Ganem D. Array-based transcript profiling and limiting-dilution reverse
transcription-PCR analysis identify additional latent genes in Kaposi's sarcoma-associated
herpesvirus. J Virol. 2010;84: 5565–5573. doi:10.1128/JVI.02723-09

45. Lukac DM, Kirshner JR, Ganem D. Transcriptional activation by the product of open
reading frame 50 of Kaposi's sarcoma-associated herpesvirus is required for lytic viral
reactivation in B cells. J Virol. 1999;73: 9348–9361.

46. Ke S, Pandya-Jones A, Saito Y, Fak JJ, Vågbø CB, Geula S, et al. m(6)A mRNA modifications
are deposited in nascent pre-mRNA and are not required for splicing but do specify
cytoplasmic turnover. Genes Dev. 2017;31: 990–1006. doi:10.1101/gad.301036.117

47. Cleary MD, Meiering CD, Jan E, Guymon R, Boothroyd JC. Biosynthetic labeling of RNA
with uracil phosphoribosyltransferase allows cell-specific microarray analysis of mRNA
synthesis and decay. Nat Biotechnol. 2005;23: 232–237. doi:10.1038/nbt1061

48. Woodford TA, Schlegel R, Pardee AB. Selective isolation of newly synthesized mammalian
mRNA after in vivo labeling with 4-thiouridine or 6-thioguanosine. Analytical
biochemistry. 1988;171: 166–172. doi:10.1016/0003-2697(88)90138-8

49. Nakamura H, Lu M, Gwack Y, Souvlis J, Zeichner SL, Jung JU. Global Changes in Kaposi's
Sarcoma-Associated Virus Gene Expression Patterns following Expression of a
Tetracycline-Inducible Rta Transactivator. J Virol. 2003;77: 4205–4220.
doi:10.1128/JVI.77.7.4205-4220.2003

50. Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z, Zhao JC. N(6)-methyladenosine modification
destabilizes developmental regulators in embryonic stem cells. Nat Cell Biol. 2014;16:
191–198. doi:10.1038/ncb2902

51. Lin S, Choe J, Du P, Triboulet R, Gregory RI. The m(6)A Methyltransferase METTL3
Promotes Translation in Human Cancer Cells. Molecular Cell. 2016;62: 335–345.

doi:10.1016/j.molcel.2016.03.021

52.   Wu R, Jiang D, Wang Y, Wang X. N(6)-Methyladenosine (m(6)A) Methylation in mRNA
652   with A Dynamic and Reversible Epigenetic Modification. Mol Biotechnol. 2016;58: 450–
653   459. doi:10.1007/s12033-016-9947-9

654   53.   Fry NJ, Law BA, Ilkayeva OR, Holley CL, Mansfield KD. N(6)-methyladenosine is required
655   for the hypoxic stabilization of specific mRNAs. RNA. 2017;23: 1444–1455.
656   doi:10.1261/rna.061044.117

657   54.   Davis ZH, Verschueren E, Jang GM, Kleffman K, Johnson JR, Park J, et al. Global mapping
658   of herpesvirus-host protein complexes reveals a transcription strategy for late genes.
659   Molecular Cell. 2015;57: 349–360. doi:10.1016/j.molcel.2014.11.026

660   55.   Abernathy E, Gilbertson S, Alla R, Glaunsinger B. Viral Nucleases Induce an mRNA
661   Degradation- Transcription Feedback Loop in Mammalian Cells. Cell Host Microbe.
662   2015;18: 243–253. doi:10.1016/j.chom.2015.06.019

663   56.   Dominissini D, Moshitch-Moshkovitz S, Salmon-Divon M, Amariglio N, Rechavi G.
664   Transcriptome-wide mapping of N(6)-methyladenosine by m(6)A-seq based on
665   immunocapturing and massively parallel sequencing. Nat Protoc. 2013;8: 176–189.
666   doi:10.1038/nprot.2012.148

667   57.   Arias C, Weisburd B, Stern-Ginossar N, Mercier A, Madrid AS, Bellare P, et al. KSHV 2.0: a
668   comprehensive annotation of the Kaposi's sarcoma-associated herpesvirus genome using
669   next-generation sequencing reveals novel genomic and functional features. PLoS Pathog.
670   2014;10: e1003847. doi:10.1371/journal.ppat.1003847

Figure Legends:

Fig 1. m^{6}A increases upon KSHV reactivation. (A) Schematic of the experimental setup.
iSLK.219 cells were induced with doxycycline for 5 days to induce the lytic cycle, and total RNA
was collected and subjected to oligo dT selection to purify poly(A) RNA. Polyadenylated RNA
was spiked with 10uM of 5-fluorouridine and digested with nuclease P1 and alkaline
phosphatase, and subjected to LC-MS/MS analysis. (B) Relative m^{6}A content in iSLK.219 cells.
The induced sample was normalized with respect to the uninduced sample (set to 1).

Fig 2. KSHV mRNA contains m^{6}A modifications. (A) Two independent replicates of iSLK.219
cells containing latent KSHV were treated with dox for 5 days to induce the viral lytic cycle
(induced) or left untreated to preserve viral latency (uninduced).   DNase-treated RNA was
isolated and subjected to m^{6}Aseq. Displayed are peaks with a fold change of four or higher,
comparing reads in the m^{6}A-IP to the corresponding input. (B) Overview of sequencing reads
from induced m^{6}A IP samples, aligned to the ORF50 transcript and the annotated GG(m^{6}A)C

consensus motifs found in exon 2 of ORF50. The red and green colored bars denote the
sequences encompassed by the FC>4 peaks in each replicate. (C) Cells were induced as in (A),
and total RNA was subjected to m^{6}A RIP, followed by RT-qPCR using primers for the indicated
viral and cellular genes. Values are displayed as fold change over input. (D) Quantification of
cellular m^{6}A peaks from m^{6}Aseq analysis.

Fig 3. m^{6}A and the reader YTHDF2 potentiate viral gene expression and virion production in
iSLK.219 cells. Cells were transfected with control scramble (scr) siRNAs or siRNAs against
Mettl3, YTHDF1, 2, or 3, then reactivated for 72 hours with doxycycline and sodium butyrate.
(A) Knockdown efficiency was measured by western blot using antibodies for the indicated
protein, with GAPDH serving as a loading control in this and all subsequent figures. (B) Viral
supernatant was collected from the reactivated iSLK.219 cells and transferred to uninfected
HEK293T recipient cells. 24 h later, the recipient cells were analyzed by flow cytometry for the
presence of GFP, indicating transfer of infectious virions. (C) ORF50, ORF37, ORF57, and K8.1
gene expression was analyzed by RT-qPCR from cells treated with the indicated siRNAs. One
representative experiment, matched to the siRNA knockdown in *(A)* is shown. (D) Expression of
the viral ORF50 and ORF59 proteins in cells treated with the indicated siRNAs was measured by
western blot 72hr post-reactivation. (E) Unreactivated iSLK.219 cells containing latent virus
were treated with control scramble (scr) siRNAs or siRNAs targeting Mettl3, YTHDF1, YTHDF2,
or YTHDF3. The cells were then reactivated with dox and sodium butyrate for 48 hr and lytic
reactivation was monitored by expression of the lytic promoter-driven red fluorescent protein.
(F) Protein was harvested from the above described cells and subjected to western blot for
ORF50 and the control GAPDH protein at 24 hr post-reactivation.

Fig 4. Depletion of the m^{6}A writer and readers does not impact ORF50 nascent transcription
in iSLK.219 cells. (A) Schematic of the experimental setup for measuring nascent RNA
synthesis. Cells were transfected with the indicated siRNAs for 48h then reactivated for 24
hours with dox. 4SU was added for 30 minutes, whereupon nascent 4SU-labeled RNA was
isolated using biotin/streptavidin affinity purification, reverse transcribed, and analyzed by RT-
qPCR using primers specific to ORF50 or ORF37. (B-C) Levels of nascent ORF50 *(B)* or ORF37
mRNA *(C)*, determined as described above. Unpaired Student's t test was used to evaluate the
statistical difference between samples. Significance was evaluated with P values <0.05. (D-E)
Uninfected iSLK.puro cells expressing DOX-inducible RTA were transfected with the indicated
siRNAs for 48 h, then treated with dox for 24 hours to induce ORF50 expression. Knockdown
efficiency *(D)* and ORF50 protein levels *(E)* were measured by western blot using antibodies for
the indicated protein, with GAPDH serving as a loading control.

Fig 5. Increased viral gene expression upon m^{6}A writer and reader depletion in TREX-BCBL1
cells. (A-B, D) TREX-BCBL1 cells were nucleofected with control scramble (scr) siRNAs or siRNAs
specific to Mettl3, YTHDF1, YTHDF2, or YTHDF3, then lytically reactivated by treatment with

dox, TPA and ionomycin for 72 hr. (A) Knockdown efficiency of the m^{6}A proteins relative to the
loading control GAPDH was quantified by western blot. (B) Levels of the KSHV ORF50 and
ORF59 proteins were assayed by western blot in the control and m^{6}A protein-depleted samples.
(C) TREX-BCBL1 cells were reactivated with dox for 72 hr, then total RNA was isolated and
subjected to m^{6}A RIP, followed by RT-qPCR for analysis of KSHV ORF50, GAPDH, and DICER.
Values are displayed as fold change over input. (D) Viral supernatant from the reactivated
control or m^{6}A protein depleted TREX-BCBL1 cells was transferred to uninfected HEK293T
recipient cells, whereupon transfer of infection was quantified by RT-qPCR for the viral LANA
transcript 48 hr post-supernatant transfer.

Supporting Information:

S1. Fig. Location of union FC>4 peaks within KSHV transcriptome. Overview of sequencing
reads aligned to regions of the KSHV transcriptome containing m^{6}A modifications. Depicted are
peaks with a fold change of four or higher in both replicates, comparing reads in the m^{6}A-IP to
the corresponding input. The blue and purple bars denote the sequences encompassed by the
FC>4 peaks in each uninduced replicate (A), while the red and green colored bars denote the
FC>4 peaks in each induced replicate (B). In the reference transcriptome, grey bars indicate
annotated 3'UTRs or ncRNAs, while burgundy arrows depict ORFs. Note that the alignment of
the ORF50 induced peaks can be found in Figure 2.

S1. Table: Full list of FC>2 peaks within KSHV transcripts in induced and uninduced samples.

S2. Table: Full list of FC>4 peaks within host transcripts in induced and uninduced samples.

S3. Table: List of RT-qPCR primers used in this study.

Primer	Sequence (5'-3')	Orientation F: Forward R: Reverse
vIL6	CGGTTCACTGCTGGTATCTG	F
vIL6	CAGTATCGTTGATGGCTGGT	R

ORF57	TTTGACGAATCGAGGGACGACG	F
ORF57	GCAGTTGAGAACGACCTTGAGAT	R
ORF37	TGGGCGAGTTTATTGGTAGTGAGG	F
ORF37	CTCCACTAGACAGCAGATGTGG	R
K8.1	TCCCTAAACGGGACCAGACT	F
K8.1	ACCCAGAGGCAGACGTATCT	R
PAN	TAATGTGAAAGGAAAGCAGCGCCC	F
PAN	CATTTAGGGCAAAGTGGCCCGATT	R
vGPCR	GTGCCTTACACGTGGAACGTT	F
vGPCR	GGTGACCAATCCATTTCCAAGA	R
K1	CCAAACGGACGAAATGAAAC	F
K1	TGTGTGGTTGCATCGCTATT	R
GAPDH	CGGAGTCAACGGATTTGGTCGTAT	F
GAPDH	AGCCTTCTCCATGGTGGTGAAGAC	R
ORF50	CGCAATGCGTTACGTTGTTG	F
ORF50	GCCCGGACTGTTGAATCG	R
DICER	TGCTATGTCGCCTTGAATGTT	F

DICER	AATTTCTCGATAGGGGTGGTCTA	R
18s	GTAACCCGTTGAACCCCATT	F
18s	CCATCCAATCGGTAGTAGCG	R
LANA	TGGCCCATCTCGCGAATA	F
LANA	AACGCGCCTCATACGAACTC	R

A

B

Total m⁶A levels by MS/MS

A

induced 1
induced 2
uninduced 1
uninduced 2

B

71412

GGAC motifs in exon 2 of ORF50

74445

ORF50
ORF49

induced 1 m6A IP

induced 2 m6A IP

C

m⁶A RIP RT-qPCR

Fold Enrichment
(Relative to GAPDH)

uninduced
induced

GAPDH Dicer vIL6 K1 vGPCR ORF50 ORF57 PAN

Cellular   Viral

D

m⁶A peaks in cellular genes

peak count

15000

10000

5000

Uninduced   Induced

A

B

No virus   Scramble   siMETTL3

0%   82%   7%

siYTHDF1   siYTHDF2   siYTHDF3

82%   <1%   58%

C

ORF50   ORF37   ORF57   K8.1

D   72 hr post-induction

E

F   24 hr post-induction

A siRNA KD → reactivate virus → → purify nascent 4SU labeled RNA → RT-qPCR
72 hr   24 hr

B ORF50

C ORF37

D iSLK.puro cells (uninfected) KD efficiency

E iSLK.puro cells (uninfected) 24 hours post-DOX

A

siMETTL3 siYTHDF1 siYTHDF2 siYTHDF3 Scramble

METTL3
GAPDH

YTHDF1
GAPDH

siMETTL3 siYTHDF1 siYTHDF2 siYTHDF3 Scramble

YTHDF2
GAPDH

YTHDF3
GAPDH

B

TREX-BCBL-1 72 hr PI

siMETTL3 siYTHDF1 siYTHDF2 siYTHDF3 Scramble

ORF50

ORF59

GAPDH

C   m⁶A RIP TREX-BCBL-1

Fold Enrichment/GAPDH

uninduced
induced

ORF50   DICER   GAPDH

D   LANA Post-Supernatant Transfer

Fold Change/18s

siMETTL3   siYTHDF2   siYTHDF3   Scramble