Acetylcholine reconfigures hippocampal circuits to enable rapid formation of overlapping memory ensembles

Luke Y. Prince[1], Krasimira Tsaneva-Atanasova[2,3], Claudia Clopath[4], and Jack R. Mellor[1]

[1] Centre for Synaptic Plasticity, School of Physiology Pharmacology, and Neuroscience, University of Bristol, BS8 1TD, UK

[2] Department of Mathematics and Living Systems Institute, University of Exeter, Exeter, EX4 4QF, UK

[3] EPRSC Centre for Predictive Modelling in Healthcare, University of Exeter, Exeter, EX4 4QJ, UK

[4] Bioengineering Department, Imperial College London, London, SW7 2AZ, UK

Abstract

In the hippocampus, episodic memories are thought to be encoded by the formation of ensembles of synaptically coupled CA3 pyramidal cells driven by sparse but powerful mossy fiber inputs from dentate gyrus granule cells. Since CA3 network capacity is finite, a mechanism for enhancing memory encoding during important events would ensure greater efficiency, but the salient signal that might determine this memory selectivity filter is unknown. Using experimental and computational methods we show that the cholinergic system can operate as a memory selectivity filter by combining 3 separate mechanisms: disinhibition of feed-forward mossy fiber inhibition enables synaptic plasticity at CA3-CA3 recurrent synapses; Increasing cellular excitability lowers the threshold for ensemble formation; Reduction of overall CA3-CA3 recurrent synapse strength allows for an increase in overlap between ensembles. Thus, we show that acetylcholine release enables the selective encoding of salient high-density episodic memories in the hippocampus.

Introduction

The hippocampus plays a central role in the formation of episodic memories by processing information from the entorhinal cortex sequentially through the dentate gyrus, CA3 and CA1 regions. Anatomical, functional and theoretical considerations propose separate computational properties for each of these regions in support of memory processing (Kesner and Rolls, 2015; Marr, 1971; McClelland and Goddard, 1996). In particular, the CA3 region is characterized by a recurrently connected set of excitatory pyramidal neurons, that are believed to encode auto-associative memories by selectively strengthening recurrent synapses between neurons in ensembles that provide a neural representation of the memory (Nakazawa et al., 2002). Configuring a recurrent network in this way is proposed to endow the network with attractor dynamics, in which the network is driven towards a stable state that minimizes energy within the network (Marr, 1971; Hopfield, 1982; Kesner and Rolls, 2015). This process is related to memory retrieval, in which external sources of input will alter the state of the network by activating subsets of neurons and through recurrent dynamics will be driven towards these stable states in which all neurons in the ensemble are reactivated – a process also referred to as pattern completion (Gold and Kesner, 2006). Within this framework, memory encoding is believed to be the procedure of altering the energy landscape through synaptic plasticity to create new attractor states, or change their position (Treves and Rolls, 1994; Tsodyks, 1999). However, not all memories are stored, indicating that there may be a gate to select which experiences should be encoded, but it is unclear how such a filter might operate.

One potential filter mechanism is the release of neuromodulators such as acetylcholine that can rapidly reconfigure neuronal networks (Prince et al., 2016). Acetylcholine is thought to promote encoding of new information by facilitating NMDA receptor function and induction of synaptic plasticity (Buchanan et al., 2010; Dennis et al., 2016; Fernandez de Sevilla and Buno, 2010; Gu and Yakel, 2011; Marino et al., 1998; Markram and Segal, 1992; Papouin et al., 2017) and selectively suppressing recurrent activity representing previously stored information in favour of feed-forward activity representing novel information (Hasselmo et al., 1995; Hummos et al., 2014). Commensurate with the concept of acetylcholine being important for encoding new information, acetylcholine release in the hippocampus and neocortex is associated with reward and arousal during periods of behavioural activity when it is beneficial to learn associated information in the

environment to gain future rewards (Hangya et al., 2015; Teles-Grilo Ruivo et al., 2017). An alternative way to conceptualize the role of acetylcholine is as a signal for uncertainty, which may be resolved by familiarization through learning (Yu and Dayan, 2005). In either scenario, salience is indicated by increasing the release of acetylcholine in the hippocampus and other brain areas.

The dentate gyrus receives excitatory glutamatergic input from layer II of the medial entorhinal cortex, and sparsifies this signal by suppressing the activity of most dentate gyrus granule cells through lateral inhibition while dramatically increasing the firing rate of a select few granule cells (O'Reilly and McClelland, 1994). By this mechanism granule cells detect salient, novel information and accentuate minor contextual details related to familiar information (a process often referred to as pattern separation). Individual granule cells provide a strong, sparse, facilitating input to a small number of CA3 pyramidal cells that can be sufficiently powerful to engage 1:1 spike transfer after multiple spikes in a granule cell burst (Acsady et al., 1998; Henze et al., 2002; Sachidhanandam et al., 2009; Vyleta et al., 2016). This focal excitation by mossy fibers drives synchronisation between subsets of CA3 pyramidal cells potentially allowing recurrent CA3-CA3 synapses to engage Hebbian plasticity mechanisms to create ensembles of strongly coupled CA3 cells thereby initiating the storage of new information (Brandalise and Gerber, 2014; Guzman et al., 2016; Kobayashi and Poo, 2004; Mishra et al., 2016; O'Reilly and McClelland, 1994; Treves and Rolls, 1994). Mossy fibers also excite a broad and diverse set of inhibitory interneurons that provide a widespread 'blanket' of feed-forward inhibition over a large population of CA3 pyramidal cells (Acsady et al., 1998; Mori et al., 2007; Szabadics and Soltesz, 2009; Toth et al., 2000). This feed-forward inhibition prevents runaway excitation and ensures tight spike timing for spike transfer (Restivo et al., 2015; Torborg et al., 2010; Zucca et al., 2017), while also enhancing memory precision (Ruediger et al., 2011) but it is not known what impact it may have on synaptic plasticity within the CA3 recurrent network.

Here we investigate the modulation by acetylcholine of ensemble creation and therefore memory encoding within the hippocampal CA3 network. Using a hierarchy of experimentally constrained computational models of mossy fiber synaptic transmission and CA3 network activity, we demonstrate that acetylcholine suppresses feed-forward inhibition enabling plasticity at recurrent CA3-CA3 synapses and the formation of

ensembles within the CA3 network. Furthermore, we show that acetylcholine increases the density of stable ensembles by enhancing permissible overlap between ensembles.

Results

Acetylcholine regulates CA3-CA3 recurrent synaptic inputs and cellular excitability
(Dasari and Gulledge, 2011; Hasselmo et al., 1995; Vogt and Regehr, 2001) both of which
are likely important for the creation of CA3 ensembles but a critical feature of ensemble
formation is the mossy fiber input from dentate gyrus. However, the summative effects of
acetylcholine on excitatory and inhibitory mossy fiber synaptic transmission are not
known. Therefore, we first recorded experimentally the effect of acetylcholine on combined
feed-forward excitatory and inhibitory synaptic transmission in the mossy fiber pathway.
Minimal stimulation of granule cells resulted in EPSCs and IPSCs that were individually
isolated by setting the membrane potential to -70mV and +10mV respectively in
accordance with experimentally determined reversal potentials for inhibitory and
excitatory transmission respectively (Supplementary Fig. 1). Mossy fibers were stimulated
with trains of 4 pulses at 20 Hz every 20 s. Application of the group II mGluR agonist
DCG-IV (1 µM) reduced EPSC amplitudes by >90% (Fig. 1B) indicating selective
activation of the mossy fiber pathway (Kamiya et al., 1996). The EPSC rise times (20-
80%), latencies, and jitter were 0.57 ± 0.11 ms, 2.1 ± 0.49 ms and 0.53 ± 0.27 ms
respectively (Supplementary Fig. 1) characteristic of mossy fiber synapses and confirming
their monosynaptic origin (Nicoll and Schmitz, 2005). DCG-IV also reduced IPSC
amplitudes by >90% and the rise times, latencies and jitter were 3.52 ± 1.08 ms, $6.20 \pm
1.82$ ms and 0.73 ± 0.25 ms respectively (Supplementary Fig. 1) indicating that IPSCs
were mediated by disynaptic feed-forward inhibitory transmission in the mossy fiber
pathway (Torborg et al., 2010). Both EPSCs and IPSCs exhibited pronounced facilitation
in response to a train of 4 stimuli at 20 Hz as previously shown for mossy fiber feed-
forward excitatory and inhibitory pathways (Torborg et al., 2010).

Acetylcholine reduces feed-forward inhibition in the mossy fiber pathway

The impact of acetylcholine on information transfer between the dentate gyrus and CA3
will depend on its effects on both excitatory and inhibitory pathways. To assess the effect
of acetylcholine on both pathways we used the broad- spectrum cholinergic receptor
agonist carbachol (CCh). Application of 5 µM CCh depressed EPSC amplitudes by a small
amount (Fig. 1C; $70.4 \pm 22.9\%$, n = 7, P = 0.035) without altering facilitation ratios (Fig.
1C; P=0.509; Measured at 1^{st} to 4^{th} pulse). This depression did not recover on washout of

CCh. In contrast, the same dose of CCh reversibly depressed IPSC amplitudes to a much greater degree (Fig. 1D; 26.7 $\pm$ 42.0%; n = 6, P = 0.012) and at the same time reduced facilitation ratios (Fig 1D; P = 0.012; measured at 1st to 4th pulse). The use of minimal stimulation meant that responses to the first stimuli were highly variable and often very small or absent due to the low basal probability of release at mossy fiber synapses (Nicoll and Schmitz, 2005; Sachidhanandam et al., 2009; Torborg et al., 2010). This was particularly true for IPSC recordings resulting in the highly variable effect of CCh on the first IPSC in a train. CCh also enhanced the excitability of neurons in the CA3 network indicated by an increase in the frequency of spontaneous EPSCs and IPSCs (sEPSCs: 14.68 $\pm$ 11.93 Hz to 34.68 $\pm$ 32.06 Hz, p = 0.077; sIPSCs 13.27 $\pm$ 3.23 Hz to 29.65 $\pm$ 21.76 Hz, p = 0.126). CA3 pyramidal cells also showed a trend towards a decrease in the input resistance (340.5 $\pm$ 176.8 MΩ to 227.4 $\pm$ 120.2 MΩ, p = 0.106) but no change in holding current at -70 mV (-141.5 $\pm$ 67.5 pA to -143.6 $\pm$ 62.2 pA) (Cheng and Yakel, 2014; Vogt and Regehr, 2001). At lower concentrations, 1 µM CCh had limited effect on IPSC amplitudes whereas at higher concentrations 10 µM CCh had similar effects to 5 µM CCh with a substantial depression of IPSC amplitudes (Supplementary Fig. 1). These results indicate that acetylcholine causes a small irreversible depression of excitatory transmission at mossy fiber synapses through a likely postsynaptic mechanism (Dickinson et al., 2009; Vogt and Regehr, 2001) whereas feed-forward inhibitory transmission is substantially depressed. Overall, this indicates a net enhancement of Excitatory-Inhibitory ratio in the mossy fiber pathway.

Effects of acetylcholine on short-term plasticity at the mossy fiber synapse

Information transfer between the dentate gyrus and CA3 network depends on bursts of high frequency activity in dentate granule cells leading to pronounced frequency facilitation of excitatory synaptic input (Henze et al., 2002; Vyleta et al., 2016; Zucca et al., 2017). This is balanced by frequency-dependent facilitation of inhibitory synaptic input (Torborg et al., 2010) but variations in the short-term plasticity dynamics between the excitatory and inhibitory pathways will lead to windows within the frequency domain when excitation dominates and action potentials are triggered in CA3 pyramidal cells (Mori et al., 2004). However, these temporal windows have not been fully characterized and, furthermore, the effect of acetylcholine on Excitatory-Inhibitory ratio over a range of

mossy fiber stimulation patterns is not known. To investigate the patterns of activity that trigger action potentials we adapted a Tsodyks-Markram based model of short-term plasticity dynamics in both excitatory and inhibitory pathways (see Methods).

Short-term plasticity models are difficult to constrain with responses evoked by regular stimulation protocols (Costa et al., 2013). Therefore, we constrained the model using responses to a stimulation pattern resembling the natural spike statistics of dentate gyrus granule cells which incorporate a broad range of inter-stimulus intervals (ISIs) (Fig 2A) (Mistry et al., 2011). Similar to the regular stimulation pattern of 4 stimuli at 20Hz, CCh depressed EPSCs and IPSCs in response to the irregular stimulation pattern across the range of ISIs but the depression was much more pronounced for IPSCs (Fig 2B). Several short-term plasticity models of increasing level of complexity for both excitatory and inhibitory synaptic responses were assessed for fit to the experimental data (see Methods for detailed description of these models). The basic form of these models includes a facitation and a depression variable, here represented as f and d respectively. Dynamics for these variables are governed by parameters for degree (a) and timecourse of facilitation (τ_f) and depression (τ_d) as well as baseline of release (f_0) and synaptic conductance (g) (Fig 2B. Parameter inference for the short-term plasticity models was carried out and the best fitting models were selected by comparing the Akaike and Bayesian Information Criteria (AIC and BIC respectively) weights. These weights represent, a normalisation of AIC and BIC values calculated by dividing the AIC and BIC values by the sum of these values across all models (log-likelihood of model given data punished for increasing complexity in two different ways). This is convenient as it allows these values to be transformed into a probability space and hence comparable across samples (Wagenmakers and Farrell, 2004). The model with the highest weight explains the data best. For excitatory mossy fiber synaptic transmission, a model containing a single facilitating variable with an exponent of 2 (f^2) best explained the experimental data (Fig 2D).

$$\text{EPSC amplitude} = g_E^{max}\, f^2\, (V_{mem} - E_{GLUT}) \qquad \text{(Equation 1)}$$

$$\frac{df}{dt} = \frac{f_0 - f}{\tau_f} + a\,(1-f)\sum_s \delta(t - t_s) \qquad \text{(Equation 2)}$$

where V_{mem} is the holding voltage of the cell in voltage clamp, g_E^{max} is the maximum excitatory conductance, E_{GLUT} is the reversal potential of glutamatergic transmission, and t_s is the timing of the s^{th} spike (or pulse). Explanation of other parameters for the short-term plasticity model is given in the Methods section.

It is noticeable that AIC and BIC weights disagree on which model best explains the data. The f^2 model had only the second highest AIC weight, but had the highest BIC weight, whereas the more complex af model had a higher AIC weight. However, the evidence ratio for BIC points favours the f^2 model ($P(f^2|\text{Data})/P(af|\text{Data}) = 8.92$ (see Kass & Raftery, 1995)), whereas the evidence ratio for AIC indicates little evidence in favour of the af model ($P(af|\text{Data})/P(f^2|\text{Data}) = 1.51$). Together this indicates that the f^2 model best explains the data.

For inhibitory feed-forward mossy fiber synaptic transmission, a complex model with facilitation (f), depression (d), and additional facilitation over the increment parameter a (afd) produced the best fit (Fig 2E) with both AIC and BIC weights convincingly pointing to the afd model as most appropriate to describe the data. Additional parameters in this model included a time course for facilitation of a (τ_a), an increment scaling factor for a (b), and baseline (a_0).

$$\text{IPSC amplitude} = g_I^{MAX} f d \left(V_{mem} - E_{GABA}\right) \qquad \text{(Equation 3)}$$

$$\frac{df}{dt} = \frac{f_0 - f}{\tau_f} + a\left(1 - f\right)\sum_s \delta(t - t_s) \qquad \text{(Equation 4)}$$

$$\frac{dd}{dt} = \frac{1 - d}{\tau_d} - f d \sum_s \delta(t - t_s) \qquad \text{(Equation 5)}$$

$$\frac{da}{dt} = \frac{a_0 - a}{\tau_a} + b\left(1 - a\right)\sum_s \delta(t - t_s) \qquad \text{(Equation 6)}$$

where g_I^{max} is maximum inhibitory conductance and E_{GABA} is the reversal potential of GABAergic transmission. See Methods for detailed explanation of the above equations.

Discrepancies between samples drawn from posterior-predictive distributions of these models indicated good fit for both models (Supplementary Fig. 2).

Using these models for the activity-dependent progression of excitatory and inhibitory synaptic weights we were then able to investigate the effect of acetylcholine on short-term plasticity by comparing normalized parameter estimates to time matched controls. Since posterior distributions for EPSC data were narrow and unimodal, maximum a posteriori (MAP) estimates were used, whereas mean parameter estimates were used for IPSC data since posterior distributions were wide and bimodal in some cases (Supplementary Fig 2). This analysis revealed the small decrease in EPSC amplitude caused by CCh resulted from a reduction in the conductance scaling parameter 'g' (Fig 2D; 49.9 $\pm$ 17.6% decreae in 'g'') indicating a postsynaptic mechanism, supporting the original suggestion based on the data in Fig 1C. The substantial decrease in IPSC amplitude caused by CCh resulted from a large reduction in the conductance scaling parameter 'g', and an increase in the baseline parameter 'f0' which also had the effect of reducing facilitation (Fig 2E; 73.1 $\pm$ 27.9% decrease in 'g '; 225.7 $\pm$ 160.1 % increase in 'p0''). Since IPSCs are disynaptic, it is not straightforward to interpret how these parameter changes reflect biophysical changes to synaptic transmission. Nevertheless, these results indicate the mechanisms underlying the effects of CCh on mossy fiber synaptic transmission and enable investigation of the granule cell spike patterns that favor excitation over inhibition.

Enhancement of mossy fiber Excitatory-Inhibitory balance by acetylcholine

Feed-forward inhibition dominates excitation in the mossy fiber pathway for the majority of spike patterns (Mori et al., 2004; Torborg et al., 2010). Since acetylcholine depresses inhibitory transmission more than excitatory transmission (Figs 1 and 2) it is expected that the Excitatory-Inhibitory balance will be shifted towards excitation but the precise spike patterns that this occurs for are unclear. To examine how Excitatory-Inhibitory balance is affected by carbachol with different spike patterns we first tested the dependence of short-term synaptic dynamics on background firing rate using the model for short-term plasticity dynamics. Spike patterns were described by two parameters: a between burst interval $\Delta t_{between}$ describing a background firing rate, and a within burst interval Δt_{within} describing the time between spikes in a burst. The steady state value of *f*,

d, and *a* given $\Delta t_{between}$ were then used to replace their baseline values ($a_0 \rightarrow a_\infty$; $f_0 \rightarrow f_\infty$; $d_0 = 1 \rightarrow d_\infty$) to set their initial values at the beginning of a burst, i.e.,

$$a_\infty = \frac{a_0 exp(\Delta t_{between}/\tau_a) - a_0 + b}{\exp(\Delta t_{between}/\tau_a) - 1 + b} \quad \text{(Equation 7)}$$

$$f_\infty = \frac{f_0 \exp(\Delta t_{between}/\tau_f) - f_0 + a_\infty}{\exp(\Delta t_{between}/\tau_f) - 1 + a_\infty} \quad \text{(Equation 8)}$$

$$d_\infty = \frac{1 - \exp(-\Delta t_{between}/\tau_d)}{1 + (1 - f_\infty) \exp(-\Delta t_{between}/\tau_d)} \quad \text{(Equation 9)}$$

By systematically varying the between and within burst intervals for both excitatory and inhibitory synaptic input we were able to simulate EPSCs and IPSCs in the presence and absence of acetylcholine (Fig 3A). The amplitudes of these responses were then used to explore the effects of acetylcholine on within burst facilitation ratios and Excitatory-Inhibitory balance across a wide range of between and within burst intervals.

Experimental data shows that mossy fiber EPSCs are exquisitely sensitive to between burst interval with facilitation revealed as between burst interval is decreased. Furthermore, it has been shown that shortening the between burst interval decreases between burst facilitation (Salin et al., 1996). Our simulations replicate this interdependence of between and within burst interval with respect to EPSC facilitation with values closely associated with the experimental data in the literature (Salin et al., 1996; Toth et al., 2000) (Fig 3B). The CCh induced depression of EPSCs was mediated by a reduction in synaptic conductance and therefore did not alter either between or within burst synaptic facilitation.

The situation for inhibitory synaptic transmission is more complex. Over the course of a burst synaptic amplitude facilitation is greatest when between and within burst intervals are largest. As between and within burst intervals reduce the facilitation morphs into a depression towards the end of the burst resulting in limited inhibition at the end of high frequency bursts (Fig 3C). CCh dramatically reduces both the initial IPSC amplitude and subsequent facilitation within bursts at all between and within burst intervals (Fig 3C).

We then combined the results from excitatory and inhibitory facilitation to predict EPSC-IPSC amplitude ratios over the course of a burst. In control conditions, excitation dominates over inhibition only after multiple spikes in a burst and when bursts occur at shorter between and within burst intervals (Fig 3D) (Mori et al., 2004; Torborg et al., 2010; Zucca et al., 2017). However, in the presence of CCh, excitation dominates over inhibition at earlier stimuli within the burst, and over longer between and within burst intervals meaning cholinergic receptor activation allows excitation to dominate over inhibition for a broader range of stimulus patterns.

We next investigated the biophysical effects of the acetylcholine-induced reduction in feed-forward inhibitory synaptic transmission at mossy fiber synapses. In particular, the modulation of back propagating action potentials and EPSPs in CA3 pyramidal cells that are critical for the induction of long-term potentiation (LTP) at recurrent CA3-CA3 synapses and therefore the formation of CA3 ensembles (Brandalise and Gerber, 2014; Guzman et al., 2016; Mishra et al., 2016). Mossy fibers provide powerful excitatory drive to the soma of CA3 pyramidal cells and have been referred to as 'conditional detonator' synapses because a single synapse can trigger postsynaptic action potentials in response to high frequency bursts, but not single, presynaptic action potentials (Henze et al., 2002; Sachidhanandam et al., 2009; Vyleta et al., 2016). We hypothesized that feed-forward inhibitory synaptic transmission reduces back propagating action potentials and mossy fiber evoked EPSPs which will inhibit or prevent the induction of LTP (Mullner et al., 2015; Tsubokawa and Ross, 1996; Wilmes et al., 2016), and that acetylcholine will relieve this inhibition by reducing feed-forward inhibition. To test this we used a well characterized multi-compartment biophysical model of a CA3 pyramidal cell (Hemond et al., 2008; Henze, Cameron, & Barrionuevo, 1996) which incorporated mossy fiber excitatory synaptic input on the very proximal portion of the apical dendrite and feed-forward inhibitory synapses distributed across both somatic and dendritic compartments (Fig 4A) (Szabadics and Soltesz, 2009). Membrane potential and the resultant intracellular calcium concentration were simulated across multiple somatic and dendritic compartments of a reconstructed CA3 pyramidal cell incorporating the thin oblique dendrites in stratum radiatum where the majority of CA3-CA3 recurrent synapses are located (Major, Larkman, Jonas, Sakmann, & Jack, 1994). With feed-forward inhibition intact, action potentials and EPSPs back propagate into the principle dendritic shafts without much change in amplitude but are rapidly attenuated on entering the thin oblique

dendrites (Fig 4B and C). This leads to minimal calcium influx through voltage- gated calcium channels at these dendritic sites (Fig 4B). However, when feed-forward inhibition is reduced by acetylcholine, there is much less attenuation of action potentials and EPSPs allowing substantial calcium influx (Fig 4B and C). The disinhibition of mossy fiber feed-forward inhibition by acetylcholine has major implications for synaptic plasticity at CA3-CA3 recurrent synapses, since spike timing-dependent plasticity is dependent on the back propagation of action potentials and EPSPs, postsynaptic calcium accumulation and activation of calcium-dependent signalling pathways (Brandalise and Gerber, 2014; Mishra et al., 2016). Our simulations indicate that disinhibition by acetylcholine is a requirement for the induction of synaptic plasticity between CA3 pyramidal cells when CA3 ensembles are activated by mossy fiber inputs.

Ensemble formation in CA3 driven by mossy fiber input

To investigate the known effects of acetylcholine on the creation of CA3 ensembles by mossy fiber input we next turned to a spiking network model of CA3. This network was comprised of point neurons with Izhikevich-type dynamics (Izhikevich, 2003) parameterized to reproduce spiking patterns for excitatory CA3 pyramidal cells and inhibitory fast spiking interneurons (Hummos et al., 2014) connected in an all-to-all fashion. Subsets of pyramidal cells were driven by excitatory mossy fiber input with short-term facilitation. CA3-CA3 recurrent synaptic connections were subject to a symmetric spike timing-dependent plasticity (STDP) rule (Fig 5A) with no short-term plasticity (Mishra et al., 2016). The symmetric STDP rule was modified to allow for depression dependent on the postsynaptic firing rate through synaptic scaling. At low firing rates, potentiation is induced with small differences in pre- and post-synaptic spike times. As the postsynaptic firing rates increase, large differences in pre- and post-synaptic spike times cause depression. At a maximum firing rate, no potentiation is possible.

Within this network we first characterized the speed and stability of ensemble creation where an ensemble was defined as being formed when all synapses between cells within the same ensemble had reached their maximum weight, and all synapses between cells not within the same ensemble had decreased to zero. In addition, the properties of mossy fiber input were studied in comparison with a more generic input reminiscent of perforant path activity during direct information transfer between entorhinal cortex and CA3 to see how

they compared in their ability to drive ensemble formation via synchronous spiking in a small population of cells (Fig 5B). Mossy fiber spike patterns were modelled as a Poisson process with brief (200 ms every 20 seconds) high intensity (50 Hz) firing rates on a very low basal firing rate (0.2 Hz), and were connected to pyramidal cells by a strong facilitating synapse (3.0 nS). This firing pattern represents a strongly separated, sparse firing pattern in a single presynaptic cell, which is expected in dentate gyrus granule cells. Perforant path spike patterns were modelled as a population of presynaptic entorhinal cells in the synchronous irregular state modelled as 120 homogeneous Poisson processes firing at 10 Hz with a correlation coefficient of 0.9 and static, weak synapses fixed at 0.1 nS, which broadly reflects entorhinal activity in a freely behaving rat (Chrobak & Buzsáki, 1998).

We initially built up the network model sequentially to investigate which components were necessary for the speed and stability of ensemble creation. At first, two CA3 pyramidal cells were connected and driven with only excitatory input. For both mossy fiber and perforant path inputs the cells quickly became connected (Fig 5B, top row). Increasing ensemble size to 10 excitatory cells destabilised the ensemble formation process (Fig 5B, middle row). The destabilization resulted from unbalanced potentiation of recurrent excitation that caused a large increase in the firing rate leading to strong depression or 'resetting' of the synaptic weight with further spiking as a result of synaptic scaling. The addition of 5 feedback inhibitory cells stabilised ensemble formation in the case of mossy fiber input, but not for perforant path input (Fig 5B, bottom row). This was because the perforant path input provided a colored noise signal amplified by recurrent excitation that caused excitatory cells to fire at high rates too often and feedback inhibition was insufficient to counter this amplification. Mossy fiber input is driven only briefly at sparse intervals, meaning there was little opportunity to exceed target firing rate, and when there was, feedback inhibition was sufficient to contain it. These results show that mossy fiber-like sparse inputs and feedback inhibition within the CA3 recurrent network are important for rapid and stable formation of CA3 ensembles.

The advantages of short-term facilitation were also explored. To model how input from multiple granule cells is transmitted to a single pyramidal cell, a single excitatory input layer network to one excitatory cell was constructed (Fig 5C). Inputs followed an inhomogeneous Poisson process, each of which was connected to the cell with a facilitating mossy fiber synapse, or with a synapse with a fixed conductance at half the maximum

weight of the facilitating synapse. Inhomogeneous rates were defined by a winner take all process (Fukai and Tanaka, 1997) with adaptation (see methods). For good transmission of a separated (winner) pattern, the postsynaptic cell should only spike (Fig 5C; red trace) when the winning input is the only input with a non-zero weight i.e., once the winner has been selected (Fig 5C; grey traces). The efficacy of transmission is defined as the rate of the winning input r_N at the time of the postsynaptic spike t_{spike} divided through by the sum of rates r of all inputs at the time of the postsynaptic spike, i.e.,

$$efficacy = \frac{\sum_{t_{spike}} r_N(t)\delta(t - t_{spike})}{\sum_i \sum_{t_{spike}} r_i(t)\delta(t - t_{spike})} \qquad \text{(Equation 10)}$$

The population of input rates were defined by two parameters: scale and separation, where the scale parameter is a scaling factor that multiples each rate in the population by the same amount, whereas the separation parameter controls the difference in input to the population of rates, determining how much the winner wins by. Increasing both the scales and separation of input rates produced a sharper transition between no transmission (efficacy = 0) and perfect transmission (efficacy = 1) when synapses were facilitating (Fig 5C). However, when synaptic conductance was fixed, there was a smoother transition from no transmission to perfect transmission. This indicates that short-term facilitation supports robust transmission of separated patterns, as weakly separated patterns can still be transmitted perfectly. In contrast, weakly separated patterns transmit more ambiguously when input is of fixed conductance. This is because facilitating synapses place very low weight on signals at the start of a winner-take-all process, but much greater weight at the end making spiking more likely once the winner has been chosen, whereas constant synapses place too much weight before a winner has been selected and can cause postsynaptic cells to fire too early.

Acetylcholine facilitates ensemble formation in CA3

We next examined how acetylcholine affects the CA3 network's ability to form ensembles. In addition to facilitating plasticity at CA3-CA3 recurrent synapses by control of feed-forward inhibition, two well established effects of acetycholine in CA3 are to increase cellular excitability and reduce the overall conductance of CA3-CA3 recurrent synapses by

reducing the probability of glutamate release (Dasari and Gulledge, 2011; Hasselmo et al., 1995; Vogt and Regehr, 2001). We implemented these effects of acetylcholine within the spiking network model by depolarizing the resting membrane potential for excitatory and inhibitory cells to -70 mV and -63 mV respectively and halved the CA3-CA3 synaptic conductance (Hummos et al., 2014). Networks contained 64 excitatory and 16 inhibitory cells with excitatory cells grouped into 8 ensembles consisting of 8 cells each (Fig 6A). Each cell within a single ensemble received the same mossy fiber input, which followed Poisson process with a low firing rate of 0.2 Hz punctuated by bursts for 250 ms every 20 s at varying frequency. No two ensembles received bursts at the same time and spike timing-dependent plasticity rules were implemented regardless of the presence of acetylcholine.

Without acetylcholine, bursts at a frequency of 30 Hz were capable of forming discrete ensembles, but these were almost completely abolished when burst frequency was reduced to 20 Hz (Fig 6B). Remarkably, acetylcholine rescued ensemble formation at the lower burst frequency. To quantify network ensemble formation performance and the impact of acetylcholine in greater detail, we used an error metric (WME) defined as the summed absolute difference between target and actual weight matrix (W_{ij}^{actual}), where the target weight matrix (W_{ij}^{target}) is maximum synaptic weights between cells i and j in the same ensemble, and zero weight otherwise, i.e.,

$$WME = \sum_{ij} |W_{ij}^{actual} - W_{ij}^{target}| \qquad \text{(Equation 11)}$$

This analysis revealed that acetylcholine lowers the frequency and increases the speed at which ensembles form (Fig 6C). To test which effects of acetylcholine were critical for these aspects of ensemble formation we removed each parameter change in turn. This revealed that the key factor was the increase in cellular excitability, as removing the parameter changes to cellular excitability abolished the effect of acetylcholine, but removing reductions in CA3-CA3 recurrent connections did not (Fig 6C and Supplementary Fig. 3).

Within the CA3 network multiple often highly overlapping ensembles may be encoded. Theoretically this increases the capacity of information encoding but reduces the fidelity of

retrieval with a necessary trade-off between these two parameters. Therefore, we investigated the impact of acetylcholine on the ability of the CA3 network to reliably encode overlapping ensembles. To incorporate overlap between ensembles the total network size was made variable whilst still containing 8 ensembles of 8 cells each and overlap was introduced by having a subset of excitatory cells receive input from two sources rather than one. Overlap was arranged in a ringed fashion such that each ensemble shared a certain number of cells with their adjacent 'neighbour' (Fig 7A). Retrieval was studied by comparing the population rates of each ensemble, with a smaller difference in rates meaning lower discrimination and more difficult retrieval.

Without acetylcholine present, stable ensembles could be formed when the degree of overlap was low but discrete ensembles could not be formed with levels of overlap >1 (Fig 7B and C). In contrast, in the presence of acetylcholine the network could safely support an overlap of 3 cells (Fig 7B and C). To test which effects of acetylcholine mediated the enhanced discrimination between overlapping ensembles we removed each parameter change in turn. This revealed that the key factor was the reduction in CA3-CA3 recurrent synapse efficacy since removing this parameter change abolished the ability for acetylcholine to allow greater ensemble overlap whereas removing the increase in cellular excitability only increased the time taken to form stable ensembles without affecting the final degree of overlap supported (Fig 7C and Supplementary Fig. 3). Interestingly, the increase in stable ensembles with significant overlap was associated with a decrease in the discrimination between ensembles during retrieval of information as more overlap produced less separation of population rates between ensembles (Fig 7D). Taken together these data indicate that acetylcholine increases the number of discrete ensembles that may be contained within a finite CA3 network but that this comes at a cost of reduced retrieval fidelity.

Discussion

In this study, we investigated the effects of acetylcholine on the formation of ensembles within the CA3 recurrent network. Experimentally, we discovered that acetylcholine dramatically reduces feed-forward inhibition in the mossy fiber pathway whilst having limited effect on mossy fiber excitatory transmission (Figs 1&2). This disinhibition causes a strong positive shift in the Excitatory-Inhibitory balance (Fig 3) resulting in a greatly enhanced potential for LTP at recurrent CA3-CA3 synapses and therefore ensemble formation (Fig 4). In addition, in a computational spiking network model, the separate effects of acetylcholine on cellular excitability and basal CA3-CA3 synaptic strength were found respectively to enhance the robustness of ensemble formation (Fig 6) and the amount of allowable overlap between ensembles (Fig 7). Together, these findings indicate a central role for acetylcholine in gating and facilitating memory formation in the CA3 network and suggest that cholinergic activity may provide an important salience cue to signal which information to encode.

The three separate mechanisms we identify as important for ensemble formation are likely supported by different cholinergic receptors at distinct cellular and subcellular locations. Our experimental data show only a small effect of cholinergic agonists at excitatory mossy fiber synapses onto CA3 pyramidal cells indicating a limited direct synaptic modulation by muscarinic or nicotinic receptors mediated by postsynaptic changes (Dickinson et al., 2009; Vogt and Regehr, 2001; Williams and Johnston, 1990). The lack of presynaptic changes indicates no indirect mechanism via enhancement of interneuron spiking and activation of presynaptic GABAB receptors by GABA spillover (Scanziani, 2000; Vogt and Regehr, 2001). For similar reasons, our data do not support a role for presynaptic nicotinic receptors ((Cheng and Yakel, 2014) but see (Vogt and Regehr, 2001)) or an increase in dentate granule cell spike frequency (Vogt and Regehr, 2001) since both would be expected to cause an increase in EPSC amplitude by presynaptic mechanisms. The large depression in feed-forward inhibitory mossy fiber transmission by cholinergic activity most likely results from a combination of an increase in feed-forward interneuron excitability and spike rate, mediated by a combination of muscarinic M1 and M3 receptors and nicotinic receptors (Cea-del Rio et al., 2010; Dasari and Gulledge, 2011; Vogt and Regehr, 2001), coupled with a strong depression of GABA release, mediated by presynaptic M2 receptors present on interneuron terminals (Szabo et al., 2010). This potentially accounts for the

observed increase in <mark>basal synaptic response</mark> but overall large reduction in synaptic conductance seen in our experimental data and short-term plasticity model. Finally, the depression in basal CA3-CA3 recurrent synaptic strength is reported to result from the activation of presynaptic M4 receptors (Dasari and Gulledge, 2011). These mechanistic conclusions predict that M2 muscarinic receptors on interneuron terminals are important for disinhibition of mossy fiber feed-forward inhibition necessary for ensemble formation, M4 muscarinic receptors on CA3-CA3 recurrent axon terminals are important for increasing the amount of permissible overlap between ensembles and M1 muscarinic receptors on CA3 pyramidal cells facilitate the rapid and stable formation of ensembles.

Feed-forward inhibition dominates excitatory transmission in the mossy fiber pathway but unlike other examples of feed-forward inhibition, such as that occurring in the neocortex or CA1 region of the hippocampus, interneurons engaged with mossy fiber feed-forward inhibition target the dendritic compartments of CA3 pyramidal cells as much if not more than the perisomatic areas (Pelkey et al., 2005; Szabadics and Soltesz, 2009; Torborg et al., 2010; Toth et al., 2000). This means that mossy fiber feed-forward inhibition strongly inhibits recurrent CA3-CA3 inputs in stratum radiatum rather than excitatory mossy fiber inputs (Miles et al., 1996; Pouille and Scanziani, 2001). We show that inhibitory input to dendritic domains in stratum radiatum strongly attenuates the back propagation of action potentials and EPSPs originating from the somatic compartment into the thin radial oblique dendrites where most of the recurrent CA3-CA3 synapses occur (Fig 4). Since these back-propagating signals are necessary for the induction of LTP at recurrent CA3-CA3 synapses (Brandalise and Gerber, 2014; Mishra et al., 2016), this indicates that mossy fiber feed-forward inhibition is well placed to control the induction of LTP at these synapses. The dominance of mossy fiber feed-forward inhibition may be partially reduced with high frequency burst stimulation where feed-forward inhibition does not facilitate as strongly as excitation enabling excitation to dominate (Mori et al., 2004) (Fig 3) although this is not the case during development of the mossy fiber pathway (Torborg et al., 2010). Here, the remarkable finding is that cholinergic activation reduces mossy fiber feed-forward inhibition by >70% (Fig 1&2) removing the attenuation of back propagating action potentials and EPSPs (Fig 4) and therefore enabling LTP at CA3-CA3 recurrent synapses.

Further investigation revealed that not only does acetylcholine enable ensemble formation by reducing mossy fiber feed-forward inhibition but it also alters the properties of the CA3

network to allow ensemble formation to occur rapidly and robustly with a high degree of overlap between ensembles. Enhancing cellular excitability within an attractor network such as the recurrent CA3 network increases the speed and robustness of synaptic plasticity due to increased spiking during ensemble activity (Fig 6) (ref). For stable ensemble formation and network configuration the increased excitability must be regulated by feedback inhibition (Fig 5). Theoretically, the reduction in CA3-CA3 synaptic efficacy caused by acetylcholine (Hasselmo et al., 1995) might be predicted to reduce the efficiency of ensemble formation but paradoxically we found that this effect of acetylcholine enabled a greater overlap between ensembles whilst still maintaining their discrete identity (Fig 7). This is important for a couple of reasons: i) it allows an increased density of discrete ensembles to be encoded which in a finite network will increase its capacity to store information (ref), and ii) an increase in overlap between ensembles has been shown to facilitate memory consolidation and generalization during reactivation of ensembles that occurs during sleep (O'Donnell and Sejnowski, 2014). Interestingly, non-linear dendritic conductances have also been shown to increase the information capacity of the CA3 network (Kaifosh and Losonczy, 2016) and future studies will be needed to investigate how acetylcholine affects non-linear conductances and therefore how these separate mechanisms combine within the hippocampal CA3 network.

Acetylcholine release within the central nervous system has classically been portrayed as a signal for arousal and attention and is strongly associated with learning (Hasselmo and Sarter, 2011; McGaughy et al., 2000; Parikh et al., 2007). This model has been adapted to propose that acetylcholine is released in response to environments where the outcome is uncertain or not as predicted by internal representations (Yu and Dayan, 2005). In such a scenario new information needs to be incorporated into the internal representation to make the environment more familiar and the outcomes more predictable (Hasselmo and Sarter, 2011). Acetylcholine supports this process by enabling the updating of internal representations of the environment or episodic memories. Our data support such a model where acetylcholine reconfigures the dentate gyrus and CA3 microcircuit to enable the formation of memory ensembles within the recurrent CA3 network.

Methods

Ethical approval

All experiments were performed in accordance with the UK Animal Scientific Procedures Act (1986) and local guidance from the Home Office Licensing Team at the University of Bristol. The protocol was approved by the Animal Welfare and Ethics Review Board at the University of Bristol.

Slice Preparation

500 μm thick transverse slices of the hippocampus were prepared from 4-6 week old C57/BL6 mice. After cervical dislocation, brains were removed and submerged in ice-cold cutting solution saturated with oxygen (in mM: 85 NaCl, 75 Sucrose, 2.5 KCl, 25 Glucose, 1.25 NaH2PO4, 4 MgCl2, 0.5 CaCl2, 24 NaHCO3). Each hippocampus was dissected out and mounted onto a cube of agar then glued to the slicing plate such that hippocampi were positioned vertically and cut using a Leica VT1200 vibratome. Slices were then transferred to a holding chamber with oxygenated aCSF (in mM: 119 NaCl, 2.5 KCl, 11 Glucose, 1 NaH2PO4, 26.5 NaHCO3, 1.3 MgSO4, 2.5 CaCl2), incubated for 30 mins at 35°C and left to rest for a further 30 minutes – 5 hours at room temperature.

Electrophysiology

Slices recordings were made in a submerged recording chamber at 33-35°C. CA3 pyramidal cells were visually identified using infra-red differential interference contrast on an Olympus BX-51W1 microscope. Recording pipettes with resistance 2-4 MΩ were pulled from borosilicate filamented glass capillaries and filled with a caesium-based intracellular solution (in mM: 130 CsMeSO3, 4 NaCl, 10 HEPES, 0.5 EGTA, 10 TEA, 1 QX-314 chloride, 2 MgATP, 0.5 NaGTP). Series resistances were continuously monitored and recordings discarded if series resistance > 25 MΩ or changed by >50%. Recordings were collected using a Multiclamp 700A amplifier (Molecular Devices) filtered at 4 kHz and sampled at 10 or 25 kHz using Signal or Spike2 acquisition software, and a CED Power 1401 data acquisition board.

Postsynaptic currents were evoked by placing monopolar stimulation electrodes in the granular layer of the dentate gyrus and applying 200μs pulses driven by a Digitimer DS2A Isolated stimulator. Stimulating the granular layer avoids the risk of contamination from perforant path inputs, associated commissurals, or monosynaptic inhibitory synapses. Excitatory currents were obtained by holding the cell in voltage-clamp mode at -70 mV, while inhibitory currents were obtained by holding at +10 mV. Candidate mossy fiber driven responses were chosen based on their latency (2-3 ms for excitatory, 5-10 ms for inhibitory). Excitatory currents were also selected based on kinetics (< 1 ms 20-80% rise time) and short-term facilitation in response to 4 pulses at 20 Hz. Stimulation strength was calibrated to the minimum strength that evoked a response (minimal stimulation). Excitatory and inhibitory responses were blocked $> 80\%$ by 1 μM DCG-IV in 100% cases (Fig 1; n=9 MF-EPSCs, n=6 MF-IPSCs), therefore it can be inferred that this approach reliably stimulated mossy fibers (Kamiya et al., 1996). Carbachol (CCh) was bath applied for 5-10 minutes before measuring effect on synaptic transmission. Effects of carbachol were normalised to control values taken in first 3 minutes of the experiment.

Short-Term Plasticity Model

The Tsodyks-Markram model (Hennig, 2013; Tsodyks & Markram, 1997) was adopted due to its widespread use, simplicity, and relation to biophysics. This model captures pre-synaptic release dynamics with two variables, a facilitating process *f* and a depressing process *d*, that represent 'resources' available to drive synaptic transmission. These evolve as follows:

$$\frac{df}{dt} = \frac{f_0 - f}{\tau_f} + a\,(1 - f) \sum_s \delta(t - t_s) \qquad \text{(Equation 12)}$$

$$\frac{dd}{dt} = \frac{1 - d}{\tau_d} - f\,d \sum_s \delta(t - t_s) \qquad \text{(Equation 13)}$$

Dynamics of *f* is governed by three parameters: f_0 (baseline value of *f*), τ_f (decay time constant of *f*), and *a* (increment scaling factor with incoming spike at time t_s). This variable loosely represents the build-up of free calcium ions in the presynaptic terminal that triggers exocytosis of neurotransmitter-containing vesicles into the synaptic cleft. Dynamics of *d* is governed by a single parameter τ_d, and loosely represents the availability

of docked vesicles at release sites. Since these variables are bounded between 0 and 1, the convert into a conductance amplitude these are multiplied by a conductance scaling factor g, i.e.,

$$\text{PSC Amplitude} = g\, f\, d \qquad \text{(Equation 14)}$$

This basic model can be extended to more complex facilitation models for example by making parameters f_0 and a time dependent (Hennig, 2013), i.e.,

$$\frac{df_0}{dt} = \frac{\widetilde{f_0} - f_0}{\tau_{f_0}} + b_{f_0}\left(1 - f_0\right) \sum_s \delta(t - t_s) \qquad \text{(Equation 15)}$$

$$\frac{da}{dt} = \frac{a_0 - a}{\tau_a} + b_a\left(1 - a\right) \sum_s \delta(t - t_s) \qquad \text{(Equation 16)}$$

where $\widetilde{f}_0$ is a baseline for f_0, a_0 is the baseline for a, τ_{f_0} is the time constant for f0, τ_a is the time constant for a, b_{f_0} is the increment scaling factor for f_0, and b_a is the increment scaling factor for a. It can also be reduced by making f or d constant. Since the mossy fiber synapse is well known for its large pool of readily releasable vesicles that can be quickly replenished, a simple reduction is to keep d constant at 1, which is true when $\tau_d <<$ min(ISI). Further complexity can be incorporated by allowing multiple independent depressing variables, or by having a time dependent scaling factor

$$\frac{dg}{dt} = -g/\tau_g + k \sum_s \delta(t - t_s) \qquad \text{(Equation 17)}$$

where τ_g is the time constant for short term changes in condutance, and k is an increment scaling factor that can take positive values for facilitation, or negative values for depression.

Naturalistic stimulation patterns

Previous research has shown that short-term facilitation models are difficult to constrain with responses evoked by regular stimulation protocols (Costa, Sjöström, & van Rossum,

2013), and that irregular or naturalistic stimulus trains allowed much better fits to data due to sampling across a broader range of inter-stimulus intervals (ISI) (Costa et al., 2013; Gundlfinger et al., 2007). Dentate gyrus granule cells in vivo have been shown to have bimodal ISI distributions, with long periods of quiescence punctuated by short bursts of action potential firing (Jung & McNaughton, 1993; Mistry, Dennis, Frerking, & Mellor, 2011). This bimodal ISI distribution was modelled as a doubly stochastic cox process to allow generation of stimuli resembling natural spike patterns (Dayan & Abbott, 2001). Each cox process i is defined by a rate parameter λ_{ii}, and a refractoriness parameter σ_i. These two processes are then mixed with responsibility π i.e.,

$$P(\text{ISI}) = \pi\, q_1 + (1 - \pi)\, q_2 \qquad \text{(Equation 18)}$$

$$q_i = r_i \quad \text{if} \quad r_i > x_i \quad \text{else} = 0 \qquad \text{(Equation 19)}$$

$$x_i \sim \mathcal{N}(\mu_i, \sigma_i^2) \qquad \text{(Equation 20)}$$

$$r_i \sim \text{Exp}(\lambda_i) \qquad \text{(Equation 21)}$$

$$\pi \sim \text{Ber}(p) \qquad \text{(Equation 22)}$$

In brief, ISIs were generated by sampling from two exponential distributions with parameters λ_i, which were rejected if they were less than a sample from a normal distribution with standard deviation σ_i. These candidate ISIs were then accepted according to a Bernoulli distribution with probability π. These parameters were set as $\lambda_1 = 3.0$, $\lambda_2 = 0.25$, $\sigma_1 = 0.12$, $\sigma_2 = 0.01$, $\pi = 0.55$. Ninety-nine ISIs were sampled to provide 100 spike times for a stimulation protocol lasting 525 seconds.

Model Fitting

The Bayesian parameter inference procedure used by (Costa et al., 2013) was used to fit parameters to the model. Each model was converted into an iterative form, integrating f and d over each ISI Δt_s between the n^{th} and $n+1^{th}$ spike to output a normalised post-synaptic conductance amplitude for the n^{th} spike in the sequence, i.e.,

$$f_{n+1} = f_0 - (f_0 - f_+) \exp(-\Delta t_s/\tau_f). \quad f_+ = f_n + a\,(1 - f_n) \quad \text{(Equation 23)}$$

$$d_{n+1} = 1 - (1 - d_+) \exp(-\Delta t_s/\tau_d), \quad d_+ = d_n - f_n\, d_n \qquad \text{(Equation 24)}$$

from the original Tsodyks-Markram formalism, however for more complex models in which a and f_0 are time dependent

$$f_{0,n+1} = \tilde{f_0} - (\tilde{f_0} - f_{0,+}) \exp(-\Delta t_s/\tau_{f_0}), \quad f_{0,+} = f_{0,n} + b_{f_0}(1 - f_{0,n}) \quad \text{(Equation 25)}$$

$$a_{n+1} = a_0 - (a_0 - a_+) \exp(-\Delta t_s/\tau_a), \quad a_+ = a_n + b_a(1 - a_n) \quad \text{(Equation 26)}$$

Additionally, when f_0 is time-dependent, Equation 22 becomes

$$f_{n+1} = \frac{1}{\tau_{f_0} - \tau_f}\left(-\tau_{f_0}\tilde{f_0}\exp(-\Delta t/\tau_{f_0}) + \tau_f\tilde{f_0}\exp(-\Delta t/\tau_f) - \tau_f\tilde{f_0} + \tau_{f_0}\tilde{f_0} - \tau_f f_+\exp(-\Delta t/\tau_f) + \tau_{f_0}f_{0+}\exp(-\Delta t/\tau_{f_0}) + \tau_{f_0}f_+\exp(-\Delta t/\tau_f) - \tau_{f_0}f_{0+}\exp(-\Delta t/\tau_f)\right)$$

(Equation 27)

These were then compared to PSC amplitude estimated by the difference between response peak and a baseline taken just before stimulus onset.

Since these equations are deterministic, a likelihood model was constructed where the amplitude was used as parameters for a normal distribution, i.e.,

$$P(D|\theta) = \mathcal{N}(A, A/2) \qquad \text{(Equation 28)}$$

where A is PSC amplitude (Equation 14).

Exponential priors were used for conductance scaling parameters, beta priors for baseline and increment parameters, and uniform priors for time constants. This was to bias conductances towards smaller values and keep baselines low as would be expected from facilitating synapses. Posterior distributions were estimated using Markov Chain Monte Carlo sampling via the Metropolis-Hastings algorithm using the pymc python module.

Model selection was conducted using AIC and BIC weights (Wagenmakers & Farrell, 2004), a transformation of AIC and BIC values into a probability space, with best model having the highest weight. This made it possible to compare the best fitting model over

the population when fitted for each individual sample. For the information criterion of a model for a given sample C_n^m this is defined as

$$w(C) = \frac{\exp(-0.5\Delta_i(C))}{\sum_{k=1}^{K} \exp(-0.5\Delta_k(C))} \qquad \text{(Equation 29)}$$

where

$$\Delta(C) = \tfrac{1}{N} \sum_n [C_n - \operatorname{argmin}_m(C^m)] \qquad \text{(Equation 30)}$$

Goodness-of-fit for the best fitting model was assessed by estimating Bayesian poster-predictive p-values, where samples are drawn from the posterior-predictive distribution and discrepancies $D(x|\theta)$ to expected values e from the model x_{sim} and to the data x_{obs} are compared (Gelman, Meng, & Stern, 1996), i.e.,

$$p = Pr[D(x_{sim}|\theta) > D(x_{obs}|\theta)] \qquad \text{(Equation 31)}$$

where

$$D(x|\theta) = \sum_j (\sqrt{x_j} - \sqrt{e_j})^2 \qquad \text{(Equation 32)}$$

If discrepancies were similar, i.e., $p > 0.025$ or < 0.975, then the model is assessed to fit well. This quantifies how easy it is to discriminate between posterior samples and actual data.

Compartmental Modelling

A 943 compartment reconstruction of a CA3 pyramidal cell with active dendrites (Hemond et al., 2008; Henze, Cameron, & Barrionuevo, 1996) was used to study whether mossy fiber feed-forward inhibition could regulate action potential backpropagation, which would be necessary for feed-forward inhibition to regulate plasticity between recurrent synapses. Simulations were carried out using NEURON.

Active conductances included voltage-gated sodium (Na_V), voltage-activated potassium conductance including delayed rectifier (K_{DR}), M-current (K_M), fast-inactivating A-type (K_A), calcium conductances including N-type (Ca_N), T-type (Ca_T), and L-type (Ca_L),

calcium-activated potassium conductances (K_C and K_{AHP}). Calcium extrusion was modelled as a 100ms decay to a resting Ca^{2+} of 50 nM. Channel kinetics were similar to those used in other hippocampal pyramidal neuron models (Migliore, 2003; Migliore, Cook, Jaffe, Turner, & Johnston, 1995; Migliore, Hoffman, Magee, & Johnston, 1999).

Somatic compartments contained all conductances, dendritic compartments contained all except K_M, and the axonal compartment contained only Na_V, K_{DR}, and K_A. For action potential generation, sodium conductance was five times higher in the axon than the rest of the neuron. Conductances were set as (in $\mu S/cm^2$): $gNa_V = 0.022$, $gK_{DR} = 0.005$, $gK_M = 0.017$, $gK_A = 0.02$, $gCa_N = 0.00001$, $gCa_T = 0.00001$, $gCa_L = 0.00001$, $gK_C = 0.00005$, $gK_{AHP} = 0.0001$.

Dendritic compartments along the apical dendrite were subdivided according to distance (in microns from soma) into those within stratum lucidum (≤ 150), stratum radiatum (>150 or ≤ 400), and stratum lacunosum moleculare (>400). Mossy fiber synapses were targeted towards compartments in stratum lucidum. Feed-forward inhibition was targeted towards somatic compartments (50%) and dendritic compartments in stratum lucidum and stratum radiatum (50%) reflecting the diversity of interneuron subtypes and their targets (Szabadics & Soltesz, 2009).

Synaptic input was modelled using bi-exponential kinetics and short-term plasticity dynamics fit to experimental data. The effect of carbachol was modelled as a three-fold reduction in feed-forward inhibitory conductance.

CA3 Network Modelling

The hippocampal CA3 region was modelled as a small all-to-all recurrent network comprised of excitatory and inhibitory point neurons with adaptive quadratic-integrate-and-fire dynamics with parameters to reflect the firing patterns in response to current injection of CA3 pyramidal cells and fast-spiking basket cells respectively (Hummos, Franklin, & Nair, 2014; Izhikevich, 2003). Continuous membrane dynamics for neuron *i* is described by two equations:

$$C_m \frac{dv_i}{dt} = k \left(v_i - v_r \right) \left(v_i - v_t \right) - u - g_E \left(v_i - v_E \right) - g_I \left(v_i - v_I \right) \quad \text{(Equation 33)}$$

$$\frac{du_i}{dt} = a[b(v_i - v_r) - u_i] \quad \text{(Equation 34)}$$

where v_i describes membrane potential, and u_i is a slow adaptation variable. Parameters for excitatory cells were: $C_m = 24$ pF, $k = 1.5$ pA/mV^2, $a = 10$ Hz, $b = 2$ nS, $c = -63$ mV, $d = 60$ pA, $v_r = -75$, $v_t = -58$ mV, $v_{peak} = 29$ mV. Parameters for inhibitory cells were: $C_m = 16$ pF, $k = 1.5$ nS/mV, $a = 900$ Hz, $b = 2$ nS, $c = -80$ mV, $d = 400$ pA, $v_r = -65$ mV, $v_t = -50$ mV, $v_{peak} = 28$ mV. Synaptic reversal potentials were set as $v_E = 10$ mV, and $v_I = -80$ mV. When $v_i \geq v_{peak}$, v_i and u_i were reset to

$$v_i \leftarrow c \quad \text{(Equation 35)}$$

$$u_i \leftarrow u_i + d \quad \text{(Equation 36)}$$

Excitatory and inhibitory cells were connected through four types of synapses: excitatory to excitatory cell (EE) synapses, excitatory to inhibitory cell synapses (EI), inhibitory to excitatory cell synapses (IE), and inhibitory to inhibitory cell (II) synapses. Kinetics were modelled as exponential synapses such that the synaptic conductance g_{syn} evolved according to:

$$\frac{dg_{syn}}{dt} = -g_{syn}/\tau_g + \widetilde{g_{syn}} w \sum_{t_s} \delta(t - t_s) \quad \text{(Equation 37)}$$

where $\widetilde{g_{syn}}$ is the maximum synaptic conductance, and $\tau_g = 10$ ms for EE and EI synapses and 20 ms for IE and II synapses. Maximum excitatory synapse condutance was 0.5 nS, and 1.0 nS for inhibitory synapses. EE and IE synapses were subject to spike timing-dependent plasticity of the form

$$\Delta w = \eta[\exp(-|t_{pre} - t_{post}|) - z] \quad \text{(Equation 38)}$$

where η is a learning rate, and z is a scaling factor. This provides a symmetric STDP rule used previously as a homeostatic means to balance excitation and inhibition through inhibitory plasticity, (Vogels, Sprekeler, Zenke, Clopath, & Gerstner, 2011) and has also recently been shown to operate at CA3 associated commissural synapses (Mishra, Kim,

Guzman, & Jonas, 2016). In the case of IE synapses, η and z are fixed, however in the case of EE synapses these evolve according to

$$\frac{d\eta}{dt} = -\eta/\tau_\eta + \xi \sum_{t_{post}} \delta(t - t_{post}) \qquad \text{(Equation 39)}$$

$$\frac{dz}{dt} = -z/\tau_z + \rho_{max}^{-1} \sum_{t_{post}} \delta(t - t_{post}) \qquad \text{(Equation 40)}$$

This makes η and z track the postsynaptic firing rate for two different purposes, and on different time scales since $\tau_\eta = 100$ ms, and $\tau_z = 1$ second; η becomes a burst detector that increases the learning rate by a factor ξ meaning STDP requires multiple post-synaptic spikes to be activated, and z scales STDP such that the postsynaptic firing rate reaches a maximum ρ_{max}, more pre- and post-synaptic spike pairs cause depression, preventing STDP from inducing unrealistically high firing rates in an excitatory recurrent network. Additionally, STDP was bounded between 0 and the maximum conductance of the synapse.

Synaptic input to cells was comprised of recurrent and feed-forward inputs, i.e.,

$$g_E^i = \sum_{ij} g_{EY}^{ij} + \sum_{ik} g_{FF}^{ik} \qquad \text{(Equation 41)}$$

$$g_I^i = \sum_{ij} g_{IY}^{ij} \qquad \text{(Equation 42)}$$

where g_{FF} is the conductance of feedforward input, g_{EY} is excitatory recurrent input, and g_{IY} is recurrent inhibition. Feedforward input was given only to excitatory cells. CA3 Network architecture and feed-forward dynamics varied according to each simulation.

Granule cell spiking was modelled as an inhomegenous Poisson process. For most simulations, it was modelled low spike rate of 0.2 Hz, punctuated by jumps to a much higher burst frequency every 20 seconds for 250 ms. In one set of simulations, a small granule cell population (N=50) was modelled as a set of Poisson processes rates defined by winner-take-all dynamics with adaptation (Fukai and Tanaka, 1997)

$$\frac{dy_i}{dt} = y(\sum_j(y_j - 50\alpha_j) + \kappa_i)/10 \qquad \text{(Equation 43)}$$

$$\frac{d\alpha_i}{dt} = -(\alpha_i - y_i)/5000 \qquad \text{(Equation 44)}$$

$$\kappa_i = h + (i/N)\sigma \qquad \text{(Equation 45)}$$

where y are granule cell rates following winner-take-all dynamics, α is rate-adaptation taking place over a longer time scale, and κ is the input to each granule cell determined by h and σ, which define the scale and separation of rates respectively, where the scale parameter is a scaling factor that multiples each rate in the population by the same amount, whereas the separation parameter controls the difference in input to the population of rates, determining how much the winner wins by. The winner granule cell receives input $\kappa = h + \sigma$.

Figure 1: **The effects of carbachol on feed-forward excitatory and inhibitory transmission in the mossy fiber pathway.** A) Left: Experimental setup indicating the location of stimulation and recording electrodes within a hippocampal slice. Right: Schema of feed-forward mossy fiber circuit. B) EPSCs and IPSCs evoked by granule cells stimulation were blocked by 1 μM DCG-IV, confirming responses were driven by mossy fiber activation. Top: Example traces of DCG-IV block (black) of EPSCs (red) and IPSCs (blue). Bottom: Time course of DCG-IV block of 4th EPSC (red, n = 9) and IPSC (blue, n = 6). C) 5 μM CCh mildly suppresses mossy fiber EPSCs. Top left: Example traces before and after bath application of 5 μM CCh. Bottom Left: Time course of CCh effect, and washout (n = 7). Top Right: Effect of CCh on response amplitudes for each pulse. Bottom Right: Effect of CCh on nth/1st Pulse Ratio. D) 5 μM CCh substantially reduces disynaptic mossy fiber driven IPSC amplitudes. Top left: Example traces before and after bath application of 5 μM CCh. Bottom Left: Time course of CCh effect, and washout (n = 5). Top Right: Effect of CCh on response amplitudes for each pulse. Bottom Right: Effect of CCh on nth/1st Pulse Ratio.

Figure 2: The mechanism of carbachol action on excitatory and inhibitory feed-forward mossy fiber transmission determined by short term plasticity models. A) Irregular stimulation protocol modelled on naturalistic granule cell spike patterns. GC spike patterns recorded *in vivo* during a spatial memory task, with a bimodal inter-spike interval (ISI) distribution (top right). Bimodal ISI distribution modelled as a doubly

stochastic cox process (middle right), with irregular stimulation protocol a sample drawn from this process (bottom right). B-C) EPSCs and IPSCs evoked by irregular stimulation protocol. Evoked peaks highlighted by white dots. An example burst is shown on expanded timescale. D) Tsodyks-Markram short term plasticity model schematic. E-F) Model selection and fitting for EPSCs and IPSCs. Top: AIC and BIC weights for each fitted model. Model selection by highest AIC and BIC weights and evidence ratios. Bottom: Modulation of CCh assessed by effect on parameter fits normalized by time-matched control.

Figure 3: **Carbachol alters the Excitatory-Inhibitory ratio within the feed-forward mossy fiber pathway in a frequency-dependent manner.** A) Simplification of bursting spike trains into two parameter spaces: between burst interval (BI) describing interval between bursts in a spike train, and within BI describing interval between spikes within a burst. B) Example synaptic waveforms of expected mossy fiber EPSCs and IPSCs given short term plasticity dynamics at three pairs of within- and between- burst intervals. C&D) Expected short term plasticity of EPSCs (C) and IPSCs (D) across a wide range of within and between BIs. E) Progression of Excitatory-Inhibitory ratio across the range of within and between BIs.

Figure 4: **Acetylcholine-mediated disinhibition facilitates back-propagation of EPSPs and action potentials into the dendrites of CA3 pyramidal cells.** A) Sketch of CA3 pyramidal cell and positioning within the layers of the hippocampus. Synaptic inputs are shown with location of contact (red - recurrent CA3-CA3 synapse, blue - feed-forward inhibitory synapse, gray - mossy fiber synapse). SLM – stratum lacunosum moleculare, SR – stratum radiatum, SL – stratum lucidum, SP – stratum pyramidale, SO – stratum oriens. B) Example traces of action potential generation at the soma from summated mossy fiber EPSPs presented at 20 Hz (top), backpropagation into the radial oblique dendrites (middle), and dendritic calcium influx (bottom), with and without acetylcholine-mediated disinhibition of feed-forward inhibition. C) Backpropagating potential amplitude and % increase with acetylcholine shown for all modelled dendritic compartments.

Figure 5: **Facilitating mossy fiber inputs generate rapid and stable ensemble formation.** A) Network properties and long-term plasticity rules. Left: A population of excitatory (red) and inhibitory (blue) cells with all-to-all connectivity and mossy fiber input (gray). Right: Recurrent excitatory CA3-CA3 spike timing-dependent plasticity rule with a symmetric window that shifted from potentiation for correlated spiking at low rates (Ppost = 0), to depression for uncorrelated spiking near the maximum postsynaptic firing rate (Ppost = Pmax). B) Comparison of mossy fiber to irregular 'perforant path' input. Weight evolution with time for two cell excitatory population (top), 10 cell excitatory population (middle) and population including 10 excitatory and 5 inhibitory cells (bottom). C) Top Left: Network setup- fifty feedforward inputs to a single granule cell. Top Right- example of winner-take-all rates and postsynaptic spiking. This instance shows strong efficacy since spiking occurs once most rates have decreased to near zero. Bottom: Efficacy of winner granule cell to CA3 pyramidal cell coupling with facilitating and constant synaptic input

Figure 6: **Acetylcholine speeds up ensemble formation and lowers input frequency requirement by increasing cellular excitability.** A) Network setup: Population of excitatory and inhibitory cells connected in all to all fashion. Subpopulations of excitatory cells receive independent feed-forward input that drives ensemble formation. B) Example weight matrix driven by input with 30 Hz, 20 Hz or 20 Hz bursts with CA3 acetylcholine effects. Stronger weights indicate robust ensemble formation. C) Evolution of ensemble formation illustrated by the weight matrix error reduction over time for different input burst frequencies. Triangles denote which effects of acetylcholine on CA3 network were included in each set of simulations.

Figure 7: Acetylcholine enables CA3 network to form stable overlapping ensembles by reducing the strength of recurrent excitatory CA3-CA3 synapses. A) Network setup: Population of excitatory and inhibitory cells connected in all to all fashion. Subpopulations of excitatory cells receive overlapping feed-forward input that drives ensemble formation. B) Example weight matrix driven by input with increasing degree of overlap [0, 2, 4]. C) Evolution of ensemble formation illustrated by the weight matrix error over time for different degrees of overlap. Triangles denote which effects of acetylcholine were included in the simulation. D) Effect of increasing overlap on the ability to discriminate between ensembles defined as the difference in ensemble population spiking rates.

Supplementary Figure 1: A-B) Latency, rise times, and jitter of mossy fiber driven EPSCs (n = 11) (A) and IPSCs (n = 12) (B) C-D) Reversal potential estimation of glutamatergic (n = 5) (C) and GABAergic (n = 6) (D) transmission at CA3 pyramidal cells. E-F) Spontaneous EPSC (n = 6) (E) and IPSC (n = 5) (F) frequency recorded before and after carbachol application G-H) CA3 pyramidal cell input resistance (n = 5) (G) and holding current at -70 mV (n = 5) (H) before and after carbachol application. I) Dose-response of carbachol effect on IPSC amplitudes (n = 3).

Supplementary Figure 2: A-B: Goodness-of-fit for EPSC (A) and IPSC (B) short term plasticity models assessed by Bayesian posterior predictive p-values. Plots show the observed data all experiments (EPSCs and IPSCs respectively) together with the expected values from the model before and after the application of 5 µM carbachol. P-values close to 0.5 indicate best fit. C-D: Posterior distributions for parameters of best fitting models given data for EPSCs (C) and IPSCs (D).

Supplementary Figure 3: A) Slices of data shown in Figure 6C along frequency (top) and time (bottom) axes. B) Slices of data shown in Figure 7C along overlap (top) and time (bottom) axes. Colour coding for plots is indicated in the legend representing inclusion of different effects of acetylcholine in CA3.

Funding

LYP, CC and JRM funded by the Wellcome Trust. The project was also supported by an IBRO & Simons fund through Grant ID # isiCNI2017. KTA gratefully acknowledges the financial support of the EPSRC via grant EP/N014391/1.

Acknowledgements

Contributions

References

Acsady, L., Kamondi, A., Sik, A., Freund, T., and Buzsaki, G. (1998). GABAergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. J Neurosci 18, 3386-3403.

Brandalise, F., and Gerber, U. (2014). Mossy fiber-evoked subthreshold responses induce timing-dependent plasticity at hippocampal CA3 recurrent synapses. Proc Natl Acad Sci U S A 111, 4303-4308.

Buchanan, K.A., Petrovic, M.M., Chamberlain, S.E., Marrion, N.V., and Mellor, J.R. (2010). Facilitation of long-term potentiation by muscarinic M(1) receptors is mediated by inhibition of SK channels. Neuron 68, 948-963.

Cea-del Rio, C.A., Lawrence, J.J., Tricoire, L., Erdelyi, F., Szabo, G., and McBain, C.J. (2010). M3 muscarinic acetylcholine receptor expression confers differential cholinergic modulation to neurochemically distinct hippocampal basket cell subtypes. J Neurosci 30, 6011-6024.

Cheng, Q., and Yakel, J.L. (2014). Presynaptic alpha7 nicotinic acetylcholine receptors enhance hippocampal mossy fiber glutamatergic transmission via PKA activation. J Neurosci 34, 124-133.

Costa, R.P., Sjostrom, P.J., and van Rossum, M.C. (2013). Probabilistic inference of short-term synaptic plasticity in neocortical microcircuits. Frontiers in computational neuroscience 7, 75.

Dasari, S., and Gulledge, A.T. (2011). M1 and M4 receptors modulate hippocampal pyramidal neurons. J Neurophysiol 105, 779-792.

Dennis, S.H., Pasqui, F., Colvin, E.M., Sanger, H., Mogg, A.J., Felder, C.C., Broad, L.M., Fitzjohn, S.M., Isaac, J.T., and Mellor, J.R. (2016). Activation of Muscarinic M1 Acetylcholine Receptors Induces Long-Term Potentiation in the Hippocampus. Cereb Cortex 26, 414-426.

Dickinson, B.A., Jo, J., Seok, H., Son, G.H., Whitcomb, D.J., Davies, C.H., Sheng, M., Collingridge, G.L., and Cho, K. (2009). A novel mechanism of hippocampal LTD involving muscarinic receptor-triggered interactions between AMPARs, GRIP and liprin-alpha. Molecular brain 2, 18.

Fernandez de Sevilla, D., and Buno, W. (2010). The muscarinic long-term enhancement of NMDA and AMPA receptor-mediated transmission at Schaffer collateral synapses develop through different intracellular mechanisms. J Neurosci 30, 11032-11042.

Fukai, T., and Tanaka, S. (1997). A simple neural network exhibiting selective activation of neuronal ensembles: from winner-take-all to winners-share-all. Neural Comput 9, 77-97.

Gold, A.E., and Kesner, R.P. (2005). The role of the CA3 subregion of the dorsal hippocampus in spatial pattern completion in the rat. Hippocampus 15, 808-814.

Gu, Z., and Yakel, J.L. (2011). Timing-dependent septal cholinergic induction of dynamic hippocampal synaptic plasticity. Neuron 71, 155-165.

Guzman, S.J., Schlogl, A., Frotscher, M., and Jonas, P. (2016). Synaptic mechanisms of pattern completion in the hippocampal CA3 network. Science 353, 1117-1123.

Hangya, B., Ranade, S.P., Lorenc, M., and Kepecs, A. (2015). Central Cholinergic Neurons Are Rapidly Recruited by Reinforcement Feedback. Cell 162, 1155-1168.

Hasselmo, M.E., and Sarter, M. (2011). Modes and models of forebrain cholinergic neuromodulation of cognition. Neuropsychopharmacology 36, 52-73.

Hasselmo, M.E., Schnell, E., and Barkai, E. (1995). Dynamics of learning and recall at excitatory recurrent synapses and cholinergic modulation in rat hippocampal region CA3. J Neurosci 15, 5249-5262.

Henze, D.A., Wittner, L., and Buzsaki, G. (2002). Single granule cells reliably discharge targets in the hippocampal CA3 network in vivo. Nat Neurosci 5, 790-795.

Hummos, A., Franklin, C.C., and Nair, S.S. (2014). Intrinsic mechanisms stabilize encoding and retrieval circuits differentially in a hippocampal network model. Hippocampus 24, 1430-1448.

Kaifosh, P., and Losonczy, A. (2016). Mnemonic Functions for Nonlinear Dendritic Integration in Hippocampal Pyramidal Circuits. Neuron 90, 622-634.

Kamiya, H., Shinozaki, H., and Yamamoto, C. (1996). Activation of metabotropic glutamate receptor type 2/3 suppresses transmission at rat hippocampal mossy fiber synapses. J Physiol (Lond) 493, 447-455.

Kesner, R.P., and Rolls, E.T. (2015). A computational theory of hippocampal function, and tests of the theory: new developments. Neurosci Biobehav Rev 48, 92-147.

Kobayashi, K., and Poo, M.M. (2004). Spike train timing-dependent associative modification of hippocampal CA3 recurrent synapses by mossy fibers. Neuron 41, 445-454.

Marino, M.J., Rouse, S.T., Levey, A.I., Potter, L.T., and Conn, P.J. (1998). Activation of the genetically defined m1 muscarinic receptor potentiates N-methyl-D-aspartate (NMDA) receptor currents in hippocampal pyramidal cells. P Natl Acad Sci USA 95, 11465-11470.

Markram, H., and Segal, M. (1992). The inositol 1,4,5-trisphosphate pathway mediates cholinergic potentiation of rat hippocampal neuronal responses to NMDA. J Physiol 447, 513-533.

Marr, D. (1971). Simple memory: a theory for archicortex. Philos Trans R Soc Lond B Biol Sci 262, 23-81.

McClelland, J.L., and Goddard, N.H. (1996). Considerations arising from a complementary learning systems perspective on hippocampus and neocortex. Hippocampus 6, 654-665.

McGaughy, J., Everitt, B.J., Robbins, T.W., and Sarter, M. (2000). The role of cortical cholinergic afferent projections in cognition: impact of new selective immunotoxins. Behav Brain Res 115, 251-263.

Miles, R., Toth, K., Gulyas, A.I., Hajos, N., and Freund, T.F. (1996). Differences between somatic and dendritic inhibition in the hippocampus. Neuron 16, 815-823.

Mishra, R.K., Kim, S., Guzman, S.J., and Jonas, P. (2016). Symmetric spike timing-dependent plasticity at CA3-CA3 synapses optimizes storage and recall in autoassociative networks. Nature communications 7, 11552.

Mistry, R., Dennis, S., Frerking, M., and Mellor, J.R. (2011). Dentate gyrus granule cell firing patterns can induce mossy fiber long-term potentiation in vitro. Hippocampus 21, 1157-1168.

Mori, M., Abegg, M.H., Gahwiler, B.H., and Gerber, U. (2004). A frequency-dependent switch from inhibition to excitation in a hippocampal unitary circuit. Nature 431, 453-456.

Mori, M., Gahwiler, B.H., and Gerber, U. (2007). Recruitment of an inhibitory hippocampal network after bursting in a single granule cell. Proc Natl Acad Sci U S A 104, 7640-7645.

Mullner, F.E., Wierenga, C.J., and Bonhoeffer, T. (2015). Precision of Inhibition: Dendritic Inhibition by Individual GABAergic Synapses on Hippocampal Pyramidal Cells Is Confined in Space and Time. Neuron 87, 576-589.

Nakazawa, K., Quirk, M.C., Chitwood, R.A., Watanabe, M., Yeckel, M.F., Sun, L.D., Kato, A., Carr, C.A., Johnston, D., Wilson, M.A., and Tonegawa, S. (2002). Requirement for hippocampal CA3 NMDA receptors in associative memory recall. Science 297, 211-218.

Nicoll, R.A., and Schmitz, D. (2005). Synaptic plasticity at hippocampal mossy fiber synapses. Nat Rev Neurosci 6, 863-876.

O'Donnell, C., and Sejnowski, T.J. (2014). Selective memory generalization by spatial patterning of protein synthesis. Neuron 82, 398-412.

O'Reilly, R.C., and McClelland, J.L. (1994). Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off. Hippocampus 4, 661-682.

Papouin, T., Dunphy, J.M., Tolman, M., Dineley, K.T., and Haydon, P.G. (2017). Septal Cholinergic Neuromodulation Tunes the Astrocyte-Dependent Gating of Hippocampal NMDA Receptors to Wakefulness. Neuron 94, 840-854 e847.

Parikh, V., Kozak, R., Martinez, V., and Sarter, M. (2007). Prefrontal acetylcholine release controls cue detection on multiple timescales. Neuron 56, 141-154.

Pelkey, K.A., Lavezzari, G., Racca, C., Roche, K.W., and McBain, C.J. (2005). mGluR7 is a metaplastic switch controlling bidirectional plasticity of feed-forward inhibition. Neuron 46, 89-102.

Pouille, F., and Scanziani, M. (2001). Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. Science 293, 1159-1163.

Prince, L.Y., Bacon, T.J., Tigaret, C.M., and Mellor, J.R. (2016). Neuromodulation of the Feed-forward Dentate Gyrus-CA3 Microcircuit. Front Synaptic Neurosci 8, 32.

Restivo, L., Niibori, Y., Mercaldo, V., Josselyn, S.A., and Frankland, P.W. (2015). Development of Adult-Generated Cell Connectivity with Excitatory and Inhibitory Cell Populations in the Hippocampus. J Neurosci 35, 10600-10612.

Ruediger, S., Vittori, C., Bednarek, E., Genoud, C., Strata, P., Sacchetti, B., and Caroni, P. (2011). Learning-related feed-forward inhibitory connectivity growth required for memory precision. Nature 473, 514-518.

Sachidhanandam, S., Blanchet, C., Jeantet, Y., Cho, Y.H., and Mulle, C. (2009). Kainate receptors act as conditional amplifiers of spike transmission at hippocampal mossy fiber synapses. J Neurosci 29, 5000-5008.

Salin, P.A., Scanziani, M., Malenka, R.C., and Nicoll, R.A. (1996). Distinct short-term plasticity at two excitatory synapses in the hippocampus. P Natl Acad Sci USA 93, 13304-13309.

Scanziani, M. (2000). GABA spillover activates postsynaptic GABA(B) receptors to control rhythmic hippocampal activity. Neuron 25, 673-681.

Szabadics, J., and Soltesz, I. (2009). Functional specificity of mossy fiber innervation of GABAergic cells in the hippocampus. J Neurosci 29, 4239-4251.

Szabo, G.G., Holderith, N., Gulyas, A.I., Freund, T.F., and Hajos, N. (2010). Distinct synaptic properties of perisomatic inhibitory cell types and their different modulation by cholinergic receptor activation in the CA3 region of the mouse hippocampus. Eur J Neurosci 31, 2234-2246.

Teles-Grilo Ruivo, L.M., Baker, K.L., Conway, M.W., Kinsley, P.J., Gilmour, G., Phillips, K.G., Isaac, J.T., Lowry, J.P., and Mellor, J.R. (2017). Coordinated Acetylcholine Release in Prefrontal Cortex and Hippocampus Is Associated with Arousal and Reward on Distinct Timescales. Cell reports 18, 905-917.

Torborg, C.L., Nakashiba, T., Tonegawa, S., and McBain, C.J. (2010). Control of CA3 output by feed-forward inhibition despite developmental changes in the excitation-inhibition balance. J Neurosci 30, 15628-15637.

Toth, K., Suares, G., Lawrence, J.J., Philips-Tansey, E., and McBain, C.J. (2000). Differential mechanisms of transmission at three types of mossy fiber synapse. J Neurosci 20, 8279-8289.

Treves, A., and Rolls, E.T. (1994). Computational analysis of the role of the hippocampus in memory. Hippocampus 4, 374-391.

Tsubokawa, H., and Ross, W.N. (1996). IPSPs modulate spike backpropagation and associated [Ca2+]i changes in the dendrites of hippocampal CA1 pyramidal neurons. J Neurophysiol 76, 2896-2906.

Vogt, K.E., and Regehr, W.G. (2001). Cholinergic modulation of excitatory synaptic transmission in the CA3 area of the hippocampus. J Neurosci 21, 75-83.

Vyleta, N.P., Borges-Merjane, C., and Jonas, P. (2016). Plasticity-dependent, full detonation at hippocampal mossy fiber-CA3 pyramidal neuron synapses. Elife 5.

Wagenmakers, E.J., and Farrell, S. (2004). AIC model selection using Akaike weights. Psychon Bull Rev 11, 192-196.

Williams, S., and Johnston, D. (1990). Muscarinic depression of synaptic transmission at the hippocampal mossy fiber synapse. J Neurophysiol 64, 1089-1097.

Wilmes, K.A., Sprekeler, H., and Schreiber, S. (2016). Inhibition as a Binary Switch for Excitatory Plasticity in Pyramidal Neurons. PLoS computational biology 12, e1004768.

Yu, A.J., and Dayan, P. (2005). Uncertainty, neuromodulation, and attention. Neuron 46, 681-692.

Zucca, S., Griguoli, M., Malezieux, M., Grosjean, N., Carta, M., and Mulle, C. (2017). Control of Spike Transfer at Hippocampal Mossy Fiber Synapses In Vivo by GABAA and GABAB Receptor-Mediated Inhibition. J Neurosci 37, 587-598.

Additional References

Chrobak, J. J., & Buzsáki, G. (1998). Gamma oscillations in the entorhinal cortex of the freely behaving rat. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *18*(1), 388–98. http://doi.org/0270-6474/97/180388-11$05.00/0

Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences of the United States of America*, *79*(8), 2554–2558. http://doi.org/10.1073/pnas.79.8.2554

Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, *90*(430), 773–795.

Major, G., Larkman, A. U., Jonas, P., Sakmann, B., & Jack, J. J. (1994). Detailed passive cable models of whole-cell recorded CA3 pyramidal neurons in rat hippocampal slices. *Journal of Neuroscience*, *14*(8), 4613–4638.