Molecular evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome

Kristen A. Panfilio[1, 2]*, Iris M. Vargas Jentzsch[1], Joshua B. Benoit[3], Deniz Erezyilmaz[4], Yuichiro Suzuki[5], Stefano Colella[6, 7], Hugh M. Robertson[8], Monica F. Poelchau[9], Robert M. Waterhouse[10, 11], Panagiotis Ioannidis[10], Matthew T. Weirauch[12], Daniel S.T. Hughes[13], Shwetha C. Murali[13, 14, 15], John H. Werren[16], Chris G.C. Jacobs[17, 18], Elizabeth J. Duncan[19, 20], David Armisén[21], Barbara M.I. Vreede[22], Patrice Baa-Puyoulet[6], Chloé S. Berger[21], Chun-che Chang[23], Hsu Chao[13], Mei-Ju M. Chen[9], Yen-Ta Chen[1], Christopher P. Childers[9], Ariel D. Chipman[22], Andrew G. Cridge[19], Antonin J.J. Crumière[21], Peter K. Dearden[19], Elise M. Didion[3], Huyen Dinh[13], HarshaVardhan Doddapaneni[13], Amanda Dolan[16, 24], Shannon Dugan[13], Cassandra G. Extavour[25, 26], Gérard Febvay[6], Markus Friedrich[27], Neta Ginzburg[22], Yi Han[13], Peter Heger[28], Christopher J. Holmes[3], Thorsten Horn[1], Yi-min Hsiao[23], Emily C. Jennings[3], J. Spencer Johnston[29], Tamsin E. Jones[25], Jeffery W. Jones[27], Abderrahman Khila[21], Stefan Koelzer[1], Viera Kovacova[30], Megan Leask[19], Sandra L. Lee[13], Chien-Yueh Lee[9], Mackenzie R. Lovegrove[19], Hsiao-ling Lu[23], Yong Lu[31], Patricia J. Moore[32], Monica C. Munoz-Torres[33], Donna M. Muzny[13], Subba R. Palli[34], Nicolas Parisot[6], Leslie Pick[31], Megan Porter[35], Jiaxin Qu[13], Peter N. Refki[21, 36], Rose Richter[16, 37], Rolando Rivera Pomar[38], Andrew J. Rosendale[3], Siegfried Roth[1], Lena Sachs[1], M. Emília Santos[21], Jan Seibert[1], Essia Sghaier[21], Jayendra N. Shukla[34, 39], Richard J. Stancliffe[40], Olivia Tidswell[19, 41], Lucila Traverso[42], Maurijn van der Zee[17], Séverine Viala[21], Kim C. Worley[13], Evgeny M. Zdobnov[10], Richard A. Gibbs[13], Stephen Richards[13]

* Correspondence: kristen.panfilio@alum.swarthmore.edu

The full list of author information is available at the end of the manuscript.

34 **ABSTRACT**
35
Background:
The Hemiptera (aphids, cicadas, and true bugs) are a key insect order, with high
diversity for feeding ecology and excellent experimental tractability for molecular
genetics. Building upon recent sequencing of hemipteran pests such as phloem-
feeding aphids and blood-feeding bed bugs, we present the genome sequence and
comparative analyses centered on the milkweed bug *Oncopeltus fasciatus*, a seed
feeder of the family Lygaeidae.
Results:
The 926-Mb *Oncopeltus* genome is well represented by the current assembly and
official gene set. We use our genomic and RNA-seq data not only to characterize the
protein-coding gene repertoire and perform isoform-specific RNAi, but also to
elucidate patterns of molecular evolution and physiology. We find ongoing, lineage-
specific expansion and diversification of repressive C2H2 zinc finger proteins. The
discovery of intron gain and turnover specific to the Hemiptera also prompted
evaluation of lineage and genome size as predictors of gene structure evolution.
Furthermore, we identify enzymatic gains and losses that correlate with feeding
biology, particularly for reductions associated with derived, fluid-nutrition feeding.
Conclusions:
With the milkweed bug, we now have a critical mass of sequenced species for a
hemimetabolous insect order and close outgroup to the Holometabola, substantially
improving the diversity of insect genomics. We thereby define commonalities among
the Hemiptera and delve into how hemipteran genomes reflect distinct feeding
ecologies. Given *Oncopeltus'*s strength as an experimental model, these new
sequence resources bolster the foundation for molecular research and highlight
technical considerations for the analysis of medium-sized invertebrate genomes.
61
62
63 **Keywords:**
64 Phytophagy; Transcription Factors; Gene Structure; Lateral Gene Transfer; RNAi;
65 Gene family evolution; Evolution of Development
66
67

BACKGROUND

The number of animals with sequenced genomes continues to increase dramatically, and there are now over 100 insect species with assembled and annotated genomes [1]. However, the majority belong to the Holometabola (*e.g.*, flies, beetles, wasps, butterflies), the group characterized by a biphasic life history with distinct larval and adult phases separated by dramatic metamorphosis during a pupal stage. The Holometabola represent only a fraction of the full morphological and ecological diversity across the Insecta: over half of all orders are hemimetabolous. Imbalance in genomic resources limits the exploration of this diversity, including the environmental and developmental requirements of a hemimetabolous life style with a progression of flightless nymphal (juvenile) instars. Addressing this paucity, we report comparative analyses based on genome sequencing of the large milkweed bug, *Oncopeltus fasciatus*, as a hemimetabolous representative of the larger diversity of insects.

Oncopeltus is a member of the Hemiptera, the most species-rich hemimetabolous order. Together with the Thysanoptera and, traditionally, the Psocodea, the Hemiptera form the hemipteroid assemblage (or Acercaria), a close outgroup to the Holometabola [2, 3]. All Hemiptera share the same piercing and sucking mouthpart anatomy [4], yet they have diversified to exploit food sources ranging from seeds and plant tissues (phytophagy) to phloem sap (mucivory) and vertebrate blood (hematophagy). For this reason, many hemipterans are agricultural pests or human disease vectors, and genome sequencing efforts to date have focused on these species (Fig. 1, [5]), including phloem-feeding aphids [6-8], psyllids [9], and planthoppers [10], and the hematophagous kissing bug, *Rhodnius prolixus* [11], a vector of Chagas disease, and bed bug, *Cimex lectularius* [12, 13]. Building on transcriptomic data, genome projects are also in progress for other pest species within the same infraorder as *Oncopeltus*, such as the stink bug *Halyomorpha halys* [14, 15].

The milkweed bug has feeding ecology traits that are both conservative and complementary to those of previously sequenced hemipterans. Its phytophagy is ancestral for the large infraorder Pentatomomorpha and representative of most extant Hemiptera [16]. Moreover, as a seed feeder *Oncopeltus* has not undergone the marked life style changes associated with fluid feeding (mucivory or hematophagy), including dependence on endosymbiotic bacteria to provide nutrients lacking in the diet. Gene loss in the pea aphid, *Acyrthosiphon pisum*, makes it reliant on the obligate endosymbiont *Buchnera aphidicola* for synthesis of essential amino acids [6, 17]. Although hematophagy arose independently in *Rhodnius* and *Cimex* [16], their respective endosymbionts, *Rhodococcus rhodnii* and *Wolbachia*, must provide vitamins lacking in a blood diet [18]. In contrast, the seed-feeding subfamily Lygaeinae, including *Oncopeltus*, is notable for the absence of prominent endosymbiotic anatomy: these bugs lack both the midgut crypts that typically house bacteria and the bacteriomes and endosymbiotic balls seen in other Lygaeidae [19].

As the native food source of *Oncopeltus* is the toxic milkweed plant, its own feeding biology has a number of interesting implications regarding detoxification and sequestration of cardenolide compounds. A prominent consequence of this diet is the bright red-orange aposematic (warning) coloration seen in *Oncopeltus* embryos, nymphs, and adults [20, 21]. Thus, diet, metabolism, and body pigmentation are functionally linked biological features for which one may expect changes in gene

118 repertoires to reflect diversity within an order, and the Hemiptera provide an excellent
119 opportunity to explore this.
120
Furthermore, *Oncopeltus* has been an established laboratory model organism
for over 60 years, with a rich experimental tradition in a wide range of studies from
physiology and development to evolutionary ecology [21-23]. It is among the few
experimentally tractable hemimetabolous insect species, and it is amenable to a range
of molecular techniques (*e.g.*, [24-26]). In fact, it was one of the first insect species to
be functionally investigated by RNA interference (RNAi, [27]). RNAi in *Oncopeltus*
is highly effective across different life history stages, which has led to a resurgence of
experimental work over the past fifteen years, with a particular focus on the evolution
of developmentally important regulatory genes (reviewed in [23]).
130
Here, we focus on these two themes – feeding biology diversity within the
Hemiptera and *Oncopeltus* as a research model for macroevolutionary genetics. Key
insights derive from a combination of global comparative genomics and detailed
computational analyses that are supported by extensive manual curation, empirical
data for gene expression, sequence validation, and new isoform-specific RNAi. We
thereby identify genes with potentially restricted life history expression in *Oncopeltus*
and that are unique to the Hemiptera, clarify evolutionary patterns of zinc finger
protein expansion, categorize predictors of insect gene structure, and identify lateral
gene transfer and amino acid metabolism features that correlate with feeding biology.
140
141
142 **RESULTS AND DISCUSSION**
143
The genome and its assembly
Oncopeltus fasciatus has a diploid chromosome number (2*n*) of 16, comprised of
seven autosomal pairs and two sex chromosomes with the XX/XY sex determination
system [28, 29]. To analyze this genetic resource, we sequenced and assembled the
genome using next-generation sequencing approaches (Table 1, see also Methods and
Supplemental Notes Sections 1-4). We measure the genome size to be 923 Mb in
females and 928 Mb in males based on flow cytometry data (Supplemental Note
2.1.a). The assembly thus contains 84% of the expected sequence, which is
comparable to other recent, medium-sized insect genomes [12, 30]. However, our
analyses of the *k*-mer frequency distribution in raw sequencing reads yielded
ambiguous estimates of genome size and heterozygosity rate, which is suggestive of
high heterozygosity and repetitive content ([31], Supplemental Note 2.1.b). In further
analyses we indeed obtained high estimates of repetitive content, although
heterozygosity does not unduly influence gene prediction (see below, based on
protein orthology assessments). These computationally challenging features may be
increasingly relevant as comparative genomics extends to insect species with larger
genomes (>1 Gb) – a common feature among hemimetabolous insects [5, 32].
161
162 As template DNA was prepared from dissected adults from which gut material
163 was removed, the resulting assembly is essentially free of contamination. Only five
164 small scaffolds had high bacterial homology, each to a different, partial bacterial
165 genome (Supplemental Note 2.2).
166
167

Table 1. *Oncopeltus fasciatus* genome metrics.

Feature	Value	
2*n* chromosomes	16	
Genome size	926 Mb (mean between males and females)	
Assembly size	1,099 Mb (contigs only: 774 Mb)	
Coverage	106.9× raw coverage, 83.7% of reads in final assembly	
Contig N50	4,047 bp	
Scaffold N50	340.0 kb	
# Scaffolds	17,222	
GC content	genome: 32.7%, protein-coding sequence (OGS v1.2): 42%	
OGS v1.1 (curated fraction)	19,690 models[1] (1,426 models, 7.2%)	19,465 genes (1,201 genes, 6.2%)
OGS v1.2 (curated fraction)	19,809 models[1] (1,697 models, 8.7%)	19,616 genes (1,518 genes, 7.7%)

[1] Individual genes may be represented by multiple models in cases of curated alternative isoforms or if exons of the gene are split across scaffolds.

The official gene set and conserved gene linkage

The official gene set (OGS) was generated by automatic annotation followed by manual curation in a large-scale effort by the research community (Supplemental Notes Sections 3-4). Curation revised automatic models, added alternative isoforms and *de novo* models, and documented multiple models for genes split across scaffolds. We found that automatic predictions were rather conservative for hemipteran gene structure (see below). Thus, manual curation often extended gene loci as exons were added, including merging discrete automatic models (Supplemental Note 4, Table S4.4). The OGS v1.1 was generated for global analyses to characterize the gene repertoire. The latest version, OGS v1.2, primarily adds chemoreceptor genes of the ionotropic and odorant receptor classes and genes encoding metabolic enzymes. Altogether, the research community curated 1,697 gene models (8.7% of OGS v1.2), including 316 *de novo* models (Table S4.1, Supplemental Notes Section 5). The majority of curated models are for genes encoding cuticular proteins (11%), chemoreceptors (19%), and developmental regulators such as transcription factors and signaling pathway components (40%, including the BMP/TGF-β, Toll/NF-κB, Notch, Hedgehog, Torso RTK, and Wnt pathways).

In addition to assessing gene model quality, manual curation of genes whose orthologs are expected to occur in syntenic clusters also validates assembly scaffolding. Complete loci could be found for single orthologs of all Hox cluster genes, where *Hox3/zen* and *Hox4/Dfd* are linked in the current assembly and have ≥99.9% nucleotide identity with experimentally validated sequences ([33-35], Supplemental Note 5.1.b). Conserved linkage was also confirmed for the homeobox genes of the Iroquois complex, the Wnt ligands *wingless* and *wnt10*, and two linked pairs from the Runt transcription factor complex (Supplemental Notes 5.1.a, 5.1.c, 5.1.i, 5.1.j). Further evidence for correct scaffold assembly comes from the curation

205 of large, multi-exonic loci. For example, the cell polarity and cytoskeletal regulator
206 encoded by the conserved *furry* gene includes 47 exons spanning a 437-kb locus,
207 which were all correctly assembled on a single scaffold.
208
209

Gene expression profiles across the milkweed bug life cycle
To augment published transcriptomic resources [36, 37], we sequenced three different
post-embryonic samples ("i5K" dataset, see Methods). We then compared the OGS
to the resulting *de novo* transcriptome and to a previously published embryonic and
maternal (ovary) transcriptome ("454" pyrosequencing dataset, [36]). Our OGS is
quite comprehensive, containing 90% of transcripts from each transcriptomic dataset
(Fig. 2a). The OGS also contains an additional 3,146 models (16% of OGS) not
represented in either transcriptome, including 163 *de novo* models encoding
chemoreceptors. Such genes are known for lineage-specific expansions and highly
tissue- and stage-specific expression ([38, 39], and see below), and our OGS captures
these genes with rare transcripts.
221

The OGS also incorporates many partial and unidentified 454 transcripts,
nearly trebling the transcripts with an assigned gene model or homology compared to
the original study (from 9% to 26%, by blastn, e<10^{-9}). This included 10,130
transcripts that primarily mapped to UTRs and previously lacked recognizable coding
sequence, such as for the *Oncopeltus brinker* ortholog, a BMP pathway component
([40], Supplemental Note 5.1.f), and the enzyme-encoding genes *CTP synthase* and
roquin. At the same time, the transcriptomes provided expression support for the
identification of multiple isoforms in the OGS, such as for the germline determinant
nanos [36]. More generally, most OGS gene models have expression support (91% of
19,690), with 74% expressed broadly in at least three of four samples (Fig. 2b). The
inclusion of a fifth dataset from a published adult library [37] provided only a 1%
gain in expression support, indicating that with the current study the expression data
volume for *Oncopeltus* is quite complete.
235

RNA-seq studies were further conducted to establish male, female, and nymph
specific gene sets (Fig. 2b-c, Supplemental Note 2.4), from which we also infer that
the published adult dataset of unspecified sex is probably male. Moreover, most
genes with stage-restricted or stage-enriched expression are in our male sample (Fig.
2b-c). For example, gustatory receptor (GR) genes show noticeable restriction to the
adult male and published adult (probable male) samples (n= 169 GRs: 40% no
expression, 27% only expressed in these two samples), with half of these expressed in
both biological replicates (52%). Interestingly, the nymphal sample is enriched for
genes encoding structural cuticular proteins (94%, which is >56% more than any
other sample). This likely reflects the ongoing molting cycles, with their cyclical
upregulation of chitin metabolism and cuticular gene synthesis [41], that are
experienced by the different instars and molt cycle stages of individuals pooled in this
sample. Lastly, gene sets with sex-specific enrichment across several hemipteroid
species substantiate known aspects of male and female reproduction (Fig. 2c: serine-
threonine kinases [42] or vitellogenin and other factors associated with oocyte
generation, respectively). Some of these enriched genes have unknown functions and
could comprise additional, novel factors associated with reproduction in *Oncopeltus*.
253
254

Protein orthology and hemipteran copy number comparisons

To further assay protein-coding gene content, we compared *Oncopeltus* with other arthropods. A phylogeny based on strictly conserved single copy orthologs correctly reconstructs the hemipteran and holometabolan clades' topologies (Fig. 3a, compare with Fig. 1a), although larger-scale insect relationships remain challenging [3].

We then expanded our appraisal to the Benchmarking Universal Single-Copy Orthologs dataset of 1,658 Insecta genes (BUSCO v3, [43]). Virtually all BUSCO genes are present in the *Oncopeltus* OGS (98.9%, Fig. 3b, Supplemental Note 6.1). Although some genes are fragmented, the assembly has a high level of BUSCO completeness (94.6%), independent of annotation prediction limitations that missed some exons from current gene models. Furthermore, BUSCO assessments can elucidate potential consequences of high heterozygosity, which could result in the erroneous inclusion of multiple alleles for a single gene. In fact, the fraction of duplicated BUSCO genes in *Oncopeltus* (1.4%) is low, compared to both the well-assembled bed bug genome (2.2%, [12]) and the pea aphid (4.8%), which is known to have lineage-specific duplications [6, 44]. Thus, by these quality metrics the *Oncopeltus* OGS and assembly are comparable to those of fellow hemipterans, strongly supporting the use of these resources in further comparisons.

We next categorized all proteins by conservation in global, clustering-based orthology analyses (OrthoDB, [1, 45]). As in most species, half of *Oncopeltus* proteins are highly conserved (Fig. 3a). Moreover, 98% of all *Oncopeltus* protein-coding genes have homology, expression, and/or curation support (Fig. 3c). Proteins without homology include species-specific chemoreceptors and antimicrobial peptides (Supplemental Note 5.1.h), as well as potentially novel or partial models. Overall, we estimate that the *Oncopeltus* protein repertoire is comparable to that of other insects in size and conservation. For the Hemiptera, *Oncopeltus* also has fewer missing orthology groups than either the kissing bug or pea aphid (Table S6.1). Indeed the pea aphid is a notable outlier, with its long branch in the phylogeny and for its large protein-coding gene content with low conservation (Fig. 3a). As more hemipteran genomes are sequenced, other species now offer less derived alternatives for phylogenomic comparisons.

Compared to the pea aphid [44], *Oncopeltus* is more conservative in presence and copy number for several signaling pathway components. In contrast to gene absences in the pea aphid, *Oncopeltus* retains orthologs of the EGF pathway component *sprouty*, the BMP receptor *wishful thinking*, and the hormone nuclear receptor *Hr96* (Supplemental Note 5.1.e). Also, whereas multiple copies were reported for the pea aphid, we find a single *Oncopeltus* ortholog for the BMP pathway components *decapentaplegic* and *Medea* and the Wnt pathway intracellular regulator encoded by *shaggy/GSK-3*, albeit with five potential isoforms of the latter (Supplemental Notes 5.1.f, 5.1.j). Duplications of miRNA and piRNA gene silencing factors likewise seem to be restricted to the pea aphid, even compared to other aphid species ([46], Supplemental Note 5.4.a). However, our survey of *Oncopeltus* and other hemimetabolous species reveals evidence for frequent, independent duplications of the Wnt pathway component *armadillo/β-catenin* ([47], Supplemental Note 5.1.j). Curiously, *Oncopeltus* appears to encode fewer histone loci than any other arthropod genome and yet exhibits a similar, but possibly independent, pattern of duplications of

304 histone acetyltransferases to those previously identified in *Cimex* and the pea aphid
305 (Supplemental Note 5.4.c).
306
On the other hand, we documented several notable *Oncopeltus*-specific
duplications. For the BMP transducer *Mad*, we find evidence for three paralogs in
Oncopeltus, where two occur in tandem and may reflect a particularly recent
duplication (Supplemental Note 5.1.f). Similarly, a tandem duplication of *wnt8*
appears to be unique to *Oncopeltus* (Supplemental Note 5.1.j). More striking is the
identification of six potential paralogs of *cactus*, a member of the Toll/NF-κB
signaling pathway for innate immunity, whereas the bed bug and kissing bug each
retain only a single copy ([48], Supplemental Note 5.1.g).
315
Lastly, we explored hemipteran-specific orthology groups against a backdrop
of 107 other insect species [1]. What makes a bug a bug in terms of protein-coding
genes? Several orthogroups contain potentially novel genes that show no homology
outside the Hemiptera and await direct experimental analysis, for which the
Hemiptera are particularly amenable (*e.g.*, [49-52, reviewed in 5]). Secondly, there
are hemipteran-specific orthogroups of proteins with recognized functional domains
and homologs in other insects, but where evolutionary divergence has led to lineage-
specific subfamilies. One example is a heteropteran-specific cytochrome P450 (CYP)
enzyme (EOG090W0V4B), which in *Oncopeltus* is expressed in all life history stages
(Fig. 2b). The expansion of CYP proteins is associated with potential insecticide
resistance, as specific P450s can confer resistance to specific chemicals (*e.g.,* [53, 54];
Supplemental Notes 5.3.b, 5.3.c). Hence, the identification of lineage-specific CYP
enzymes can suggest potential targets for integrated pest management approaches.
329
330
Transcription factor repertoires and homeobox gene evolution
Having explored the global protein repertoire, we next focused specifically on
transcription factors (TFs), which comprise a major class of proteins that has been
extensively studied in *Oncopeltus*. This is a class of key regulators of development
whose functions can diverge substantially during evolution and for which RNAi-
based experimental investigations have been particularly fruitful in the milkweed bug
(*e.g.*, [33, 34, 55-57], Supplemental Notes 5.1.a-e).
338
To systematically evaluate the *Oncopeltus* TF repertoire, we used a pipeline to
scan all predicted proteins and assign them to TF families, including orthology
assignments where DNA binding motifs could be predicted (see Methods, [58]). We
identified 762 putative TFs in *Oncopeltus*, which is similar to other insects for total
TF count and for the size of each TF family (Fig. 4a: note that the heatmap also
reflects the large, duplicated repertoire in the pea aphid, see also Tables S6.3-S6.5).
345
We were able to infer DNA binding motifs for 25% (n=189) of *Oncopeltus*
TFs, mostly based on data from *Drosophila melanogaster* (121 TFs) but also from
distantly related taxa such as mammals (56 TFs). Such high conservation is further
reflected in explicit orthology assignments for most proteins within several large TF
families, including the homeodomain (53 of 85, 62%), basic helix-loop-helix (bHLH,
35 of 45, 78%), and forkhead box (16 of 17, 94%) families. In contrast, most C2H2
zinc finger proteins lack orthology assignment (only 22 of 360, 6%). Across species,
the homeodomain and C2H2 zinc finger proteins are the two largest TF superfamilies

(Fig. 4a). Given their very different rates of orthology assignment, we probed further
into their pipeline predictions and the patterns of evolutionary diversification.
356
The number of homeodomain proteins identified by the pipeline displays a
narrow normal distribution across species (Fig. 4b, mean ± standard deviation: 97 ±
9), consistent with a highly conserved, slowly evolving protein family. Supporting
this, many *Oncopeltus* homeodomain proteins that were manually curated also
received a clear orthology assignment (Fig. 4c: pink), with only four exceptions (Fig.
4c: yellow). Only one case suggests a limitation of a pipeline that is not specifically
tuned to hemipteran proteins (Goosecoid). Manual curation of partial or split models
identified a further 11 genes encoding homeodomains, bringing the actual tally in
Oncopeltus to 96. Overall, we find the TF pipeline results to be a robust and
reasonably comprehensive representation of these gene classes in *Oncopeltus*.
367
These analyses also uncovered a correction to the published *Oncopeltus*
literature for the developmental patterning proteins encoded by the paralogs *engrailed*
and *invected*. These genes arose from an ancient tandem duplication prior to the
hexapod radiation. Their tail-to-tail orientation enables ongoing gene conversion
[59], making orthology discrimination particularly challenging. For *Oncopeltus*, we
find that the genes also occur in a tail-to-tail orientation and that *invected* retains a
diagnostic alternative exon [59]. These new data reveal that the purported *Oncopeltus*
engrailed ortholog in previous developmental studies (*e.g.*, [55, 60-63]) is in fact
invected (Supplemental Note 5.1.a).
377
378
Independent expansions of C2H2 zinc fingers within the Hemiptera
Unlike homeodomain proteins, C2H2 zinc finger (C2H2-ZF) repertoires are
prominent for their large family size and variability throughout the animal kingdom
[64], and this is further supported by our current analysis in insects. With >350
C2H2-ZFs, *Oncopeltus*, the pea aphid, termite, and some mosquito species have 1.5×
more members than the insect median (Fig. 4b). This is nearly half of all *Oncopeltus*
TFs. While the expansion in mosquitoes could have a single origin within the
Culicinae, the distribution in the Hemiptera, where *Cimex* has only 227 C2H2-ZFs,
suggests that independent expansions occurred in *Oncopeltus* and the pea aphid. Prior
to the sequencing of other hemipteran genomes, the pea aphid's large C2H2-ZF
repertoire was attributed to the expansion of a novel subfamily, APEZ, also referred
to as zinc finger 271-like [44].
391
392   In fact*,* manual curation in *Oncopeltus* confirms the presence of a subfamily
393 with similar characteristics to APEZ (Fig. 4c: yellow fraction). In *Oncopeltus* we find
394 >115 proteins of the ZF271 class that are characterized by numerous tandem repeats
395 of the C2H2-ZF domain and its penta-peptide linker, with 3-45 repeats per protein.
396
Intriguingly, we find evidence for ongoing evolutionary diversification of this
subfamily. A number of *Oncopeltus* ZF271-like genes occur in tandem clusters of 4-
8 genes – suggesting recent duplication events. Yet, clustered genes differ in gene
structure (number and size of exons), and we identified a number of probable ZF271-
like pseudogenes whose open reading frames have become disrupted – consistent with
high turnover. *Oncopeltus* ZF271-like proteins also differ in the sequence and length
of the zinc finger domains among themselves and compared to aphid proteins

(WebLogo analysis, [65]), similar to zinc finger array shuffling seen in humans [66].
Furthermore, whole-protein phylogenetic analysis supports independent, rapid
expansions in the pea aphid and *Oncopeltus* (Fig. 4d).
407

Clustered zinc finger gene expansion has long been recognized in mammals,
with evidence for strong positive selection to increase both the number and diversity
of zinc finger domains per protein as well as the total number of proteins [67]. This
was initially found to reflect an arms-race dynamic of co-evolution between selfish
transposable elements and the C2H2-ZF proteins that would repress them [68]. In
vertebrates, these C2H2-ZF proteins bind to the promoters of transposable elements
via their zinc finger arrays and use their Krüppel-associated box (KRAB) domain to
bind the chromatin-remodeling co-repressor KAP-1, which in turn recruits
methyltransferases and deacetylases that silence the targeted promoter [69].
417

Insects do not have a direct ortholog of vertebrate KAP-1 (Supplemental Note
5.4.d), and neither the aphid nor *Oncopeltus* ZF271-like subfamilies possess a KRAB
domain or any other domain besides the zinc finger arrays. However, close molecular
outgroups to this ZF271-like subfamily include the developmental repressor Krüppel
[70] and the insulator protein CTCF [71] (data not shown). Like these outgroups, the
Oncopeltus ZF271-like genes are strongly expressed: 98% have expression support,
with 86% expressed in at least three different life history stages (Fig. 2b). Thus, the
insect ZF271-like proteins may also play prominent roles in repressive DNA binding.
Indeed, we find evidence for a functional methylation system in *Oncopeltus*
(Supplemental Note 5.4.c), like the pea aphid, which would provide a means of gene
silencing by chromatin remodeling, albeit via mediators other than KAP-1.
429

However, an arms race model need not be the selective pressure that favors
insect ZF271-like family expansions. Recent analyses in vertebrates identified
sophisticated, additional regulatory potential by C2H2-ZF proteins, building upon
original transposable element binding for new, lineage-specific and even positive
gene regulation roles [66, 72, 73]. Moreover, although *Cimex* has half as many long
terminal repeat (LTR) repetitive elements as *Oncopeltus* and the pea aphid, overall we
do not find a correlation between relative or absolute repetitive content and ZF271-
like family expansion within the Hemiptera (see next section).
438
439

Proportional repeat content across hemipterans
With the aim of reducing assembly fragmentation and to obtain a better picture of
repeat content, we performed low coverage, long read PacBio sequencing in
Oncopeltus (Supplemental Note 2.3). Using PacBio reads in a gap-filling assay on
the Illumina assembly raised the total detected repetitive content from 25% to 32%,
while repeat estimations based on simultaneous assessment of Illumina and PacBio
reads nearly doubled this value to 58%. As expected, the capacity to identify repeats
is strongly dependent on assembly quality and sequencing technology, with the
Oncopeltus repetitive content underrepresented in the current (Illumina-only)
assembly. Furthermore, as increasing genome size compounds the challenge of
assembling repeats, the repeat content of the current assembly is lower than in species
with smaller genome sizes (Fig. 5a, with the sole exception of the honey bee), and we
therefore used our gap-filled dataset as a more accurate basis for further comparisons.
453

To support direct comparisons among hemipterans, we also performed our
RepeatModeler analysis on the bed bug and pea aphid assemblies. Repeats comprised
36% and 31% of the respective assemblies, similar to the gap-filled value of 32% in
Oncopeltus. Nevertheless, given the smaller sizes of these species' assemblies – 651
Mb in the bed bug and 542 Mb in the pea aphid – the absolute repeat content is much
higher in *Oncopeltus* (Fig. 5b). Excluding unknown repeats, the most abundant
transposable elements in *Oncopeltus* are LINE retrotransposons, covering 10% of the
assembly (Table S2.5). This is also the case in the bed bug (12%), while in the pea
aphid DNA transposons with terminal inverted repeats (TIRs) are the most abundant
(2% of the assembly identified here, and 4% reported from manual curation in the pea
aphid genome paper, [6]). Across species, the remaining repeat categories appear to
grow proportionally with assembly size, except for simple repeats, which were the
category with the largest relative increase in size after gap-filling in *Oncopeltus*
(Supplemental Note 2.3). However, given the mix of data types (Illumina [12] and
Sanger [6]), these patterns should be treated as hypotheses for future testing.
469
470
Lineage and genome size-related trends in insect gene structure
Both our manual curation work and BUSCO analyses highlighted the fact that
Oncopeltus genes are often comprised of many, small exons. We thus undertook a
comparative analysis to determine whether this is a general feature to be considered
for structural annotation of hemipteran genomes. We find that both lineage and
genome size can serve as predictors of gene structure.
477
Firstly, we created a high quality dataset of 30 functionally diverse, large
genes whose manual curation could reasonably ensure complete gene models across
seven species from four insect orders (Fig. 6a, Supplemental Note 6.3). Most species
encode the same total number of amino acids for these conserved proteins, with the
thrips *Frankliniella occidentalis* and the fruit fly being notable exceptions with larger
proteins (Fig. 6a: blue plot line). However, the means of encoding this information
differs between lineages, with hemipteroid orthologs comprised of twice as many
exons as their holometabolous counterparts (Fig. 6a: orange plot line). Thus, there is
an inverse correlation between exon number and exon size (Fig. 6a: orange vs. red
plot lines). This analysis corroborates and extends previous probabilistic estimates of
intron density, where the pea aphid as a sole hemipteran representative had the
highest intron density of ten insect species [74].
490
To test these trends, we next expanded our analysis to all manually curated
exons in two species from each of three orders (Hemiptera, Coleoptera, Diptera).
Here, we expect that curated exon sizes are accurate, without the need to assume that
entire gene models are complete. This large dataset corroborates our original
findings, with bugs having small exons while both the median and Q3 quartile reflect
larger exon sizes in beetles and flies (Fig. 6b). Notably, the median and median
absolute deviation are highly similar between species pairs within the Hemiptera and
Coleoptera. Meanwhile, the exon metrics within the Diptera support large protein
sizes as a drosophilid-specific, rather than dipteran-wide, feature.
500
501   Does the high exon count in the Hemiptera reflect an ancient, conserved
502 increase at the base of this lineage, or ongoing remodeling of gene structure with high
503 turnover? To assess the exact nature of evolutionary changes, we annotated intron

positions within multiple sequence alignments of selected proteins and plotted gains and losses onto the phylogeny, providing a total sample of 165 evolutionary changes at 148 discrete splice sites (Fig. 7, see also Supplemental Note 6.3 for gene selection and method). These data reveal several major correlates with intron gain or loss. The bases of both the hemipteroid and hemipteran radiations show the largest gains, while most losses occur in the dipteran lineage (Fig. 7: orange and purple shading, respectively). Furthermore, we find progressive gains across hemipteroid nodes, and it is only in this lineage that we additionally find species-specific splice changes for the highly conserved *epimerase* gene (Fig. 7: orange outline). Thus, we find evidence for both ancient intron gain and ongoing gene structure remodeling in this lineage.

Surprisingly, both *hemocytin* and *epimerase* – our exemplar genes with many (up to 74) and few exons (3-8 per species), respectively – show independent losses of the same splice sites in *Drosophila* and *Tribolium*. One feature these species share is a genome size 2.4-6.0× smaller than in the other species examined here (Fig. 7: red shading). Pairwise comparisons within orders also support this trend, as the beetle and fly species with larger genomes exhibit species-specific gains compared to intron loss in their sister taxa (Fig. 7: red outlines). Thus, genome size seems to positively correlate with intron number. However, lineage is a stronger predictor of gene structure: the coleopteran and dipteran species pairs have highly similar exon size metrics despite differences in genome size (Fig. 6b). A global computational analysis over longer evolutionary distances also supports a link between genome size and intron number in arthropods, although chelicerates and insects may experience different rates of evolutionary change in these features [75]. It will be interesting to see if the correlation with genome size is borne out in other invertebrate taxa.

The selective pressures and mechanisms of intron gain in the Hemiptera will be a challenge to uncover. While median exon size (Fig. 6b) could reflect species-specific nucleosome sizes [76, 77], this does not explain why only the Hemiptera seldom exceed this (Fig. 6b: Q3 quartile). Given gaps in draft genome assemblies, we remain cautious about interpreting (large) intron lengths but note that many hemipteran introns are too small to have harbored a functional transposase gene (*e.g.,* median intron size of 429 bp, n=69 introns in *hemocytin* in *Cimex*). Such small introns could be consistent with proliferation of non-autonomous short interspersed nuclear elements (SINEs). However, characterization of such highly divergent non-coding elements would require curated SINE libraries for insects, comparable to those generated for vertebrates and plants [76, 77]. Meanwhile, it appears that hemipteran open reading frames ≥160 bp are generally prevented by numerous in-frame stop codons just after the donor splice site. Most stop codons are encoded by the triplet TAA in both *Oncopeltus* and *Cimex* (data not shown), although these species' genomes are not particularly AT-rich (Table 1).

Even if introns are small, having gene loci comprised of numerous introns and exons adds to the cost of gene expression in terms of both transcription duration and mRNA processing. One could argue that a gene like *hemocytin*, which encodes a clotting agent, would require rapid expression in the case of wounding – a common occurrence in adult *Cimex* females due to the traumatic insemination method of reproduction [12]. Thus, as our molecular understanding of comparative insect and particularly hemipteran biology deepens, we will need to increasingly consider how

553　life history traits are manifest in genomic signatures at the structural level (*e.g.*, Figs.
554　5-7), as well as in terms of protein repertoires (Figs. 3-4).
555
556

Expansion after a novel lateral gene transfer (LGT) event in phytophagous bugs
In addition to the need for cuticle repair, traumatic insemination may be responsible
for the numerous LGT events predicted in the bed bug [12]. In contrast, the same
pipeline analyses [78] followed by manual curation predicted very few LGTs in
Oncopeltus, which lacks this unusual mating behavior. Here, we have identified 11
strong LGT candidates, and we confirmed the incorporation of bacterial DNA into the
milkweed bug genome for all five candidates chosen for empirical testing (Table
S2.4). Curiously, we find several LGTs potentially involved in bacterial or plant cell
wall metabolism that were acquired from different bacterial sources at different times
during hemipteran lineage evolution, including two distinct LGTs that are unique to
Oncopeltus and implicated in the synthesis of peptidoglycan, a bacterial cell wall
constituent (Supplemental Note 2.2).
569
Conversely, two other validated LGT candidates are implicated in cell wall
degradation. We find two strongly expressed, paralogous copies in *Oncopeltus* of a
probable bacterial-origin gene encoding an endo-1,4-beta-mannosidase enzyme
(MAN4, EC 3.2.1.78). Inspection of genome assemblies and protein accessions
reveals that this LGT event occurred after the infraorder Pentatomomorpha, including
the stink bug *Halyomorpha halys*, diverged from other hemipterans, including the bed
bug (Fig. 8a). Independent duplications then led to the two copies in *Oncopeltus* and
an astonishing nine tandem copies in *Halyomorpha* (Figs. 8b, S2.6). Since the
original LGT event, the *mannosidase* genes have gained introns that are unique to
each species and to subsets of paralogs (Fig. 8c). Thus, the "domestication" [79] of
mannosidase homologs as multi-exonic genes further illustrates the hemipteran
penchant for intron introduction and maintenance of small exons. The retention and
subsequent expansion of these genes implies their positive selection, consistent with
the phytophagous diet of these species. It is tempting to speculate that copy number
proliferation in the stink bug correlates with the breadth of its diet, as this agricultural
pest feeds on a number of different tissues in a range of host plants [80].
586
587

Cuticle development, structure, and warning pigmentation
Body cuticle is another trait associated with feeding ecology, particularly for
pigmentation. Furthermore, the milkweed bug has been a powerful model for
endocrine studies of hemimetabolous molting and metamorphosis since the 1960's
[22, 81-84]. Therefore, we next focused on the presence and function of genes
involved in cuticle development and structure.
594
Molting is triggered by the release of ecdysteroids, steroid hormones that are
synthesized from cholesterol by cytochrome P450 enzymes of the Halloween family
[85], and we were able to identify these in the *Oncopeltus* genome (Supplemental
Notes 5.2.b, 5.3.b). From the ecdysone response cascade defined in *Drosophila* [86],
we identified *Oncopeltus* orthologs of both early and late-acting factors, including
ecdysteroid hormones and their receptors. It will be interesting to see if the same
regulatory relationships are conserved in the context of hemimetabolous molting in
Oncopeltus. For example, *E75A* is required for reactivation of ecdysteroid production

during the molt cycle in *Drosophila* larvae [87] and likely operates similarly in
Oncopeltus, since *Of-E75A* RNAi prevents fourth-instar nymphs from molting to the
fifth instar (H. Kelstrup and L. Riddiford, unpublished data).
606
607   In hemipterans, activation of juvenile hormone (JH) signaling at molts
determines whether the insect progresses to another nymphal instar or, if lacking,
becomes an adult [50]. We were able to identify many components of the JH signal
transduction pathway in the *Oncopeltus* genome, including orthologs of *Methoprene-*
tolerant (*Met*), the JH receptor [50, 88], and the JH-response gene *Kr-h1* [50, 89, 90].
JH acts to determine cuticle identity through regulation of the *broad* gene in a wide
variety of insects, where different isoforms direct specific aspects of metamorphosis
in *Drosophila* [91, 92]. In *Oncopeltus*, *broad* expression directs progression through
the nymphal stages [93], but the effect of each isoform was unknown. We identified
three isoforms in *Oncopeltus* – *Z2*, *Z3*, and *Z4* – and performed isoform-specific
RNAi. In contrast to *Drosophila*, Broad isoform functions appear to be more
redundant in *Oncopeltus*, as knockdown of isoforms *Z2* and *Z3* has similar effects on
survival to adulthood as well as adult wing size and morphology (Fig. 9).
620
621   Regulators such as Broad initiate the transcription of a large battery of genes
that encode the structural components of the cuticle needed at each molt, consistent
with our expression analyses (Fig. 2b-c, discussed above). We identified 173 genes
encoding putative cuticle structural proteins in *Oncopeltus* (Supplemental Note 5.2.c).
Similar to other insects, the CPR family, with the RR-1 (soft cuticle), RR-2 (hard
cuticle), and unclassifiable types, constituted the largest cuticle protein family. While
several protein families are similar in size to those of other insects (CPAP1, CPAP3,
and TWDL: Table S5.12), we found a slight expansion in the *Oncopeltus* CPF family
(Fig. S5.14). For cuticle production, similar to the bed bug and the Asian longhorned
beetle [12, 30], we identified a single *chitin synthase* gene with conserved alternative
splice isoforms, which suggests that *chitin synthase 2* is a duplication specific to only
certain beetle and fly lineages within the Holometabola [94].
633
634   A major characteristic of the milkweed bug is the distinctive red-orange and
black aposematic (warning) coloration within the cuticle and epidermis that deters
predators (*e.g.*, Figs. 1, 9, [20, 21]). For black coloration, the melanin synthesis
pathway known from holometabolous insects (*e.g.*, [95, 96]) is conserved at the
sequence (Fig. S5.15) and functional [97, 98] level in *Oncopeltus*, supporting
conservation in hemimetabolous lineages as well. In contrast, production of the
primary warning coloration, pteridine red erythropterin [99], has not been as
extensively studied and remains an open avenue for hemimetabolous research. Pterin
pigments are synthesized from GTP through a series of enzymatic reactions [100].
Thus far in *Oncopeltus* we could identify orthologs of *punch*, which encodes a GTP
cyclohydrolase [101], and *sepia*, which is required for the synthesis of the red eye
pigment drosopterin [102]. The bright red color of *Oncopeltus* eggs may in part
reflect chemical protection transmitted parentally [103]. Thus, further identification
of pigmentation genes will provide fitness indicators for maternal contributions to
developmental success under natural conditions (*i.e.*, the presence of egg predators).
649
650

Chemoreception and metabolism in relation to feeding biology

Chemoreception and metabolism in relation to feeding biology
Aposematic pigmentation advertises the fact that toxins in the milkweed seed diet are
incorporated into the insects themselves, a metabolic feat that was independently
acquired in *Oncopeltus* and the monarch butterfly (*Danaus plexippus*), which shares
this food source and body coloration [37, 104]. Moreover, given the fundamental
differences between phytophagous, mucivorous, and hematophagous diets, we
investigated to what extent differences in feeding ecology across hemipterans are
represented in their chemoreceptor and metabolic enzyme repertoires.

659
660
661
Table 2. Numbers of chemoreceptor genes/proteins per family in selected insect
species. In some cases the number of proteins is higher than the number of genes due
to an unusual form of alternative splicing, which is particularly notable for the
Oncopeltus GRs. Data are shown for four Hemiptera as well as *Drosophila*
melanogaster, the body louse *Pediculus humanus*, and the termite *Zootermopsis*
nevadensis [11, 12, 105-109].
668

Species	Odorant	Gustatory	Ionotropic
Oncopeltus fasciatus [1]	120/121	115/169	37/37
Cimex lectularius [1,2]	48/49	24/36	30/30
Rhodnius prolixus [1,2]	116/116	28/30	33/33
Acyrthosiphon pisum [3]	79/79	77/77	19/19
Pediculus humanus [2]	12/13	6/8	14/14
Zootermopsis nevadensis	70/70	87/90	150/150
Drosophila melanogaster	60/62	60/68	65/65

669
670 [1] Hemiptera: Heteroptera
671 [2] independent acquisitions of hematophagy [16]
672 [3] Hemiptera, phloem-feeding
673
674
675
Insects must smell and taste their environment to locate and identify food,
mates, oviposition sites, and other essential cues. Perception of the enormous
diversity of environmental chemicals is primarily mediated by the odorant (OR),
gustatory (GR), and ionotropic (IR) families of chemoreceptors, which each encode
tens to hundreds of distinct proteins [109-112]. Chemoreceptor family size appears to
correlate with feeding ecology. *Oncopeltus* retains a moderate complement of each
family, while species with derived fluid nutrition diets (sap or blood) have relatively
depauperate OR and GR families (Table 2, Supplemental Note 5.3.f). In detail, a few
conserved orthologs such as the OrCo protein and a fructose receptor are found across
species, but other subfamilies are lineage specific. *Oncopeltus* and *Acyrthosiphon*
retain a set of sugar receptors that was lost independently in the blood-feeding bugs
(*Rhodnius* [11], *Cimex* [12]) and body louse (*Pediculus* [108]). Conversely,
Oncopeltus and *Cimex* retain a set of candidate carbon dioxide receptors, a gene
lineage lost from *Rhodnius*, *Acyrthosiphon*, and *Pediculus* [11, 12, 105], but which is
similar to a GR subfamily expansion in the more distantly related hemimetabolous
termite (Isoptera, [106]). Comparable numbers of IRs occur across the Heteroptera.
In addition to a conserved set of orthologs primarily involved in sensing temperature
and certain acids and amines, *Oncopeltus* has a minor expansion of IRs distantly
related to those involved in taste in *Drosophila*. The major expansions in each insect

lineage are the candidate "bitter" GRs ([113], Supplemental Note 5.3.f, Fig. S5.19).
In summary, *Oncopeltus* exhibits moderate expansion of specific subfamilies likely to
be involved in host plant recognition, consistent with it being a preferentially
specialist feeder with a potentially patchy food source [21, 114].
699
As host plant recognition is only the first step, we further explored whether
novel features of the *Oncopeltus* gene set are directly associated with its diet. We
therefore used the CycADS annotation pipeline [115] to reconstruct the *Oncopeltus*
metabolic network. The resulting BioCyc metabolism database for *Oncopeltus*
("OncfaCyc") was then compared with those for 26 other insect species ([116],
http://arthropodacyc.cycadsys.org/), including three other hemipterans: the pea aphid,
the green peach aphid, and the kissing bug (Tables 3-4). For a global metabolism
analysis, we detected the presence of 1085 Enzyme Commission (EC) annotated
reactions with at least one protein in the *Oncopeltus* genome (Supplemental Note 6.4,
Table S6.10). Among these, 10 enzyme classes (represented by 17 genes) are unique
and 17 are missing when compared to the other insects (Table 4, Table S6.11).
711
We then specifically compared amino acid metabolism in the four hemipterans
representing the three different diets. Eight enzymes are present only in *Oncopeltus*
(Table 4), including the arginase that degrades arginine (Arg) into urea and ornithine,
a precursor of proline (Pro). Given this difference, we extended our analysis to assess
species' repertoires for the entire urea cycle (Fig. 10a, Table S6.13). *Oncopeltus* and
six other species can degrade Arg but cannot synthesize it (Fig. 10b). Only the other
three hemipterans can neither synthesize nor degrade Arg via this cycle (Fig. 10c),
while most species have an almost complete cycle (Fig. 10d). This suggests that the
ability to synthesize Arg was lost at the base of the Hemiptera, with subsequent,
independent loss of Arg degradation capacity in the aphid and *Rhodnius* lineages.
Retention of Arg degradation in *Oncopeltus* might be linked to the milkweed seed
food source, as most seeds are very rich in Arg [117], and Arg is indeed among the
metabolites detected in *Oncopeltus* [118]. However, the monarch butterfly is one of
only a handful of species that retains the complete Arg pathway (Fig. 10d: blue text).
Despite a shared food source, these species may therefore differ in their overall Arg
requirements, or – in light of a possible group benefit of *Oncopeltus* aggregation
during feeding ([21]; *e.g.*, Fig. 1b) – in their efficiency of Arg uptake.
729
Other enzymes are also present only in the milkweed bug compared to the
other examined hemipterans (Table S6.12). Like other insects [116], *Oncopeltus*
retains the ability to degrade tyrosine (Tyr). This pathway was uniquely lost in the
aphids, where this essential amino acid is jointly synthesized – and consumed – by the
aphid host and its endosymbiotic bacteria [6, 7, 17, 119]. Conversely, a gain specific
to the milkweed bug lineage was the duplication of the Na+/K+ ATPase alpha
subunits whose amino acid substitutions confer resistance to milkweed cardenolides
[37, 120]. In the *Oncopeltus* genome, we find support for the recent nature of these
duplications: the genes encoding subunits ATPα1B and ATPα1C occur as a tandem
duplication, notably on a scaffold that also harbors one of the clustered ZF271-like
gene expansions (see above).
741
742

Table 3. Hemipteran ArthropodaCyc database summaries.
Overview statistics for the newly created database for *Oncopeltus fasciatus* (Ofas) in comparison with public databases for *Rhodnius prolixus* (Rpro), *Acyrthosiphon pisum* (Apis), and *Myzus persicae* (Mper) available from [116]. Based on OGS v1.1.

Species ID	*Ofas*	*Rpro*	*Apis*	*Mper*	*Mper*
Gene set ID	OGS v1.1	RproC1.1 (Built on RproC1 assembly)	OGS v2.1b (Built on Acyr_2.0 assembly)	Clone G006 v1.0	Clone O v1.0
CycADS Database ID	OncfaCyc	RhoprCyc	AcypiCyc v2.1b	Myzpe_G006 Cyc	Myzpe_O Cyc
Total mRNA[1]	19,673	15,437	36,195	24,814	24,770
Pathways	294	312	307	319	306
Enzymatic Reactions	2,192	2,366	2,339	2,384	2,354
Polypeptides	19,820	15,471	36,228	24,849	24,805
Enzymes	3,050	2,660	5,087	4,646	4,453
Compounds	1,506	1,665	1,637	1,603	1,655

[1] In the BioCyc databases all splice variants are counted in the summary tables for genes.

Table 4. Hemipteran ArthropodaCyc annotations of metabolic genes.
Taxonomic abbreviations are as in Table 3.

	Ofas	*Rpro*	*Apis*	*Mper*
Global metabolism				
EC[1] present in the genome	1085	1241	1288	1222
EC unique to this genome[2]	10	13	23	5
EC missing only in this genome[2]	17[4]	8	2	6
Amino acid metabolism (KEGG)				
EC present in the genome	169	188	195	185
EC unique to this genome[2]	2	1	6	1
EC missing only in this genome[2]	5	2	0	2
EC unique to this genome[3]	8	10	12	8
EC missing only in this genome[3]	14	5	0	2

[1] "EC" refers to the number of proteins, as represented by their unique numerical designations within the Enzyme Commission (EC) classification system for enzymes and their catalytic reactions.
[2] in comparison to all other insects from ArthropodaCyc
[3] in comparison among the four hemipterans
[4] includes three EC categories added in OGS v1.2 (see also Table S6.11)

CONCLUSIONS

The integrated genomic and transcriptomic resources presented here for the milkweed bug *Oncopeltus fasciatus* (Figs. 2,5) underpin a number of insights into evolutionary and developmental genomics. Our macroevolutionary comparisons across insect orders, now extended to the hemimetabolous Hemiptera, reveal unexpected patterns of molecular evolution. We also show how hemipteran feeding ecology and suites of related biological characters are reflected in the genome.

The gene structure trends we identified, with lineage predominating over genome size as a predictor and with many intron gains in the hemipteroid lineage (Figs. 6,7), offer initial parameters and hypotheses for the Hemiptera, Coleoptera, and Diptera. Such ordinal-level parameters can be evaluated against new species' data and also inform customized pipelines for automated gene model predictions. At the same time, it will be interesting to explore the ramifications of hemipteroid intron gains, as there are few documented lineages with episodic intron gain [77]. For example, possessing more, small exons may provide greater scope to generate protein modularity via isoforms based on alternative exon usage [121]. Furthermore, with the larger genome sizes and lower gene densities of hemipteroids compared to the well-studied Hymenoptera, it remains open whether hemipteroid gene and intron size may also correlate with recombination rates [122].

Our analyses also highlight new directions for future experimental research, building on *Oncopeltus*'s long-standing history as a laboratory model and its active research community in the modern molecular genetics era (*e.g.*, Fig. 9, [25-27]). Functional testing will clarify the roles of genes we have identified as unique to the Hemiptera, including those implicated in chemical protection, bacterial and plant cell wall metabolism, or encoding wholly novel proteins (Figs. 3,8, see also Supplemental Note 2.2). Meanwhile, the prominent and species-specific expansions specifically of ZF271-like zinc fingers (Fig. 4), combined with the absence of the co-repressor KAP-1 in insects, argues for investigation into alternative interaction partners, which could clarify the nature of these zinc fingers' regulatory roles and their binding targets.

One key output of this study is the generation of a metabolism database for *Oncopeltus*, contributing to the ArthropodaCyc collection (Table 3). In addition to comparisons with other species (Fig. 10), this database can also serve as a future reference for studies that use *Oncopeltus* as an ecotoxicology model (*e.g.*, [123]). While we have primarily focused on feeding ecology in terms of broad comparisons between phytophagy and fluid feeding, *Oncopeltus* is also poised to support future work on nuances among phytophagous species. Despite its milkweed diet in the wild, the lab strain of *Oncopeltus* has long been adapted to feed on sunflower seeds, demonstrating a latent capacity for more generalist phytophagy [114]. This potential may also be reflected in a larger gustatory receptor repertoire than would be expected for an obligate specialist feeder (Table 2). Thus, *Oncopeltus* can serve as a reference species for promiscuously phytophagous pests such as the stink bug. Finally, we have identified a number of key genes implicated in life history trade-offs in *Oncopeltus*, for traits such as cardenolide tolerance, pigmentation, and plasticity in reproduction under environmental variation. The genome data thus represent an important tool to elucidate the proximate mechanisms of fundamental aspects of life history evolution in both the laboratory and nature.

METHODS

(More information is available in the supplementary materials, Additional file 1.)

Milkweed bug strain, rearing, and DNA/RNA extraction

The milkweed bug *Oncopeltus fasciatus* (Dallas), Carolina Biological Supply strain (Burlington, North Carolina, USA), was maintained in a laboratory colony under standard husbandry conditions (sunflower seed and water diet, 25 ℃, 12:12 light-dark photoperiod). Voucher specimens for an adult female (record # ZFMK-TIS-26324) and adult male (record # ZFMK-TIS-26325) have been preserved in ethanol and deposited in the Biobank of the Centre for Molecular Biodiversity Research, Zoological Research Museum Alexander Koenig, Bonn, Germany (https://www.zfmk.de/en/biobank).

Genomic DNA was isolated from individual, dissected adults using the Blood & Cell Culture DNA Midi Kit (G/100) (Qiagen Inc., Valencia, California, USA). Total RNA was isolated from individual, dissected adults and from pooled, mixed-instar nymphs with TRIzol Reagent (Invitrogen/ Thermo Fisher Scientific, Waltham, Massachusetts, USA). Dissection improved accessibility of muscle tissue by disrupting the exoskeleton, and gut material was removed.

Genome size calculations (flow cytometry, *k*-mer estimation)

Genome size estimations were obtained by flow cytometry with Hare and Johnston's protocol [124]. Four to five females and males each from the Carolina Biological Supply lab strain and a wild strain (collected from Athens, Georgia, USA; GPS coordinates: 33° 56' 52.8216" N, 83° 22' 38.3484'' W) were measured (see also Supplemental Note 2.1.a). At the bioinformatic level, we attempted to estimate genome size by *k*-mer spectrum distribution analysis for a range of *k*=15 to 34 counted with Jellyfish 2.1.4 [125] and bbmap [126], graphing these counts against the frequency of occurrence of *k*-mers (depth), and calculating genome size based on the coverage at the peak of the distribution (Supplemental Note 2.1.b).

Genome sequencing, assembly, annotation, and official gene set overview

Library preparation, sequencing, assembly, and automatic gene annotation were conducted at the Baylor College of Medicine Human Genome Sequencing Center (as in [12, 30]). About 1.1 billion 100-bp paired-end reads generated on an Illumina HiSeq2000s machine were assembled using ALLPATHS-LG [127], from two paired-end (PE) and two mate pair (MP) libraries specifically designed for this algorithm (Supplemental Note 1). Three libraries were sequenced from an individual adult male (180- and 500-bp PE, 3-kb MP), with the fourth from an individual adult female (8-10-kb MP). The final assembly (see metrics in Table 1) has been deposited in GenBank (accession GCA_000696205.1).

Automated annotation of protein-coding genes was performed using a Maker 2.0 annotation pipeline [128] tuned specifically for arthropods (Supplemental Note 3). These gene predictions were used as the starting point for manual curation via the Apollo v.1.0.4 web browser interface [129], and automatic and manual curations were compiled to generate the OGS (see also Supplemental Note 4). Databases of the genome assembly, Maker automatic gene predictions, and OGS v1.1 are available through the i5K Workspace@NAL [130], and the Ag Data Commons data access

system of the United States Department of Agriculture's (USDA) National
Agricultural Library as individual citable databases [131-133]. The current version of
the gene set, OGS v1.2, is deposited at GenBank as an 'annotation-only' update to the
Whole Genome Shotgun project (accession JHQO00000000). Here, we describe
version JHQO02000000. The annotations can be downloaded from NCBI's ftp site,
ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/000/696/205/GCA_000696205.1_Ofas_1.0/.
876
Repeat content analysis
Repetitive regions were identified in the *Oncopeltus* genome assembly with
RepeatModeler Open-1.0.8 [134] based on a species-specific repeat library generated
de novo with RECON [135], RepeatScout [136], and Tandem Repeats Finder [137].
Then, RepeatMasker Open-4.0 [138] was used to mask repeat sequences based on the
RepeatModeler library. Given the fragmented nature of the assembly, we attempted
to fill and close assembly gaps by sequencing additional material, generating long
reads with single molecule real time sequencing on a PacBio RS II machine
(estimated coverage of 8x). Gap filling on the Illumina assembly scaffolds was
performed with PBJelly version 13.10.22, and the resulting assembly was used for
repeat content estimation and comparison with *Cimex lectularius* and *Acyrthosiphon*
pisum (see also Supplemental Note 2.3).
889
Transcriptome resources
Total RNA from three distinct life history samples (pooled, mixed-instar nymphs; an
adult male; an adult female) was also sequenced on an Illumina HiSeq2000s machine,
producing a total of 72 million 100-bp paired-end reads (Supplemental Note 1.3,
Table S1.1; GenBank Bioproject: PRJNA275739). These expression data were used
to support the generation of the OGS at different stages of the project: as input for the
evidence-guided automated annotation with Maker 2.0 (Supplemental Note 3), as
expression evidence tracks in the Apollo browser to support the community curation
of the OGS, and, once assembled into a *de novo* transcriptome, as a point of
comparison for quality control of the OGS.
900
The raw RNA-seq reads were pre-processed by filtering out low quality bases
(phred score <30) and Truseq adapters with Trimmomatic-0.30. Further filtering
removed ribosomal and mitochondrial RNA sequences with Bowtie 2 [139], based on
a custom library built with all hemipteran ribosomal and mitochondrial RNA
accessions from NCBI as of 7[th] February 2014 (6,069 accessions). The pooled,
filtered reads were mapped to the genome assembly with Tophat2-PE on CyVerse
[140]. A second set of RNA-seq reads from an earlier study ("published adult"
dataset, [37]) was also filtered and mapped in the same fashion, and both datasets
were loaded into the *Oncopeltus* Apollo instance as evidence tracks (under the track
names "pooled RNA-seq - cleaned reads" and "RNA-seq raw PE reads Andolfatto et
al", respectively).
912
Additionally, a *de novo* transcriptome was generated from our filtered RNA-
seq reads (pooled from all three samples prepared in this study) using Trinity [141]
and TransDecoder [142] with default parameters. This transcriptome is referred to as
"i5K", to distinguish it from a previously published maternal and early embryonic
transcriptome for *Oncopeltus* (referred to as "454", [36]). Both the i5K and 454
transcriptomes were mapped to the genome assembly with GMAP v. 2014-05-15 on

CyVerse. These datasets were also loaded into the Apollo browser as evidence tracks
to assist in manual curation.
921
Life history stage-specific and sex-specific expression analyses in hemipteroids
Transcript expression of the OGS v1.1 genes was estimated by running RSEM2 [143]
on the filtered RNA-seq datasets for the three i5K postembryonic stages against the
OGS v1.1 cDNA dataset. Transcript expression was then based on the transcripts per
million (TPM) value. The TPM values were processed by adding a value of 1 (to
avoid zeros) and then performing a log2-transformation. The number of expressed
genes per RNA-seq library was compared for TPM cutoffs of >1, >0.5, and >0.25. A
>0.25 cutoff was chosen, which reduced the number of expressed genes by 6.6%
compared to a preliminary analysis based on a simple cutoff of $\geq$10 mapped reads per
transcript, while the other TPM cutoffs were deemed too restrictive (reducing the
expressed gene set by >10%). This analysis was also applied to the "published adult"
dataset [37]. To include embryonic stages in the comparison, transcripts from the 454
transcriptome were used as blastn queries against the OGS v1.1 cDNA dataset (cutoff
e-value $<10^{-5}$). The results from all datasets were converted to binary format to
generate Venn diagrams (Fig. 2b).
937
Statistically significant sex-specific and developmental stage-specific gene
enrichment was determined from RNA-seq datasets according to published methods
[144, 145], with modifications. Data from *Oncopeltus* (see previous methods section,
Bioproject: PRJNA275739) were compared between stages and pairwise with the
hemipterans *Cimex lectularius*, PRJNA275741; *Acyrthosiphon pisum*, PRJNA209321;
and *Pachypsylla venusta*; PRJNA275248; as well as with the hemipteroid
Frankliniella occidentalis (Thysanoptera), PRJNA203209 (see also Fig. 2c,
Supplemental Note 2.4).
946
947 **Protein gene orthology assessments via OrthoDB and BUSCO analyses**
948 These analyses follow previously described approaches and with the current database
949 and pipeline versions [1, 43, 45, 146]. See Supplemental Note 6.1 for further details.
950
Global transcription factor identification
Likely transcription factors (TFs) were identified by scanning the amino acid
sequences of predicted protein-coding genes for putative DNA binding domains
(DBDs), and when possible, the DNA binding specificity of each TF was predicted
using established procedures [58]. Briefly, all protein sequences were scanned for
putative DBDs using the 81 Pfam [147] models listed in Weirauch and Hughes [148]
and the HMMER tool [149], with the recommended detection thresholds of Per-
sequence Eval < 0.01 and Per-domain conditional Eval < 0.01. Each protein was
classified into a family based on its DBDs and their order in the protein sequence
(*e.g.*, bZIPx1, AP2x2, Homeodomain+Pou). The resulting DBD amino acid
sequences were then aligned within each family using Clustal Omega [150], with
default settings. For protein pairs with multiple DBDs, each DBD was aligned
separately. From these alignments, the sequence identity was calculated for all DBD
sequence pairs (*i.e.*, the percent of amino acid residues that are identical across all
positions in the alignment). Using previously established sequence identity thresholds
for each family [58], the predicted DNA binding specificities were mapped by simple
transfer. For example, the DBD of OFAS001246-RA is 98% identical to the
Drosophila melanogaster Bric a Brac 1 (Bab1) protein. Since the DNA binding

specificity of Bab1 has already been experimentally determined, and the cutoff for the
Pipsqueak family TFs is 85%, we can infer that OFAS001246-RA will have the same
binding specificity as *Drosophila* Bab1.
972

RNA interference
Double-stranded RNA (dsRNA) was designed to target the final, unique exon of the
broad isoforms *Z2, Z3*, and *Z4*. A portion of the coding sequence for the zinc finger
region from these exons (179 bp, 206 bp, and 216 bp, respectively) was cloned into a
plasmid vector and used as template for *in vitro* RNA synthesis, using the gene-
specific primer pairs: Of-Z2_fwd: 5′-ATGTGGCAGACAAGCATGCT-3′; Of-
Z2_rev: 5′-CTAAAATTTGACATCAGTAGGC-3′; Of-Z3_fwd: 5′-
ccttctcctgttactactcac-3′; Of-Z3_rev: 5′-ttatatgggcggctgtccaa-3′; Of-Z4_fwd: 5′-
AACACTGACCTTGGTTACACA-3′; Of-Z4_rev: 5′-
TAGGTGGAGGATTGCTAAAATT-3′. Two separate transcription reactions (one
for each strand) were performed using the Ambion MEGAscript kit (Ambion, Austin,
Texas, USA). The reactions were purified by phenol/chloroform extraction followed
by precipitation as described in the MEGAscript protocol. The separate strands were
re-annealed in a thermocycler as described previously [33]. Nymphs were injected
with a Hamilton syringe fitted with a 32-gauge needle as described [55]. The
concentration of *Of-Z2*, *Of-Z3* and *Of-Z4* dsRNA was 740 ng/µl, 1400 ng/µl, and
1200 ng/µl, respectively. All nymphs were injected within 8 hours of the molt to the
fourth (penultimate juvenile) instar (n ≥12 per treatment: see Fig. 9). Fore- and
hindwings were then dissected from adults and photographed at the same scale as
wings from wild type, uninjected controls.
993

CycADS annotation and OncfaCyc database generation
We used the Cyc Annotation Database System (CycADS, [115]), an automated
annotation management system, to integrate protein annotations from different
sources into a Cyc metabolic networks reconstruction that was integrated into the
ArthropodaCyc database. Using our CycADS pipeline, *Oncopeltus fasciatus* proteins
from the official gene set OGS v1.1 were annotated using different methods –
including KAAS [151], PRIAM [152], Blast2GO [153, 154], and InterProScan with
several approaches [155] – to obtain EC and GO numbers. All annotation
information data were collected in the CycADS SQL database and automatically
extracted to generate appropriate input files to build or update BioCyc databases [156]
using the Pathway Tools software [157]. The OncfaCyc database, representing the
metabolic protein-coding genes of *Oncopeltus*, was thus generated and is now
included in the ArthropodaCyc database, a collection of arthropod metabolic network
databases ([116], http://arthropodacyc.cycadsys.org/).
1008

1009

1010   **FIGURE LEGENDS**
1011

Fig. 1. The large milkweed bug, *Oncopeltus fasciatus*, shown in its phylogenetic
and environmental context.
(a) Species tree of selected Hemiptera with genomic and transcriptomic resources,
based on phylogenetic analyses and divergence time estimates in [3]. Species marked
with an asterisk (*) have published resources; those with the appellation "i5K" are
part of a current pilot project supported by the Baylor College of Medicine Human
Genome Sequencing Center and the National Agricultural Library of the USDA.

Note that recent analyses suggest the traditional infraorder Cimicomorpha, to which
Rhodnius and *Cimex* belong, may be paraphyletic [16].
(b-c) Milkweed bugs on their native food source, the milkweed plant: gregarious
nymphs of different instars on a milkweed seed pod (b), and pale, recently eclosed
adults and their shed exuvia (c). Images were taken at Avalon Park and Preserve,
Stony Brook, New York, USA, courtesy of Deniz Erezyilmaz, used with permission.
(d) Individual bugs, shown from left to right: first instar nymphs (ventral and dorsal
views) and adults (dorsal and lateral views); images courtesy of Kristen Panfilio
(nymphs) and Jena Johnson (adults), used with permission. The arrow labels the
labium (the "straw"), part of the hemipteran mouthpart anatomy adapted for feeding
by piercing and sucking.
1030
Fig. 2. Comparisons of the official gene set and transcriptomic resources for
Oncopeltus fasciatus.
(a) Area-proportional Venn diagram comparing the OGS v1.1 ("OGS"), a Trinity *de*
novo transcriptome from the three post-embryonic RNA-seq samples ("i5K"), and the
maternal and embryonic transcriptome from 454 data ("454" [36]). Sample sizes and
the fraction of each transcriptome represented in the OGS are indicated (for the 454
dataset, only transcripts with homology identification were considered). The unique
fraction of each set is also specified (%). Dataset overlaps were determined by blastn
(best hit only, e-value $<10^{-9}$).
(b) Venn diagram of gene model expression support across four life history samples.
Values are numbers of gene models, with percentages also given for the largest
subsets. Note that the "Embryo/Maternal" sample derives from 454 pyrosequencing
data and therefore has a smaller data volume than the other, Illumina-based samples.
(c) Summary of sex- and developmental stage-specific RNA-seq comparisons across
hemipteroid species: *Acyrthosiphon pisum, Apis; Cimex lectularius, Clec;*
Frankliniella occidentalis, Focc (thysanopteran outgroup)*; Oncopeltus fasciatus,*
Ofas; Pachypsylla venusta, Pven. n.d., not determined. For complete numerical
details see Supplemental Note 2.4
Analyses based on OGS v1.1.
1050
Fig. 3. Orthology comparisons and phylogenetic placement of *Oncopeltus*
***fasciatus* among other Arthropoda.**
(a) Comparisons of protein-coding genes in 12 arthropod species, with the Hemiptera
highlighted in red text. The bar chart shows the number of proteins per conservation
level (see legend), based on OrthoDB orthology clustering analyses. To the left is a
maximum likelihood phylogeny based on concatenation of 395 single-copy orthologs
(all nodes have 100% support unless otherwise noted; branch length unit is
substitutions per site). The inset pie chart shows the proportion of proteins per
conservation level in *Oncopeltus* (*"Ofas"*). See also Supplemental Note 6.1.
(b) BUSCO-based analysis of *Oncopeltus* compared to other hemipterans for ortholog
presence and copy number in both the assembly and OGS resources, using four-letter
species abbreviations (full names in panel a).
(c) Proportion of *Oncopeltus* proteins that have expression and/or curation validation
support per conservation level (same color legend as in (a)). Expression support is
based on the life history stage data in Fig. 2b.
Analyses based on OGS v1.1.
1067

Fig. 4. Distribution of transcription factor families across insect genomes.
(a) Heatmap depicting the abundance of 74 transcription factor (TF) families across
16 insect genomes (Hemiptera highlighted in red text), with *Daphnia* as an outgroup,
based on the presence of predicted DNA binding domains (see Methods). The color
key has a log (base 2) scale (light blue means the TF family is completely absent).
Values are in Table S6.3.
(b) Bar graph showing the number of proteins of each of the two most abundant TF
families, homeodomains and C2H2 zinc fingers (ZFs), per species using four-letter
abbreviations (full names in panel a). Solid lines demarcate insect orders: Hemiptera
(Hemipt.), Hymenoptera (Hym.), Coleoptera (Col.), and Diptera (Dipt.). The dashed
line demarcates the dipteran family Culicidae (mosquitoes).
(c) Proportions of *Oncopeltus* homeodomain (HD) and C2H2 zinc finger proteins
with orthology assignment (predicted DNA binding specificity) and/or manual
curation. "Classified" refers to automated classification of a protein to a TF family,
but without a specific orthology assignment.
(d) Maximum likelihood phylogeny of representative subsets of the zinc finger 271-
like family in *Oncopeltus* (49 proteins, blue text) and the pea aphid (55 proteins, black
text), with chelicerate (red text) and holometabolan (yellow text) outgroups (16
proteins, 7 species), based on the *Oncopeltus* OGS and GenBank protein accessions.
Gaps were removed during sequence alignment curation; all nodes have ≥50%
support; branch length unit is substitutions per site [158]. Key nodes are circled for
the clades containing all aphid or all *Oncopeltus* proteins (82% support each), and for
each 'core' clade comprised exclusively of proteins from each species (97% and
100%, respectively; triangles shown to scale for branch length and number of clade
members). Branch length unit is substitutions per site.
Analyses based on OGS v1.1.
1094
Fig. 5. Comparison of repeat content estimations.
(a) Comparison of total repetitive content among insect genomes. The three values
for *Oncopeltus* are shown (in ascending order: original Illumina assembly, gap-filled
assembly, Illumina-PacBio hybrid estimate). Values for the three hemipterans labeled
in red text are from RepeatModeler (gold bars for the pea aphid and bed bug; blue and
gold bars for *Oncopeltus*). All other values are from the respective genome papers,
including a second value corresponding to the published repeat content for the first
version of the aphid genome [6, 10, 106, 159-164]. Species abbreviations as in Fig. 4,
and additionally: *Nlug, Nilaparvata lugens; Lmig, Locusta migratoria; Bmor, Bombyx*
mori; Aalb, Aedes albopictus.
(b) Comparison of repetitive element categories between three hemipteran genomes,
based on results from RepeatModeler. Here we present assembly coverage as actual
sequence length (Mb) to emphasize the greater repeat content in *Oncopeltus* (based on
the gap-filled assembly, see also Supplemental Note 2.3).
1109
Figure 6. Trends in gene structure show hemipteroid-specific tendencies.
(a) Median values per species for protein size, exon size, and exon number for a
curated set of highly conserved genes encoding large proteins of diverse functional
classes (see also Supplemental Note 6.3). Sample sizes are indicated, with 11 genes
for which orthologs were evaluated in all species. Where it was not possible to
analyze all 30 genes for a given species, equal sampling was done across the range of
protein sizes of the complete dataset, based on the *Cimex* ortholog sizes (1:1:1
sampling from big:medium:small subcategories of 10 genes each).

(b) Box plot representations of coding sequence exon size (aa) for two species from
each of three insect orders, based on datasets of unique coding sequence exons (one
isoform per gene) and excluding terminal exons <10 aa (as most of those exons may
rather be UTRs or a small placeholder N-terminal exon based on automated Maker
model predictions). Only manually curated gene models were considered for the i5K
species, including *Oncopeltus*; the entire OGS was used for *Tribolium* and
Drosophila. For clarity, outliers are omitted; whiskers represent 1.5× the value of the
Q3 (upper) or Q2 (lower) quartile range. MAD, median absolute deviation.
Species are represented by their four-letter abbreviations, with their ordinal
relationships given below the phylogeny in panel (a): Hemip., Hemiptera; Thys.,
Thysanoptera; Col., Coleoptera; Dipt., Diptera. Species abbreviations as in Figs. 2,4
and additionally: *Gbue, Gerris buenoi* [165]; *Agla, Anoplophora glabripennis* [30];
Ccap, Ceratitis capitata [166].
1131
Fig 7. Splice site evolution correlates with both lineage and genome size.
Splice site changes are shown for *hemocytin* (blue text), *Tenascin major* (*Ten-m*,
turquoise text), and *UDP-galactose 4′-epimerase* (brown text), mapped onto a species
tree of eight insects. Patterns of splice site evolution were inferred based on the most
parsimonious changes that could generate the given pattern within a protein sequence
alignment of all orthologs (see also Supplemental Note 6.3 for methodology and data
sources). If inferred gains or losses were equally parsimonious, we remained agnostic
and present a range for the ancestral number of splice sites present at the base of the
tree, where the bracketed number indicates how many ancestral positions are still
retained in all species. Along each lineage, subsequent changes are indicated in
brackets, with the sign indicating gains (+) or losses (-). Values shown to the right are
species-specific changes. The values shown between the *D. melanogaster* and *T.*
castaneum lineages denote changes that have occurred independently in both species.
Colored boxes highlight the largest sources of change, as indicated in the legend.
Species are represented by their four-letter abbreviations (as in Fig. 6), and estimated
genome sizes are indicated parenthetically (measured size: [12, 30, 163, 166, 167];
draft assembly size: GenBank Genome IDs 14741 and 17730). Divergence times are
shown in gray and given in millions of years [3]. Abbreviations as in Figs. 4,6, and
also: Col., Coleoptera; Dipt., Diptera; Hemip., Hemiptera; Hemipt., hemipteroid
assemblage (including *F. occidentalis*); n.d., no data.
1152
Fig. 8. Lateral gene transfer introduction and subsequent evolution within the
Hemiptera for mannosidase-encoding genes.
(a) Species tree summary of evolutionary events. Stars represent the original LGT
introduction and subsequent copy number gains (see legend).
(b) Maximum likelihood phylogeny of mannosidase proteins, including bacterial
sequences identified among the best GenBank blastp hits for *Oncopeltus* and
Halyomorpha (accession numbers as indicated, and for "Other bacteria" are:
ACB22214.1, AEE17431.1, AEI12929.1, AEO43249.1, AFN74531.1, CDM56239.1,
CUA67033.1, KOE98396.1, KPI24888.1, OAN41395.1, ODP26899.1, ODS11151.1,
OON18663.1, PBD05534.1, SIR54690.1, WP096035621.1, YP001327394.1). All
nodes have ≥50% support from 500 bootstrap replicates [168]. Triangles are shown
to scale for branch length and number of clade members; branch length unit is
substitutions per site. See also Fig. S2.6.
(c) Manually curated protein sequence alignment for the N-terminal region only.
Splice sites ("|" symbol) are shown, where one position is ancestral and present in all

paralogs of a given species (magenta) and one position occurs in a subset of paralogs
and is presumed to be younger (cyan, within the 5′ UTR in *Halyomorpha*). Residues
highlighted in yellow are conserved between the two hemipteran species. The
Oncopeltus paralog represented in the OGS as OFAS017153-RA is marked with an
asterisk to indicate that this version of the gene model is incomplete and lacks the
initial exon (gray text in the alignment). For clarity, only the final three digits of the
Halyomorpha GenBank accessions are shown (full accessions: XP_014289XXX).
1175
Fig. 9. Isoform-specific RNAi based on new genome annotations affects the
molting and cuticle identity gene *broad*.
(a) Genomic organization of the cuticle identity gene *broad*. The regions used as
template to generate isoform-specific dsRNA are indicated (red asterisks: the final,
unique exons of each isoform). Previous RNAi studies targeted sequence within
exons 1-5 that is shared among all isoforms (dashed red box, [93]).
(b) Knock down of the *Oncopeltus Z2* or *Z3 broad* isoforms at the onset of the
penultimate instar resulted in altered nymphal survival and morphogenesis that was
reflected in the size and proportion of the fore and hind wings at the adult stage
(upper and lower images, respectively, shown to the same scale for all wings). We
did not detect any effect on the wing phenotype when targeting the *Z4*-specific exon,
demonstrating the specificity of the zinc finger coding region targeted by RNAi.
Experimental statistics are provided in the figure inset, including for the buffer-
injected negative control.
1190
Fig. 10. Comparison of the urea cycle of *Oncopeltus* with 26 other insect species.
(a) Detailed diagram of the urea cycle (adapted from KEGG).
(b) Group of 7 species, including *Oncopeltus*, for which Arg degradation via arginase
(3.5.3.1), but not synthesis, is possible.
(c) Group of 3 species for which neither the degradation nor synthesis of arginine via
the urea cycle is possible (the three other hemipterans in this analysis).
(d) Group of 17 species sharing a complete (or almost complete) urea cycle.
Hemiptera are identified in red text and the milkweed-feeding monarch butterfly is in
blue text. Enzyme names corresponding to EC numbers: 1.5.1.2 = pyrroline-5-
carboxylate reductase; 1.14.13.39 = nitric-oxide synthase; 2.1.3.3 = ornithine
carbamoyltransferase; 2.6.1.13 = ornithine aminotransferase; 3.5.3.1 = arginase;
4.3.2.1 = argininosuccinate lyase; 6.3.4.5 = argininosuccinate synthase.
Analyses based on OGS v1.1.
1204
1205
1206 **TABLES (see above within relevant manuscript sections)**
1207
1208 **Table 1. *Oncopeltus fasciatus* genome metrics.**
1209
1210 **Table 2. Numbers of chemoreceptor genes/proteins per family in selected insect**
1211 **species.**
1212
1213 **Table 3. Hemipteran ArthropodaCyc database summaries.**
1214
1215 **Table 4. Hemipteran ArthropodaCyc annotations of metabolic genes.**
1216

1217 **DECLARATIONS**
1218
1219 **ADDITIONAL FILES**
1220 Additional file 1: Supplementary figures, tables, methods, and other text. (PDF)
1221 Additional file 2: Large supporting tables. (XLSX)
1222 Additional file 3: Chemoreceptor sequences in FASTA format. (TXT)
1223

ACKNOWLEDGEMENTS
We thank Dorith Rotenberg (Kansas State University, currently North Carolina State
University, USA) and Michael Sparks (Agricultural Research Service, United States
Department of Agriculture, USA) for generously making available the unpublished
genome assemblies of the fellow hemipteroid i5K species *Frankliniella occidentalis*
and *Halyomorpha halys*, respectively, for use in specific analyses presented here.
Similarly, we thank Hans Kelstrup and Lynn Riddiford (Janelia Farm Research
Campus, HHMI, USA) for sharing unpublished data on *Of-E75A* RNAi. We thank
George Coupland (Max Planck Institute for Plant Breeding Research, Cologne,
Germany) as well as Lisa Czaja, Kurt Steuber, and Bruno Huettel (Max Planck
Genome Centre Cologne, Germany) for conducting the PacBio sequencing and
providing support with data handling. We also thank Oliver Niehuis (Albert Ludwig
University, Freiburg, Germany) and Alexander Klassmann (University of Cologne,
Germany) for discussions on *k*-mer and gene structure analyses, respectively, Sarah
Kingan (University of Rochester, USA) for assistance with LGT phylogenies, as well
as Jeanne Wilbrandt (Zoologisches Forschungsmuseum Alexander Koenig, Bonn,
Germany) for comments on the manuscript.
1241

FUNDING
Funding for genome sequencing, assembly and automated annotation was provided by
the National Institutes of Health (NIH) grant U54 HG003273 (NHGRI) to RAG. The
i5K pilot project (https://www.hgsc.bcm.edu/arthropods) assisted in sequencing of the
Oncopeltus fasciatus genome. We also acknowledge funding for the project from
German Research Foundation (DFG) grants PA 2044/1-1 and SFB 680 project A12 to
KAP. Support for specific analyses was provided by the Swiss National Science
Foundation with grant 31003A_143936 to EMZ and PP00P3_170664 to RMW; the
European Research Council grant ERC-CoG #616346 to AK; DFG grant SFB 680
project A1 to SiR; the National Science Foundation with grant US NSF DEB1257053
to JHW; and by NIH grants 5R01GM080203 (NIGMS) and 5R01HG004483
(NHGRI) and by the Director, Office of Science, Office of Basic Energy Sciences,
U.S. Department of Energy, Contract No. DE-AC02-05CH11231 to MCMT.
1255

1256 **AVAILABILITY OF DATA AND MATERIALS**
1257 All sequence data are publically available at the NCBI, bioproject number
1258 PRJNA229125. In addition, gene models and a browser are available at the National
1259 Agricultural Library ([131-133], https://i5k.nal.usda.gov/Oncopeltus_fasciatus).
1260
AUTHORS' CONTRIBUTIONS
KAP and StR conceived the project. KAP managed and coordinated the project.
KAP and SK provided specimens for sequencing and performed DNA and RNA
extractions. StR, SD, SLL, HC, HVD, HD, YH, JQ, SCM, DSTH, KCW, DMM, and
RAG constructed libraries and performed sequencing. StR, SCM, and DSTH
performed the genome assembly and automated gene prediction. IMVJ, JSJ, and PJM

analyzed genome size. IMVJ, VK, PH, and KAP contributed to repetitive content analyses. AD, RR, JHW, KAP, and SK performed bacterial scaffold detection and LGT analyses. MCMT developed Apollo software. KAP, IMVJ, MCMT, CPC, C-YL, and MFP implemented Apollo-based manual curation. KAP, IMVJ, JBB, DE, YS, HMR, DA, CGCJ, BMIV, EJD, CSB, C-CC, Y-TC, ADC, AGC, AJJC, PKD, EMD, CGE, MF, NG, TH, Y-MH, ECJ, TEJ, JWJ, AK, ML, MRL, H-LL, YL, SRP, LP, MP, PNR, RRP, SiR, LS, MES, JS, ES, JNS, OT, LT, MVDZ, SV, and AJR participated in manual curation and contributed to the Supplemental Notes. IMVJ, KAP, DSTH, M-JMC, CPC, C-YL, and MFP performed curation quality control and generated the OGS. IMVJ, KAP, CJH, and JBB generated *de novo* transcriptomes and performed life history stage expression analyses. RMW, PI, KAP, and EMZ performed orthology and phylogenomic analyses. MTW, KAP, IMVJ, PH, and BMIV performed transcription factor analyses. EJD conducted analyses of DNA methylation. KAP, PH, and RJS contributed to comparative analyses of gene structure. DE conducted the RNAi experiments. SC, PB-P, GF, and NP generated and performed comparative analyses on the OncfaCyc database. KAP, IMVJ, JBB, DE, YS, SC, HMR, and MTW wrote the manuscript. KAP, IMVJ, JBB, DE, YS, SC, HMR, MFP, RMW, PI, MTW, StR, PJM, and AK edited the manuscript. IMVJ and KAP organized the Supplementary Materials. All authors approved the final manuscript.

COMPETING INTERESTS
The authors declare that they have no competing interests.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE
Not applicable.

AUTHOR DETAILS
[1] Institute for Zoology: Developmental Biology, University of Cologne, Zülpicher Str. 47b, 50674 Cologne, Germany
[2] School of Life Sciences, University of Warwick, Gibbet Hill Campus, Coventry CV4 7AL, UK
[3] Department of Biological Sciences, University of Cincinnati, Cincinnati, Ohio 45221, USA
[4] Department of Biochemistry and Cell Biology and Center for Developmental Genetics, Stony Brook University, Stony Brook, New York 11794, USA
[5] Department of Biological Sciences, Wellesley College, 106 Central St., Wellesley, Massachusetts 02481, USA
[6] Univ Lyon, INSA-Lyon, INRA, BF2I, UMR0203, F-69621, Villeurbanne, France
[7] Current address: LSTM, Laboratoire des Symbioses Tropicales et Méditerranéennes, INRA, IRD, CIRAD, SupAgro, University of Montpellier, Montpellier, France
[8] Department of Entomology, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA
[9] National Agricultural Library, Beltsville, Maryland 20705, USA
[10] Department of Genetic Medicine and Development and Swiss Institute of Bioinformatics, University of Geneva, Geneva 1211, Switzerland
[11] Current address: Department of Ecology and Evolution, University of Lausanne, Lausanne 1015, Switzerland
[12] Center for Autoimmune Genomics and Etiology, Division of Biomedical Informatics, and Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Department of Pediatrics, College of Medicine, University of Cincinnati, Cincinnati, Ohio 45229, USA
[13] Human Genome Sequencing Center, Department of Human and Molecular Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, Texas 77030, USA
[14] Current address: Department of Genome Sciences, University of Washington School of Medicine, Seattle, Washington 98195, USA

[15] Current address: Howard Hughes Medical Institute, University of Washington, Seattle, Washington 98195, USA

[16] Department of Biology, University of Rochester, Rochester, New York 14627, USA

[17] Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE Leiden, Netherlands

[18] Max Planck Institute for Chemical Ecology, Hans-Knöll Strasse 8, 07745 Jena, Germany

[19] Department of Biochemistry and Genomics Aotearoa, University of Otago, Dunedin 9054, New Zealand

[20] School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK

[21] Institut de Génomique Fonctionnelle de Lyon, Université de Lyon, Université Claude Bernard Lyon 1, CNRS UMR 5242, École Normale Supérieure de Lyon, 46 Allée d'Italie, 69364 Lyon, France

[22] Department of Ecology, Evolution and Behavior, The Alexander Silberman Institute of Life Sciences, The Hebrew University of Jerusalem, Edmond J. Safra Campus, Givat Ram 91904, Jerusalem, Israel

[23] Department of Entomology/Institute of Biotechnology, College of Bioresources and Agriculture, National Taiwan University, Taipei, Taiwan

[24] Current address: School of Life Sciences, Rochester Institute of Technology, Rochester, New York 14623, USA

[25] Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts 02138, USA

[26] Department of Molecular and Cellular Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts 02138, USA

[27] Department of Biological Sciences, Wayne State University, Detroit, Michigan 48202, USA

[28] Institute for Genetics, University of Cologne, Zülpicher Straße 47a, 50674 Cologne, Germany

[29] Department of Entomology, Texas A&M University, College Station, Texas 77843, USA

[30] CECAD, University of Cologne, Cologne, Germany

[31] Department of Entomology and Program in Molecular & Cell Biology, University of Maryland, College Park, Maryland 20742, USA

[32] Department of Entomology, University of Georgia, 120 Cedar St., Athens, Georgia 30602, USA

[33] Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA

[34] Department of Entomology, College of Agriculture, Food and Environment, University of Kentucky, Lexington, Kentucky 40546, USA

[35] Department of Biology, University of Hawai'i at Mānoa, Honolulu, Hawaii 96822, USA

[36] Current address: Department of Evolutionary Genetics, Max-Planck-Institut für Evolutionsbiologie, August-Thienemann-Straße 2, 24306 Plön, Germany

[37] Current address: Earthworks Institute, 185 Caroline Street, Rochester, New York 14620, USA

[38] Centro de Bioinvestigaciones, Universidad Nacional del Noroeste de Buenos Aires, Argentina

[39] Current address: Department of Biotechnology, Central university of Rajasthan (CURAJ), NH-8, Bandarsindri, Ajmer- 305801, India

[40] Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

[41] Current address: Department of Zoology, University of Cambridge, Cambridge CB2 3DT, UK

[42] Centro Regional de Estudios Genómicos, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, La Plata, Argentina

REFERENCES

1. Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, Seppey M, Loetscher A, Kriventseva EV: **OrthoDB v9.1: cataloging evolutionary and functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs.** *Nucleic Acids Res* 2017, **45**:D744-D749.

2. Huang DY, Bechly G, Nel P, Engel MS, Prokop J, Azar D, Cai CY, van de Kamp T, Staniczek AH, Garrouste R, et al: **New fossil insect order Permopsocida elucidates major radiation and evolution of suction feeding in hemimetabolous insects (Hexapoda: Acercaria).** *Sci Rep* 2016, **6**:23004.

3. Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG, et al: **Phylogenomics resolves the timing and pattern of insect evolution.** *Science* 2014, **346**:763-767.

4. Grimaldi D, Engel MS: *Evolution of the Insects.* Cambridge: Cambridge University Press; 2005.

5. Panfilio KA, Angelini DR: **By land, air, and sea: hemipteran diversity through the genomic lens.** *Curr Opin Insect Sci* 2018, **25**:in press, DOI: 10.1016/j.cois.2017.1012.1005.

6. The International Aphid Genomics Consortium: **Genome sequence of the pea aphid** *Acyrthosiphon pisum* PLoS Biol 2010, **8**:e1000313.

7. Mathers TC, Chen Y, Kaithakottil G, Legeai F, Mugford ST, Baa-Puyoulet P, Bretaudeau A, Clavijo B, Colella S, Collin O, et al: **Rapid transcriptional plasticity of duplicated gene clusters enables a clonally reproducing aphid to colonise diverse plant species.** *Genome Biol* 2017, **18**:27.

8. Wenger JA, Cassone BJ, Legeai F, Johnston JS, Bansal R, Yates AD, Coates BS, Pavinato VA, Michel A: **Whole genome sequence of the soybean aphid,** *Aphis glycines*. *Insect Biochem Mol Biol* 2017.

9. Sloan DB, Nakabachi A, Richards S, Qu J, Murali SC, Gibbs RA, Moran NA: **Parallel histories of horizontal gene transfer facilitated extreme reduction of endosymbiont genomes in sap-feeding insects.** *Mol Biol Evol* 2014, **31**:857-871.

10. Xue J, Zhou X, Zhang C-X, Yu L-L, Fan H-W, Wang Z, Xu H-J, Xi Y, Zhu Z-R, Zhou W-W, et al: **Genomes of the rice pest brown planthopper and its endosymbionts reveal complex complementary contributions for host adaptation.** *Genome Biol* 2014, **15**:521.

11. Mesquita RD, Vionette-Amaral RJ, Lowenberger C, Rivera-Pomar R, Monteiro FA, Minx P, Spieth J, Carvalho AB, Panzera F, Lawson D, et al: **Genome of** *Rhodnius prolixus***, an insect vector of Chagas disease, reveals unique adaptations to hematophagy and parasite infection.** *Proc Natl Acad Sci USA* 2015, **112**:14936-14941.

12. Benoit JB, Adelman ZN, Reinhardt K, Dolan A, Poelchau M, Jennings EC, Szuter EM, Hagan RW, Gujar H, Shukla JN, et al: **Unique features of a global human ectoparasite identified through sequencing of the bed bug genome.** *Nat Commun* 2016, **7**:10165.

13. Rosenfeld JA, Reeves D, Brugler MR, Narechania A, Simon S, Durrett R, Foox J, Shianna K, Schatz MC, Gandara J, et al: **Genome assembly and geospatial phylogenomics of the bed bug** *Cimex lectularius*. *Nat Commun* 2016, **7**:10164.

14. Sparks ME, Shelby KS, Kuhar D, Gundersen-Rindal DE: **Transcriptome of the invasive brown marmorated stink bug,** *Halyomorpha halys* **(Stal) (Heteroptera: Pentatomidae).** *PLoS One* 2014, **9**:e111646.

15. Ioannidis P, Lu Y, Kumar N, Creasy T, Daugherty S, Chibucos MC, Orvis J, Shetty A, Ott S, Flowers M, et al: **Rapid transcriptome sequencing of an invasive pest, the brown marmorated stink bug,** *Halyomorpha halys*. *BMC Genomics* 2014, **15**:738.

16. Li H, Leavengood JM, Jr., Chapman EG, Burkhardt D, Song F, Jiang P, Liu J, Zhou X, Cai W: **Mitochondrial phylogenomics of Hemiptera reveals adaptive innovations driving the diversification of true bugs.** *Proc Biol Sci* 2017, **284**.

17. Wilson ACC, Ashton PD, Charles H, Colella S, Febvay G, Jander G, Kushlan PF, Macdonald SJ, Schwartz JF, Thomas GH, Douglas AE: **Genomic insight into the amino acid relations of the pea aphid,** *Acyrthosiphon pisum***, with its symbiotic bacterium** *Buchnera aphidicola*. *Insect Mol Biol* 2010, **19 Suppl 2**:249-258.

18. Eichler S, Schaub GA: **Development of symbionts in triatomine bugs and the effects of infections with trypanosomatids.** *Exp Parasitol* 2002, **100**:17-27.

19. Matsuura Y, Kikuchi Y, Hosokawa T, Koga R, Meng X-Y, Kamagata Y, Nikoh N, Fukatsu T: **Evolution of symbiotic organs and endosymbionts in lygaeid stinkbugs.** *The ISME Journal* 2012, **6**:397–409.

20.   Berenbaum MR, Miliczky E: **Mantids and milkweed bugs - efficacy of aposematic**
coloration against invertebrate predators. *American Midland Naturalist* 1984, **111:**64-68.
21.   Burdfield-Steel ER, Shuker DM: **The evolutionary ecology of the Lygaeidae.** *Ecol Evol*
2014, **4:**2278-2301.
22.   Lawrence PA: **Mitosis and the cell cycle in the metamorphic moult of the milkweed bug**
Oncopeltus fasciatus; a radioautographic study. *J Cell Sci* 1968, **3:**391-404.
23.   Chipman AD: *Oncopeltus fasciatus* **as an evo-devo research organism.** *Genesis* 2017, **55**.
24.   Panfilio KA: **Late extraembryonic development and its** *zen-RNAi*-**induced failure in the**
milkweed bug *Oncopeltus fasciatus*. *Dev Biol* 2009, **333:**297-311.
25.   Panfilio KA, Roth S: **Epithelial reorganization events during late extraembryonic**
development in a hemimetabolous insect. *Dev Biol* 2010, **340:**100-115.
26.   Sharma AI, Yanes KO, Jin L, Garvey SL, Taha SM, Suzuki Y: **The phenotypic plasticity of**
developmental modules. *Evodevo* 2016, **7:**15.
27.   Hughes CL, Kaufman TC: **RNAi analysis of** *Deformed, proboscipedia* **and** *Sex combs*
reduced **in the milkweed bug** *Oncopeltus fasciatus*: **novel roles for Hox genes in the**
hemipteran head. *Development* 2000, **127:**3683-3694.
28.   Wolfe SL, John B: **The organization and ultrastructure of male meiotic chromosomes in**
Oncopeltus fasciatus. *Chromosoma* 1965, **17:**85-103.
29.   Messthaler H, Traut W: **Phases of Sex Chromosome Inactivation in** *Oncopeltus fasciatus*
and *Pyrrhocoris apterus* **(Insecta, Heteroptera).** *Caryologia* 1975, **28:**501-510.
30.   McKenna DD, Scully ED, Pauchet Y, Hoover K, Kirsch R, Geib SM, Mitchell RF,
Waterhouse RM, Ahn SJ, Arsala D, et al: **Genome of the Asian longhorned beetle**
(*Anoplophora glabripennis***), a globally significant invasive species, reveals key functional**
and evolutionary innovations at the beetle-plant interface. *Genome Biol* 2016, **17:**227.
31.   Simpson JT: **Exploring genome characteristics and sequence quality without a reference.**
Bioinformatics 2014, **30:**1228-1235.
32.   Hanrahan SJ, Johnston JS: **New genome size estimates of 134 species of arthropods.**
Chromosome Res 2011, **19:**809-823.
33.   Hughes CL, Kaufman TC: **RNAi analysis of** *Deformed, proboscipedia* **and** *Sex combs*
reduced **in the milkweed bug** *Oncopeltus fasciatus*: **novel roles for Hox genes in the**
Hemipteran head. *Development* 2000, **127:**3683-3694.
34.   Panfilio KA, Liu PZ, Akam M, Kaufman TC: *Oncopeltus fasciatus zen* **is essential for**
serosal tissue function in katatrepsis. *Dev Biol* 2006, **292:**226-243.
35.   Tian X, Xie Q, Li M, Gao C, Cui Y, Xi L, Bu W: **Phylogeny of pentatomomorphan bugs**
(Hemiptera-Heteroptera:Pentatomomorpha) based on six Hox gene fragments. *Zootaxa*
2011, **2888:**57-68.
36.   Ewen-Campen B, Shaner N, Panfilio KA, Suzuki Y, Roth S, Extavour CG: **The maternal**
and early embryonic transcriptome of the milkweed bug *Oncopeltus fasciatus*. *BMC*
Genomics 2011, **12:**61.
37.   Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P: **Parallel molecular evolution**
in an herbivore community. *Science* 2012, **337:**1634-1637.
38.   Robertson HM: **The insect chemoreceptor superfamily in Drosophila pseudoobscura:**
Molecular evolution of ecologically-relevant genes over 25 million years *J Insect Sci* 2009,
9:18.
39.   Robertson HM: **Taste: Independent origins of chemoreception coding systems?** *Curr Biol*
2001, **11:**R560-R562.
40.   Jazwinska A, Rushlow C, Roth S: **The role of** *brinker* **in mediating the graded response to**
Dpp in early *Drosophila* **embryos.** *Development* 1999, **126:**3323-3334.
41.   Togawa T, Dunn WA, Emmons AC, Nagao J, Willis JH: **Developmental expression**
patterns of cuticular protein genes with the R&R Consensus from *Anopheles gambiae.*
Insect Biochem Mol Biol 2008, **38:**508-519.
42.   Karr TL: **Fruit flies and the sperm proteome.** *Hum Mol Genet* 2007, **16 Spec No. 2:**R124-
133.
43.   Waterhouse RM, Seppey M, Simao FA, Manni M, Ioannidis P, Klioutchnikov G, Kriventseva
EV, Zdobnov EM: **BUSCO applications from quality assessments to gene prediction and**
phylogenomics. *Mol Biol Evol* 2017.
44.   Shigenobu S, Bickel RD, Brisson JA, Butts T, Chang CC, Christiaens O, Davis GK, Duncan
EJ, Ferrier DE, Iga M, et al: **Comprehensive survey of developmental genes in the pea**
aphid, *Acyrthosiphon pisum*: **frequent lineage-specific duplications and losses of**
developmental genes. *Insect Mol Biol* 2010, **19 Suppl 2:**47-62.

45.   Kriventseva EV, Tegenfeldt F, Petty TJ, Waterhouse RM, Simão FA, Pozdnyakov IA,
1488   Ioannidis P, Zdobnov EM: **OrthoDB v8: update of the hierarchical catalog of orthologs**
1489   **and the underlying free software.** *Nucl Acids Res* 2015, **43:**D250-D256.

1490   46.   Bansal R, Michel AP: **Core RNAi Machinery and *Sid1*, a component for systemic RNAi,**
1491   **in the hemipteran insect, *Aphis glycines*.** *Int J Mol Sci* 2013, **14:**3786-3801.

1492   47.   Bao R, Fischer T, Bolognesi R, Brown SJ, Friedrich M: **Parallel duplication and partial**
1493   **subfunctionalization of beta-catenin/armadillo during insect evolution.** *Mol Biol Evol*
1494   2012, **29:**647-662.

1495   48.   Sachs L, Chen YT, Drechsler A, Lynch JA, Panfilio KA, Lassig M, Berg J, Roth S: **Dynamic**
1496   **BMP signaling polarized by Toll patterns the dorsoventral axis in a hemimetabolous**
1497   **insect.** *eLife* 2015, **4:**e05502.

1498   49.   Armisen D, Refki PN, Crumiere AJ, Viala S, Toubiana W, Khila A: **Predator strike shapes**
1499   **antipredator phenotype through new genetic interactions in water striders.** *Nat Commun*
1500   2015, **6:**8153.

1501   50.   Konopova B, Smykal V, Jindra M: **Common and distinct roles of juvenile hormone**
1502   **signaling genes in metamorphosis of holometabolous and hemimetabolous insects.** *PLoS*
1503   *One* 2011, **6:**e28728.

1504   51.   Vellichirammal NN, Gupta P, Hall TA, Brisson JA: **Ecdysone signaling underlies the pea**
1505   **aphid transgenerational wing polyphenism.** *Proc Natl Acad Sci U S A* 2017, **114:**1419-
1506   1423.

1507   52.   Wulff JP, Sierra I, Sterkel M, Holtof M, Van Wielendaele P, Francini F, Broeck JV, Ons S:
1508   **Orcokinin neuropeptides regulate ecdysis in the hemimetabolous insect Rhodnius**
1509   **prolixus.** *Insect Biochem Mol Biol* 2017, **81:**91-102.

1510   53.   Chiu TL, Wen Z, Rupasinghe SG, Schuler MA: **Comparative molecular modeling of**
1511   **Anopheles gambiae CYP6Z1, a mosquito P450 capable of metabolizing DDT.** *Proc Natl*
1512   *Acad Sci U S A* 2008, **105:**8855-8860.

1513   54.   Gong Y, Li T, Feng Y, Liu N: **The function of two P450s, CYP9M10 and CYP6AA7, in**
1514   **the permethrin resistance of Culex quinquefasciatus.** *Sci Rep* 2017, **7:**587.

1515   55.   Liu PZ, Kaufman TC: ***hunchback* is required for suppression of abdominal identity, and**
1516   **for proper germband growth and segmentation in the intermediate germband insect**
1517   ***Oncopeltus fasciatus*.** *Development* 2004, **131:**1515-1527.

1518   56.   Schaeper ND, Pechmann M, Damen WGM, Prpic N-M, Wimmer EA: **Evolutionary**
1519   **plasticity of *collier* function in head development of diverse arthropods.** *Dev Biol* 2010,
1520   **344:**363-376.

1521   57.   Aspiras AC, Smith FW, Angelini DR: **Sex-specific gene interactions in the patterning of**
1522   **insect genitalia.** *Dev Biol* 2011, **360:**369-380.

1523   58.   Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi HS,
1524   Lambert SA, Mann I, Cook K, et al: **Determination and inference of eukaryotic**
1525   **transcription factor sequence specificity.** *Cell* 2014, **158:**1431-1443.

1526   59.   Peel AD, Telford MJ, Akam M: **The evolution of hexapod engrailed-family genes:**
1527   **evidence for conservation and concerted evolution.** *Proc Biol Sci* 2006, **273:**1733-1742.

1528   60.   Ben-David J, Chipman AD: **Mutual regulatory interactions of the trunk gap genes during**
1529   **blastoderm patterning in the hemipteran *Oncopeltus fasciatus*.** *Dev Biol* 2010, **346:**140-
1530   149.

1531   61.   Erezyilmaz DF, Kelstrup HC, Riddiford LM: **The nuclear receptor E75A has a novel pair-**
1532   **rule-like function in patterning the milkweed bug, *Oncopeltus fasciatus*.** *Dev Biol* 2009,
1533   **334:**300-310.

1534   62.   Liu PZ, Kaufman TC: ***even-skipped* is not a pair-rule gene but has segmental and gap-like**
1535   **functions in *Oncopeltus fasciatus*, an intermediate germband insect.** *Development* 2005,
1536   **132:**2081-2092.

1537   63.   Weisbrod A, Cohen M, Chipman AD: **Evolution of the insect terminal patterning system--**
1538   **insights from the milkweed bug, *Oncopeltus fasciatus*.** *Dev Biol* 2013, **380:**125-131.

1539   64.   Albertin CB, Simakov O, Mitros T, Wang ZY, Pungor JR, Edsinger-Gonzales E, Brenner S,
1540   Ragsdale CW, Rokhsar DS: **The octopus genome and the evolution of cephalopod neural**
1541   **and morphological novelties.** *Nature* 2015, **524:**220-224.

1542   65.   Crooks GE, Hon G, Chandonia J-M, Brenner SE: **WebLogo: A sequence logo generator.**
1543   *Genome Res* 2004, **14:**1188-1190.

1544   66.   Najafabadi HS, Mnaimneh S, Schmitges FW, Garton M, Lam KN, Yang A, Albu M,
1545   Weirauch MT, Radovani E, Kim PM, et al: **C2H2 zinc finger proteins greatly expand the**
1546   **human regulatory lexicon.** *Nat Biotechnol* 2015, **33:**555-562.

67. Emerson RO, Thomas JH: **Adaptive evolution in zinc finger transcription factors.** *PLoS Genet* 2009, **5:**e1000325.

68. Thomas JH, Schneider S: **Coevolution of retroelements and tandem zinc finger genes.** *Genome Res* 2011, **21:**1800-1812.

69. Garcia-Perez JL, Widmann TJ, Adams IR: **The impact of transposable elements on mammalian development.** *Development* 2016, **143:**4101-4114.

70. Liu PZ, Kaufman TC: *Krüppel* **is a gap gene in the intermediate insect** *Oncopeltus fasciatus* **and is required for development of both blastoderm and germband-derived segments.** *Development* 2004, **131:**4567-4579.

71. Heger P, Marin B, Bartkuhn M, Schierenberg E, Wiehe T: **The chromatin insulator CTCF and the emergence of metazoan diversity.** *Proc Natl Acad Sci USA* 2012, **109:**17507-17512.

72. Liu H, Chang L-H, Sun Y, Lu X, Stubbs L: **Deep vertebrate roots for mammalian zinc finger transcription factor subfamilies.** *Genome Biol Evol* 2014, **6:**510-525.

73. Imbeault M, Helleboid P-Y, Trono D: **KRAB zinc-finger proteins contribute to the evolution of gene regulatory networks.** *Nature* 2017, **543:**550-554.

74. Csuros M, Rogozin IB, Koonin EV: **A detailed history of intron-rich eukaryotic ancestors inferred from a global survey of 100 complete genomes.** *PLoS Comput Biol* 2011, **7:**e1002150.

75. Hoy MA, Waterhouse RM, Wu K, Estep AS, Ioannidis P, Palmer WJ, Pomerantz AF, Simao FA, Thomas J, Jiggins FM, et al: **Genome sequencing of the phytoseiid predatory mite** *Metaseiulus occidentalis* **reveals completely atomized Hox genes and superdynamic intron evolution.** *Genome Biol Evol* 2016, **8:**1762-1775.

76. Seibt KM, Wenke T, Muders K, Truberg B, Schmidt T: **Short interspersed nuclear elements (SINEs) are abundant in Solanaceae and have a family-specific impact on gene structure and genome organization.** *Plant J* 2016, **86:**268-285.

77. Huff JT, Zilberman D, Roy SW: **Mechanism for DNA transposons to generate introns on genomic scales.** *Nature* 2016, **538:**533-536.

78. Wheeler D, Redding AJ, Werren JH: **Characterization of an ancient lepidopteran lateral gene transfer.** *PLoS One* 2012, **8:**e59262.

79. Da Lage JL, Binder M, Hua-Van A, Janecek S, Casane D: **Gene make-up: rapid and massive intron gains after horizontal transfer of a bacterial alpha-amylase gene to Basidiomycetes.** *BMC Evol Biol* 2013, **13:**40.

80. Lee DH, Short BD, Joseph SV, Bergh JC, Leskey TC: **Review of the biology, ecology, and management of** *Halyomorpha halys* **(Hemiptera: Pentatomidae) in China, Japan, and the Republic of Korea.** *Environ Entomol* 2013, **42:**627-641.

81. Lawrence PA: **Cellular differentiation and pattern formation during metamorphosis of the milkweed bug Oncopeltus.** *Dev Biol* 1969, **19:**12-40.

82. Riddiford LM: **Prevention of Metamorphosis by Exposure of Insect Eggs to Juvenile Hormone Analogs.** *Science* 1970, **167:**287-&.

83. Willis JH, Lawrence PA: **Deferred Action of Juvenile Hormone.** *Nature* 1970, **225:**81-83.

84. Masner P, Bowers WS, Kalin M, Muhle T: **Effect of precocene II on the endocrine regulation of development and reproduction in the bug,** *Oncopeltus fasciatus.* *Gen Comp Endocrinol* 1979, **37:**156-166.

85. Rewitz K, O'Connor M, Gilbert L: **Molecular evolution of the insect Halloween family of cytochrome P450s: phylogeny, gene organization and functional conservation.** *Insect Biochem Mol Biol* 2007, **37:**741-753.

86. Huet F, Ruiz C, Richards G: **Sequential gene activation by ecdysone in Drosophila melanogaster: the hierarchical equivalence of early and early late genes.** *Development* 1995, **121:**1195-1204.

87. Bialecki M, Shilton A, Fichtenberg C, Segraves WA, Thummel CS: **Loss of the ecdysteroid-inducible E75A orphan nuclear receptor uncouples molting from metamorphosis in Drosophila.** *Dev Cell* 2002, **3:**209-220.

88. Charles JP, Iwema T, Epa VC, Takaki K, Rynes J, Jindra M: **Ligand-binding properties of a juvenile hormone receptor, Methoprene-tolerant.** *Proceedings of the National Academy of Sciences of the United States of America* 2011, **108:**21128-21133.

89. Minakuchi C, Zhou X, Riddiford L: **Kruppel homolog 1 (Kr-h1) mediates juvenile hormone action during metamorphosis of Drosophila melanogaster.** *Mech Dev* 2008, **125:**91-105.

90. Minakuchi C, Namiki T, Shinoda T: **Kruppel homolog 1, an early juvenile hormone-**
response gene downstream of Methoprene-tolerant, mediates its anti-metamorphic
action in the red flour beetle Tribolium castaneum. *Dev Biol* 2009, **352:**341-350.
91. DiBello PR, Withers DA, Bayer CA, Fristrom JW, Guild GM: **The Drosophila Broad-**
Complex encodes a family of related proteins containing zinc fingers. *Genetics* 1991,
129:385-397.
92. Karim F, Guild G, Thummel C: **The Drosophila Broad-Complex plays a key role in**
controlling ecdysone-regulated gene expression at the onset of metamorphosis.
Development 1993, **118:**977-988.
93. Erezyilmaz DF, Riddiford LM, Truman JW: **The pupal specifier broad directs progressive**
morphogenesis in a direct-developing insect. *Proceedings of the National Academy of*
Sciences of the United States of America 2006, **103:**6925-6930.
94. Arakane Y, Hogenkamp DG, Zhu YC, Kramer KJ, Specht CA, Beeman RW, Kanost MR,
Muthukrishnan S: **Characterization of two chitin synthase genes of the red flour beetle,**
***Tribolium castaneum*, and alternate exon usage in one of the genes during development.**
Insect Biochem Mol Biol 2004, **34:**291-304.
95. True JR: **Insect melanism: the molecules matter.** *Trends in Ecology & Evolution* 2003,
18:640-647.
96. Zhan SA, Guo QH, Li MH, Li MW, Li JY, Miao XX, Huang YP: **Disruption of an N-**
acetyltransferase gene in the silkworm reveals a novel role in pigmentation. *Development*
2010, **137:**4083-4090.
97. Liu J, Lemonds TR, Popadic A: **The genetic control of aposematic black pigmentation in**
hemimetabolous insects: insights from Oncopeltus fasciatus. *Evolution & Development*
2014, **16:**270-277.
98. Liu J, Lemonds TR, Marden JH, Popadic A: **A Pathway Analysis of Melanin Patterning in**
a Hemimetabolous Insect. *Genetics* 2016, **203:**403-413.
99. Lawrence PA: **Some new mutants of large milkweed bug *Oncopeltus fasciatus* Dall.**
Genetical Research 1970, **15:**347-350.
100. Morgan ED: *Biosynthesis in Insects: Advanced Edition.* London: Royal Society of Chemistry,;
2010.
101. McLean JR, Krishnakumar S, O'Donnell JM: **Multiple mRNAs from the Punch locus of**
Drosophila melanogaster encode isoforms of GTP cyclohydrolase I with distinct N-
terminal domains. *J Biol Chem* 1993, **268:**27191-27197.
102. Wiederrecht GJ, Paton DR, Brown GM: **Enzymatic Conversion of Dihydroneopterin**
Triphosphate to the Pyrimidodiazepine Intermediate Involved in the Biosynthesis of the
Drosopterins in Drosophila-Melanogaster. *Journal of Biological Chemistry* 1984,
259:2195-2200.
103. Newcombe D, Blount JD, Mitchell C, Moore AJ: **Chemical egg defence in the large**
milkweed bug, Oncopeltus fasciatus, derives from maternal but not paternal diet.
Entomologia Experimentalis et Applicata 2013, **149:**197-205.
104. Zhan S, Merlin C, Boore JL, Reppert SM: **The monarch butterfly genome yields insights**
into long-distance migration. *Cell* 2011, **147:**1171-1185.
105. Smadja C, Shi P, Butlin RK, Robertson HM: **Large gene family expansions and adaptive**
evolution for odorant and gustatory receptors in the pea aphid, Acyrthosiphon pisum.
Mol Biol Evol 2009, **26:**2073-2086.
106. Terrapon N, Li C, Robertson HM, Ji L, Meng X, Booth W, Chen Z, Childers CP, Glastad KM,
Gokhale K, et al: **Molecular traces of alternative social organization in a termite genome.**
Nat Commun 2014, **5:**3636.
107. Croset V, Rytz R, Cummins SF, Budd A, Brawand D, Kaessmann H, Gibson TJ, Benton R:
Ancient protostome origin of chemosensory ionotropic glutamate receptors and the
evolution of insect taste and olfaction. *PLoS Genet* 2010, **6:**e1001064.
108. Kirkness EF, Haas BJ, Sun W, Braig HR, Perotti MA, Clark JM, Lee SH, Robertson HM,
Kennedy RC, Elhaik E, et al: **Genome sequences of the human body louse and its primary**
endosymbiont provide insights into the permanent parasitic lifestyle. *Proc Natl Acad Sci*
U S A 2010, **107:**12168-12173.
109. Robertson HM, Warr CG, Carlson JR: **Molecular evolution of the insect chemoreceptor**
gene superfamily in Drosophila melanogaster. *Proc Natl Acad Sci U S A* 2003, **100 Suppl**
2:14537-14542.
110. Joseph RM, Carlson JR: **Drosophila Chemoreceptors: A Molecular Interface Between the**
Chemical World and the Brain. *Trends Genet* 2015, **31:**683-695.

111. Benton R: **Multigene Family Evolution: Perspectives from Insect Chemoreceptors.** *Trends Ecol Evol* 2015, **30:**590-600.
112. Rytz R, Croset V, Benton R: **Ionotropic receptors (IRs): chemosensory ionotropic glutamate receptors in Drosophila and beyond.** *Insect Biochem Mol Biol* 2013, **43:**888-897.
113. Xu W, Papanicolaou A, Zhang HJ, Anderson A: **Expansion of a bitter taste receptor family in a polyphagous insect herbivore.** *Sci Rep* 2016, **6:**23666.
114. Feir D: ***Oncopeltus fasciatus:* A research animal.** *Annu Rev Entomol* 1974, **19:**81-96.
115. Vellozo AF, Véron AS, Baa-Puyoulet P, Huerta-Cepas J, Cottret L, Febvay G, Calevro F, Rahbe Y, Douglas AE, Gabaldón T, et al: **CycADS: an annotation database system to ease the development and update of BioCyc databases.** *Database* 2011, **2011:**bar008-bar008.
116. Baa-Puyoulet P, Parisot N, Febvay G, Huerta-Cepas J, Vellozo AF, Gabaldón T, Calevro F, Charles H, Colella S: **ArthropodaCyc: a CycADS powered collection of BioCyc databases to analyse and compare metabolism of arthropods.** *Database (Oxford)* 2016, **pii:**baw081.
117. Hojilla-Evangelista MP, Evangelista RL: **Characterization of milkweed (Asclepias spp.) seed proteins.** *Industrial crops and ...* 2009.
118. Dean CAE, Teets NM, Koštál V, Šimek P, Denlinger DL: **Enhanced stress responses and metabolic adjustments linked to diapause and onset of migration in the large milkweed bug *Oncopeltus fasciatus*.** *Physiol Entomol* 2016, **DOI: 10.1111/phen.12140**.
119. Rabatel A, Febvay G, Gaget K, Duport G, Baa-Puyoulet P, Sapountzis P, Bendridi N, Rey M, Rahbé Y, Charles H, et al: **Tyrosine pathway regulation is host-mediated in the pea aphid symbiosis during late embryonic and early larval development.** *BMC Genomics* 2013, **14:**235.
120. Dobler S, Petschenka G, Wagschal V, Flacht L: **Convergent adaptive evolution – how insects master the challenge of cardiac glycoside-containing host plants.** *Entomologia Experimentalis et Applicata* 2015, **157:**30-39.
121. Grau-Bove X, Ruiz-Trillo I, Irimia M: **Origin of exon skipping-rich transcriptomes in animals driven by evolution of gene architecture.** *Genome Biol* 2018, **19:**135.
122. Niehuis O, Gibson JD, Rosenberg MS, Pannebakker BA, Koevoets T, Judson AK, Desjardins CA, Kennedy K, Duggan D, Beukeboom LW, et al: **Recombination and its impact on the genome of the haplodiploid parasitoid wasp Nasonia.** *PLoS One* 2010, **5:**e8597.
123. Ferrero A, Torreblanca A, Garcera MD: **Assessment of the effects of orally administered ferrous sulfate on Oncopeltus fasciatus (Heteroptera: Lygaeidae).** *Environ Sci Pollut Res Int* 2017, **24:**8551-8561.
124. Hare EE, Johnston JS: **Genome size determination using flow cytometry of propidium iodide-stained nuclei.** *Methods Mol Biol* 2011, **772:**3-12.
125. Marcais G, Kingsford C: **A fast, lock-free approach for efficient parallel counting of occurrences of k-mers.** *Bioinformatics* 2011, **27:**764-770.
126. Bushnell B: **BBMap short read aligner.** 2016.
127. Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, et al: **High-quality draft assemblies of mammalian genomes from massively parallel sequence data.** *Proc Natl Acad Sci U S A* 2011, **108:**1513-1518.
128. Holt C, Yandell M: **MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects.** *BMC Bioinformatics* 2011, **12:**491.
129. Lee E, Helt G, Reese J, Munoz-Torres M, Childers C, Buels R, Stein L, Holmes I, Elsik C, Lewis S: **Web Apollo: a web-based genomic annotation editing platform.** *Genome Biology* 2013, **14**.
130. Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee CY, Lin H, Lin JW, Hackett K: **The i5k Workspace@NAL--enabling genomic data access, visualization and curation of arthropod genomes.** *Nucleic Acids Res* 2015, **43:**D714-719.
131. Hughes DST, Koelzer S, Panfilio KA, Richards S: ***Oncopeltus fasciatus* genome annotations v0.5.3.** *Ag Data Commons (Database)* 2015:http://dx.doi.org/10.15482/USDA.ADC/1173237.
132. Murali SC, The i5k genome assembly team (29 additional authors), Han Y, Richards S, Worley K, Muzny D, Gibbs R, Koelzer S, Panfilio KA: ***Oncopeltus fasciatus* genome assembly 1.0.** *Ag Data Commons (Database)* 2015:http://dx.doi.org/10.15482/USDA.ADC/1173238.
133. Vargas Jentzsch IM, Hughes DST, Poelchau M, Robertson HM, Benoit JB, Rosendale AJ, Armisén D, Duncan EJ, Vreede BMI, Jacobs CGC, et al: ***Oncopeltus fasciatus* Official Gene Set OGS_v1.1 for genome assembly Oncopeltus fasciatus v1.0.** *Ag Data Commons (Database)* 2015:http://dx.doi.org/10.15482/USDA.ADC/1173142.

134.   **RepeatModeler Open-1.0.8** [http://www.repeatmasker.org]

135.   Bao Z, Eddy SR: **Automated de novo identification of repeat sequence families in**
sequenced genomes. *Genome Res* 2002, **12:**1269-1276.

136.   Price AL, Jones NC, Pevzner PA: **De novo identification of repeat families in large**
genomes. *Bioinformatics* 2005, **21 Suppl 1:**i351-358.

137.   Benson G: **Tandem repeats finder: a program to analyze DNA sequences.** *Nucleic Acids*
Res 1999, **27:**573-580.

138.   **RepeatMasker Open-4.0.** [http://www.repeatmasker.org]

139.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods* 2012,
9:357-359.

140.   Goff S, Vaughn M, McKay S, Lyons E, Stapleton A, Gessler D, Matasci N, Wang L, Hanlon
M, Lenards A, et al: **The iPlant Collaborative: Cyberinfrastructure for Plant Biology.**
Frontiers in plant science 2011, **2**.

141.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, et al: **Full-length transcriptome assembly from RNA-Seq data**
without a reference genome. *Nat Biotechnol* 2011, **29:**644-652.

142.   Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles
D, Li B, Lieber M, et al: **De novo transcript sequence reconstruction from RNA-seq using**
the Trinity platform for reference generation and analysis. *Nat Protoc* 2013, **8:**1494-1512.

143.   Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data with or**
without a reference genome. *BMC Bioinformatics* 2011, **12:**323.

144.   Schoville SD, Chen YH, Andersson MN, Benoit JB, Bhandari A, Bowsher JH, Brevik K,
Cappelle K, Chen MM, Childers AK, et al: **A model species for agricultural pest genomics:**
the genome of the Colorado potato beetle, *Leptinotarsa decemlineata* **(Coleoptera:**
Chrysomelidae). *Sci Rep* 2018, **8:**1931.

145.   Scolari F, Benoit JB, Michalkova V, Aksoy E, Takac P, Abd-Alla AM, Malacrida AR, Aksoy
S, Attardo GM: **The spermatophore in** *Glossina morsitans morsitans***: Insights into male**
contributions to reproduction. *Sci Rep* 2016, **6:**20334.

146.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM: **BUSCO: assessing**
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*
2015, **31:**3210-3212.

147.   Finn RD, Mistry J, Tate J, Coggill P, Heger A, Pollington JE, Gavin OL, Gunasekaran P,
Ceric G, Forslund K, et al: **The Pfam protein families database.** *Nucleic Acids Res* 2010,
38:D211-222.

148.   Weirauch MT, Hughes TR: **A catalogue of eukaryotic transcription factor types, their**
evolutionary origin, and species distribution. *Subcell Biochem* 2011, **52:**25-73.

149.   Eddy SR: **A new generation of homology search tools based on probabilistic inference.**
Genome Inform 2009, **23:**205-211.

150.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Soding J, et al: **Fast, scalable generation of high-quality protein multiple**
sequence alignments using Clustal Omega. *Mol Syst Biol* 2011, **7:**539.

151.   Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M: **KAAS: an automatic genome**
annotation and pathway reconstruction server. *Nucleic Acids Research* 2007, **35:**W182-
185.

152.   Claudel-Renard C, Chevalet C, Faraut T, Kahn D: **Enzyme-specific profiles for genome**
annotation: PRIAM. *Nucleic Acids Research* 2003, **31:**6633-6639.

153.   Conesa A, Götz S: **Blast2GO: A Comprehensive Suite for Functional Analysis in Plant**
Genomics. *International journal of plant genomics* 2008, **2008:**619832.

154.   Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M: **Blast2GO: a universal**
tool for annotation, visualization and analysis in functional genomics research.
Bioinformatics (Oxford, England) 2005, **21:**3674-3676.

155.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
Mitchell A, Nuka G, et al: **InterProScan 5: genome-scale protein function classification.**
Bioinformatics 2014, **30:**1236-1240.

156.   Karp PD, Ouzounis CA, Moore-Kochlacs C, Goldovsky L, Kaipa P, Ahrén D, Tsoka S,
Darzentas N, Kunin V, López-Bigas N: **Expansion of the BioCyc collection of**
pathway/genome databases to 160 genomes. *Nucleic Acids Research* 2005, **33:**6083-6089.

157.   Karp PD, Paley SM, Krummenacker M, Latendresse M, Dale JM, Lee TJ, Kaipa P, Gilham F,
Spaulding A, Popescu L, et al: **Pathway Tools version 13.0: integrated software for**

pathway/genome informatics and systems biology. *Briefings in Bioinformatics* 2010,
11:40-79.
158.   Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S,
Lefort V, Lescot M, et al: **Phylogeny.fr: robust phylogenetic analysis for the non-**
specialist. *Nucleic Acids Res* 2008, **36:**W465-469.
159.   Wang X, Fang X, Yang P, Jiang X, Jiang F, Zhao D, Li B, Cui F, Wei J, Ma C, et al: **The**
locust genome provides insight into swarm formation and long-distance flight. *Nat*
Commun 2014, **5:**2957.
160.   The International Silkworm Genome Consortium: **The genome of a lepidopteran model**
insect, the silkworm Bombyx mori. *Insect Biochem Mol Biol* 2008, **38:**1036-1045.
161.   Elsik CG, Worley KC, Bennett AK, Beye M, Camara F, Childers CP, de Graaf DC, Debyser
G, Deng J, Devreese B, et al: **Finding the missing honey bee genes: lessons learned from a**
genome upgrade. *BMC Genomics* 2014, **15:**86.
162.   Honeybee Genome Sequencing Consortium: **Insights into social insects from the genome of**
the honeybee Apis mellifera. *Nature* 2006, **443:**931-949.
163.   Richards S, Gibbs RA, Weinstock GM, Brown SJ, Denell R, Beeman RW, Gibbs R, Bucher
G, Friedrich M, Grimmelikhuijzen CJ, et al: **The genome of the model beetle and pest**
Tribolium castaneum. *Nature* 2008, **452:**949-955.
164.   Chen XG, Jiang X, Gu J, Xu M, Wu Y, Deng Y, Zhang C, Bonizzoni M, Dermauw W, Vontas
J, et al: **Genome sequence of the Asian Tiger mosquito, Aedes albopictus, reveals insights**
into its biology, genetics, and evolution. *Proc Natl Acad Sci U S A* 2015, **112:**E5907-5915.
165.   Armisen D, Rajakumar R, Friedrich M, Benoit JB, Robertson HM, Panfilio KA, Ahn S-J,
Poelchau MF, Chao H, Dinh H, et al: **The genome of the water strider** *Gerris buenoi*
reveals expansions of gene repertoires associated with adaptations to life on the water.
BMC Genomics in press**:**acceptance e-mail 12 Oct. 2018.
166.   Papanicolaou A, Schetelig MF, Arensburger P, Atkinson PW, Benoit JB, Bourtzis K,
Castañera P, Cavanaugh JP, Chao H, Childers C, et al: **The whole genome sequence of the**
Mediterranean fruit fly, *Ceratitis capitata* **(Wiedemann), reveals insights into the biology**
and adaptive evolution of a highly invasive pest species. *Genome Biol* 2016, **17:**192.
167.   Ellis LL, Huang W, Quinn AM, Ahuja A, Alfrejd B, Gomez FE, Hjelmen CE, Moore KL,
Mackay TF, Johnston JS, Tarone AM: **Intrapopulation genome size variation in D.**
melanogaster reflects life history variation and plasticity. *PLoS Genet* 2014, **10:**e1004522.
168.   Kumar S, Stecher G, Tamura K: **MEGA7: Molecular Evolutionary Genetics Analysis**
Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016, **33:**1870-1874.
1818

Fig 1. The large milkweed bug, *Oncopeltus fasciatus*, shown in its phylogenetic and environmental context.

a Dataset comparisons

OGS
n= 19,690

16%

4%

8%

454
n= 12,180
(90% in OGS)

i5K
n= 24,174
(90% in OGS)

c Stage-specific enrichment

Ofas

Clec

Focc

Apis

Pven

b Expression support across the life cycle

n= 19,690

143   273

186   1,081
                       (6%)

            10,646
            (59%)   510
249   (3%)

            3,380
207   (19%)

106   197

410   222

Embryo/
Maternal
(454)

Nymph   Female   Male

	Nymph	Female	Male
Ofas upregulated (#)	400	594	1,180
in ≥3 species (%)	n.d.	23.6%	6.8%
in Heteroptera (%)	n.d.	7.7%	14.6%
unique to *Ofas* (%)	n.d.	51.5%	68.5%
Exemplar classes	Chitin metabolism	Vitellogenin	Ser/Thr kinases
	Cuticle structural proteins	Sugar metabolism	Putative secreted peptides

Fig 2. Comparisons of the official gene set and transcriptomic resources.

Fig 3. Orthology comparisons and phylogenetic placement of *Oncopeltus fasciatus* among other Arthropoda.

Fig 4. Distribution of transcription factor families across insect genomes.

Fig 5. Comparison of repeat content estimations.

Fig 6. Trends in gene structure show hemipteroid-specific tendencies.

Fig 7. Splice site evolution correlates with both lineage and, independently, genome size.

Fig 8. Lateral gene transfer introduction and subsequent evolution within the Hemiptera for mannosidase-encoding genes.

Fig 9. Isoform-specific RNAi based on new genome annotations affects the molting and cuticle identity gene _broad_.

Fig 10. Comparison of the urea cycle of *Oncopeltus* with 26 other insect species.