Molecular evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome

Kristen A. Panfilio[1, 2]*, Iris M. Vargas Jentzsch[1], Joshua B. Benoit[3], Deniz Erezyilmaz[4], Yuichiro Suzuki[5], Stefano Colella[6, 7], Hugh M. Robertson[8], Monica F. Poelchau[9], Robert M. Waterhouse[10, 11], Panagiotis Ioannidis[10], Matthew T. Weirauch[12], Daniel S.T. Hughes[13], Shwetha C. Murali[13, 14, 15], John H. Werren[16], Chris G.C. Jacobs[17, 18], Elizabeth J. Duncan[19, 20], David Armisén[21], Barbara M.I. Vreede[22], Patrice Baa-Puyoulet[6], Chloé S. Berger[21], Chun-che Chang[23], Hsu Chao[13], Mei-Ju M. Chen[9], Yen-Ta Chen[1], Christopher P. Childers[9], Ariel D. Chipman[22], Andrew G. Cridge[19], Antonin J.J. Crumière[21], Peter K. Dearden[19], Elise M. Didion[3], Huyen Dinh[13], HarshaVardhan Doddapaneni[13], Amanda Dolan[16, 24], Shannon Dugan[13], Cassandra G. Extavour[25, 26], Gérard Febvay[6], Markus Friedrich[27], Neta Ginzburg[22], Yi Han[13], Peter Heger[28], Thorsten Horn[1], Yi-min Hsiao[23], Emily C. Jennings[3], J. Spencer Johnston[29], Tamsin E. Jones[25], Jeffery W. Jones[27], Abderrahman Khila[21], Stefan Koelzer[1], Viera Kovacova[30], Megan Leask[19], Sandra L. Lee[13], Chien-Yueh Lee[9], Mackenzie R. Lovegrove[19], Hsiao-ling Lu[23], Yong Lu[31], Patricia J. Moore[32], Monica C. Munoz-Torres[33], Donna M. Muzny[13], Subba R. Palli[34], Nicolas Parisot[6], Leslie Pick[31], Megan Porter[35], Jiaxin Qu[13], Peter N. Refki[21, 36], Rose Richter[16, 37], Rolando Rivera Pomar[38], Andrew J. Rosendale[3], Siegfried Roth[1], Lena Sachs[1], M. Emília Santos[21], Jan Seibert[1], Essia Sghaier[21], Jayendra N. Shukla[34, 39], Richard J. Stancliffe[40], Olivia Tidswell[19, 41], Lucila Traverso[42], Maurijn van der Zee[17], Séverine Viala[21], Kim C. Worley[13], Evgeny M. Zdobnov[10], Richard A. Gibbs[13], Stephen Richards[13]*

* Correspondence: kristen.panfilio@alum.swarthmore.edu; stephenr@bcm.edu

The full list of author information is available at the end of the manuscript.

34   **ABSTRACT**
35
Background:
The Hemiptera (aphids, cicadas, and true bugs) are a key insect order whose members
offer a close outgroup to the Holometabola, with high diversity within the order for
feeding ecology and excellent experimental tractability for molecular genetics.
Sequenced genomes have recently become available for hemipteran pest species such
as phloem-feeding aphids and blood-feeding bed bugs. To complement and build
upon these resources, we present the genome sequence and comparative analyses
centered on the large milkweed bug, *Oncopeltus fasciatus*, a seed feeder of the family
Lygaeidae.
Results:
The 926-Mb genome of *Oncopeltus* is relatively well represented by the current
assembly and official gene set, which supports *Oncopeltus* as a fairly conservative
hemipteran species for anchoring molecular comparisons. We use our genomic and
RNA-seq data not only to characterize features of the protein-coding gene repertoire
and perform isoform-specific RNAi, but also to elucidate patterns of molecular
evolution and physiology. We find ongoing, lineage-specific expansion and
diversification of repressive C2H2 zinc finger proteins and of intron gain and turnover
in the Hemiptera. These analyses also weigh the relative importance of lineage and
genome size as predictors of gene structure evolution in insects. Furthermore, we
identify enzymatic gains and losses that correlate with hemipteran feeding biology,
particularly for reductions in chemoreceptor family size and loss of metabolic
reactions within species with derived, fluid-nutrition feeding modes.
Conclusions:
With the milkweed bug genome, for the first time we have a critical mass of
sequenced species representing a hemimetabolous insect order, substantially
improving the diversity of insect genomics beyond holometabolans such as flies and
ants. We use this addition to define commonalities among the Hemiptera and then
delve into how hemipteran species' genomes reflect their feeding ecology types. Our
novel and detailed analyses integrate global and rigorous manual approaches,
generating hypotheses and identifying specific sets of genes for future investigation.
Given *Oncopeltus*'s strength as an experimental research model, we take particular
care to evaluate the sequence resources presented here, augmenting its foundation for
molecular research and highlighting potentially general considerations exemplified in
the assembly and annotation of this medium-sized genome.
70
71
72   **Keywords:**
73   Phytophagy; Transcription Factors; Gene Structure; Lateral Gene Transfer; RNAi;
74   Gene family evolution; Evolution of Development
75
76

BACKGROUND

In the past few years, the number of animals with sequenced genomes has increased dramatically, and there are now over 100 insect species with assembled and annotated genomes [1]. However, the majority belong to the Holometabola (*e.g.*, flies, beetles, wasps, butterflies), the group characterized by a biphasic life history with distinct larval and adult phases separated by a dramatic metamorphosis during a pupal stage. With fewer than half of all orders, the Holometabola represent only a fraction of the full morphological and ecological diversity across the Insecta. This imbalance in genomic resources limits the exploration of this diversity, including the environmental and developmental requirements of a hemimetabolous life style with a progression of flightless nymphal (juvenile) instars. Addressing this paucity, we report here comparative analyses based on genome sequencing of the large milkweed bug, *Oncopeltus fasciatus*, as a hemimetabolous representative of the larger diversity of insects.

The Hemiptera, the order to which *Oncopeltus* belongs, comprise the most species-rich hemimetabolous order and a close outgroup to the Holometabola as part of the hemipteroid assemblage (or Acercaria), with the Thysanoptera as a sister order and the Psocodea also traditionally included in this clade [2, 3]. All Hemiptera share the same piercing and sucking mouthpart anatomy [4], yet they have diversified to exploit food sources ranging from seeds and plant tissues (phytophagy) to phloem sap (mucivory) and mammalian blood (hematophagy). For this reason, many hemipterans are agricultural pests or human disease vectors, and genome sequencing efforts to date have focused on these species (Fig. 1), including phloem-feeding aphids [5-7], psyllids [8], and planthoppers [9], and the hematophagous kissing bug, *Rhodnius prolixus* [10], a vector of Chagas disease, and bed bug, *Cimex lectularius* [11, 12]. Building on transcriptomic data, genome projects are also in progress for other pest species within the same infraorder as *Oncopeltus*, such as the stink bug *Halyomorpha halys* [13, 14].

In this context, *Oncopeltus* represents a relatively benign species with conservative life history traits, affording a baseline against which other species can be compared. As a seed feeder, *Oncopeltus* has not undergone the marked life style changes that are associated with fluid feeding (mucivory or hematophagy), including dependence on endosymbiotic bacteria to provide needed complements lacking in the diet. For example, in the pea aphid, *Acyrthosiphon pisum*, its obligate endosymbiont, *Buchnera aphidicola,* provides essential amino acids and vitamins: previous analysis of the two genomes revealed a complementation of the two organisms' amino acid metabolism systems [5, 15]. Similarly, although hematophagy arose independently in *Rhodnius* and *Cimex* [16], their respective endosymbionts, *Rhodococcus rhodnii* and *Wolbachia*, provide vitamins lacking in the blood diet [17]. In contrast, the seed-feeding subfamily Lygaeinae, including *Oncopeltus*, is notable for the absence of prominent endosymbiotic anatomy: these bugs lack not only the midgut crypts that typically house bacteria but also the bacteriomes and endosymbiotic balls seen even in other Lygaeidae [18].

Nonetheless, as the native food source of *Oncopeltus* is the milkweed plant, its own feeding biology has a number of interesting implications associated with detoxification and sequestration of cardenolide compounds, including the bright red-

127 orange aposematic (warning) coloration seen in *Oncopeltus* embryos, nymphs, and
128 adults [19, 20]. Thus, diet, metabolism, and body pigmentation are functionally
129 linked biological features for which one may expect changes in gene repertoires to
130 reflect diversity across species of the same order, and the Hemiptera provide an
131 excellent opportunity to explore this.
132
Furthermore, *Oncopeltus* has been an established laboratory model organism
for over 60 years, with a rich experimental tradition in a wide range of studies from
physiology and development to evolutionary ecology [20-22]. It is among the few
experimentally tractable hemimetabolous insect species, and it is amenable to a range
of molecular techniques (*e.g.*, [23-25]). In fact, it was one of the first insect species to
be functionally investigated by RNA interference (RNAi, [26]). RNAi in *Oncopeltus*
is highly effective across different life history stages, which has led to a resurgence of
experimental work over the past fifteen years, with a particular focus on the evolution
of developmentally important regulatory genes (reviewed in [22]).
142
Focusing on these two avenues – feeding biology diversity within the
Hemiptera and *Oncopeltus* as a research model for macroevolutionary genetics – we
present here key insights derived from a combination of global comparative genomics
and detailed computational analyses supported by extensive manual curation,
empirical data for gene expression, sequence validation, and new isoform-specific
RNAi. Namely, we identify sets of genes with potentially restricted life history
expression in *Oncopeltus* and that are unique to the Hemiptera, clarify evolutionary
patterns of zinc finger protein expansion, identify predictors of insect gene structure,
and identify lateral gene transfer and amino acid metabolism features that correlate
with feeding biology.
153
154

RESULTS AND DISCUSSION

The genome and its assembly

Oncopeltus fasciatus has a diploid chromosome number (2*n*) of 16, comprised of seven autosomal pairs and two sex chromosomes with the XX/XY sex determination system [27, 28]. To analyze this genetic resource, we sequenced and assembled the genome using next-generation sequencing approaches (Table 1; see also Methods and Supplemental Notes Sections 1-4). We measure the genome size to be 923 Mb in females and 928 Mb in males based on flow cytometry data (see also Supplemental Note 2.1.a), such that the assembly contains 84% of the expected sequence in assembled contigs, which is comparable to that of other recent, medium-sized insect genomes [11, 29]. However, our analyses of the *k*-mer frequency distribution in raw sequencing reads yielded ambiguous estimates of genome size and heterozygosity rate, which is suggestive of both high heterozygosity and high repetitive content ([30], see also Supplemental Note 2.1.b). Consistent with this, in further analyses we obtained high estimates of repetitive content (see below), which would imply a large proportion of potentially redundant sequence and possible misassembly within contigs of the current assembly. This phenomenon may be increasingly relevant as comparative genomics based on short read sequencing extends to additional insect species with genomes in the 1-Gb range.

As template DNA was prepared from dissected adults from which gut material was removed, the resulting assembly is essentially free of contamination, with only five small scaffolds with high bacterial homology (each to a different, partial bacterial genome, see also Supplemental Note 2.2), which either represent trace bacterial contamination or lateral gene transfers that have not assembled to flanking eukaryotic DNA, making their confirmation in the current assembly difficult.

Table 1. *Oncopeltus fasciatus* genome metrics.

Feature	Value	
2*n* chromosomes	16	
Genome size	926 Mb (mean between males and females)	
Assembly size	1,099 Mb (contigs only: 774 Mb)	
Coverage	106.9× raw coverage, 83.7% of reads in final assembly	
Contig N50	4,047 bp	
Scaffold N50	340.0 kb	
# Scaffolds	17,222	
GC content	genome: 32.7%, protein-coding sequence (OGS v1.2): 42%	
OGS v1.1 (curated fraction)	19,690 models[1] (1,426 models, 7.2%)	19,465 genes (1,201 genes, 6.2%)
OGS v1.2 (curated fraction)	19,809 models[1] (1,697 models, 8.7%)	19,616 genes (1,518 genes, 7.7%)

[1] Individual genes may be represented by multiple models in cases of curated alternative isoforms or if the gene is split across scaffolds.

The official gene set and conserved gene linkage

The official gene set (OGS) was generated by automatic annotation followed by manual curation in a large-scale effort by the research community (see also Supplemental Notes Sections 3-4). Curation revised automatic annotation models, added alternative isoforms and *de novo* models, and documented multiple discrete models for genes whose exons were split across scaffolds. We found that automatic predictions were somewhat conservative for hemipteran gene structure (see below), and manual curation primarily resulted in larger gene loci as exons were added and/or extended, including merging discrete automatic models (see also Supplemental Note 4, Table S4.4). The OGS v1.1 was generated for global, pipeline analyses to characterize the gene repertoire. The latest version, OGS v1.2, represents a minor update, primarily for the addition of chemoreceptor genes of the ionotropic and odorant receptor classes and curation of genes encoding metabolic enzymes. Altogether, the research community curated 1,697 gene models (8.7% of OGS v1.2), including 316 *de novo* models (see also Table S4.1). Reflecting the primary research interests of the community (see also Supplemental Notes Section 5), the majority of curated models are for genes encoding cuticular proteins (11%), chemoreceptors (19%), and developmental regulators such as transcription factors and signaling pathway components (40%, including the BMP/TGF-β, Toll/NF-κB, Notch, Hedgehog, Torso RTK, and Wnt pathways).

In addition to assessing gene model quality, manual curation of genes whose orthologs are expected to occur in syntenic clusters also validates assembly scaffolding. Complete loci could be found for single orthologues of all Hox cluster genes, where *Hox3*/*zen* and *Hox4*/*Dfd* are linked in the current assembly and have ≥99.9% nucleotide identity with experimentally validated sequences ([31-33], Supplemental Note 5.1.b). Conserved linkage was also confirmed for the homeobox genes of the Iroquois complex, the Wnt ligands *wingless* and *wnt10*, and two linked pairs from the Runt transcription factor complex (Supplemental Notes 5.1.a, 5.1.c, 5.1.i, 5.1.j). Further evidence for correct scaffold assembly comes from the curation of large, multi-exonic loci. For example, the cell polarity and cytoskeletal regulator encoded by the conserved *furry* gene includes 47 exons spanning a 437-kb locus, which were all correctly assembled on a single scaffold.

Transcriptomic resources and gene expression profiles across the milkweed bug life cycle

To augment published transcriptomic resources [34, 35], we sequenced three different post-embryonic samples ("i5K" dataset, see Methods). We then compared the OGS to the resulting *de novo* transcriptome and to a previously published embryonic and maternal (ovary) transcriptome ("454" pyrosequencing dataset, [34]). Our OGS is quite comprehensive, containing 90% of transcripts from each transcriptomic dataset and an additional 3,146 models (16% of OGS: Fig. 2a). Among the additional models, 274 (9%) were manually validated, including 163 *de novo* models for odorant and gustatory receptors. These gene classes are known for lineage-specific expansions and highly tissue- and stage-specific expression, with usually only one receptor expressed per sensory neuron ([36, 37], and see below).

Furthermore, the OGS does a good job of "mopping up" partial and
unidentified 454 transcripts. We could substantially improve orthologous gene
discovery by mapping the 454 transcripts to the OGS (blastn, $e<10^{-9}$), nearly trebling
the proportion of transcripts with an assigned gene model or homology compared to
the original study (from 9% to 26%). This included 10,130 transcripts that primarily
mapped to UTRs and could not have been identified by coding sequence homology,
such as the 654-bp transcript for the *Oncopeltus brinker* ortholog, which encodes a
putative inhibitor of the BMP pathway ([38], see also Supplemental Note 5.1.f), and
four unassembled transcripts each from the 3′ UTRs of the enzyme-encoding genes
CTP synthase and *roquin*. At the same time, the transcriptomes provided expression
support for the identification of multiple isoforms in the OGS. For example, we could
confirm previously described isoforms for the germline determinant encoded by
nanos [34]. Where assembly limitations curtailed OGS gene models, full-length
transcripts are represented in the transcriptomes, such as for the ecdysis regulator
CCAP-R [39] and the chromatin linker Histone H1.
256
We then took advantage of our stage-specific RNA datasets to provide an
initial survey of gene expression profiles across biological samples and across the life
cycle. Most OGS gene models have expression support (91% of 19,690), with 74%
expressed broadly in at least three of four samples (Fig. 2b). The inclusion of a fifth
dataset from a published adult library [35] provided only a 1% gain in expression
support (218 gene models), indicating that with the current study the expression data
volume for *Oncopeltus* is quite complete. At the same time, direct comparison of the
three adult samples suggests that the published adult dataset of unspecified sex is
probably male, as it shares 4.6× more expressed genes with our male than our female
sample.
267
As these data derive from limited biological sampling, we remain cautious
about true stage specificity and do not quantify expression levels. We do, however,
note that most genes with stage-restricted expression are in sets involving our male
sample (Fig. 2b: male-only or male and nymph), although this sample does not
contain more reads or more expressed genes. Furthermore, we also find stage-
specific patterns for some of our most abundant curated gene classes. Gustatory
receptor (GR) genes show noticeable restriction to the adult male and published adult
(probable male) samples (n= 169 GRs: 40% no expression, 27% only expressed in
these two samples), with half of these expressed in both biological replicates (52%).
Interestingly, the nymphal sample is enriched for genes encoding structural cuticular
proteins (94%, which is >56% more than any other sample). This likely reflects the
ongoing molting cycles, with their cyclical upregulation of cuticular gene synthesis
[40], that are experienced by the different instars and molt cycle stages of individuals
pooled in this sample.
282
283
Protein orthology and hemipteran copy number comparisons
To further assay protein-coding gene content, we then compared *Oncopeltus* with
eleven other arthropod species. A phylogeny based on single copy orthologs correctly
reconstructs the hemipteran and holometabolan clades' topologies (Fig. 3a, compare
with Fig. 1a), although larger-scale insect relationships remain challenging [3]. In
expanding this to the Benchmarking Universal Single-Copy Orthologs (BUSCO,
[41]) dataset of 2,675 Arthropoda genes, we also found that most BUSCO genes are

present in the *Oncopeltus* OGS, although with additional genes identified on genomic
scaffolds but not yet incorporated into the gene set (see also Supplemental Note 6.1).
We next categorized all proteins by conservation in global, clustering-based orthology
analyses [42]. As in most species, half of *Oncopeltus* proteins (51%) falls within the
top three conservation levels (Fig. 3a). Moreover, 98% of all *Oncopeltus* protein-
coding genes has homology, expression, and/or curation support (Fig. 3b), including
support for 80% of proteins without homology, such as a few species-specific
chemoreceptors and antimicrobial peptides (see also Supplemental Note 5.1.h), while
some unsupported models may be split or partial. Overall, we estimate that the
Oncopeltus protein repertoire is comparable to that of other insects in size and degree
of conservation.
302
In contrast, the pea aphid, *Acyrthosiphon pisum*, is a notable outlier even
among fellow Hemiptera, where we provide a side-by-side comparison with
Oncopeltus as well as the recently-sequenced kissing bug, *Rhodnius prolixus* [10],
and bed bug, *Cimex lectularius* [11, 12]. The pea aphid is striking for its long branch
in phylogenetic comparisons and for its large protein-coding gene content with low
conservation (Fig. 3a), consistent with the observation of numerous lineage-specific
duplications [5]. As the first hemipteran to have its genome sequenced, the pea aphid
has often been used to boost taxonomic sampling in phylogenomic comparisons (*e.g.*,
[29]). However, the pea aphid may not be the best representative, and as more
hemipteran genomes are sequenced, other species now offer less derived alternatives.
313
314
Compared to the pea aphid [43], *Oncopeltus* is more conservative in both
presence and copy number for several signaling pathway components. In contrast to
gene absences described for the pea aphid, *Oncopeltus* retains orthologs of the EGF
pathway component *sprouty*, the BMP receptor *wishful thinking*, and the hormone
nuclear receptor *Hr96* (see also Supplemental Note 5.1.e). Similarly, whereas
multiple copies were reported for the pea aphid, we find a single *Oncopeltus* ortholog
for the BMP pathway components *decapentaplegic* and *Medea* and the Wnt pathway
intracellular regulator encoded by *shaggy/GSK-3*, albeit with five potential isoforms
of the latter (see also Supplemental Notes 5.1.f, 5.1.j). Duplications of miRNA and
piRNA gene silencing components likewise seem to be restricted to the pea aphid
compared to other hemipterans – including other aphid species ([44], see also
Supplemental Note 5.4.a). However, our survey of *Oncopeltus* and several other
hemimetabolous species reveals evidence for frequent parallel duplications of the Wnt
pathway component *armadillo/β-catenin*, yet without the sequence and functional
divergence previously observed independently in the pea aphid and *Tribolium* ([45],
see also Supplemental Note 5.1.j). Curiously, *Oncopeltus* appears to encode fewer
histone loci than any other arthropod genome and yet exhibits a similar, but possibly
independent, pattern of duplications of histone acetyltransferases to those previously
identified in *Cimex* and the pea aphid (see also Supplemental Note 5.4.c).
334
On the other hand, we documented several notable *Oncopeltus*-specific
duplications. Whereas two copies of the BMP transducer *Mad* were reported in the
pea aphid [43], we find evidence for three paralogs in *Oncopeltus*, where two of these
genes occur in tandem and may reflect a particularly recent duplication (see also
Supplemental Note 5.1.f). Similarly, a tandem duplication of *wnt8* appears to be
unique to *Oncopeltus* (see also Supplemental Note 5.1.j). More striking is the

identification of six potential paralogs of *cactus*, a member of the Toll/NF-κB
signaling pathway for innate immunity, whereas the bed bug and kissing bug
each retain only a single copy ([46], see also Supplemental Note 5.1.g).
344
345    We then took advantage of broader comparative datasets [42] to identify
lineage-specific features of the Hemiptera. In other words, what makes a bug a bug in
terms of protein-coding genes? To address this, we partitioned an orthology analysis
of 64 insect species into three broad taxonomic groups (Fig. 3c). Highlights for the
Hemiptera, which are further corroborated in an updated dataset with 116 insect
species (OrthoDB v.9.1, [1]), fall into two classes. The first class contains potentially
new genes that show no homology outside the Hemiptera. We identified three such
instances with orthologous protein members present in at least four hemipterans, and
where no conserved functional domains were recognized. Interpretation of these
intriguing "uncharacterized proteins" will have to await direct experimental analyses,
for which the Hemiptera in general are particularly amenable (*e.g.*, [47-50]). The
second class comprises proteins with recognized functional domains and homologs in
other insects, but where evolutionary divergence has led to hemipteran-specific
subfamilies. For example, one protein orthology group ("orthogroup"
EOG090W0V4B) is comprised of a heteropteran-specific cytochrome P450 (CYP)
enzyme that in *Oncopeltus* is expressed in all life history stages. The expansion of
this protein family is associated with a species' potential scope for insecticide
resistance, as specific P450s can confer resistance to specific chemicals (*e.g.,* [51, 52];
see also Supplemental Notes 5.3.b, 5.3.c). Hence, the identification of lineage-
specific CYP enzymes can suggest potential targets for integrated pest management
approaches.
366
367
Transcription factor repertoires and homeobox gene evolution
Having explored the global protein repertoire, we next focused specifically on
transcription factors (TFs), which comprise a major class of proteins that has been
extensively studied in *Oncopeltus*. This is a class of key regulators of development
whose functions can diverge substantially during evolution and for which RNAi-
based experimental investigations have been particularly fruitful in the milkweed bug
(*e.g.*, [31, 32, 53-55], see also Supplemental Notes 5.1.a-e).
375
376    To systematically evaluate the *Oncopeltus* TF repertoire, we used a pipeline to
scan all predicted proteins and assign them to TF families, including orthology
assignments in cases where DNA binding motifs could be predicted (see Methods,
[56]). We identified 762 putative TFs in *Oncopeltus*, which is similar to other insects
of diverse orders for total TF count and for the size of each TF family (Fig. 4a: note
that the heatmap also reflects the large, duplicated repertoire in the pea aphid; see also
Tables S6.2-6.4).
383
384    We were able to infer DNA binding motifs for 25% (n=189) of *Oncopeltus*
TFs, mostly based on data from *D. melanogaster* (121 TFs) but also from distantly
related taxa such as mammals (56 TFs). Such high conservation is further reflected in
the fact that most proteins within several large TF families have inferred motifs and
therefore explicit orthology assignments, including for the homeodomain (53 of 85,
62%), basic helix-loop-helix (bHLH, 35 of 45, 78%), and forkhead box (16 of 17,
94%) families. In contrast, most C2H2 zinc finger proteins lack orthology assignment

(only 22 of 360, 6%). Across species, the homeodomain and C2H2 zinc finger
proteins are the two largest TF superfamilies (Fig. 4a). Given their very different
rates of orthology assignment in *Oncopeltus*, we probed further into the pipeline
predictions and the patterns of evolutionary diversification of these proteins.
395
The number of homeodomain proteins identified by the pipeline displays a
narrow normal distribution across species (Fig. 4b, mean ± standard deviation: 97 ±
9), consistent with a highly conserved, slowly evolving protein family. Supporting
this, many *Oncopeltus* homeodomain proteins that were manually curated also
received a clear orthology assignment (Fig. 4c: pink), with only four exceptions (Fig.
4c: yellow). Only one case suggests a limitation of a pipeline that is not specifically
tuned to hemipteran proteins (Goosecoid), while an incomplete gene model received
homeodomain classification but without explicit orthology assignment (Distal-less).
Manual curation of other partial or split models identified a further 11 genes encoding
homeodomains, bringing the actual tally in *Oncopeltus* to 96, which is comparable to
the mean across species. Overall, we find the TF pipeline results to be a robust and
reasonably comprehensive representation of these gene classes in *Oncopeltus*.
408
These analyses also uncovered a correction to the published *Oncopeltus*
literature for the key developmental patterning proteins encoded by the closely related
paralogs *engrailed* and *invected*. These genes arose from an ancient tandem
duplication prior to the hexapod radiation, where their tail-to-tail orientation enables
ongoing gene conversion [57], making orthology discrimination particularly
challenging. For *Oncopeltus*, we find that the genes also occur in a tail-to-tail
orientation and that *invected* retains a diagnostic alternative exon [57]. These new
genomic and expression data reveal that the *Oncopeltus* gene used as the purported
engrailed ortholog in previous developmental studies (*e.g.*, [53, 58-61]) is in fact
invected (see also Supplemental Note 5.1.a).
419
420
Independent expansions of C2H2 zinc fingers within the Hemiptera
Unlike homeodomain proteins, C2H2 zinc finger (C2H2-ZF) repertoires are
prominent for their large family size and variability throughout the animal kingdom
[62], and this is further supported by our current analysis in insects. With >350
C2H2-ZFs, *Oncopeltus*, the pea aphid, the termite, and several mosquito species have
1.5× more members than the insect median (Fig. 4b). This is nearly half of all
Oncopeltus TFs. While the expansion in mosquitoes could have a single origin after
the Culicinae diverged from the Anophelinae, the distribution in the Hemiptera, where
Cimex has only 227 C2H2-ZFs, suggests that independent expansions occurred in
Oncopeltus and the pea aphid. Prior to the sequencing of other hemipteran genomes,
the pea aphid's large C2H2-ZF repertoire was attributed to the expansion of a novel
subfamily, APEZ, also referred to as zinc finger 271-like [43].
433
In fact, manual curation in *Oncopeltus* confirms the presence of a subfamily
with similar characteristics to APEZ (Fig. 4c: 42% of all C2H2-ZFs were curated,
including 38% of those without orthology assignment, yellow). Specifically, in
Oncopeltus we find >115 proteins of the ZF271 class that are characterized by
numerous tandem repeats of the C2H2-ZF domain and its penta-peptide linker, with
3-45 repeats per protein.
440

However, at both the gene and protein levels we find evidence for ongoing
evolutionary diversification of the *Oncopeltus* ZF271-like subfamily. A number of
Oncopeltus ZF271-like genes occur in tandem clusters of 4-8 genes, suggesting recent
duplication events. Yet, at the same time, gene structure (number and size of exons)
is not shared between genes within clusters, and we identified a number of probable
ZF271-like pseudogenes whose open reading frames have become disrupted –
consistent with high turnover. At the domain level, *Oncopeltus* ZF271-like proteins
differ in the sequence and length of the zinc finger domains amongst themselves and
compared to aphid proteins (WebLogo analysis [63]), similar to zinc finger array
shuffling seen in humans [64]. Furthermore, whole-protein phylogenetic analysis
supports independent, rapid expansions in the pea aphid and *Oncopeltus* (Fig. 4d).
452
Clustered zinc finger gene expansion has long been recognized in mammals,
with evidence for strong positive selection to increase both the number and diversity
of zinc finger domains per protein as well as the total number of proteins [65]. This
was initially found to reflect an arms-race dynamic of co-evolution between selfish
transposable elements and the C2H2-ZF proteins that would repress them [66]. In
vertebrates, these C2H2-ZF proteins bind to the promoters of transposable elements
via their zinc finger arrays and use their Krüppel-associated box (KRAB) domain to
bind the chromatin-remodeling co-repressor KAP-1, which in turn recruits
methyltransferases and deacetylases that silence the targeted promoter [67].
462
Insects do not have a direct ortholog of vertebrate KAP-1 (see also
Supplemental Note 5.4.d), and neither the aphid nor *Oncopeltus* ZF271-like
subfamilies possess a KRAB domain or any other domain besides the zinc finger
arrays. However, close molecular outgroups to this ZF271-like subfamily include the
developmental repressor Krüppel [68] and the insulator protein CTCF [69] (data not
shown). Like these outgroups, the *Oncopeltus* ZF271-like genes are strongly
expressed: 98% have expression support, with 86% expressed in at least three
different life history stages (Fig. 2b). Thus, the insect ZF271-like proteins may also
play prominent roles in repressive DNA binding. Indeed, we find evidence for a
functional methylation system in *Oncopeltus* (see also Supplemental Note 5.4.c), like
the pea aphid, which would provide a means of gene silencing by chromatin
remodeling, albeit via mediators other than KAP-1.
475
However, an arms race model need not be the selective pressure that favors
insect ZF271-like family expansions. Recent analyses in vertebrates identified
sophisticated, additional regulatory potential by C2H2-ZF proteins, building upon
original transposable element binding for new, lineage-specific and even positive
gene regulation roles [64, 70, 71]. Moreover, although *Cimex* has half as many long
terminal repeat (LTR) repetitive elements as *Oncopeltus* and the pea aphid, they
constitute only a minor fraction of these species' transposable elements, and overall
we do not find a correlation between relative or absolute repetitive content and
ZF271-like family expansion within the Hemiptera (Fig. 5, and see below).
485
486
Proportional repeat content across hemipterans
With the aim of reducing assembly fragmentation and to obtain a better picture of
repeat content, we performed low coverage, long read PacBio sequencing in
Oncopeltus (see also Supplemental Note 2.3). Using PacBio reads in a gap-filling

assay on the Illumina assembly raised the total detected repetitive content from 25%
to 32%, while repeat estimations based on simultaneous assessment of Illumina and
PacBio reads nearly doubled this value to 58%. As expected, the capacity to identify
repeats is strongly dependent on assembly quality and sequencing technology, with
the *Oncopeltus* repetitive content underrepresented in the current (Illumina-only)
assembly. Furthermore, as increasing genome size compounds the challenge of
assembling repeats, the repeat content of the current assembly is lower than in species
with smaller genome sizes (Fig. 5a, with the sole exception of the honey bee), and we
therefore used our gap-filled dataset for further repeat profile comparisons.
500
To allow for direct comparisons among hemipterans, we also performed our
RepeatModeler analysis on the bed bug and pea aphid assemblies. In these analyses,
36% and 31% of the respective assemblies were covered by repeats, similar to the
gap-filled value of 32% in *Oncopeltus*. Nevertheless, given the smaller sizes of these
species' assemblies – 651 Mb in the bed bug and 542 Mb in the pea aphid – the
absolute repeat content is much higher in *Oncopeltus* (Fig. 5b). Excluding unknown
repeats, the most abundant transposable elements in *Oncopeltus* are LINE
retrotransposons, covering 10% of the assembly (see also Table S2.5). This is also
the case in the bed bug (12%), while in the pea aphid DNA transposons with terminal
inverted repeats (TIRs) are the most abundant (2% of the assembly identified here,
and 4% reported from manual curation in the pea aphid genome paper, [5]). Across
species, the remaining repeat categories appear to grow proportionally with assembly
size, except for simple repeats, which were the category with the largest relative
increase in size after gap-filling in *Oncopeltus* (see also Supplemental Note 2.3).
However, given the mix of sequence data types (Illumina only in the bed bug [11],
Sanger in the pea aphid [5]), these patterns should be treated as hypotheses for future
testing, until the assembly of repetitive regions becomes more feasible.
518
519
Lineage and genome size-related trends in insect gene structure
During manual curation, we noticed that *Oncopeltus* genes were often comprised of
many, small exons. Furthermore, sequence conservation among the Hemiptera
supported terminal coding sequence exons that were small and separated from the rest
of the gene model by large introns. To explore patterns of gene structure across the
insects, we undertook a broader comparative analysis. We find that both lineage and
genome size can serve as predictors of gene structure.
527
Firstly, we created a high quality ("gold standard") dataset of 30 functionally
diverse, large genes whose manual curation could reasonably ensure complete gene
models across seven species from four insect orders (Fig. 6a; see also Supplemental
Note 6.3). Most species encode the same total number of amino acids for these
conserved proteins, with the thrips *Frankliniella occidentalis* and *Drosophila* being
notable exceptions with larger proteins (Fig. 6a: blue plot line). However, the means
of encoding this information differs between lineages, with hemipteroid orthologs
comprised of twice as many exons as their holometabolous counterparts (Fig. 6a:
orange plot line). Thus, there is an inverse correlation between exon number and
exon size (Fig. 6a: orange vs. red plot lines). This analysis corroborates and extends
previous probabilistic estimates of intron density, where the pea aphid as a sole
hemipteran representative had the highest intron density of ten insect species [72].
540

To test these trends, we next expanded our analysis to all manually curated
exons in two species from each of three orders, including the Mediterranean fruit fly,
Ceratitis capitata [73], as a second dipteran alongside *Drosophila melanogaster*.
Here, we expect that curated exon sizes are accurate, without the need to assume that
entire gene models are complete. This large dataset supports our original findings,
with bugs having small exons, and with both the median and Q3 quartile reflecting
larger exon sizes in beetles and flies (Fig. 6b). Notably, the median and median
absolute deviation are highly similar between species pairs within the Hemiptera and
Coleoptera, irrespective of sample size. Meanwhile, the different exon metrics
between *Ceratitis* and *Drosophila* suggest that the large protein sizes we initially
observed in *Drosophila* (Fig. 6a: blue plot line) are a general but drosophilid-specific,
rather than dipteran-wide, feature.
553
Does the high exon count in the Hemiptera reflect an ancient, conserved
increase at the base of this lineage, or ongoing remodeling of gene structure with high
turnover? To assess the exact nature of evolutionary changes, we annotated intron
positions within multiple sequence alignments of selected proteins and plotted gains
and losses onto the phylogeny, providing a total sample of 165 evolutionary changes
at 148 discrete splice sites (Fig. 7; see also Supplemental Note 6.3 for gene selection
and method). These data reveal several major correlates with intron gain or loss. The
bases of both the hemipteroid and hemipteran radiations show the largest gains, while
most losses occur in the dipteran lineage (Fig. 7: orange and purple shading,
respectively). Furthermore, we find progressive gains across the hemipteroid nodes,
and it is only in these species that we additionally find species-specific splice changes
for the highly conserved *epimerase* gene (Fig. 7: orange outline). Thus, we find
evidence for both ancient intron gain and ongoing gene structure remodeling in this
lineage.
568
Surprisingly, both *hemocytin* and *epimerase* – our exemplar genes with many
(up to 74) and very few exons (3-8 per species), respectively – show independent
losses of the same splice sites in *Drosophila* and *Tribolium*. One feature these species
share is a genome size 2.4-6.0× smaller than all other species examined here (Fig. 7:
red shading). Pairwise comparisons within orders also support this trend, as the beetle
and fly species with larger genomes exhibit species-specific gains compared to intron
loss in their sister taxa (Fig. 7: red outlines). Thus, while lineage is a stronger
predictor of gene structure evolution (the coleopteran and dipteran species include one
each with a big or small genome and yet have highly similar metrics in Fig. 6b),
genome size seems to positively correlate with intron number (*e.g.,* the common
dipteran ancestor lost introns before *Ceratitis*, with a larger genome, experienced
subsequent gains: Fig. 7). A global computational analysis over longer evolutionary
distances also supports a link between genome size and intron number within
arthropods, but where chelicerates and insects may experience different rates of
evolutionary change in these features [74]. As new insect species' genomes are
sequenced, gene structure expectations at the ordinal level can help customize
parameters for automatic gene annotation, while it will be interesting to see if the
correlation with genome size is borne out in other taxa.
587
The selective pressures and mechanisms of intron gain in the Hemiptera will
be a challenge to uncover. While median exon size (Fig. 6b) could reflect species-
specific nucleosome sizes [75, 76], this does not account for the fact that most

hemipteran exons do not occur in multiples greater than a single nucleosome. Given
gaps in draft genome assemblies, we remain cautious about interpreting (large) intron
lengths but note that many hemipteran introns are too small to have harbored a
functional transposase gene (*e.g.*, median intron size of 429 bp, n=69 introns in
hemocytin in *Cimex*). Such small introns could be consistent with proliferation of
non-autonomous short interspersed nuclear elements (SINEs), although as highly
divergent non-coding elements their characterization in insects would require curated
SINE libraries comparable to those generated for vertebrates and plants [75, 76].
Meanwhile, it appears that hemipteran open reading frames ≥160 bp are generally
prevented by numerous in-frame stop codons just after the donor splice site. Most
stop codons are encoded by the triplet TAA in both *Oncopeltus* and *Cimex* (data not
shown), although these species' genomes are not particularly AT-rich (Table 1).

Even if introns are small, having gene loci comprised of numerous introns and
exons adds to the cost of gene expression in terms of both transcription duration and
mRNA processing. One could argue that a gene like *hemocytin*, which encodes a
clotting agent, would require rapid expression in the case of wounding – a common
occurrence in adult *Cimex* females due to the traumatic insemination method of
reproduction [11]. Thus, as our molecular understanding of comparative insect and
particularly hemipteran biology deepens, we will need to increasingly consider how
life history traits are manifest in genomic signatures at the structural level (*e.g.*, Figs.
5-7), as well as in terms of protein repertoires (Figs. 3-4).

Expansion after a novel lateral gene transfer (LGT) event in phytophagous bugs
In addition to the need for cuticle repair, traumatic insemination may be responsible
for the numerous LGT events predicted in the bed bug [11]. In contrast, the same
pipeline analyses [77] followed by manual curation predicted very few LGTs in
Oncopeltus, which lacks this unusual mating behavior. Here, we have identified 11
strong LGT candidates, and we confirmed the incorporation of bacterial DNA into the
milkweed bug genome for all five candidates chosen for empirical testing (see also
Table S2.4). Curiously, we find several LGTs potentially involved in bacterial or
plant cell wall metabolism that were acquired from different bacterial sources at
different times during hemipteran lineage evolution, including two distinct LGTs that
are unique to *Oncopeltus* and implicated in the synthesis of peptidoglycan, a bacterial
cell wall constituent (see also Supplemental Note 2.2).

Conversely, two further validated LGT candidates encode enzymes rather
known for their roles in degradation of bacterial cell walls: we find two strongly
expressed, paralogous copies in *Oncopeltus* of a probable bacterial-origin gene
encoding an endo-1,4-beta-mannosidase enzyme (MAN4, EC 3.2.1.78). This likely
ancient LGT event provides an interesting vignette that further illustrates gene
structure evolution processes within the Hemiptera. Inspection of genome assemblies
and predicted protein accessions reveals that this LGT event is shared with the stink
bug *Halyomorpha halys*, a member of the same infraorder (Pentatomomorpha), but
was introduced after this lineage diverged from other hemipterans, including the bed
bug (Fig. 8a). Furthermore, whereas *Oncopeltus* now has two copies of this gene,
independently the original *Halyomorpha* gene underwent a series of tandem
duplications leading to nine extant copies (Fig. 8b, see also Fig. S2.6). Since the
original LGT event, the *mannosidase* genes in both bug species have become

"domesticated" as multi-exonic genes (as in [78]). Moreover, as the splice site pattern
is unique to each species and evinces subsequent splice introductions in subsets of
paralogs (Fig. 8c), *mannosidase* genes further illustrate the hemipteran penchant for
intron introduction and maintenance of small exons. The retention and subsequent
expansion of these genes implies their positive selection, consistent with the
phytophagous diet of these hemipteran species. In this context, it is tempting to
speculate further that the marked proliferation of this enzyme in the stink bug
correlates with the breadth of its diet, as this agricultural pest feeds on a number of
different tissues in a range of host plants [79].
650
651

Cuticle development, structure, and warning pigmentation
Given the milkweed bug's history as a powerful model for endocrine studies of
hemimetabolous molting and metamorphosis since the late 1960's [21, 80-83], we
next focused on genes underlying the development and structural properties of the
Oncopeltus cuticle. Molting is triggered by the release of ecdysteroids, steroid
hormones that are synthesized from cholesterol in the prothoracic gland by
cytochrome P450 enzymes of the Halloween family [84], and we were able to identify
these in the *Oncopeltus* genome (see also Supplemental Notes 5.2.b, 5.3.b for these
and following metamorphosis gene details). From the ecdysone response cascade
defined in *Drosophila* [85], we identified *Oncopeltus* orthologs of both early and late-
acting factors. It will be interesting to see if the same regulatory relationships are
conserved in the context of hemimetabolous molting in *Oncopeltus*. For example,
E75A is required for reactivation of ecdysteroid production during the molt cycle in
Drosophila larvae [86] and likely operates similarly in *Oncopeltus*, since *Of-E75A*
RNAi prevents fourth-instar nymphs from molting to the fifth instar (H. Kelstrup and
L. Riddiford, unpublished data). In holometabolous insects, a declining titer of
ecdysteroids leads to the release of a series of neuropeptides that ultimately causes the
insect to molt, or ecdyse [87, 88]. Orthologs of these hormones and their receptors
are also present in the *Oncopeltus* genome or transcriptomic data.
671

In hemipterans, activation of juvenile hormone (JH) signaling at molts
determines whether the insect progresses to another nymphal instar or, if lacking,
becomes an adult [47]. We were able to identify many components of the JH signal
transduction pathway in the *Oncopeltus* genome, including an ortholog of
Methoprene-tolerant (*Met*), the JH receptor [47, 89], and the JH-response gene *Kr-h1*
[47, 90, 91]. JH acts to determine cuticle identity through regulation of the *broad*
gene in a wide variety of insects, where different isoforms direct specific aspects of
metamorphosis in *Drosophila* [92, 93]. In *Oncopeltus*, *broad* expression directs
progression through each of the nymphal stages [94], but the effect of each isoform
was unknown. We identified three isoforms in *Oncopeltus* – Z2, Z3, and Z4 – and
performed isoform-specific RNAi. In contrast to *Drosophila*, Broad isoform
functions appear to be more redundant in *Oncopeltus*, as knockdown of isoforms Z2
and Z3 have similar effects on survival to adulthood as well as adult wing size and
morphology (Fig. 9).
686

Regulators such as Broad initiate the transcription of a large battery of genes
that encode the structural components of the cuticle needed at each molt, consistent
with our expression analyses (Fig. 2b, discussed above). We identified 173 genes
encoding putative cuticle structural proteins in the milkweed bug, using established

sequence motifs (see also Supplemental Note 5.2.c). Similar to other insects, the CPR
family, with the RR-1 (soft cuticle), RR-2 (hard cuticle), and unclassifiable types,
constituted the largest of the cuticle protein gene groups. While several protein
families are similar in size to those of other insects (CPAP1, CPAP3, and TWDL: see
also Table S5.12), we found a slight expansion in the *Oncopeltus* CPF family (see
also Fig. S5.14). For cuticle production, similar to the bed bug and the Asian
longhorned beetle [11, 29], we identified a single *chitin synthase* gene with conserved
alternative splice isoforms, which suggests that *chitin synthase 2* is a duplication
specific to only certain beetle and fly lineages within the Holometabola [95].
700
One of the major characteristics of the milkweed bug is the distinctive red-
orange and black aposematic (warning) coloration within the cuticle and epidermis
that has been shown to act as a deterrent to predators (*e.g.*, Figs. 1, 9, [19, 20]). For
black coloration, we were able to identify the key melanin synthesis enzymes (see
also Fig. S5.15). The melanin synthesis pathway is conserved across holometabolous
insects (*e.g.*, [96, 97]), and recent work in *Oncopeltus* [98, 99] supports functional
conservation of melanin production in hemimetabolous lineages as well. In contrast,
production of the primary warning coloration produced by the pteridine red
erythropterin [100] remains an open avenue for hemimetabolous research. This
pigment, along with other pterins, is synthesized from GTP through a series of
enzymatic reactions [101]. The genes encoding enzymes that convert GTP into
pterins have not been as extensively studied as melanins, and thus far in *Oncopeltus*
we were only able to identify orthologs of *punch*, which encodes a GTP
cyclohydrolase [102], and *sepia*, which is required for the synthesis of the red eye
pigment drosopterin [103]. The bright red color of *Oncopeltus* eggs may in part
reflect chemical protection transmitted parentally [104]. Thus, further identification
of pigmentation genes will provide fitness indicators for maternal contributions to
developmental success under natural conditions (*i.e.*, the presence of egg predators).
719
720
Chemoreception and metabolism in relation to feeding biology
The aposematic pigmentation of the milkweed bug advertises the fact that toxins in its
milkweed seed diet are incorporated into the bugs themselves, a metabolic feat that
was independently acquired in *Oncopeltus* and in the similarly colored monarch
butterfly (*Danaus plexippus*), which shares this food source [35, 105]. Moreover,
given the fundamental differences in metabolic pathways between phytophagous,
mucivorous, and hematophagous species, we investigated to what extent differences
in feeding ecology across hemipterans are represented in the chemoreceptor and
metabolic enzyme repertoires of these species.
730
The ability of insects to smell and taste the enormous diversity of chemicals
important to them for locating and identifying food, mates, oviposition sites, and other
aspects of their environment is primarily mediated by three large gene families. The
closely related Odorant Receptor (OR) and Gustatory Receptor (GR) families, and the
distinct Ionotropic Receptor (IR) family [106-109], commonly encode tens to
hundreds of chemoreceptors in arthropods. Consistent with having a less derived
feeding ecology than species with phloem-restricted or obligate hematophagous diets,
Oncopeltus retains a moderate complement of chemoreceptors from the different
classes (Table 2, see also Supplemental Note 5.3.f). The hematophagous *Cimex* and
Rhodnius have relatively depauperate OR and GR families compared to *Oncopeltus*.

While a few conserved orthologs such as the OrCo protein and a fructose receptor are found across species, *Oncopeltus* and *Acyrthosiphon* retain a set of sugar receptors, a gene lineage lost independently from the blood-feeding bugs (*Rhodnius* [10], *Cimex* [11]) and body louse (*Pediculus* [110]). Conversely, *Oncopeltus* has, like *Cimex*, a set of candidate carbon dioxide receptors, a gene lineage lost from *Rhodnius*, *Acyrthosiphon*, and *Pediculus* [10, 11, 111], but which is similar to a GR subfamily expansion in the more distantly related hemimetabolous termite (Isoptera, [112]). Comparable numbers of IRs occur across the heteropterans, where in addition to a conserved set of orthologs primarily involved in sensing temperature and certain acids and amines, *Oncopeltus* has a minor expansion of IRs distantly related to those involved in taste in *Drosophila*. The major expansions in each insect lineage are the candidate "bitter" GRs ([113], see also Supplemental Note 5.3.f, Fig. S5.19). In summary, *Oncopeltus* exhibits moderate expansion of specific subfamilies likely to be involved in host plant recognition, consistent with it being a preferentially specialist feeder with a potentially patchy food source [20, 114].

Table 2. Numbers of chemoreceptor genes/proteins per family in selected insect species. In some cases the number of proteins is higher than the number of genes due to an unusual form of alternative splicing, which is particularly notable for the *Oncopeltus* GRs. Data are shown for four Hemiptera as well as *Drosophila melanogaster*, the body louse *Pediculus humanus*, and the termite *Zootermopsis nevadensis* [10, 11, 108, 110-112, 115].

Species	Odorant	Gustatory	Ionotropic
Oncopeltus fasciatus [1]	120/121	115/169	37/37
Cimex lectularius [1,2]	48/49	24/36	30/30
Rhodnius prolixus [1,2]	116/116	28/30	33/33
Acyrthosiphon pisum [3]	79/79	77/77	19/19
Pediculus humanus [2]	12/13	6/8	14/14
Zootermopsis nevadensis	70/70	87/90	150/150
Drosophila melanogaster	60/62	60/68	65/65

[1] Hemiptera: Heteroptera
[2] independent acquisitions of hematophagy [16]
[3] Hemiptera, phloem-feeding

As host plant recognition is only the first step, we further explored whether novel features of the *Oncopeltus* gene set may be directly associated with its diet. We therefore used the CycADS annotation pipeline [116] to reconstruct the *Oncopeltus* metabolic network. The resulting BioCyc metabolism database for *Oncopeltus* ("OncfaCyc") was then compared with those for 26 other insect species in the current ArthropodaCyc collection ([117], http://arthropodacyc.cycadsys.org/), including three other hemipterans: the pea aphid, the green peach aphid, and the kissing bug (Tables 3-4). For a global metabolism analysis, we detected the presence of 1085 Enzyme Commission (EC) annotated reactions with at least one protein in the *Oncopeltus* genome (see also Supplemental Note 6.4, Table S6.9). Among these, 10 enzyme

783 classes (represented by 17 genes) are unique and 17 are missing when compared to
784 the other insects (Table 4, Table S6.10).
785
786

787
788 **Table 3. Hemipteran ArthropodaCyc database summaries.**
789 Overview statistics for the newly created database for *Oncopeltus fasciatus* (Ofas) in
790 comparison with public databases for *Rhodnius prolixus* (Rpro), *Acyrthosiphon pisum*
791 (Apis), and *Myzus persicae* (Mper) available from [117]. Based on OGS v1.1.
792

Species ID	*Ofas*	*Rpro*	*Apis*	*Mper*	*Mper*
Gene set ID	OGS v1.1	RproC1.1 (Built on RproC1 assembly)	OGS v2.1b (Built on Acyr_2.0 assembly)	Clone G006 v1.0	Clone O v1.0
CycADS Database ID	OncfaCyc	RhoprCyc	AcypiCyc v2.1b	Myzpe_G006 Cyc	Myzpe_O Cyc
Total mRNA[1]	19,673	15,437	36,195	24,814	24,770
Pathways	294	312	307	319	306
Enzymatic Reactions	2,192	2,366	2,339	2,384	2,354
Polypeptides	19,820	15,471	36,228	24,849	24,805
Enzymes	3,050	2,660	5,087	4,646	4,453
Compounds	1,506	1,665	1,637	1,603	1,655

793
794 [1] In the BioCyc databases all splice variants are counted in the summary tables for
795 genes.
796

797
798

799
800 **Table 4. Hemipteran ArthropodaCyc annotations of metabolic genes.**
801 Taxonomic abbreviations are as in Table 3.
802

	Ofas	*Rpro*	*Apis*	*Mper*
Global metabolism				
EC[1] present in the genome	1085	1241	1288	1222
EC unique to this genome[2]	10	13	23	5
EC missing only in this genome[2]	17[4]	8	2	6
Amino acid metabolism (KEGG)				
EC present in the genome	169	188	195	185
EC unique to this genome[2]	2	1	6	1
EC missing only in this genome[2]	5	2	0	2
EC unique to this genome[3]	8	10	12	8
EC missing only in this genome[3]	14	5	0	2

803
[1] "EC" refers to the number of proteins, as represented by their unique numerical
designations within the Enzyme Commission (EC) classification system for enzymes
and their catalytic reactions.
[2] in comparison to all other insects from ArthropodaCyc
[3] in comparison among the four hemipterans
[4] includes three EC categories added in OGS v1.2 (see also Table S6.10)
810

We then looked specifically at amino acid metabolism in four hemipterans
representing the three different diets (see also Table S6.11). Among eight EC
annotated enzymes present in the milkweed bug genome but not the other three, we
identified the arginase (E.C. 3.5.3.1) that degrades arginine (Arg) into urea and
ornithine, a precursor of proline (Pro). Given this difference, we extended our
analysis to the entire urea cycle (Fig. 10a). Across all 26 insects present in the
database, we identified three distinct groups (see also Table S6.12): (i) *Oncopeltus*
and six other non-hemipteran insects that are able to degrade Arg but cannot
synthesize it (Fig. 10b); (ii) the other three hemipterans that uniquely can neither
synthesize nor degrade Arg via this cycle (Fig. 10c); and (iii) the other 17 insects
that, with some minor differences, have an almost complete cycle (Fig. 10d). This
suggests that loss of the ability to synthesize Arg may already have occurred at the
base of the Hemiptera, with subsequent, independent loss of Arg degradation capacity
in the aphid and *Rhodnius* lineages. Retention of Arg degradation in *Oncopeltus*
might be linked to the milkweed seed food source, as most seeds are very rich in Arg
[118], and Arg is indeed among the metabolites detected in *Oncopeltus* [119].
However, the monarch butterfly is one of only a handful of species that retains the
complete Arg pathway (Fig. 10d: blue text). Despite a shared food source, these
species may therefore differ in their overall Arg requirements, or – in light of a
possible group benefit of *Oncopeltus* aggregation during feeding [20] – in their
efficiency of Arg uptake.

Other enzymes are also present only in the milkweed bug in comparison with
the other hemipterans (see also Table S6.11). As would be expected, *Oncopeltus*, like
other insects [117], has the ability to degrade tyrosine (Tyr), a pathway that was
uniquely lost in the aphids. Given the variable yields of Tyr from a mucivorous diet
[120], this amino acid needed for cuticle maturation (sclerotization) is jointly
synthesized – and consumed – by the aphid host and its endosymbiotic bacteria [5, 6,
15, 121]. Meanwhile, we find support for the recent nature of milkweed bug lineage-
specific duplications that led to three copies of the Na+/K+ ATPase alpha subunits
whose amino acid substitutions confer increased resistance to milkweed cardenolides
[35, 122]. In the *Oncopeltus* genome, the genes encoding subunits ATPα1B and
ATPα1C occur as a tandem duplication, notably on a scaffold that also harbors one of
the clustered ZF271-like gene expansions (see above).

CONCLUSIONS

The integrated genomic and transcriptomic resources presented here for the milkweed
bug *Oncopeltus fasciatus* (Figs. 2,5) underpin new insights into molecular evolution
and suites of related biological characters within the Hemiptera. The gene structure
trends we identified, with lineage predominating over genome size as a predictor and
with many intron gains in the hemipteroid lineage (Figs. 6,7), offer initial parameters
and hypotheses for the Hemiptera, Coleoptera, and Diptera. Such ordinal-level
parameters can be evaluated against new species' data and also inform customized
pipelines for future automated gene model predictions. At the same time, it will be
interesting to explore the ramifications of hemipteroid intron gains. For example,
while possessing more, small exons brings an increased transcriptional cost, it may

also provide greater scope to generate protein modularity via isoforms based on
alternative exon usage. Furthermore, with the larger genome sizes and lower gene
densities of hemipteroids compare to the well-studied Hymenoptera, it will also be
interesting to see whether and in which direction hemipteroid gene and intron size
may correlate with recombination rates [123].
866
Our analyses also highlight new directions for future experimental research,
building on *Oncopeltus*'s long-standing history as a laboratory model and its active
research community in the modern molecular genetics era (*e.g.*, Fig. 9, [24-26]).
Functional testing will clarify the roles of genes we have identified as unique to the
Hemiptera, including those implicated in chemical protection, bacterial and plant cell
wall metabolism, or encoding wholly novel proteins (Figs. 3,8, see also Supplemental
Note 2.2). Meanwhile, the prominent and species-specific expansions specifically of
ZF271-like zinc fingers (Fig. 4), combined with the absence of the co-repressor KAP-
1 in insects, argues for investigation into alternative possible interaction partners,
which could clarify the nature of these zinc fingers' regulatory role and their binding
targets.
878
One key output of this study is the generation of a metabolism database for
Oncopeltus, contributing to the ArthropodaCyc collection (Table 3). In addition to
comparisons with other species (Fig. 10), this database can also serve as a future
reference for studies that use *Oncopeltus* as an ecotoxicology model species (*e.g.*,
[124]). While we have primarily focused on feeding ecology in terms of broad
comparisons between phytophagy and fluid feeding, *Oncopeltus* is also poised to
support future work on nuances among phytophagous species. Despite its milkweed
diet in the wild, the lab strain of *Oncopeltus* has long been adapted to feed on
sunflower seeds, demonstrating a latent capacity for more generalist phytophagy
[114]. This potential may also be reflected in a larger gustatory receptor repertoire
than would be expected for an obligate specialist feeder (Table 2). Thus, *Oncopeltus*
can serve as a reference species for promiscuously phytophagous pest species such as
the stink bug. Finally, given that we have identified a number of key genes
implicated in life history trade-offs, the genome data represent an important tool to
explore the proximate mechanisms of fundamental aspects of life history evolution in
an organism in which the ultimate explanations for traits such as cardenolide
tolerance, pigmentation, and plasticity in reproduction under environmental variation
have been elucidated in both the laboratory and nature.
897
898
899

METHODS

(More information is available in the supplementary materials, Additional file 1.)

Milkweed bug strain, rearing, and DNA/RNA extraction

The milkweed bug *Oncopeltus fasciatus* (Dallas), Carolina Biological Supply strain (Burlington, North Carolina, USA), was maintained in a laboratory colony under standard husbandry conditions (sunflower seed and water diet, 25 ºC, 12:12 light-dark photoperiod). Voucher specimens for an adult female (record # ZFMK-TIS-26324) and adult male (record # ZFMK-TIS-26325) have been preserved in ethanol and deposited in the Biobank of the Centre for Molecular Biodiversity Research, Zoological Research Museum Alexander Koenig, Bonn, Germany (https://www.zfmk.de/en/biobank).

Genomic DNA was isolated from individual, dissected adults using the Blood & Cell Culture DNA Midi Kit (G/100) (Qiagen Inc., Valencia, California, USA). Total RNA was isolated from individual, dissected adults and from pooled, mixed-instar nymphs with TRIzol Reagent (Invitrogen/ Thermo Fisher Scientific, Waltham, Massachusetts, USA). Dissection improved accessibility of muscle tissue by disrupting the exoskeleton, and gut material was removed.

Genome size calculations (flow cytometry, *k*-mer estimation)

Genome size estimations were obtained by flow cytometry with Hare and Johnston's protocol [125]. Four to five females and males each from the Carolina Biological Supply lab strain and a wild strain (collected from Athens, Georgia, USA; GPS coordinates: 33° 56' 52.8216" N, 83° 22' 38.3484'' W) were measured (see also Supplemental Note 2.1.a). At the bioinformatic level, we attempted to estimate genome size by *k*-mer spectrum distribution analysis for a range of *k*=15 to 34 counted with Jellyfish 2.1.4 [126] and bbmap [127], graphing these counts against the frequency of occurrence of *k*-mers (depth), and calculating genome size based on the coverage at the peak of the distribution (see also Supplemental Note 2.1.b).

Genome sequencing, assembly, annotation, and official gene set overview

Library preparation, sequencing, assembly, and automatic gene annotation were conducted at the Baylor College of Medicine Human Genome Sequencing Center (as in [11, 29]). About 1.1 billion 100-bp paired-end reads generated on an Illumina HiSeq2000s machine were assembled using ALLPATHS-LG [128], from two paired-end (PE) and two mate pair (MP) libraries specifically designed for this algorithm (see also Supplemental Note 1). Three libraries were sequenced from an individual adult male (180- and 500-bp PE, 3-kb MP), with the fourth from an individual adult female (8-10-kb MP). The final assembly (see metrics in Table 1) has been deposited in GenBank (accession GCA_000696205.1).

Automated annotation of protein-coding genes was performed using a Maker 2.0 annotation pipeline [129] tuned specifically for arthropods (see also Supplemental Note 3). These gene predictions were used as the starting point for manual curation via the Apollo v.1.0.4 web browser interface [130], and automatic and manual curations were compiled to generate the OGS (see also Supplemental Note 4). Databases of the genome assembly, Maker automatic gene predictions, and OGS v1.1 are available through the i5K Workspace@NAL [131], and the Ag Data Commons data access system of the United States Department of Agriculture's (USDA) National

950 Agricultural Library as individual citable databases [132-134]. The current version of
951 the gene set, OGS v1.2, will be deposited in NCBI under accession number XXX.
952

Repeat content analysis
Repetitive regions were identified in the *Oncopeltus* genome assembly with
RepeatModeler Open-1.0.8 [135] based on a species-specific repeat library generated
de novo with RECON [136], RepeatScout [137], and Tandem Repeats Finder [138].
Then, RepeatMasker Open-4.0 [139] was used to mask repeat sequences based on the
RepeatModeler library. Given the fragmented nature of the assembly, we attempted
to fill and close assembly gaps by sequencing additional material, generating long
reads with single molecule real time sequencing on a PacBio RS II machine (34
SMRT cells to an expected coverage of 8x, see also Supplemental Note 2.3). Gap
filling on the Illumina assembly scaffolds was performed with PBJelly version
13.10.22, and the resulting assembly was used for repeat content estimation and
comparison with *Cimex lectularius* and *Acyrthosiphon pisum.*
965

Transcriptome resources
Total RNA from three distinct life history samples (pooled, mixed-instar nymphs; an
adult male; an adult female) was also sequenced on an Illumina HiSeq2000s machine,
producing a total of 72 million 100-bp paired-end reads (see also Supplemental Note
1.3, Table S1.1). These expression data were used to support the generation of the
OGS at different stages of the project: as input for the evidence-guided automated
annotation with Maker 2.0 (see also Supplemental Note 3), as expression evidence
tracks in the Apollo browser to support the community curation of the OGS, and,
once assembled into a *de novo* transcriptome, as a point of comparison for quality
control of the OGS.
976

The raw RNA-seq reads were pre-processed by filtering out low quality bases
(phred score <30) and Truseq adapters with Trimmomatic-0.30. Further filtering
removed ribosomal and mitochondrial RNA sequences with Bowtie 2 [140], based on
a custom library built with all hemipteran ribosomal and mitochondrial RNA
accessions from NCBI as of 7[th] February 2014 (6,069 accessions). The pooled,
filtered reads were mapped to the genome assembly with Tophat2-PE on CyVerse
[141]. A second set of RNA-seq reads from an earlier study ("published adult"
dataset, [35]) was also filtered and mapped in the same fashion, and both datasets
were loaded into the *Oncopeltus* Apollo instance as evidence tracks (under the track
names "pooled RNA-seq - cleaned reads" and "RNA-seq raw PE reads Andolfatto et
al", respectively).
988

Additionally, a *de novo* transcriptome was generated from our filtered RNA-
seq reads (pooled from all three samples prepared in this study) using Trinity [142]
and TransDecoder [143] with default parameters. This transcriptome is referred to as
"i5K", to distinguish it from a previously published maternal and early embryonic
transcriptome for *Oncopeltus* (referred to as "454", [34]). Both the i5K and 454
transcriptomes were mapped to the genome assembly with GMAP v. 2014-05-15 on
CyVerse. These datasets were also loaded into the Apollo browser as evidence tracks
to assist in manual curation.
997

Life stage specific expression analyses

Transcript expression of the OGS v1.1 genes was estimated by running RSEM2 [144] on the filtered RNA-seq datasets for the three postembryonic stages against the OGS v1.1 cDNA dataset. Transcript expression was then based on the transcripts per million (TPM) value. The TPM values were processed by adding a value of 1 (to avoid zeros) and then performing a log2-transformation. The number of expressed genes per RNA-seq library was compared for TPM cutoffs of >1, >0.5, and >0.25. For this first-pass expression assessment, a >0.25 cutoff was chosen, which reduced the number of expressed genes by 6.6% compared to the first analysis, while the other TPM cutoffs were deemed too restrictive (reducing the expressed gene set by 10.3% and 16.6%, respectively, with the >0.5 and >1 cutoffs). This analysis was also applied to the "published adult" dataset [35]. To include embryonic stages in the comparison, transcripts from the 454 transcriptome were used as blastn queries against the OGS v1.1 cDNA dataset (cutoff e-value $<10^{-5}$). The results from all datasets were converted to binary format to generate Venn diagrams (Fig. 2b).

Protein gene orthology assessments via OrthoDB and BUSCO analyses

These analyses follow previously described approaches [41, 42]. See Supplemental Note 6.1 for further details.

Global transcription factor identification

Likely transcription factors (TFs) were identified by scanning the amino acid sequences of predicted protein-coding genes for putative DNA binding domains (DBDs), and when possible, the DNA binding specificity of each TF was predicted using established procedures [56]. Briefly, all protein sequences were scanned for putative DBDs using the 81 Pfam [145] models listed in Weirauch and Hughes [146] and the HMMER tool [147], with the recommended detection thresholds of Per-sequence Eval < 0.01 and Per-domain conditional Eval < 0.01. Each protein was classified into a family based on its DBDs and their order in the protein sequence (e.g., bZIPx1, AP2x2, Homeodomain+Pou). The resulting DBD amino acid sequences were then aligned within each family using Clustal Omega [148], with default settings. For protein pairs with multiple DBDs, each DBD was aligned separately. From these alignments, the sequence identity was calculated for all DBD sequence pairs (*i.e.*, the percent of amino acid residues that are identical across all positions in the alignment). Using previously established sequence identity thresholds for each family [56], the predicted DNA binding specificities were mapped by simple transfer. For example, the DBD of OFAS001246-RA is 98% identical to the *Drosophila melanogaster* Bric a Brac 1 (Bab1) protein. Since the DNA binding specificity of Bab1 has already been experimentally determined, and the cutoff for the Pipsqueak family TFs is 85%, we can infer that OFAS001246-RA will have the same binding specificity as *Drosophila* Bab1.

RNA interference

Double-stranded RNA (dsRNA) was designed to target the final, unique exon of the *broad* isoforms *Z2, Z3*, and *Z4*. A portion of the coding sequence for the zinc finger region from these exons (179 bp, 206 bp, and 216 bp, respectively) was cloned into a plasmid vector and used as template for *in vitro* RNA synthesis, using the gene-specific primer pairs: Of-Z2_fwd: 5′-ATGTGGCAGACAAGCATGCT-3′; Of-Z2_rev: 5′-CTAAAATTTGACATCAGTAGGC-3′; Of-Z3_fwd: 5′-ccttctcctgttactactcac-3′; Of-Z3_rev: 5′-ttatatgggcggctgtccaa-3′; Of-Z4_fwd: 5′-

AACACTGACCTTGGTTACACA-3′; Of-Z4_rev: 5′-
TAGGTGGAGGATTGCTAAAATT-3′. Two separate transcription reactions (one
for each strand) were performed using the Ambion MEGAscript kit (Ambion, Austin,
Texas, USA). The reactions were purified by phenol/chloroform extraction followed
by precipitation as described in the MEGAscript protocol. The separate strands were
re-annealed in a thermocycler as described previously [31]. Nymphs were injected
with a Hamilton syringe fitted with a 32-gauge needle as described [53]. The
concentration of *Of-Z2*, *Of-Z3* and *Of-Z4* dsRNA was 740 ng/µl, 1400 ng/µl, and
1200 ng/µl, respectively. All nymphs were injected within 8 hours of the molt to the
fourth (penultimate juvenile) instar (n ≥12 per treatment: see Fig. 9). Fore- and
hindwings were then dissected from adults and photographed at the same scale as
wings from wild type, uninjected controls.
1060
CycADS annotation and OncfaCyc database generation
We used the Cyc Annotation Database System (CycADS, [116]), an automated
annotation management system, to integrate protein annotations from different
sources into a Cyc metabolic networks reconstruction that was integrated into the
ArthropodaCyc database. Using our CycADS pipeline, *Oncopeltus fasciatus* proteins
from the official gene set OGS v1.1 were annotated using different methods –
including KAAS [149], PRIAM [150], Blast2GO [151, 152], and InterProScan with
several approaches [153] – to obtain EC and GO numbers. All annotation
information data were collected in the CycADS SQL database and automatically
extracted to generate appropriate input files to build or update BioCyc databases [154]
using the Pathway Tools software [155]. The OncfaCyc database, representing the
metabolic protein-coding genes of *Oncopeltus*, was thus generated and is now
included in the ArthropodaCyc database, a collection of arthropod metabolic network
databases ([117], http://arthropodacyc.cycadsys.org/).
1075
1076
1077   **FIGURE LEGENDS**
1078
Fig. 1. The large milkweed bug, *Oncopeltus fasciatus*, shown in its phylogenetic
and environmental context.
(a) Species tree of selected Hemiptera with genomic and transcriptomic resources,
based on phylogenetic analyses and divergence time estimates in [3]. Species marked
with an asterisk (*) have published resources; those with the appellation "i5K" are
part of a current pilot project supported by the Baylor College of Medicine Human
Genome Sequencing Center and the National Agricultural Library of the USDA.
Note that recent analyses suggest the traditional infraorder Cimicomorpha, to which
Rhodnius and *Cimex* belong, may be paraphyletic [16].
(b-c) Milkweed bugs on their native food source, the milkweed plant: gregarious
nymphs of different instars on a milkweed seed pod (b), and pale, recently eclosed
adults and their shed exuvia (c). Images were taken at Avalon Park and Preserve,
Stony Brook, New York, USA, courtesy of Deniz Erezyilmaz, used with permission.
(d) Individual bugs, shown from left to right: first instar nymphs (ventral and dorsal
views) and adults (dorsal and lateral views); images courtesy of Kristen Panfilio
(nymphs) and Jena Johnson (adults), used with permission. The arrow labels the
labium (the "straw"), part of the hemipteran mouthpart anatomy adapted for feeding
by piercing and sucking.
1097

Fig. 2. Comparisons of the official gene set and transcriptomic resources.
(a) Area-proportional Venn diagram comparing the OGS v1.1 ("OGS"), a Trinity *de novo* transcriptome from the three post-embryonic RNA-seq samples ("i5K"), and the maternal and embryonic transcriptome from 454 data ("454" [34]). Sample sizes and the fraction of each transcriptome represented in the OGS are indicated (for the 454 dataset, only transcripts with homology identification were considered). The unique fraction of each set is also specified (%). Dataset overlaps were determined by blastn (best hit only, e-value $<10^{-9}$).
(b) Four-set Venn diagram representation of OGS v1.1 gene model expression across four different life history samples. Values are counts of gene models, with percentages also given for the largest subsets. Note that the "Embryo/Maternal" sample derives from 454 pyrosequencing data and therefore has a smaller data volume than the other samples, which were generated with Illumina sequencing.

Fig. 3. Orthology comparisons and phylogenetic placement of *Oncopeltus fasciatus* among other Arthropoda.
(a) Comparisons of protein-coding gene content in 12 arthropod species, with the Hemiptera highlighted in red text. The bar chart shows the number of proteins per conservation level (see legend), based on OrthoDB orthology clustering analyses. To the left is a maximum likelihood phylogeny based on concatenation of 395 single-copy orthologs (all nodes have 100% support unless otherwise noted; branch length unit is substitutions per site). The inset pie chart shows the proportion of proteins per conservation level in *Oncopeltus* (*"Ofas"*). See also Supplemental Note 6.1.
(b) Proportion of *Oncopeltus* proteins that have expression and/or curation validation support per conservation level (same color legend as in (a)). Expression support is based on the life history stage data in Fig. 2b.
(c) Protein orthology data evaluated by taxonomic grouping, based on the OrthoDB v8 i5K "Insecta" analysis with 64 species, a subset of the recently released OrthoDB v9 (http://www.orthodb.org): Hemiptera (red, 8 species); other hemimetabolous species (paraphyletic, yellow, 6 species); Holometabola (purple, 50 species). Values are given for both orthogroups (black text, defined in [42]) and protein-coding genes (blue text). As analyses that require all species to be represented in a given orthogroup are limited by the quality of every species' OGS, the cutoff for orthogroup presence in a given Venn diagram set was rather that roughly half of all relevant species are included, and strictly no species from a different set are permitted. Set [A] contains ≥10 hemimetabolous species (allowing for 4-8 Hemiptera and 2-6 other hemimetabolous species). Set [B] contains ≥4 Hemiptera and ≥25 Holometabola. Set [C] contains ≥2 other hemimetabolous species and for the Holometabola at least one representative from each of the Hymenoptera, Coleoptera, Lepidoptera, and Diptera. Analyses based on OGS v1.1.

Fig. 4. Distribution of transcription factor families across insect genomes.
(a) Heatmap depicting the abundance of transcription factor (TF) families across 17 insect genomes (Hemiptera highlighted in red text), with *Daphnia* as an outgroup. Each entry indicates the number of TF genes for the given family in the given genome, based on the presence of predicted DNA binding domains (see Methods). The color key has a log (base 2) scale (light blue means the TF family is completely absent). Values are in Supplementary Table S6.2.
(b) Bar graph showing the number of proteins of each of the two most abundant TF families, homeodomains and C2H2 zinc fingers (ZFs), per species. Solid lines

demarcate insect orders: Hemiptera (Hemipt.), Hymenoptera (Hym.), Coleoptera
(Col.), and Diptera (Dipt.). The dashed line demarcates the dipteran family Culicidae
(mosquitoes).
(c) Proportions of *Oncopeltus* homeodomain (HD) and C2H2 zinc finger proteins
with orthology assignment (predicted DNA binding specificity) and/or manual
curation. "Classified" refers to automated classification of a protein to a TF family,
but without a specific orthology assignment.
(d) Maximum likelihood phylogeny of representative subsets of the zinc finger 271-
like family in *Oncopeltus* (49 proteins, blue text) and protein accessions retrieved
from GenBank for the pea aphid (55 proteins, black text) as well as chelicerate (red
text) and holometabolan (yellow text) outgroups (16 proteins, 7 species). Gaps were
removed during sequence alignment curation, with default pipeline settings [156]. All
nodes have ≥50% support. Key nodes are circled for the distinct clades containing all
aphid or all *Oncopeltus* proteins (82% support each), and for each 'core' clade
comprised exclusively of proteins from each species (97% and 100%, respectively;
triangles shown to scale for branch length and number of clade members). Branch
length unit is substitutions per site.
Analyses based on OGS v1.1.
1166
Fig. 5. Comparison of repeat content estimations.
(a) Comparison of total repetitive content among insect genomes. The three values
for *Oncopeltus* are shown (in ascending order: original Illumina assembly, gap-filled
assembly, Illumina-PacBio hybrid estimate). Values for the three hemipterans labeled
in red text are from RepeatModeler (gold bars for the pea aphid and bed bug; blue and
gold bars for *Oncopeltus*). All other values are from the respective genome papers,
including a second value corresponding to the published repeat content for the first
version of the aphid genome [5, 9, 112, 157-162]. Species abbreviations as in Fig. 4
(compare panels a and b), and additionally: *Nlug, Nilaparvata lugens; Lmig, Locusta*
migratoria; Bmor, Bombyx mori; Aalb, Aedes albopictus.
(b) Comparison of repetitive element categories between three hemipteran genomes,
based on results from RepeatModeler. Here we present assembly coverage as actual
sequence length (Mb) to emphasize the greater repeat content in *Oncopeltus* (based on
the gap-filled assembly: see also Supplemental Note 2.3).
1181
Figure 6. Trends in gene structure show hemipteroid-specific tendencies.
(a) Trends in protein size, exon size, and exon number are shown for a highly
conserved set of genes encoding large proteins of diverse functional classes ("gold
standard", curated gene set). Median values are plotted. Sample sizes are indicated
for each species, with 11 genes for which orthologs were evaluated in all species.
Where it was not possible to analyze all 30 genes for a given species, equal sampling
was done across the range of protein sizes of the complete dataset, based on the
Cimex ortholog sizes (1:1:1 sampling from big:medium:small subcategories of 10
genes each). See also Supplemental Note 6.3.
(b) Box plot representations of coding sequence exon size (aa) for two species from
each of three insect orders, based on datasets of unique coding sequence exons (one
isoform per gene) and excluding terminal exons <10 aa (as most of those exons may
rather be UTRs or a small placeholder N-terminal exon, as a byproduct of the Maker
gene annotation pipeline's requirement to create nominally complete protein coding
genes with in-frame start codons). Only manually curated gene models were
considered for *Oncopeltus* and the other recent i5K species; the entire OGS was used

for *Tribolium* and *Drosophila*. For clarity, outliers are omitted; whiskers represent
1.5× the value of the Q3 (upper) or Q2 (lower) quartile range. MAD, median absolute
deviation.
Species are represented by their four-letter taxonomic abbreviations, with their ordinal
relationships given below the phylogeny in panel (a): Hemip., Hemiptera; Thys.,
Thysanoptera; Col., Coleoptera; Dipt., Diptera. Species abbreviations as in Fig. 4 and
additionally: *Gbue, Gerris buenoi*; *Focc, Frankliniella occidentalis*; *Agla,*
Anoplophora glabripennis; Ccap, Ceratitis capitata.
1206

Fig 7. Splice site evolution correlates with both lineage and, independently,
genome size.
Splice site changes are shown for *hemocytin* (blue text), *Tenascin major* (*Ten-m*,
turquoise text), and *UDP-galactose 4′-epimerase* (brown text), mapped onto a species
tree of eight insects. Patterns of splice site evolution were inferred based on the most
parsimonious changes that could generate the given pattern within a protein sequence
alignment of all orthologs (see also Supplemental Note 6.3 for complete methodology
and data sources). In instances where an equal number of lineage specific gains or
losses was possible, we remained agnostic and present a range for the ancestral
number of splice sites indicated at the base of the tree, where the bracketed number
indicates how many ancestral positions are still retained in all species. Along each
lineage, subsequent changes are indicated in brackets, with the sign indicating gains
(+) or losses (-). Values shown to the right are species-specific changes. Note that
the values shown between the *D. melanogaster* and *T. castaneum* lineages denote
changes that have occurred independently in both. Colored boxes highlight the
largest sources of change, as indicated in the legend and discussed in the main text.
Species are represented by their four-letter abbreviations (as defined in Fig. 6), and
the estimated genome sizes are indicated parenthetically (measured size: [11, 29, 73,
161, 163]; draft assembly size: GenBank Genome IDs 14741 and 17730). Divergence
times are shown in gray and given in millions of years [3]. Abbreviations as in Figs.
4,6, and: Col., Coleoptera; Dipt., Diptera; Hemip., Hemiptera; Hemipt., hemipteroid
assemblage (including *F. occidentalis*); n.d., no data.
1229

Fig. 8. Lateral gene transfer introduction and subsequent evolution within the
Hemiptera for mannosidase-encoding genes.
(a) Species tree summary of evolutionary events. Stars represent the original LGT
introduction and subsequent copy number gains (see legend).
(b) Maximum likelihood phylogeny of mannosidase proteins, including bacterial
sequences identified among the best GenBank blastp hits for *Oncopeltus* and
Halyomorpha (accession numbers as indicated, and for "Other bacteria" are:
ACB22214.1, AEE17431.1, AEI12929.1, AEO43249.1, AFN74531.1, CDM56239.1,
CUA67033.1, KOE98396.1, KPI24888.1, OAN41395.1, ODP26899.1, ODS11151.1,
OON18663.1, PBD05534.1, SIR54690.1, WP096035621.1, YP001327394.1). All
nodes have ≥50% support from 500 bootstrap replicates [164]. Triangles are shown
to scale for branch length and number of clade members; branch length unit is
substitutions per site.
(c) Manually curated protein sequence alignment for the N-terminal region, showing
the position of splice sites ("|" symbol), where one position is ancestral and present in
all paralogs of a given species (magenta), and one position occurs in a subset of
paralogs and is presumed to be younger (cyan, note the position is within the 5′ UTR
in the case of *Halyomorpha*). Residues highlighted in yellow are conserved between

the two species. The *Oncopeltus* paralog represented in the OGS as OFAS017153-RA is marked with an asterisk to indicate that this version of the gene model is incomplete and lacks the initial exon (gray text in the alignment). For clarity, only the final three digits of the *Halyomorpha* GenBank accessions are shown (full accessions: XP_014289XXX).

Fig. 9. Isoform-specific RNAi based on new genome annotations affects the molting and cuticle identity gene *broad*.
(a) Genomic organization of the cuticle identity gene *broad*. The regions used as template to generate isoform-specific dsRNA are indicated (red asterisks: the final, unique exons of each isoform). Previous RNAi studies targeted sequence within exons 1-5 that is shared among all isoforms (dashed red box, [94]).
(b) Knock down of the *Oncopeltus Z2* or *Z3 broad* isoforms at the onset of the penultimate instar resulted in altered nymphal survival and morphogenesis that was reflected in the size and proportion of the fore and hind wings at the adult stage (upper and lower images, respectively, shown to the same scale for all wings). We did not detect any effect on the wing phenotype when targeting the *Z4*-specific exon, demonstrating the specificity of the zinc finger coding region targeted by RNAi. Experimental statistics are provided in the figure inset, including for the buffer-injected negative control.

Fig. 10. Comparison of the urea cycle of *Oncopeltus* with 26 other insect species.
(a) Detailed diagram of the urea cycle (adapted from KEGG).
(b) Group of 7 species, including *Oncopeltus*, for which Arg degradation via arginase (3.5.3.1), but not synthesis, is possible.
(c) Group of 3 species for which neither the degradation nor synthesis of arginine via the urea cycle is possible (all 3 other hemipterans in this analysis).
(d) Group of 17 species sharing a complete (or almost complete) urea cycle. Hemiptera are identified in red text and the milkweed-feeding monarch butterfly is in blue text. Enzyme names corresponding to EC numbers: 1.5.1.2 = pyrroline-5-carboxylate reductase; 1.14.13.39 = nitric-oxide synthase; 2.1.3.3 = ornithine carbamoyltransferase; 2.6.1.13 = ornithine aminotransferase; 3.5.3.1 = arginase; 4.3.2.1 = argininosuccinate lyase; 6.3.4.5 = argininosuccinate synthase. Analyses based on OGS v1.1.

TABLES (see above within relevant manuscript sections)

Table 1. *Oncopeltus fasciatus* genome metrics.

Table 2. Numbers of chemoreceptor genes/proteins per family in selected insect species.

Table 3. Hemipteran ArthropodaCyc database summaries.

Table 4. Hemipteran ArthropodaCyc annotations of metabolic genes.

DECLARATIONS

ADDITIONAL FILES

Additional file 1: Supplementary figures, tables, methods, and other text. (PDF)

Additional file 2: Large supporting tables. (XLSX)

Additional file 3: Chemoreceptor sequences in FASTA format. (TXT)

ACKNOWLEDGEMENTS

We thank Dorith Rotenberg (Kansas State University, currently North Carolina State University, USA), Abderrahman Khila on behalf of the Water Strider Genome Consortium (Institute of Functional Genomics and École Normale Supérieure de Lyon, France), and Michael Sparks (Agricultural Research Service, United States Department of Agriculture, USA) for generously making available the unpublished genome assemblies of the fellow hemipteroid i5K species *Frankliniella occidentalis*, *Gerris buenoi*, and *Halyomorpha halys*, respectively, for use in specific analyses presented here. Similarly, we thank Hans Kelstrup and Lynn Riddiford (Janelia Farm Research Campus, HHMI, USA) for sharing unpublished data on *Of-E75A* RNAi. We thank George Coupland (Max Planck Institute for Plant Breeding Research, Cologne, Germany) as well as Lisa Czaja, Kurt Steuber, and Bruno Huettel (Max Planck Genome Centre Cologne, Germany) for conducting the PacBio sequencing and providing support with data handling. We also thank Oliver Niehuis (Albert Ludwig University, Freiburg, Germany) and Alexander Klassmann (University of Cologne, Germany) for discussions on *k*-mer and gene structure analyses, respectively, Sarah Kingan (University of Rochester, USA) for assistance with LGT phylogenies, as well as Jeanne Wilbrandt (Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany) for comments on the manuscript.

FUNDING

Funding for genome sequencing, assembly and automated annotation was provided by the National Institutes of Health (NIH) grant U54 HG003273 (NHGRI) to RAG. The i5K pilot project (https://www.hgsc.bcm.edu/arthropods) assisted in sequencing of the *Oncopeltus fasciatus* genome. We also acknowledge funding for the project from German Research Foundation (DFG) grants PA 2044/1-1 and SFB 680 project A12 to KAP. Support for specific analyses was provided by the Swiss National Science Foundation with grant 31003A_143936 to EMZ and PP00P3_170664 to RMW; the European Research Council grant ERC-CoG #616346 to AK; DFG grant SFB 680 project A1 to SiR; the National Science Foundation with grant US NSF DEB1257053 to JHW; and by NIH grants 5R01GM080203 (NIGMS) and 5R01HG004483 (NHGRI) and by the Director, Office of Science, Office of Basic Energy Sciences, U.S. Department of Energy, Contract No. DE-AC02-05CH11231 to MCMT.

AVAILABILITY OF DATA AND MATERIALS

All sequence data are publically available at the NCBI, bioproject number PRJNA229125. In addition, gene models and a browser are available at the National Agricultural Library ([132-134], https://i5k.nal.usda.gov/Oncopeltus_fasciatus).

AUTHOR'S CONTRIBUTIONS

KAP and StR conceived, managed, and coordinated the project. KAP and SK provided specimens for sequencing and performed DNA and RNA extractions. StR, SD, SLL, HC, HVD, HD, YH, JQ, SCM, DSTH, KCW, DMM, and RAG constructed

libraries and performed sequencing. StR, SCM, and DSTH performed the genome assembly and automated gene prediction. IMVJ, JSJ, and PJM analyzed genome size. IMVJ, VK, PH, and KAP contributed to repetitive content analyses. AD, RR, JHW, KAP, and SK performed bacterial scaffold detection and LGT analyses. MCMT developed Apollo software. KAP, IMVJ, MCMT, CPC, C-YL, and MFP implemented Apollo-based manual curation. KAP, IMVJ, JBB, DE, YS, HMR, DA, CGCJ, BMIV, EJD, CSB, C-CC, Y-TC, ADC, AGC, AJJC, PKD, EMD, CGE, MF, NG, TH, Y-MH, ECJ, TEJ, JWJ, AK, ML, MRL, H-LL, YL, SRP, LP, MP, PNR, RRP, SiR, LS, MES, JS, ES, JNS, OT, LT, MVDZ, SV, and AJR participated in manual curation and contributed to the Supplemental Notes. IMVJ, KAP, DSTH, M-JMC, CPC, C-YL, and MFP performed curation quality control and generated the OGS. IMVJ, KAP, and JBB generated *de novo* transcriptomes and performed life history stage expression analyses. RMW, PI, KAP, and EMZ performed orthology and phylogenomic analyses. MTW, KAP, IMVJ, PH, and BMIV performed transcription factor analyses. EJD conducted analyses of DNA methylation. KAP, PH, and RJS contributed to comparative analyses of gene structure. DE conducted the RNAi experiments. SC, PB-P, GF, and NP generated and performed comparative analyses on the OncfaCyc database. KAP, IMVJ, JBB, DE, YS, SC, HMR, and MTW wrote the manuscript. KAP, IMVJ, JBB, DE, YS, SC, HMR, MFP, RMW, PI, MTW, StR, PJM, and AK edited the manuscript. IMVJ and KAP organized the Supplementary Materials. All authors approved the final manuscript.

COMPETING INTERESTS
The authors declare that they have no competing interests.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE
Not applicable.

AUTHOR DETAILS
[1] Institute for Zoology: Developmental Biology, University of Cologne, Zülpicher Str. 47b, 50674 Cologne, Germany
[2] School of Life Sciences, University of Warwick, Gibbet Hill Campus, Coventry CV4 7AL, UK
[3] Department of Biological Sciences, University of Cincinnati, Cincinnati, Ohio 45221, USA
[4] Department of Biochemistry and Cell Biology and Center for Developmental Genetics, Stony Brook University, Stony Brook, New York 11794, USA
[5] Department of Biological Sciences, Wellesley College, 106 Central St., Wellesley, Massachusetts 02481, USA
[6] Univ Lyon, INSA-Lyon, INRA, BF2I, UMR0203, F-69621, Villeurbanne, France
[7] Current address: LSTM, Laboratoire des Symbioses Tropicales et Méditerranéennes, INRA, IRD, CIRAD, SupAgro, University of Montpellier, Montpellier, France
[8] Department of Entomology, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA
[9] National Agricultural Library, Beltsville, Maryland 20705, USA
[10] Department of Genetic Medicine and Development and Swiss Institute of Bioinformatics, University of Geneva, Geneva 1211, Switzerland
[11] Current address: Department of Ecology and Evolution, University of Lausanne, Lausanne 1015, Switzerland
[12] Center for Autoimmune Genomics and Etiology, Division of Biomedical Informatics, and Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Department of Pediatrics, College of Medicine, University of Cincinnati, Cincinnati, Ohio 45229, USA
[13] Human Genome Sequencing Center, Department of Human and Molecular Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, Texas 77030, USA

[14] Current address: Department of Genome Sciences, University of Washington School of
Medicine, Seattle, Washington 98195, USA
[15] Current address: Howard Hughes Medical Institute, University of Washington, Seattle,
Washington 98195, USA
[16] Department of Biology, University of Rochester, Rochester, New York 14627, USA
[17] Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE Leiden, Netherlands
[18] Max Planck Institute for Chemical Ecology, Hans-Knöll Strasse 8, 07745 Jena, Germany
[19] Department of Biochemistry and Genomics Aotearoa, University of Otago, Dunedin 9054, New
Zealand
[20] School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK
[21] Institut de Génomique Fonctionnelle de Lyon, Université de Lyon, Université Claude Bernard
Lyon 1, CNRS UMR 5242, École Normale Supérieure de Lyon, 46 Allée d'Italie, 69364 Lyon,
France
[22] Department of Ecology, Evolution and Behavior, The Alexander Silberman Institute of Life
Sciences, The Hebrew University of Jerusalem, Edmond J. Safra Campus, Givat Ram 91904,
Jerusalem, Israel
[23] Department of Entomology/Institute of Biotechnology, College of Bioresources and
Agriculture, National Taiwan University, Taipei, Taiwan
[24] Current address: School of Life Sciences, Rochester Institute of Technology, Rochester, New
York 14623, USA
[25] Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street,
Cambridge, Massachusetts 02138, USA
[26] Department of Molecular and Cellular Biology, Harvard University, 26 Oxford Street,
Cambridge, Massachusetts 02138, USA
[27] Department of Biological Sciences, Wayne State University, Detroit, Michigan 48202, USA
[28] Institute for Genetics, University of Cologne, Zülpicher Straße 47a, 50674 Cologne, Germany
[29] Department of Entomology, Texas A&M University, College Station, Texas 77843, USA
[30] CECAD, University of Cologne, Cologne, Germany
[31] Department of Entomology and Program in Molecular & Cell Biology, University of Maryland,
College Park, Maryland 20742, USA
[32] Department of Entomology, University of Georgia, 120 Cedar St., Athens, Georgia 30602, USA
[33] Environmental Genomics and Systems Biology Division, Lawrence Berkeley National
Laboratory, Berkeley, California, USA
[34] Department of Entomology, College of Agriculture, Food and Environment, University of
Kentucky, Lexington, Kentucky 40546, USA
[35] Department of Biology, University of Hawai'i at Mānoa, Honolulu, Hawaii 96822, USA
[36] Current address: Department of Evolutionary Genetics, Max-Planck-Institut für
Evolutionsbiologie, August-Thienemann-Straße 2, 24306 Plön, Germany
[37] Current address: Earthworks Institute, 185 Caroline Street, Rochester, New York 14620, USA
[38] Centro de Bioinvestigaciones, Universidad Nacional del Noroeste de Buenos Aires, Argentina
[39] Current address: Department of Biotechnology, Central university of Rajasthan (CURAJ), NH-
8, Bandarsindri, Ajmer- 305801, India
[40] Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn,
Germany
[41] Current address: Department of Zoology, University of Cambridge, Cambridge CB2 3DT, UK
[42] Centro Regional de Estudios Genómicos, Facultad de Ciencias Exactas, Universidad Nacional
de La Plata, La Plata, Argentina
1444
1445
1446

REFERENCES

1. Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, Seppey M, Loetscher A, Kriventseva EV: **OrthoDB v9.1: cataloging evolutionary and functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs.** *Nucleic Acids Res* 2017, **45:**D744-D749.

2. Huang DY, Bechly G, Nel P, Engel MS, Prokop J, Azar D, Cai CY, van de Kamp T, Staniczek AH, Garrouste R, et al: **New fossil insect order Permopsocida elucidates major radiation and evolution of suction feeding in hemimetabolous insects (Hexapoda: Acercaria).** *Sci Rep* 2016, **6:**23004.

3. Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG, et al: **Phylogenomics resolves the timing and pattern of insect evolution.** *Science* 2014, **346:**763-767.

4. Grimaldi D, Engel MS: *Evolution of the Insects.* Cambridge: Cambridge University Press; 2005.

5. The International Aphid Genomics Consortium: **Genome sequence of the pea aphid** *Acyrthosiphon pisum* *PLoS Biol* 2010, **8:**e1000313.

6. Mathers TC, Chen Y, Kaithakottil G, Legeai F, Mugford ST, Baa-Puyoulet P, Bretaudeau A, Clavijo B, Colella S, Collin O, et al: **Rapid transcriptional plasticity of duplicated gene clusters enables a clonally reproducing aphid to colonise diverse plant species.** *Genome Biol* 2017, **18:**27.

7. Wenger JA, Cassone BJ, Legeai F, Johnston JS, Bansal R, Yates AD, Coates BS, Pavinato VA, Michel A: **Whole genome sequence of the soybean aphid, Aphis glycines.** *Insect Biochem Mol Biol* 2017.

8. Sloan DB, Nakabachi A, Richards S, Qu J, Murali SC, Gibbs RA, Moran NA: **Parallel histories of horizontal gene transfer facilitated extreme reduction of endosymbiont genomes in sap-feeding insects.** *Mol Biol Evol* 2014, **31:**857-871.

9. Xue J, Zhou X, Zhang C-X, Yu L-L, Fan H-W, Wang Z, Xu H-J, Xi Y, Zhu Z-R, Zhou W-W, et al: **Genomes of the rice pest brown planthopper and its endosymbionts reveal complex complementary contributions for host adaptation.** *Genome Biol* 2014, **15:**521.

10. Mesquita RD, Vionette-Amaral RJ, Lowenberger C, Rivera-Pomar R, Monteiro FA, Minx P, Spieth J, Carvalho AB, Panzera F, Lawson D, et al: **Genome of *Rhodnius prolixus*, an insect vector of Chagas disease, reveals unique adaptations to hematophagy and parasite infection.** *Proc Natl Acad Sci USA* 2015, **112:**14936-14941.

11. Benoit JB, Adelman ZN, Reinhardt K, Dolan A, Poelchau M, Jennings EC, Szuter EM, Hagan RW, Gujar H, Shukla JN, et al: **Unique features of a global human ectoparasite identified through sequencing of the bed bug genome.** *Nat Commun* 2016, **7:**10165.

12. Rosenfeld JA, Reeves D, Brugler MR, Narechania A, Simon S, Durrett R, Foox J, Shianna K, Schatz MC, Gandara J, et al: **Genome assembly and geospatial phylogenomics of the bed bug *Cimex lectularius*.** *Nat Commun* 2016, **7:**10164.

13. Sparks ME, Shelby KS, Kuhar D, Gundersen-Rindal DE: **Transcriptome of the invasive brown marmorated stink bug, Halyomorpha halys (Stal) (Heteroptera: Pentatomidae).** *PLoS One* 2014, **9:**e111646.

14. Ioannidis P, Lu Y, Kumar N, Creasy T, Daugherty S, Chibucos MC, Orvis J, Shetty A, Ott S, Flowers M, et al: **Rapid transcriptome sequencing of an invasive pest, the brown marmorated stink bug, *Halyomorpha halys*.** *BMC Genomics* 2014, **15:**738.

15. Wilson ACC, Ashton PD, Charles H, Colella S, Febvay G, Jander G, Kushlan PF, Macdonald SJ, Schwartz JF, Thomas GH, Douglas AE: **Genomic insight into the amino acid relations of the pea aphid, Acyrthosiphon pisum, with its symbiotic bacterium Buchnera aphidicola.** *Insect Molecular Biology* 2010, **19 Suppl 2:**249-258.

16. Li H, Leavengood JM, Jr., Chapman EG, Burkhardt D, Song F, Jiang P, Liu J, Zhou X, Cai W: **Mitochondrial phylogenomics of Hemiptera reveals adaptive innovations driving the diversification of true bugs.** *Proc Biol Sci* 2017, **284**.

17. Eichler S, Schaub GA: **Development of symbionts in triatomine bugs and the effects of infections with trypanosomatids.** *Exp Parasitol* 2002, **100:**17-27.

1503   18.   Matsuura Y, Kikuchi Y, Hosokawa T, Koga R, Meng X-Y, Kamagata Y, Nikoh N,
1504   Fukatsu T: **Evolution of symbiotic organs and endosymbionts in lygaeid stinkbugs.**
1505   *The ISME Journal* 2012, **6:**397–409.
1506   19.   Berenbaum MR, Miliczky E: **Mantids and milkweed bugs - efficacy of aposematic
1507   coloration against invertebrate predators.** *American Midland Naturalist* 1984, **111:**64-
1508   68.
1509   20.   Burdfield-Steel ER, Shuker DM: **The evolutionary ecology of the Lygaeidae.** *Ecol Evol*
1510   2014, **4:**2278-2301.
1511   21.   Lawrence PA: **Mitosis and the cell cycle in the metamorphic moult of the milkweed
1512   bug Oncopeltus fasciatus; a radioautographic study.** *J Cell Sci* 1968, **3:**391-404.
1513   22.   Chipman AD: **Oncopeltus fasciatus as an evo-devo research organism.** *Genesis* 2017,
1514   **55**.
1515   23.   Panfilio KA: **Late extraembryonic development and its *zen-RNAi*-induced failure in
1516   the milkweed bug *Oncopeltus fasciatus*.** *Dev Biol* 2009, **333:**297-311.
1517   24.   Panfilio KA, Roth S: **Epithelial reorganization events during late extraembryonic
1518   development in a hemimetabolous insect.** *Dev Biol* 2010, **340:**100-115.
1519   25.   Sharma AI, Yanes KO, Jin L, Garvey SL, Taha SM, Suzuki Y: **The phenotypic
1520   plasticity of developmental modules.** *Evodevo* 2016, **7:**15.
1521   26.   Hughes CL, Kaufman TC: **RNAi analysis of *Deformed, proboscipedia* and *Sex combs
1522   reduced* in the milkweed bug *Oncopeltus fasciatus*: novel roles for Hox genes in the
1523   hemipteran head.** *Development* 2000, **127:**3683-3694.
1524   27.   Wolfe SL, John B: **The organization and ultrastructure of male meiotic chromosomes
1525   in *Oncopeltus fasciatus*.** *Chromosoma* 1965, **17:**85-103.
1526   28.   Messthaler H, Traut W: **Phases of Sex Chromosome Inactivation in *Oncopeltus
1527   fasciatus* and *Pyrrhocoris apterus* (Insecta, Heteroptera).** *Caryologia* 1975, **28:**501-
1528   510.
1529   29.   McKenna DD, Scully ED, Pauchet Y, Hoover K, Kirsch R, Geib SM, Mitchell RF,
1530   Waterhouse RM, Ahn SJ, Arsala D, et al: **Genome of the Asian longhorned beetle
1531   (*Anoplophora glabripennis*), a globally significant invasive species, reveals key
1532   functional and evolutionary innovations at the beetle-plant interface.** *Genome Biol*
1533   2016, **17:**227.
1534   30.   Simpson JT: **Exploring genome characteristics and sequence quality without a
1535   reference.** *Bioinformatics* 2014, **30:**1228-1235.
1536   31.   Hughes CL, Kaufman TC: **RNAi analysis of Deformed, proboscipedia and Sex combs
1537   reduced in the milkweed bug Oncopeltus fasciatus: novel roles for Hox genes in the
1538   Hemipteran head.** *Development* 2000, **127:**3683-3694.
1539   32.   Panfilio KA, Liu PZ, Akam M, Kaufman TC: ***Oncopeltus fasciatus zen* is essential for
1540   serosal tissue function in katatrepsis.** *Dev Biol* 2006, **292:**226-243.
1541   33.   Tian X, Xie Q, Li M, Gao C, Cui Y, Xi L, Bu W: **Phylogeny of pentatomomorphan
1542   bugs (Hemiptera-Heteroptera:Pentatomomorpha) based on six Hox gene fragments.**
1543   *Zootaxa* 2011, **2888:**57-68.
1544   34.   Ewen-Campen B, Shaner N, Panfilio KA, Suzuki Y, Roth S, Extavour CG: **The
1545   maternal and early embryonic transcriptome of the milkweed bug *Oncopeltus
1546   fasciatus*.** *BMC Genomics* 2011, **12:**61.
1547   35.   Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P: **Parallel molecular
1548   evolution in an herbivore community.** *Science* 2012, **337:**1634-1637.
1549   36.   Robertson HM: **The insect chemoreceptor superfamily in Drosophila pseudoobscura:
1550   Molecular evolution of ecologically-relevant genes over 25 million years** *J Insect Sci*
1551   2009, **9:**18.
1552   37.   Robertson HM: **Taste: Independent origins of chemoreception coding systems?** *Curr
1553   Biol* 2001, **11:**R560-R562.
1554   38.   Jazwinska A, Rushlow C, Roth S: **The role of *brinker* in mediating the graded
1555   response to Dpp in early *Drosophila* embryos.** *Development* 1999, **126:**3323-3334.
1556   39.   Ewer J, Truman JW: **Increases in cyclic 3',5'-guanosine monophosphate (cGMP)
1557   occur at ecdysis in an evolutionarily conserved crustacean cardioactive peptide-
1558   immunoreactive insect neuronal network.** *Journal of Comparative Neurology* 1996,
1559   **370:**330-341.

40. Togawa T, Dunn WA, Emmons AC, Nagao J, Willis JH: **Developmental expression patterns of cuticular protein genes with the R&R Consensus from *Anopheles gambiae*.** *Insect Biochem Mol Biol* 2008, **38:**508-519.

41. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.** *Bioinformatics* 2015, **31:**3210-3212.

42. Kriventseva EV, Tegenfeldt F, Petty TJ, Waterhouse RM, Simão FA, Pozdnyakov IA, Ioannidis P, Zdobnov EM: **OrthoDB v8: update of the hierarchical catalog of orthologs and the underlying free software.** *Nucl Acids Res* 2015, **43:**D250-D256.

43. Shigenobu S, Bickel RD, Brisson JA, Butts T, Chang CC, Christiaens O, Davis GK, Duncan EJ, Ferrier DE, Iga M, et al: **Comprehensive survey of developmental genes in the pea aphid, *Acyrthosiphon pisum*: frequent lineage-specific duplications and losses of developmental genes.** *Insect Mol Biol* 2010, **19 Suppl 2:**47-62.

44. Bansal R, Michel AP: **Core RNAi Machinery and *Sid1*, a component for systemic RNAi, in the hemipteran insect, *Aphis glycines*.** *Int J Mol Sci* 2013, **14:**3786-3801.

45. Bao R, Fischer T, Bolognesi R, Brown SJ, Friedrich M: **Parallel duplication and partial subfunctionalization of beta-catenin/armadillo during insect evolution.** *Mol Biol Evol* 2012, **29:**647-662.

46. Sachs L, Chen YT, Drechsler A, Lynch JA, Panfilio KA, Lassig M, Berg J, Roth S: **Dynamic BMP signaling polarized by Toll patterns the dorsoventral axis in a hemimetabolous insect.** *eLife* 2015, **4:**e05502.

47. Konopova B, Smykal V, Jindra M: **Common and distinct roles of juvenile hormone signaling genes in metamorphosis of holometabolous and hemimetabolous insects.** *PLoS One* 2011, **6:**e28728.

48. Armisen D, Refki PN, Crumiere AJ, Viala S, Toubiana W, Khila A: **Predator strike shapes antipredator phenotype through new genetic interactions in water striders.** *Nat Commun* 2015, **6:**8153.

49. Wulff JP, Sierra I, Sterkel M, Holtof M, Van Wielendaele P, Francini F, Broeck JV, Ons S: **Orcokinin neuropeptides regulate ecdysis in the hemimetabolous insect Rhodnius prolixus.** *Insect Biochem Mol Biol* 2017, **81:**91-102.

50. Vellichirammal NN, Gupta P, Hall TA, Brisson JA: **Ecdysone signaling underlies the pea aphid transgenerational wing polyphenism.** *Proc Natl Acad Sci U S A* 2017, **114:**1419-1423.

51. Chiu TL, Wen Z, Rupasinghe SG, Schuler MA: **Comparative molecular modeling of Anopheles gambiae CYP6Z1, a mosquito P450 capable of metabolizing DDT.** *Proc Natl Acad Sci U S A* 2008, **105:**8855-8860.

52. Gong Y, Li T, Feng Y, Liu N: **The function of two P450s, CYP9M10 and CYP6AA7, in the permethrin resistance of Culex quinquefasciatus.** *Sci Rep* 2017, **7:**587.

53. Liu PZ, Kaufman TC: ***hunchback* is required for suppression of abdominal identity, and for proper germband growth and segmentation in the intermediate germband insect *Oncopeltus fasciatus*.** *Development* 2004, **131:**1515-1527.

54. Schaeper ND, Pechmann M, Damen WGM, Prpic N-M, Wimmer EA: **Evolutionary plasticity of *collier* function in head development of diverse arthropods.** *Dev Biol* 2010, **344:**363-376.

55. Aspiras AC, Smith FW, Angelini DR: **Sex-specific gene interactions in the patterning of insect genitalia.** *Dev Biol* 2011, **360:**369-380.

56. Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi HS, Lambert SA, Mann I, Cook K, et al: **Determination and inference of eukaryotic transcription factor sequence specificity.** *Cell* 2014, **158:**1431-1443.

57. Peel AD, Telford MJ, Akam M: **The evolution of hexapod engrailed-family genes: evidence for conservation and concerted evolution.** *Proc Biol Sci* 2006, **273:**1733-1742.

58. Ben-David J, Chipman AD: **Mutual regulatory interactions of the trunk gap genes during blastoderm patterning in the hemipteran *Oncopeltus fasciatus*.** *Dev Biol* 2010, **346:**140-149.

59. Erezyilmaz DF, Kelstrup HC, Riddiford LM: **The nuclear receptor E75A has a novel pair-rule-like function in patterning the milkweed bug, *Oncopeltus fasciatus.*** *Dev Biol* 2009, **334:**300-310.

60. Liu PZ, Kaufman TC: ***even-skipped* is not a pair-rule gene but has segmental and gap-like functions in *Oncopeltus fasciatus*, an intermediate germband insect.** *Development* 2005, **132:**2081-2092.

61. Weisbrod A, Cohen M, Chipman AD: **Evolution of the insect terminal patterning system--insights from the milkweed bug, *Oncopeltus fasciatus.*** *Dev Biol* 2013, **380:**125-131.

62. Albertin CB, Simakov O, Mitros T, Wang ZY, Pungor JR, Edsinger-Gonzales E, Brenner S, Ragsdale CW, Rokhsar DS: **The octopus genome and the evolution of cephalopod neural and morphological novelties.** *Nature* 2015, **524:**220-224.

63. Crooks GE, Hon G, Chandonia J-M, Brenner SE: **WebLogo: A sequence logo generator.** *Genome Res* 2004, **14:**1188-1190.

64. Najafabadi HS, Mnaimneh S, Schmitges FW, Garton M, Lam KN, Yang A, Albu M, Weirauch MT, Radovani E, Kim PM, et al: **C2H2 zinc finger proteins greatly expand the human regulatory lexicon.** *Nat Biotechnol* 2015, **33:**555-562.

65. Emerson RO, Thomas JH: **Adaptive evolution in zinc finger transcription factors.** *PLoS Genet* 2009, **5:**e1000325.

66. Thomas JH, Schneider S: **Coevolution of retroelements and tandem zinc finger genes.** *Genome Res* 2011, **21:**1800-1812.

67. Garcia-Perez JL, Widmann TJ, Adams IR: **The impact of transposable elements on mammalian development.** *Development* 2016, **143:**4101-4114.

68. Liu PZ, Kaufman TC: ***Krüppel* is a gap gene in the intermediate insect *Oncopeltus fasciatus* and is required for development of both blastoderm and germband-derived segments.** *Development* 2004, **131:**4567-4579.

69. Heger P, Marin B, Bartkuhn M, Schierenberg E, Wiehe T: **The chromatin insulator CTCF and the emergence of metazoan diversity.** *Proc Natl Acad Sci USA* 2012, **109:**17507-17512.

70. Liu H, Chang L-H, Sun Y, Lu X, Stubbs L: **Deep vertebrate roots for mammalian zinc finger transcription factor subfamilies.** *Genome Biol Evol* 2014, **6:**510-525.

71. Imbeault M, Helleboid P-Y, Trono D: **KRAB zinc-finger proteins contribute to the evolution of gene regulatory networks.** *Nature* 2017, **543:**550-554.

72. Csuros M, Rogozin IB, Koonin EV: **A detailed history of intron-rich eukaryotic ancestors inferred from a global survey of 100 complete genomes.** *PLoS Comput Biol* 2011, **7:**e1002150.

73. Papanicolaou A, Schetelig MF, Arensburger P, Atkinson PW, Benoit JB, Bourtzis K, Castañera P, Cavanaugh JP, Chao H, Childers C, et al: **The whole genome sequence of the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann), reveals insights into the biology and adaptive evolution of a highly invasive pest species.** *Genome Biol* 2016, **17:**192.

74. Hoy MA, Waterhouse RM, Wu K, Estep AS, Ioannidis P, Palmer WJ, Pomerantz AF, Simao FA, Thomas J, Jiggins FM, et al: **Genome sequencing of the phytoseiid predatory mite *Metaseiulus occidentalis* reveals completely atomized Hox genes and superdynamic intron evolution.** *Genome Biol Evol* 2016, **8:**1762-1775.

75. Seibt KM, Wenke T, Muders K, Truberg B, Schmidt T: **Short interspersed nuclear elements (SINEs) are abundant in Solanaceae and have a family-specific impact on gene structure and genome organization.** *Plant J* 2016, **86:**268-285.

76. Huff JT, Zilberman D, Roy SW: **Mechanism for DNA transposons to generate introns on genomic scales.** *Nature* 2016, **538:**533-536.

77. Wheeler D, Redding AJ, Werren JH: **Characterization of an ancient lepidopteran lateral gene transfer.** *PLoS One* 2012, **8:**e59262.

78. Da Lage JL, Binder M, Hua-Van A, Janecek S, Casane D: **Gene make-up: rapid and massive intron gains after horizontal transfer of a bacterial alpha-amylase gene to Basidiomycetes.** *BMC Evol Biol* 2013, **13:**40.

79.   Lee DH, Short BD, Joseph SV, Bergh JC, Leskey TC: **Review of the biology, ecology,**
and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) in China,
Japan, and the Republic of Korea. *Environ Entomol* 2013, **42:**627-641.
80.   Lawrence PA: **Cellular differentiation and pattern formation during metamorphosis**
of the milkweed bug Oncopeltus. *Dev Biol* 1969, **19:**12-40.
81.   Riddiford LM: **Prevention of Metamorphosis by Exposure of Insect Eggs to Juvenile**
Hormone Analogs. *Science* 1970, **167:**287-&.
82.   Willis JH, Lawrence PA: **Deferred Action of Juvenile Hormone.** *Nature* 1970, **225:**81-
83.
83.   Masner P, Bowers WS, Kalin M, Muhle T: **Effect of precocene II on the endocrine**
regulation of development and reproduction in the bug, *Oncopeltus fasciatus.* *Gen*
Comp Endocrinol 1979, **37:**156-166.
84.   Rewitz K, O'Connor M, Gilbert L: **Molecular evolution of the insect Halloween family**
of cytochrome P450s: phylogeny, gene organization and functional conservation.
Insect Biochem Mol Biol 2007, **37:**741-753.
85.   Huet F, Ruiz C, Richards G: **Sequential gene activation by ecdysone in Drosophila**
melanogaster: the hierarchical equivalence of early and early late genes.
Development 1995, **121:**1195-1204.
86.   Bialecki M, Shilton A, Fichtenberg C, Segraves WA, Thummel CS: **Loss of the**
ecdysteroid-inducible E75A orphan nuclear receptor uncouples molting from
metamorphosis in Drosophila. *Dev Cell* 2002, **3:**209-220.
87.   Truman J, Rountree D, Reiss S, Schwartz L: **Ecdysteroids regulate the release and**
action of eclosion hormone in the tobacco hornworm, Manduca sexta (L.) *J Insect*
Physiol 1983, **29:**895–900.
88.   Zitnan D, Kingan TG, Hermesman JL, Adams ME: **Identification of ecdysis-triggering**
hormone from an epitracheal endocrine system. *Science* 1996, **271:**88-91.
89.   Charles JP, Iwema T, Epa VC, Takaki K, Rynes J, Jindra M: **Ligand-binding properties**
of a juvenile hormone receptor, Methoprene-tolerant. *Proceedings of the National*
Academy of Sciences of the United States of America 2011, **108:**21128-21133.
90.   Minakuchi C, Zhou X, Riddiford L: **Kruppel homolog 1 (Kr-h1) mediates juvenile**
hormone action during metamorphosis of Drosophila melanogaster. *Mech Dev* 2008,
125:91-105.
91.   Minakuchi C, Namiki T, Shinoda T: **Kruppel homolog 1, an early juvenile hormone-**
response gene downstream of Methoprene-tolerant, mediates its anti-metamorphic
action in the red flour beetle Tribolium castaneum. *Dev Biol* 2009, **352:**341-350.
92.   DiBello PR, Withers DA, Bayer CA, Fristrom JW, Guild GM: **The Drosophila Broad-**
Complex encodes a family of related proteins containing zinc fingers. *Genetics* 1991,
129:385-397.
93.   Karim F, Guild G, Thummel C: **The Drosophila Broad-Complex plays a key role in**
controlling ecdysone-regulated gene expression at the onset of metamorphosis.
Development 1993, **118:**977-988.
94.   Erezyilmaz DF, Riddiford LM, Truman JW: **The pupal specifier broad directs**
progressive morphogenesis in a direct-developing insect. *Proceedings of the National*
Academy of Sciences of the United States of America 2006, **103:**6925-6930.
95.   Arakane Y, Hogenkamp DG, Zhu YC, Kramer KJ, Specht CA, Beeman RW, Kanost MR,
Muthukrishnan S: **Characterization of two chitin synthase genes of the red flour**
beetle, *Tribolium castaneum*, and alternate exon usage in one of the genes during
development. *Insect Biochem Mol Biol* 2004, **34:**291-304.
96.   True JR: **Insect melanism: the molecules matter.** *Trends in Ecology & Evolution* 2003,
18:640-647.
97.   Zhan SA, Guo QH, Li MH, Li MW, Li JY, Miao XX, Huang YP: **Disruption of an N-**
acetyltransferase gene in the silkworm reveals a novel role in pigmentation.
Development 2010, **137:**4083-4090.
98.   Liu J, Lemonds TR, Popadic A: **The genetic control of aposematic black pigmentation**
in hemimetabolous insects: insights from Oncopeltus fasciatus. *Evolution &*
Development 2014, **16:**270-277.

99.   Liu J, Lemonds TR, Marden JH, Popadic A: **A Pathway Analysis of Melanin**
Patterning in a Hemimetabolous Insect. *Genetics* 2016, **203:**403-413.

100.   Lawrence PA: **Some new mutants of large milkweed bug** *Oncopeltus fasciatus* **Dall.**
Genetical Research 1970, **15:**347-350.

101.   Morgan ED: *Biosynthesis in Insects: Advanced Edition.* London: Royal Society of
Chemistry,; 2010.

102.   McLean JR, Krishnakumar S, O'Donnell JM: **Multiple mRNAs from the Punch locus of**
Drosophila melanogaster encode isoforms of GTP cyclohydrolase I with distinct N-
terminal domains. *J Biol Chem* 1993, **268:**27191-27197.

103.   Wiederrecht GJ, Paton DR, Brown GM: **Enzymatic Conversion of Dihydroneopterin**
Triphosphate to the Pyrimidodiazepine Intermediate Involved in the Biosynthesis of
the Drosopterins in Drosophila-Melanogaster. *Journal of Biological Chemistry* 1984,
259:2195-2200.

104.   Newcombe D, Blount JD, Mitchell C, Moore AJ: **Chemical egg defence in the large**
milkweed bug, Oncopeltus fasciatus, derives from maternal but not paternal diet.
Entomologia Experimentalis et Applicata 2013, **149:**197-205.

105.   Zhan S, Merlin C, Boore JL, Reppert SM: **The monarch butterfly genome yields**
insights into long-distance migration. *Cell* 2011, **147:**1171-1185.

106.   Joseph RM, Carlson JR: **Drosophila Chemoreceptors: A Molecular Interface Between**
the Chemical World and the Brain. *Trends Genet* 2015, **31:**683-695.

107.   Benton R: **Multigene Family Evolution: Perspectives from Insect Chemoreceptors.**
Trends Ecol Evol 2015, **30:**590-600.

108.   Robertson HM, Warr CG, Carlson JR: **Molecular evolution of the insect**
chemoreceptor gene superfamily in Drosophila melanogaster. *Proc Natl Acad Sci U S*
A 2003, **100 Suppl 2:**14537-14542.

109.   Rytz R, Croset V, Benton R: **Ionotropic receptors (IRs): chemosensory ionotropic**
glutamate receptors in Drosophila and beyond. *Insect Biochem Mol Biol* 2013,
43:888-897.

110.   Kirkness EF, Haas BJ, Sun W, Braig HR, Perotti MA, Clark JM, Lee SH, Robertson HM,
Kennedy RC, Elhaik E, et al: **Genome sequences of the human body louse and its**
primary endosymbiont provide insights into the permanent parasitic lifestyle. *Proc*
Natl Acad Sci U S A 2010, **107:**12168-12173.

111.   Smadja C, Shi P, Butlin RK, Robertson HM: **Large gene family expansions and**
adaptive evolution for odorant and gustatory receptors in the pea aphid,
Acyrthosiphon pisum. *Mol Biol Evol* 2009, **26:**2073-2086.

112.   Terrapon N, Li C, Robertson HM, Ji L, Meng X, Booth W, Chen Z, Childers CP, Glastad
KM, Gokhale K, et al: **Molecular traces of alternative social organization in a termite**
genome. *Nat Commun* 2014, **5:**3636.

113.   Xu W, Papanicolaou A, Zhang HJ, Anderson A: **Expansion of a bitter taste receptor**
family in a polyphagous insect herbivore. *Sci Rep* 2016, **6:**23666.

114.   Feir D: *Oncopeltus fasciatus*: **A research animal.** *Annu Rev Entomol* 1974, **19:**81-96.

115.   Croset V, Rytz R, Cummins SF, Budd A, Brawand D, Kaessmann H, Gibson TJ, Benton
R: **Ancient protostome origin of chemosensory ionotropic glutamate receptors and**
the evolution of insect taste and olfaction. *PLoS Genet* 2010, **6:**e1001064.

116.   Vellozo AF, Véron AS, Baa-Puyoulet P, Huerta-Cepas J, Cottret L, Febvay G, Calevro F,
Rahbe Y, Douglas AE, Gabaldón T, et al: **CycADS: an annotation database system to**
ease the development and update of BioCyc databases. *Database* 2011, **2011:**bar008-
bar008.

117.   Baa-Puyoulet P, Parisot N, Febvay G, Huerta-Cepas J, Vellozo AF, Gabaldón T, Calevro
F, Charles H, Colella S: **ArthropodaCyc: a CycADS powered collection of BioCyc**
databases to analyse and compare metabolism of arthropods. *Database (Oxford)*
2016, **pii:**baw081.

118.   Hojilla-Evangelista MP, Evangelista RL: **Characterization of milkweed (Asclepias**
spp.) seed proteins. *Industrial crops and ...* 2009.

119.   Dean CAE, Teets NM, Koštál V, Šimek P, Denlinger DL: **Enhanced stress responses**
and metabolic adjustments linked to diapause and onset of migration in the large
milkweed bug *Oncopeltus fasciatus*. *Physiol Entomol* 2016, **DOI: 10.1111/phen.12140.**

120. Sandström J, Moran N: **How nutritionally imbalanced is phloem sap for aphids?**
Entomologia Experimentalis et Applicata 1999, **91:**203-210.

121. Rabatel A, Febvay G, Gaget K, Duport G, Baa-Puyoulet P, Sapountzis P, Bendridi N,
Rey M, Rahbé Y, Charles H, et al: **Tyrosine pathway regulation is host-mediated in**
the pea aphid symbiosis during late embryonic and early larval development. *BMC*
Genomics 2013, **14:**235.

122. Dobler S, Petschenka G, Wagschal V, Flacht L: **Convergent adaptive evolution – how**
insects master the challenge of cardiac glycoside-containing host plants. *Entomologia*
Experimentalis et Applicata 2015, **157:**30-39.

123. Niehuis O, Gibson JD, Rosenberg MS, Pannebakker BA, Koevoets T, Judson AK,
Desjardins CA, Kennedy K, Duggan D, Beukeboom LW, et al: **Recombination and its**
impact on the genome of the haplodiploid parasitoid wasp Nasonia. *PLoS One* 2010,
5:e8597.

124. Ferrero A, Torreblanca A, Garcera MD: **Assessment of the effects of orally**
administered ferrous sulfate on Oncopeltus fasciatus (Heteroptera: Lygaeidae).
Environ Sci Pollut Res Int 2017, **24:**8551-8561.

125. Hare EE, Johnston JS: **Genome size determination using flow cytometry of propidium**
iodide-stained nuclei. *Methods Mol Biol* 2011, **772:**3-12.

126. Marcais G, Kingsford C: **A fast, lock-free approach for efficient parallel counting of**
occurrences of k-mers. *Bioinformatics* 2011, **27:**764-770.

127. Bushnell B: **BBMap short read aligner.** 2016. http://sourceforge.net/projects/bbmap

128. Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall
G, Shea TP, Sykes S, et al: **High-quality draft assemblies of mammalian genomes**
from massively parallel sequence data. *Proc Natl Acad Sci U S A* 2011, **108:**1513-
1518.

129. Holt C, Yandell M: **MAKER2: an annotation pipeline and genome-database**
management tool for second-generation genome projects. *BMC Bioinformatics* 2011,
12:491.

130. Lee E, Helt G, Reese J, Munoz-Torres M, Childers C, Buels R, Stein L, Holmes I, Elsik
C, Lewis S: **Web Apollo: a web-based genomic annotation editing platform.** *Genome*
Biology 2013, **14**.

131. Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee CY, Lin H, Lin JW,
Hackett K: **The i5k Workspace@NAL--enabling genomic data access, visualization**
and curation of arthropod genomes. *Nucleic Acids Res* 2015, **43:**D714-719.

132. Hughes DST, Koelzer S, Panfilio KA, Richards S: *Oncopeltus fasciatus* **genome**
annotations v0.5.3. *Ag Data Commons (Database)*
2015**:**http://dx.doi.org/10.15482/USDA.ADC/1173237.

133. Murali SC, The i5k genome assembly team (29 additional authors), Han Y, Richards S,
Worley K, Muzny D, Gibbs R, Koelzer S, Panfilio KA: *Oncopeltus fasciatus* **genome**
assembly 1.0. *Ag Data Commons (Database)*
2015**:**http://dx.doi.org/10.15482/USDA.ADC/1173238.

134. Vargas Jentzsch IM, Hughes DST, Poelchau M, Robertson HM, Benoit JB, Rosendale
AJ, Armisén D, Duncan EJ, Vreede BMI, Jacobs CGC, et al: *Oncopeltus fasciatus*
Official Gene Set OGS_v1.1 for genome assembly Oncopeltus fasciatus v1.0. *Ag Data*
Commons (Database) 2015**:**http://dx.doi.org/10.15482/USDA.ADC/1173142.

135. **RepeatModeler Open-1.0.8** [http://www.repeatmasker.org]

136. Bao Z, Eddy SR: **Automated de novo identification of repeat sequence families in**
sequenced genomes. *Genome Res* 2002, **12:**1269-1276.

137. Price AL, Jones NC, Pevzner PA: **De novo identification of repeat families in large**
genomes. *Bioinformatics* 2005, **21 Suppl 1:**i351-358.

138. Benson G: **Tandem repeats finder: a program to analyze DNA sequences.** *Nucleic*
Acids Res 1999, **27:**573-580.

139. **RepeatMasker Open-4.0.** [http://www.repeatmasker.org]

140. Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods*
2012, **9:**357-359.

141.   Goff S, Vaughn M, McKay S, Lyons E, Stapleton A, Gessler D, Matasci N, Wang L,
1839   Hanlon M, Lenards A, et al: **The iPlant Collaborative: Cyberinfrastructure for Plant**
1840   **Biology.** *Frontiers in plant science* 2011, **2**.
1841   142.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
1842   Raychowdhury R, Zeng Q, et al: **Full-length transcriptome assembly from RNA-Seq**
1843   **data without a reference genome.** *Nat Biotechnol* 2011, **29**:644-652.
1844   143.   Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB,
1845   Eccles D, Li B, Lieber M, et al: **De novo transcript sequence reconstruction from**
1846   **RNA-seq using the Trinity platform for reference generation and analysis.** *Nat*
1847   *Protoc* 2013, **8**:1494-1512.
1848   144.   Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data**
1849   **with or without a reference genome.** *BMC Bioinformatics* 2011, **12**:323.
1850   145.   Finn RD, Mistry J, Tate J, Coggill P, Heger A, Pollington JE, Gavin OL, Gunasekaran P,
1851   Ceric G, Forslund K, et al: **The Pfam protein families database.** *Nucleic Acids Res*
1852   2010, **38**:D211-222.
1853   146.   Weirauch MT, Hughes TR: **A catalogue of eukaryotic transcription factor types, their**
1854   **evolutionary origin, and species distribution.** *Subcell Biochem* 2011, **52**:25-73.
1855   147.   Eddy SR: **A new generation of homology search tools based on probabilistic**
1856   **inference.** *Genome Inform* 2009, **23**:205-211.
1857   148.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
1858   Remmert M, Soding J, et al: **Fast, scalable generation of high-quality protein multiple**
1859   **sequence alignments using Clustal Omega.** *Mol Syst Biol* 2011, **7**:539.
1860   149.   Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M: **KAAS: an automatic genome**
1861   **annotation and pathway reconstruction server.** *Nucleic Acids Research* 2007,
1862   **35**:W182-185.
1863   150.   Claudel-Renard C, Chevalet C, Faraut T, Kahn D: **Enzyme-specific profiles for genome**
1864   **annotation: PRIAM.** *Nucleic Acids Research* 2003, **31**:6633-6639.
1865   151.   Conesa A, Götz S: **Blast2GO: A Comprehensive Suite for Functional Analysis in**
1866   **Plant Genomics.** *International journal of plant genomics* 2008, **2008**:619832.
1867   152.   Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M: **Blast2GO: a**
1868   **universal tool for annotation, visualization and analysis in functional genomics**
1869   **research.** *Bioinformatics (Oxford, England)* 2005, **21**:3674-3676.
1870   153.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
1871   Mitchell A, Nuka G, et al: **InterProScan 5: genome-scale protein function**
1872   **classification.** *Bioinformatics* 2014, **30**:1236-1240.
1873   154.   Karp PD, Ouzounis CA, Moore-Kochlacs C, Goldovsky L, Kaipa P, Ahrén D, Tsoka S,
1874   Darzentas N, Kunin V, López-Bigas N: **Expansion of the BioCyc collection of**
1875   **pathway/genome databases to 160 genomes.** *Nucleic Acids Research* 2005, **33**:6083-
1876   6089.
1877   155.   Karp PD, Paley SM, Krummenacker M, Latendresse M, Dale JM, Lee TJ, Kaipa P,
1878   Gilham F, Spaulding A, Popescu L, et al: **Pathway Tools version 13.0: integrated**
1879   **software for pathway/genome informatics and systems biology.** *Briefings in*
1880   *Bioinformatics* 2010, **11**:40-79.
1881   156.   Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon
1882   S, Lefort V, Lescot M, et al: **Phylogeny.fr: robust phylogenetic analysis for the non-**
1883   **specialist.** *Nucleic Acids Res* 2008, **36**:W465-469.
1884   157.   Wang X, Fang X, Yang P, Jiang X, Jiang F, Zhao D, Li B, Cui F, Wei J, Ma C, et al: **The**
1885   **locust genome provides insight into swarm formation and long-distance flight.** *Nat*
1886   *Commun* 2014, **5**:2957.
1887   158.   The International Silkworm Genome Consortium: **The genome of a lepidopteran model**
1888   **insect, the silkworm Bombyx mori.** *Insect Biochem Mol Biol* 2008, **38**:1036-1045.
1889   159.   Elsik CG, Worley KC, Bennett AK, Beye M, Camara F, Childers CP, de Graaf DC,
1890   Debyser G, Deng J, Devreese B, et al: **Finding the missing honey bee genes: lessons**
1891   **learned from a genome upgrade.** *BMC Genomics* 2014, **15**:86.
1892   160.   Honeybee Genome Sequencing Consortium: **Insights into social insects from the**
1893   **genome of the honeybee Apis mellifera.** *Nature* 2006, **443**:931-949.

161. Richards S, Gibbs RA, Weinstock GM, Brown SJ, Denell R, Beeman RW, Gibbs R,
Bucher G, Friedrich M, Grimmelikhuijzen CJ, et al: **The genome of the model beetle**
and pest Tribolium castaneum. *Nature* 2008, **452:**949-955.
162. Chen XG, Jiang X, Gu J, Xu M, Wu Y, Deng Y, Zhang C, Bonizzoni M, Dermauw W,
Vontas J, et al: **Genome sequence of the Asian Tiger mosquito, Aedes albopictus,**
reveals insights into its biology, genetics, and evolution. *Proc Natl Acad Sci U S A*
2015, **112:**E5907-5915.
163. Ellis LL, Huang W, Quinn AM, Ahuja A, Alfrejd B, Gomez FE, Hjelmen CE, Moore KL,
Mackay TF, Johnston JS, Tarone AM: **Intrapopulation genome size variation in D.**
melanogaster reflects life history variation and plasticity. *PLoS Genet* 2014,
10:e1004522.
164. Kumar S, Stecher G, Tamura K: **MEGA7: Molecular Evolutionary Genetics Analysis**
Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016, **33:**1870-1874.
1907

Fig 1. The large milkweed bug, *Oncopeltus fasciatus*, shown in its phylogenetic and environmental context.

Fig 2. Comparisons of the official gene set and transcriptomic resources.

Fig 3. Orthology comparisons and phylogenetic placement of *Oncopeltus fasciatus* **among other Arthropoda.**

Fig 4. Distribution of transcription factor families across insect genomes.

Fig 5. Comparison of repeat content estimations.

Fig 6. Trends in gene structure show hemipteroid-specific tendencies.

Fig 7. Splice site evolution correlates with both lineage and, independently, genome size.

Fig 8. Lateral gene transfer introduction and subsequent evolution within the Hemiptera for mannosidase-encoding genes.

Treatment	Injected Nymphs	Surviving Adults	RNAi phenotype
Z2	33	12 (36%)	8 (67%)
Z3	29	16 (55%)	9 (56%)
Z4	28	22 (79%)	0 (0%)
Control	12	11 (92%)	0 (0%)

Fig 9. Isoform-specific RNAi based on new genome annotations affects the molting and cuticle identity gene _broad_.

Fig 10. Comparison of the urea cycle of *Oncopeltus* with 26 other insect species.