Combining genotype, phenotype, and environmental data to delineate site-adjusted provenance strategies for ecological restoration

Carolina S. Carvalho[1], Brenna R. Forester[2], Simone K. Mitre[1], Ronnie Alves[1], Vera L. Imperatriz-Fonseca[1], Silvio J. Ramos[1], Luciana C. Resende-Moreira[1], José O. Siqueira[1], Leonardo C. Trevelin[1], Cecilio F. Caldeira[1], Markus Gastauer[1], Rodolfo Jaffé[1,3*]

[1] Instituto Tecnológico Vale, Belém-PA, Brazil.

[2] Colorado State University, Fort Collins-CO, USA

[3] Departamento de Ecologia, Universidade de São Paulo, São Paulo-SP, Brazil.

[*] **Correspondence:**

Rodolfo Jaffé

email: r.jaffe@ib.usp.br

phone: +55 (91) 3213 5523

Abstract

Despite the importance of climate-adjusted provenancing to mitigate the effects of

environmental change, climatic considerations alone are insufficient when restoring highly

degraded sites. Here we propose a comprehensive landscape genomic approach to assist the

restoration of moderately disturbed and highly degraded sites. To illustrate it we employ

genomic datasets comprising thousands of single nucleotide polymorphisms from two plant

species suitable for the restoration of iron-rich Amazonian Savannas. We first use a subset of

neutral loci to assess genetic structure and determine the genetic neighborhood size. We then

identify genotype-phenotype-environment associations, map adaptive genetic variation, and

predict adaptive genotypes for restoration sites. Whereas local provenances were found

optimal to restore a moderately disturbed site, a mixture of genotypes seemed the most

promising strategy to recover a highly degraded mining site. We discuss how our results can

help define site-adjusted provenancing strategies, and argue that our methods can be more

broadly applied to assist other restoration initiatives.

Keywords: Genotype-environment associations (GEA), genotype-phenotype associations

(GPA), landscape genomics, local adaptation, RAD sequencing, restoration genomics, single

nucleotide polymorphisms (SNP).

Introduction

In spite of the broadly recognized importance of genetic provenance for restoration initiatives, the use of genomic tools to define provenance strategies is still uncommon. Choosing provenances based on genetic knowledge can help increase genetic diversity and adaptability, thereby contributing to the success of restoration initiatives (Broadhurst et al., 2008; Mijangos et al., 2015; Weeks et al., 2011). Fortunately, the use of genetic provenancing is increasing, as advances in Next-Generation-Sequencing technologies have made possible large-scale assessments of neutral and adaptive genetic variation (Breed et al., 2019; Mijangos et al., 2015; Williams et al., 2014). For instance, neutral loci (i.e., those which are not subject to natural selection) can be used to identify independent demographic units, assess fine-scale spatial genetic structure, and quantify genetic diversity (Allendorf et al., 2013; Balkenhol et al., 2017); whereas adaptive loci (under natural selection) are relevant to detect adaptations to local environmental conditions and delineate adaptive units (Funk et al., 2012; Rellstab et al., 2015).

Restoration genomic studies published so far have assessed the effect of multiple environmental variables on genetic composition in order to identify which individuals or populations are "pre-adapted" to future climates (Gugger et al., 2018; Lu et al., 2019; Martins et al., 2018; Rossetto et al., 2019; Shryock et al., 2017, 2015; Steane et al., 2014; Supple et al., 2018). Although this information is essential to inform predictive and climate-adjusted provenancing schemes (Prober et al., 2015), the emphasis on climate has overshadowed the application of genomic methods to restore extremely degraded sites (Bucharova et al., 2019; Lesica and Allendorf, 1999). Such site-adjusted provenancing (targeting the restoration of

specific sites considering their current environmental conditions) may not even incorporate

climate change considerations, as highly degraded sites have unique characteristics that make

them extremely challenging to restore. First, highly degraded sites usually require immediate

restoration or rehabilitation, so adaptations to current environmental conditions are more

suitable to guide provenance strategies than those based on future climate (Gastauer et al.,

2019). Second, environmental protection agencies usually require the restitution of

ecosystems to conditions as close to a pre-disturbance baseline as possible, as well as regular

(and costly) monitoring until rehabilitation goals have been achieved (Gastauer et al., 2019).

The main efforts thus lay in the quick establishment of viable populations that will restore

ecosystem functions and processes, prevent soil erosion, and protect biological diversity.

Third, highly degraded sites such as exhausted open-pit mines have radically different

environmental characteristics than natural habitats (Gastauer et al., 2019), so site-specific

characteristics are of primary importance to define provenance strategies. Such site-specific

variables generally need to be measured *in situ* and at fine spatial resolution, since they may

not be available as spatial layers in open-access repositories (and if they are, their spatial

resolution may be too coarse to reflect the reality of environmental conditions on the ground).

Finally, site-adjusted provenancing strategies need to consider local adaptations to climate,

soil, terrain, and even biological interactions, whereas climate-centered provenancing

strategies focus exclusively on climate.

The degree of disturbance can play an important role in determining site-adjusted

provenancing strategies (Breed et al., 2013; Lesica and Allendorf, 1999). Whereas local

genotypes are generally the best suited to restore sites where the degree of disturbance is low,

adaptations found in distant populations may facilitate establishment in highly degraded sites

to which local genotypes may not be adapted (Breed et al., 2013; Broadhurst et al., 2008;

Lesica and Allendorf, 1999)). Mixtures of genotypes from different populations have been

suggested as the best strategy to recover highly degraded sites, given that enhanced genetic

variation is more likely to rapidly generate local adaptations to novel ecological challenges

(Lesica and Allendorf, 1999). In any case, determining the most appropriate site-adjusted

provenancing strategy will require the delineation of local provenances and the spatial

distribution of local adaptations (Breed et al., 2019).

Although the use of neutral genetic markers to identify independent demographic units

is now common practice (Coates et al., 2018), few restoration studies have delineated seed

sourcing strategies based on population genetic structure (Durka et al., 2017) or the genetic

neighborhood size (the distance at which genetic composition stops being spatially

autocorrelated) (Krauss et al., 2013; Krauss and Koch, 2004; Rossetto et al., 2019). On the

other hand, only three restoration genomic studies so far have identified putative adaptive loci

and then mapped adaptive genetic variation (Martins et al., 2018; Shryock et al., 2015; Steane

et al., 2014). While the assessment of phenotype through common garden and reciprocal

transplant experiments to identify local adaptations has a long history (Aitken and Bemmels,

2016), no study has yet combined genotype-environment associations (GEA) with genotype-

phenotype associations (GPA) to delineate seed sourcing areas. This approach could improve

the inference of potential candidate genes and provide important insights into genes

underlying fitness-related traits (Mahony et al., 2019; Talbot et al., 2016; Vangestel et al.,

2018).

Here we propose a comprehensive landscape genomic approach to assist the

restoration of moderately disturbed and highly degraded sites. Relying on genotyping-by-

sequencing we identified thousands of single nucleotide polymorphisms in two plant species

of special interest for the restoration of exhausted mining sites from the Carajás Mineral

Province, located in the Eastern Amazon (Skirycz et al., 2014; Souza-Filho et al., 2019). We

first used a subset of neutral loci to assess broad and fine-scale genetic structure and

determine the genetic neighborhood size. Subsequently, we combined univariate and

multivariate methods to identify GEA and GPA and employed spatial principal components

analyses (sPCA) to map adaptive genetic variation while accounting for spatial

autocorrelation in genetic composition. Finally, we predicted the adaptive genotypes

associated with the environmental conditions of restoration sites (Fig. 1).

We focus on the Carajás Mineral Province, which harbors one of the world's largest

deposits of iron ore and huge iron ore mining projects, with operations dating back to the

1980s (Poveromo, 1999; Souza-Filho et al., 2019). The banded ironstone formations known

as *Cangas* (where iron-ore deposits are concentrated) are characterized by shallow, acidic,

nutrient-depleted and metal-rich soils and marked by high solar radiation, hot temperatures,

and a severe drought period (Skirycz et al., 2014), which pose severe challenges to plant

growth. Environmental legislation in Brazil requires the rehabilitation of disabled iron ore

mining sites (Gastauer et al., 2019), which constitute extremely degraded and difficult to

restore environments. These mining sites are characterized by extensive vegetation and soil

removal, compacted, nutrient-poor soils and steep slopes (Boyer and Wratten, 2010; Garris et

al., 2016; Whiting et al., 2004). The successful restoration and rehabilitation of these highly

degraded sites thus requires an appropriate selection of plant species (Giannini et al.,

2017) and seed sourcing zones to ensure that the introduced plants can effectively colonize

and establish viable populations. However, mine rehabilitation programs in the region employ

seed mixtures of exotic plants, due to the scarcity of native seeds and their lower germination

and growth rates (Silva et al., 2018). Additionally, some natural Canga environments from the

region have been repeatedly disturbed by fires, illegal cattle ranching and the introduction of

alien plant species, such as grasses and ferns. This is the case in Serra da Bocaina, which

became a National Park in 2017 and has since been protected (Mota et al., 2018). No

restoration initiatives have yet been implemented to recover the original ecosystems found in

Serra da Bocaina, but provenance strategies are expected to differ substantially from those

required to rehabilitate disabled mines.

Being native species, dominant in Canga environments, and once found at the sites

being restored, *Mimosa acutistipula var. ferrea* Barneby and *Dioclea apurensis* Kunth (both

legumes) are among the most promising plants for use in Canga restoration and mineland

rehabilitation programs (Giannini et al., 2017). Considered metallophyte species, both exhibit

biological mechanisms to tolerate and thrive in metalliferous soils (Preite et al., 2019; Whiting

et al., 2004). Moreover, they are extremely abundant in pristine Cangas ecosystems and

interact symbiotically with nitrogen-fixing bacteria, thus contributing to soil enrichment and

acting as pioneer species in restoration sites (Nunes et al., 2015; Ramos et al., 2019a; Silva et

al., 2018). *Mimosa acutistipula* is drought tolerant and well adapted to the low nutrient

content of Canga soils (Silva et al., 2018). On the other hand, *D. apurensis* requires low

nutrient inputs and shows high nutrient use efficiency (Ramos et al., 2019b). Moreover, this

species is a fast-growing liana with a ground-covering growth form, enabling the revegetation

and stabilization of mine pits and waste piles. Both species have high germination rates

(Ramos et al., 2019a), can be observed growing on minelands, and seem to be central in plant-

pollinator networks (unpublished data). Considering the heterogeneous and hostile

environment where both species occur (Mitre et al., 2018), and their life-history similarities,

we expected to find similar patterns of neutral and adaptive genetic structure in both species

across our study area. We also expected that local populations would not be adapted to the

environmental conditions found in exhausted mining sites given they drastically differ from

pre-mining conditions, whereas populations from Serra da Bocaina would show adaptations to

local environmental conditions. Based on our results, we propose two different site-adjusted

provenance strategies for the restoration of a degraded mine site and a disturbed but unmined

Canga environment. We discuss the merits of our approach and argue that it can be more

broadly applied to define site-adjusted provenancing strategies.

Material and Methods

Sampling

We followed a stratified sampling design, seeking to ensure high statistical power in GEA and

GPA analyses by maximizing environmental variability within different genetic clusters. We

collected samples of 180 individuals of *M. acutistipula* var. *ferrea* and 167 individuals of *D.*

apurensis between February and May of 2018 (SISBIO collection permit N. 48272-6), across

the three major Canga highlands of the Carajás Mineral Province (Fig. 2). For each individual

plant, we collected a sample of root-proximal soil (0-5 cm) for chemical characterization and

leaflet samples for phenotype and genotype analyses. These Canga ecosystems are composed

of several physiognomies, comprising grasslands, scrublands, wetlands and forest formations

(Mota et al., 2015), which differ in terms of the plant communities they support as well as in

their soil chemistry (Mitre et al., 2018). To ensure sampling across environmental gradients,

individuals were collected in each one of these physiognomies within each highland. We also

scattered samples to cover the full extent of each highland (Fig. 2). A minimum distance of 20

meters between samples was used to minimize sampling related individuals. In addition to the

soil samples gathered along with plant tissue, we collected 50 extra soil samples from a highly

degraded mine site (the mine pit from an exhausted mine) and seven soil samples from a

never mined, but moderately disturbed site (scattered across Serra da Bocaina, Fig. 2). These

soil samples were used to predict adaptive genotypes associated with the environmental

conditions of both restoration sites (see details below).

Environmental data

Soil samples were air-dried and sieved using 2 mm mesh, and once dry they were sent to

LABRAS (http://labrasambientaiseagricolas.com.br/) for chemical analyses. These included

pH, organic matter, available P, K, and Na, exchangeable Ca, Mg, and Al, exchangeable S and

available B, Cu, Fe, Mn and Zn (see details in Supporting Information Methods S1). To

reduce the number of chemical parameters describing soil composition, we selected those

known to most affect plant physiology in metalliferous ecosystems (organic matter, Fe, Mn, P,

pH, S, and B (Bothe, 2011; Mitre et al., 2018; Whiting et al., 2004)), along with a set of

orthogonal variables explaining most variation in soil composition across our study area. To

identify this set of orthogonal variables we first used the function *imputePCA* from the

missMDA R package (Josse and Husson, 2016) to impute missing data (20 samples contained

missing data for at least one parameter) using the regularized iterative PCA algorithm

recommended to avoid overfitting (Josse and Husson, 2016). We then ran separate principal

component analyses (PCA) for each species using all the centered and scaled chemical

parameters and selected the three variables showing the strongest correlation with the first, second and third PC axes (each showing eigenvalues > 1 and > 10% of total variance explained, and all three explaining > 50% of total variance; Table S1). The selected soil variables were organic matter, Zn and Na in both species).

We also retrieved climatic data from WorldClim version 1 [1950-2000; (Hijmans et al., 2005)], using the sample coordinates to extract all bioclimatic variables. We followed a similar protocol to obtain the set of orthogonal variables explaining most climatic variance across our study region (Table S1). The first three PCA axes explained 80% of total climatic variance in both species, and the bioclimatic variables most strongly correlated with those axes were isothermality (bio03), minimum temperature of coldest month (bio06) and precipitation of driest quarter (bio17) for *M. acutistipula*; and isothermality (bio03), minimum temperature of coldest month (bio06) and maximum temperature of warmest month (bio05) for *D. apurensis* (see Fig. S1 for maps of these layers). Correlations between these environmental variables were all below |r| < 0.6 (Fig. S2).

Phenotypic data

For each leaf sample we determined macro- and micro-nutrient content and the specific leaf area (SLA), using standard methods (see details in Methods S1). We selected those phenotypic variables known to affect plant physiology in metalliferous ecosystems (SLA, N, B, Fe, Mn, P, N/P, (Bothe, 2011; Mitre et al., 2018; Pérez-Harguindeguy et al., 2013). As described above, we also selected the three orthogonal phenotypic variables explaining most of phenotypic variance (the first three PCA axes explained more than 50% of total variance in both species, Table S1) after imputing missing data (11 samples contained missing data in at

least one parameter). Selected phenotypic variables were Zn, N and B for *M. acutistipula* and

P, Mn and K for *D. apurensis* (correlations between these phenotypic variables were all below

|r| < 0.6; Fig. S2).

Genome size estimation, DNA extraction, genotype-by-sequencing and bioinformatic

processing

We used flow cytometry to estimate haploid genome size in both species (1C DNA content

was 712 Mbp in *M. acutistipula* and 642 Mbp in *D. apurensis,* see details in Methods S1).

Total DNA was extracted using Qiagen's DNeasy Plant Mini Kit. DNA concentration was

quantified using Qubit High Sensitivity Assay kit (Invitrogen), and DNA integrity assessed

through 1.2% agarose gel electrophoresis. Samples with concentrations below 5 ng/µL or

showing no clean bands were excluded from all analyses, and selected samples were

normalized to a concentration of 5 ng/µL and a total volume of 30 µL. These were then

shipped to SNPSaurus (http://snpsaurus.com/) for sequencing and raw data bioinformatic

processing (see details in Methods S1). Briefly, genomic DNA was converted into nextRAD

genotyping-by-sequencing libraries (SNPsaurus, LLC) as in (Russello et al., 2015),

considering the estimated genome size of each species. Genomic DNA was first fragmented

with the Nextera DNA Flex reagent (Illumina, Inc), which also ligates short adapter sequences

to the ends of the fragments. The Nextera reaction was scaled for fragmenting 14 ng and 20

ng of genomic DNA for *M. acutistipula* and *D. apurensis*, respectively. Fragmented DNA was

then amplified for 25 cycles at 75 degrees, with one of the primers matching the adapter and

extending 8 nucleotides into the genomic DNA with the selective sequence TGCAGGAG.

Thus, only fragments starting with a sequence that can be hybridized by the selective

sequence of the primer were efficiently amplified. The nextRAD libraries were then

sequenced on a HiSeq 4000 with six and five lanes of 150 bp reads for *M. acutistipula* and *D.*

apurensis, respectively (University of Oregon). Reads were trimmed using *BBMap tools*

(http://sourceforge.net/projects/bbmap/) to exclude Nextera adapters and a reference contig

was created by collecting 10 million reads in total, evenly distributed from the samples, and

excluding reads that had counts fewer than 6 or more than 800. The remaining loci were then

aligned to each other to identify allele loci and collapse allelic haplotypes to a single

representative. All reads were mapped to the reference contig with an alignment identity

threshold of 95% using *BBMap tools*. Genotype calling was done using *callvariants* (*BBMap*

tools), and the resulting set of genotypes were filtered to remove alleles with population

frequency of less than 3%. Loci that were heterozygous in all samples and loci that contained

more than 2 alleles in a sample (suggesting collapsed paralogs) were removed. A total of

7,165 RAD-tag sequences were obtained for *M. acutistipula* and 4,325 for *D. apurensis.*

Considering the genome size of each species and a linkage block size of 378 Kbp (mean value

for the Fabaceae family, (Lowry et al., 2017)), we estimated a maximum proportion of

genome coverage (assuming one RAD-tag per block) of 100% (McKinney et al., 2017). From

those RAD-tags, 17,403 SNPs were generated for *M. acutistipula* and 9,857 SNPs for *D.*

apurensis (minimum sequencing depth of 14 and 9, respectively).

Genetic diversity and neutral genetic structure

The R package *r2vcftools* (https://github.com/nspope/r2vcftools) - a wrapper for VCFtools

(Danecek et al., 2011) - was used to perform final quality control on the genotype data. To

assess neutral genetic structure and genetic diversity, we used a series of filters to obtain a set

of neutral and independent loci. Filtering criteria included quality (Phred score > 30), read

depth (20 – 800), minor allele frequency (MAF > 0.05), linkage disequilibrium (r^2 < 0.8,

(Xuereb et al., 2018)), Hardy-Weinberg Equilibrium (HWE, p > 0.0001), and loci and

individuals with less than 20% missing data (an example filtering script can be seen in https://

github.com/rojaff/r2vcftools_basics). Additionally, we removed loci potentially under

selection using genome scans. These accounted for population structure (assessed using the

snmf function from the *LEA* package, as described below), and controlled for false discovery

rates by adjusting p-values with the genomic inflation factor (λ) and setting false discovery

rates to q=0.05, using the Benjamini-Hochberg algorithm (François et al., 2016) (see details

below).

We used two complementary genetic clustering approaches to assess neutral

population structure: the *snmf* function from the *LEA* package (Frichot and François, 2015),

and Discriminant Analysis of Principal Components - DAPC from the *adegenet* package

(Jombart and Ahmed, 2011). The *snmf* model implements a fast yet accurate likelihood

algorithm (Frichot et al., 2014), while DAPC is a robust genetic clustering method with no

assumption about the underlying population genetic model (Jombart and Ahmed, 2011).

Based on previous population genomic studies for other co-occurring plant species (Carvalho

et al., 2019; Lanes et al., 2018; Silva et al., 2020), we tested from one to ten ancestral

populations (k). In the case of *snmf* we performed ten replicate runs for each value of k,

choosing the most likely k based on minimized cross-entropy. For DAPC, we inferred optimal

k using k-means clustering and the Bayesian Information Criterion (BIC). Considering the

ancestry coefficients assigned by *snmf*, we then estimated expected heterozygosity (H_E),

inbreeding coefficients (F), and nucleotide diversity (π) for each genetic cluster. We also

estimated pairwise F_{ST} using dartR R package (Gruber et al., 2018), and effective population

sizes (N_e) employing the linkage disequilibrium method implemented in NeEstimator 2.1 and

a lowest allele frequency value of 0.05 (Do et al., 2014). Finally, we assessed fine-scale

spatial genetic structure in each species within each genetic cluster through local polynomial

fitting (LOESS) of Yang's genetic relatedness between pairs of individuals (Yang et al.,

2010) and pairwise geographic distance, as in (Carvalho et al., 2019).

Assessing genotype-environment associations (GEA) and genotype-phenotype associations

(GPA)

To assess GEA and GPA (Fig. 1) we first filtered loci for quality (Phred score > 30), read

depth (20-800), minor allele frequency (MAF > 0.05), linkage disequilibrium ($r^2 < 0.8$), and

loci and individuals with less than 20% of missing data. We then combined univariate and

multivariate methods, namely Latent Factor Mixed Models (LFMM) and Redundancy

Analysis (RDA). While LFMM identifies associations between single loci and single

predictors, RDA can detect multilocus signatures of selection as a function of a multivariate

set of predictors (Caye et al., 2019; Forester et al., 2018). Both methods assume a linear

relationship between allele frequency and environmental variables, have been used

extensively (Ahrens et al., 2018), provide a good compromise between detection power and

error rates, and are robust to a variety of sampling designs and underlying demographic

models (Forester et al., 2018; Rellstab et al., 2015). Since both methods require complete data

sets (without missing values), we performed an imputation of missing genotypes (7.6% and

7% missing genotypes for *M. acutistipula* and *D. apurensis* respectively) based on the *snmf*

population assignments from the previous step, using the *impute* function and the mode

method from the *LEA* package (Frichot and François, 2015). This function imputes missing

genotypes using ancestry and genotype frequency estimates from the *snmf* run.

LFMM analysis were performed using the *lfmm* package (Caye et al., 2019) and ridge

estimates, which minimize regularized least-squares with a L_2 penalty (see example script

here: https://bcm-uga.github.io/lfmm/articles/lfmm). Instead of using raw predictor variables,

we employed the first four axes resulting from a Principal Components Analysis (PCA) on all

predictor variables in order to minimize the number of tests. These four axes explained more

than 60% of total environmental and phenotypic variance in both species, and were strongly

correlated (|r| > 0.7) with organic material, B, Fe, Bio06, Bio17, Zn, S and Na (environmental

variables), and N/P, P, Fe (phenotypic variables) in both species. We ran LFMM using the

previously identified number of genetic clusters (*k*=3, see results) as latent factors, to account

for the underlying neutral genetic structure. We then calculated the genomic inflation factor

(λ) and modified it until a calibrated distribution of adjusted *p*-values was found, and set false

discovery rates at a rate of *q*=0.05 using the Benjamini–Hochberg algorithm (François et al.,

2016).

We performed RDA using the *rda* function from the *vegan* package (Oksanen et al.,

2019) as implemented in Forester et al. (2018), modeling genotypes as a function of predictor

variables, and producing as many constrained axes as predictors (see example script here:

https://popgen.nescent.org/2018-03-27_RDA_GEA.html). Multicollinearity between

predictors was assessed using the variance inflation factor (VIF) and since all predictor

variables showed VIF < 3 none were excluded. Raw predictor variables were scaled and

centered prior to analyses and the population assignments from *snmf* (population ID) were

used to control for population structure by running a partial RDA. Significance of RDA

constrained axes was assessed using the *anova.cca* function and significant axes were then

used to identify candidate loci in both species. Candidate loci were identified using a

Mahalanobis distance-based approach (Capblancq et al., 2018), which made RDA result

comparable with those obtained with LFMM, since it allowed adjusting *p*-values using the

genomic inflation factor (λ) and setting false discovery rates to *q*=0.05, as described above

(calculated and modified genomic inflation factors and *p*-value distributions for LFMM and

RDA tests are provided in Figs. S3-S8). To assess the impact of population genetic structure

on our number of detections we ran additional cluster-level GEA analyses (LFMM and RDA),

using only individuals belonging to the same genetic cluster (setting *k*=1 in LFMM and

omitting population ID in RDA). Finally, to visualize patterns of GEA and GPA, we ran

additional RDA models excluding neutral loci, using the combined candidate adaptive loci

detected using the general RDA and LFMM analyses.

In order to search for the proteins coded by the genes contained in the flanking regions

of our candidate SNPs (found in GEA and GPA analyses), contig sequences containing

candidate loci were first submitted to the EMBOSS Transeq

(http://www.ebi.ac.uk/Tools/st/emboss_transeq/) to obtain corresponding protein sequences.

We used all six frames with standard code (codon table), regions (start-end), trimming (yes),

and reverse (no). We then ran a functional analysis using InterPro

(https://www.ebi.ac.uk/interpro/; interproscan.sh -dp –appl PfamA, TIGRFAM, PRINTS,

PrositePatterns, Gene3d –goterms –pathways -f tsv -o MySequences.tsv -i MySequences.faa),

searching for gene ontology terms and pathways along a variety of annotation databases (i.e.,

Interpro, Pfam, Tigrfam, Prints, PrositePattern and Gene3d).

 Mapping adaptive genetic variation

To map adaptive genetic variation, we used the *adegenet* package (Jombart and Ahmed, 2011)

to run a Spatial Principal Component Analysis (sPCA) on the combined candidate adaptive

loci detected in GEA and GPA analyses using general LFMM and RDA (results for intersected

loci are presented in Fig. S15). sPCA is a spatially explicit multivariate method that yields

scores summarizing genetic variability and spatial structure among individuals (Jombart et al.,

2008). Spatial structure is estimated using a Moran's Index that relies on the comparison of

allelic frequencies observed in one individual to the values observed in neighboring

individuals. These neighboring individuals can be defined by distinct connection networks,

which in our case was set to a distance-based neighborhood, as indicated for aggregated

distributions (Jombart et al., 2008). The Moran's Index generates two types of spatial

structuring: global structure, which reflects positive spatial autocorrelation, and local

structure, that reflects negative spatial autocorrelation (Jombart et al., 2008). To decide if

global and/or local structures should be interpreted and thus retained in sPCA analyses, we

used the global and local tests proposed by Jombart & Ahmed (2011). The first three retained

axes were then interpolated on 10 meter resolution grids covering our study area, and the

resulting rasters used to create an RGB composite, using the Merge function in QGIS 3.4 (see

example scripts here: https://github.com/rojaff/LanGen_pipeline). The resulting color patterns

represent the similarity in adaptive genetic composition.

To predict the adaptive genotypes associated with environmental data collected from

restoration sites (the highly degraded exhausted mine and the moderately disturbed Serra da

Bocaina, Fig. 2), we employed the GEA-RDA models fitted on the combined candidate

adaptive loci detected by global LFMM and RDA (see previous section), and ran the

predict.cca function from the *vegan* package. Environmental samples (soil and climate) from

these sites were thus used to predict RDA scores, based on the fitted GEA-RDA models. We

then performed a *k*-means clustering analysis (using Euclidean distances) on observed and

predicted RDA scores for individuals from each species, using all significant constrained axes

and allowing the number of clusters to vary between two and five (three Canga highlands and

two restoration sites). We used the *NbClust* package (Charrad et al., 2014) to obtain the

optimal number of clusters chosen by 30 different algorithms. Observed and predicted RDA

scores grouping together, suggest that our sampled individuals possess adaptations

associated with the environmental conditions of restoration sites. Observed and predicted

RDA scores placed in different clusters, on the other hand, indicate that none of our sampled

individuals seems adapted to the environmental conditions of restoration sites.

Results

Genetic diversity and neutral genetic structure

After filtering for quality, read depth, minor allele frequencies, missing data, linkage

disequilibrium, Hardy-Weinberg Equilibrium, and outlier loci, we retained 7,376 and 3,496

neutral and independent SNPs and 177 and 163 individuals for *M. acutistipula* and *D.*

apurensis, respectively, which were then used to assess genetic diversity and population

structure. Both genetic clustering approaches (*snmf* and DAPC) indicated the presence of

three clusters in the two study species (Fig. S9). Admixture levels were low, all individuals

were correctly assigned to their source Canga highland (Fig. 2), and there was genetic

differentiation between genetic clusters (pairwise F_{ST} values were significant and ranged

between 0.11 and 0.13 in *M. acutistipula* and between 0.16 and 0.27 in *D. apurensis*).

Expected heterozygosity and nucleotide diversity were similar in both species, but inbreeding

coefficients were lower and effective population sizes larger in *M. acutistipula* (Table 1). Both

species showed significant inbreeding coefficients in all genetic clusters and exhibited the

largest effective population sizes in Serra Sul (Table 1). We detected spatial autocorrelation in

genetic relatedness within genetic clusters in each species (Fig. S10-S11). In both, the strength

of spatial autocorrelation was highest in Serra Sul, where genetic neighborhood size was

larger (~5km, Fig. S10-S11).

Genotype-environment and genotype-phenotype associations

After filtering for quality, read depth, minor allele frequencies, missing data, and linkage

disequilibrium we retained 9,480 and 4,720 SNPs and 177 and 163 individuals for *M.*

acutistipula and *D. apurensis*, respectively. Using LFMM we identified a total of 198 and 154

contigs (RAD-tags) containing GEA, and 94 and 185 contigs containing GPA in *M.*

acutistipula and *D. apurensis*, respectively (Tables S2 and S3). Only the first two constrained

axes from RDA analyses were significant (ANOVA's $p < 0.05$) in GEA and GPA analyses for

both species. RDA revealed a total of 403 and 225 contigs containing significant GEA and

281 and 119 contigs containing significant GPA in *M. acutistipula* and *D. apurensis*

respectively (Fig. 3, Fig. S12 and Tables S2 and S3). In *M. acutistipula* 344 contigs were most

correlated to climatic variables and 69 to soil variables, while in *D. apurensis* 203 contings

were most correlated to climatic and 23 to soil variables. Combining both methods (LFMM

and RDA), we found a total of 588 contigs showing GEA in *M. acutistipula* and 360 in *D.*

apurensis, and 368 contigs showing GPA in *M. acutistipula* and 288 in *D. apurensis*. Only

108 contigs contained both GEA and GPA in *M. acutistipula* and 65 in *D. apurensis.* Finally,

cluster-level GEA analyses revealed many cluster-exclusive detections in both species (Fig.

S13).

Subsequent RDA models using the combined candidate adaptive loci detected using

general LFMM and RDA analyses revealed population-level patterns of GEA and GPA (Fig.

4). In *M. acutistipula* GEA and GPA models explained 17% and 5% of total variance

respectively, while in *D. apurensis* GEA and GPA models explained 31% and 9% of total

variance. In both species, axes loadings were higher for climatic variables (0.01-0.89 for *M.*

acutistipula and 0.01-0.83 for *D. apurensis*) than for soil variables (0.005-0.47 for *M.*

acutistipula and 0.003-0.59 for *D. apurensis*). In *M. acutistipula*, the first and second axes

split individuals into three large GEA groups corresponding to their sampling location. While

individuals from Serra Norte showed associations with higher isothermality (bio03) and

higher winter temperatures (bio06), individuals from Serra Sul showed associations with

warmer winter temperatures and wetter dry season precipitation (bio17). Individuals from

Serra da Bocaina exhibited associations with higher pH (less acidic soils) and drier dry season

precipitation (Fig. 4a). Interestingly, *Dioclea apurensis* showed similar GEA patterns based on

isothermality (bio03), winter temperatures (bio06), and pH, despite using a slightly different

set of predictors (Fig. 4b). On the other hand, the first and second constrained axes divided

individuals into two large GPA groups in *M. acutistipula* (Fig. 4c), the first one encompassing

individuals from Serra Norte (which showed associations with higher SLA and Mn, and lower

P), and the second individuals from Serra Sul and Serra da Bocaina (showing associations

with lower SLA and Mn). In *D. apurensis*, the first and second axes split individuals into three

GPA groups, with individuals from Serra Norte showing associations with a higher leaf

content of Fe and Mn and a lower content of P, while those from Serra Sul showed

associations with higher N and those from Serra da Bocaina with lower SLA and N/P (Fig.

4d). Leaf-level nutrients were weakly correlated with soil-level nutrients (Pearson's

correlation coefficients ranged between -0.07 and 0.39 for *M. acutistipula* and between -0.04

to 0.24 for *D. apurensis*).

A subset of the contigs containing candidate SNPs showed InterPro annotations (105

contigs in *M. acutistipula* and 59 in *D. apurensis*). Candidate adaptive genes were associated

to different functions, including intracellular transport, catalytic activity, synthesis of

hormones, metabolic and oxidation-reduction processes, and plant defense response (a full list

of candidate genes with InterPro annotations is presented in Table S4). Only 17 putative

adaptive genes containing InterPro annotations were shared between both species (Table S5).

Mapping adaptive genetic variation

The combination (union) of candidate adaptive loci detected through GEA and GPA resulted

in 914 loci for *M. acutistipula* and 614 loci for *D. apurensis*. Since none of the sPCA local

structure tests were significant, we retained the first three positive global axes, which

explained most variance in both species (51% and 81% of total variance for *M. acutistipula*

and *D. apurensis* respectively, Fig. S14). These revealed a similar adaptive genetic structure in

both species (Fig. 5), with two adaptive units in Serra Norte and one in Serra da Bocaina.

Mimosa acutistipula nevertheless exhibited a clinal adaptive pattern in Serra Sul, whereas *D.*

apurensis did not. Similar spatial patterns were found when using the intersected loci (i.e.

those shared by GEA and GPA; Fig. S15). Finally, predicted genotypes associated with

climatic and soil characteristics from a highly degraded mining site did not cluster together

with any of our study populations in either species (Fig. 6a and 6b). In contrast, most predicted genotypes for the environmental conditions from the moderately disturbed Serra da Bocaina clustered together with individuals collected in the same location, revealing local provenances are putatively adapted to local environmental conditions (Fig. 6c and 6d).

Discussion

The delineation of seed sourcing areas requires accounting for evolutionary history, genetic diversity, and how likely individuals will adapt to the environmental conditions of the targeted restoration sites (Breed et al., 2019). Here we employ a comprehensive landscape genomic approach to characterize neutral and adaptive genetic variation and provide insights to assist the restoration of a highly degraded mining site and a moderately disturbed Canga highland from the Carajás Mineral Province. We discuss how our results can help define site-adjusted provenancing strategies and argue that our methods can be more broadly applied to assist other restoration and rehabilitation initiatives.

Several studies have stressed the importance of avoiding inbreeding, increasing genetic diversity to maintain evolutionary potential, and minimizing outbreeding depression in restored populations (Broadhurst et al., 2008; Hufbauer et al., 2015; Mijangos et al., 2015; Weeks et al., 2011). The assessment of neutral genetic structure provides information on how to minimize outbreeding depression by avoiding mixing individuals from different evolutionary lineages (Mijangos et al., 2015). Estimates of the genetic neighborhood size, on the other hand, provide clues on how to sample unrelated individuals within seed sourcing areas to increase genetic diversity and reduce the risk of inbreeding depression in restored

populations (Breed et al., 2019; Krauss and Koch, 2004). Our initial assessment of neutral

genetic structure identified three demographically independent units (or Management Units

sensu (Funk et al., 2012)), which could be considered distinct provenances to minimize the

risk of outbreeding depression (Frankham et al., 2017). Within these zones, our estimates of

genetic neighborhood size provide information on within-cluster seed sourcing strategies to

maximize genetic diversity. In Serra Sul, for example, seed sources located 5 Km apart are not

expected to be related (Fig. S10-S11), and would comprise a better representation of standing

genetic variation than individuals collected across smaller spatial scales. Effective population

size estimates (Table 1) nevertheless indicate that none of our observed genetic clusters is

likely to experience inbreeding depression in the near future based on the 50/500 rule

(Jamieson and Allendorf, 2012). Inbreeding levels observed in both species were nonetheless

significantly different from zero, suggesting some level of selfing or mating between related

individuals is taking place.

Patterns of local adaptation will ultimately determine the ability of plants to

effectively colonize and quickly recover disturbed sites (Mijangos et al., 2015). By using the

combined candidate loci detected in GEA and GPA, using both univariate and multivariate

methods, we improved the detection of single-locus and multi-locus adaptive signals.

(Mahony et al., 2019; Talbot et al., 2016; Vangestel et al., 2018). Interestingly, more

intersections between GEA and GPA were found when using RDA than when using LFMM

(Fig. 3), indicating that most adaptations to local environmental conditions expressing

differential phenotypes are polygenic (Forester et al., 2018). Indeed most fitness-related traits

in plants have a polygenic basis (Falke et al., 2013), including tolerance to soil with

phytotoxic levels of heavy metals (Arnold et al., 2016). We nevertheless note that other genes

occurring in the flanking regions of our candidate SNPs could be responsible for the detected

adaptive signals, and that many sequences did not match translated proteins, or found matches

with uncharacterized proteins. Still, the most frequent amongst our identified candidate

proteins are involved in plant defense and stress responses (including reverse transcriptase,

ribonuclease H-like domain, P-loop NTPase fold, leucine-rich repeat, and thaumatin) or basic

metabolic processes (pentatricopeptide repeat, protein kinase domain, Nitrite/Sulfite reductase

ferredoxin-like domain, major intrinsic protein, and kinesin motor domain), suggestive of

adaptations to harsh environments (Tables S4 and S5). Some of these putative adaptive genes

were shared between the two species as well as with other co-occurring species (Table S5),

indicating convergent evolution to similar environmental pressures (Arnold et al., 2016;

Yeaman et al., 2016). These shared genes thus constitute primary targets for functional studies

investigating the molecular basis of adaptation to Canga environments and minelands.

Cluster-level GEA analyses revealed many cluster-exclusive detections in both species

(Fig. S13), suggesting that microgeographic adaptation may play a role in driving genetic

patterns within highlands (Richardson et al., 2014). To visualize and better understand the

mechanisms behind the observed GEA and GPA we ran additional RDAs using the combined

candidate adaptive loci detected in our general LFMM and RDA analyses. As expected, we

found similar patterns of GEA in both species (Figs. 4a and 4b). Interestingly, the strongest

GEA were found with climatic variables in both species, in spite of the coarse resolution of

WorldClim data and the narrow climatic variation found across our study area (Fig. S1). Our

results thus suggest that local climate constitutes an important environmental filter driving

local adaptation, as found in other species from Canga environments (Lanes et al., 2018) and

temperate climates (Pais et al., 2017; Pluess et al., 2016). In *M. acutistipula,* Serra Norte

populations showed associations with higher SLA, suggesting climatic or soil conditions in

Serra Norte are more favorable to plant growth (He et al., 2018). Individuals from Serra da

Bocaina and Serra Sul showed associations with lower SLA and lower concentration of

several micro- and macro-nutrients, suggesting that increasing leaf thickness in these

individuals avoids dissection or better preserves scarce nutrients (Costa-Saura et al., 2016). In

contrast, SLA-associations in *D. apurensis* did not separate Canga highlands, showing that the

influence of climatic variation on SLA is different across species (Gong and Gao, 2019; Liu et

al., 2017). In *D. apurensis*, different genotype associations with leaf micro and macronutrients

separated highlands (Fig. 4d), suggesting different physiological requirements or nutrient

availability at each site. Low correlations between leaf and soil Fe and Mn concentrations,

suggest our study species are controlling nutrient absorption, which makes them suitable for

the restoration of areas with a high concentration of these metals. Controlled common-garden

or reciprocal transplant experiments are nevertheless needed to assess growth and overall

performance of different genotypes (sources) in different soils and climates (Aitken and

Bemmels, 2016; Rellstab et al., 2015).

Our local adaptation maps reveal areas containing similar local adaptations (colors) in

each species (Fig. 5, Fig. S15), which could be used to delineate seed sourcing strategies. In

contrast to the commonly employed Generalized Dissimilarity Models (GDM) (Gugger et al.,

2018; Rossetto et al., 2019; Shryock et al., 2015; Supple et al., 2018), our mapping approach

based on sPCA allows incorporating GPA and predicting adaptive genetic variation from site-

level data, which is particularly useful for areas lacking high-resolution environmental layers.

Moreover, sPCA explicitly account for spatial autocorrelation in genetic composition, which

is likely to play an important role explaining patterns of local adaptation (Lesica and

Allendorf, 1999; Richardson et al., 2014) (see Fig. S16 for alternative adaptation maps

generated using GDM). Our adaptation maps showed similar adaptations across Serra da

Bocaina (i.e. a single adaptive unit), and most predicted genotypes associated with local

environmental samples (climate and soil) clustered together with individuals sampled in Serra

da Bocaina. This result indicates that local provenances are probably best adapted to local

environmental conditions at this site under contemporary climates (Fig. 6c and 6d), and

supports the recommendations made by Lesica & Allendorf (1999) for the restoration of

moderately disturbed sites. Since genetic neighborhood size in Serra da Bocaina was roughly

3 Km for both species, our results suggest that local seeds collected in areas separated by at

least 3 Km would maximize genetic diversity at this location.

In contrast, predicted genotypes for the environmental data collected at the degraded

mine site did not cluster with any of our study populations in either species (Fig. 6a and 6b).

This indicates that none of the genotypes we sampled from natural habitats overlap with the

multivariate environmental conditions present at the mine site. In this case, mixing genotypes

containing different local adaptations could be regarded as the best option to maximize

evolutionary potential and facilitate adaptation to novel environments (Lesica and Allendorf,

1999). Seeds could be sourced from all the identified adaptive units (colors in sPCA maps);

and within these units they could be sampled in areas separated by the genetic neighborhood

size to further enhance genetic diversity. Although mixing individuals from different

management units could result in outbreeding depression (Hufford and Mazer, 2003; Weeks et

al., 2011), the risk is likely marginal for these study species, which are widely distributed

across the continent (Dutra and Morim, 2015; Queiroz, 2015). Moreover, environmental

conditions and plant communities show remarkable similarities across the Carajás Mineral

Province when compared to other *campo rupestre* formations (Zappi et al., 2019). Such a

regional admixture provenancing approach (Bucharova et al., 2019) would represent a

significant improvement over introducing exotic species, which are currently being used in

mine reclamation programs due to the availability of seeds in large quantities and their ability

to quickly colonize mine environments to prevent soil erosion (D'Antonio and Meyerson,

2002; Gastauer et al., 2019; Silva et al., 2018). Our results could guide the establishment of

seed production areas for both native species, aiming to overcome shortfalls in seed

availability while capturing standing neutral and adaptive genetic variation (Nevill et al.,

2016).

Our work illustrates how neutral and adaptive genetic variation can be used to provide

evidence-based recommendations for provenance schemes aimed to effectively restore sites

ranging between moderately disturbed and highly degraded. In our two study species, local

provenances were found optimal to restore a moderately disturbed site, whereas a mixture of

genotypes was suggested as the most promising strategy to recover a highly degraded mining

site, to which local provenances were not adapted. Our proposed methodological approach

(Fig. 1) can be more broadly applied to define site-adjusted provenance strategies in other

locations and for other disturbance regimes. We recognize that the high costs associated with

genomic analyses and the complexity of bioinformatic and statistical analyses represent

important barriers for practitioners (Breed et al., 2019; Shafer et al., 2015). Still, as genomic

data becomes available for more species exhibiting different life-history characteristics,

restoration genomic initiatives using similar methods coupled with visually appealing and

user-friendly interfaces (Rossetto et al., 2019), are likely to substantially improve restoration

outcomes.

Acknowledgments

Funding was provided by Instituto Tecnológico Vale, Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq) grants 301616/2017-5 (RJ), 153535/2018-0 (MG) and

316067/2018-0 (LCRM), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior

(CAPES) grant 88887.156652/2017-00 (CSC). We thank Cesar Neto and Eder Lanes for

assistance in the field, Nelson Carvalho and Santelmo Vasconcelos for assistance in genome

size estimation, Eder Lanes and Manoel Lopes for help in the laboratory, and three

anonymous referees for improving earlier versions of this manuscript.

References

Ahrens CW, Rymer PD, Stow A, Bragg J, Dillon S, Umbers KDL, Dudaniec RY. 2018. The
search for loci under selection: trends, biases and progress. *Mol Ecol* **27**:1342–1356.
doi:10.1111/mec.14549

Aitken SN, Bemmels JB. 2016. Time to get moving: Assisted gene flow of forest trees. *Evol*
Appl **9**:271–290. doi:10.1111/eva.12293

Allendorf FW, Luikart GH, Aitken SN. 2013. Conservation and the genetics of populations.
Oxford: Wiley-Blackwell.

Arnold BJ, Lahner B, DaCosta JM, Weisman CM, Hollister JD, Salt DE, Bomblies K, Yant L.
2016. Borrowed alleles and convergence in serpentine adaptation. *Proc Natl Acad Sci*
113:8320–8325. doi:10.1073/pnas.1600405113

Balkenhol N, Dudaniec RY, Krutovsky K V, Johnson JS, Cairns DM, Segelbacher G, Selkoe
KA, von der Heyden S, Wang IJ, Selmoni O, Joost S. 2017. Landscape Genomics:
Understanding Relationships Between Environmental Heterogeneity and Genomic
Characteristics of Populations. Cham: Springer International Publishing. pp. 261–322.
doi:10.1007/13836_2017_2

Bothe H. 2011. Plants in Heavy Metal SoilsDetoxification of Heavy Metals. pp. 35–57.
doi:10.1007/978-3-642-21408-0_2

Boyer S, Wratten SD. 2010. The potential of earthworms to restore ecosystem services after opencast mining - A review. *Basic Appl Ecol* **11**:196–203. doi:10.1016/j.baae.2009.12.005

Breed MF, Harrison PA, Blyth C, Byrne M, Gaget V, Gellie NJC, Groom SVC, Hodgson R, Mills JG, Prowse TAA, Steane DA, Mohr JJ. 2019. The potential of genomics for restoring ecosystems and biodiversity. *Nat Rev Genet*. doi:10.1038/s41576-019-0152-0

Breed MF, Stead MG, Ottewell KM, Gardner MG, Lowe AJ. 2013. Which provenance and where? Seed sourcing strategies for revegetation in a changing environment. *Conserv Genet* **14**:1–10. doi:10.1007/s10592-012-0425-z

Broadhurst LM, Lowe A, Coates DJ, Cunningham SA, McDonald M, Vesk PA, Yates C. 2008. Seed supply for broadscale restoration: Maximizing evolutionary potential. *Evol Appl* **1**:587–597. doi:10.1111/j.1752-4571.2008.00045.x

Bucharova A, Bossdorf O, Hölzel N, Kollmann J, Prasse R, Durka W. 2019. Mix and match: regional admixture provenancing strikes a balance among different seed-sourcing strategies for ecological restoration. *Conserv Genet* **20**:7–17. doi:10.1007/s10592-018-1067-6

Capblancq T, Luu K, Blum MGB, Bazin E. 2018. Evaluation of redundancy analysis to identify signatures of local adaptation. *Mol Ecol Resour* **18**:1223–1233. doi:10.1111/1755-0998.12906

Carvalho CS, Lanes ÉCM, Silva AR, Caldeira CF, Carvalho-Filho N, Gastauer M, Imperatriz-Fonseca VL, Nascimento Júnior W, Oliveira G, Siqueira JO, Viana PL, Jaffé R. 2019. Habitat Loss Does Not Always Entail Negative Genetic Consequences. *Front Genet* **10**. doi:10.3389/fgene.2019.01101

Caye K, Jumentier B, Lepeule J, François O. 2019. LFMM 2: fast and accurate inference of gene-environment associations in genome-wide studies. *Mol Biol Evol* **36**:852–860.

Charrad M, Ghazzali N, Boiteux V, Niknafs A. 2014. NbClust: An R Package for Determining the Relevant Number of Clusters in a Data Set. *J Stat Softw* **61**:1–36.

Coates DJ, Byrne M, Moritz C. 2018. Genetic Diversity and Conservation Units: Dealing With the Species-Population Continuum in the Age of Genomics. *Front Ecol Evol* **6**:1–13. doi:10.3389/fevo.2018.00165

Costa-Saura JM, Martínez-Vilalta J, Trabucco A, Spano D, Mereu S. 2016. Specific leaf area and hydraulic traits explain niche segregation along an aridity gradient in Mediterranean woody species. *Perspect Plant Ecol Evol Syst* **21**:23–30. doi:10.1016/j.ppees.2016.05.001

D'Antonio C, Meyerson LA. 2002. Exotic Plant Species as Problems and Solutions in
Ecological Restoration: A Synthesis. *Restor Ecol* **10**:703–713. doi:10.1046/j.1526-
100X.2002.01051.x

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter
G, Marth GT, Sherry ST, McVean G, Durbin R, Group 1000 Genomes Project Analysis.
2011. The variant call format and VCFtools. *Bioinformatics* **27**:2156–2158. doi:10.1093/
bioinformatics/btr330

Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR. 2014. NeEstimator v2: Re-
implementation of software for the estimation of contemporary effective population size
(Ne) from genetic data. *Mol Ecol Resour* **14**:209–214. doi:10.1111/1755-0998.12157

Durka W, Michalski SG, Berendzen KW, Bossdorf O, Bucharova A, Hermann J-M, Hölzel N,
Kollmann J. 2017. Genetic differentiation within multiple common grassland plants
supports seed transfer zones for ecological restoration. *J Appl Ecol* **54**:116–126.
doi:10.1111/1365-2664.12636

Dutra VF, Morim MP. 2015. Mimosa in Lista de Espécies da Flora do Brasil. Jardim Botânico
do Rio de Janeiro. *http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB116253*.

Falke KC, Glander S, He F, Hu J, de Meaux J, Schmitz G. 2013. The spectrum of mutations
controlling complex traits and the genetics of fitness in plants. *Curr Opin Genet Dev*
23:665–671. doi:10.1016/j.gde.2013.10.006

Forester BR, Lasky JR, Wagner HH, Urban DL. 2018. Comparing methods for detecting
multilocus adaptation with multivariate genotype–environment associations. *Mol Ecol*
27:2215–2233. doi:10.1111/mec.14584

François O, Martins H, Caye K, Schoville SD. 2016. Controlling false discoveries in genome
scans for selection. *Mol Ecol* **25**:454–469.

Frankham R, Ballou JD, Ralls K, Eldridge M, Dudash MR, Fenster CB, Lacy RC, Sunnucks
P. 2017. Genetic management of fragmented animal and plant populations. Oxford
University Press.

Frichot E, François O. 2015. LEA: An R package for landscape and ecological association
studies. *Methods Ecol Evol* **6**:925–929. doi:10.1111/2041-210X.12382

Frichot E, Mathieu F, Trouillon T, Bouchard G, François O. 2014. Fast and Efficient
Estimation of Individual Ancestry Coefficients. *Genetics* **196**:973–983.
doi:10.1534/genetics.113.160572

Funk WC, McKay JK, Hohenlohe PA, Allendorf FW. 2012. Harnessing genomics for
delineating conservation units. *Trends Ecol Evol* **27**:489–496.
doi:http://dx.doi.org/10.1016/j.tree.2012.05.012

Garris HW, Baldwin SA, Van Hamme JD, Gardner WC, Fraser LH. 2016. Genomics to assist
mine reclamation: A review. *Restor Ecol* **24**:165–173. doi:10.1111/rec.12322

Gastauer M, Souza Filho PWM, Ramos SJ, Caldeira CF, Silva JR, Siqueira JO, Furtini Neto
AE. 2019. Mine land rehabilitation in Brazil: Goals and techniques in the context of legal
requirements. *Ambio* **48**:74–88. doi:10.1007/s13280-018-1053-8

Giannini TC, Giulietti AM, Harley RM, Viana PL, Jaffe R, Alves R, Pinto CE, Mota NFO,
Caldeira Jr CF, Imperatriz-Fonseca VL, Furtini AE, Siqueira JO. 2017. Selecting plant
species for practical restoration of degraded lands using a multiple-trait approach.
Austral Ecol **42**:510–521. doi:10.1111/aec.12470

Gong H, Gao J. 2019. Soil and climatic drivers of plant SLA (specific leaf area). *Glob Ecol*
Conserv **20**:e00696. doi:10.1016/j.gecco.2019.e00696

Gruber B, Unmack PJ, Berry OF, Georges A. 2018. dartr: An r package to facilitate analysis
of SNP data generated from reduced representation genome sequencing. *Mol Ecol*
Resour **18**:691–699.

Gugger PF, Liang CT, Sork VL, Hodgskiss P, Wright JW. 2018. Applying landscape genomic
tools to forest management and restoration of Hawaiian koa (Acacia koa) in a changing
environment. *Evol Appl* **11**:231–242. doi:10.1111/eva.12534

He D, Chen Y, Zhao K, Cornelissen JHC, Chu C. 2018. Intra- and interspecific trait variations
reveal functional relationships between specific leaf area and soil niche within a
subtropical forest. *Ann Bot* **121**:1173–1182. doi:10.1093/aob/mcx222

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution
interpolated climate surfaces for global land areas. *Int J Climatol* **25**:1965–1978.
doi:10.1002/joc.1276

Hufbauer RA, Szucs M, Kasyon E, Youngberg C, Koontz MJ, Richards C, Tuff T, Melbourne
BA. 2015. Three types of rescue can avert extinction in a changing environment. *Proc*
Natl Acad Sci **112**:10557–10562. doi:10.1073/pnas.1504732112

Hufford KM, Mazer SJ. 2003. Plant ecotypes: Genetic differentiation in the age of ecological
restoration. *Trends Ecol Evol* **18**:147–155. doi:10.1016/S0169-5347(03)00002-8

Jamieson IG, Allendorf FW. 2012. How does the 50 / 500 rule apply to MVPs? *Trends Ecol*
Evol **27**:578–584. doi:10.1016/j.tree.2012.07.001

Jombart T, Ahmed I. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP
data. *Bioinformatics* **27**:3070–3071.

Jombart T, Devillard S, Dufour A-B, Pontier D. 2008. Revealing cryptic spatial patterns in
genetic variability by a new multivariate method. *Heredity (Edinb)* **101**:92–103.
doi:10.1038/hdy.2008.34

Josse J, Husson F. 2016. missMDA: A Package for Handling Missing Values in Multivariate
Data Analysis. *J Stat Softw* **70**:1–31. doi:10.18637/jss.v070.i01

Krauss SL, Koch JM. 2004. Rapid genetic delineation of provenance for plant community
restoration. *J Appl Ecol* **41**:1162–1173. doi:10.1111/j.0021-8901.2004.00961.x

Krauss SL, Sinclair EA, Bussell JD, Hobbs RJ. 2013. An ecological genetic delineation of
local seed-source provenance for ecological restoration. *Ecol Evol* **3**:2138–2149.
doi:10.1002/ece3.595

Lanes ÉC, Pope NS, Alves R, Carvalho Filho NM, Giannini TC, Giulietti AM, Imperatriz-
Fonseca VL, Monteiro W, Oliveira G, Silva AR, Siqueira JO, Souza-Filho PW,
Vasconcelos S, Jaffé R. 2018. Landscape Genomic Conservation Assessment of a
Narrow-Endemic and a Widespread Morning Glory From Amazonian Savannas. *Front
Plant Sci* **9**:532. doi:10.3389/fpls.2018.00532

Lesica P, Allendorf FW. 1999. Ecological Genetics and the Restoration of Plant Communities:
Mix or Match? *Restor Ecol* **7**:42–50. doi:10.1046/j.1526-100X.1999.07105.x

Liu M, Wang Z, Li S, Lü X, Wang X, Han X. 2017. Changes in specific leaf area of dominant
plants in temperate grasslands along a 2500-km transect in northern China. *Sci Rep*
7:10780. doi:10.1038/s41598-017-11133-z

Lowry DB, Hoban S, Kelley JL, Lotterhos KE, Reed LK, Antolin MF, Storfer A. 2017.
Breaking RAD: an evaluation of the utility of restriction site-associated DNA sequencing
for genome scans of adaptation. *Mol Ecol Resour* **17**:142–152. doi:10.1111/1755-
0998.12635

Lu M, Krutovsky K V, Loopstra CA. 2019. Predicting Adaptive Genetic Variation of Loblolly
Pine (Pinus taeda L.) Populations Under Projected Future Climates Based on
Multivariate Models. *J Hered* **110**:857–865. doi:10.1093/jhered/esz065

Mahony CR, MacLachlan IR, Lind BM, Yoder JB, Wang T, Aitken SN. 2019. Evaluating
genomic data for management of local adaptation in a changing climate: A lodgepole
pine case study. *Evol Appl* eva.12871. doi:10.1111/eva.12871

Martins K, Gugger PF, Llanderal-Mendoza J, González-Rodríguez A, Fitz-Gibbon ST, Zhao
JL, Rodríguez-Correa H, Oyama K, Sork VL. 2018. Landscape genomics provides

evidence of climate-associated genetic variation in Mexican populations of Quercus
rugosa. *Evol Appl* **11**:1842–1858. doi:10.1111/eva.12684

McKinney GJ, Larson WA, Seeb LW, Seeb JE. 2017. RADseq provides unprecedented
insights into molecular ecology and evolutionary genetics: comment on Breaking RAD
by Lowry et al . (2016). *Mol Ecol Resour* **17**:356–361. doi:10.1111/1755-0998.12649

Mijangos JL, Pacioni C, Spencer PBS, Craig MD. 2015. Contribution of genetics to
ecological restoration. *Mol Ecol* **24**:22–37. doi:10.1111/mec.12995

Mitre SK, Mardegan SF, Caldeira CF, Ramos SJ, Furtini Neto AE, Siqueira JO, Gastauer M.
2018. Nutrient and water dynamics of Amazonian canga vegetation differ among
physiognomies and from those of other neotropical ecosystems. *Plant Ecol* **219**:1341–
1353. doi:10.1007/s11258-018-0883-6

Mota NF de O, Silva LVC, Martins FD, Viana PL. 2015. Vegetação sobre sistemas
ferruginosos da Serra dos Carajás. *Geossistemas Ferruginosos no Bras Inst Pr{\'\i}stino,*
Belo Horiz 289–315.

Mota NF de O, Watanabe MTC, Zappi DC, Hiura AL, Pallos J, Viveros RS, Giulietti AM,
Viana PL. 2018. Cangas da Amazônia: a vegetação única de Carajás evidenciada pela
lista de fanerógamas. *Rodriguésia* **69**:1435–1488. doi:10.1590/2175-7860201869336

Nevill PG, Tomlinson S, Elliott CP, Espeland EK, Dixon KW, Merritt DJ. 2016. Seed
production areas for the global restoration challenge. *Ecol Evol* **6**:7490–7497.
doi:10.1002/ece3.2455

Nunes JA, Schaefer CEGR, Júnior WGF, Neri A V., Correa GR, Enright NJ. 2015. Soil-
vegetation relationships on a banded ironstone 'island', Carajás Plateau, Brazilian
Eastern Amazonia. *An Acad Bras Cienc* **87**:2097–2110. doi:10.1590/0001-
376520152014-0106

Oksanen J, Blanchet F, Friendly, Kindt R, Legendre P, McGlinn D, Minchin P, O'Hara R,
Simpson G, Solymos P, Stevens M, Szoecs E, Wagner H. 2019. vegan: Community
Ecology Package.

Pais AL, Whetten RW, Xiang Q-YJ. 2017. Ecological genomics of local adaptation in Cornus
florida L. by genotyping by sequencing. *Ecol Evol* **7**:441–465. doi:10.1002/ece3.2623

Pérez-Harguindeguy N, Diaz S, Gamier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte
MS, Comwell WK, Craine JM, Gurvich DE, others. 2013. New handbook for stand-
ardised measurement of plant functional traits worldwide. *Aust J Bot* **61**:167–234.

Pluess AR, Frank A, Heiri C, Lalagüe H, Vendramin GG, Oddou-Muratorio S. 2016. Genome-
environment association study suggests local adaptation to climate at the regional scale
in Fagus sylvatica. *New Phytol* **210**:589–601. doi:10.1111/nph.13809

Poveromo JJ. 1999. Iron OresThe Making, Shaping, and Treating of Steel: Ironmaking
Volume. Pittsburg, PA: The AISE Steel Foundation. pp. 547–550.

Preite V, Sailer C, Syllwasschy L, Bray S, Ahmadi H, Krämer U, Yant L. 2019. Convergent
evolution in Arabidopsis halleri and Arabidopsis arenosa on calamine metalliferous soils.
Philos Trans R Soc B Biol Sci **374**. doi:10.1098/rstb.2018.0243

Prober SM, Byrne M, McLean EH, Steane DA, Potts BM, Vaillancourt RE, Stock WD. 2015.
Climate-adjusted provenancing: a strategy for climate-resilient ecological restoration.
Front Ecol Evol **3**. doi:10.3389/fevo.2015.00065

Queiroz LP. 2015. Dioclea in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de
Janeiro. *http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB83091*.

Ramos SJ, Caldeira CF, Gastauer M, Costa DLP, Furtini Neto AE, de Souza FBM, Souza-
Filho PWM, Siqueira JO. 2019a. Native leguminous plants for mineland revegetation in
the eastern Amazon: seed characteristics and germination. *New For*. doi:10.1007/s11056-
019-09704-1

Ramos SJ, Gastauer M, Mitre SK, Caldeira CF, Silva JR, Furtini Neto AE, Oliveira G, Souza
Filho PWM, Siqueira JO. 2019b. Plant growth and nutrient use efficiency of two native
Fabaceae species for mineland revegetation in the eastern Amazon. *J For Res*.
doi:10.1007/s11676-019-01004-w

Rellstab C, Gugerli F, Eckert AJ, Hancock AM, Holderegger R. 2015. A practical guide to
environmental association analysis in landscape genomics. *Mol Ecol* **24**:4348–4370.
doi:10.1111/mec.13322

Richardson JL, Urban MC, Bolnick DI, Skelly DK. 2014. Microgeographic adaptation and the
spatial scale of evolution. *Trends Ecol Evol* **29**:165–176. doi:10.1016/j.tree.2014.01.002

Rossetto M, Bragg J, Kilian A, McPherson H, van der Merwe M, Wilson PD. 2019. Restore
and Renew: a genomics-era framework for species provenance delimitation. *Restor Ecol*
27:538–548. doi:10.1111/rec.12898

Russello MA, Waterhouse MD, Etter PD, Johnson EA. 2015. From promise to practice:
pairing non-invasive sampling with genomics in conservation. *PeerJ* **3**:e1106.
doi:10.7717/peerj.1106

Shafer ABA, Wolf JBW, Alves PC, Bergström L, Bruford MW, Brännström I, Colling G,
Dalén L, De Meester L, Ekblom R, Fawcett KD, Fior S, Hajibabaei M, Hill JA, Hoezel

AR, Höglund J, Jensen EL, Krause J, Kristensen TN, Krützen M, McKay JK, Norman AJ, Ogden R, Österling EM, Ouborg NJ, Piccolo J, Popović D, Primmer CR, Reed FA, Roumet M, Salmona J, Schenekar T, Schwartz MK, Segelbacher G, Senn H, Thaulow J, Valtonen M, Veale A, Vergeer P, Vijay N, Vilà C, Weissensteiner M, Wennerström L, Wheat CW, Zieliński P. 2015. Genomics and the challenging translation into conservation practice. *Trends Ecol Evol* **30**:78–87. doi:10.1016/j.tree.2014.11.009

Shryock DF, Havrilla CA, DeFalco LA, Esque TC, Custer NA, Wood TE. 2017. Landscape genetic approaches to guide native plant restoration in the Mojave Desert. *Ecol Appl* **27**:429–445. doi:10.1002/eap.1447

Shryock DF, Havrilla CA, DeFalco LA, Esque TC, Custer NA, Wood TE. 2015. Landscape genomics of Sphaeralcea ambigua in the Mojave Desert: a multivariate, spatially-explicit approach to guide ecological restoration. *Conserv Genet* **16**:1303–1317. doi:10.1007/s10592-015-0741-1

Silva AR, Resende-Moreira LC, Carvalho CS, Lanes ECM, Ortiz-Vera MP, Viana PL, Jaffé R. 2020. Range-wide neutral and adaptive genetic structure of an endemic herb from Amazonian Savannas. *AoB Plants* **12**. doi:10.1093/aobpla/plaa003

Silva JR, Gastauer M, Ramos SJ, Mitre SK, Furtini Neto AE, Siqueira JO, Caldeira CF. 2018. Initial growth of Fabaceae species: Combined effects of topsoil and fertilizer application for mineland revegetation. *Flora Morphol Distrib Funct Ecol Plants* **246–247**:109–117. doi:10.1016/j.flora.2018.08.001

Skirycz A, Castilho A, Chaparro C, Carvalho N, Tzotzos G, Siqueira JO. 2014. Canga biodiversity, a matter of mining. *Front Plant Sci* **5**:653. doi:10.3389/fpls.2014.00653

Souza-Filho PWM, Giannini TC, Jaffé R, Giulietti AM, Santos DC, Nascimento Jr. WR, Guimarães JTF, Costa MF, Imperatriz- Fonseca VL, Siqueira JO. 2019. Mapping and quantification of ferruginous outcrop savannas in the Brazilian Amazon: A challenge for biodiversity conservation. *PLoS One* **14**:e0211095.

Steane DA, Potts BM, McLean E, Prober SM, Stock WD, Vaillancourt RE, Byrne M. 2014. Genome-wide scans detect adaptation to aridity in a widespread forest tree species. *Mol Ecol* **23**:2500–2513. doi:10.1111/mec.12751

Supple MA, Bragg JG, Broadhurst LM, Nicotra AB, Byrne M, Andrew RL, Widdup A, Aitken NC, Borevitz JO. 2018. Landscape genomic prediction for restoration of a Eucalyptus foundation species under climate change. *Elife* **7**:1–22. doi:10.7554/eLife.31835

Talbot B, Chen T-W, Zimmerman S, Joost S, Eckert AJ, Crow TM, Semizer-Cuming D, Seshadri C, Manel S. 2016. Combining Genotype, Phenotype, and Environment to Infer Potential Candidate Genes. *J Hered* **108**:esw077. doi:10.1093/jhered/esw077

Vangestel C, Eckert AJ, Wegrzyn JL, St. Clair JB, Neale DB. 2018. Linking phenotype,
genotype and environment to unravel genetic components underlying cold hardiness in
coastal Douglas-fir (Pseudotsuga menziesii var. menziesii). *Tree Genet Genomes* **14**:10.
doi:10.1007/s11295-017-1225-x

Weeks AR, Sgro CM, Young AG, Frankham R, Mitchell NJ, Miller KA, Byrne M, Coates DJ,
Eldridge MDB, Sunnucks P, Breed MF, James EA, Hoffmann AA. 2011. Assessing the
benefits and risks of translocations in changing environments: A genetic perspective.
Evol Appl **4**:709–725. doi:10.1111/j.1752-4571.2011.00192.x

Whiting SN, Reeves RD, Richards D, Johnson MS, Cooke JA, Malaisse F, Paton A, Smith
JAC, Angle JS, Chaney RL, Ginocchio R, Jaffre T, Johns R, McIntyre T, Purvis OW, Salt
DE, Schat H, Zhao FJ, Baker AJM. 2004. Research Priorities for Conservation of
Metallophyte Biodiversity and their Potential for Restoration and Site Remediation.
Restor Ecol **12**:106–116. doi:10.1111/j.1061-2971.2004.00367.x

Williams A V., Nevill PG, Krauss SL. 2014. Next generation restoration genetics:
Applications and opportunities. *Trends Plant Sci* **19**:529–537.
doi:10.1016/j.tplants.2014.03.011

Xuereb A, Kimber CM, Curtis JMR, Bernatchez L, Fortin MJ. 2018. Putatively adaptive
genetic variation in the giant California sea cucumber (Parastichopus californicus) as
revealed by environmental association analysis of restriction-site associated DNA
sequencing data. *Mol Ecol* **27**:5035–5048. doi:10.1111/mec.14942

Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath
AC, Martin NG, Montgomery GW, Goddard ME, Visscher PM. 2010. Common SNPs
explain a large proportion of the heritability for human height. *Nat Genet* **42**:565–569.
doi:10.1038/ng.608

Yeaman S, Hodgins KA, Lotterhos KE, Suren H, Nadeau S, Degner JC, Nurkowski KA,
Smets P, Wang T, Gray LK, Liepe KJ, Hamann A, Holliday JA, Whitlock MC, Rieseberg
LH, Aitken SN. 2016. Convergent local adaptation to climate in distantly related
conifers. *Science (80-)* **353**:1431–1433. doi:10.1126/science.aaf7812

Zappi DC, Moro MF, Walker B, Meagher T, Viana PL, Mota NFO, Watanabe MTC, Nic
Lughadha E. 2019. Plotting a future for Amazonian canga vegetation in a campo rupestre
context. *PLoS One* **14**:e0219753. doi:10.1371/journal.pone.0219753
907
908
909

Data Availability Statement

Geographic coordinates, genotypes in Variant Call Format, and sequences in FASTA format

for both species are available in Figshare: https://doi.org/10.6084/m9.figshare.12185235.v1.

All the mentioned R scripts have been deposited in GitHub and their url addresses provided in

the text.

Author contributions

RJ conceived, designed and coordinated the project. CSC, MG and RJ coordinated the field

work and sampling. CSC, SM and MG performed laboratory work. CSC, LCRM, LT, BF, MG

and RJ performed the data analysis. The first draft of the paper was written by CSC and RJ

and all authors contributed to discussing the results and editing the paper.

Tables
922

923 **Table 1:** Genetic diversity measures for *Mimosa acutistipula* var. *ferrea* and *Dioclea*
924 *apurensis* within genetic clusters (highlands). Sample sizes (*N*) are followed by mean
925 expected heterozygosity (H_E), mean inbreeding coefficient (*F*), nucleotide diversity (π) and
926 effective population size (N_e). Values represent 95% confidence intervals.

Species	Highland	N	H_E	F	π	N_e
M. acutistipula	Serra Sul	80	[0.28/0.28]	[0.12/0.15]	[0.26/0.26]	[1601.0/1778.0]
	Serra Norte	61	[0.28/0.28]	[0.10/0.14]	[0.23/0.24]	[554.7/581.2]
	Serra da Bocaina	36	[0.28/0.28]	[0.12/0.17]	[0.25/0.26]	[1008.0/1243.5]
	Total	177	[0.28/0.28]	[0.10/0.17]	[0.23/0.26]	[554.7/1778.0]
D. apurensis	Serra Sul	81	[0.25/0.25]	[0.10/0.14]	[0.22/0.23]	[1070.1/1278.5]
	Serra Norte	45	[0.29/0.29]	[0.14/0.20]	[0.24/0.25]	[193.2/204.6]
	Serra da Bocaina	37	[0.27/0.27]	[0.22/0.31]	[0.21/0.22]	[193.7/212.2]
	Total	163	[0.25/0.29]	[0.10/0.31]	[0.21/0.25]	[193.2/1278.5]

927
928

929 **Figures**
930

Figure 1: Diagram summarizing our methodological approach. We used a subset of neutral
and independent loci to assess broad and fine-scale genetic structure and determine the
genetic neighborhood size. Subsequently, we combined genotype, phenotype, and
environmental data to identify loci under selection and then employed all candidate loci to
map patterns of adaptive genetic variability and predict adaptive genotypes for restoration
sites. The graphs show results for *Mimosa acutistipula.*
937

Figure 2: Maps of our study region depicting restoration sites and neutral genetic structure for *Mimosa acutistipula* var. *ferrea* and *Dioclea apurensis.* **a**: Location of the Carajás Mineral Province within the Amazon biome. **b:** Location of the major Canga highlands (Montane Savanna) where samples were collected. White continuous lines represent the Carajás National Forest (FLONA Carajás), white dashed lines the Campos Ferruginosos National Park (PNCF), and stars depict restoration sites. Ancestry coefficients for all samples from *M. acutistipula* var. *ferrea* (**c**) and *D. apurensis* (**d**), determined using the *snmf* function from the LEA package.

946 **Figure 3:** Venn diagram showing the intersection of contigs (RAD-tags) containing candidate
947 SNPs for *Mimosa acutistipula* var. *ferrea* (**a**) and *Dioclea apurensis* (**b**). Genotype-
948 environment associations (GEA) and genotype-phenotype association (GPA) were assessed
949 using Redundant Analysis (RDA) and Latent Factor Mixed Model (LFMM). The number of
950 detections by method for each species are presented in Tables S2 and S3.

Figure 4: Redundancy analysis (RDA) using only the candidate loci identified in genotype-environment (upper panels) and genotype-phenotype association analyses (lower panels) in *Mimosa acutistipula* var. *ferrea* (**a**, **c**) and *Dioclea apurensis* (**b**, **d**). The plots show the first and second constrained axes from RDA, with SNPs represented as gray filled circles, environment and phenotype variables as blue arrows, and individuals from different Canga highlands in color circles (bio03: isothermality; bio05: maximum temperature of warmest month; bio06: minimum temperature of coldest month; bio17: precipitation of driest quarter; MO: organic material; and SLA: specific leaf area).

Figure 5: Spatial distribution of adaptive genetic variation in *Mimosa acutistipula* var. *ferrea* (**a**) and *Dioclea apurensis* (**b**). The maps represent an RGB composite made using interpolated principal components from a sPCA, ran on the combined candidate loci found in GEA and GPA. Regions with similar colors within each panel represent analogous genetic composition and areas with diagonal lines were not sampled (i.e., adaptive genetic composition was extrapolated from neighboring areas).

Figure 6: Clustering of predicted and observed adaptive genetic variation in *Mimosa acutistipula* var. *ferrea* (a, c) and *Dioclea apurensis* (b, d). A *k*-means clustering approach was employed to examine the similarity between observed (filled symbols) and predicted genotypes (open symbols) associated with the environmental conditions of highly degraded (upper panels) and moderately disturbed (lower panels) sites. Colors indicate different clusters and symbols the locations where samples were collected.