Element content and distribution has limited, tolerance metric dependent, impact on salinity tolerance in cultivated sunflower (*Helianthus annuus*)

A.A. Temme*, V.A. Burns*, L.A. Donovan.

*co-first

Abstract

Disruption of ion homeostasis is a major component of salinity stress's effect on crop yield. In cultivated sunflower prior work revealed a trade-off between vigor and salinity tolerance. Here we determined the association of elemental content/distribution traits with salinity tolerance, both with and without taking this trade-off into account. We grew seedlings of twelve *Helianthus annuus* genotypes in two treatments (0/100 mM NaCl). Plants were measured for biomass (+allocation), and element content (Na, P, K, Ca, Mg, S, Fe, B, Mn, Cu, Zn) in leaves (young and mature), stem, and roots. Genotype tolerance was determined by the proportional decline in biomass and as the deviation from the expected vigor/tolerance trade-off. Genotype rankings on these metrics were not the same. Elemental content and allocation/distribution were highly correlated both at the plant and organ level. Suggestive associations between tolerance and elemental traits were fewer and weaker than expected and differed by tolerance metric. Given the highly correlated nature of elemental content, it remains difficult to pinpoint specific traits underpinning tolerance. Results do show that taking vigor related trade-offs into account is important in determining traits related to tolerance and that the multivariate nature of associated traits should be considered.

keywords: Salt stress, sunflower, sodium, potassium, elemental content, elemental allocation

Running title: Sunflower element content and distribution and salinity tolerance

Introduction

High soil salinity is a major abiotic stress impacting crop yield worldwide. Due to natural processes or as the result of land clearing, unsustainable irrigation practices, and high evaporation high levels of salt (generally NaCl) are present in the soil (Munns and Gilliham, 2015). In these salinized environments, plants experience the twin stresses of a high soil osmotic potential, limiting water uptake, and a high concentration of sodium, leading to ionic stress (Munns, 2002). This ionic stress is caused by plants taking up sodium at the expense of essential elements such as potassium and calcium (Hawkesford *et al.*, 2012). The accumulation of Na to toxic levels, in combination with elemental imbalances within tissues, disrupts cellular and enzymatic functions (Chapin, 1980; Negrão *et al.*, 2017; Morton *et al.*, 2018). With 20% of the worlds irrigated agricultural land affected by salinity (FAO 2005), it is imperative that we improve crop salt tolerance to continue feeding a growing global population.

The effect of salt stress on plant elemental status has been a central focus of research. Given sodium's central role in salt stress, lower Na concentrations in shoot tissues has been shown to correlate with salt tolerance in crops (Flowers and Yeo, 1995; Flowers, 2004). Additionally, given the antagonistic effect of

high sodium concentrations in the soil on potassium uptake, maintenance of potassium levels is
correlated with tolerance as well (Akram *et al.*, 2009; Hawkesford *et al.*, 2012; Demidchik, 2014; Temme
et al., 2019). However, most of these studies are limited to leaf or shoot versus root comparisons,
restricting analyses to correlations between individual elemental concentrations and salt tolerance. By
taking a whole-plant and multiple tissue approach, and incorporating a broader range of elements, we
can infer active mechanisms of elemental allocation, such as exclusion or accumulation (Abrahamson
and Caswell, 1982; Romero and Maranon, 1996) and relate those mechanism to salt tolerance.

While the role of sodium and potassium during salinity stress has been extensively studied (Shabala and
Cuin, 2008; Shabala, 2017; Wu *et al.*, 2018) less focus has been placed on other elements (Broadley *et*
al., 2012; Lewis, 2018). Over the past decade inductively coupled plasma mass spectroscopy (ICP-MS)
has opened up the door for relatively inexpensive analysis of elemental content of plant tissue (Salt *et*
al., 2008), leading to the identification of a host of genes involved in plant elemental content (Whitt *et*
al., 2018). In previous work on cultivated sunflower ICP-MS analysis revealed leaf sulfur content, besides
potassium, to play a role in salinity tolerance (Temme *et al.*, 2019). Given the impacts of salinity stress
on plant nutrient stoichiometry, a comprehensive picture of plant elemental content has the potential
to shed light on mechanisms of salt sensitivity and tolerance (Negrão *et al.*, 2017).

Stress tolerance can be difficult to quantify and numerous metrics for tolerance exist (Zhu *et al.*, 2016;
Morton *et al.*, 2018)). Generally, salinity tolerance is defined as having a high relative performance (i.e.
low proportional decline in performance between control and saline conditions, hereafter
"proportional-decline-tolerance" metric) (Munns and Tester, 2008; Negrão *et al.*, 2017). An ideal crop
cultivar would combine this low proportional decline with high vigor under benign conditions. However,
studies have shown there can be a tradeoff between growth and stress tolerance (Mayrose *et al.*, 2011;
Koziol *et al.*, 2012), and indeed for cultivated sunflower there appears to be a negative trade-off
between vigor (growth in benign conditions) and the proportional effect of salt stress. More vigorous
genotypes have a greater proportional decline in biomass under salt stress (Temme *et al.*, 2019). Given
this trade-off, selecting for the ideal cultivar with high vigor and high tolerance is challenging using the
proportional-decline-tolerance metric.

Fortunately, it is possible to construct an additional tolerance metric that takes this vigor/"tolerance"
trade-off into account. By scoring genotypes on their deviation from the expected effect of salinity
stress based on their biomass, we can redefine tolerance by taking vigor into account. This metric allows
us to disentangle traits related to vigor from traits related to tolerance (performing better than
expected under stress). Finding traits related to this metric of tolerance could allow us to incorporate
them in vigorous genotypes to select for the ideal high-vigor/high-tolerance genotype. However, the
question remains on whether this "expectation-deviation-tolerance" relates to the more conventional
proportional-decline-tolerance metric.

Cultivated sunflower (*Helianthus annuus*), a major oilseed crop, is a moderately salt tolerant species
(Katerji et al. 2000), and previous studies have shown there is genotypic variation in response to salt
stress and elemental status (Ashraf and Tufail, 1995; Shi and Sheng, 2005; Temme *et al.*, 2019). We ask
the following questions in order to determine the effect of taking a known vigor/salinity effect trade-off
into account when determining tolerance and develop a more comprehensive picture of the effects of
salt stress on elemental status and tolerance. (1) Do genotypes differ in the effect of salinity on biomass
and how do genotypes compare for salinity tolerance measured as either proportional-reduction-

tolerance or expectation-deviation-tolerance? (2) Are there genotypic differences in the effect of salinity
on whole plant and organ (leaf, stem, root) level elemental content? (3) Are whole plant and organ level
elemental content and allocation associated with either metric of salinity tolerance?

Materials and Methods

Plant Growth

Achenes from twelve genetically diverse lines of *H. annuus* (Table 3.S1) were germinated at the
University of Georgia greenhouses in a 3:1 sand to turface MVP® (Turface Athletics, PROFILE Products,
LLC, Buffalo Grove, IL) mixture. Ten days after germination, seedlings were transplanted into 30 cm tall,
5L pots filled with the same growth substrate, supplemented with 15ml of lime (Austinville Limestone,
Austinville, VA), 15ml of gypsum (Performance Minerals Corporation, Birmingham, AL), and 39g of 15-9-
12 (N-P-K) Osmocote Plus blend (Osmocote, The Scotts Company, Marysville, OH) per pot. Plants were
arranged in a split plot design with two treatment ponds per each of 4 plots. Three replicate plants per
genotype were placed in each pond. Shallow treatment ponds enabled the bottom 8-10cm of the pots
to be submerged in the pond solution. Open water surfaces were covered with black plastic to reduce
evaporation and algal growth.

After a week of exposure to fresh water to facilitate establishment, each pond was assigned a salinity
treatment of either 0 or 100 mM of sodium chloride (NaCl). All ponds were drained and refilled every
day until the treatment ponds reached the target concentration (due to dilution in the soil water). Once
treatment concentrations were reached after seven days, ponds were drained and refilled every three
days to account for evaporation and maintain a stable concentration within the ponds. To homogenize
the distribution of salt within the pots and to prevent salt crystallization on the surface of the soil, pots
were top-watered with a ~200 ml of pond water every time the water was replaced. After treatment
initiation, plants were grown for eighteen days and then harvested.

Measurements and sample collection

At harvest, plants were measured for height and stem diameter and then separated into young leaves
(leaves above, and including the most recent fully expanded leaf and bud if present), mature leaves
(leaves below the most recent fully expanded leaf, senesced leaves, and cotyledons), stem, and roots.
Roots were thoroughly rinsed with fresh water during the root washing. Biomass samples of these four
tissue types were dried at 60°C and weighed for dry biomass.

In order to have sufficient tissue for analysis of element content, genotype replicate samples within a
pond were pooled for each tissue type, resulting in four biological replicates per tissue type, genotype
and treatment. Biomass samples were coarsely ground using a Wiley ® Mini Mill (Thomas Scientific,
Swedesboro, NJ) and finely ground with a Qiagen TissueLyser (Qiagen, Germantown, MD) for leaf tissue,
and an 8000M Mixer/Mill® High-Energy Ball Mill (SPEX, Metuchen, NJ) for tougher stem and root tissue.
Samples were sent to Midwest Laboratories (Midwest Laboratories, Omaha NB) for Inductively Coupled
Argon Plasma Optical Emission (ICP) Analysis. Analysis provided the elemental content of boron (B),
calcium (Ca), copper (Cu), iron (Fe), potassium (K), magnesium (Mg), manganese (Mn), phosphorus (P),
sodium (Na), and sulfur (S), zinc (Zn) for young leaf, mature leaf, stem, and root tissues.

For each biological replicate, the whole plant elemental content was calculated by multiplying the
elemental content of each tissue sample by the mass of that tissue, summing all tissue type amounts,
and then dividing by the total biomass. The fraction of the element budget allocated to each tissue (e.g.

what percentage of total plant sodium was present in the roots) was then calculated from dividing the
tissue type element amount by the whole plant element amounts. To compare genotypes that could
differ in their organ level element content due to differences in whole plant elemental content we
devised a mass allocation based normalization (after (Romero and Maranon, 1996). For each plant we
calculated the ratio of the organ mass fraction to the fraction of the element budget in that organ. In
this mass relative allocated amount (MRAA) metric, values above one indicate preferential allocation
(more of the element is in that tissue then would be based on neutral distribution) and values below
one indicate exclusion (less of the element is in that tissue then would be expected based on neutral
distribution. To allow for comparison of ratios above and below one we $\log_2$ transformed the ratios so
that a halving and doubling would have the same magnitude (-1, +1). Thus, this MRAA value represents
preferential allocation or exclusion of elements from particular tissue types independent of elemental
status at the whole plant level. We've provided an interactive data sheet that demonstrates the
relationship between mass allocation, organ level elemental content, whole plant elemental content
and MRAA (**Supplemental datasheet 1**).

Data Analysis

All analyses were carried out using R v3.5 (R Foundation for Statistical Computing, Vienna, Austria). For
biomass measurements, to combat issues relating to mortality (19 out of 288 plants died) and missing
samples (8%, due to labeling error) we averaged tissue types per genotype per treatment pond (taking
the mean of natural log transformed weights to minimize skew). This allowed us to construct a complete
dataset based on the "average" plant per genotype per treatment pond. Total biomass and biomass
fractions of tissue types was then calculated based on the pond average of the tissue types.

Genotype means for all traits in control and stressed environments were then calculated using a mixed-
model approach (r package *lme4 (Bates et al., 2015)*) with genotype and treatment as fixed factor and
pond within block as random factor. Then from this model estimated marginal means without the
random factor were calculated (R package *emmeans,* Lenth, 2018). Significance of main effects and their
interaction was determined using Wald's Chi-square test on the mixed model (R package *car,* Fox and
Weisberg, 2011). Using the genotype means per treatment we additionally calculated the change in trait
value between control and treatment as the difference in means.

Given the likely correlations between elements we determined common axes of variation using principal
component analysis. Principal components were calculated for elemental content at the whole plant
level, within each tissue type, and between tissue types at control conditions, salt treatment, and the
difference between treatments. Additionally, we calculate the same principal components for MRAA.
Differences between tissue types in element content and MRAA were calculated using Hotelling's-T test
on the first two principal components.

Salt tolerance of each genotype was calculated in two ways. First, tolerance was calculated as the
proportional change in biomass between both control and salt treatment (taken as the difference
between the natural log of total plant biomass in control and salt treatment). Using this metric,
genotypes having a smaller negative proportional change have a greater salt tolerance. Then, to follow
up on the previous observation that genotypes with high biomass in control conditions have a greater
proportional decrease (Temme *et al.,* 2019), a second measure of tolerance was defined as whether a
genotype performing better or worse than expected based on size in the control treatment. To quantify
this, we fitted a linear regression through the proportional change in biomass vs the natural log of total

biomass at control. Using this metric, genotypes with a more positive residual from this linear fit
perform better than expected and have a greater salt tolerance.

To determine the effect of trait variation on variation in tolerance we ran a series of linear regressions
between tolerance (both metrics) and the first two principal components of whole plant and within
tissue type element content and element MRAA at control, salt stressed, and the difference between
them. For completeness, all individual component traits of the PC analyses were regressed against
tolerance as well. Given the issues of multiple comparisons we limited our interpretation of tolerance
associations to the element principal components and only show individual elements as suggestive
evidence for an effect on tolerance.

All data visualizations were made using ggplot (Wickham, 2009) and ggbiplot
(https://github.com/vqv/ggbiplot)

Results

Genotypic differences in salinity tolerance

Genotypes differed in their absolute biomass and response to salinity (**fig 1a, Table 1**), However,
proportionally (when comparing log transformed values) the difference between genotypes in the effect
of salinity was much more nuanced (**fig 1b**). Across all genotypes we found no evidence for a differential
response to salinity. Proportionally all genotypes had a comparable effect of salinity stress. This
conclusion is however somewhat simplistic. Although there were no differences among genotypes that
rose to the level of significance, there was a significant relationship between vigor (biomass at control
conditions) and the proportional effect of salinity (**fig 1c**). Genotypes with a higher vigor had a greater
proportional decline in biomass due to salinity stress. Thus, we used two different metrics to
characterize tolerance. The first metric was the proportional reduction in biomass: hereafter
"proportional-reduction-tolerance." The second metric took this vigor/salinity effect relationship into
account and was the deviation (residual) from the expected effect: hereafter "expectation-deviation-
tolerance."

The extremes in the rankings on both metrics lined up. Genotypes with the highest or lowest
proportional-reduction-tolerance tended to also be the genotypes with the highest or lowest
expectation-deviation-tolerance. However, genotypes in the middle of the ranking differed substantially
between tolerance metrics leading to no significant (r_s 0.50, *p*=0.099**)** relationship between tolerance
metric ranks (**Fig S1**).

Height and stem diameter, traits strongly related to biomass showed comparable results with biomass
(**Table 1, Figure S2**). Allocation of mass to different plant organs (young leaf, mature leaf, stem, and
root) showed a mixed response with no effect of treatment on leaf mass fraction (LMF) and no
interaction between genotype and treatment on LMF and stem mass fraction. Root mass fraction did
differ significantly between genotypes and treatment including a significant interaction. While not a trait
of focus we did measure specific leaf area (SLA) which had a significant effect of treatment, genotype
and an interaction between them. (**Table 1, Figure S2**)

Whole plant and organ level elements

Depending on element, genotypes differed in their whole plant elemental content and in the effect of salinity thereon. (**Table 1, Figure S3**) Given the highly correlated nature of elemental content, we sought to capture the variation in elemental content in principal component analysis.

At the whole plant level principal component analysis revealed that under control and salt treatment over 70% of the variation in elemental content could be captured by the first two principal components (**Fig 2a,b**). Results showed strong, negative, relationships between whole plant sodium and potassium content and between boron and zinc content. Depending on treatment there were strong positive relationships between elements though these shifted between treatments. Additionally, over 50% of the variation in the response of whole plant elemental content, assessed as the difference between control and salt treatment, was captured in the first two principal components. The positive and negative relationships among shifts in elemental content were similar to those of elemental content itself (**Fig 2c**).

Within organs, over 60% in the variation in element content and MRAA under saline conditions could be captured in the first two principal components (**Figure S4, S5**). Contrasting individual plant organs, again, depending on the element, genotypes largely differed in their organ level elemental content (**Table 2**) and in the mass relative allocated amount (MRAA)(**Table 3**) to their organs (**Figure S6**). Notably, under salt stress, sodium content increased the most in stems and roots and the least in leaves. Potassium content showed the opposite pattern. (**Table 2**). In terms of mass relative allocated amount (MRAA), genotypes were capable of keeping the majority of sodium constrained to the roots and out of the leaves under control conditions (**Table 3, Figure S3**). However, under salt stress the MRAA in leaves became less negative indicating sodium inevitably reached these sensitive tissues. Due to this the level of preferential allocation to the roots also decreased. Surprisingly, MRAA in stems greatly increased, going from nearly neutral to positive showing increased allocation of sodium to stems under saline conditions (**Table 3, Figure S3**). Potassium MRAA showed an opposite pattern.

While genotypes differed in the elemental content and MRAA in a given organ, the difference among organs was far greater than the difference among genotypes. Over 70% of the variation in elemental content between organs could be captured in the first two principal components with organs having a significantly different suite of elemental content. Under benign conditions, genotypes had a greater sodium, copper and iron content in roots and more manganese, boron and calcium in leaves (both young and mature) (**Figure 3a**). Under saline conditions, this changed such that stems and then roots had a greater increase in sodium content and that leaves had a lower decrease in potassium content (**Figure 3b,c**). Separation of tissue types on element MRAA was comparable to element content (**Figure 3d-f**).

Relationships between mass allocation, elements and salinity tolerance.

We performed a series of linear regressions to determine the association between elemental traits and tolerance metrics. Given the large number of comparisons this brings up the issue of false discoveries. Thus, while we present these results, care should be taken in interpreting causative mechanisms from weakly significant results. To partially combat this here we focus on the relationship between the principal components of elemental content and allocation with tolerance instead of each individual element. It must be noted however that even these relationships do not hold up to stringent multiple comparisons correction and should only be interpreted as indicative of potential associations.

At the whole plant level, elemental content was not strongly associated with tolerance (**Table S1**). Only
the first principal component of elemental content in control conditions (**Fig 2a**) was associated with
proportional-reduction-tolerance. The elements most strongly tied to this axis were calcium, sulfur, and
copper (**Table 4**). This result likely recapitulates the vigor/effect of salinity relationship since this same
PC axis is suggestively correlated with vigor under control conditions ($p<0.1$).

As a contributor to variation in MRAA we related changes in biomass allocation to tolerance as well.
Biomass allocation to plant organs was differentially associated with both metrics of tolerance **(Table S2.**
Proportional-reduction-tolerance was negatively correlated with leaf mass fraction at control conditions.
Although this simply recapitulates the correlation found in fig 1c due to the fact that genotypes with
more biomass had a lower leaf mass fraction. Expectation-deviation-tolerance was correlated with the
change in stem mass fraction (genotypes with a lower reduction in stem mass fraction had a lower than
expected decline).

When we related within organ principal components (**Figure S4, S5**) to both metrics of tolerance (**Table
S3**) a contrasting picture emerged. Proportional-reduction-tolerance was associated with the first
principal component of stem elemental content under saline conditions (**Figure 4a**) (with highest
contribution to manganese, phosphorus and sulfur content (**Table 4**)) as well as the second principal
component of the change in MRAA of elements to mature leaves (**Figure 4b**) (with highest contribution
to boron, phosphorous, and calcium MRAA change (**Table 4**)). Expectation-deviation-tolerance was
associated with the second principal component of root elemental content under saline conditions
(**Figure 4c**)(highest contribution to potassium, sodium and boron (**Table 4**)) as well as second principal
component of the change in root elemental content (**Figure 4d**) (highest contributions to sodium,
potassium and sulfur (**Table 4**)).

Surprisingly, the elements that contributed the most to the principal components associated with either
tolerance metric weren't necessarily the most significantly associated with tolerance when looked at all
elements individually. We found putatively significant relationships between element content, MRAA
and both metrics of tolerance but the sheer number of multiple comparisons (>100) lead to these being
weakly suggestive at best. **(Figure S7, Table S4**)

Discussion

Salinity is a major abiotic stress decreasing crop yield by limiting water uptake and disrupting cell
metabolism and ion homeostasis. Mitigating some of the adverse effects requires some combination of
energy intensive sodium exclusion or sequestration in vacuoles and/or non-photosynthesizing tissues
(Tyerman *et al.*, 2019; Munns *et al.*, 2019). Indeed, in cultivated sunflower prior work has shown a
trade-off where genotypes more vigorous in benign conditions are proportionally more affected by
salinity than less vigorous genotypes (Temme *et al.*, 2019). High tolerance could come at the cost of
vigor under benign conditions (Munns and Gilliham, 2015). However, here we show that tolerance can
be redefined by explicitly taking this trade-off into account by scoring genotypes on their deviation from
the expected proportional decline. This expectation-deviation-tolerance allows us to identify different
traits associated with genotype performing better or worse than expected and potentially allows for the
selection of varieties that break away from this vigor/effect of salinity trade-off.

Across twelve cultivated sunflower genotypes, we replicated the vigor/proportional decline in biomass
trade-off (Temme *et al.*, 2019). It should be noted that while this relationship is significant and explains a

moderate portion of the variance in proportional decline (**Fig 1c**, $p<0.05$, R^2_{adj} 0.35) the magnitude of
variation in proportional decline between genotypes is such that it doesn't rise to the level of
significance when simply contrasting genotypes (**Fig 1b**). When contrasting genotypes on two metrics of
tolerance, one based solely on the proportional decline, proportional-decline-tolerance, and one on
deviation from the expected proportional decline based on a genotypes vigor, expectation-deviation-
tolerance, we found genotypes ranking to be uncorrelated (**Fig S7**). Given this discrepancy in ranking we
further explored the link between variation in genotypes elemental traits and the connection with both
tolerance metrics.

Element content and allocation was assessed for eleven elements (Na, P, K, Ca, Mg, S, Fe, B, Mn, Cu, Zn)
in both control and saline conditions as well as the change between them. We found surprisingly strong
correlations between elemental content at the whole plant level (**Fig 2a-b**) and clear differentiation in
elemental content between organs (**Table 2**, **Fig 3a-b**). At the whole plant level, under both benign and
stressed conditions, there were strong negative correlations between sodium and potassium. This likely
reflects variation in uptake selectivity and the antagonistic nature of sodium and potassium uptake
(Blumwald, 2000; Mäser *et al.*, 2002). Additionally, there was a similar negative relationship between
boron and zinc which have a negative relationship in maize (Hosseini *et al.*, 2007) and wheat (Singh *et*
al., 1990). Out of eleven elements, only whole plant zinc content was unaffected by salinity (**Table 1, Fig**
S1). These results show the strong effects of high soil salinity on plant elemental content beyond those
typically measured (generally Na and K (Mäser *et al.*, 2002; Wu *et al.*, 2018).

We further investigated the impact of salinity stress on the content and distribution of elements to
leaves (young and mature), stem, and roots. Prior research has shown elemental distribution to be
element specific and impacted by stress (Conn and Gilliham, 2010). Within organs elemental content
was strongly correlated as well with differences between organs being far greater than differences
between genotypes (**Fig 3a-b**). Our results support the notion that in multivariate space, variation
between tissue types (even between young and mature leaves) is far greater than the variation between
genotypes (Agren and Weih, 2012). Besides elemental content, we investigated the effect of salinity on
the preferential allocation of elements to tissue types. In this mass relative allocated amount (MRAA),
we assessed the deviation from a neutral distribution by calculating the fold change of the proportion of
mass allocated to an organ and the proportion of whole plant elemental amount allocated to the organ
(after (Romero and Maranon, 1996). Given the predominant role of sodium and potassium, we highlight
these elements.

For sodium, this revealed that under benign conditions plants preferentially keep sodium out of the
leaves and keep it in the roots (**Table 3, Fig S3**). Under salinity stress, mechanisms maintaining Na into
the root appear unable to cope and Na content rises throughout the plant (**Table 2**, **Fig S3**). Mass
relative allocation changes as well with less exclusion of sodium from the leaves and relatively less
preferential allocation to the roots. Interestingly, MRAA of sodium to the stem increases (**Table 3, Fig**
S3). These results indicate that mechanisms pulling sodium out of the transpiration stream at the root
level (liken to HKT1 transporters (Munns and Tester, 2008; Hasegawa, 2013)) are overwhelmed but that
mechanisms at the stem or leaf level kick into gear. Possible mechanisms could include high activity of
tonoplast transporters sequestering Na into the vacuole in stem tissue (Bassil *et al.*, 2019), transporters
pulling out sodium in the petiole (Jeschke and Pate, 1991), or active phloem loading, moving Na out of
the leaf (Zhong *et al.*, 1998).

Potassium distribution broadly mirrors sodium with elevated MRAA of potassium to leaves and lower
MRAA in roots under benign conditions. Under saline conditions, this pattern is magnified with even
greater potassium MRAA to leaves and less to roots (**Table 3, Fig S3**). As at the whole plant level
potassium content decreases, this heightened differentiation between plant organs showcases the key
role of potassium in leaf processes ((Hawkesford *et al.*, 2012). In the stems, the greater accumulation
and allocation of sodium is not matched by an increase in potassium MRAA. Since potassium and
sodium have antagonistic effects in the cytosol maintaining a proper balance between these ions is vital
(Munns and Tester, 2008). The discrepancy between stem sodium and potassium content and allocation
would thus suggest sequestration in the vacuole to be a possible mechanism of sodium accumulation
here to maintain cytosolic Na:K ratio's while elevating Na sequestration in the stem tissue (Blumwald,
2000; Zeng *et al.*, 2018; Bassil *et al.*, 2019).

Despite the effects of salinity stress on whole plant elemental content, the mass relative allocated
amount (MRAA) was not dramatically affected for elements other than sodium or potassium.
Interestingly, copper, iron, sodium and zinc have a positive MRAA in roots (both under control and
stressed conditions) whereas for all other elements it's neutral or negative. This is opposite findings in
wheat (Garnett and Graham, 2005) and carrot (Inal *et al.*, 2009). It could be that this reflects ions bound
to the root apoplast from the soil solution (Negrão *et al.*, 2017) but given our thorough washing and the
lower MRAA of other elements in roots, that seems not to be a major factor. Given sunflower's status as
a moderate heavy metal accumulator (Adesodun *et al.*, 2010), this points to active exclusion of these
elements from the shoot or a functional role in the root for these elements.

Linking variation in elemental traits to variation in tolerance has some inherent difficulties in our
experimental design. Given the large number of traits versus a comparatively small number of
genotypes, multiple comparisons corrections are a large dampener on the significance of our findings.
Care must be taken in interpreting our findings, as chance playing a role in our findings cannot be ruled
out. For future work testing hypotheses generated by our results we strongly recommend a larger
number of genotypes to reach greater power to detect links between a continuum of tolerance values
and a host of elemental traits.

Bearing in mind multiple comparisons issues on significance, we found that the associations between
tolerance and elemental traits differed by tolerance metric. Associated with proportional-decline-
tolerance were shifts in mature leaf element allocation and stem elemental content under salt stress.
Associated with expectation-deviation-tolerance were root elemental content and it's shifts under salt
stress. (**Fig 4**) Top elements loading onto these multivariate axes were manganese and boron for
proportional-reduction-tolerance and potassium and sodium for expectation-deviation-tolerance (**Table
4**).

Strikingly, these top loading elements were not necessarily the elements that were associated with
tolerance when looked at individually. Though fraught with multiple comparisons correction issues
(>100 comparisons), when looking at each individual element in each plant part in turn, results support
the role of root potassium content for expectation-deviation-tolerance but also suggest a role for root
and plant sulfur as well as leaf and plant manganese content. Sulfur content has been previously found
to play a role in sunflower salt tolerance (Temme *et al.*, 2019). Root sulpholipid content has been linked
to salinity tolerance (Erdei *et al.*, 1980). Maintenance of manganese content is linked to salt tolerance in
brassica (Chakraborty *et al.*, 2016). Manganese is tied to salinity stress in it's component of manganese

superoxide dismutase which functions as a reactive oxygen species (ROS) scavenger and the oxidative
component of salt stress (Hernández *et al.*, 1994; Wang *et al.*, 2010).

Given the large number of traits increasing the number of genotypes assessed for these elemental traits
will shed more light on significant association with salt tolerance. Moreover, given the short duration of
this experiment a next step will be to track element status across developmental stage and it's impact
on salinity tolerance beyond the seedling stage. Developmental status has a large impact on element
distribution in wheat (Weih *et al.*, 2016), barley (Birsin *et al.*, 2010). With the strong trade-off between
plant vigor (biomass in control conditions) and proportional-reduction-tolerance and the limited overlap
between traits connected with both metrics of tolerance, these results suggest the possibility for
decoupling vigor related traits and tolerance. By decoupling vigor from tolerance, we could select for
high yielding genotypes that are able to maintain their growth under saline conditions. A key open
question then is whether these tolerance and vigor traits share a common genetic basis or if tolerance
related genes could be pyramided into high vigor varieties (Morton *et al.*, 2018).

Here we found that increased sodium sequestration in stems and roots and increased potassium
allocation to leaves appears to be a common response due to salinity. However, given the highly
correlated nature of organ elemental content, it is difficult to pinpoint whether variation in these
specific traits underpins variation in tolerance. What is clear is that the multivariate nature of tolerance
associated traits needs to be considered especially for elemental content. Our results suggest that taking
vigor related trade-offs into account can be vital in determining the traits related to tolerance. By
selecting on traits that allow a genotype to perform better than expected, we could potentially break
free from these trade-offs and maintain high vigor in a saline environment.

Acknowledgements

389 We would like to thank the University of Georgia greenhouse staff and several undergraduate students
390 for their aid in set up and harvest of the experiment. This work was financially supported by grant
391 NSF1444522 to LAD.

References

Abrahamson WG, Caswell H. 1982. On the Comparative Allocations of Biomass, Energy, and Nutrients in Plants. Ecology **63**, 982–991.

Adesodun JK, Atayese MO, Agbaje TA, Osadiaye BA, Mafe OF, Soretire AA. 2010. Phytoremediation Potentials of Sunflowers (Tithonia diversifolia and Helianthus annuus) for Metals in Soils Contaminated with Zinc and Lead Nitrates. Water, air, and soil pollution **207**, 195–201.

Agren GI, Weih M. 2012. Plant stoichiometry at different scales: element concentration patterns reflect environment more than genotype. The New phytologist **194**, 944–952.

Akram MS, Ashraf M, Akram NA. 2009. Effectiveness of potassium sulfate in mitigating salt-induced adverse effects on different physio-biochemical attributes in sunflower (Helianthus annuus L.). Flora **204**, 471–483.

Ashraf M, Tufail M. 1995. VARIATION IN SALINITY TOLERANCE IN SUNFLOWER (HELIANTHUS-ANNUUS L). Journal of Agronomy and Crop Science-Zeitschrift Fur Acker Und Pflanzenbau **174**, 351–362.

Bassil E, Zhang S, Gong H, Tajima H, Blumwald E. 2019. Cation Specificity of Vacuolar NHX-Type Cation/H+ Antiporters. Plant physiology **179**, 616–629.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software, Articles **67**, 1–48.

Birsin MA, Adak MS, Inal A, Aksu A, Gunes A. 2010. MINERAL ELEMENT DISTRIBUTION AND ACCUMULATION PATTERNS WITHIN TWO BARLEY CULTIVARS. Journal of plant nutrition **33**, 267–284.

Blumwald E. 2000. Sodium transport and salt tolerance in plants. Current opinion in cell biology **12**, 431–434.

Broadley M, Brown P, Cakmak I, Ma JF, Rengel Z, Zhao F. 2012. Beneficial Elements. In: Marschner P, ed. Marschner's Mineral Nutrition of Higher Plants (Third Edition). San Diego: Academic Press, 249–269.

Chakraborty K, Sairam RK, Bhaduri D. 2016. Effects of different levels of soil salinity on yield attributes, accumulation of nitrogen, and micronutrients in Brassica spp. Journal of plant nutrition **39**, 1026–1037.

Chapin SFI. 1980. The Mineral Nutrition of Wild Plants. Annual Review of Ecology and Systematics **11**, 233–260.

Conn S, Gilliham M. 2010. Comparative physiology of elemental distributions in plants. Annals of botany **105**, 1081–1102.

Demidchik V. 2014. Mechanisms and physiological roles of K+ efflux from root cells. Journal of plant physiology **171**, 696–707.

Erdei L, Stuiver B (C EE)., Kuiper PJC. 1980. The effect of salinity on lipid composition and on activity of Ca2+- and Mg2+-stimulated ATPases in salt-sensitive and salt-tolerant Plantago species. Physiologia plantarum **49**, 315–319.

Flowers TJ. 2004. Improving crop salt tolerance. Journal of experimental botany **55**, 307–319.

Flowers TJ, Yeo AR. 1995. Breeding for Salinity Resistance in Crop Plants: Where Next? Functional plant biology: FPB **22**, 875–884.

Fox J, Weisberg S. 2011. *An R Companion to Applied Regression, Second Edition*. Thousand Oaks (CA): Sage.

Garnett TP, Graham RD. 2005. Distribution and remobilization of iron and copper in wheat. Annals of botany **95**, 817–826.

Hasegawa PM. 2013. Sodium (Na+) homeostasis and salt tolerance of plants. Environmental and experimental botany **92**, 19–31.

Hawkesford M, Horst W, Kichey T, Lambers H, Schjoerring J, Møller IS, White P. 2012. Functions of Macronutrients. In: Marschner P,, In: Marschner H, eds. Marschner's Mineral Nutrition of Higher Plants. San Diego, UNITED KINGDOM: Elsevier, 135–189.

Hernández JA, del Río LA, Sevilla F. 1994. Salt Stress-Induced Changes in Superoxide Dismutase Isozymes in Leaves and Mesophyll Protoplasts from Vigna unguiculata (L.) Walp. The New phytologist **126**, 37–44.

Hosseini SM, Maftoun M, Karimian N, Ronaghi A, Emam Y. 2007. Effect of Zinc × Boron Interaction on Plant Growth and Tissue Nutrient Concentration of Corn. Journal of plant nutrition **30**, 773–781.

Inal A, Gunes A, Pilbeam DJ, Kadioglu YK, Eraslan F. 2009. Concentrations of essential and nonessential elements in shoots and storage roots of carrot grown in NaCl and Na 2 SO 4 salinity. X-Ray Spectrometry **38**, 45–51.

Jeschke WD, Pate JS. 1991. Ionic Interactions of Petiole and Lamina During the Life of a Leaf of Castor Bean (Ricinus communis L.) Under Moderately Saline Conditions. Journal of experimental botany **42**, 1051–1064.

Koziol L, Rieseberg LH, Kane N, Bever JD. 2012. Reduced drought tolerance during domestication and the evolution of weediness results from tolerance-growth trade-offs. Evolution; international journal of organic evolution **66**, 3803–3814.

Lenth RV. 2018. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.1.

Lewis DH. 2018. Boron: the essential element for vascular plants that never was. The New phytologist.

Mäser P, Gierth M, Schroeder JI. 2002. Molecular mechanisms of potassium and sodium uptake in plants. Plant and soil, 43–54.

Mayrose M, Kane NC, Mayrose I, Dlugosch KM, Rieseberg LH. 2011. Increased growth in sunflower correlates with reduced defences and altered gene expression in response to biotic and abiotic stress. Molecular ecology **20**, 4683–4694.

Morton MJL, Awlia M, Al-Tamimi N, Saade S, Pailles Y, Negrão S, Tester M. 2018. Salt stress under the scalpel - Dissecting the genetics of salt tolerance. The Plant journal: for cell and molecular biology.

Munns R. 2002. Comparative physiology of salt and water stress. Plant, cell & environment **25**, 239–250.

Munns R, Gilliham M. 2015. Salinity tolerance of crops – what is the cost? The New phytologist **208**, 668–673.

Munns R, Passioura JB, Colmer TD, Byrt CS. 2019. Osmotic adjustment and energy limitations to plant growth in saline soil. The New phytologist.

Munns R, Tester M. 2008. Mechanisms of salinity tolerance. Annual review of plant biology **59**, 651–681.

Negrão S, Schmöckel SM, Tester M. 2017. Evaluating physiological responses of plants to salinity stress. Annals of botany **119**, 1–11.

Romero JM, Maranon T. 1996. Allocation of biomass and mineral elements in Melilotus segetalis (annual sweetclover): effects of NaCl salinity and plant age. The New phytologist **132**, 565–573.

Salt DE, Baxter I, Lahner B. 2008. Ionomics and the study of the plant ionome. Annual review of plant

biology **59**, 709–733.

Shabala S. 2017. Signalling by potassium: another second messenger to add to the list? Journal of experimental botany **68**, 4003–4007.

Shabala S, Cuin TA. 2008. Potassium transport and plant salt tolerance. Physiologia plantarum **133**, 651–669.

Shi D, Sheng Y. 2005. Effect of various salt–alkaline mixed stress conditions on sunflower seedlings and analysis of their stress factors. Environmental and experimental botany **54**, 8–21.

Singh JP, Dahiya DJ, Narwal RP. 1990. Boron uptake and toxicity in wheat in relation to zinc supply. Fertilizer Research **24**, 105–110.

Temme AA, Kerr KL, Donovan LA. 2019. Vigour/tolerance trade-off in cultivated sunflower (Helianthus annuus) response to salinity stress is linked to leaf elemental composition. Journal of Agronomy and Crop Science **205**, 508–518.

Tyerman SD, Munns R, Fricke W, *et al.* 2019. Energy costs of salinity tolerance in crop plants. The New phytologist **221**, 25–29.

Wang YC, Qu GZ, Li HY, Wu YJ, Wang C, Liu GF, Yang CP. 2010. Enhanced salt tolerance of transgenic poplar plants expressing a manganese superoxide dismutase from Tamarix androssowii. Molecular biology reports **37**, 1119–1124.

Weih M, Pourazari F, Vico G. 2016. Nutrient stoichiometry in winter wheat: Element concentration pattern reflects developmental stage and weather. Scientific reports **6**, 35958.

Whitt L, Ricachenevsky F, Ziegler G, Clemens S, Walker E, Maathuis FJM, Kear P, Baxter I. 2018. A curated list of genes that control elemental accumulation in plants. bioRxiv, 456384.

Wickham H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wu H, Zhang X, Giraldo JP, Shabala S. 2018. It is not all about sodium: revealing tissue specificity and signalling roles of potassium in plant responses to salt stress. Plant and soil **431**, 1–17.

Zeng Y, Li Q, Wang H, Zhang J, Du J. 2018. Two NHX-type transporters from Helianthus tuberosus improve the tolerance of rice to salinity and nutrient deficiency stress. Plant biotechnology .

Zhong WJ, Kaiser W, Köhler J, Bauer-Ruckdeschel H, Komor E. 1998. Phloem loading of inorganic cations and anions by the seedling of Ricinus communis L. Journal of plant physiology **152**, 328–335.

Zhu M, Shabala S, Shabala L, Fan Y, Zhou MX. 2016. Evaluating Predictive Values of Various Physiological Indices for Salinity Stress Tolerance in Wheat. Journal of Agronomy and Crop Science **202**, 115–124.

Table 1. Overview of plant level traits. Median and range (in parentheses) of trait values of 12 genotypes. Stars note p-value significance of Wald's Chi-square test on treatment (T), genotype (G), and their interaction (TxG). *<0.05, **<0.01, ***<0.001.

Trait	Control (0mM NaCl)	Salt (100mM NaCl)	T	G	TxG
Whole plant mass (g)	5.15 (2.02 - 8.96)	2.94 (1.67 - 6.16)	***	***	***
Young leaf mass (g)	1.26 (0.58 - 2.59)	0.77 (0.47 - 1.85)	***	***	ns
Mature leaf mass (g)	0.9 (0.37 - 1.82)	0.58 (0.22 - 1.12)	***	***	*
Stem mas (g)	1.82 (0.28 - 3)	0.95 (0.29 - 1.92)	***	***	***
Root mass (g)	1 (0.29 - 2.27)	0.76 (0.39 - 1.44)	**	***	**
YLMF	0.26 (0.22 - 0.38)	0.27 (0.22 - 0.35)	ns	***	ns
MLMF	0.21 (0.17 - 0.27)	0.19 (0.13 - 0.24)	ns	ns	ns
LMF	0.46 (0.4 - 0.64)	0.45 (0.38 - 0.55)	ns	***	ns
SMF	0.34 (0.11 - 0.4)	0.28 (0.1 - 0.37)	***	***	ns
RMF	0.2 (0.12 - 0.27)	0.23 (0.19 - 0.36)	***	***	***
Height (cm)	52.4 (11.36 - 61.32)	36.57 (11.92 - 43.13)	***	***	***
Stem diameter (mm)	8.1 (6.63 - 10.08)	6.22 (5.3 - 8.88)	***	***	ns
SLA (cm^2 g^{-1})	368.46 (327.13 - 410.37)	326.95 (252.61 - 355.63)	***	***	**
B (ppm)	7.49 (5.93 - 8.3)	7.76 (6.84 - 10.34)	***	***	*
Ca (%)	0.79 (0.65 - 1.09)	0.7 (0.56 - 0.98)	***	***	*
Cu (ppm)	2.84 (2.58 - 3.7)	3.4 (2.82 - 5.1)	***	***	**
Fe (ppm)	27.48 (18.92 - 45.39)	30.6 (23.61 - 54.35)	**	***	*
K (%)	5.93 (5.54 - 6.82)	4.58 (3.86 - 5.49)	***	***	***
Mg (%)	0.4 (0.36 - 0.58)	0.33 (0.26 - 0.5)	***	***	ns
Mn (ppm)	25.25 (20.05 - 37.96)	34.83 (31.62 - 55.88)	**	***	**
Na (%)	0.12 (0.08 - 0.35)	1.75 (1.21 - 2.59)	***	***	***
P (%)	0.4 (0.33 - 0.58)	0.36 (0.31 - 0.46)	**	***	*
S (%)	0.53 (0.39 - 0.67)	0.44 (0.34 - 0.58)	***	***	***
Zn (ppm)	15.46 (7.42 - 23.73)	15.11 (6.15 - 26.83)	ns	***	ns

Table 2. Organ level elemental content. Median and range (in parentheses) of trait values of 12 genotypes. Stars note p-value significance of Wald's Chi-square test on treatment (T), genotype (G), organ (O) and their interactions. .<0.1, *<0.05, **<0.01, ***<0.001. Letters note significant Tukey post hoc groups based on contrasting all eight organ/treatment groups.

Element	Treatment	Young Leaves	Mature Leaves	Stem	Root	T	G	O	GxT	GxO	TxO	GxTxO
B (ppm)	control	9.31 (7.47 - 14.88) [a]	15.53 (12.5 - 18.35) [c]	2.47 (2.17 - 3.12) [e]	3.04 (2.45 - 3.33) [e]	***	***	***	ns	***	***	ns
	salt	10.9 (8.53 - 15.58) [b]	17.68 (14.83 - 20.03) [d]	2.8 (2.15 - 4.67) [e]	2.86 (2.32 - 3.1) [e]							
Ca (%)	control	0.88 (0.74 - 1.16) [a]	1.12 (0.76 - 1.79) [c]	0.59 (0.52 - 0.72) [d]	0.57 (0.4 - 0.61) [e]	***	***	***	***	***	***	***
	salt	0.82 (0.61 - 0.99) [b]	1.12 (0.77 - 1.8) [c]	0.47 (0.4 - 0.52) [e]	0.51 (0.42 - 0.66) [de]							
Cu (ppm)	control	2.41 (2.05 - 2.6) [ab]	2.03 (1.78 - 2.52) [bc]	1.23 (0.65 - 2.77) [d]	7.28 (6.2 - 9.38) [e]	***	***	***	**	***	ns	*
	salt	2.81 (2.02 - 3.5) [a]	2.72 (1.86 - 3.4) [a]	1.26 (0.95 - 3.17) [cd]	8.15 (5.92 - 9.89) [f]							
Fe (ppm)	control	11.07 (8.9 - 70.53) [a]	11.74 (9.57 - 32.02) [a]	5.31 (3.1 - 12.67) [a]	87.76 (61.95 - 169.98) [b]	ns	***	***	**	***	**	***
	salt	10.27 (6.8 - 13.35) [a]	12.42 (7.92 - 35) [a]	7.31 (3.83 - 15.02) [a]	101.86 (80.33 - 149.54) [b]							
K (%)	control	5.32 (5 - 6.34) [a]	7.37 (6.17 - 8.77) [b]	6.94 (5.45 - 8.5) [c]	3.37 (3.02 - 3.99) [e]	***	***	***	***	***	***	***
	salt	5.47 (4.83 - 6.27) [a]	7.32 (4.98 - 9.24) [bc]	4.62 (3.24 - 5.02) [d]	2.05 (1.49 - 2.51) [f]							
Mg (%)	control	0.38 (0.31 - 0.58) [ab]	0.4 (0.28 - 0.9) [cd]	0.53 (0.46 - 0.66) [e]	0.18 (0.14 - 0.2) [f]	***	***	***	*	***	***	**
	salt	0.38 (0.28 - 0.58) [ac]	0.47 (0.33 - 0.94) [d]	0.33 (0.23 - 0.43) [b]	0.17 (0.13 - 0.19) [f]							
Mn (ppm)	control	29.94 (23.58 - 41.2) [ab]	37.4 (25.7 - 59.25) [cd]	19.35 (15.35 - 23.83) [f]	20.7 (13.15 - 30.9) [f]	**	***	***	***	***	***	***
	salt	39.62 (32.5 - 61.07) [ac]	49.4 (37.1 - 103.07) [e]	26.79 (22.45 - 31.73) [bdf]	26.86 (22.9 - 38.47) [bdf]							
Na (%)	control	0.01 (0 - 0.19) [a]	0.01 (0 - 0.02) [a]	0.12 (0.05 - 0.79) [ab]	0.43 (0.28 - 0.93) [b]	***	***	***	***	***	***	***
	salt	0.42 (0.11 - 1.25) [b]	1.01 (0.37 - 1.84) [c]	2.96 (1.71 - 5.75) [d]	2.58 (1.81 - 2.78) [e]							
P (%)	control	0.61 (0.5 - 0.78) [a]	0.34 (0.28 - 0.53) [c]	0.25 (0.2 - 0.45) [d]	0.42 (0.31 - 0.6) [e]	**	***	***	***	***	***	ns
	salt	0.52 (0.38 - 0.57) [b]	0.29 (0.24 - 0.35) [d]	0.24 (0.2 - 0.38) [d]	0.4 (0.31 - 0.51) [ce]							
S (%)	control	0.64 (0.6 - 0.77) [a]	0.73 (0.62 - 0.94) [c]	0.26 (0.14 - 0.39) [d]	0.51 (0.41 - 0.69) [b]	***	***	***	***	***	***	ns
	salt	0.5 (0.42 - 0.57) [b]	0.65 (0.46 - 0.85) [a]	0.22 (0.17 - 0.29) [d]	0.46 (0.36 - 0.62) [b]							
Zn (ppm)	control	5.68 (4.65 - 8.47) [a]	18.46 (12.25 - 33.37) [bc]	8.77 (2.67 - 36.42) [b]	26.16 (5.7 - 64.67) [cd]	ns	***	***	ns	***	*	***
	salt	5.5 (4.54 - 7.88) [a]	14.1 (9.17 - 33.73) [b]	8.54 (2.04 - 43.06) [b]	35.23 (5.63 - 57.93) [d]							

Table 3. Element mass relative allocated amount (MRAA) per organ. Median and range (in parentheses) of trait values of 12 genotypes. Stars note p-value significance of Wald's Chi-square test on treatment (T), genotype (G), organ (O) and their interactions. .<0.1, *<0.05, **<0.01, ***<0.001. Letters note significant Tukey post hoc groups based on contrasting all eight organ/treatment groups.

Element	Treatment	Young Leaves	Mature Leaves	Stem	Root	T	G	O	GxT	GxO	TxO	GxTxO
B	control	0.36 (0.06 - 0.74) [a]	1.18 (0.74 - 1.26) [b]	-1.5 (-1.79 - -1.27) [c]	-1.28 (-1.49 - -0.98) [d]	ns	*	***	**	***	***	ns
	salt	0.48 (-0.02 - 0.67) [a]	1.23 (0.72 - 1.37) [b]	-1.53 (-1.88 - -1.18) [c]	-1.48 (-1.75 - -1.27) [c]							
Ca	control	0.2 (0.01 - 0.43) [a]	0.52 (0.18 - 0.71) [b]	-0.36 (-1.01 - -0.09) [d]	-0.51 (-1.05 - -0.23) [e]	ns	***	***	ns	***	***	***
	salt	0.2 (-0.08 - 0.42) [a]	0.72 (0.41 - 0.88) [c]	-0.53 (-1.08 - -0.37) [e]	-0.45 (-0.83 - -0.18) [d]							
Cu	control	-0.28 (-0.76 - -0.04) [a]	-0.58 (-0.95 - -0.13) [b]	-1.24 (-2 - -0.44) [c]	1.27 (0.98 - 1.51) [d]	***	***	***	**	***	***	*
	salt	-0.41 (-1 - 0.06) [ab]	-0.37 (-0.98 - 0.02) [ab]	-1.46 (-1.82 - -0.7) [c]	1.13 (0.82 - 1.32) [d]							
Fe	control	-1.11 (-1.98 - 0.14) [a]	-1.03 (-1.87 - -0.5) [a]	-2.41 (-3.36 - -1.54) [c]	1.7 (0.82 - 2.11) [d]	**	***	***	***	***	***	***
	salt	-1.81 (-2.13 - -1.17) [b]	-1.11 (-2.22 - -0.79) [a]	-2.21 (-3.04 - -1.17) [c]	1.66 (1.18 - 1.85) [d]							
K	control	-0.15 (-0.31 - 0.09) [a]	0.31 (0.11 - 0.5) [b]	0.24 (-0.08 - 0.47) [b]	-0.81 (-0.96 - -0.64) [d]	*	***	***	.	***	***	***
	salt	0.23 (0.06 - 0.49) [b]	0.62 (0.29 - 0.76) [c]	-0.11 (-0.45 - 0.19) [a]	-1.17 (-1.64 - -0.83) [e]							
Mg	control	-0.08 (-0.35 - 0.09) [a]	0.07 (-0.39 - 0.64) [c]	0.38 (0.18 - 0.75) [d]	-1.17 (-1.67 - -0.9) [e]	***	***	***	ns	***	***	***
	salt	0.22 (0.02 - 0.37) [b]	0.47 (0.02 - 0.91) [d]	-0.02 (-0.66 - 0.48) [ac]	-1.04 (-1.55 - -0.68) [f]							
Mn	control	0.26 (-0.2 - 0.56) [a]	0.53 (0.27 - 0.66) [b]	-0.44 (-0.94 - -0.13) [c]	-0.38 (-0.99 - -0.05) [c]	ns	***	***	ns	***	ns	***
	salt	0.21 (-0.12 - 0.45) [a]	0.56 (0.07 - 0.87) [b]	-0.48 (-0.81 - -0.21) [c]	-0.37 (-0.84 - -0.05) [c]							
Na	control	-4.09 (-5.56 - -2.61) [a]	-4.17 (-5.87 - -2.7) [a]	-0.18 (-1.26 - 0.39) [d]	1.58 (1.13 - 2.04) [f]	***	***	***	*	***	***	ns
	salt	-2.38 (-3.8 - -0.74) [b]	-1.11 (-2.47 - 0.03) [c]	0.72 (-0.51 - 1.71) [e]	0.5 (-0.48 - 1.16) [e]							
P	control	0.57 (0.42 - 0.77) [a]	-0.16 (-0.44 - 0.18) [c]	-0.63 (-0.98 - -0.41) [e]	0.05 (-0.3 - 0.33) [f]	ns	*	***	ns	***	***	**
	salt	0.43 (0.14 - 0.72) [b]	-0.33 (-0.58 - -0.12) [d]	-0.63 (-0.74 - -0.19) [e]	0.15 (-0.09 - 0.31) [f]							
S	control	0.33 (-0.1 - 0.66) [a]	0.55 (0.26 - 0.78) [c]	-1.04 (-1.53 - -0.52) [d]	-0.06 (-0.19 - 0.37) [e]	ns	***	***	ns	***	***	*
	salt	0.16 (-0.17 - 0.54) [b]	0.55 (0.32 - 0.83) [c]	-0.99 (-1.28 - -0.57) [d]	0.01 (-0.15 - 0.31) [e]							
Zn	control	-1.45 (-1.81 - -0.28) [a]	0.08 (-0.44 - 1.67) [bc]	-0.7 (-1.54 - 0.61) [d]	0.27 (-0.61 - 1.65) [bc]	ns	ns	***	ns	***	*	***
	salt	-1.36 (-2.12 - -0.1) [a]	-0.17 (-1.23 - 1.66) [b]	-0.75 (-1.38 - 0.66) [d]	0.97 (-0.91 - 1.63) [c]							

Table 4. Element loadings on PCA axes linked with tolerance. Fraction of variation in element content or MRAA that loads onto the PCA axes that suggestively (non-significant after multiple comparison correction) associate with variation in tolerance (*Table S4*). Rank of fraction explained in parentheses. WP: whole plant, M leaf: mature leaf,

	Proportional-reduction-tolerance			Expectation-deviation-tolerance	
	WP content	**Stem content**	**M leaf MRAA**	**Root content**	**Root content**
Element	**PC1 (control)**	**PC1 (salt)**	**PC1 (delta)**	**PC2 (salt)**	**PC2 (delta)**
B	0.19 (9)	0.30 (7)	**0.61** (1)	**0.29** (3)	0.02 (9)
Ca	**0.88** (1)	>0.01 (10)	**0.53** (3)	0.01 (10)	0.06 (8)
Cu	**0.76** (3)	0.60 (4)	0.43 (5)	0.05 (6)	0.06 (7)
Fe	0.34 (7)	0.01 (9)	>0.01 (11)	0.06 (5)	0.36 (4)
K	0.09 (11)	0.50 (6)	0.02 (9)	**0.75** (1)	**0.44** (2)
Mg	0.75 (4)	>0.01 (11)	0.17 (8)	0.23 (4)	0.12 (6)
Mn	0.75 (5)	**0.78** (1)	0.53 (4)	0.05 (7)	0.02 (10)
Na	0.18 (10)	0.53 (5)	0.40 (6)	**0.42** (2)	**0.60** (1)
P	0.69 (6)	**0.73** (2)	**0.55** (2)	0.02 (9)	>0.01 (11)
S	**0.81** (2)	**0.63** (3)	0.01 (10)	0.05 (8)	**0.38** (3)
Zn	0.25 (8)	0.14 (8)	0.19 (7)	>0.01 (11)	0.17 (5)

Figure captions

Figure 1. Effect of salinity on biomass and tolerance to salinity. (a) Genotype mean biomass at harvest in control and saline (100 mM NaCl) conditions. (b) Natural log transformed biomass values at control and salt treatment. (c) Relationship between biomass under control conditions and the proportional decline in biomass under salt treatment. Genotypes with more biomass under control conditions have a greater proportional decline in biomass under salt treatment (p<0.05). Different colors denote different genotypes.

Figure 2. Multivariate view of whole plant elemental content. First and second principal components of whole plant elemental content under (a) control treatment, (b) salt treatment, and (c) the difference between treatments. Under both treatments >70% of the variability in elemental content for 11 macro and micro nutrients was captured in the first two principal components. >60% of the variability in the change of elemental content was captured in the first two principal components. Different colors denote different genotypes.

Figure 3. Multivariate view of organ level elemental content. First and second principal component of organ level elemental content under (a) control treatment, (b) salt treatment, and (c) the difference between treatments. Organ types, leaf (young and mature), stem, and root, were significantly differentiated with the first two PC axes capturing >70% of the variability between tissues. Letters denote Hottelings-T significant differences on the first two PC axes.

Figure 4. Relationship between "significant" organ level elemental content principal components and tolerance. Proportional-reduction tolerance and (a) stem element content under salt treatment PC1 and (b) the change in mature leaf element MRAA PC2. Expectation-deviation-tolerance and (c) root element content under salt treatment PC2 and (d) the change in root element content PC2. Panel insets show the PCA graph corresponding to the panel. Note that while all slopes are p<0.05, these are non-significant after multiple comparisons correction (Table 4).

Figue S1. Genotypes ranking on tolerance metrics. Each genotype's rank on expectation-deviation-tolerance versus proportional-reduction-tolerance. Spearman's correlation of ranks is non-significant (r_s=0.50, p=0.099), dashed line notes 1:1 line.

Figure S2. Phenotypic traits measured at harvest. Stem Diam (stem diameter), CCI (chlorophyll content index), Plant height, Specific Leaf Area (SLA), Young Leaf Mass Fraction (YLMF), Mature Leaf Mass Fraction (MLMF), Leaf Mass Fraction (LMF), Stem Mass Fraction (SMF), and Root Mass Fraction (RMF). Different colors note different genotypes. Insets show genotype, treatment and their interaction p-value (Wald's chi-square). .<0.1, *<0.05, **<0.01, ***<0.001.

Figure S3. Whole elemental content reaction norms between control and salt (100mM NaCl) treatment. Different colors note different genotypes. Insets show genotype, treatment and their interaction p-value (Wald's chi-square). .<0.1, *<0.05, **<0.01, ***<0.001.

Figure S4. Within organ multivariate view of elemental content. First and second principal components of per organ elemental content under salt treatment and of the difference between control and salt treatment. Different colors denote different genotypes.

Figure S5. Within organ multivariate view of element mass relative allocated amount (MRAA). MRAA shows the log2 fold difference between the proportion of element allocated to an organ and the proportion of mass allocated to an organ. First and second principal components of per organ MRAA under salt treatment and of the difference between control and salt treatment. Different colors denote different genotypes.

Figure S6. Organ specific reaction norms of elemental content. Set of figures (one per element) displaying organ level (Young Leaf, Mature Leaf, Stem, Root) reaction norms of elemental content (top row), element distribution (middle row), and Mass Relative Allocated Amount (MRAA, bottom row). MRAA shows the log2 fold difference between the proportion of element allocated to an organ and the proportion of mass allocated to an organ. Different colors note different genotypes. Insets show genotype, treatment and their interaction p-value (Wald's chi-square). .<0.1, *<0.05, **<0.01, ***<0.001.

Figure S7. Tolerance and element correlation. Multi page supplement on the correlation between tolerance, defined as the residuals of the log biomass/proportional decrease in biomass relationship and organ level elemental content under salt treatment (page 1), the change in elemental content (page 2), and the proportional decrease in biomass and organ level elemental content under salt treatment (page 3), the change in elemental content (page 4). Individual elements are arranged in rows with organ classes in columns. Young leaf (YL), mature leaf (ML), stem (S), Root (R), and whole plant (Total).

Whole Plant Element fraction – control
Whole Plant Element fraction – salt
Whole Plant Element fraction – delta

PC1 (51.8% explained var.)
PC2 (21.9% explained var.)

PC1 (50.6% explained var.)
PC2 (23.7% explained var.)

PC1 (35.2% explained var.)
PC2 (25.0% explained var.)

groups

22   178   233   263
65   185   237   271
73   191   250   278